Title: Naked mole-rat cortical neurons are resistant to acid-induced cell

death

Authors: Zoé Husson[1], Ewan St John Smith[1]

[1] Department of Pharmacology, University of Cambridge, Tennis Court Road CB2 1PD, United

Kingdom

Corresponding author: Ewan St. John Smith, Department of Pharmacology, University of

Cambridge, Tennis Court Road, Cambridge, CB2 1PD, United Kingdom, Tel.: +44 1223 334048;

Fax: +44 1223 334100; E-mail: es336@cam.ac.uk

Co-author contact details: Zoé Husson, Department of Pharmacology, University of Cambridge,

Tennis Court Road, Cambridge, CB2 1PD, United Kingdom, Tel.: +44 1223 334006; Fax: +44

1223 334100; E-mail: zmah2@cam.ac.uk

Abstract

Regulation of brain pH is a critical homeostatic process and changes in brain pH modulate various ion channels and receptors and thus neuronal excitability. Tissue acidosis, resulting from hypoxia or hypercapnia, can activate various proteins and ion channels, among which acid-sensing ion channels (ASICs) a family of primarily Na^+ permeable ion channels, which alongside classical excitotoxicity causes neuronal death. Naked mole-rats (NMRs, *Heterocephalus glaber*) are long-lived, fossorial, eusocial rodents that display remarkable behavioral/cellular hypoxia and hypercapnia resistance. In the central nervous system, ASIC subunit expression is similar between mouse and NMR with the exception of much lower expression of ASIC4 throughout the NMR brain. However, ASIC function and neuronal sensitivity to sustained acidosis has not been examined in the NMR brain. Here, we show with whole-cell patch-clamp electrophysiology of cultured NMR and mouse cortical and hippocampal neurons that NMR neurons have smaller voltage-gated Na^+ channel currents and more hyperpolarized resting membrane potentials. We further demonstrate that acid-mediated currents in NMR neurons are of smaller magnitude than in mouse, and that all currents in both species are fully blocked by the ASIC antagonist benzamil. We further demonstrate that NMR neurons show greater resistance to acid-induced cell death than mouse neurons. In summary, NMR neurons show significant cellular resistance to acidotoxicity compared to mouse neurons, contributing factors likely to be smaller ASIC-mediated currents and reduced NaV activity.

Key words

ASIC, acid-induced currents, acidotoxicity, naked mole-rat, hippocampus

Background

Acid-sensing channels (ASICs) are ion channels of the ENaC/Deg superfamily and most

subunits are activated by extracellular protons [1,2]. Six different ASIC subunits are encoded by

4 ASIC genes (ASIC1a, ASIC1b, ASIC2a, ASIC2b, ASIC3 and ASIC4), which assemble as homo-

or heterotrimers [3]; neither ASIC2b nor ASIC4 form proton-sensitive homotrimers. ASICs are

primarily permeable to Na^+, although ASIC1a homomeric channels are also Ca^{2+} permeable [2].

In the central nervous system (CNS), neurons have been shown to primarily express ASIC1a

homomers and heteromers of ASIC1a/2a and ASIC1a/2b [4–8], where they have been

demonstrated to have key roles in synaptic plasticity [9–11] and fear conditioning [12–14], as well

as being major players in neuronal death resulting from brain ischemia [5,15–17], and

neurodegenerative diseases [18–20].

Regulation of brain pH is a highly complex and important process [21]. Brain tissue

acidosis can result either from an increase in tissue partial pressure of carbon dioxide (PCO_2)

during hypercapnia, or from the accumulation of the byproducts of anaerobic metabolism, such

as lactate and protons, during hypoxia [22]. During periods of tissue acidosis, activation of ASICs

by extracellular acidification is worsened by the release of allosteric modulators such as lactate

[23], spermine [16] and arachidonic acid [24,25]. In addition to the activation of the Ca^{2+} permeable

ASIC1a channel [15,16], a drop in pH also modulates the activity of numerous others ion

channels, including voltage-gated ion channels [26–29] and glutamate receptors [30,31],

therefore leading to disturbance in ion homeostasis, excitotoxicity and ultimately neuronal death

[5,15,16].

Naked mole-rats (NMRs, *Heterocephalus glaber*) are subterranean rodents belonging to

the Bathyergidae African mole-rat family found in East Africa [32]. Unusually for a mammal, NMRs

are eusocial [33,34]. However, NMRs also display a range of remarkable physiological

peculiarities, which is beginning to make a significant impact on biomedical research [35]. The unusual physiology of the NMR includes: extreme longevity with no increased risk of death with ageing [36], an apparent absence of age-related neurodegenerative disorders [37,38], resistance to cancer [39–41], insensitivity to certain noxious and irritant stimuli [42–45] and hypoxia/hypercapnia resistance resulting from altered NMDA receptor function and an ability to utilize fructose as an energy source [46–49]. It is striking that NMRs are resistant to many pathological conditions known to involve ASICs. Recordings of ASIC-mediated currents in dorsal root ganglion (DRG) sensory neurons demonstrated an increased frequency and magnitude of ASIC responses in NMR neurons compared to mouse neurons [43], with APETx2, an inhibitor of ASIC3-containing ASICs, demonstrating a key role for ASIC3, even though nmrASIC3 does not appear to form functional homotrimers [50]. Previously we mapped out ASIC expression in different NMR brain regions and observed similar expression between mouse and NMR, a key exception being much lower ASIC4 levels throughout the NMR brain [51], however, no one has yet studied the function of ASICs in NMR brain neurons.

In this study, we investigated acid-induced currents in mouse and NMR neurons using whole-cell patch clamp recording of cultured neonatal hippocampal and cortical neurons. We find that NMR neurons have ASIC-mediated currents of significantly smaller peak current amplitude than those recorded from mouse neurons and that NMR neurons are resistant to acid-induced cell death. Overall, these results suggest that the reduced acid-induced cell death in NMR neurons may be neuroprotective.

Methods

Animals

All experiments were conducted in accordance with the United Kingdom Animal (Scientific Procedures) Act 1986 Amendment Regulations 2012 under a Project License (70/7705) granted

to E. St. J. S. by the Home Office; the University of Cambridge Animal Welfare Ethical Review Body also approved procedures. Breeding couples of 1 male and 2 female C57/bl6 mice were conventionally housed with nesting material and a red plastic shelter; the holding room was temperature-controlled (21 °C) and mice were on a normal 12-hour light/dark cycle with food and water available *ad libitum*. Naked mole-rats were bred in house and maintained in a custom-made caging system with conventional mouse/rat cages connected by different lengths of tunnel. Bedding and nesting material were provided along with running wheels and chew blocks. The room was warmed to 28 °C and humidified, with a heat cable to provide extra warmth running under 2–3 cages, and red lighting (08:00–16:00) was used.

Neuronal cultures

P0-P2 mice and P0-P5 naked mole-rats were used to prepare cortical and hippocampal neuronal cultures. Multiple pups (2-4) were used to prepare a single culture. Following decapitation, heads were immediately placed in dishes containing ice-cold Hank's Balanced Salt Solution (HBSS) solution (20 mM HEPES, 30 mM glucose in HBSS, Life Technologies). Brains were removed, transferred to a new dish and the two hippocampi and cortices were isolated. Tissues were subsequently incubated in an enzymatic digestion solution: 2 mg/mL papain (Worthington Biochemical Corporation) in Hibernate-Ca^{2+} solution (Brain Bits), activated by 0.5 mM Glutamax (Life Technologies) at 37 °C for 30 min in a 5 % CO_2 incubator. The digestion solution was then replaced by HBSS solution supplemented with DNAse I (250 Kunitz units/mL, Sigma Aldrich) and tissues were slowly triturated (5-7 times) using a P1000 pipette. Neuronal suspensions were filtered through a 100 μm nylon cell strainer (Corning) to remove non-dissociated pieces of tissues before centrifugation for 5 mins at 1100 rpm at room temperature. Supernatants were discarded and the pellets carefully resuspended in HBSS solution. After further centrifugation for 5 min 1100 rpm at room temperature, pellets were resuspended in MEM/HS solution: 10 % heat-inactivated horse serum (Life Technologies), 2 mM glucose, 0.0025 % Glutamax, and 0.2 mg/mL primocin (InVivogen). Hippocampal and cortical neurons were plated on 35 mm plastic dishes (Fisher

Scientific), previously coated with 100 mg/mL poly-L-lysine (Sigma-Aldrich), rinsed with water and dried, at a density of 300 000 cells/mL (2 mL/dish). After a 4-hour incubation in a 37 °C / 5 % CO_2 incubator, the MEM/HS solution was removed and the dishes were flooded with Neurobasal/B27 solution (1X B27 Supplement, 0.0025 % Glutamax, and 0.2 mg/mL primocin). Naked mole-rat neurons were kept at 33 °C in 5 % CO_2 incubator, whereas mouse neurons were kept at 37 °C in a 5 % CO_2 incubator; this is due to NMRs being cold-blooded and NMR cells do not withstand 37 °C for long periods of time [52]. Half of the medium was exchanged for fresh medium every 2-3 days until the cultures were used for experiments.

Electrophysiology

Hippocampal and cortical neurons from mouse and NMR were used for whole-cell patch-clamp recordings at 9-12 days *in* vitro (DIV9-12). Recordings were performed at room temperature using the following solutions: extracellular (in mM) – 140 NaCl, 4 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 4 glucose, 10 HEPES, adjusted to pH 7.4 with NaOH and 300–310 mOsm with sucrose; intracellular (in mM) – 110 KCl, 10 NaCl, 1 $MgCl_2$, 1 EGTA, 10 HEPES, 2 Na_2ATP, 0.5 Na_2GTP, adjusted to pH 7.3 with KOH and to 310–315 mOsm with sucrose. Acidic extracellular solutions were made using MES (pH 5.0). Patch pipettes were pulled (Model P-97, Flaming/Brown puller; Sutter Instruments,) from borosilicate glass capillaries (Hilgenberg GmbH) and had a resistance of 6-10 MΩ. Data were acquired using an EPC10 amplifier and Patchmaster software (HEKA). Whole-cell currents were recorded at 20 kHz, pipette and membrane capacitance were compensated using Patchmaster macros, and series resistance was compensated by >60 %. Cell capacitances and resting membrane potentials were measured just after cell opening in whole-cell configuration. To study macroscopic voltage-gated currents, a standard voltage-step protocol was used whereby cells were held at -120 mV for 200 msecs before stepping to the test potential (-80 mV - +65 mV in 5 mV increments) for 50 msecs, returning to the holding potential (-60 mV) for 200 msecs between sweeps. In some experiments, tetrodotoxin (300 nM, Alomone Labs) was perfused for 30 seconds before repeating the voltage-step protocol. To measure neuronal acid-sensitivity, cells were

exposed to the following protocol: 5 seconds of pH 7.4; 5-seconds of pH 5; and 5 seconds of pH

7.4. ASIC antagonists (100 µM Benzamil, Sigma) were perfused during 30 seconds before

applying another 5 second pulse of pH 5. After 90 second wash time with pH 7.4 solution, a 5

second pulse of pH 5 was applied to check for reversal of any block observed. Current amplitude

was measured in Fitmaster (HEKA) by taking the maximum peak response and subtracting the

mean baseline amplitude in the preceding 50 msec (voltage-gated currents) or ~2.5 sec (ASIC

currents); current amplitude was normalized for cell size by dividing by cell capacitance. Using

Igor Pro, for each individual cell that underwent the voltage-step protocol, the following equation

was fitted to the normalized inward currents:

10
$$i(x) = \Gamma \cdot x \cdot \frac{1 - e^{-\frac{x - Erev}{25mV}}}{1 - e^{-\frac{x}{25mV}}} \cdot \frac{1}{(1 - e^{-\frac{x - VHalf}{slope}})^3}$$

11   where Erev is the reversal potential; Vhalf the half-activating potential; Γ a constant and x the

12   command potential. Similarly, a Boltzmann equation was fitted to the normalized outward

13   currents:

14
$$\frac{I}{Imax} = \frac{1}{1 + e^{\frac{(VHalf - Vm)}{slope}}}$$

where Vm is the membrane voltage and VHalf the voltage at half-maximal activation. To

determine the inactivation time of the ASIC-mediated currents, a single exponential was fitted.

Data are expressed as mean ± standard error of the mean (SEM). Using Prism (GraphPad),

paired t-tests were used to compare the effects of ASIC antagonists on proton-gated currents

within both mouse and NMR neuron datasets; unpaired t-tests were used to compare parameters,

such as neuronal resting membrane potential and capacitance and ASIC-mediated current

amplitude, between mouse and NMR neuron datasets.

Acid-induced cell death assays

Mouse and NMR neurons were used to measure acid-induced cell death at DIV9-12. The pH 7.4

and pH 5 extracellular solutions used were the same as those described above for

electrophysiology experiments. Neuronal cultures from both mouse and NMR were rinsed twice

with 37 °C solution (pH 7.4 or pH 5) and then incubated with pH 7.4 or pH 5 solution for 2 hours

at 37 °C. Cultures from both conditions were then rinsed with warm pH 7.4 solution and incubated

during 30 min with pH 7.4 solution containing 1.5 mM propidium iodide (PI, Sigma Aldrich) to stain

necrotic cells and Hoechst 33342 (dilution 1/2500, Sigma) to label all nuclei. Labelled cultures

were imaged using an epifluorescence microscope (Olympus) equipped with a 20X objective

(Olympus) and a QImaging camera. To determine the percentage of dead necrotic PI-positive

cells, we used the software Fiji to count the total number of cells per field of view by counting

nuclei on the Hoechst images, and subsequently counting the number of PI-positive nuclei on the

PI images. One to three dishes per condition were used, and three different images per dish were

taken. Data were collected from three different cultures and each culture was prepared from

multiple animals. A one-way ANOVA test (Prism, GraphPad) corrected for multiple comparisons

(Tukey test) was used to compare the percentage of cell death in each field of view at pH 7.4 and

pH 5 in both species. Data are expressed as mean ± standard error of the mean (SEM).

16　Results

Basic electrophysiological properties and voltage-gated Na$^+$ channel activity differ between NMR

and mouse neurons

Electrophysiological recordings from NMR neurons have been performed in both DRG

sensory neurons [43,50] and CNS neurons [46,47,53]. However, neuronal activity from NMR CNS

has only been recorded in brain slices, in the form of field excitatory postsynaptic potentials, and

the basic electrophysiological properties of NMR neurons in hippocampal and cortical cultures

have not yet been described.

We first compared the capacitance and resting membrane potential of NMR and mouse

neurons from both cortical and hippocampal neuronal cultures (Fig. 1). The capacitance of NMR

neurons was significantly smaller than in mouse neurons in both cortical and hippocampal

cultures (cortex: 17.27 ± 1.02 pF versus 28.72 ± 2.32 pF for NMR (n = 30) and mouse (n = 24)

neurons, respectively; hippocampus: 17.41 ± 1.03 pF versus 38.13 ± 3.27 pF for NMR (n = 18)

and mouse (n = 26) neurons, respectively; unpaired two-sided t-tests, **** $p < 0.0001$, Fig. 1a).

Resting membrane potentials were measured as soon as the whole-cell configuration was

established and NMR neurons were significantly more hyperpolarized than mouse neurons, in

both cortical and hippocampal cultures (cortex: -57.03 ± 2.64 mV versus -44.05 ± 2.84 mV for

NMR (n = 30) and mouse (n = 21) neurons, respectively; hippocampus: -55.11 ± 4.79 mV versus

-43.85 ± 2.59 mV for NMR (n = 18) and mouse (n = 26) neurons, respectively; unpaired two-sided

t-tests; ** $p < 0.01$; * $p < 0.05$, Fig. 1b).

We then investigated macroscopic voltage-gated currents in NMR and mouse neurons,

using a voltage-step whereby cells were held at -120 mV for 200 msecs before stepping to the

test potential (-80 mV to +65 mV in 5 mV increments) for 50 msecs, and returning to the holding

potential (-60 mV) for 200 msecs between sweeps (Fig. 1c). Both NMR and mouse neurons

showed inward and outward currents (Fig. 1c-e). In some experiments, the voltage-step protocol

was run twice, the second time after 300 nM tetrodotoxin (TTX) had been applied for 30 seconds

to investigate the contribution of TTX-sensitive voltage-gated Na^+ channels (NaVs) to the

macroscopic voltage-gated inward currents recorded in NMR neurons (Fig. 1c, right panel). In

both cortical and hippocampal NMR neurons, the voltage-gated inward currents were fully blocked

by 300 nM TTX (n = 8 and n = 2 for cortical and hippocampal neurons, respectively). The fact that

no inward current remained after application of 300 nM TTX indicates that only NaVs were

activated with our voltage-step protocol and that there was no measurable contribution of voltage-

gated Ca^{2+} channels to the inward currents recorded. Moreover, these results indicate that cortical

and hippocampal NMR neurons only express TTX-sensitive NaVs.

Strikingly, NMR neurons had a significantly smaller inward peak current density in both

cortical and hippocampal cultures (cortex: 28.79 ± 3.83 pA/pF versus 202.40 ± 21.35 pA/pF for

NMR (n = 19) and mouse (n = 17) neurons, respectively; hippocampus: 39.00 ± 6.39 pA/pF versus

196.70 ± 28.28 pA/pF for NMR (n = 13) and mouse (n = 16) neurons, respectively; unpaired two-

sided t-tests, **** p < 0.0001, Fig. 1d.i-ii). Additionally, the voltage of half-activation (Vhalf) and

the peak inward current amplitude potential (peak Vm) were more depolarized in NMR neurons

compared to mouse neurons (cortex: Vhalf: -36.12 ± 2.80 mV versus -45.42 ± 1.99 mV and peak

Vm: -12.43 ± 3.02 mV versus -31.27 ± 2.61 mV for NMR (n = 19) and mouse (n = 17) neurons,

respectively; hippocampus: Vhalf: -37.10 ± 2.34 mV versus -45.71 ± 1.30 mV and peak Vm: -

17.62 ± 4.00 mV versus -30.68 ± 3.09 mV for NMR (n = 13) and mouse (n = 16) neurons,

respectively; unpaired two-sided t-tests, **** p < 0.0001, ** p < 0.01; * p < 0.05, Fig. 1d.iii-iv).

These results suggest that NMR neurons may be less excitable compared to mouse neurons,

with more hyperpolarized resting membrane potentials and smaller voltage-gated inward currents

that are activated at more depolarized potentials, i.e. a greater depolarizing stimulus is required

to activate NMR voltage-gated inward currents that produce much smaller currents.

By contrast, current-voltage curves for voltage-gated outward currents were similar

between NMR and mouse neurons from both cortical and hippocampal neurons (Fig. 1e.i) and

Vhalfs were not significantly different (cortex: 7.13 ± 1.91 mV versus 10.77 ± 2.18 mV for NMR (n

= 24) and mouse (n = 17) neurons, respectively; hippocampus: 5.15 ± 2.50 mV versus 11.59 ±

2.26 mV for NMR (n = 14) and mouse (n = 16) neurons, respectively; unpaired two-sided t-tests,

p = 0.221 and p = 0.094 for cortex and hippocampus respectively, Fig. 1e.ii).

Acid-induced currents are mediated by ASICs in both NMR and mouse neurons

The expression profile of ASIC subunits is similar in mouse and NMR brains, with the

exception of lower levels of ASIC4 throughout the NMR brain [51], results suggesting that

functional ASIC-mediated currents should be present in NMR as others have shown in mouse

[4,5,7,8,15].

A 5 second pulse of pH 5 was applied to NMR and mouse neurons from both hippocampal and cortical neurons and rapidly activating and inactivating acid-induced responses were recorded in every cell of both species (Fig. 2a). However, the peak current density of acid-mediated responses recorded in NMR neurons was significantly smaller than in mouse neurons, in both hippocampal and cortical neurons (cortex: 15.41 ± 1.82 pA/pF versus 85.66 ± 10.90 pA/pF for NMR (n = 31) and mouse (n = 24) neurons, respectively; hippocampus: 20.54 ± 2.86 pA/pF versus 100.90 ± 17.68 pA/pF for NMR (n = 22) and mouse (n = 26) neurons, respectively; unpaired two-sided t-tests, **** p < 0.0001, *** p < 0.001, Fig. 2b). By contrast, the inactivation time constant of the acid-mediated currents was similar between NMR and mouse neurons (cortex: 0.35 ± 0.012 s versus 0.44 ± 0.056 s for NMR (n = 27) and mouse (n = 18) neurons, respectively, unpaired two-sided t-test, p=0.0516; hippocampus: 0.47 ± 0.06 s versus 0.35 ± 0.04 s for NMR (n = 13) and mouse (n = 19) neurons, respectively; unpaired two-sided t-test, p = 0.0861, Fig. 2c).

The transient nature of the acid-mediated inward currents in both NMR and mouse neurons is characteristic of ASIC-mediated currents [3] and to confirm the involvement of ASICs we utilized the non-selective ASIC antagonist benzamil [54]. After a first application of pH 5 for 5 seconds, we applied 100 µM benzamil for 30 seconds before a second pH 5 pulse, then followed by a wash period of 90 seconds (Fig. 2d, 2g). In cortical and hippocampal mouse neurons, the acid-induced currents were reversibly blocked by 100 µM benzamil (cortex: pH 5: 133.60 ± 19.01 pA/pF; benzamil: 16.77 ± 5.28 pA/pF; wash: 133.80 ± 19.88 pA/pF (n = 9); hippocampus: pH 5: 151.00 ± 33.64 pA/pF; benzamil: 7.84 ± 2.67 pA/pF; wash: 89.26 ± 20.48 pA/pF (n = 10); one-way paired ANOVA test, Tukey's multiple comparison test, *** p < 0.001, ** p < 0.01, Fig. 2e-f). This result concurs with previous studies indicating that acid-evoked currents in mouse hippocampal and cortical neurons are mediated by ASICs [4,8,15].

Similarly, in cortical and hippocampal NMR neurons, acid-induced currents were reversibly inhibited by 100 µM benzamil (cortex: pH 5: 9.66 ± 1.09 pA/pF; benzamil: 2.78 ± 0.49

pA/pF; wash: 9.13 ± 0.89 pA/pF (n = 7); hippocampus: pH 5: 11.13 ± 1.88 pA/pF; benzamil: 3.50

± 0.53 pA/pF; wash: 8.15 ± 1.30 pA/pF (n = 6); one-way paired ANOVA test, Tukey's multiple

comparison test, *** p < 0.001, ** p < 0.01, Fig. 2h-i). This is the first demonstration of functional

ASIC-mediated currents in CNS NMR neurons.

NMR neurons are resistant to acid-induced cell death

ASICs are involved in acid-induced cell death, so-called acidotoxicity, which can occur

during periods of ischemia [5,15,16]. Because NMR neurons exhibit significantly smaller ASIC

currents (Fig. 2), we hypothesized that this could be neuroprotective when neurons are in an

acidic environment. We exposed neuronal cultures from mouse and NMR cortices to a pH 7.4 or

a pH 5 solution for 2 hours at 37 °C. Nuclei were stained by Hoechst 33342 and necrotic (dead)

neurons were labeled using propidium iodide (PI) (Fig. 3). Percentages of dead neurons were

calculated by counting the number of necrotic PI-positive cells over the total number of Hoechst

33342-positive neurons in the field of view, and a one-way ANOVA test corrected for multiple

comparisons (see *Methods*) was used to compare neuronal death at pH 5 between both species

(Fig. 3c). At pH 7.4, percentages of dead cells were comparable between mouse and NMR cortical

cultures (mouse - pH 7.4: 20.31 ± 3.50 %, n = 18 fields of view; NMR – pH 7.4: 13.31 ± 1.53 %,

n = 27 fields of view; from 3 independent experiments), suggesting no differences in neuronal

death under basal conditions between NMR and mouse cultures. When incubated for 2 hours in

pH 5 solution, mouse cortical neurons exhibited a significantly increased percentage of cell death

compared to all other conditions (mouse – pH 5: 60.55 ± 4.87 %, n = 18 fields of view; NMR – pH

5: 22.29 ± 2.30 %, n = 28 fields of view, from 3 independent experiments, p < 0.0001, ****). By

contrast, when NMR cortical neurons were exposed to pH 5 for 2 hours, no difference in the level

of cell death was observed compared to incubation at pH 7.4. Similar results were obtained using

mouse and NMR hippocampal neurons (data not shown). This indicates that NMR neurons are

resistant to acid-induced cell death, possibly due to their reduced ASIC currents compared to mouse neurons.

Discussion

In this study, we recorded for the first time from cultured NMR brain neurons and described their basic electrophysiological properties (Fig. 1). We showed that NMR neuronal capacitance was smaller than in mouse neurons and that the resting membrane potential of NMR neurons is more hyperpolarized than in mouse neurons. One point to consider is that neurons were recorded from blindly and so we can only broadly compare hippocampal and cortical neurons between species and cannot comment on if these differences occur in all types of neuron or specific neuronal subpopulations, such as interneurons. Although capacitance and resting membrane potential values obtained from mouse neurons were similar to those reported by others using cultured rodent neurons [55,56], for NMR neurons however, previous data from others found their resting membrane potential to be more hyperpolarized (NMR cortical neurons: -57.03 ± 2.64 mV, NMR hippocampal neurons: -55.11 ± 4.79 mV, Fig. 1; NMR hippocampal pyramidal neurons (4 months old): -70.3 ± 6.1 mV, NMR hippocampal dentate granule cells (4 months old): -75.1 ± 3.6 mV, [53]). Explanations for differences observed in resting membrane potentials between studies are the recording conditions (*in vitro* vs. *in vivo*), the developmental stage of the animal from which neurons are obtained and the constituents of the intracellular and extracellular solutions. When comparing our study with that of Penz and colleagues, who reported more hyperpolarized resting membrane potentials, two factors that could contribute to the difference observed is that we made recordings from cultured neurons, whereas they used brain slices, and secondly, we isolated neurons from neonatal animals, whereas slice recordings were made from animals aged at least 4-months.

A standard voltage-step protocol that has been already successfully used in NMR sensory neurons to predominantly isolate NaV activity [43] was used to measure macroscopic voltage-gated current activity in mouse and NMR neurons. NaV currents recorded from mouse neurons were not different to what have been recorded in similar experimental conditions (for example, mouse cortical neurons at DIV 10-12: Vhalf: -41.62 ± 1.46 mV in [57] versus this study : -45.42 ± 1.99 mV). Similarly, inward currents recorded in NMR brain neurons were also very similar to currents recorded in NMR sensory neurons (NMR cortex: Vhalf: -36.12 ± 2.80 mV; NMR hippocampus: Vhalf: -37.10 ± 2.34 mV; NMR DRG neurons: Vhalf: from -34.2 ± 0.1 mV to -44.6 ± 0.5 mV [43]). However, NMR inward currents significantly differed from mouse currents: the peak current density was significantly smaller and the Vhalf and peak Vm values were more depolarized. These differences in NMR NaV activity likely result from differences in amino acid sequence of NaV subunits and/or differential expression of accessory subunits and warrant further investigation. We also found that addition of 300 nM TTX completely abolished all voltage-gated inward currents demonstrating an absence of TTX resistant NaV subunits in the NMR brain, this result aligns with the observation that the TTX-resistant NaV subunits, NaV1.5, NaV1.8 and NaV1.9 are not expressed in the central nervous system [58]. It should also be noted that although NMR neurons were cultured at 32 °C under hypoxic (3 % O_2) conditions, recordings under normoxic, standard laboratory conditions, whereas NMRs live in a hypoxic and hypercapnic environment [59,60], which may influence channel activity *in vivo*.

ASIC activity in brain neurons is now accepted as a key factor in numerous physiological and pathological conditions [61]. However, nothing is known about acid-induced responses in NMR brain neurons, which is of considerable interest considering the behavioral hypoxia/hypercapnia resistance and lack of acid-induced nocifensive behavior display by NMR that are likely adaptations to adapting to a safe, but relatively hypoxic and hypercapnic habitat [35]. Recently, we described ASIC subunit expression in the NMR CNS [62], which is similar to that in the mouse CNS, with the exception of lowered ASIC4 levels in the NMR brain. However,

evidence for functional ASIC activity is lacking and is of particular interest in light of our recent

finding that nmrASIC3 forms non-functional homomers [50]. Here, we find that NMR brain neurons

produce ASIC-mediated currents in response to acid stimulation, in both hippocampus and cortex,

as demonstrated by acid-induced responses being fully, reversibly blocked by 100 μM benzamil

(Fig. 2). However, the peak current density of NMR ASIC-mediated responses was significantly

reduced compared to responses recorded in mouse neurons. The reasons for such reduction in

ASIC currents in NMR neurons is not known and additional research is needed to determine what

underpins this different, e.g. are regulators of ASIC plasma membrane trafficking different in

NMR? Is there a different developmental expression profile of ASICs between mouse and NMR?

With regard to the ASIC currents themselves, they have similar decay kinetics in both NMR and

mouse neurons (Fig. 2c), which suggests that a similar mixture of ASIC subunits are expressed,

as our previous mRNA based analysis suggested [62].

Incubation with a pH 5.0 solution showed that unlike mouse neurons, NMR neurons do

not undergo any significant acid-induced cell death (Fig. 3). This is the first demonstration of

resistance to acid-induced neuronal death in NMR neurons. In rodent neurons, some factors

shown to be protective against acid-induced neuronal injury, include lower temperature [63],

pharmacological blockade or genetic deletion of ASIC activity [15] or ASIC trafficking [64], i.e. it

is established that ASICs play a key role in acidotoxicity. Considering the similar prevalence of

ASIC currents, we suggest that the reduced ASIC-mediated current amplitude observed in NMR

neurons may be an additional neuroprotective mechanism in NMR brains, alongside the

previously described increased hypoxia-inducible transcription factor (HIF1-α) expression [65] or

more efficient *in vivo* CO_2 buffering [47]. One possible mechanism for the decreased amplitude

observed is reduced ASIC plasma membrane trafficking which is known to be modulated by an

extracellular acidic environment [64]. However, it is also possible that the reduced acid-induced

cell death observed is not ASIC-dependent because although ASIC activation appears to play a

major role in neuronal injury [61], several other molecular players are also modulated by a drop

in extracellular pH, such as NaVs [27,28] and glutamate receptors [31] and may contribute to the lowered acid-induced cell death observed, for example, here we also show that NMR neurons also have smaller NaV-mediated currents, which may also add a layer of neuroprotection.

Conclusions

In this work, we describe for the first time the basic electrophysiological properties of NMR neurons in culture and showed that the resting membrane potential of NMR neurons is more hyperpolarized, as well as the amplitude of NaVs being smaller than that of mouse neurons. We then demonstrated that acid-induced currents are present in NMR neurons and are, as in mouse, ASIC-mediated. The key result is that acid-induced cell death is virtually absent in NMR neurons, with reduced ASIC and NaV amplitudes likely contributing to this observation, and thus this is a further adaptation enabling NMR to live in a subterranean, hypercapnic/hypoxic environment.

Abbreviations

ASIC, acid-sensing ion channel; CNS, central nervous system; DRG, dorsal root ganglion, NaV, voltage-gated Na$^+$ channel; NMR, naked mole-rat; TTX, tetrodotoxin

Declarations

Ethics

All experiments were conducted in accordance with the United Kingdom Animal (Scientific Procedures) Act 1986 Amendment Regulations 2012 under a Project License (70/7705) granted to E. St. J. S. by the Home Office; the University of Cambridge Animal Welfare Ethical Review Body also approved procedures.

Consent for publication

N/A

Availability of data and material

The datasets supporting the conclusions of this article are available in the University of Cambridge

Apollo repository: https://doi.org/10.17863/CAM.18704

Competing interests

The authors have no competing interests.

Funding

This work was supported by an Isaac Newton Trust Research Grant from the University of

Cambridge. ZH was funded by a EMBO Long-Term Fellowship (ALTF1565-2015).

Authors' contributions

ZH and ESS designed the study. ZH performed all the experiments and analysed the data. ZH

and ESS wrote the manuscript.

Acknowledgements

Thanks to all members of ESS lab for their useful discussions, and in particular to the assistant

staff in the Department of Pharmacology and Animal Facility for their technical help and animal

husbandry.

References

1. Gründer S, Pusch M. Biophysical properties of acid-sensing ion channels (ASICs).

Neuropharmacology. 2015;94:9–18.

2. Waldmann R, Champigny G, Bassilana F, Heurteaux C, Lazdunski M. A proton-gated cation

channel involved in acid-sensing. Nature. 1997. p. 173–7.

3. Hesselager M, Timmermann DB, Ahring PK. pH Dependency and Desensitization Kinetics of

Heterologously Expressed Combinations of Acid-sensing Ion Channel Subunits. J. Biol. Chem.

2004;279:11006–15.

4. Baron A, Waldmann R, Lazdunski M. ASIC-like , proton-activated currents in rat hippocampal neurons. J. Physiol. 2002;539:485–94.

5. Sherwood TW, Lee KG, Gormley MG, Askwith CC. Heteromeric Acid-Sensing Ion Channels (ASICs) Composed of ASIC2b and ASIC1a Display Novel Channel Properties and Contribute to Acidosis-Induced Neuronal Death. J. Neurosci. 2011;31:9723–34.

6. Wu L-J, Duan B, Mei Y-D, Gao J, Chen J-G, Zhuo M, et al. Characterization of Acid-sensing Ion Channels in Dorsal Horn Neurons of Rat Spinal Cord. J. Biol. Chem. 2004;279:43716–24.

7. Gao J, Wu LJ, Xu L, Xu T Le. Properties of the proton-evoked currents and their modulation by Ca 2+ and Zn2+ in the acutely dissociated hippocampus CA1 neurons. Brain Res. 2004;1017:197–207.

8. Askwith CC, Wemmie J a., Price MP, Rokhlina T, Welsh MJ. Acid-sensing Ion Channel 2 (ASIC2) Modulates ASIC1 H+-activated Currents in Hippocampal Neurons. J. Biol. Chem. 2004;279:18296–305.

9. Wemmie J a., Chen J, Askwith CC, Hruska-Hageman AM, Price MP, Nolan BC, et al. The acid-activated ion channel ASIC contributes to synaptic plasticity, learning, and memory. Neuron. 2002;34:463–77.

10. Liu MG, Li HS, Li WG, Wu YJ, Deng SN, Huang C, et al. Acid-sensing ion channel 1a contributes to hippocampal LTP inducibility through multiple mechanisms. Sci. Rep. 2016;6:1–14.

11. Mango D, Braksator E, Battaglia G, Marcelli S, Mercuri NB, Feligioni M, et al. Acid-sensing ion channel 1a is required for mGlu receptor dependent long-term depression in the hippocampus. Pharmacol. Res. 2017;119:12–9.

12. Vralsted VC, Price MP, Du J, Schnizler M, Wunsch AM, Ziemann AE, et al. Expressing acid-sensing ion channel 3 in the brain alters acid-evoked currents and impairs fear conditioning. Genes, Brain Behav. 2011;10:444–50.

13. Taugher RJ, Lu Y, Fan R, Ghobbeh A, Kreple CJ, Faraci FM, et al. ASIC1A in neurons is

critical for fear-related behaviors. Genes, Brain Behav. 2017;16:745–55.

14. Wemmie JA, Coryell MW, Askwith CC, Lamani E, Leonard AS, Sigmund CD, et al. Overexpression of acid-sensing ion channel 1a in transgenic mice increases acquired fear-related behavior. Proc. Natl. Acad. Sci. 2004;101:3621–6.

15. Xiong Z-G, Zhu X-M, Chu X-P, Minami M, Hey J, Wei W-L, et al. Neuroprotection in Ischemia: Blocking Calcium-Permeable Acid-Sensing Ion Channels. Cell. 2004;118:687–98.

16. Duan B, Wang Y-Z, Yang T, Chu X-P, Yu Y, Huang Y, et al. Extracellular Spermine Exacerbates Ischemic Neuronal Injury through Sensitization of ASIC1a Channels to Extracellular Acidosis. J. Neurosci. 2011;31:2101–12.

17. Gu L, Liu X, Yang Y, Luo D, Zheng X. ASICs aggravate acidosis-induced injuries during ischemic reperfusion. Neurosci. Lett. 2010;479:63–8.

18. Friese MA, Craner MJ, Etzensperger R, Vergo S, Wemmie JA, Welsh MJ, et al. Acid-sensing ion channel-1 contributes to axonal degeneration in autoimmune inflammation of the central nervous system. 2007;13:1483–9.

19. Chu X-P, Xiong Z-G. Physiological and Pathological Functions of Acid-Sensing Ion Channels in the Central Nervous System. Curr. Drug Targets. 2012;13:263–71.

20. Gonzales EB, Sumien N. Acidity and Acid-Sensing Ion Channels in the Normal and Alzheimer's Disease Brain. J. Alzheimer's Dis. 2017;57:1137–44.

21. Chesler M. Regulation and Modulation of pH in the Brain. 2003;1183–221.

22. Rehncrona S. Brain acidosis. Ann. Emerg. Med. 1985;14:770–6.

23. Immke DC, McCleskey EW. Lactate enhances the acid-sensing Na+ channel on ischemia-sensing neurons. Nat. Neurosci. 2001;4:869–70.

24. Allen NJ, Attwell D. Modulation of ASIC channels in rat cerebellar Purkinje neurons by ischaemia-related signals. J. Physiol. 2002;543:521–9.

25. Smith ES, Cadiou H, McNaughton P a. Arachidonic acid potentiates acid-sensing ion channels in rat sensory neurons by a direct action. Neuroscience. 2007;145:686–98.

26. Tombaugh GC, Somjen GG. Effects of extracellular pH on voltage-gated Na+, K+ and Ca2+

currents in isolated rat CA1 neurons. J. Physiol. 1996;493:719–32.

27. Nakamura M, Jang I-S. Acid modulation of tetrodotoxin-resistant Na+ channels in rat

nociceptive neurons. Neuropharmacology. Elsevier Ltd; 2015;90:82–9.

28. Nakamura M, Kim DY, Jang IS. Acid modulation of tetrodotoxin-sensitive Na+ channels in

large-sized trigeminal ganglion neurons. Brain Res. 2016;1651:44–52.

29. Sepulveda F V., Pablo Cid L, Teulon J, Niemeyer MI. Molecular Aspects of Structure,

Gating, and Physiology of pH-Sensitive Background K2P and Kir K+-Transport Channels.

Physiol. Rev. 2015;95:179–217.

30. Mcdonald JW, Bhattacharyya T, Sensi SL, Lobner D, Ying HS, Canzoniero LMT, et al.

Extracellular Acidity Potentiates AMPA Receptor-Mediated Cortical Neuronal Death.

1998;18:6290–9.

31. Traynelis SF, Cull-Candy SG. Proton inhibition of N-methyl-D-aspartate receptors in

cerebellar neurons. Lett. to Nat. 1990;345:347–50.

32. Bennett NC, Faulkes CG. African mole-rats: Ecology and Eusociality. Cambridge University

Press; 2000.

33. Brett RA. The ecology of naked mole-rat colonies: burrowing, food and limiting factors. In:

Sherman P, Jarvis J, Alexander R, editors. Biol. Naked Mole-Rat. Princeton University Press;

1991. p. 137–84.

34. Jarvis JUM. Reproduction of naked mole-rats. In: Sherman P, Jarvis J, Alexander M,

editors. Biol. Naked Mole-Rat. Princeton University Press; 1991. p. 384–425.

35. Schuhmacher L, Husson Z, Smith ES. The naked mole-rat as an animal model in biomedical

research: current perspectives. Open Access Anim. Physiol. 2015;7:137.

36. Ruby JG, Smith M, Buffenstein R. Naked Mole-Rat mortality rates defy gompertzian laws by

not increasing with age. Elife. 2018;7:1–18.

37. Edrey YH, Medina DX, Gaczynska M, Osmulski P a., Oddo S, Caccamo A, et al. Amyloid

beta and the longest-lived rodent: The naked mole-rat as a model for natural protection from

alzheimer's disease. Neurobiol. Aging. 2013;34:2352–60.

38. Orr ME, Garbarino VR, Salinas A, Buffenstein R. Sustained high levels of neuroprotective,

high molecular weight, phosphorylated tau in the longest-lived rodent. Neurobiol. Aging. Elsevier

Inc; 2015;36:1496–504.

39. Seluanov A, Hine C, Azpurua J, Feigenson M, Bozzella M, Mao Z, et al. Hypersensitivity to

contact inhibition provides a clue to cancer resistance of naked mole-rat. Proc. Natl. Acad. Sci.

U. S. A. 2009;106:19352–7.

40. Tian X, Azpurua J, Hine C, Vaidya A, Myakishev-Rempel M, Ablaeva J, et al. High-

molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. Nature.

2013;499:346–9.

41. Liang S, Mele J, Wu Y, Buffenstein R, Hornsby PJ. Resistance to experimental

tumorigenesis in cells of a long-lived mammal, the naked mole-rat (Heterocephalus glaber).

Aging Cell. 2010;9:626–35.

42. Park TJ, Lu Y, Jüttner R, Smith ESJ, Hu J, Brand A, et al. Selective inflammatory pain

insensitivity in the African naked mole-rat (Heterocephalus glaber). PLoS Biol. 2008;6:0156–70.

43. Smith ESJ, Omerbašić D, Lechner SG, Anirudhan G, Lapatsina L, Lewin GR. The molecular

basis of acid insensitivity in the African naked mole-rat. Science. 2011;334:1557–60.

44. Smith ESJ, Blass GRC, Lewin GR, Park TJ. Absence of histamine-induced itch in the

African naked mole-rat and "rescue" by Substance P. Mol. Pain. 2010;6:29.

45. Lavinka PC, Brand A, Landau VJ, Wirtshafter D, Park TJ. Extreme tolerance to ammonia

fumes in African naked mole-rats: Animals that naturally lack neuropeptides from trigeminal

chemosensory nerve fibers. J. Comp. Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol.

2009;195:419–27.

46. Larson J, Park TJ. Extreme hypoxia tolerance of naked mole-rat brain. Neuroreport.

2009;20:1634–7.

47. Park TJ, Reznick J, Peterson BL, Blass G, Omerba D, Bennett NC, et al. Fructose-driven glycolysis supports anoxia resistance in the naked mole-rat. 2017;311:307–11.

48. Peterson BL, Larson J, Buffenstein R, Park TJ, Fall CP. Blunted neuronal calcium response to hypoxia in naked mole-rat hippocampus. PLoS One. 2012;7:1–8.

49. Peterson BL, Park TJ, Larson J. Adult naked mole-rat brain retains the NMDA receptor subunit GluN2D associated with hypoxia tolerance in neonatal mammals. Neurosci. Lett. 2012;506:342–5.

50. Schuhmacher L-N, Callejo G, Srivats S, Smith ESJ. Naked mole-rat acid-sensing ion channel 3 forms nonfunctional homomers, but functional heteromers. J. Biol. Chem. 2018;293:1756–66.

51. Schuhmacher L, Smith ESJ. Expression of acid-sensing ion channels and selection of reference genes in mouse and naked mole rat. Mol. Brain. Molecular Brain; 2016;1–12.

52. Omerbašić D, Smith ESJ, Moroni M, Homfeld J, Eigenbrod O, Bennett NC, et al. Hypofunctional TrkA Accounts for the Absence of Pain Sensitization in the African Naked Mole-Rat. Cell Rep. 2016;17:748–58.

53. Penz OK, Fuzik J, Kurek AB, Romanov R, Larson J, Park TJ, et al. Protracted brain development in a rodent model of extreme longevity. Sci. Rep. 2015;5:11592.

54. Baron A, Lingueglia E. Pharmacology of acid-sensing ion channels – Physiological and therapeutical perspectives. Neuropharmacology. Elsevier Ltd; 2015;94:19–35.

55. Evans MS, Collings M a, Brewer GJ. Electrophysiology of embryonic, adult and aged rat hippocampal neurons in serum-free culture. J. Neurosci. Methods. 1998;79:37–46.

56. Yang J, Thio LL, Clifford DB, Zorumski CF. Electrophysiological properties of identified postnatal rat hippocampal pyramidal neurons in primary culture. Dev. Brain Res. 1993;71:19–26.

57. Wang X, Zhang XG, Zhou TT, Li N, Jang CY, Xiao ZC, et al. Elevated neuronal excitability due to modulation of the voltage-gated sodium channel Nav1.6 by Aβ1-42. Front. Neurosci.

2016;10:1–9.

58. Catterall WA. Voltage-gated sodium channels at 60 : structure , function and

pathophysiology. 2012;11:2577–89.

59. Shams I, Avivi A, Nevo E. Oxygen and carbon dioxide fluctuations in burrows of

subterranean blind mole rats indicate tolerance to hypoxic – hypercapnic stresses.

2005;142:376–82.

60. Mcnab BK. The Metabolism of Fossorial Rodents: A Study of Convergence. Ecol. Soc. Am.

1966;47:712–33.

61. Huang Y, Jiang N, Li J, Ji Y-H, Xiong Z-G, Zha X. Two aspects of ASIC function: Synaptic

plasticity and neuronal injury. Neuropharmacology. Elsevier Ltd; 2015;1–6.

62. Schuhmacher L, St E, Smith J. Expression of acid-sensing ion channels and selection of

reference genes in mouse and naked mole rat. Mol. Brain. Molecular Brain; 2016;1–12.

63. Ding D, Moskowitz SI, Li R, Lee SB, Esteban M, Tomaselli K, et al. Acidosis induces

necrosis and apoptosis of cultured hippocampal neurons. Exp. Neurol. 2000;162:1–12.

64. Zeng W-Z, Liu D-S, Buan B, Song X-L, Wang X, Wei D, et al. Molecular Mechanism of

Constitutive Endocytosis Of Acid-Sensing Ion Channel 1a and Its Protective Function in

Acidosis-Induced Neuronal Death. J Neurosci. 2013;33:7066–78.

65. Xiao B, Wang S, Yang G, Sun X, Zhao S, Lin L, et al. HIF-1α contributes to hypoxia

adaptation of the naked mole rat. Oncotarget. 2017;8:109941–51.

Figure Legends

Fig. 1. Electrophysiological properties of NMR neurons differ from mouse neurons. NMR cortical

and hippocampal neurons have significantly smaller capacitance (**a**) and more hyperpolarized

resting membrane potential (**b**) than mouse neurons. **c.** Example of whole-cell patch-clamp

recordings made from a mouse (left, black) and a NMR (middle, blue) hippocampal neuron in

response to the voltage step protocol and the inhibition of response in NMR neurons by 300 nM

TTX (right, blue). Cortical and hippocampal NMR neurons have significantly smaller voltage-gated

inward currents (**a, b**), a more depolarized Vhalf (**c**) and a more depolarized peak activation

membrane potential Vm (**d**) compared to mouse neurons. **e.** Voltage-gated outward currents in

NMR cortical and hippocampal show no significant difference to those in mouse neurons. * p <

0.5; ** p < 0.01; **** p< 0.0001, unpaired t-tests within each structure. Numbers in brackets

indicate the number of recorded cells.

Fig. 2. ASICs mediate acid-induced currents in NMR and mouse CNS neurons. **a.** Both mouse

(black trace, left) and NMR (blue trace, right) neurons respond to a pH 5 solution with a transient

inward current. **b.** Acid-induced currents are of significantly smaller amplitude in NMR neurons

compared to mouse neurons. **c.** Inactivation time constants of the acid-induced responses are

similar between NMR and mouse neurons. **d.** Example trace of an acid-induced current elicited

by a pH 5 solution and the effect of 100 µM benzamil in a mouse cortical neuron. **e-f.** Acid-induced

currents in both cortical and hippocampal mouse neurons were reversibly blocked by 100 µM

benzamil (n = 9 and n = 10 cortical and hippocampal neurons, respectively). **g.** Example trace of

an acid-induced current evoked by a pH 5 solution in a cortical NMR neuron showing inhibition

by 100 µM benzamil. **h-i.** In both cortical (n = 7) and hippocampal (n = 6) NMR neurons, acid-

induced currents were reversibly blocked by 100 µM benzamil. ** p < 0.01; **** p< 0.0001, one-

way ANOVA paired tests, Tukey's multiple comparison tests. Numbers into brackets indicates the

number of recorded cells.

Fig. 3. NMR cortical neurons show resistance to acid-induced cell death. Mouse (**a**) and NMR (**b**)

cortical neurons were incubated for 2 hours with either a pH 7.4 or pH 5 solution and then stained

with Hoechst 33342 to label nuclei and PI to label necrotic dead cells. **c.** Percentages of PI-

positive neurons (i.e. dead) at pH 7.4 were similar between mouse and NMR neurons. The

percentage of mouse dead neurons was higher when cells were incubated in a pH 5 solution,

compared to pH 7.4 condition and compared to the percentage of cell death obtained with NMR

neurons at pH 5. NMR cortical neurons did not exhibit any significant difference with regard to cell

death when exposed to pH 5.0 compared to pH 7.4. **** $p < 0.0001$, one-way ANOVA test, Tukey's

multiple comparison test. Percentages of each field of view were used to compare acidotoxicity

between species. The number of fields of view analyzed for each condition are the following: n =

18 for mouse neurons (for both pH 7.4 and pH 5), n = 27 and 28 for NMR neurons at pH 7.4 and

at pH 5, respectively; obtained from 3 independent cultures for both species.

FIG. 1

FIG. 3