Title: Genome-wide study identifies 611 loci associated with risk tolerance and risky behaviors

All authors and their affiliations are listed at the end of the manuscript.

Abstract:

Humans vary substantially in their willingness to take risks. In a combined sample of over one million individuals, we conducted genome-wide association studies (GWAS) of general risk tolerance, adventurousness, and risky behaviors in the driving, drinking, smoking, and sexual domains. We identified 611 approximately independent genetic loci associated with at least one of our phenotypes, including 124 with general risk tolerance. We report evidence of substantial shared genetic influences across general risk tolerance and risky behaviors: 72 of the 124 general risk tolerance loci contain a lead SNP for at least one of our other GWAS, and general risk tolerance is moderately to strongly genetically correlated ($|\hat{r}_g| \sim 0.25$ to 0.50) with a range of risky behaviors. Bioinformatics analyses imply that genes near general-risk-tolerance-associated SNPs are highly expressed in brain tissues and point to a role for glutamatergic and GABAergic neurotransmission. We find no evidence of enrichment for genes previously hypothesized to relate to risk tolerance.

Main Text:

Choices in important domains of life, including health, fertility, finance, employment, and social relationships, rarely have consequences that can be anticipated perfectly. The degree of variability in possible outcomes is called risk. Risk tolerance—defined as the willingness to take risks, typically to obtain some reward—varies substantially across humans and has been actively studied in the behavioral and social sciences. An individual's risk tolerance may vary across domains, but survey-based measures of *general* risk tolerance (e.g., "Would you describe yourself as someone who takes risks?") have been found to be good all-around predictors of risky behaviors such as portfolio allocation, occupational choice, smoking, drinking alcohol, and starting one's own business[1–3].

Twin studies have established that various measures of risk tolerance are moderately heritable ($h^2 \sim 30\%$, although estimates in the literature vary[3–5]). Discovery of specific genetic variants associated with general risk tolerance could advance our understanding of how genetic influences are amplified and dampened by environmental factors; provide insights into underlying biological pathways; enable the construction of polygenic scores (indexes of many genetic variants) that can be used as overall measures of genetic influences on individuals; and help distinguish genetic variation associated with general versus domain-specific risk tolerance.

Although risk tolerance has been one of the most studied phenotypes in social science genetics, most claims of positive findings have been based on small-sample candidate gene studies (**Supplementary Table 11.1**), whose limitations are now appreciated[6]. To date, only two loci associated with risk tolerance have been identified in genome-wide association studies (GWAS)[7,8].

Here, we report results from large-scale GWAS of self-reported general risk tolerance (our primary phenotype) and six supplementary phenotypes: "adventurousness" (defined as the self-reported tendency to be adventurous vs. cautious); four risky behaviors: "automobile speeding propensity" (the tendency to drive faster than the speed limit), "drinks per week" (the average number of alcoholic drinks consumed per week), "ever smoker" (whether one has ever been a smoker), and "number of sexual partners" (the lifetime number of sexual partners); and the first principal component (PC) of these four risky behaviors, which we interpret as capturing the general tendency to take risks across domains. All seven phenotypes are coded such that higher

phenotype values are associated with higher risk tolerance or risk taking. **Table 1** lists, for each

GWAS, the datasets we analyzed and the GWAS sample size.

Association analyses

All seven GWAS were performed in European-ancestry subjects, following procedures described

in a pre-specified analysis plan (https://osf.io/cjx9m/) and in **Supplementary Information**

section 2.

In the discovery phase of our GWAS of general risk tolerance (n = 939,908), we performed a

sample-size-weighted meta-analysis of results from the UK Biobank (UKB, n = 431,126) and a

sample of research participants from 23andMe (n = 508,782). The UKB measure of general risk

tolerance is based on the question: "Would you describe yourself as someone who takes risks?

Yes / No." The 23andMe measure is based on a question about overall comfort taking risks, with

five response options ranging from "very comfortable" to "very uncomfortable." The genetic

correlation[9] between the UKB and 23andMe cohorts ($\hat{r}_g$ = 0.77, SE = 0.02) is smaller than one

but high enough to justify our approach of pooling the two cohorts[10].

The Q-Q plot (**Extended Data Fig. 3.2a**) from the discovery GWAS exhibits substantial

inflation (λ_{GC} = 1.41). According to the estimated intercept from a linkage disequilibrium (LD)

Score regression[11], only a small share of this inflation (~5%) in test statistics is due to bias. To

account for this bias, we inflated GWAS standard errors by the square root of the LD Score

regression intercept.

We identified 124 approximately independent SNPs (pairwise r^2 < 0.1) that attained genome-

wide significance (P < 5×10^{-8}). These 124 "lead SNPs" are listed in **Supplementary Table 3.1**

and shown in **Fig. 1a**. All have coefficients of determination (R^2's) below 0.02%, and the SNP

with the largest per-allele effect is estimated to increase general risk tolerance by ~0.026

standard deviations in our discovery sample (**Extended Data Fig. 3.3**).

In the replication phase of our GWAS of general risk tolerance (combined n = 35,445), we meta-

analyzed summary statistics from ten smaller cohorts. Additional details on cohort-level

phenotype measures are provided in **Supplementary Table 1.2**. The questions differ in terms of

their exact wording and number of response categories, but all questions ask subjects about their

overall or general attitudes toward risk. The genetic correlation[9] between the discovery and

replication GWAS is 0.83 (SE = 0.13). 123 of the 124 lead SNPs were available or well proxied

by an available SNP in the replication GWAS results. Out of the 123 SNPs, 94 have a

concordant sign (P = 1.7×10^{-9}) and 23 are significant at the 5% level in one-tailed t tests (P =

4.5×10^{-8}) (**Extended Data Fig. 5.1**). This empirical replication record matches theoretical

projections that take into account sampling variation and the winner's curse (**Supplementary**

Information section 5).

Our six supplementary GWAS—of adventurousness, four risky behaviors, and their principal

component (n = 315,894 to 557,923; **Supplementary Tables 1.1-1.2**)—were conducted using

methods comparable to those in the primary GWAS, but without a replication phase. **Extended**

Data Fig. 3.2 (**c** to **h**) shows Q-Q plots and **Extended Data Fig. 3.1** (**a** to **f**) shows Manhattan

plots.

Table 1 provides a summary overview of the seven GWAS. We identified a total of 865 lead

SNPs across the seven GWAS. The lead SNPs are located in 611 approximately independent

loci, where a locus is defined as the set of all SNPs in weak LD (pairwise $r^2 > 0.1$) with a lead

SNP. The SNP heritabilities of the seven phenotypes range from ~0.05 (for general risk

tolerance) to ~0.16 (for the first PC of the four risky behaviors).

<u>Genetic overlap</u>

There is substantial overlap across the results of our GWAS. For example, 72 of the 124 general-

risk-tolerance lead SNPs are in loci that also contain lead SNPs for at least one of the other

GWAS, including 45 for adventurousness and 49 for at least one of the four risky behaviors or

their first PC. To empirically assess if this overlap could be attributed to chance, we conducted a

resampling exercise under the null hypothesis that the lead SNPs of our supplementary GWAS

are distributed independently of the 124 general-risk-tolerance lead loci. We strongly rejected

this null hypothesis ($P < 0.0001$; **Supplementary Information section 3.3.3**).

Several regions of the genome stand out for being associated both with general risk tolerance and

with all or most of the supplementary phenotypes. We tested whether the signs of the lead SNPs

located in these regions tend to be concordant across our primary and supplementary GWAS. We

strongly rejected the null hypothesis of no concordance ($P < 3\times10^{-30}$; **Supplementary**

Information section 3.2.3), suggesting that these regions represent shared genetic influences,

rather than colocalization of causal SNPs. **Fig. 1b** and **Extended Data Fig. 3.4** show local

Manhattan plots for some of these regions. The long-range LD region[12] on chromosome 3 (~83.4

to 86.9 Mb) contains lead SNPs from all seven GWAS as well as the most significant lead SNP

from the general risk tolerance GWAS, rs993137 ($P = 2.14\times10^{-40}$), which is located in the gene

CADM2. Another long-range LD region, on chromosome 6 (~25.3 to 33.4 Mb), covers the HLA-

complex and contains lead SNPs from all GWAS except drinks per week. Three candidate

inversions (i.e., genomic regions that are highly prone to inversion polymorphisms;

Supplementary Information section 2.9.2) on chromosomes 7 (~124.6 to 132.7 Mb), 8 (~7.89

to 11.8 Mb), and 18 (~49.1 to 55.5 Mb) contain lead SNPs from six, five, and all seven of our

GWAS, respectively. Finally, four other LD blocks[13] that do not overlap known long-range LD

or candidate inversion regions each contain lead SNPs from five of our GWAS (including

general risk tolerance). The two long-range LD regions and the three candidate inversions have

previously been found to be associated with numerous phenotypes, including many cognitive and

neuropsychiatric phenotypes[14].

To investigate genetic overlap at the genome-wide level, we estimated genetic correlations with

self-reported general risk tolerance using bivariate LD Score regression[9]. (For this and all

subsequent analyses involving general risk tolerance, we used the summary statistics from the

combined meta-analysis of our discovery and replication GWAS.) The estimated genetic

correlations with our six supplementary phenotypes are all positive, larger than ~0.25, and highly

significant ($P < 2.3\times10^{-30}$; **Fig. 2**), indicating that SNPs associated with higher general risk

tolerance also tend to be associated with riskier behavior. The largest estimated genetic

correlations are with adventurousness ($\hat{r}_g = 0.83$, $SE = 0.01$), number of sexual partners (0.52, *SE*

= 0.02), automobile speeding propensity (0.45, $SE = 0.02$), and the first PC of the four risky

behaviors (0.50, $SE = 0.02$).

Our estimates of the genetic correlations between general risk tolerance and the supplementary

risky behaviors are substantially higher than the corresponding phenotypic correlations

(**Supplementary Tables 1.3** and **7.1**). Although measurement error partly accounts for the low

phenotypic correlations, the genetic correlations remain considerably higher even after

adjustment of the phenotypic correlations for measurement error. The comparatively large
genetic correlations support the view that a general factor of risk tolerance partly accounts for
cross-domain variation in risky behavior[15,16] and imply that this factor is genetically influenced.
The lower phenotypic correlations suggest that environmental factors are more important
contributors to domain-specific risky behavior[17,18].

To increase the precision of our estimates of the SNPs' effects on general risk tolerance, we
leveraged the high degree of genetic overlap across our phenotypes by conducting Multi-Trait
Analysis of GWAS (MTAG)[19]. We used as inputs the summary statistics of our GWAS of
general risk tolerance, of our first five supplementary GWAS (i.e., not including the first PC of
the four risky behaviors), and of a previously published GWAS on lifetime cannabis use[20]
(**Supplementary Information section 9**). MTAG increased the number of general-risk-tolerance
lead SNPs from 124 to 312 (**Extended Data Fig. 9.1**, **Supplementary Table 9.1**).

We also estimated genetic correlations between general risk tolerance and 28 additional
phenotypes (**Fig. 2** and in **Supplementary Table 7.1**). These included phenotypes for which we
could obtain summary statistics from previous GWAS, as well as five phenotypes for which we
conducted new GWAS. The estimated genetic correlations for the personality traits extraversion
($\hat{r}_g = 0.51$, *SE* = 0.03), neuroticism (-0.42, *SE* = 0.04), and openness to experience (0.33, *SE* =
0.03) are substantially larger in magnitude than previously reported phenotypic correlations[21],
pointing to substantial shared genetic influences among general risk tolerance and these traits.
After Bonferroni correction, we also find significant positive genetic correlations with the
neuropsychiatric phenotypes ADHD, bipolar disorder, and schizophrenia. Viewed in light of the
genetic correlations we find with risky behaviors classified as externalizing (e.g., substance use,
elevated sexual behavior, and fast driving), these results suggest the hypothesis that the overlap
with the neuropsychiatric phenotypes is driven by their externalizing component.

<u>Biological annotation</u>

To gain insights into the biological mechanisms through which genetic variation influences
general risk tolerance, we conducted a number of analyses. First, we systematically reviewed the
literature that aimed to link risk tolerance to biological pathways (**Supplementary Information**
section 11). Our review covered studies based on candidate genes (i.e., specific genetic variants

used as proxies for biological pathways), pharmacological manipulations, biochemical assays,

genetic manipulations in rodents, as well as other research designs. Our review identified 132

articles that matched our search criteria (**Supplementary Table 11.1**).

Previous work has focused on five main biological pathways: the steroid hormone cortisol, the

monoamines dopamine and serotonin, and the steroid sex hormones estrogen and testosterone.

Using a MAGMA[22] competitive gene-set analysis, we found no evidence that SNPs within genes

associated with these five pathways tend to be more associated with general risk tolerance than

SNPs in other genes (**Supplementary Table 11.3**). Further, none of the other bioinformatics

analyses we report below point to these pathways.

We also examined the 15 most commonly tested autosomal genes within the dopamine and

serotonin pathways, which were the focus of most of the 34 candidate-gene studies identified by

our literature review. We verified that the SNPs available in our GWAS results tag most of the

genetic variants typically used to test the 15 genes. Across one SNP-based test and two gene-

based tests, we found no evidence of non-negligible associations between those genes and

general risk tolerance (**Fig. 1c** and **Supplementary Table 11.4**). (We note, however, that some

brain regions identified in analyses we report below are areas where dopamine and serotonin

play important roles.)

Second, we performed a MAGMA[22] gene analysis to test each of ~18,000 protein-coding genes

for association with general risk tolerance (**Supplementary Information section 12.2**). After

Bonferroni correction, 285 genes were significant (**Extended Data Fig. 12.1** and

Supplementary Table 12.3). To gain insight into the functions and expression patterns of these

285 genes, we looked up these genes in the Gene Network[23] co-expression database. Third, to

identify relevant biological pathways and identify tissues in which genes near general-risk-

tolerance-associated SNPs are expressed, we applied the software tool DEPICT[24] to the SNPs

with P values less than 10^{-5} in our GWAS of general risk tolerance (**Supplementary**

Information section 12.4).

Both the Gene Network and the DEPICT analyses separately point to a role for glutamate and

GABA neurotransmitters, which are the main excitatory and inhibitory neurotransmitters in the

brain, respectively[25] (**Fig. 3a** and **Supplementary Tables 12.4** and **12.8**). To our knowledge, no

published large-scale GWAS of cognition, personality, or neuropsychiatric phenotypes has

pointed to clear roles both for glutamate and GABA (although glutamatergic neurotransmission
has been implicated in recent GWAS of schizophrenia[26] and major depression[27]). Our results
suggest that the balance between excitatory and inhibitory neurotransmission may contribute to
variation in general risk tolerance across individuals.

The Gene Network and the DEPICT tissue enrichment analyses also both separately point to
enrichment of the prefrontal cortex and the basal ganglia (**Fig. 3b** and **Supplementary Tables**
12.4, **12.6**, and **12.7**). The cortical and subcortical regions highlighted by DEPICT include some
of the major components of the cortical-basal ganglia circuit, which is known as the reward
system in human and non-human primates and is critically involved in learning, motivation, and
decision-making, notably under risk and uncertainty[28,29]. We caution, however, that our results
do not point exclusively to the reward system.

Lastly, we used stratified LD Score regression[30] to test for the enrichment of SNPs associated
with histone marks in 10 tissue or cell types (**Supplementary Information section 12.1**).
Central nervous system tissues are the most enriched, accounting for 44% ($SE = 3\%$) of the
heritability while comprising only 15% of the SNPs (**Extended Data Fig. 12.3a** and
Supplementary Table 12.2). Immune/hematopoietic tissues are also significantly enriched.
While a role for the immune system in modulating risk tolerance is plausible given prior
evidence of its involvement in several neuropsychiatric disorders[26,27], future work is needed to
confirm this result and to uncover specific pathways that might be involved.

<u>Polygenic prediction</u>

We constructed polygenic scores of general risk tolerance to gauge their potential usefulness in
empirical research (**Supplementary Information section 10**). We used the Add Health, HRS,
NTR, STR, UKB-siblings, and Zurich cohorts as validation cohorts (**Supplementary Table 1.1**
provides an overview of these cohorts; the UKB-siblings cohort comprised individuals with at
least one full sibling in the UKB). For each validation cohort, we constructed the score using
summary statistics from a meta-analysis of our discovery and replication GWAS that excluded
the cohort. Our measure of predictive power is the incremental R^2 (or pseudo-R^2) from adding
the score to a regression of the phenotype on sex, birth year, and the top ten principal
components of the genetic relatedness matrix.

Our preferred score was constructed with LDpred[31]. In the UKB-siblings cohort, which is our

largest validation cohort ($n \sim 35{,}000$), the score's predictive power is 1.6% for general risk

tolerance, 1.0% for the first PC of the four risky behaviors, 0.8% for number of sexual partners,

0.6% for automobile speeding propensity, and ~0.15% for drinks per week and ever smoker.

Across our validation cohorts, the score is also predictive of several personality phenotypes and a

suite of real-world measures of risky behaviors in the health, financial, career, and other domains

(**Extended Data Figs. 10.1-10.2** and **Supplementary Tables 10.1-10.3**). The incremental R^2 we

observe for general risk tolerance is consistent with the theoretical prediction, given the SNP

heritability of general risk tolerance (**Table 1**) and the imperfect genetic correlations across the

GWAS and validation cohorts[32,33] (**Supplementary Information section 10.4**).

<u>Discussion</u>

Our results provide insights into biological mechanisms that influence general risk tolerance. Our

bioinformatics analyses point to the role of gene expression in brain regions that have been

identified by neuroscientific studies on decision-making, notably the prefrontal cortex, basal

ganglia, and midbrain, thereby providing convergent evidence with that from neuroscience[28,29].

Yet our analyses failed to find evidence for the main biological pathways that had been

previously hypothesized to influence risk tolerance. Instead, our analyses implicate genes

involved in glutamatergic and GABAergic neurotransmission, which were heretofore not

generally believed to play a role in risk tolerance.

Although our focus has been on the genetics of general risk tolerance and risky behaviors,

environmental and demographic factors account for a substantial share of these phenotypes'

variation. We observe sizeable effects of sex and age on general risk tolerance in the UKB data

(**Extended Data Fig. 1.1**), and life experiences have been shown to affect both measured risk

tolerance and risky behaviors (e.g., refs. 34,35). The data we have generated will allow

researchers to construct and use polygenic scores of general risk tolerance to measure how

environmental, demographic, and genetic factors interact with one another.

For the behavioral sciences, our results bear on the ongoing debate about the extent to which risk

tolerance is a "domain-general" as opposed to a "domain-specific" trait. Low phenotypic

correlations in risk tolerance across decision-making domains have been interpreted as

supporting the domain-specific view[17,18]. Across the risky behaviors we study, we find that the

genetic correlations are considerably higher than the phenotypic correlations (even after the latter

are corrected for measurement error) and that many lead SNPs are shared across our phenotypes.

These observations suggest that the low phenotypic correlations across domains are due to

environmental factors that dilute the effects of a genetically-influenced domain-general factor of

risk tolerance.

References:

1.   Dohmen, T. *et al.* Individual risk attitudes: Measurement, determinants, and behavioral
consequences. *J. Eur. Econ. Assoc.* **9,** 522–550 (2011).

2.   Falk, A., Dohmen, T., Falk, A. & Huffman, D. The nature and predictive power of
preferences: Global evidence. *IZA Discussion Papers* (2015).

3.   Beauchamp, J. P., Cesarini, D. & Johannesson, M. The psychometric and empirical
properties of measures of risk preferences. *J. Risk Uncertain.* **54,** 203–237 (2017).

4.   Cesarini, D., Dawes, C. T., Johannesson, M., Lichtenstein, P. & Wallace, B. Genetic
variation in preferences for giving and risk taking. *Q. J. Econ.* **124,** 809–842 (2009).

5.   Harden, K. P. *et al.* Beyond dual systems: A genetically-informed, latent factor model of
behavioral and self-report measures related to adolescent risk-taking. *Dev. Cogn.*
Neurosci. **25,** 221–234 (2017).

6.   Hewitt, J. K. Editorial policy on candidate gene association and candidate gene-by-
environment interaction studies of complex traits. *Behav. Genet.* **42,** 1–2 (2012).

7.   Day, F. R. *et al.* Physical and neurobehavioral determinants of reproductive onset and
success. *Nat. Genet.* **48,** 617–623 (2016).

8.   Strawbridge, R. J. *et al.* Genome-wide analysis of risk-taking behaviour and cross-
disorder genetic correlations in 116 255 individuals from the UK Biobank cohort. *bioRxiv*
(2017). doi:http://dx.doi.org/10.1101/177014

9.   Bulik-Sullivan, B. K. *et al.* An atlas of genetic correlations across human diseases and
traits. *Nat. Genet.* **47,** 1236–1241 (2015).

10.   Okbay, A. *et al.* Genetic variants associated with subjective well-being, depressive
symptoms, and neuroticism identified through genome-wide analyses. *Nat. Genet.* **48,**
624–633 (2016).

11.   Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from
polygenicity in genome-wide association studies. *Nat. Genet.* **47,** 291–295 (2015).

12.   Price, A. L. *et al.* Long-range LD can confound genome scans in admixed populations.
Am. J. Hum. Genet. **83,** 132–139 (2008).

13.   Berisa, T. & Pickrell, J. K. Approximately independent linkage disequilibrium blocks in
human populations. *Bioinformatics* **32,** 283–285 (2016).

14.   Welter, D. *et al.* The NHGRI GWAS Catalog, a curated resource of SNP-trait

associations. *Nucleic Acids Res.* **42,** D1001-1006 (2014).

15. Einav, B. L., Finkelstein, A., Pascu, I. & Cullen, M. R. How general are risk preferences?

Choices under uncertainty in different domains. *Am. Econ. Rev.* **102,** 2606–2638 (2016).

16. Frey, R., Pedroni, A., Mata, R., Rieskamp, J. & Hertwig, R. Risk preference shares the

psychometric structure of major psychological traits. *Sci. Adv.* **3,** e1701381 (2017).

17. Weber, E. U., Blais, A. E. & Betz, N. E. A domain-specific risk-attitude scale: Measuring

risk perceptions and risk behaviors. *J. Behav. Decis. Mak. J. Behav. Dec. Mak.* **15,** 263–

290 (2002).

18. Hanoch, Y., Johnson, J. G. & Wilke, A. Domain specificity in experimental measures and

participant recruitment: an application to risk-taking behavior. *Psychol. Sci.* **17,** 300–304

(2006).

19. Turley, P. *et al.* MTAG: Multi-Trait Analysis of GWAS. *bioRxiv* (2017).

doi:https://doi.org/10.1101/118810

20. Stringer, S. *et al.* Genome-wide association study of lifetime cannabis use based on a large

meta-analytic sample of 32 330 subjects from the International Cannabis Consortium.

Transl. Psychiatry **6,** e769 (2016).

21. Becker, A., Deckers, T., Dohmen, T., Falk, A. & Kosse, F. The relationship between

economic preferences and psychological personality measures. *Annu. Rev. Econom.* **4,**

453–478 (2012).

22. de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: Generalized gene-

set analysis of GWAS data. *PLoS Comput. Biol.* **11,** 1–19 (2015).

23. Fehrmann, R. S. N. *et al.* Gene expression analysis identifies global gene dosage

sensitivity in cancer. *Nat. Genet.* **47,** 115–125 (2015).

24. Pers, T. H. *et al.* Biological interpretation of genome-wide association studies using

predicted gene functions. *Nat. Commun.* **6,** 5890 (2015).

25. Petroff, O. A. C. GABA and glutamate in the human brain. *Neurosci.* **8,** 562–573 (2002).

26. Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci. *Nature*

511, 421–427 (2014).

27. Wray, N. R. *et al.* Genome-wide association analyses identify 44 risk variants and refine

the genetic architecture of major depression. *bioRxiv* (2017).

doi:https://doi.org/10.1101/167577

28.   Haber, S. N. & Knutson, B. The reward circuit: linking primate anatomy and human

imaging. *Neuropsychopharmacology* **35,** 4–26 (2010).

29.   Tobler, P. N. & Weber, E. U. Valuation for Risky and Uncertain Choices. in

Neuroeconomics 149–172 (Elsevier, 2014). doi:10.1016/B978-0-12-416008-8.00009-7

30.   Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-

wide association summary statistics. *Nat. Genet.* **47,** 1228–1235 (2015).

31.   Vilhjálmsson, B. J. *et al.* Modeling linkage disequilibrium increases accuracy of polygenic

risk scores. *Am. J. Hum. Genet.* **97,** 576–592 (2015).

32.   Daetwyler, H. D., Villanueva, B. & Woolliams, J. A. Accuracy of predicting the genetic

risk of disease using a genome-wide approach. *PLoS One* **3,** e3395 (2008).

33.   de Vlaming, R. *et al.* Meta-GWAS Accuracy and Power (MetaGAP) calculator shows that

hiding heritability is partially due to imperfect genetic correlations across studies. *PLoS*

Genet. **13,** e1006495 (2017).

34.   Sahm, C. R. How much does risk tolerance change? *Q. J. Financ.* **2,** 1250020 (2012).

35.   Malmendier, U. & Nagel, S. Depression babies: Do macroeconomic experiences affect

risk taking? *Q. J. Econ.* **126,** 373–416 (2011).

36.   Frey, B. J. & Dueck, D. Clustering by passing messages between data points. *Science*

315, (2007).

37.   Furberg, H. *et al.* Genome-wide meta-analyses identify multiple loci associated with

smoking behavior. *Nat. Genet.* **42,** 441–447 (2010).

38.   Shi, H., Kichaev, G. & Pasaniuc, B. Contrasting the genetic architecture of 30 complex

traits from summary association data. *Am. J. Hum. Genet.* **99,** 139–153 (2016).

Acknowledgments This research was carried out under the auspices of the Social Science Genetic Association Consortium (SSGAC). The research has also been conducted using the UK Biobank Resource under Application Number 11425. The study was supported by funding from the Ragnar Söderberg Foundation (E9/11 and E42/15), the Swedish Research Council (421-2013-1061), The Jan Wallander and Tom Hedelius Foundation, an ERC Consolidator Grant to Philipp Koellinger (647648 EdGe), the Pershing Square Fund of the Foundations of Human Behavior, and the NIA/NIH through grants P01-AG005842, P01-AG005842-20S2, P30-AG012810, and T32-AG000186-23 to NBER, and R01-AG042568-02 to the University of Southern California. We thank the International Cannabis Consortium, the Psychiatric Genomics Consortium, and the eQTLgen Consortium for sharing summary statistics from the GWAS of lifetime cannabis use, summary statistics from the GWAS of ADHD, and eQTL summary statistics, respectively. A full list of acknowledgments is provided in **Supplementary Information section 13**.

Author Contributions A full list of author contributions is included in **Supplementary Information section 13**.

Author Information Adam Auton, Pierre Fontanillas, David A Hinds, and Aaron Kleinman are employees of 23andMe, and Ronald Kessler has had ties to various companies in the past three years; the authors declare no other competing financial interests. Further details are provided in **Supplementary Information section 13**. Correspondence and requests for materials should be addressed to J.P.B (jonathan.pierre.beauchamp@gmail.com), R.K.L. (r.karlssonlinner@vu.nl), D.J.B (daniel.benjamin@gmail.com) or P.D.K. (p.d.koellinger@vu.nl).

Data Availability Upon publication, results can be downloaded from the SSGAC website (http://thessgac.org/data).

Figure 1 | Manhattan plots. In all panels, the *x*-axis is chromosomal position; the *y*-axis is the significance on a −log₁₀ scale; the horizontal dashed line marks the threshold for genome-wide significance ($P = 5 \times 10^{-8}$); and each approximately independent (pairwise $r^2 < 0.1$) genome-wide significant association ("lead SNP") is marked by a red ×. **a**, Manhattan plots for the discovery GWAS of general risk tolerance. **b**, Local Manhattan plots of two genomic regions that contain lead SNPs for all seven of our GWAS. The gray background marks the locations of long-range LD or candidate inversion regions. **c**, Local Manhattan plots of the loci around the 15 most commonly tested candidate genes in the prior literature on the genetics of risk tolerance. Each locus comprises all SNPs within 500 kb of the gene's borders that are in LD ($r^2 > 0.1$) with a SNP in the gene. The 15 plots are concatenated and shown together in the panel, divided by the black vertical lines. The 15 genes are not particularly strongly associated with general risk tolerance or the risky behaviors, as can be seen by comparing the results within each row across panels **b** and **c** (the three rows correspond to the GWAS of general risk tolerance, adventurousness, and the first PC of the four risky behaviors).

Figure 2 | Genetic correlations with general risk tolerance. The genetic correlations were estimated using bivariate LD Score (LDSC) regression[9]. Error bars show 95% confidence intervals. For the supplementary phenotypes and the additional risky behaviors, green bars represent significant estimates with the expected signs, where higher risk tolerance is associated with riskier behavior. For the other phenotypes, blue bars represent significant estimates. Light green and light blue bars represent genetic correlations that are statistically significant at the 5% level, and dark green and dark blue bars represent correlations that are statistically significant after Bonferroni correction for 34 tests (the total number of phenotypes tested). Grey bars represent correlations that are not statistically significant at the 5% level.

Figure 3 | Results from selected biological analyses. a, DEPICT gene-set enrichment diagram.
We identified 93 reconstituted gene sets that are significantly enriched (FDR < 0.01) for genes
overlapping loci associated with general risk tolerance; using the Affinity Propagation method[36],
these were grouped into the 13 clusters displayed in the graph. Each cluster was named after the
most significant gene set it contained, and each cluster's color represents the permutation P value
of its most significant gene set. The "synapse part" cluster includes the gene set "glutamate
receptor activity," and several members of the "GABA$_A$ receptor activation" cluster are defined
by gamma-aminobutyric acid signaling. Overlap between the named representatives of two

clusters is represented by an edge. Edge width represents the Pearson correlation ρ between the

two respective vectors of gene membership scores ($\rho < 0.3$, no edge; $0.3 \leq \rho < 0.5$, thin edge; 0.5

$\leq \rho < 0.7$, intermediate edge; $\rho \geq 0.7$, thick edge). **b**, Results of DEPICT tissue enrichment

analysis using GTEx data. The panel shows whether the genes overlapping loci associated with

general risk tolerance are significantly overexpressed (relative to genes in random sets of loci

matched by gene density) in various tissues. Tissues are grouped by organ or tissue type. The

orange bars correspond to tissues with significant overexpression (FDR < 0.01). The *y*-axis is the

significance on a $-\log_{10}$ scale.

Table 1 | GWAS results

GWAS	Cohorts analyzed	n	Mean χ^2	LD Score intercept (*SE*)	# lead SNPs	SNP h^2 (*SE*)
General risk tolerance (disc. GWAS)	UKB; 23andMe	939,908	1.85	1.04 (0.01)	124	0.046 (0.001)
General risk tolerance (rep. GWAS)	10 indep. cohorts	35,445	1.03	1.00 (0.07)	0	--
General risk tolerance (disc. + rep.)	UKB; 23andMe; 10 indep. cohorts	975,353	1.87	1.04 (0.01)	132	0.045 (0.001)
Adventurousness	23andMe	557,923	1.98	1.05 (0.01)	167	0.098 (0.002)
Automobile speeding propensity	UKB	404,291	1.53	1.03 (0.01)	42	0.079 (0.003)
Drinks per week	UKB	414,343	1.61	1.03 (0.01)	85	0.085 (0.003)
Ever smoker	UKB; TAG Consortium [37]	518,633	1.97	1.05 (0.01)	223	0.109 (0.003)
Number of sexual partners	UKB	370,711	1.77	1.04 (0.01)	118	0.128 (0.003)
First PC of the four risky behaviors	UKB	315,894	1.77	1.05 (0.01)	106	0.156 (0.004)

The table provides an overview of the GWAS of our primary and supplementary phenotypes. "n": GWAS sample size; "Mean χ^2": mean GWAS chi-squared statistics across HapMap3 SNPs with minor allele frequency (MAF) greater than 0.01; "LD Score intercept": estimate of the intercept from a LD Score regression[11] using HapMap3 SNPs with MAF greater than 0.01; "# lead SNPs": number of lead SNPs, calculated after the associated statistics have been adjusted using the estimated LD score intercept; "SNP h^2": SNP heritability estimated with the Heritability Estimator from Summary Statistics (HESS) method[38] using 1000 Genomes phase 3 SNPs with MAF greater than 0.05; "disc.": discovery; "rep.": replication; "indep.": independent.

Authors:

Richard Karlsson Linnér[1,2,3]*, Pietro Biroli[4]*, Edward Kong[5]*, S Fleur W Meddens[1,2,3]*, Robbee Wedow[6,7,8]*, Mark Alan Fontana[9,10], Maël Lebreton[11,12], Abdel Abdellaoui[13], Anke R Hammerschlag[1], Michel G Nivard[13], Aysu Okbay[1,3], Cornelius A Rietveld[14,2,15], Pascal N Timshel[16,17], Stephen P Tino[18], Maciej Trzaskowski[19], Ronald de Vlaming[1,2,3], Christian L Zünd[4], Yanchun Bao[20], Laura Buzdugan[21,22], Ann H Caplin[23], Chia-Yen Chen[24,25,26], Peter Eibich[27,28,29], Pierre Fontanillas[30], Juan R Gonzalez[31,32,33], Peter K Joshi[34], Ville Karhunen[35], Aaron Kleinman[30], Remy Z Levin[36], Christina M Lill[37,38], Gerardus A Meddens[39], Gerard Muntané[40], Sandra Sanchez-Roige[41], Frank J van Rooij[15], Erdogan Taskesen[1], Yang Wu[19], Futao Zhang[19], *23andMe Research Team*, *eQTLgen Consortium*, *International Cannabis Consortium*, *Psychiatric Genomics Consortium*, *Social Science Genetic Association Consortium*, Adam Auton[30], Jason D Boardman[8,7], David W Clark[34], Andrew Conlin[42], Conor C Dolan[13], Urs Fischbacher[43,44], Patrick JF Groenen[45,2], Kathleen Mullan Harris[46,47], Gregor Hasler[48], Albert Hofman[15,49], Mohammad A Ikram[15], Sonia Jain[50], Robert Karlsson[51], Ronald C Kessler[52], Maarten Kooyman[53], James MacKillop[54,55], Minna Männikkö[35], Carlos Morcillo-Suarez[40], Matthew B McQueen[56], Klaus M Schmidt[57], Melissa C Smart[20], Matthias Sutter[58,59], A Roy Thurik[14,2,60], Andre G Uitterlinden[61], Jon White[62], Harriet de Wit[63], Jian Yang[19,64], Lars Bertram[37,65,66], Dorret Boomsma[13], Tõnu Esko[67], Ernst Fehr[22], David A Hinds[30], Magnus Johannesson[68], Meena Kumari[20], David Laibson[5], Patrik KE Magnusson[51], Michelle N Meyer[69], Arcadi Navarro[40,70], Abraham A Palmer[41], Tune H Pers[16,17], Danielle Posthuma[1,71], Daniel Schunk[72], Murray B Stein[41,50], Rauli Svento[42], Henning Tiemeier[15], Paul RHJ Timmers[34], Patrick Turley[25,26,73], Robert J Ursano[74], Gert G Wagner[75,28], James F Wilson[34,76], Jacob Gratten[19], James J Lee[77], David Cesarini[78], Daniel J Benjamin[9,73,79]§, Philipp D Koellinger[1,3,80]§, Jonathan P Beauchamp[18]*§

* Lead analysts; authors 2-5 listed alphabetically.
§ These authors oversaw this work.

[1] Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research, Amsterdam Neuroscience, VU University Amsterdam, De Boelelaan 1085, Amsterdam, 1081 HV, The Netherlands

[2] Erasmus University Rotterdam Institute for Behavior and Biology, Erasmus School of Economics, Erasmus University Rotterdam, Burgemeester Oudlaan 50, Rotterdam, 3062 PA Rotterdam, The Netherlands

[3] School of Business and Economics, VU University Amsterdam, De Boelelaan 1105, Amsterdam, 1081 HV, The Netherlands

[4] Department of Economics, University of Zurich, Schönberggasse 1, Zurich, 8001, Switzerland

[5] Economics, Harvard University, 1805 Cambridge St, Cambridge, MA 02138, USA

[6] Institute for Behavioral Genetics, University of Colorado Boulder, 1480 30th Street, Boulder, CO 80309, USA

[7] Institute of Behavioral Science, University of Colorado Boulder, 1440 15th Street, Boulder, CO 80309, USA

[8] Department of Sociology, University of Colorado Boulder, UCB 327, Boulder, CO 80309, USA

[9] Center for the Advancement of Value in Musculoskeletal Care, Hospital for Special Surgery, 535 East 70th Street, New York, NY 10021, USA

[10] Center for Economic and Social Research, University of Southern California, 635 Downey Way, Los Angeles, CA 90089, USA

[11] Amsterdam School of Economics, University of Amsterdam, Roetersstraat 11, Amsterdam, 1018 WB, The Netherlands

[12] Amsterdam Brain and Cognition, University of Amsterdam, Nieuwe Achtergracht 129B, Amsterdam, 1001 NK, Netherlands

[13] Biological psychology, Vrije Universiteit Amsterdam, van der Boechorstraat 1, Amsterdam, 1087 BT, The Netherlands

[14] Department of Applied Economics, Erasmus School of Economics, Erasmus University Rotterdam, Burgemeester Oudlaan 50, Rotterdam, 3062 PA, The Netherlands

[15] Department of Epidemiology, Erasmus Medical Center, Wytemaweg 80, Rotterdam, 3015 GE, The Netherlands

[16] Novo Nordisk Foundation Centre for Basic Metabolic Research, University of Copenhagen, Copenhagen, 2200, Denmark

[17] Department of Epidemiology Research, Statens Serum Institut, Copenhagen, 2300, Denmark

[18] Department of Economics, University of Toronto, 150 St. George St., Toronto, Ontario, M5S 3G7, Canada

[19] Institute for Molecular Bioscience, The University of Queensland, Brisbane, 4072, Australia

[20] Institute for Social & Economic Research, University of Essex, Wivenhoe Park, Colchester, CO4 3SQ, UK

[21] Seminar for Statistics, Department of Mathematics, ETH Zurich, Raemistrasse 101, Zurich, 8092, Switzerland

[22] Department of Economics, University of Zurich, Bluemlisalpstrasse 10, Zurich, 8006, Switzerland

[23] Stuyvesant High School, New York, NY 10282, USA

[24] Psychiatric & Neurodevelopmental Genetics Unit, Massachusetts General Hospital, Boston, MA 02114, USA

[25] Analytical and Translational Genetics Unit, Massachusetts General Hospital, Richard B. Simches Research building, 185 Cambridge St, CPZN-6818, Boston, MA 02114, USA

[26] Stanley Center for Psychiatric Genomics, The Broad Institute at Harvard and MIT, 75 Ames St, Cambridge, MA 02142, USA

[27] Nuffield Department of Population Health, University of Oxford, Old Road Campus, Oxford, OX3 7LF, UK

[28] Socio-Economic Panel Study, DIW Berlin, Mohrenstr. 58, Berlin, 10117, Germany

[29] Max Planck Institute for Demographic Research, Konrad-Zuse-Str. 1, Rostock, 18057, Germany

[30] Research, 23andMe, Inc., 899 W Evelyn Ave, Mountain View, CA 94041, USA

[31] Barcelona Institute for Global Health (ISGlobal), Barcelona, 8003, Spain

[32] Universitat Pompeu Fabra (UPF), Barcelona, 8003, Spain

[33] CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, 280229, Spain

[34] Centre for Global Health Research, Usher Institute for Population Health Sciences and Informatics, University of Edinburgh, Teviot Place, Edinburgh, EH8 9AG, Scotland

[35] Center for Life Course Health Research, University of Oulu, Aapistie 5, Oulu, 90220, Finland

[36] Department of Economics, University of California San Diego, La Jolla, CA 92093, USA

[37] Institute of Neurogenetics, University of Lübeck, Maria-Goeppert-Str. 1, Lübeck, 23562, Germany

[38] Institute of Human Genetics, University of Lübeck, Ratzeburger Allee 160, Lübeck, 23538, Germany

[39] Team Loyalty BV, Debbemeerstraat 25, Hoofddorp, 2131 HE, The Netherlands

[40] Departament de Ciències Experimentals i de la Salut, Universitat Pompeu Fabra, Dr. Aiguader, 88, Barcelona, 8003, Spain

[41] Psychiatry, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

[42] Department of Economics and Finance, Oulu Business School, University of Oulu, Oulu, 90570, Finland

[43] Department of Economics, University of Konstanz, Universitätssstraße 10, Konstanz, 78457, Germany

[44] Thurgau Institute of Economics, Hauptstrasse 90, Kreuzlingen, 8280, Switzerland

[45] Department of Econometrics, Erasmus University Rotterdam, Burgemeester Oudlaan 50, Rotterdam, 3062 PA Rotterdam, The Netherlands

[46] Department of Sociology , University of North Carolina at Chapel Hill, 201 Hamilton Hall CB #3210, Chapel Hill, NC 27599, USA

[47] Carolina Population Center, University of North Carolina at Chapel Hill, 123 West Franklin St., Suite 2190, Chapel Hill, NC 27516, USA

[48] Division of Molecular Psychiatry, Translational Research Center, University Hospital of Psychiatry, University of Bern, Bolligenstrasse 111, Bern, 3000, Switzerland

[49] Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA

[50] Family Medicine and Public Health, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

[51] Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Nobels väg 12a, Stockholm, 171 77, Sweden

[52] Department of Health Care Policy, Harvard Medical School, 180 Longwood Ave., Boston, MA 02115, USA

[53] SURFsara, Science Park 140, Amsterdam, 1098 XG, The Netherlands

[54] Peter Boris Centre for Addictions Research, McMaster University/St. Joseph's Healthcare Hamilton, 100 West 5th Street, Hamilton, Ontario, L8P 3R2, Canada

[55] Homewood Research Institute, 150 Delhi Street, Riverslea Building, Guelph, Ontario, N1E 6K9, Canada

[56] Department of Integrative Physiology, University of Colorado Boulder, 354 UCB, Boulder, CO 80309, USA

[57] Department of Economics, University of Munich, Ludwigstrasse 28, Munich, 80539, Germany

[58] Department of Economics, University of Cologne, Albert-Magnus-Platz, Cologne, 50923, Germany

[59] Experimental Economics Group, Max Planck Institute for Research into Collective Goods, Schumacher Strasse 10, Bonn, D-51113, Germany

[60] Montpellier Business School, Montpellier, 34080, France

[61] Internal Medicine, Erasmus Medical Center, Wytemaweg 80, Rotterdam, 3015 CN, The Netherlands

[62] UCL Genetics Institute, Department of Genetics, Evolution and Environment, University College London, Darwin Building, Gower Street, London, WC1E 6BT, UK

[63] Psychiatry and Behavioral Neuroscience, University of Chicago, 5841 S. Maryland Ave., MC 3077, Rm L485, Chicago, IL 60637, USA

[64] Queensland Brain Institute, The University of Queensland, Brisbane, 4072, Australia

[65] Institute of Cardiogenetics, University of Lübeck, Maria-Goeppert-Str. 1, Lübeck, 23562, Germany

[66] School of Public Health, Imperial College London, South Kensington Campus, London, SW7 2AZ, UK

[67] Estonian Genome Center, University of Tartu, Tartu, 51010, Estonia

[68] Department of Economics, Stockholm School of Economics, Box 6501, Stockholm, 113 83, Sweden

[69] Center for Translational Bioethics and Health Care Policy, Geisinger Health System, Danville, PA 17822, USA

[70] Institució Catalana de Recerca i Estudis Avançats (ICREA), Parc de Recerca Bomèdica. Carrer Dr. Aiguader, 88, Barcelona, 8003, Spain

[71] Department of Clinical Genetics, VU Medical Centre, De Boelelaan 1085, Amsterdam, 1081 HV, The Netherlands

[72] Department of Economics, Johannes Gutenberg University, Jakob-Welder-Weg 4, Mainz, 55128, Germany

[73] Behavioral and Health Genomics Center, University of Southern California, 635 Downey Way, Los Angeles, CA 90089, USA

[74] Psychiatry, Uniformed Services University Health Science, Bethesda, MD 20814, USA

[75] Max Planck Institute for Human Development, Lentzeallee 94, Berlin, 14195, Germany

[76] MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Edinburgh, EH4 2XU, Scotland

[77] Department of Psychology, University of Minnesota Twin Cities, 75 East River Parkway, Minneapolis, MN 55455, USA

[78] Department of Economics, New York University, 19 W 4th St, 6 FL, NY 10012, USA

[79] National Bureau of Economic Research, 1050 Massachusetts Ave, Cambridge, MA 02138, USA

[80] German Institute for Economic Research, DIW Berlin, Mohrenstr. 58, Berlin, 10117, Germany