Baldspot/ELOVL6 is a conserved modifier of disease and the ER stress response

**ELOVL6* modifies ER stress

Rebecca A. S. Palu and Clement Y. Chow*

Department of Human Genetics, University of Utah School of Medicine, Salt Lake City, UT

84112 USA

* Corresponding author
Clement Y. Chow
Department of Human Genetics
University of Utah School of Medicine
EIHG 5150
15 North 2030 East
Salt Lake City, UT 84112
cchow@genetics.utah.edu
16

ABSTRACT

Endoplasmic reticulum (ER) stress is an important modifier of human disease. Genetic variation

in genes involved in the ER stress response has been linked to inter-individual differences in this

response. However, the mechanisms and pathways by which genetic modifiers are acting on the

ER stress response remain unclear. In this study, we characterize the role of the long chain fatty

acid elongase Baldspot (ELOVL6) in modifying ER stress and disease. We demonstrate that loss

of *Baldspot* rescues degeneration and reduces IRE1-dependent *Xbp1* splicing and cell death in

a *Drosophila* model of retinitis pigmentosa and ER stress ($Rh1^{G69D}$). Dietary supplementation of

stearate bypasses the need for Baldspot activity. Finally, we demonstrate that *Baldspot* regulates

ER stress across different tissues and induction methods. Our findings suggest that *ELOVL6* is a

promising target in the treatment of not only retinitis pigmentosa, but a number of different ER

stress related disorders.

INTRODUCTION

Phenotypic heterogeneity is common in simple and complex diseases and is the driving force behind the Precision Medicine Initiative [1-4]. Genetic variation among individuals accounts for much of this heterogeneity, but the identity and nature of modifying genes or variants is largely unknown [4, 5]. These modifying variants are often cryptic and may not influence the physiology or visible phenotypes of healthy individuals, but can alter the expression of disease phenotypes. Understanding the role of modifiers and the pathways in which they function will enable the development of patient-specific therapeutic approaches.

One such process is the response to endoplasmic reticulum (ER) stress, which occurs when misfolded proteins accumulate in the lumen of the ER. This activates the Unfolded Protein Response (UPR) in an attempt to return the cell to homeostasis. Failure to do so will eventually result in apoptosis [6]. Conditions of chronic ER stress lead to excess apoptosis, tissue degeneration, and dysfunction. In rare cases, primary mutations of key ER stress pathway components cause Mendelian syndromes [7-11]. More commonly, ER stress can exacerbate disease, including obesity [12], neurological disease [13], retinal degeneration [14], and some cancers [15]. Experimental genetic or pharmacological manipulation of ER stress levels can alter disease phenotypes, suggesting that inter-individual differences in ER stress responses may influence disease severity. Indeed, human [16], mouse [17], and *Drosophila* [18, 19] show extensive genetic variation in their response to ER stress. Understanding the role of genetic variation in modulating ER stress pathways may provide more therapeutic targets and improve the accurate identification of high risk patients.

In a previous study, we crossed the $Rh1^{G69D}$ model of retinitis pigmentosa (RP) into the *Drosophila* Genetic Reference Panel (DGRP) to identify genetic modifiers of this disease [19]. In this model, retinal degeneration is induced by overexpression of misfolded rhodopsin in the developing larval

eye disc [20], resulting in chronic ER stress and apoptosis. The DGRP is used to study the genetic

architecture underlying complex traits. It is comprised of approximately 200 inbred strains, such

that each strain represents a single, wild-derived genome. Thus, the DGRP captures genetic

variation that is present in a natural, wild population. Importantly, we have the whole genome

sequence of each strain, allowing phenotype-genotype studies [21]. The degree of degeneration

induced by this model varied substantially across the strains of the DGRP. Using an association

analysis, we identified a number of promising candidates that modify this RP phenotype and have

putative roles in ER stress [19].

In this study, we demonstrate that *Drosophila Baldspot,* one of the candidate RP modifiers

identified in our genetic variation screen, is both a disease modifier of ER stress-induced retinal

degeneration and a more general modifier of ER stress across different physiological contexts.

Baldspot, which is orthologous to mammalian *ELOVL6,* is an ER-associated long chain fatty acid

elongase which catalyzes the extension of palmitate to stearate [22, 23]. We show that loss of

Baldspot rescues retinal degeneration by reducing ER stress signaling. Loss of Baldspot elongase

activity reduces IRE1/Xbp1 activity and downstream apoptosis, without affecting the RIDD activity

of IRE1 or the misfolded protein load. Strikingly, in the absence of Baldspot activity, increasing

levels of stearate through dietary supplementation bypasses the need for Baldspot to induce

IRE1/Xbp1 signaling. Finally, we show that Baldspot regulates ER stress in a number of different

genetic and chemically-induced ER stress models across multiple tissue types, making *Baldspot*

a potentially wide-reaching modifier of ER stress-associated diseases. Understanding how

modifier genes alter disease outcomes is the first step to developing personalized therapies.

RESULTS

A SNP in *Baldspot* is associated with *Rh1*^{G69D}-induced retinal degeneration

In previous work, we overexpressed a misfolded rhodopsin protein in the developing eye disc

(*GMR-GAL4>UAS-Rh1^{G69D}*), the larval precursor to the *Drosophila* retina and eye, in the 200

strains from the DGRP [19]. Using a genome-wide association approach, we identified SNPs that

are associated with degree of retinal degeneration. One of these candidate SNPs,

Chr3L:16644000 (BDGP R5/dm3), is located in intron four of *Baldspot*. *Baldspot* is the only

Drosophila orthologue of mammalian *ELOVL6*, a member of the very long chain fatty acid

elongase family of enzymes (Figure 1A) [24]. Strains carrying the minor allele (C, 18880 ± 3010

pixels) exhibit more severe *Rh1^{G69D}*–induced retinal degeneration as compared to strains carrying

the major allele (T, 21822 ± 2294 pixels) (Figure 1B; $p < 0.005$). This is also apparent when

examining the range of degeneration in strains containing each of these alleles (Figure 1C).

Based on these results, we hypothesized that altering the expression or activity of *Baldspot* may

impact degeneration.

Loss of *Baldspot* rescues *Rh1^{G69D}*-induced retinal degeneration.

To test whether eliminating *Baldspot* expression can modify *Rh1^{G69D}*-induced retinal

degeneration, we expressed an RNAi construct targeted against *Baldspot* in the *GMR-*

GAL4>UAS-Rh1^{G69D} background. We found that loss of *Baldspot* expression results in partial

rescue of Rh1^{G69D}-induced retinal degeneration (*Rh1^{G69D}/Baldspoti*) as compared to controls

expressing only the Rh1^{G69D} misfolded protein and no RNAi (*Rh1^{G69D}* controls) (Figure 2A).

Quantification of eye size demonstrates a significantly larger, less degenerate eye in

Rh1^{G69D}/Baldspoti flies (14043 ± 960 pixels) as compared to *Rh1^{G69D}* controls (11942 ± 473 pixels)

($P = 7.4 \times 10^{-6}$) (Figure 2B). Loss of *Baldspot* on a wild-type background resulted in no change in

eye size or phenotype (Figure 2C). We conclude that loss of *Baldspot* expression provides

significant rescue of *Rh1^{G69D}*-induced retinal degeneration.

Loss of *Baldspot* in the *Rh1^{G69D}* eye disc reduces ER stress and apoptosis

Rescue of eye size in the $Rh1^{G69D}$ model is often accompanied by reduction in apoptosis [25, 26].

Based on this and the rescue effect we observed, we hypothesized that apoptosis in the $Rh1^{G69D}$

eye discs would also be reduced upon loss of *Baldspot* expression. Indeed, we found that

apoptosis, as measured by terminal deoxynucleotidyl transferase dUTP nick end labeling

(TUNEL) staining, is reduced in $Rh1^{G69D}$/*Baldspoti* eye discs as compared to $Rh1^{G69D}$

controls. (Figure 2D).

Excessive ER stress in the $Rh1^{G69D}$ eye discs leads to apoptosis and degeneration in the adult

eyes. To determine if the decrease in apoptosis we observed was due to changes in the ER stress

response, we measured *Xbp1* splicing. Under conditions that induce ER stress, *Xbp1* transcript

is spliced by IRE1, one of three transmembrane sensors of misfolded proteins in the ER. The in-

frame *Xbp1* transcript is translated and acts as a transcription factor, inducing genes involved in

the UPR [27, 28]. The level of spliced *Xbp1* transcript is directly proportional to the degree of

activation of the ER stress response. To measure IRE1 signaling and *Xbp1* splicing in the absence

of *Baldspot* expression, we used a previously characterized transgenic *Xbp1-EGFP* marker [20,

29-31]. In this transgene, the 5' end of the *Xbp1* mRNA is expressed under the control of the *UAS*

promoter. The 3' end of the gene, downstream of the IRE1 splice site, has been replaced with

sequence encoding *EGFP*. As with the endogenous *Xbp1* transcript, the in-frame, spliced *Xbp1-*

EGFP transcript, and subsequently the Xbp1-EGFP fusion protein, increases as ER stress

signaling increases. Thus, EGFP signal increases as ER stress signaling increases. As previously

reported, expression of *UAS-Xbp1-EGFP* in $Rh1^{G69D}$ controls results in increased EGFP,

indicating high levels of ER stress (Figure 2E) [20, 29-31]. When *UAS-Xbp1-EGFP* is expressed

in $Rh1^{G69D}$/*Baldspoti* flies, we see a significant reduction in EGFP signal (3.8 ± 0.8 fluorescent

units vs. 6.5 ± 1.1 in controls; $P = 9.6 \times 10^{-4}$), suggestive of reduced ER stress signaling (Figure

2E and F). Strikingly, the $Rh1^{G69D}$/*Baldspoti* eyes discs display no substantial change in rhodopsin

protein levels (Figure 2E), suggesting that loss of *Baldspot* expression reduces IRE1/Xbp1 ER

stress signaling without affecting the accumulation or degradation of misfolded Rh1 protein.

Stearate supplementation bypasses the requirement for *Baldspot*

Increased cellular concentration of stearate is associated with increased activation of the ER

stress response [32, 33]. It is likely that the modifying effect of Baldspot is linked to its fatty acid

elongation activity. Baldspot converts palmitate to stearate and the absence of Baldspot function

should result in lower levels of stearate. We hypothesized that lower levels of stearate underlies

the rescuing effect, reduced ER stress signaling, and retinal degeneration we observed, and that

this signaling could be restored if stearate were supplemented during development. To test this,

we raised $Rh1^{G69D}/Baldspoti$ and $Rh1^{G69D}$ control flies on media with or without 10% stearate and

measured eye size in adults. Stearate supplementation resulted in increased degeneration and

reduced eye size in $Rh1^{G69D}/Baldspoti$ flies (14033 ± 1553 pixels) as compared to those raised on

standard media with no stearate (16846 ± 1699 pixels) (P = 1 x 10^{-7}) (Figure 3A). This increase

in degeneration in $Rh1^{G69D}/Baldspoti$ flies was accompanied by an increase in IRE1 activity and

spliced *Xbp1*, as evidenced by an increase in Xbp1-EGFP expression in eye discs (Figure 3B).

Stearate supplementation had no effect on eye size, retinal degeneration, or Xbp1-EGFP signal

in $Rh1^{G69D}$ control flies (13478 ± 1197 pixels vs. 12873 ± 1006 pixels) (Figure 3 A-B), likely

because of the high level of ER stress signaling already present. High stearate levels are sufficient

to induce ER stress signaling and bypasses the need for Baldspot activity.

***Baldspot* is a general modifier of ER stress signaling**

Because *Baldspot* is ubiquitously expressed (www.flyatlas2.org/), it is possible that *Baldspot's*

modifying role in ER stress is generalizable across different contexts and tissues. We first tested

whether Baldspot modifies ER stress in another tissue. Because wing size is a visible and easily

scorable trait, we used the *MS1096-GAL4* driver to express $Rh1^{G69D}$ in the developing larval wing

disc (wing-*Rh1^{G69D}*). The Rh1^{G69D} protein acts as a misfolded protein, inducing ER stress only in

that tissue. The wing disc gives rise to the adult wing, and perturbations in cellular homeostasis

in this tissue result in small, misshapen wings [34]. Expression of wing-*Rh1^{G69D}* produces a

phenotype similar to a vestigial wing (Figure 4A). When the RNAi construct targeted against

Baldspot was concurrently expressed in the wing disc (wing-*Rh1^{G69D}/Baldspoti*), this vestigial

phenotype was partially rescued. The adult wings showed an increase in size and an increase in

unfolding upon eclosion (Figure 4A). Similar to what we observed in the eye discs, this partial

rescue was associated with a decrease in the level of Xbp1-EGFP in the wing disc (Figure 4B). It

appears that loss of *Baldspot* modifies IRE1/Xbp1 signaling across different tissues and is not

eye disc-specific.

To test whether Baldspot modifies ER stress signaling by pharmacological induction, we used

tunicamycin to induce ER stress. Tunicamycin inhibits N-linked glycosylation in the ER and results

in massive misfolding, ER stress, and a robust UPR [35]. Treatment with tunicamycin in

Drosophila larvae results in significant developmental lethality. We acutely exposed 2nd instar

control larvae (*Tub-GAL4*, no RNAi) and larvae with ubiquitous knockdown of *Baldspot* (*Tub-*

GAL4>Baldspot RNAi) to tunicamycin or control DMSO and scored survival to pupation. To

account for any larval lethality due to loss of *Baldspot*, we normalized the survival of larvae on

tunicamycin to the DMSO treatment. Strikingly, larvae lacking *Baldspot* were less susceptible to

ER stress-induced lethality than control larvae (Figure 4D), as evidenced by increased survival to

pupation. The modifying effect of *Baldspot* expression on the ER stress response is generalizable

across different tissues and methods of ER stress induction.

Baldspot does not regulate IRE1 RIDD function

Because our data indicates that Baldspot regulates Ire1/Xbp1 signaling, we next tested whether

Baldspot regulates another key IRE1 function: Regulated IRE1 Dependent Degradation of

mRNAs (RIDD) [36-38]. In addition to *Xbp1* splicing, activated IRE1 is also important for the

degradation of a number of ER-associated transcripts through RIDD. To test if RIDD activity is

affected by loss of *Baldspot* expression, we measured levels of known RIDD target mRNAs, under

ER stress, in control cells and tissues with or without *Baldspot* expression. We employed S2 cell

culture and treated S2 cells with dsRNA targeting either *Baldspot* or *EGFP* as a control. This

treatment resulted in a near complete reduction in *Baldspot* transcript levels (~7.8% of control; P

= 1.3 x 10^{-9}) (Figure 5A). We used Dithiothreitol (DTT) to induce ER stress in S2 cells. DTT is a

reducing agent that breaks disulfide bonds in proteins, resulting in misfolded proteins and a strong

ER stress response [39]. As expected, treatment of cells with DTT increased *Xbp1* splicing ~3-

fold, indicative of ER stress. In agreement with our *in vivo* studies in the *Rh1^{G69D}* fly, this splicing

was significantly reduced in the absence of *Baldspot*, with only ~2-fold increase (P = 0.023),

indicating that IRE1 signaling is disrupted (Figure 5B).

We found that under DTT-induced ER stress, there was a similar degree of reduction (~81% of

control) in RIDD target expression in the cells treated with DsRNA against *Baldspot* or control

DsRNA against EGFP (Figure 5C). Non-RIDD targets, such as *Actin5C*, were unaffected (Figure

5C). It appears that loss of *Baldspot* preferentially affects IRE1/Xbp1 signaling, while RIDD

function remains intact.

DISCUSSION

The ER stress response has the potential to be an important modifier of human disease. ER

stress is activated in a wide variety of diseases ranging from Mendelian degenerative diseases to

complex metabolic disorders [14]. The role of the ER stress response is complex and in some

cases, a stronger response is beneficial and in other cases, a strong response underlies the

disease. In some forms of retinitis pigmentosa, like the focus of this study, loss of vision is a direct

result of the ER stress-induced cell death [40]. Neurodegenerative diseases such as Parkinson's,

Alzheimer's, Amyotrophic Lateral Sclerosis (ALS), and prion disease are all associated with the

accumulation of cytoplasmic misfolded proteins that can activate a secondary ER stress response

and subsequent cell death [15]. In diabetes and obesity, misfolded proteins and the ER stress

response are at least partially responsible for the cellular dysfunction that is a hallmark of the

disease state [41]. In some cancers, however, cells selectively upregulate ER stress pathways

associated with survival, while inhibiting those associated with cell death [42]. Genetic variation

that alters the degree to which the ER stress response is activated may modify disease in the

population.

We and others have shown that genetic variation can influence how and to what extent the ER

stress response is activated. Genetic variation in *Drosophila* [18, 19], mouse [17], and human

cells [16] can impact the expression of genes involved in ER stress pathways. Furthermore, this

variation in expression has been linked to differences in the UPR and the downstream activation

of cell death [17-19]. It is apparent that cryptic genetic variation drives these extensive inter-

individual differences in the ER stress response [17]. Stress or disease conditions reveal the effect

of this otherwise silent genetic variation. The same variation that drives differences in this

important stress response might also influence disease heterogeneity. Indeed, modulating the ER

stress response is an effective way of modulating disease outcome in models of human disease.

[43-48].

In this study we show that loss of *Baldspot/ELOVL6* activity can affect the outcome of retinal

degeneration and has the potential to be a broader regulator of ER stress-associated diseases.

As a 16-to-18 carbon fatty acid elongase in the ER membrane, this enzyme is responsible for

increasing the concentration of C18 stearate, a saturated long chain fatty acid [24, 33]. Increases

in stearate concentrations in the ER membrane can induce and enhance ER stress through the

activation of IRE1/Xbp1 [32, 33]. Here, we show that reducing the activity of Baldspot/ELOVL6

reduces the degree to which the UPR and apoptosis are induced in models of ER stress. Our

study suggests that Baldspot/ELOVL6 has a modifying effect on the IRE1 branch of the ER stress

response. IRE1 has several roles, including catalyzing *Xbp1* signaling and RNA degradation

through RIDD. Strikingly, we found that Baldspot/ELOVL6 specifically influences *Xbp1* signaling

and not RIDD function. Despite utilizing the same endonuclease domain, *Xbp1* splicing and RIDD

have different roles in development, cellular homeostasis, and stress responses [49]. This is also

consistent with observations that the activation of various arms of the ER stress response can be

alternately beneficial or detrimental, dependent upon the disease and tissue context [50]. It is

unclear how these functions are partitioned by IRE1, but further studies exploring the

Baldspot/ELOVL6 interaction with IRE1 may clarify the differential regulation of *Xbp1* signaling

and RIDD function.

Therapeutics that reduce the activity of this key fatty acid elongase could conceivably be used to

treat or delay progression in a variety of ER stress-associated diseases. In RP patients with

mutations that cause the accumulation of misfolded rhodopsin, similar to our *Drosophila* model,

reduction of ELOVL6 expression might reduce cell death, delaying vision loss. We present

evidence that the modulation of the ER stress response through Baldspot/ELOVL6 activity may

impact the severity of diseases beyond RP. Particularly intriguing is that *Baldspot/ELOVL6* can

alter the ER stress response when it is induced by different conditions. Regardless of the

mechanism by which ER stress is induced, inhibiting ELOVL6 activity reduces IRE1/Xbp1

signaling. In fact, in a recent study of the *db/db* mouse model of diabetes, loss of *ELOVL6* alters

insulin sensitivity and the expression of ER stress genes [22]. It is likely that reducing

Baldspot/ELOVL6 expression would also alter complex diseases where ER stress contributes to

degeneration and cell death. In some cases, even a small delay in degeneration could

dramatically improve the quality of life.

We have identified *Baldspot/ELOVL6* in *Drosophila* as a novel modifier of the ER stress response.

Its potential as a therapeutic target extends beyond RP to any number of ER stress-associated

disorders. The identification of *Baldspot* in a natural genetic variation screen [19] suggests that

variation in expression of this gene is well tolerated under healthy conditions and might be

amenable to modulation. Identifying variable elements of the ER stress response pathway may

prove to be a successful strategy to nominating new modifiers with broad applications to many

ER stress-associated diseases.

METHODS

Fly stocks and maintenance

Flies were raised at room temperature on standard diet based on the Bloomington Stock Center

standard medium with malt. The strain containing *GMR-GAL4* and *UAS-Rh1^{G69D}* on the second

chromosome (*GMR>Rh1^{G69D}*) has been previously described [19]. The following strains are from

the Bloomington Stock Center: *MS1096-GAL4* (8696), *Baldspot* RNAi (44101), control attP40

(36304). The strain containing the *UAS-Xbp1-EGFP* transgene were a gift from Don Ryoo (NYU)

[20].

Eye/wing imaging

For eye and wing images, adult females were collected under CO_2 anesthesia and aged to 2-7

days, then flash frozen on dry ice. Eyes were imaged at 3X magnification using a Leica EC3

camera. Wings were dissected away from the body, then imaged at 4.5X magnification using the

same camera. Eye area was measured in ImageJ as previously described [19].

Immunohistochemistry

Eye discs and wing discs were dissected from wandering L3 larvae in cold 1X PBS, then

immediately transferred to cold 4% PFA on ice. Tissues were fixed in 4% PFA for 15-20 min, then

washed in 1XPAT (0.1% TritonX100) prior to blocking with 5% normal donkey serum. Tissues

were stained with primary antibodies for rhodopsin (1:50, Developmental Studies Hybridoma

Bank #4C5) and GFP (1:2000, Thermo-Fisher #A6455). Apoptosis was monitored using the

ApopTag Red *In Situ* Apoptosis Detection Kit (Millipore #S7165). Tissues were mounted in

Slowfade™ Diamond Antifade Mountant (ThermoFisher #S36967) and imaged with an Olympus

FV1000 confocal microscope.

Stearate supplementation

Crosses were set up on the control, standard media alone, or standard media with 10% stearate

supplementation. To make stearate supplemented media, standard media was melted in the

microwave and kept at 98°C on a stir plate. Stearate (10% final concentration) was added to this

media and mixed until lipids were homogenous with the media. The standard control diet received

the same treatment, but no additional stearate was added. To determine retinal degeneration,

flies were aged and imaged as described above. Antibody staining was performed on the eye

imaginal discs of L3 larvae, also as described above.

Tunicamycin treatment

Crosses to generate the indicated genotypes were set up on egg caps containing yeast paste. L2

larvae were then treated with either 10 µg/mL Tunicamycin (diluted 1:1000 from a 10 mg/mL stock

solution) or 1:1000 DMSO in Schneider's media for 30 minutes at room temperature. The larvae

were then washed in 1XPBS twice and placed on solid media containing 6% yeast and 1% agar.

Viability was determined by survival to pupation. Survival for each genotype was normalized to

the DMSO-treated control condition. Each replicate represents ~25-60 larvae per genotype; a

total 180 to 250 larvae per genotype were analyzed across all replicates.

S2 cells

DsRNA was generated using the MEGAscript™ T7 Transcription kit (ThermoFisher #AM1334),

with primers for *EGFP* (F: TTAATACGACTCACTATAGGGAGACCACAAGTTCAGCGTGTCC

and R: TTAATACGACTCACTATAGGGAGAGGGGTGTTCTGCTGGTAGTG) and *Baldspot* (F:

TTAATACGACTCACTATAGGGAGAATCCGCCCAGGTTCATCTCG and R:

TTAATACGACTCACTATAGGGAGAGTCACATCTCGCAGCGCAAC). S2 cells were treated with

DsRNA against *EGFP* (as a control) or against *Baldspot* at a density of approximately 2×10^6

cells/mL in a 24-well plate. Cells were incubated with DsRNA for 4-7 days before treatment with

0.5 mM DTT for 60 min for *Xbp1* splicing or 1.0 mM DTT for 20 min for RIDD target analysis.

Control cells were treated with DMSO. RNA was isolated from cells using Trizol/chloroform

extraction and used to generate cDNA. Knockdown of *Baldspot* was confirmed using qPCR

(primers: F: TGCTGGTCATCTTCGGTGGTC and R: ACGCAGACGGAGTGGAAGAG). RIDD

target levels were also confirmed by qPCR (*sparc* primers: F: AAAATGGGCTGTGTCCTAACC

and R: TGCAGCACAATCTACTCAATCC). *Xbp1* splicing was evaluated from the cDNA using

PCR (primers for *Xbp1:* F: TCAGCCAATCCAACGCCAG and R:

TGTTGTATACCCTGCGGCAG). The spliced and unspliced bands were separated on a 12%

acrylamide gel and the proportion of these bands quantified using ImageJ software.

Statistics

Statistics were calculated using R software. P-values were determined using ANOVA for eye size,

fluorescence levels, and transcript levels from S2 cells. Tukey multiple testing correction was

used for the fatty acid feeding experiment. Bonferroni correction was used for *Xbp1* splicing in S2

cells. A pairwise T-test was performed for larval tunicamycin treatment. A cutoff of P = 0.05 was

used for significance.

Acknowledgements

We thank Drs. Aylin Rodan and Carl Thummel for use of equipment and sharing reagents. We thank

Emily Coelho for helpful comments.

Funding

This research was supported by an NIH/NIGMS R35 award (1R356M124780) and a Glenn Award

from the Glenn Foundation for Medical Research to CYC. CYC is the Mario R. Capecchi

Endowed Chair in Genetics. R.A.S.P. was supported by a NIDDK T32 fellowship

(5T32DK091217).

REFERENCES

1. Nadeau JH. Modifier genes in mice and humans. Nat Rev Genet. 2001;2(3):165-174.
353
354 2. Scriver CR, Waters PJ. Monogenic traits are not simple: lessons from phenylketonuria. Trends
355 Genet. 1999;15(7):267-272.
356
357 3. Gallati S. Disease-modifying genes and monogenic disorders: experience in cystic fibrosis.
358 Appl Clin Genet. 2014;7:133-146.
359
360 4. Queitsch C, Carlson KD, Girirajan S. Lessons from model organisms: phenotypic robustness
361 and missing heritability in complex disease. PLoS Genet. 2012;8(11):e1003041.
362
363 5. Chow CY. Bringing genetic background into focus. Nat Rev Genet. 2016;17(2):63-64.
364
365 6. Schroder M, Kaufman RJ. The mammalian unfolded protein response. Annu Rev Biochem.
366 2005;74:739-789.
367
368 7. Ansar M, Santos-Cortez RL, Saqib MA, Zulfiqar F, Lee K, Ashraf NM, et al. Mutation of ATF6
369 causes autosomal recessive achromatopsia. Hum Genet. 2015;134(9):941-950.
370
371 8. Eura Y, Yanamoto H, Arai Y, Okuda T, Miyata T, Kokame K. Derlin-1 deficiency is embryonic
372 lethal, Derlin-3 deficiency appears normal, and Herp deficiency is intolerant to glucose load and
373 ischemia in mice. PLoS One. 2012;7(3):e34298.
374
375 9. Delepine M, Nicolino M, Barrett T, Golamaully M, Lathrop GM, Julier C. EIF2AK3, encoding
376 translation initiation factor 2-alpha kinase 3, is mutated in patients with Wolcott-Rallison
377 syndrome. Nat Genet. 2000;25(4):406-409.
378
379 10. Kohl S, Zobor D, Chiang WC, Weisschuh N, Staller J, Gonzalez Menendez I, et al. Mutations
380 in the unfolded protein response regulator ATF6 cause the cone dysfunction disorder
381 achromatopsia. Nat Genet. 2015;47(7):757-765.
382

11. Anttonen AK, Mahjneh I, Hamalainen RH, Lagier-Tourenne C, Kopra O, Waris L, et al. The gene disrupted in Marinesco-Sjogren syndrome encodes SIL1, an HSPA5 cochaperone. Nat Genet. 2005;37(12):1309-1311.

12. Yilmaz E. Endoplasmic Reticulum Stress and Obesity. Adv Exp Med Biol. 2017;960:261-276.

13. Scheper W, Hoozemans JJ. The unfolded protein response in neurodegenerative diseases: a neuropathological perspective. Acta Neuropathol. 2015;130(3):315-331.

14. Sano R, Reed JC. ER stress-induced cell death mechanisms. Biochim Biophys Acta. 2013;1833(12):3460-3470.

15. Hetz C, Saxena S. ER stress and the unfolded protein response in neurodegeneration. Nat Rev Neurol. 2017;13(8):477-491.

16. Dombroski BA, Nayak RR, Ewens KG, Ankener W, Cheung VG, Spielman RS. Gene expression and genetic variation in response to endoplasmic reticulum stress in human cells. Am J Hum Genet. 2010;86(5):719-729.

17. Chow CY, Wang X, Riccardi D, Wolfner MF, Clark AG. The genetic architecture of the genome-wide transcriptional response to ER stress in the mouse. PLoS Genet. 2015;11(2):e1004924.

18. Chow CY, Wolfner MF, Clark AG. Using natural variation in Drosophila to discover previously unknown endoplasmic reticulum stress genes. Proc Natl Acad Sci U S A. 2013;110(22):9013-9018.

19. Chow CY, Kelsey KJ, Wolfner MF, Clark AG. Candidate genetic modifiers of retinitis pigmentosa identified by exploiting natural variation in Drosophila. Hum Mol Genet. 2016;25(4):651-659.

20. Ryoo HD, Domingos PM, Kang MJ, Steller H. Unfolded protein response in a Drosophila model for retinal degeneration. EMBO J. 2007;26(1):242-252.

21. Mackay TF, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The Drosophila melanogaster Genetic Reference Panel. Nature. 2012;482(7384):173-178.

22. Matsuzaka T, Shimano H, Yahagi N, Kato T, Atsumi A, Yamamoto T, et al. Crucial role of a long-chain fatty acid elongase, Elovl6, in obesity-induced insulin resistance. Nat Med. 2007;13(10):1193-1202.

23. Senyilmaz D, Virtue S, Xu X, Tan CY, Griffin JL, Miller AK, et al. Regulation of mitochondrial morphology and function by stearoylation of TFR1. Nature. 2015;525(7567):124-128.

24. Jung A, Hollmann M, Schafer MA. The fatty acid elongase NOA is necessary for viability and has a somatic role in Drosophila sperm development. J Cell Sci. 2007;120(Pt 16):2924-2934.

25. Kang MJ, Chung J, Ryoo HD. CDK5 and MEKK1 mediate pro-apoptotic signalling following endoplasmic reticulum stress in an autosomal dominant retinitis pigmentosa model. Nat Cell Biol. 2012;14(4):409-415.

434 26. Morishita J, Kang MJ, Fidelin K, Ryoo HD. CDK7 regulates the mitochondrial localization of a
435 tail-anchored proapoptotic protein, Hid. Cell Rep. 2013;5(6):1481-1488.
436
437 27. Cox JS, Walter P. A novel mechanism for regulating activity of a transcription factor that
438 controls the unfolded protein response. Cell. 1996;87(3):391-404.
439
440 28. Sidrauski C, Walter P. The transmembrane kinase Ire1p is a site-specific endonuclease that
441 initiates mRNA splicing in the unfolded protein response. Cell. 1997;90(6):1031-1039.
442
443 29. Ryoo HD, Li J, Kang MJ. Drosophila XBP1 expression reporter marks cells under endoplasmic
444 reticulum stress and with high protein secretory load. PLoS One. 2013;8(9):e75774.
445
446 30. Sone M, Zeng X, Larese J, Ryoo HD. A modified UPR stress sensing system reveals a novel
447 tissue distribution of IRE1/XBP1 activity during normal Drosophila development. Cell Stress
448 Chaperones. 2013;18(3):307-319.
449
450 31. Huang HW, Zeng X, Rhim T, Ron D, Ryoo HD. The requirement of IRE1 and XBP1 in resolving
451 physiological stress during Drosophila development. J Cell Sci. 2017;130(18):3040-3049.
452
453 32. Wei Y, Wang D, Topczewski F, Pagliassotti MJ. Saturated fatty acids induce endoplasmic
454 reticulum stress and apoptosis independently of ceramide in liver cells. Am J Physiol Endocrinol
455 Metab. 2006;291(2):E275-281.
456
457 33. Green CD, Olson LK. Modulation of palmitate-induced endoplasmic reticulum stress and
458 apoptosis in pancreatic beta-cells by stearoyl-CoA desaturase and Elovl6. Am J Physiol
459 Endocrinol Metab. 2011;300(4):E640-649.
460
461 34. O'Brochta D, Bryant PJ. Cell degeneration and elimination in the imaginal wing disc, caused
462 by the mutationsvestigial andUltravestigial ofDrosophila melanogaster. Wilehm Roux Arch Dev
463 Biol. 1983;192(5):285-294.
464
465 35. Samali A, Fitzgerald U, Deegan S, Gupta S. Methods for monitoring endoplasmic reticulum
466 stress and the unfolded protein response. Int J Cell Biol. 2010;2010:830307.
467
468 36. Hollien J, Lin JH, Li H, Stevens N, Walter P, Weissman JS. Regulated Ire1-dependent decay
469 of messenger RNAs in mammalian cells. J Cell Biol. 2009;186(3):323-331.
470
471 37. Gaddam D, Stevens N, Hollien J. Comparison of mRNA localization and regulation during
472 endoplasmic reticulum stress in Drosophila cells. Mol Biol Cell. 2013;24(1):14-20.
473
474 38. Hollien J, Weissman JS. Decay of endoplasmic reticulum-localized mRNAs during the
475 unfolded protein response. Science. 2006;313(5783):104-107.
476
477 39. Jamsa E, Simonen M, Makarow M. Selective retention of secretory proteins in the yeast
478 endoplasmic reticulum by treatment of cells with a reducing agent. Yeast. 1994;10(3):355-370.
479
480 40. Hartong DT, Berson EL, Dryja TP. Retinitis pigmentosa. Lancet. 2006;368(9549):1795-1809.
481
482 41. Ariyasu D, Yoshida H, Hasegawa Y. Endoplasmic Reticulum (ER) Stress and Endocrine
483 Disorders. Int J Mol Sci. 2017;18(2).
484

485 42. Jain BP. An Overview of Unfolded Protein Response Signaling and Its Role in Cancer. Cancer
486 Biother Radiopharm. 2017;32(8):275-281.
487
488 43. Gonzalez-Perez P, Woehlbier U, Chian RJ, Sapp P, Rouleau GA, Leblond CS, et al.
489 Identification of rare protein disulfide isomerase gene variants in amyotrophic lateral sclerosis
490 patients. Gene. 2015;566(2):158-165.
491
492 44. Woehlbier U, Colombo A, Saaranen MJ, Perez V, Ojeda J, Bustos FJ, et al. ALS-linked protein
493 disulfide isomerase variants cause motor dysfunction. EMBO J. 2016;35(8):845-865.
494
495 45. Agyemang AF, Harrison SR, Siegel RM, McDermott MF. Protein misfolding and dysregulated
496 protein homeostasis in autoinflammatory diseases and beyond. Semin Immunopathol.
497 2015;37(4):335-347.
498
499 46. Araki E, Oyadomari S, Mori M. Endoplasmic reticulum stress and diabetes mellitus. Intern
500 Med. 2003;42(1):7-14.
501
502 47. Nishi M, Nanjo K. Insulin gene mutations and diabetes. J Diabetes Investig. 2011;2(2):92-100.
503
504 48. Gorbatyuk MS, Knox T, LaVail MM, Gorbatyuk OS, Noorwez SM, Hauswirth WW, et al.
505 Restoration of visual function in P23H rhodopsin transgenic rats by gene delivery of BiP/Grp78.
506 Proc Natl Acad Sci U S A. 2010;107(13):5961-5966.
507
508 49. Coelho DS, Cairrao F, Zeng X, Pires E, Coelho AV, Ron D, et al. Xbp1-independent Ire1
509 signaling is required for photoreceptor differentiation and rhabdomere morphogenesis in
510 Drosophila. Cell Rep. 2013;5(3):791-801.
511
512 50. Athanasiou D, Aguila M, Bellingham J, Kanuga N, Adamson P, Cheetham ME. The role of the
513 ER stress-response protein PERK in rhodopsin retinitis pigmentosa. Hum Mol Genet.
514 2017;26(24):4896-4905.
515
516

Figure 1. A SNP in *Baldspot* is associated with degeneration in a model of retinitis pigmentosa. A. *Baldspot* is the only *Drosophila* orthologue of mammalian *ELOVL6* (*). Phylogeny of long chain fatty acid elongases shows the relationship between *Drosophila* (black) and human (red) genes. **B.** The 3L:16644000SNP in *Baldspot* (BDGP R5/dm3) is associated with *Rh1^{G69D}*–induced degeneration in the DGRP. *Rh1^{G69D}* DGRP eye size is plotted by allele identity. Strains carrying the minor C allele (18880 ± 3010 pixels) have significantly smaller eyes than those carrying the major, T allele (21822 ± 2294 pixels). **C.** Images from *Rh1^{G69D}* DGRP strains that carry either the T or the C allele. Images were chosen from strains representing the range of eye sizes for the two alleles. All data from B and C were taken from Chow et.al 2016. ** $P < 0.005$

Figure 2. Loss of *Baldspot* in the *Rh1^{G69D}* eye disc alleviates ER stress and apoptosis. A. Degeneration caused by overexpression of *Rh1^{G69D}* is partially rescued by RNAi knockdown of *Baldspot* (*Rh1^{G69D}/Baldspoti* vs *Rh1^{G69D}* controls). **B.** This rescue was significant when eye size was quantified, as indicated by a larger eye in *Rh1^{G69D}/Baldspoti* flies (14043 ± 960 pixels vs. controls 11942 ± 473 pixels). **C.** Loss of *Baldspot* had no effect on eye phenotype in an otherwise wild-type background. **D.** *Rh1^{G69D}/Baldspoti* eye discs also display reduced apoptosis levels compared to *Rh1^{G69D}* controls, as measured by TUNEL staining. **E.** *Rh1^{G69D}/Baldspoti* eye discs had reduced expression of Xbp1-EGFP as compared to *Rh1^{G69D}* controls, while rhodopsin levels were unchanged. Eye discs were dissected from wandering L3 larvae expressing *Rh1^{G69D}* and *UAS-Xbp1-EGFP*, stained for rhodopsin and GFP as an indicator of ER stress, and counterstained with 4',6-diamidino-2-phenylindole (DAPI). F. Xbp1-EGFP levels are significantly reduced in *Rh1^{G69D}/Baldspoti* eye discs (3.8 ± 0.8 fluorescent units) as compared to *Rh1^{G69D}* controls (6.5 ± 1.1 fluorescent units). Fluorescence was quantified using mean fluorescence from the image stack as calculated in Image J. Scale bars = 0.1 mm. *** $P < 0.0005$, **** $P < 0.0001$.

Figure 3. Stearate supplementation increases degeneration and activation of ER stress pathways in *Rh1^{G69D}/Baldspoti* flies. A. Eye size was significantly reduced in *Rh1^{G69D}/Baldspoti* flies (*Baldspot* RNAi) when they were fed 10% stearate (14033 ± 1553 pixels) as compared to standard media (16846 ± 1699 pixels). Eye size was not significantly changed in *Rh1^{G69D}* control flies (No RNAi) fed 10% stearate (13479 ± 1197 pixels) as compared to the standard media (12873 ± 1006 pixels). **B.** The 10% stearate diet also increased Xbp1-EGFP levels in eye discs from *Rh1^{G69D}/Baldspoti* flies to a similar level seen in *Rh1^{G69D}* controls. Scale bars = 0.1 mm. **** P < 0.0001.

Figure 4. *Baldspot* is a general modifier of the ER stress response. A. Loss of *Baldspot* partially rescues the adult vestigial wing phenotype caused by expression of *Rh1^{G69D}* in the wing disc (wing-*Rh1^{G69D}*/Baldspoti vs wing-*Rh1^{G69D}* controls). **B**. Wing discs from wing-*Rh1^{G69D}*/Baldspoti flies had lower Xbp1-EGFP expression, as compared to wing discs from wing-*Rh1^{G69D}* controls. Wing discs were dissected from wandering L3 larvae expressing *Rh1^{G69D}* and *UAS-Xbp1-EGFP* and stained for GFP and DAPI as a counterstain. The wing disc is outlined in red in the merged image. **C.** Larvae with ubiquitous knockdown of *Baldspot* are significantly more resistant to Tunicamycin-induced ER stress than control larvae. L2 larvae were treated with tunicamycin or DMSO and survival to pupation was calculated as a measure of resistance to ER stress. Five representative experiments are shown. Scale bars = 0.1 mm. ** P < 0.005.

Figure 5. Baldspot regulates *Xbp1* splicing but not RIDD activity of IRE1. A. S2 Cells treated with DsRNA targeted against *Baldspot* had near complete reduction in *Baldspot* transcript levels (7.8%) as compared to control cells treated with DsRNA against *EGFP*. **B.** DTT treatment increased *Xbp1* splicing in control cells (2.96-fold increase), while splicing was reduced in cells treated with DsRNA against *Baldspot* (2.12-fold increase). **C.** Levels of the RIDD target *sparc* are reduced in control cells treated with DTT when compared to DMSO-treated controls (~81% of controls). This reduction is unaffected by treatment with *Baldspot* DsRNA. *Actin5c* transcript levels were unaffected by DTT treatment. * P < 0.05, **** P < 0.0001.