Learning multiple concurrent higher-order visual sequences in human primary visual cortex

Clive R. Rosenthal[1], Indira Mallik[2], Cesar Caballero-Gaudes[3],
Martin I. Sereno[4], and David Soto[3,5]

[1] *Nuffield Department of Clinical Neurosciences, University of Oxford, UK*
[2] *Division of Brain Sciences, Imperial College London, UK*
[3] *Basque Center on Cognition, Brain and Language, San Sebastian, Spain*
[4] *Department of Psychology, San Diego State University, USA*
[5] *Ikerbasque, Basque Foundation for Science, Bilbao, Spain*

February 21, 2018

Correspondence to: d.soto@bcbl.eu, Basque Center on Cognition, Brain and Language, Paseo Mikeletegi 69, 2nd Floor 20009 San Sebastian

Abstract

Learning and memory are supported by a network involving the medial temporal lobe and linked neocortical regions. Emerging evidence indicates that primary sensory regions (i.e., V1) may contribute to recognition memory, but this has been only tested only with a single visuospatial sequence as the target memorandum. The present study used functional magnetic resonance imaging to test the capacity of human V1 to support the learning of multiple, concurrent and complex visual sequences involving discontinous (second-order) associations. Two peripheral, task-irrelevant but structured sequences of orientated gratings appeared simultaneously in right and left visual fields alongside a central, task-relevant sequence that was in the focus of spatial attention. Pseudorandom sequences were introduced at multiple intervals during the presentation of the three structured visual sequences to provide an online measure of sequence-specific knowledge at each retinotopic location. We found that a network involving the precuneus and V1 was involved in learning the structured sequence presented at central fixation, whereas right V1 was modulated by repeated exposure to the concurrent structured sequence presented in the left visual field. The same result was not found in left V1. These results indicate for the first time that human V1 can support the learning of multiple concurrent sequences involving complex discontinuous inter-item associations, even those that are goal-irrelevant.

1 Introduction

Primary visual cortex (V1) is typically thought to carry out primitive low-level visual computations and perceptual learning of low-level visual features (Sasaki, Nanez, & Watanabe, 2010), but there is now emerging evidence that V1 can play a role in higher-order functions such as the learning and recognition of visuo-spatial sequences in humans (Rosenthal, Andrews, Antoniades, Kennard, & Soto, 2016), and in model organisms (Cooke, Komorowski, Kaplan, Gavornik, & Bear, 2015; Gavornik & Bear, 2014). Functional coupling between V1 and putative memory substrates has been identified during non-conscious recognition (Rosenthal et al., 2016), incidental statistical learning of visible items (Turk-Browne, Scholl, Johnson, & Chun, 2010), conscious recall of tone-grating pairs (Bosch, Jehee, Fernández, & Doeller, 2014) and cue-action associations (Hindy, Ng, & Turk-Browne, 2016). However, the capacity limitations of V1 activity during the learning of complex memoranda remain to be systematically evaluated.

The aim of the present functional magnetic resonance imaging (fMRI) study was to assess the capacity of human V1 to support concurrent learning and recognition memory of three complex visual sequences presented at attended (task-relevant) locations and unattended (task-irrelevant) peripheral regions of the visual field subtended by V1. Prior research has implicated V1 in learning and memory for single or simple visual sequences (Rosenthal et al., 2016; Cooke et al., 2015; Gavornik & Bear, 2014). For instance, a recent fMRI-study by Rosenthal et al. (2016) showed a role of V1 in learning a single, goal-relevant non-conscious second-order conditional visuospatial sequence of targets specified at four locations. This study however could not address whether multiple concurrent, both goal-relevant and goal-irrelevant sequences can be encoded in V1. Also, it is also unknown whether or not higher-order learning effects for goal-irrelevant sequences occur in retinotopic V1 areas.

The present study tackled these questions. In particular, the current study examined he capacity of human V1 for learning multiple sequences based on discontinuous associations of visual events was tested at three different positions in the visual field, both task-relevant and task-irrelevant sequences at fixed retinotopic locations. Manipulating the relevance of the sequences for the task goal of the observer may be critical to characterize the role of V1 in higher-order learning. This is because activity in V1 may reflect the learning computation in question or V1 may be driven by strategic factors, such as the orientation of attention based on goals(Li, Piëch, & Gilbert, 2004; Gandhi, Heeger, & Boynton, 1999; Liu, Larsson, & Carrasco, 2007). Modulations in blood oxygen-level dependen (BOLD) signal change for structured against random sequences presented in peripheral locations that are irrelevant for current goals would indicate that learning-related modulations in V1 can occur outside of strategic factors (e.g.) related to attentional orientation.

We thus assessed the brain responses in V1 associated with the repeated presentation of a visual sequence presented at fixation and also tested whether right and left V1 exhibited learning-related activity associated with the repeated pre-

sentation of the peripheral, task-irrelevant, but structured, higher order visual sequences in its corresponding contralateral visual receptive fields. Evidence of V1 modulation in either unattended location would be consistent with a model of V1 at the interface between perception and memory functions, which can operate independently of goal-directed attention control factors.

2 Methods

2.1 Participants

Seventeen undergraduate students were recruited (mean age of 22 years; nine female). All participants gave informed written consent to take part in accordance with the terms of approval granted by the local research ethics committee and the principles expressed in the Declaration of Helsinki, and were naive to the purpose of the experiment. All participants reported normal or corrected-to-normal vision, and had no history of neurological disease.

2.2 Experimental procedure

Participants underwent fMRI scanning as they viewed a visual display containing three different visual sequences. The sequence presented in the center of the screen involved changes in the color of a circle (white, light grey, dark grey and black). The sequences presented in the left and right visual fields involved changes in the orientation of a grating (i.e., 0, 45, 90 or 135 degrees from the vertical). Each sequence was comprised of 12 stimuli following a second-order conditional (SOC) rule (Reed & Johnson, 1994), whereby at the lowest structural level the current stimulus is contingent on the two preceding stimulus. The sequences used were as follows: SOC1: 3 4 2 3 1 2 1 4 3 2 4 1; SOC2: 3 4 1 2 4 3 1 4 2 1 3 2; SOC3: 2 4 2 1 3 4 1 2 3 1 4 3, where 1 2 3 4 represents a stimulus type).

Participants were instructed to discriminate the current color of the central sequence by pressing one of four buttons. The central sequence was goal-relevant and hence in the focus of attention. The peripheral sequences were designated task-irrelevant by instructing participant to ignore the changes in orientations of the gratings and only respond to the colour changes in the central sequence and not to respond to the orientations of these peripheral sequences, which were therefore goal-irrelevant. The phase of the gratings was varied independently of the sequences of orientations, and varied randomly across trials to preclude neural adaptation effects of the phase dimension. The three SOC sequences at each of the three locations were presented on a loop, but were asynchronously interspersed with the presentation of the 12 element pseudorandom sequence (i.e., containing no immediate repetitions of a colour or grating orientation). Pseudorandom sequences were always repeated 4 times, hence comprising 48 stimulus presentations or trials. Stimuli at all three locations were each presented for 1 second with an interval of 0.25 seconds between each stimulus. The

pseudorandom sequence appeared twice per run at different times for each of the 3 sequences. The temporal position within a run of each of the pseudorandom sequences was consistent across participants. The Table below illustrates the structure of each of the 3 training runs. There were 672 trials on each of the three training runs and each lasted for 14 minutes. The experimental task is depicted in Figure 1.

Figure 1: Illustration of the sequence training protocol inside the MRI scanner. The right panel indicates the structure of each of the 3 training runs (S: for the visual second-order conditional sequence; R: for pseudorandom blocks, comprised of a 48 trials each). The top row corresponds to the left sequence, the middle to the central sequence and the bottom row within each run corresponds to the right sequence. Note that each of the pseudorandom sequences were presented asynchronously across the three sequences and scanning runs.

2.3 Recognition test outside of the scanner

After training, the participants performed three recognition tests outside the scanner, one for each of the three 12-element second-order conditional sequences presented at central fixation and left/right positions of the visual field and counterbalanced across participants. On each trial of the recognition test, participants were shown either an old or a new sequence which served as a recognition cue (six-element sequences; 12 old and 12 new). Old sequences were structured sequences that were presented during the training phase inside the scanner and were presented in the same way as during the training protocol. New sequences were also based on a SOC rule, but had not been presented to the participant during the training phase. The participant was asked to indicate whether each of the sequences presented on each trial was 'old' or 'new'. Then, participants indicated the confidence in their response (from 1-guess; 2-fairly confident; 3-sure). There were no restrictions on the participants when viewing the central sequences and hence they were allowed to simulate the motor responses given

during training.

Receiver-operating-characteristic (ROC) analyses were used to derive measures of type-1 sensitivity (i.e., ability to distinguish old/new sequences) and type-2 sensitivity (i.e. how confidence relates to memory accuracy). For the type-1 analyses, the 'signal' and 'noise' were defined as old and new retrieval cues, respectively. A 'hit' was, therefore, a correct response ('old') to an old trained (six-element sequence) retrieval cue and a 'correct rejection' was a correct response ('new') to a new untrained (six-element sequence) retrieval cue. A 'false alarm' was an incorrect response ('old') to a new retrieval cue and a 'miss' was an incorrect response ('new') to an old retrieval. To obtain the type-1 ROC curves, we plotted the probability of hits as a function of the probability of false alarms for all possible decision criteria. The different points in the ROC curve were obtained by calculating cumulative probabilities for hits and false alarms along a recoded confidence continuum ranging from 3 (i.e., certain) to 1 (i.e.. least certain) in signal present trials (old retrieval cues), and from 3 (i.e., certain) to 1 (i.e., least certain) in noise trials (new retrieval cues). On the basis of simple geometry, we computed the area under the ROC curve as a distribution-free measure of the discriminability. For the type-2 sensitivity analyses, the area under the ROC curve was calculated by plotting the cumulative probability of correct responses (either hit or correct rejection) and the cumulative probability of incorrect responses (either false alarm or miss) across the different levels of confidence. The area under the ROC curve was estimated as distribution-free measure of metacognitive ability (Fleming, Weil, Nagy, Dolan, & Rees, 2010; Kornbrot, 2006). Data processing and analyses were performed in R (version 3.2.2).

2.4   MRI data acquisition

MRI scanning was performed in a Siemens Avanto 1.5T MRI scanner using a receive-only 32-channel head coil and body transmit. A screen was mounted at the end of the scanner bore and visual stimuliwere projected onto it from the console room through a wave guide. Subjects viewed the screen using a 45 deg mirror. Functional volumes for both the training phase, functional localiser, and retinotopy consisted of multi-slice T2*-weighted echoplanar images (EPI) with blood oxygenation level dependent (BOLD) contrast with a multiband acceleration factor of 4. We used the following scanning parameters to achieve whole brain coverage: TR=1000 ms, TE=54.8 ms, 40 coronal slices, 3.2 mm slice thickness, no interslice gap, and FoV=205x205 mm (3.2 x 3.2 mm in-plane voxels). There were 844 whole-brain scans per training run. To facilitate anatomical localization and cross-participant alignment, a high-resolution 1x1x1 mm whole-brain structural T1-weighted, magnetization prepared rapid gradient echo (MP-RAGE) scan was acquired for each participant (FOV=256x224 mm, 176 partitions, TR=8.4 ms, TE=3.57 ms, TI=1000 ms, inversion spacing=2730 ms, BW=190 Hz/pixel).

2.5   Functional MRI preprocessing

We used FEAT (fMRI Expert Analysis Tool) Version 6.0, as part of FSL (www.fmrib.ox.ac.uk/fsl). The first 10 EPI volumes were removed to account for T1 equilibrium effects. Non-brain removal was performed using Brain Extraction Tool. Volume realigment of functional scans was carried out using FMRIB's Linear Image Registration Tool MCFLIRT. Scans were realigned relative to the middle scan. We applied a 50 s high-pass temporal filtering to remove low frequency noise, and spatial smoothing using a FWHM Gaussian kernel of 6 mm.

2.6   Retinotopic mapping

Retinotopic mapping was performed to identify right and left human V1 in each individual. We used standard phase-encoded polar angle retinotopic mapping (Sereno et al., 1995). Participants fixated centrally while viewing a clockwise or counterclockwise rotating pie-shaped 'wedge' with a polar eccentricity. The duration of the scan was 8 min 32 secs (512 volumes). Fourier methods were applied to obtain polar angle maps of the cortex and delineate the borders of right and left V1.

2.7   Functional localizer

A functional localiser scan was conducted in all participants and involved the presentation of items that were flashed at the same three different locations. A block design was used. At each location, there were 24 presentations of gratings (250 ms each with a 250 ms inter-grating interval) in pseudorandom orientations. Following each block, there was a rest period of 12 seconds. The phase of the gratings also varied randomly across trials. We obtained contrast of parameter estimates of brain responses to the spatial position of the peripheral sequences (left>right and right>left positions).

   The parameter estimates from the localiser were used to define another mask comprising the voxels in the V1 retinotopic masks that responded to the position of the gratings during the training phase (e.g., define the voxels within the right V1 mask that showed a response to the left-side position of the gratings and likewise for left V1). This area in V1 was delineated by finding the intersection between the V1 retinotopy and the functional localiser maps using fslmaths -mas tool. Hence, we isolated the area of V1 that was most selective of the location of the right and left sequences. We initially derived masks comprising the 10 most responsive voxels and repeated the procedure considering the 5, 20 and 40 most responsive voxels to further assess the reliability of the results.

2.8   MRI Statistical Analyses

Time-series statistical analyses were conducted using FILM (FMRIB's Improved Linear Model) with local autocorrelation correction. The data were analyzed

using voxelwise time series analysis within the framework of the general linear model. A design matrix was generated in which each individual presentation of a grating was modelled with a double-gamma hemodynamic response function. Explanatory regressors were created for each of the structured and pseudorandom sequences. Standard motion realignment parameters were included in each individual subject's general linear model as nuisance regressors. We included further nuisance regressors for those volumes corresponding to motion outliers using the FSL motion outliers function. The root mean squared (RSM) head position difference to the reference volume was used as metric to detect fMRI timepoints corrupted by large motion. The threshold was set to the FSL default, namely, the 75 percentile + 1.5 inter quartile range of the distribution of RSM values of each run. A confound matrix was generated and used in the GLM to completely remove the effects of these timepoints on the analysis. This is intended to deal with the effects of intermediate to large motions, which corrupt images beyond anything that the linear motion parameter regression methods can fix.

Sequence blocks were modeled in chunks of 48 trials in order to contrast activity between structured and pseudorandom baseline blocks (S and R in Figure 1); 48 trials were selected as a multiple of 12-element unit of SOC sequence that allowed the number of trials on structured sequence blocks to be equated with the pseudorandom blocks, as per our prior study (Rosenthal et al., 2016).

For the analysis of the central sequence, we first derived contrast of parameter estimates between the structured relative to the pseudorandom sequences (S<R) within each run. S<R contrast of parameter estimates were obtained as follows. First, we compared the first central pseudorandom sequence with the preceding structured central sequences that were not flanked by random sequences in the right and left visual fields (e.g. run 3, see Figure 1). Therefore, the two structured blocks preceding the first central pseudorandom were not considered in the S<R contrast. Instead, we selected the preceding structured block (i.e., the third one preceding the first central pseudorandom sequence, see Figure 1). This was done to preclude the possibility that BOLD responses to the critical central structured sequence were counfounded by hemodynamic responses changes from the peripheral pseudorandom sequence presented concurrently. Secondly, we performed the same analyses by comparing the structured sequence block immediately preceding a pseudorandom sequence block at similar timepoint across the runs. The results obtained from these two analysis strategies were similar.

Following the computation of within-run contrast of parameter estimates (S<R), we performed across-run within-subject (fixed-effects) analyses using lower level parameter estimates to test for a linear training effects across the 3 runs (e.g., -1 0 1) and likewise exponential training effects (e.g., -2 -0.5 2.5). Finally, the parameter estimates from these linear and exponential contrasts across the three learning runs were converted to standard MNI 2 mm template with linear transformation and passed to a higher-level across subjects one-sample t-test to assess brain regions consistent across participants that were associated with increased training effects in the structured relative to the pseudorandom

sequence using FLAME (Local Analysis of Mixed Effect). Statistical maps were thresholded using clusters determined by Z > 2.3 and a corrected cluster extent significance threshold of p=0.05, using Gaussian Random Field Theory. Across participants one-sample t-tests were also performed separately on each of the 3 runs to assess S<R brain actitivy differences on each run. These considerations applied also to the psychophysiological interaction (PPI) analyses described below.

We also used non-parametric permutation-based one-sample t-tests to further assess the statistical significance of the results. The FSL Randomise program was used (Winkler, Ridgway, Webster, Smith, & Nichols, 2014) with threshold-free cluster enhancement (TFCE). TFCE is a method that enhances values within cluster-like structures, while preserving voxel-level inference (Smith & Nichols, 2009). We performd 6 mm spatial smoothing of the variance which is typically included to increase study power.

For the analysis of the peripheral right and left sequence, we adopted the following approach. Parameter estimates were derived in native functional space for each structured and pseudorandom sequence block (Figure 1) for each training run and participant. We then used the V1 masks obtained from the retinotopy and the functional localiser to extract the parameter estimates of V1 activity across the training phase, separately for the right and left V1 masks. For each run, we then performed paired-sample t-tests comparing V1 activity estimates in the first pseudorandom sequence and the inmediately preceding structured sequence.

2.9   Psychophysiological interaction (PPI) analyses

A seed-voxel PPI based approach was used to identify signals of interest that would be missed in standard subtraction based analyses. A 6 mm radius posterior cingulate/precuneus mask was drawn and based on the peak voxels of the posterior cingulate/precuneus that showed effects of training. This mask was used to define the seed region's time course for the PPI analyses. The aim was to assess whether the temporal correlation between posterior precuneus and primary visual cortex was modulated during the learning of the task-relevant, second-order conditional sequence.

We then estimated a model for each participant that included the same psychological regressors as outlined above, for the onsets of each sequence, and, critically, for the PPI model, a physiological regressor for the time course of the region-of-interest, and psychological x physiological interaction regressors for the PPI. These new regressors were, therefore, added to the previous first-level model for each participant/run. Parameter estimates for structured versus pseudorandom conditions based on the PPI regressors were derived in a similar fashion to the analyses outlined above (i.e., using both fixed-effects analysis across runs followed by mixed-effect analysis across participants), which, here, directly compared the changes in functional coupling associated with training. Given our a priori interests in V1(Rosenthal et al., 2016), the higher-level analyses (across subjects) were performed using a region-of-interest approach, with

a target occipital mask in standard space derived from the FSL Harvard-Oxford Atlas.

3 Results

3.1 Behavioural performance during training

We assessed the response time (RT) and the accuracy of manual responses to each item color of the central sequence as a function of the nature of the sequence (structured vs pseudorandom). Overall, the results indicate that the repeated exposure to the deterministic discontinuous associations of the structured sequences affected somatomotor performance relative to each novel pseudorandom sequence. We conducted an ANOVA on the manual RTs with run(1,2,3) and sequence type (S,R) as factors. We compared the first R sequence and the preceding S sequence of each run. There was no main effect of run on RTs ($F(2,32)=1.95$, $p=0.159$). The main effect of sequence type was significant ($F(2,32)=11.67$, $p=0.004$), with slower performance on R relative to S sequence; however this effect did not interact with run. ($F(2,32=0.607$, $p=0.551$). The accuracy data showed a consonant pattern with the RTs. There was no main effect of run on accuracy ($F(2,32)=0.177$, $p=0.838$). The main effect of sequence type was significant ($F(2,32)=14.913$, $p=0.001$), with impaired accuracy on R relative to S sequence blocks; however this effect did not interact with run ($F(2,32=2.287$, $p=0.118$). Figure 2 illustrates these results. Taken together, these results indicate that the statistical structure of the central goal-relevant sequence was learned.

Figure 2: Behavioural results of the learning phase collapsed across runs (A) Mean proportion of correct responses in S and R blocks (B) Mean RTs in S and R conditions.

3.2 Recognition performance outside the scanner

Following the fMRI training task, a surprise recognition test was administered to test the memory of participants for each of the 3 sequences. To summarise the main results, participants memory confidence ratings discriminated whether the

central (goal-relevant) sequence retrieval cues were 'old' (repeateadly studied during the training runs) or 'new', but there was no evidence of significant recognition memory for the unattended left and right sequences.

To derive sensitivity measures of recognition performance we computed the area under the type-1 ROC for 'old'/'new' discrimination. For the sequence presented at central fixation, memory sensitivity was marginally above chance ($t(16)$=1.882, p=0.078, mean=0.565). Type-2 sensitivity performance, namely, the extent to which confidence ratings predicted the accuracy of the recognition response showed a similar trend ($t(16)$=1.911, p=0.074, mean=0.544).

However, additional analyses of recognition performance based on memory confidence evinced that old and new probes were processed differently. Notably, participants were more confident in their responses when judging old sequences relative to new sequences ($t(16)$=2.2706, p=0.036, mean confidence old=1.926; mean new=1.784), indicating that confidence was diagnostic of the memory status (old/new) of the sequences. The memory sensitivity and confidence results are presented in Figure 3.

Figure 3: Behavioural results of the recognition test outside the scanner (A) Type-1 and type-2 memory sensitivity to old and new sequences (B) Memory confidence for old and new sequences.

Further results presented next indicated that participants were unable to recognise either the left or the right sequences.

Analyses of the recognition test performance for the left sequence showed that the area under the type-1 and type-2 ROC was no different from chance ($t(16)$=0.543, p=0.594, mean= 0.484 for type-1 ROC, and $t(16)$=0.352, p=0.729, mean=0.492 for type-2 ROC performance). Moreover, the results showed that participants were not more confident in responding to old vs. new left sequences ($t(16)$=1.0371, p=0.31, mean confidence old=1.593; mean new=1.647).

There was also no evidence of recognition memory for the right sequence. Type-1 and type-2 performance were no different from chance ($t(16)$=1.075, p=0.298, mean 0.483, and $t(16)$=0.101, p=0.92, mean=0.498). Participants confidence ratings did not discriminate between old and new left sequences ($t(16)$=1.165, p=0.261, mean confidence old=1.598; mean new=1.544).

3.3 fMRI Results

A key question of the present study is whether the V1 is involved in supporting the learning of complex deterministic discountinous associations in the structured peripheral sequences of oriented gratings (unattended, task-irrelevant) that participants were unable to recognise in a subsequent test. The behavioural results showed that participants could only learn and recognise the central goal-relevant sequence of color changes.

For the analysis of the central sequence, we computed statistical contrasts of parameter estimates between structured (S) and pseudorandom (R) sequences (i.e., comparing S<R) on each of the three training runs (see Methods). This S<R contrast predicts increased response to the presentation of (novel) pseudorandom sequences relative to the repeated representation of the structured sequence and this contrast was motivated by our prior fMRI perceptual sequence learning study (Rosenthal et al., 2016), and is in keeping with a plethora of evidence for novelty effects in memory paradigms which are reflected by increases in hemodynamic responses (Ranganath & Ritchey, 2012).

First, BOLD responses to the first R central sequence were compared to a preceding structured sequence that was not flanked by pseudorandom sequences (see Methods) in order to avoid that BOLD responses to the critical central structured sequence were not counfounded by hemodynamic responses changes from the peripheral pseudorandom sequences presented concurrently. We then tested for linear and exponential effects of training on BOLD responses (e.g., increased differential S<R response in run 3 relative to run 1). Learning the structure of the repeated sequence ought to be reflected in brain activity changes across the training runs whenever a non-structured, pseudorandom sequence was presented.

The analyses showed significant clusters in the posterior cingulate gyrus (MNI 0 -48 16, Z=4.26) extending into the retrosplenial cortex, and the posterior precuneus in the the vicinity of intracalcarine sulcus, the ventromedial prefrontal cortex (MNI, 2 64 2, Z=3.41). All of these regions displayed significant linear and exponential changes in activity across runs (Z > 2.3. p < 0.05 whole brain corrected using cluster-based random-field theory with family wise correction).

Secondly, we performed the same analyses by comparing the structured sequence block immediately preceding a pseudorandom sequence block at similar timepoint across the runs. The results obtained from this second analysis were similar to the first except that the ventromedial prefrontal cortex did not survive the threshold for statistical significance. Activity was restricted to the posterior cingulate and posterior precuneus.

To examine further the effects of training, we performed group-analyses separately for each of the three runs. On the basis of prior studies of learning SOC sequences (Schendan, Searl, Melrose, & Stern, 2003; Albouy et al., 2008), training effects were predicted to be most evident on the later third training run. Differences between the structured and the pseudorandom sequence blocks in run 1 and in run 2 did not survive the above significance threshold. Significant differences in the posterior cingulate, precuneus and ventromedial PFC emerged

only in run 3, and were evident in both linear and exponential learning contrasts ($p < 0.05$ whole brain corrected and also following non-parametric tests using threshold-free cluster enhacement).

Following prior reports that putative mnemonic regions (e.g. hippocampus) show increased temporal correlation with V1 during memory recall (Bosch et al., 2014)and implicit statistical learning of visible stimuli (Turk-Browne et al., 2010), we conducted similar analyses considering the posterior precuneus cluster that was observed during learning, as this region is also a key node of the memory network (Ranganath & Ritchey, 2012). This region has connections to higher level visual areas (Margulies et al., 2009) and thus has access to V1 via feedback projections.

The psychophysiological interaction (PPI) analysis used a 6 mm seed mask centered on the posterior precuneus area identified in the learning contrast (MNI 0 -48 12) at the cluster peak in the S<R contrast in run 3 reported above. This was used to define the seed region's time course for the PPI. Given our a priori interest in primary visual cortex, training effects across participants were assessed within a standardized occipital mask as target of the PPI.

We found a cluster of activity in V1 peaking at the intracalcarine sulcus (MNI: 8 -92 0, Z=3.34) showing linear and exponential increases in temporal correlation with posterior precuneus across the training runs on structured relative to the pseudorandom sequences (see figure 4). This result was observed in the comparison between first pseudorandom sequence block and the structure sequence immediately preceding it. This suggests that learning a representation of the structured sequence involves the coupling between putative memory substrates and V1. For the position of anatomically defined V1 in the calcarine, see (Horton & Hoyt, 1991); also, the activity in the intracalcarine cortex and its vicinity coincides with probabilistic V1.

Figure 4: Brain regions in the precuneus and ventromedial prefrontal cortex exhibited linear changes in BOLD signal for the structured<pseudorandom sequence contrast across the three training runs (in yellow, $Z > 2.3$, $p < 0.05$, whole-brain corrected). The green areas in primary visual cortex indicate the results of the PPI analysis from the posterior precuneus cluster ($Z > 2.3$, $p < 0.05$, corrected for the occipital mask in blue shading).

Right and left V1 were mapped using independent retinotopy and functional

localiser scans of each participant (see Methods). On the basis that learning effects (S<R) for the central structured sequence were expressed in the last training run, it was predicted that learning of the peripheral sequences would be also evident in run 3.

Accordingly, the BOLD signal difference between structured and pseudo-random sequences (S<R)in right V1 in the last training period (run 3) was significantly different from chance (Figure 5; $t(16)=2.480$, $p=0.0246$, two-tailed paired t-test). This result was qualified by a repeated measures ANOVA with run (1, 2, 3) as factor that tested for linear and quadratic trends in the learning effect (S<R for the left sequence) across the runs. There was a significant quadratic trend in the learning effect for right V1 showing that learning effects were expressed in run3 ($F(1,16)=6.32$, $p<0.023$, Figure 5). The linear trend was not significant ($F(1,16)=1.39$, $p<0.25$). This pattern of results is consistent with the non-linear learning effects observed for the central sequence.

These results were consistent across a wide range of V1 masks. In particular, we initially tested our hypothesis with masks that comprised the 10 most responsive voxels to the location of the left sequence, as shown by the combination of retinotopy and the functional localiser, and then verified that the result held with masks containing 5, 20 and 40 most responsive voxels. Figure 5 illustrates the representative results from the 10 most responsive voxels.

Figure 5: Right V1 responses as a function of the structured (S) or pseudorandom (R) status of the sequences on each of the three training runs. The red * illustrates significant BOLD signal change between structured and pseudorandom sequences in run 3 (see text).

For left V1, there was no evidence of learning the structure of the deterministic discontinuous associations of the structured sequence in the predicted last training run ($t(16)=0.66$, $p=0.518$). Neither the linear nor the quadratic learning trends across runs were observed ($F(1,16)=.422$, $p<0.525$; ($F(1,16)=.829$, $p<0.376$).

Figure 6 below illustrates these results.

Figure 6: Left V1 responses as a function of the structured (S) or pseudorandom (R) status of the sequences on each of the three training runs. There was no evidence of BOLD signal differences between structured and pseudorandom sequences in any of the runs (see text).

4   Discussion

The results indicate that V1 operates as part of a network involving the precuneus during the learning of task-relevant visual sequences involving discontinuous associations and which are within the focus of spatial attention. The precuneus, posterior cingulate and ventromedial prefrontal cortex have been identified as part of a memory network (Ranganath & Ritchey, 2012) which can be modulated by the temporal structure of goal-relevant memoranda (Hasson, Chen, & Honey, 2015), with higher-order regions exhibiting sensitivity to increasingly longer temporal structures, ranging from tens of seconds (e.g. written sentences) in more posterior areas to minutes (e.g., a movie).

This cluster of learning related activity in the posterior cingulate/precuneus also extended into the retrosplenial cortex. This area is activated by spatial cognition tasks (Epstein, 2008), episodic memory demands (Vann, Aggleton, & Maguire, 2009) and it has been also heavily implicated in both spatial and object-based forms of contextual/predictive processing and associative learning in scene perception, alongside the ventromedial prefrontal cortex (Bar & Aminoff, 2003; Bar, 2004). V1 coupling with posterior precuneus cortex during repeated exposure to the structured sequences is consonant with a mnemonic feedback signal from higher-order brain regions involved in contextual/predictive processing (Rao & Ballard, 1999; Alink, Schwiedrzik, Kohler, Singer, & Muckli, 2010).

In addition to the contribution of V1 in encoding the central attended sequence, the results revealed that retinotopic V1 areas can also support learning of the statistical regularities associated with goal-irrelevant items presented at non-attended spatial positions. Note that performance in the central attention task based on 4-color discrimination had to be carried out at a fast pace and performance was not at ceiling (i.e. around 0.8 proportion correct). This means that the central task was challenging enough to keep the focus of the observers' attention engaged, and thus observers were very unlikely to spare attentional resources to process the peripheral 12-element sequences. This is

also indicated by the fact that observers performed at chance in the subsequent recognition test. However there was evidence both from the manual response times and the accuracy rates that observers learned the structure of the repeating second-order conditional sequences, as performance was impaired by the introduction of pseudorandom sequences, and there was evidence of recognition in the subsequent memory test performed outside the scanner. We also note that the behavioural expression of the learning effect both in manual RTs and response accuracy followed a different time course to the learning effects found in the brain responses. The former did not significantly increase across the training runs, while learning effects on brain response were most evident in the third training run. It is worth noting that sequence learning effects may reflect a combination of response-response, stimulus-response and response-stimulus bindings (Ziessler & Nattkemper, 2001; Ziessler, 1998) which may have different time courses, however this was not addressed in the present neuroimaging study.

The present results go beyond a recent demonstration that V1 can support the learning of a single second-order conditional sequence (Rosenthal et al., 2016); this study could not determine the capacity of V1 to support multiple co-ocurring sequences and did not address whether learning can occur in retinotopic areas. Furthermore, the learning effects in V1 in Rosenthal et al. were found under conditions in which the elements of the sequence were goal-relevant for a counting test. Thus, the present results show for the first time that V1 can support learning of both goal-relevant and irrelevant visual sequences presented at retinotopic areas.

The results also indicate that spatial relational encoding is not necessary for the concurrent learning second-order conditional sequences in V1, in keeping with the study of Gavornik and Bear (2014) using a more simple and single visual sequence. Rosenthal et al. (2016) study of non-conscious visual sequence learning employed a single second-order conditional visuospatial sequence specified at four locations. Here each of the three second-order conditional sequence were presented at one location. Hence, sequence learning in Rosenthal et al. were characterized by spatial relational encoding, while object-based relational encoding defined the present study. This may be relevant to the presence of basal ganglia activity in Rosenthal et al. and its absence in the present study, since here there is no need for allocentric/egocentric coding of spatial relations, which may be relevant to drive sequence learning effects in the basal ganglia (Kermadi & Joseph, 1995).

Retinotopic activity in right V1 was associated with learning the SOC sequence presented in the left visual field for which right V1 exhibited the greater spatial selectivity. However, there was no evidence of analogous effects in left V1. It is possible that visual competition amongst the three concurrent sequences reduced overall the capacity of left and right V1 areas to encode the multiple co-ocurring sequences, but there is no immediate reason to explain the preferential capacity of right V1 activity to support sequence-specific learning alongside a null sensitivity in left V1. Prior work has suggested the existence of lateralised perceptual biases (Railo, Tallus, & Hamalainen, 2011; Nicholls, Mat-

tingley, & Bradshaw, 2005) in which perceptual processing is biased towards the left side of space. If such biases were operating here that could explain the preferential engagement of right V1 for processing the left sequence. Another possibility to explain the significant learning effect in right V1 and its apparent absence in left V1 may be found on a model of visuospatial hemispheric specialization (Kosslyn, 1987). This model proposes that the right hemisphere supports object-oriented processing, which here it would include the serial dependencies between sequence orientations. On the other hand, the model proposed by Kosslyn (1987) argues that processing in the left hemisphere is more general and less object-oriented. In line with this model, memory-based fMRI studies showed that encoding-related activity during study of visual objects in the right fusiform were associated with later remembering of precise object-based relations, while left fusiform activity during encoding predicted subsequent, non-specific recognition (Garoff, Slotnick, & Schacter, 2005).

The learning effects reported in this study are in keeping with recent reports that V1 can be subject to experience-dependent changes that support higher-order mnemonic processes (Rosenthal et al., 2016; Ekman, Kok, & de Lange, 2017; Cooke et al., 2015; Gavornik & Bear, 2014). Our findings further indicate that V1 processing capacity is flexible enough to be deployed even when attention is constrained by visual competition and task-relevance factors. Top-down processes related to attention (Kastner, Pinsk, De Weerd, Desimone, & Ungerleider, 1999) and predictive processing (Kok, Jehee, & De Lange, 2012) are known to modulate V1. However V1 seems capable of operating autonomously during higher-order processing, so that complex knowledge at goal-irrelevant peripheral locations can nevertheless be encoded. This is consistent with a recent observation that V1 can automatically 'preplay' an expected sequence of events, even when attention resources are constrained (Ekman et al., 2017).

Finally, it is worth noting that the mnemonic feedback signals we have observed in V1 may come from different laminae than feedforward signals; experiments at higher resolution will be needed to make this determination.

5   Acknowledgements

D.S. acknowledges support from the Spanish Ministry of Economy and Competitiveness (MINECO), through the 'Severo Ochoa' Programme for Centres/Units of Excellence in R&D (SEV-2015-490) and project grants PSI2016-76443-P from MINECO and PI-2017-25 from the Basque Government.

References

Albouy, G., Sterpenich, V., Balteau, E., Vandewalle, G., Desseilles, M., Dang-Vu, T., . . . others (2008). Both the hippocampus and striatum are involved in consolidation of motor sequence memory. *Neuron*, *58*(2), 261–272.

Alink, A., Schwiedrzik, C. M., Kohler, A., Singer, W., & Muckli, L. (2010). Stimulus predictability reduces responses in primary visual cortex. *Journal of Neuroscience*, *30*(8), 2960–2966.

Bar, M. (2004). Visual objects in context. *Nature Reviews Neuroscience*, *5*(8), 617.

Bar, M., & Aminoff, E. (2003). Cortical analysis of visual context. *Neuron*, *38*(2), 347–358.

Bosch, S. E., Jehee, J. F., Fernández, G., & Doeller, C. F. (2014). Reinstatement of associative memories in early visual cortex is signaled by the hippocampus. *Journal of Neuroscience*, *34*(22), 7493–7500.

Cooke, S. F., Komorowski, R. W., Kaplan, E. S., Gavornik, J. P., & Bear, M. F. (2015). Visual recognition memory, manifested as long-term habituation, requires synaptic plasticity in v1. *Nature neuroscience*, *18*(2), 262–271.

Ekman, M., Kok, P., & de Lange, F. P. (2017). Time-compressed preplay of anticipated events in human primary visual cortex. *Nature Communications*, *8*.

Epstein, R. A. (2008). Parahippocampal and retrosplenial contributions to human spatial navigation. *Trends in cognitive sciences*, *12*(10), 388–396.

Fleming, S. M., Weil, R. S., Nagy, Z., Dolan, R. J., & Rees, G. (2010). Relating introspective accuracy to individual differences in brain structure. *Science*, *329*(5998), 1541–1543.

Gandhi, S. P., Heeger, D. J., & Boynton, G. M. (1999). Spatial attention affects brain activity in human primary visual cortex. *Proceedings of the National Academy of Sciences*, *96*(6), 3314–3319.

Garoff, R. J., Slotnick, S. D., & Schacter, D. L. (2005). The neural origins of specific and general memory: The role of the fusiform cortex. *Neuropsychologia*, *43*(6), 847–859.

Gavornik, J. P., & Bear, M. F. (2014). Learned spatiotemporal sequence recognition and prediction in primary visual cortex. *Nature neuroscience*, *17*(5), 732–737.

Hasson, U., Chen, J., & Honey, C. J. (2015). Hierarchical process memory: memory as an integral component of information processing. *Trends in cognitive sciences*, *19*(6), 304–313.

Hindy, N. C., Ng, F. Y., & Turk-Browne, N. B. (2016). Linking pattern completion in the hippocampus to predictive coding in visual cortex. *Nature neuroscience*, *19*(5), 665.

Horton, J. C., & Hoyt, W. F. (1991). The representation of the visual field in human striate cortex: a revision of the classic holmes map. *Archives of ophthalmology*, *109*(6), 816–824.

Kastner, S., Pinsk, M. A., De Weerd, P., Desimone, R., & Ungerleider, L. G. (1999). Increased activity in human visual cortex during directed attention in the absence of visual stimulation. *Neuron*, *22*(4), 751–761.

Kermadi, I., & Joseph, J. (1995). Activity in the caudate nucleus of monkey during spatial sequencing. *Journal of Neurophysiology*, *74*(3), 911–933.

Kok, P., Jehee, J. F., & De Lange, F. P. (2012). Less is more: expectation sharpens representations in the primary visual cortex. *Neuron*, *75*(2),

265–270.

Kornbrot, D. E. (2006). Signal detection theory, the approach of choice: Model-based and distribution-free measures and evaluation. *Attention, Perception, & Psychophysics*, *68*(3), 393–414.

Kosslyn, S. M. (1987). Seeing and imagining in the cerebral hemispheres. *Psychological Review*, *94*(2), 148–175.

Li, W., Piëch, V., & Gilbert, C. D. (2004). Perceptual learning and top-down influences in primary visual cortex. *Nature neuroscience*, *7*(6), 651.

Liu, T., Larsson, J., & Carrasco, M. (2007). Feature-based attention modulates orientation-selective responses in human visual cortex. *Neuron*, *55*(2), 313–323.

Margulies, D. S., Vincent, J. L., Kelly, C., Lohmann, G., Uddin, L. Q., Biswal, B. B., . . . Petrides, M. (2009). Precuneus shares intrinsic functional architecture in humans and monkeys. *Proceedings of the National Academy of Sciences*, *106*(47), 20069–20074.

Nicholls, M. E., Mattingley, J. B., & Bradshaw, J. L. (2005, September). The effect of strategy on pseudoneglect for luminance judgements. *Cognitive Brain Research*, *25*(1), 71–77. Retrieved 2017-10-07, from `http://linkinghub.elsevier.com/retrieve/pii/S092664100500114X` doi: 10.1016/j.cogbrainres.2005.04.006

Railo, H., Tallus, J., & Hamalainen, H. (2011). Right visual field advantage for perceived contrast: Correlation with an auditory bias and handedness. *Brain and Cognition*, *77*(3), 391–400.

Ranganath, C., & Ritchey, M. (2012). Two cortical systems for memory-guided behaviour. *Nature reviews. Neuroscience*, *13*(10), 713.

Rao, R. P., & Ballard, D. H. (1999). Predictive coding in the visual cortex: a functional interpretation of some extra-classical receptive-field effects. *Nature neuroscience*, *2*(1).

Reed, J., & Johnson, P. (1994). Assessing implicit learning with indirect tests: Determining what is learned about sequence structure. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *20*(3), 585–594. Retrieved 2017-12-04, from `http://doi.apa.org/getdoi.cfm?doi=10.1037/0278-7393.20.3.585` doi: 10.1037/0278-7393.20.3.585

Rosenthal, C. R., Andrews, S. K., Antoniades, C. A., Kennard, C., & Soto, D. (2016). Learning and recognition of a non-conscious sequence of events in human primary visual cortex. *Current Biology*, *26*(6), 834–841.

Sasaki, Y., Nanez, J. E., & Watanabe, T. (2010). Advances in visual perceptual learning and plasticity. *Nature reviews. Neuroscience*, *11*(1), 53.

Schendan, H. E., Searl, M. M., Melrose, R. J., & Stern, C. E. (2003, March). An fMRI Study of the Role of the Medial Temporal Lobe in Implicit and Explicit Sequence Learning. *Neuron*, *37*(6), 1013–1025. Retrieved 2017-12-04, from `http://linkinghub.elsevier.com/retrieve/pii/S0896627303001235` doi: 10.1016/S0896-6273(03)00123-5

Sereno, M., Dale, A., Reppas, J., Kwong, K., Belliveau, J., Brady, T., & Rosen BR, R., Tootell. (1995). Borders of multiple visual areas in hu-

mans revealed by functional magnetic resonance imaging. *Science*, *268*, 889–93.

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage*, *44*(1), 83–98.

Turk-Browne, N. B., Scholl, B. J., Johnson, M. K., & Chun, M. M. (2010). Implicit perceptual anticipation triggered by statistical learning. *Journal of Neuroscience*, *30*(33), 11177–11187.

Vann, S. D., Aggleton, J. P., & Maguire, E. A. (2009). What does the retrosplenial cortex do? *Nature Reviews Neuroscience*, *10*(11), 792.

Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., & Nichols, T. E. (2014). Permutation inference for the general linear model. *Neuroimage*, *92*, 381–397.

Ziessler, M. (1998). Response–effect learning as a major component of implicit serial learning. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *24*(4), 962.

Ziessler, M., & Nattkemper, D. (2001). Learning of event sequences is based on response-effect learning: further evidence from a serial reaction task. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *27*(3), 595.