Computational elucidation of regulatory network responding to acid stress in *Lactococcus lactis* MG1363

Xin Chen[1,*], Anjun Ma[5,6,*], Hanyuan Zhang[2], Chao Liu[3], Huansheng Cao[4], Qin Ma,[5,6,§]

[1]Center for Applied Mathematics, Tianjin University, Tianjin, 300072, China.

[2]College of computer science and engineering, University of Nebraska Lincoln, NE 68588, USA.

[3]Shandong Provincial Hospital affiliated to Shandong University, Jinan, 250021, China

[4]Center for Fundamental and Applied Microbiomics, Biodesign Institute, Arizona State University, Tempe, Arizona 85287, USA.

[5]Bioinformatics and Mathematical Biosciences Lab, Department of Agronomy, Horticulture and Plant Science, South Dakota State University, Brookings, SD 57006, USA.

[6]Department of Mathematics and Statistics, South Dakota State University, Brookings, SD 57006, USA.

*The authors wish it to be known that, in their opinion, the first two authors should be regarded as Joint First Authors

§To whom correspondence should be addressed. Email: qin.ma@sdstate.edu (QM).

(Words count: 3900, 3 figures, 2 tables.)

ABSTRACT

Acid stress caused by lactate increment can lead to the growth inhibition of bacteria and yes has not been fully defined. Regulons, serve as co-regulated gene groups contribute to the transcriptional regulation of microbe genome, have the potential in understanding the underlying regulatory mechanism. *Lactococcus lactis* is one of the most important Gram-positive lactic acid-producing bacteria, widely used in food industry and has been proved to have advantages in oral delivery of drug and vaccine. In this study, we designed a novel computational pipeline, RECTA, for regulon prediction. The pipeline carried out differentially expressed gene prediction, gene co-expression analysis, *cis*-regulatory motif finding, and comparative genomic study to

predict and validate regulons related to acid stress response in *Lactococcus lactis*
MG1363. A total of 51 regulons were identified, and 14 of them have computational
verified significance. Among these 14 regulons, five of them were computationally
predicted to be connected with acid stress response with (i) known transcriptional
factors in MEME suite database successfully mapped in *Lactococcus lactis* MG1363;
and (ii) differentially expressed genes between pH values of 6.5 (control) and 5.1
(treatment). Validated by 36 literature confirmed acid stress response related proteins
and genes, 33 genes in *Lactococcus lactis* MG1363 were found having orthologous
genes using BLAST, associated to six regulons. An acid response related regulatory
network was constructed, involving two trans-membrane proteins, eight regulons
(*llrA*, *llrC*, *hllA*, *ccpA*, NHP6A, *rcfB*, regulons #8 and #39), nine functional modules,
and 33 genes with orthologous genes known to be associated to acid stress. Our
RECTA pipeline provides an effective way to construct a reliable gene regulatory
network based on regulon elucidation. The predicted resistance pathways could serve
as promising candidates for better acid tolerance engineering in *Lactococcus lactis*. It
has a strong application power and can be effectively applied to other bacterial
genomes, where the elucidation of the transcriptional regulation network is needed.

KEYWORDS

Lactococcus lactis MG1363; Acid stress response; Differential expressed gene; Gene
co-expression; *Cis*-regulatory motif finding; Regulon; Gene regulatory network.

INTRODUCTION

Lactococcus lactis (*L. lactis*) is one of the mesophilic Gram-positive lactic acid-
producing bacteria. It is has been widely applied in dairy fermentations such as cheese
and milk product (Wegmann et al. 2007). Several studies have provided evidence of
its essential roles in desecrating and delivering proteins or vaccine for immune
treatment such as diabetes (Ma, Liu, et al. 2014), malaria (Ramasamy et al. 2006),
tumor (Bermudez-Humaran et al. 2005, Zhang et al. 2016) and infections (Hanniffy et
al. 2007). Holding the advantage of higher acid tolerance to protect vectors from
resolving during delivery inside of the animal body, *L. lactis* has more potential and
safety in oral drug development (Hols et al. 1999). Moreover, it has been found that *L.
lactis*, along with some *Lactobacillus*, *Bifidobacterium* and other gut microbiota, were
associated with obesity (Million et al. 2012). Such studies lead to the possibility and
availability of *L. lactis* in metagenome study to investigate the effect of microbial
interaction between *L. lactis* and other species in the human body. The optimal pH
range for their growth is between 6.3–6.9 (Hutkins and Nannen 1993). The production
of lactic acid, a weak organic acid with a pK_a of 3.86, leads to pH drop in the
extracellular media, reduced metabolic capability, decreased growth rate and

ultimately reduced cell viability (van de Guchte et al. 2002). It is now well established
that *Lactococcus* have evolved stress-sensing systems, which enable them to tolerate
harsh environmental conditions (Carvalho et al. 2013, Hutkins and Nannen 1993, van
de Guchte, Serror, Chervaux, Smokvina, Ehrlich and Maguin 2002).

The reason that bacteria maintain the protection mechanism to against acid stress is to
withstand the deleterious effects caused by the harmful high level of protons in the
exposed environment. Acid stress is known to change the level of the alarmones
(guanosine tetraphosphate and guanosine pentaphosphate, collectively referred to as
(p)ppGpp) (Hauryliuk et al. 2015) and leads to stringent response (Rallu et al. 2000).
Many mechanisms or genes related to the acid stress response (ASR) have been
identified. Proton–pumping activity as the direct regulator to acid stress response
controls the intracellular pH level by pumping extra protons out of the cell
(Koebmann et al. 2000, Lund et al. 2014), and the increase of alkaline compounds
level also counters the acidification found in streptococci (Shabayek and Spellerberg
2017). Cell repairing for acid damage by chaperones or proteases, such as GroES,
GroEL, GrpE, HrcA, DnaK, DnaJ and Clp (Frees et al. 2003, Jayaraman et al. 1997),
and hdeA/B and Hsp31 in *Escherichia coli* (*E. coli*) (Kern et al. 2007, Mujacic and
Baneyx 2007), the arginine deiminase system (ADI) (Budin-Verneuil et al. 2006,
Ryan et al. 2009, Sun et al. 2012, Zuniga et al. 2002) and glutamate decarboxylases
(GAD) pathways, etc. (Hoskins et al. 2001, Nomura et al. 1999, Sanders et al. 1998)
have been proved to be associated with the acid response. Additionally,
transcriptional regulators, sigma factors, and two-component signal transduction
system (TCSs) have also been demonstrated to be responsible for ASR by modifying
gene expression (Cotter and Hill 2003). These genes or pathways suggest low pH has
widespread or global adverse effects on cell functions and inflicts response at
genomic, metabolic, and macromolecular levels. To better understand the mechanism
that controls the acid tolerance and responds to the acid stress in *L. lactis*, we
considered MG1363, a strain extensively studied for acid resistance, to carry out
computational analyses (Carvalho, Turner, Fonseca, Solopova, Catarino, Kuipers,
Voit, Neves and Santos 2013, Hartke et al. 1996, Linares et al. 2010, Sanders et al.
1999).

To adequately describe the transcriptional state and gene regulation responsible for
ASR in *L. lactis*, a gene regulatory network (GRN) integrating all individual pathways
is needed. Genomic and transcriptomic analyses have been widely used for
elucidating GRN hierarchies and offering insight into the coordination of response
capabilities (Arnoldini et al. 2012, Carvalho, Turner, Fonseca, Solopova, Catarino,
Kuipers, Voit, Neves and Santos 2013, Levine et al. 2013, Locke et al. 2011).

However, an obstacle to this kind of genome-scale network is the lack of well-
documented transcription factors (TFs) and the regulatory mechanisms at the
transcription unit level in MG1363. One way to study the mechanism of
transcriptional regulation in microbe genomics is the regulon prediction. A regulon is
a group of co-regulated operons, which contains single or multiple consecutive genes
along the genome (Cao, Ma, et al. 2017, Mao et al. 2015, Zhou et al. 2014). Genes in
the same operon are controlled by the same promoter and are co-regulated by one or a
set of TFs (Jacob et al. 1960). The elucidation of regulons can improve the
identification of transcriptional genes, and thus, reliably predict the gene transcription
regulation networks (Liu, Zhou, et al. 2016).

There are three ways for regulon prediction, which includes: (i) Predicting new
operons for a known regulon (Kumka and Bauer 2015, Tan et al. 2001). This method
combines motif profiling with a comparative genomic strategy to search for related
regulon members and carries out systematical gene regulation study. (ii) Integrating
cis-regulatory motif (*motif* for short) comparison and clustering to find significantly
enriched motif candidates (Gupta et al. 2007, Ma et al. 2013). The candidate motifs
are then assembled into regulons. (iii) Performing *ab initio* novel regulon inference
using *de novo* motif finding strategy (Novichkov et al. 2010). The approach uses
phylogenetic footprinting technique which mostly relies on the reference verification
(Blanchette et al. 2002, Katara et al. 2012, Liu, Zhang, et al. 2016), and can perform a
horizontal sequential comparison to predict regulons in target organisms by searching
known functional related regulons or TFs from other relevant species. One algorithm
for phylogenetic footprinting analysis called Motif Prediction by Phylogenetic
Footprinting (MP3) has been developed by Dr. Liu etc. and used for regulon
prediction in *E. coli* (Liu, Zhang, Zhou, Li, Fennell, Wang, Kang, Liu and Ma 2016).
MP3 was then integrated into DMINDA web server along with other algorithms like
the Database of Prokaryotic Operons 2.0 (DOOR2) (Cao, Ma, Chen and Xu 2017,
Mao et al. 2014), BOttleneck BROken (BoBro) (Li, Ma, et al. 2011) and BoBro-based
motif Comparison (BBC) (Ma, Liu, Zhou, Yin, Li and Xu 2013), to construct a
complete pipeline for regulon prediction. In the latest research, a newly developed
pipeline called Single-cell Regulatory Network Inference and Clustering (SCENIC)
combines motif finding from co-expression gene modules (CEMs) with regulon
prediction for single-cell clustering and analysis (Aibar et al. 2017). Such mentality
builds up a way of regulon application in single-cell and metagenomic research.
Nevertheless, without a suitable regulon database, researchers need to build up the
library first through operon identification, CEM analysis, motif prediction and
comparison (Jensen et al. 2005).

Here, we designed a computational framework of regulon identification based on
comparative genomics and transcriptomics analysis (**RECTA**) to elucidate the GRN
responding to acid stress in MG1363. The general framework is showcased in Figure
1 with six steps: (i) MG1363 co-expression gene modules (CEMs) and differentially
expressed genes (DEG) were generated from microarray data by hcluster package and
Wilcoxon test in R, respectively. MG1363 operons were predicted from genome
sequence using DOOR2 webserver and assigned into each CEM; (ii) For each CEM,
the 300bp upstream to the promoter was extracted and the sequences were used to
find motifs using DMINDA2.0; (iii) The top five significant motifs in each CEM were
reassembled by their similarity comparison and clustering to predict regulons; (iv)
The motifs were compared to known transcription factor binding sites (TFBSs) in the
MEME suite, and the TFs corresponding to these TFBSs were mapped to MG1363
using BLAST. Only regulons with DEGs and mapped TF were kept as ASR-related
regulons; (v) Experimentally identified ASR-related genes in other organisms were
mapped to MG1363 using BLAST and allocated to corresponding regulons for further
verification; and (vi) The relationship between regulons and functional gene modules
were established to elucidate the overall ASR mechanism in MG1363.

Figure 1 should be placed here.

As a result, 14 regulons are identified, literature verified or putative, to be connected
to ASR. Eight regulons, related to nine functional modules and 33 associated genes,
which are considered as the essential elements in acid resistance in MG1363. This
proposed computational pipeline and the above results significantly expand the
current understanding of the ASR system, providing a new method to predict
systematic regulatory network based on regulon clustering.

METHODS

Data Acquisition

L. lactic MG1363 genome sequence was downloaded from NCBI (GenBank
accession number: AM406671). The microarray dataset contains eight samples under
different acid stress conditions for MG1363 was downloaded from the Gene
Expression Omnibus (GEO) Database (Series number: GSE47012). The data has been
treated with LOWESS normalization by the provider. The details on cell culture
preparation and data processing can be found in the previous study (Carvalho, Turner,
Fonseca, Solopova, Catarino, Kuipers, Voit, Neves and Santos 2013). This dataset has

all the bacteria grown in the basic conditions: a 2-liter fermenter in chemically
defined medium containing 1% (w/v) glucose at 30°C. The control and treatment
samples were grown at a pH of 6.5 and 5.1, respectively.

Several TFBS databases integrating in the MEME suite, including DPINTERACT (*E.*
coli) (http://arep.med.harvard.edu/dpinteract/), JASPAR (Anthony Mathelier 2016),
RegTransBase (Prokaryotes) (Kazakov et al. 2007), Prodoric Release (Prokaryotes)
(Munch et al. 2003), and Yeastract (Yeast) (Teixeira et al. 2014), were utilized for
regulon filtering in known TF templates to find homologous TFs and corresponding
genes in MG1363 using BLAST with default parameters. In literature validation part,
all ASR-related transporters and genes were collected from published articles, and
their sequences were obtained from NCBI and UniProt databases.

Operon identification

The genome-scale operons of MG1363 were identified by DOOR2. It is a one-stop
operon-centered resource including operons, alternative transcriptional units, motifs,
terminators, conserved operons information across multiple species (Mao, Ma, Zhou,
Chen, Zhang, Yang, Mao, Lai and Xu 2014). Operons were predicted by the back-end
prediction algorithm, with a prediction accuracy of 90-95%, based on the features of
intergenic distance, neighborhood conservation, short DNA motifs, length ratio
between gene pairs, and newly developed transcriptomic features trained from the
strand-specific RNA-seq dataset (Chou et al. 2015, Li, Liu, et al. 2011).

Gene differential expression analysis and co-expression analysis

DEGs were identified based on the Wilcoxon signed-rank test (Bauer 1972) between
the control and treatment, which was performed in R. The gene co-expression analysis
was performed using a hierarchical clustering method ('hcluster' package in R
(Antoine Lucas 2006)) to detect the CEMs under the acid stress in MG1363.

Motif finding and regulon prediction

Genes from each CEM were first mapped to the identified operons to retrieve the
basic transcription units. Next, 300 bps in the upstream of the translation starting sites
for each operon were extracted, in which motif finding was carried out using the web
server DMINDA (Ma, Zhang, et al. 2014, Yang et al. 2017). DMINDA is a dominant

motif prediction tool, embraced five analytical algorithms to find, scan, and compare
motifs (Li, Liu, Ma and Xu 2011, Liu et al. 2017, Ma, Liu, Zhou, Yin, Li and Xu
2013), including a phylogenetic footprint framework to elucidate the mechanism of
transcriptional regulation at a system level in prokaryotic genomes (Li, Ma, Mao, Yin,
Zhu and Xu 2011, Liu, Zhang, Zhou, Li, Fennell, Wang, Kang, Liu and Ma 2016, Liu,
Zhou, Li, Zhang, Zeng, Liu and Ma 2016). All sequences were uploaded to the server
and default parameters were used to find the top five significant motifs (p-value <
0.05) in each cluster. The identified motifs were subjected to motif comparison and
grouped into regulons based a sequence similarity cutoff using the BBC program in
DMINDA (Ma, Liu, Zhou, Yin, Li and Xu 2013).

Regulon validation based on TF BLAST and DEG filtering.

Each highly conserved motif was considered to contain the same TFBS among
species. Therefore, a comparison study was performed using TOMTOM in the
MEME Suite (Bailey et al. 2009) between identified motif and public-domain TFBS
databases, including DPINTERACT, JASPAR, RegTransBase, Prodoric Release and
Yeastract, to find TFBSs and corresponding TFs with significant p-values in other
prokaryotic species. Those TFs were then mapped to MG1363 using BLAST by
default parameters to predict the connection between regulons and TFs in MG1363.
On the other hand, since genes without differential expression were supposed not to
react to pH changes, and thus, irrelevant to ASR, regulons without DEG were
supposed not involved in the GRN, and thus, excluded from the following steps.

Regulon validation based on known ASR proteins from the literature.

To validate the performance of the above computational pipeline for regulon
prediction, a literature-based validation was performed. Thirty-six ASR-related
proteins and genes in other organisms including *L. lactis, E. coli*, *Streptococci*, *etc.*
were first manually collected from literature, and their sequence was retrieved from
the NCBI and UniProt databases. They were used to examine the existing known
mechanisms in response to pH changes in MG1363 using the BLAST program by
default parameters on NCBI. Such literature-based validation can either confirm the
putative regulons when known ASR-related genes can be found in the significant
regulons or expand our results to some insufficiently significant regulons, which
indicate both false positive and true negative rate to evaluate the computational
pipeline.

RESULTS

Predicted operons and CEM generation

A total of 1,565 operons with 2,439 coding genes of MG1363 (Dataset S1) were
retrieved from the DOOR2 database. By co-expression analysis, the 1,565 operons
were grouped into 124 co-expressed clusters. Among these clusters, two large ones
contain more than 200 operons. Each of them was removed from the subsequent
analyses as larger clusters may have higher chances to induce false positive operons
which were connected with true operons by co-expression analysis. For the rest 122
clusters covering 2,122 genes, 26 (21%) of them contain operons no more than ten;
the smallest cluster had two operons, and most of the clusters (90%) contained
operons in a range of 10 to 50 (Dataset S2 and Figure S1).

Predicted regulons based on motif finding and clustering.

Using BoBro in the DMINDA web server, multiple motif sequences were identified
from the 300 bps in the upstream of the translation start sites for each operon. Only
the top five significant motifs (p-value < 0.05) were selected in each cluster, giving
rise to a total of 610 (122×5) identified. The motif comparison-and-clustering
analysis was then performed on the 610 motifs, and 51 motif clusters were identified
with a motif similarity 0.8 as a cutoff. Intuitively, the operons sharing highly similar
motifs in each motif cluster are supposed to be regulated by the same TF and tend to
be in the same regulon. Hence, these 51 motif clusters correspond to 51 regulons
(Dataset S3).

Computationally verified regulon based on TF BLAST and DEG analysis.

Among the above 51 regulons, 14 were found containing motifs significantly (E-value
< 0.05) matched to known TFBSs using TOMTOM in the MEME suite,
representatively. The motif logos are shown in Figure S2, and more details can be
found in Dataset S4. The 14 corresponding TFs were then mapped to MG1363 using
BLAST as templates to find the TFs/genes regulate of each regulon. As a result, eight
known TFs, spo0A, lhfB, GAL80, CovR, c4494, ihfA, CovR, and RHE_PF00288,
were successfully mapped to MG1363 resulting in eight TFs with duplicates. *llrA*
(llmg_0908) regulates regulons #12 and #37, *ccpA* (llmg_0775) regulated regulons
#15 and #47, *hllA* (llmg_0496) regulates regulons #7 and #31 (Table 1). *CcpA*
(Abranches et al. 2008, Zomer et al. 2007), *llrA* (O'Connell-Motherway et al. 2000),
llrC (O'Connell-Motherway, van Sinderen, Morel-Deville, Fitzgerald, Ehrlich and
Morel 2000), and *hllA* (Bolotin et al. 1999) were known to be ASR-related genes in *L.*

lactis; gene (llmg_0271), without any related known TF, was found similar to
template TF GAL80 in yeast, which has not been associated with any ASR regulation
pathways yet. For all 14 significant regulons, regulons #3, #4, #20, #28, #40, and #44
are potential candidates as, currently, no related TFs in *L. lactis* have been found.

Table 1 should be placed here.

Additionally, 86 down-regulated genes and 55 up-regulated genes (Dataset S5),
resulted from DEG analysis, were integrated into the regulons. Regulons #10, #37,
#44 and #47 were found lack of DEGs. Thus gene (llmg_0271), related to regulon
#10, was not likely to respond to acid stress in MG1363 even has successfully
mapped to MG1363, and was then grouped into the potential candidate. On the
contrary, *ccpA* and *llrA* were still retained due to their involvements in regulons #15
and #12 with DEGs, respectively.

By the end of the computational pipeline, we predicted that regulons #2, #7, #12, #15
and #31 were related to GRN in MG1363. Merging regulon #7 and #31 as one, we
referred their TF names (*ccpA, llrA, llrC,* and *hllA*) to represent the five regulons for
convenience.

Verified regulons based on literature verification

Altogether, 36 literature-supported ASR-related transporters were successfully
mapped to MG1363 using blast with an E-value cutoff as 1e-10 and resulted in a total
of 33 mapped genes. All the 36 transporters were categorized into nine modules based
on their biological functions or regulated pathways, including L-lactate
dehydrogenase (LDH), GAD, ADI, urea degradation, F1/F0ATPase, acid stress,
protein repair and protease, envelope alterations and DNA repair. The 33 mapped
genes compose of 22 operons and six regulons: *llrA*, *llrC*, *hllA*, NHP6A, regulon #8
and #39, which were subjected, one or more, to each functional module (Table 2).

Regulon *llrA*, *llrC*, and *hllA* have already been computationally identified in Table 1
and supported again by literature verification results. NHP6A which interestingly has
a homologous TF in human and fungi but not in *L. lactis* (Kolodrubetz and Burgum
1990, Stillman 2010), yet failed to map in MG1363. Here, we are using NHP6A to

represent regulon #20, as their relationship has been predicted computationally in Table 1. Regulon #39 was identified to be regulated by *llrD*, one of the six two-component regulatory systems in MG1363 (O'Connell-Motherway, van Sinderen, Morel-Deville, Fitzgerald, Ehrlich and Morel 2000). Regulons #8 (llmg_1803) and #39 (*llrD*) were not included in the 14 significant regulons in Table 1. For NHP6A, regulons #8 and #39 were the enrichment by literature validation as it expanded regulon results of RECTA pipeline. Among the nine functional modules, *llrA* were found connected to five of them, and NHP6A related to three. On the other hand, the GAD and urea degradation functional modules were failed to connect to any previous regulons.

Table 2 should be placed here.

Compare to the regulon verification based on TF blast and DEG, the literature verification identified two more regulons (#8 and #39) that lay in the insignificant group, however, with no sign of *ccpA* regulon. Thus, such result indicated a possible false positive rate of 1/5 and a true negative rate of 2/37 of our computational pipeline, indicating a reliability and feasibility of using RECTA to predict the ASR-related regulons. In Figure 2, we are showing the processes and results for both literature verification and computational pipeline in details. The final eight regulons predicted from both parts were then compared to construct a GRN response to acid stress, integrating with other information found in the literature.

Figure 2 should be placed here.

A model of regulatory network in response to pH change

According to the results outlined above, we are presenting a working model of the transcriptional regulatory network in response to acid stress response in MG1363 (Figure 3). The network consists of two transmembrane proteins (Dataset S6), eight regulons, nine functional modules, and 33 orthologous genes known for ASR in other bacteria that are also contributing in MG1363.

Figure 3 should be placed here.

The network is subjected to respond to the change of intracellular proton level. The signal is captured by H+ sensor and regulons are initiated to be regulated. Although significance was not shown for *rcfB* in our computational result, it has been reported to recognize and regulate promoter P170 (Madsen et al. 1999), P1 and P3 (Hindre et al. 2004, Rince et al. 1994), which are activated by the ACiD box and essential to acid response (Madsen et al. 2005). With the ACiD box, operons like *groESL*, *lacticin 481* and *lacZ* have been proved to be regulated by *rcfB*, while *als*, *aldB*, etc. have not (Madsen, Hindre, Le Pennec, Israelsen and Dufour 2005). The homologous comparative study also predicted the existence of the ACiD box in *llrA* (Akyol et al. 2008, O'Connell-Motherway, van Sinderen, Morel-Deville, Fitzgerald, Ehrlich and Morel 2000). With such evidence, we separated *rcfB* from regulon 39 and predicted that *rcfB* is first triggered by H+ sensor and act as the global initiator that controls the other seven regulons. It was reasonable that *rcfB*-related regulon #39 failed to show significant TF matching result after CEM treatment in the operon clustering step. *RcfB* worked as a trustworthy global factor; its differential expression should be less significant than regulons directly respond to acid stress, thus lead to the failure of being predicted by the RECTA pipeline. Nevertheless, the less amount of microarray data sets (eight) also limited the real performance to the ASR. However, the mechanism of how H+ sensor is activating and regulating the GRN and *rcfB* remains unclear. In the seven regulons, three of them, *llrA*, *llrC* and *hllA*, were documentary verified to be related to ASR; regulons #8 and #39 showed less significant in regulon prediction; NHP6A was considered as putative regulon due to its failure mapping in MG1363; and *ccpA* was another putative regulon without literature support.

The six downstream regulons (*llrA*, *llrC*, *hllA*, NHP6A, regulon #39, and regulon #8) other than *ccpA*, interact with each other to regulate six ASR-related functional modules, including ADI system, DNA repair, LDH, protein repair, envelope alterations, and F0/F1ATPase. The ADI pathway, which generates ATP and protects cells from acid stress (Zuniga, Perez and Gonzalez-Candelas 2002), is under the regulation of NHP6A, *llrC*, *llrA*, and *hllA*. Another important pathway is the LDH (EC 1.1.1.27) under the regulation of NHP6A and *llrA*, which converts pyruvate and H+ to lactate which is exported outside of cells (Dennis and Kaplan 1960). Chaperons which take part in macromolecule protection and repairing are subjected to regulon *llrA*. The function of chaperons includes providing protection to against environmental stress, helping protein folding, and repairing damaged proteins, and have been demonstrated to show clear linkage with acid stress in numerous Gram-positive bacteria (Frees, Vogensen and Ingmer 2003, Jayaraman, Penders and Burne 1997, Kern, Malki, Abdallah, Tagourti and Richarme 2007, Mujacic and Baneyx 2007). F0/F1ATPase, controlled by *llrA* and regulon #8, also plays an important role

in maintaining normal cellular pH, which pumps H+ out of cells at the expense of
ATP (Amachi et al. 1998, Koebmann, Nilsson, Kuipers and Jensen 2000, Lund,
Tramonti and De Biase 2014, O'Sullivan and Condon 1999). The GAD (Nomura,
Nakajima, Fujita, Kobayashi, Kimoto, Suzuki and Aso 1999, Sanders, Leenhouts,
Burghoorn, Brands, Venema and Kok 1998) and urea degradation (Cotter and Hill
2003) functional modules are missing reliable associations with the regulons in
MG1363 while maintaining functions in ASR mechanism in other species.

DISCUSSION

In this study, we designed a computational framework, RECTA, to construct an eight-
regulons enrolled ASR regulatory network. The framework provides a useful tool and
will be a starting point toward a more systems-level understanding of the question
(Cao, Wei, et al. 2017). The identified motifs and regulons suggest the acid resistance
is a coordinated response regarding regulons, although most of which have not been
identified or experimentally verified. From the three well-identified regulons—*llrA*,
llrC, and *hllA*, it appears the gene regulation is also complex, as these regulons also
interact with other proteins and TFs. F0/F1ATPase is directly involved in the
concentration regulations of the intracellular proton. Other pathways are responsible
for repairing the damage caused by acid stress, such as DNA repair, protein repair,
and cell envelops alterations. However, there were also several reported ASR-related
genes or transporters such as *htrA* in *Clostridium spp*. (Alsaker et al. 2010),
CovS/CovR acid response regulator in *Streptococcus* (Cumley et al. 2012),
cyclopropane fatty acid synthase (cfa) for cell-membrane modification (Budin-
Verneuil et al. 2005), and oxidative damage protectant genes like *sodA*, *nox-1* and
nox-2 (Santi et al. 2009) that failed to map to MG1363. Using more gene expression
datasets for CEM and DEG analyses could be a way to strengthen the result of our
computational pipeline, which might cover more significant regulons to construct a
more solid and complete regulatory network.

Homology mapping at the genomic level showed very long distance in evolution
between MG1363 and currently well-annotated model species. Hence, the functional
analysis for MG1363 is limited, and it is hard to apply gene functional enrichment to
verify our prediction results. With more expression dataset and experiments about
protein-protein interactions, the ASR mechanism can be largely improved in *L. lactic*
MG1363.

In summary, we found that the ASR at the transcriptome level in MG1363 is an
orchestrated complex network. Functional annotation shows these regulons are
involved in many levels of biological processes, including but not limited to DNA
expression, transcription, and metabolism. Our method builds a TF-regulons-GRN
relationship so that the new ASR-related genes can be predicted. Besides, the low
false positive and true negative rate indicate the RECTA pipeline as sensitive and
reasonable. In fact, considering the high accuracy, we regarded *ccpA* as the putative
regulon, though not connected to any related functional modules, while more robust
proves are required. Such results expand current pathways to those that can
corroborate cell structures—cell wall, cell membranes—and related functions. Our
findings suggest that acid has profound adverse effects and inflict systems-level
response. Such predicted response pathways can inform better resistance design.

Looking forward to the acid tolerance advantage of *L. lactis*, which makes its
prospective application in drug and vaccine delivery, the effects on anti-obesity, and
the metagenomic study, the ASR-related GRN in *L. lactis* shows an excellent research
value. Fully understanding of its theory may contribute to the development of
Lactococcus therapy, and can even expand to other close species by genetic
modification. Furthermore, our computational pipeline provides an effective way to
construct reliable GRN based on regulon prediction, integrating CEMs, DEG analysis,
motif finding, and comparative genomics study. It has a durable application power
and can be effectively applied to other bacterial genomes, where the elucidation of the
transcriptional regulation network is needed.

LIST OF ABBREVIATIONS

ADI: arginine deiminase
BoBro-based motif Comparison (BBC)
BOttleneck BROken (BoBro)
CEM: co-expression (gene) module
DEG: differentially expressed gene
E. coli: Escherichia coli
GAD: glutamate decarboxylases
GRN: gene regulatory network
LDH: lactate dehydrogenase
L. lactis: *Lactococcus lactis*
MG1363: *Lactococcus lactis* MG1363
Motif: cis-regulatory motif

RECTA: regulon identification based on comparative genomics and transcriptomics
analysis
TF: transcription factors
TFBS: transcriptional factor binding site

DECLARATIONS

Ethics Approval and Consent to Participate

Not applicable

Conflict of Interest Statement

The authors declare that they have no competing interests

Data Availability Statements

All datasets analyzed and generated for this study are included in the manuscript and
the supplementary files.

Funding Statements

This work was supported by the State of South Dakota Research Innovation Center,
the Agriculture Experiment Station of South Dakota State University, National
Science Foundation of United States (1546869), Sanford Health – SDSU
Collaborative Research Seed Grant Program, and China National 863 High-Tech
Program (Grant No. 2015AA020101). This work used the Extreme Science and
Engineering Discovery Environment (XSEDE), which is supported by National
Science Foundation grant number ACI-1548562.

Authors' Contributions

XC and QM carried out the framework of the paper and are responsible for the
regulon prediction work. HZ carried out the operon prediction and motif finding
work. CL was responsible for DEG analysis. HC and AM drafted and revised the
manuscript. All authors submitted comments, read and approved the final manuscript.

REFERENCES

Abranches J, Nascimento MM, Zeng L, Browngardt CM, Wen ZT, Rivera MF, Burne RA. 2008. CcpA
regulates central metabolism and virulence gene expression in Streptococcus mutans. J Bacteriol.
Apr;190:2340-2349. Epub 2008/01/29.

Aibar S, Gonzalez-Blas CB, Moerman T, Huynh-Thu VA, Imrichova H, Hulselmans G, Rambow F, Marine
JC, Geurts P, Aerts J, et al. 2017. SCENIC: single-cell regulatory network inference and clustering. Nat
Methods. Nov;14:1083-1086. Epub 2017/10/11.

Akyol I, Comlekcioglu U, Karakas A, Serdaroglu K, Ekinci MS, Ozkose E. 2008. Regulation of the acid
inducible *rcfB* promoter in *Lactococcus lactis* subsp. lactis. Annals of Microbiology. 2008//;58:269.

Alsaker KV, Paredes C, Papoutsakis ET. 2010. Metabolite stress and tolerance in the production of
biofuels and chemicals: gene-expression-based systems analysis of butanol, butyrate, and acetate
stresses in the anaerobe Clostridium acetobutylicum. Biotechnol Bioeng. Apr 15;105:1131-1147. Epub
2009/12/10.

Amachi S, Ishikawa K, Toyoda S, Kagawa Y, Yokota A, Tomita F. 1998. Characterization of a mutant of
Lactococcus lactis with reduced membrane-bound ATPase activity under acidic conditions. Biosci
Biotechnol Biochem. Aug;62:1574-1580. Epub 1998/10/03.

Anthony Mathelier OF, David J. Arenillas, Chih-yu Chen, Grégoire Denay, Jessica Lee, Wenqiang Shi,
Casper Shyr, Ge Tan, Rebecca Worsley-Hunt Allen W. Zhang, François Parcy, Boris Lenhard,
Albin Sandelin, Wyeth W. Wasserman. 2016. JASPAR 2016: a major expansion and update of the open-
access database of transcription factor binding profiles Nucleic Acids Research. 03 November
2015;44:D110–D115.

Antoine Lucas SJ. 2006. Using amap and ctc Packages for Huge Clustering. R News.6:58-60.

Arnoldini M, Mostowy R, Bonhoeffer S, Ackermann M. 2012. Evolution of Stress Response in the Face
of Unreliable Environmental Signals. PLoS Comput Biol.8:e1002627.

Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble WS. 2009. MEME Suite:
tools for motif discovery and searching. Nucleic Acids Research. July 1, 2009;37:W202-W208.

Bauer DF. 1972. Constructing Confidence Sets Using Rank Statistics. Journal of the American Statistical
Association.67:687-690.

Bermudez-Humaran LG, Cortes-Perez NG, Lefevre F, Guimaraes V, Rabot S, Alcocer-Gonzalez JM,
Gratadoux JJ, Rodriguez-Padilla C, Tamez-Guerra RS, Corthier G, et al. 2005. A novel mucosal vaccine
based on live Lactococci expressing E7 antigen and IL-12 induces systemic and mucosal immune
responses and protects mice against human papillomavirus type 16-induced tumors. J Immunol. Dec
01;175:7297-7302. Epub 2005/11/23.

Blanchette M, Schwikowski B, Tompa M. 2002. Algorithms for phylogenetic footprinting. J Comput
Biol.9:211-223. Epub 2002/05/23.

Bolotin A, Mauger S, Malarme K, Ehrlich SD, Sorokin A. 1999. Low-redundancy sequencing of the entire
Lactococcus lactis IL1403 genome. Antonie Van Leeuwenhoek. Jul-Nov;76:27-76. Epub 1999/10/26.

Budin-Verneuil A, Maguin E, Auffray Y, Ehrlich DS, Pichereau V. 2006. Genetic structure and
transcriptional analysis of the arginine deiminase (ADI) cluster in Lactococcus lactis MG1363. Can J
Microbiol. Jul;52:617-622.

Budin-Verneuil A, Maguin E, Auffray Y, Ehrlich SD, Pichereau V. 2005. Transcriptional analysis of the
cyclopropane fatty acid synthase gene of Lactococcus lactis MG1363 at low pH. FEMS Microbiol Lett.
Sep 15;250:189-194. Epub 2005/08/16.

Cao H, Ma Q, Chen X, Xu Y. 2017. DOOR: a prokaryotic operon database for genome analyses and
functional inference. Brief Bioinform. Jul 28. Epub 2017/10/03.

Cao H, Wei D, Yang Y, Shang Y, Li G, Zhou Y, Ma Q, Xu Y. 2017. Systems-level understanding of ethanol-
induced stresses and adaptation in E. coli. Scientific Reports. 03/16/online;7:44150.

Carvalho AL, Turner DL, Fonseca LL, Solopova A, Catarino T, Kuipers OP, Voit EO, Neves AR, Santos H.
2013. Metabolic and transcriptional analysis of acid stress in *Lactococcus lactis*, with a focus on the
kinetics of lactic acid pools. PLOS ONE.8:e68470.

Chou WC, Ma Q, Yang S, Cao S, Klingeman DM, Brown SD, Xu Y. 2015. Analysis of strand-specific RNA-
seq data using machine learning reveals the structures of transcription units in Clostridium
thermocellum. Nucleic Acids Res. May 26;43:e67. Epub 2015/03/15.

Cotter PD, Hill C. 2003. Surviving the acid test: responses of gram-positive bacteria to low pH. Microbiol
Mol Biol Rev. Sep;67:429-453, table of contents. Epub 2003/09/11.

Cumley NJ, Smith LM, Anthony M, May RC. 2012. The CovS/CovR acid response regulator is required for
intracellular survival of group B Streptococcus in macrophages. Infect Immun. May;80:1650-1661. Epub
2012/02/15.

Dennis D, Kaplan NO. 1960. D- and L-lactic acid dehydrogenases in Lactobacillus plantarum. J Biol Chem.
Mar;235:810-818. Epub 1960/03/01.

Frees D, Vogensen FK, Ingmer H. 2003. Identification of proteins induced at low pH in Lactococcus lactis.
International Journal of Food Microbiology. 11/1/;87:293-300.

Gupta S, Stamatoyannopoulos JA, Bailey TL, Noble WS. 2007. Quantifying similarity between motifs.
Genome Biol.8:R24. Epub 2007/02/28.

Hanniffy SB, Carter AT, Hitchin E, Wells JM. 2007. Mucosal delivery of a pneumococcal vaccine using
Lactococcus lactis affords protection against respiratory infection. J Infect Dis. Jan 15;195:185-193. Epub
2006/12/28.

Hartke A, Bouché S, Giard J-C, Benachour A, Boutibonnes P, Auffray Y. 1996. The lactic acid stress
response of *Lactococcus lactis* subsp. lactis. Current Microbiology.33:194-199.

Hauryliuk V, Atkinson GC, Murakami KS, Tenson T, Gerdes K. 2015. Recent functional insights into the

role of (p)ppGpp in bacterial physiology. Nat Rev Micro. 05//print;13:298-309.

Hindre T, Le Pennec JP, Haras D, Dufour A. 2004. Regulation of lantibiotic lacticin 481 production at the
transcriptional level by acid pH. FEMS Microbiol Lett. Feb 16;231:291-298. Epub 2004/02/28.

Hols P, Kleerebezem M, Schanck AN, Ferain T, Hugenholtz J, Delcour J, de Vos WM. 1999. Conversion of
Lactococcus lactis from homolactic to homoalanine fermentation through metabolic engineering. Nat
Biotechnol. Jun;17:588-592. Epub 1999/06/29.

Hoskins J, Alborn WE, Jr., Arnold J, Blaszczak LC, Burgett S, DeHoff BS, Estrem ST, Fritz L, Fu DJ, Fuller W,
et al. 2001. Genome of the bacterium Streptococcus pneumoniae strain R6. J Bacteriol. Oct;183:5709-
5717. Epub 2001/09/07.

Hutkins RW, Nannen NL. 1993. pH homeostasis in lactic acid bacteria. Journal of Dairy Science.
1993/08/01;76:2354-2365.

Jacob F, Perrin D, Sanchez C, Monod J. 1960. [Operon: a group of genes with the expression coordinated
by an operator]. C R Hebd Seances Acad Sci. Feb 29;250:1727-1729. Epub 1960/02/29.

Jayaraman GC, Penders JE, Burne RA. 1997. Transcriptional analysis of the Streptococcus mutans hrcA,
grpE and dnaK genes and regulation of expression in response to heat shock and environmental
acidification. Mol Microbiol. Jul;25:329-341. Epub 1997/07/01.

Jensen ST, Shen L, Liu JS. 2005. Combining phylogenetic motif discovery and motif clustering to predict
co-regulated genes. Bioinformatics. Oct 15;21:3832-3839. Epub 2005/08/18.

Katara P, Grover A, Sharma V. 2012. Phylogenetic footprinting: a boost for microbial regulatory genomics.
Protoplasma. Oct;249:901-907. Epub 2011/11/25.

Kazakov AE, Cipriano MJ, Novichkov PS, Minovitsky S, Vinogradov DV, Arkin A, Mironov AA, Gelfand MS,
Dubchak I. 2007. RegTransBase--a database of regulatory sequences and interactions in a wide range of
prokaryotic genomes. Nucleic Acids Res. Jan;35:D407-412. Epub 2006/12/05.

Kern R, Malki A, Abdallah J, Tagourti J, Richarme G. 2007. Escherichia coli HdeB is an acid stress
chaperone. J Bacteriol. Jan;189:603-610. Epub 2006/11/07.

Koebmann BJ, Nilsson D, Kuipers OP, Jensen PR. 2000. The membrane-bound H^+-ATPase complex is
essential for growth of *Lactococcus lactis*. Journal of Bacteriology. September 1, 2000;182:4738-4743.

Kolodrubetz D, Burgum A. 1990. Duplicated NHP6 genes of Saccharomyces cerevisiae encode proteins
homologous to bovine high mobility group protein 1. J Biol Chem. Feb 25;265:3234-3239. Epub
1990/02/25.

Kumka JE, Bauer CE. 2015. Analysis of the FnrL regulon in Rhodobacter capsulatus reveals limited
regulon overlap with orthologues from Rhodobacter sphaeroides and Escherichia coli. BMC Genomics.
Nov 04;16:895. Epub 2015/11/06.

Levine JH, Lin Y, Elowitz MB. 2013. Functional roles of pulsing in genetic circuits. Science.342:1193-1200.

Li G, Liu B, Ma Q, Xu Y. 2011. A new framework for identifying cis-regulatory motifs in prokaryotes.
Nucleic Acids Res. Apr;39:e42. Epub 2010/12/15.

Li G, Ma Q, Mao X, Yin Y, Zhu X, Xu Y. 2011. Integration of sequence-similarity and functional association
information can overcome intrinsic problems in orthology mapping across bacterial genomes. Nucleic
Acids Res. Dec;39:e150. Epub 2011/10/04.

Linares DM, Kok J, Poolman B. 2010. Genome sequences of *Lactococcus lactis* MG1363 (revised) and
NZ9000 and comparative physiological studies. Journal of Bacteriology. November 1, 2010;192:5806-
5812.

Liu B, Yang J, Li Y, McDermaid A, Ma Q. 2017. An algorithmic perspective of de novo cis-regulatory motif
finding based on ChIP-seq data. Brief Bioinform. Mar 08. Epub 2017/03/24.

Liu B, Zhang H, Zhou C, Li G, Fennell A, Wang G, Kang Y, Liu Q, Ma Q. 2016. An integrative and applicable
phylogenetic footprinting framework for cis-regulatory motifs identification in prokaryotic genomes.
BMC genomics. Aug 09;17:578. Epub 2016/08/11.

Liu B, Zhou C, Li G, Zhang H, Zeng E, Liu Q, Ma Q. 2016. Bacterial regulon modeling and prediction based
on systematic cis regulatory motif analyses. Sci Rep. Mar 15;6:23030. Epub 2016/03/16.

Locke JCW, Young JW, Fontes M, Jiménez MJH, Elowitz MB. 2011. Stochastic pulse regulation in bacterial
stress response. Science.334:366-369.

Lund P, Tramonti A, De Biase D. 2014. Coping with low pH: molecular strategies in neutralophilic bacteria.
FEMS Microbiol Rev. Nov;38:1091-1125. Epub 2014/06/06.

Ma Q, Liu B, Zhou C, Yin Y, Li G, Xu Y. 2013. An integrated toolkit for accurate prediction and analysis of
cis-regulatory motifs at a genome scale. Bioinformatics. Sep 15;29:2261-2268. Epub 2013/07/13.

Ma Q, Zhang H, Mao X, Zhou C, Liu B, Chen X, Xu Y. 2014. DMINDA: an integrated web server for DNA
motif identification and analyses. Nucleic Acids Research. April 21, 2014.

Ma Y, Liu J, Hou J, Dong Y, Lu Y, Jin L, Cao R, Li T, Wu J. 2014. Oral administration of recombinant
Lactococcus lactis expressing HSP65 and tandemly repeated P277 reduces the incidence of type I
diabetes in non-obese diabetic mice. PLoS One.9:e105701. Epub 2014/08/27.

Madsen SM, Arnau J, Vrang A, Givskov M, Israelsen H. 1999. Molecular characterization of the pH-
inducible and growth phase-dependent promoter P170 of Lactococcus lactis. Mol Microbiol. Apr;32:75-
87. Epub 1999/04/27.

Madsen SM, Hindre T, Le Pennec JP, Israelsen H, Dufour A. 2005. Two acid-inducible promoters from
Lactococcus lactis require the cis-acting ACiD-box and the transcription regulator RcfB. Mol Microbiol.
May;56:735-746. Epub 2005/04/12.

Mao X, Ma Q, Liu B, Chen X, Zhang H, Xu Y. 2015. Revisiting operons: an analysis of the landscape of
transcriptional units in E. coli. BMC Bioinformatics. Nov 04;16:356. Epub 2015/11/06.

Mao X, Ma Q, Zhou C, Chen X, Zhang H, Yang J, Mao F, Lai W, Xu Y. 2014. DOOR 2.0: presenting operons
and their functions through dynamic and integrated views. Nucleic Acids Research. January 1,
2014;42:D654-D659.

Million M, Maraninchi M, Henry M, Armougom F, Richet H, Carrieri P, Valero R, Raccah D, Vialettes B,
Raoult D. 2012. Obesity-associated gut microbiota is enriched in Lactobacillus reuteri and depleted in
Bifidobacterium animalis and Methanobrevibacter smithii. Int J Obes (Lond). Jun;36:817-825. Epub
2011/08/11.

Mujacic M, Baneyx F. 2007. Chaperone Hsp31 contributes to acid resistance in stationary-phase
Escherichia coli. Appl Environ Microbiol. Feb;73:1014-1018. Epub 2006/12/13.

Munch R, Hiller K, Barg H, Heldt D, Linz S, Wingender E, Jahn D. 2003. PRODORIC: prokaryotic database
of gene regulation. Nucleic Acids Res. Jan 01;31:266-269. Epub 2003/01/10.

Nomura M, Nakajima I, Fujita Y, Kobayashi M, Kimoto H, Suzuki I, Aso H. 1999. Lactococcus lactis
contains only one glutamate decarboxylase gene. Microbiology. Jun;145 (Pt 6):1375-1380.

Novichkov PS, Rodionov DA, Stavrovskaya ED, Novichkova ES, Kazakov AE, Gelfand MS, Arkin AP, Mironov
AA, Dubchak I. 2010. RegPredict: an integrated system for regulon inference in prokaryotes by
comparative genomics approach. Nucleic Acids Res. Jul;38:W299-307. Epub 2010/06/15.

O'Connell-Motherway M, van Sinderen D, Morel-Deville F, Fitzgerald GF, Ehrlich SD, Morel P. 2000. Six
putative two-component regulatory systems isolated from Lactococcus lactis subsp. cremoris MG1363.
Microbiology. Apr;146 (Pt 4):935-947. Epub 2000/04/28.

O'Sullivan E, Condon S. 1999. Relationship between acid tolerance, cytoplasmic pH, and ATP and H+-
ATPase levels in chemostat cultures of Lactococcus lactis. Appl Environ Microbiol. Jun;65:2287-2293.
Epub 1999/05/29.

Rallu F, Gruss A, Ehrlich SD, Maguin E. 2000. Acid- and multistress-resistant mutants of Lactococcus
lactis : identification of intracellular stress signals. Molecular Microbiology.35:517-528.

Ramasamy R, Yasawardena S, Zomer A, Venema G, Kok J, Leenhouts K. 2006. Immunogenicity of a
malaria parasite antigen displayed by Lactococcus lactis in oral immunisations. Vaccine. May
01;24:3900-3908. Epub 2006/03/21.

Rince A, Dufour A, Le Pogam S, Thuault D, Bourgeois CM, Le Pennec JP. 1994. Cloning, expression, and
nucleotide sequence of genes involved in production of lactococcin DR, a bacteriocin from lactococcus
lactis subsp. lactis. Appl Environ Microbiol. May;60:1652-1657. Epub 1994/05/01.

Ryan S, Begley M, Gahan CG, Hill C. 2009. Molecular characterization of the arginine deiminase system

in Listeria monocytogenes: regulation and role in acid tolerance. Environ Microbiol. Feb;11:432-445.
Epub 2009/02/07.

Sanders JW, Leenhouts K, Burghoorn J, Brands JR, Venema G, Kok J. 1998. A chloride-inducible acid
resistance mechanism in Lactococcus lactis and its regulation. Mol Microbiol. Jan;27:299-310. Epub
1998/03/04.

Sanders JW, Venema G, Kok J. 1999. Environmental stress responses in *Lactococcus lactis*. FEMS
Microbiology Reviews.23:483-501.

Santi I, Grifantini R, Jiang SM, Brettoni C, Grandi G, Wessels MR, Soriani M. 2009. CsrRS regulates group
B Streptococcus virulence gene expression in response to environmental pH: a new perspective on
vaccine development. J Bacteriol. Sep;191:5387-5397. Epub 2009/06/23.

Shabayek S, Spellerberg B. 2017. Acid Stress Response Mechanisms of Group B Streptococci. Front Cell
Infect Microbiol.7:395. Epub 2017/09/25.

Stillman DJ. 2010. Nhp6: a small but powerful effector of chromatin structure in Saccharomyces
cerevisiae. Biochim Biophys Acta. Jan-Feb;1799:175-180. Epub 2010/02/04.

Sun Y, Fukamachi T, Saito H, Kobayashi H. 2012. Adenosine deamination increases the survival under
acidic conditions in Escherichia coli. J Appl Microbiol. Apr;112:775-781. Epub 2012/01/27.

Tan K, Moreno-Hagelsieb G, Collado-Vides J, Stormo GD. 2001. A comparative genomics approach to
prediction of new members of regulons. Genome Res. Apr;11:566-584. Epub 2001/04/03.

Teixeira MC, Monteiro PT, Guerreiro JF, Goncalves JP, Mira NP, dos Santos SC, Cabrito TR, Palma M, Costa
C, Francisco AP, et al. 2014. The YEASTRACT database: an upgraded information system for the analysis
of gene and genomic transcription regulation in Saccharomyces cerevisiae. Nucleic Acids Res.
Jan;42:D161-166. Epub 2013/10/31.

van de Guchte M, Serror P, Chervaux C, Smokvina T, Ehrlich SD, Maguin E. 2002. Stress responses in
lactic acid bacteria. Antonie van Leeuwenhoek.82:187-216.

Wegmann U, O'Connell-Motherway M, Zomer A, Buist G, Shearman C, Canchaya C, Ventura M,
Goesmann A, Gasson MJ, Kuipers OP, et al. 2007. Complete Genome Sequence of the Prototype Lactic
Acid Bacterium Lactococcus lactis subsp. cremoris MG1363. Journal of Bacteriology. April 15,
2007;189:3256-3270.

Yang J, Chen X, McDermaid A, Ma Q. 2017. DMINDA 2.0: integrated and systematic views of regulatory
DNA motif identification and analyses. Bioinformatics. Apr 13.

Zhang B, Li A, Zuo F, Yu R, Zeng Z, Ma H, Chen S. 2016. Recombinant Lactococcus lactis NZ9000 secretes
a bioactive kisspeptin that inhibits proliferation and migration of human colon carcinoma HT-29 cells.
Microb Cell Fact. Jun 10;15:102. Epub 2016/06/12.

Zhou C, Ma Q, Li G. 2014. Elucidation of operon structures across closely related bacterial genomes.
PLoS One.9:e100999. Epub 2014/06/25.

Zomer AL, Buist G, Larsen R, Kok J, Kuipers OP. 2007. Time-resolved determination of the CcpA regulon
of Lactococcus lactis subsp. cremoris MG1363. J Bacteriol. Feb;189:1366-1381. Epub 2006/10/10.

Zuniga M, Perez G, Gonzalez-Candelas F. 2002. Evolution of arginine deiminase (ADI) pathway genes.
Mol Phylogenet Evol. Dec;25:429-444. Epub 2002/11/27.

FIGURE LEGEND

Figure 1: The flowchart of constructing global ASR transcriptional network in
MG1363. Step 1: microarray data was used to generate co-expressed gene clusters
and DEG, and MG1363 genome sequence was used to find operons. Step 2: a motif
finding progress was carried out to identify all statistically significant motifs in each
of the CEMs. Step 3: a regulon finding procedure was designed to identify all the
possible regulon candidates encoded in the genome based on motif comparison and
clustering. Step 4: the motifs of each of these regulons were compared to known
TFBSs, and DEG analysis between low pH condition and normal condition was used
to figure out the ASR-related regulons. Step 5: regulon validation based on literature
information verified the significant putative regulons and expanded the results to
some insufficiently significant regulons. Step 6: the ASR-related GRN in MG1363
was predicted and described with eight regulons, nine functional modules, and 33
genes. The combination of the above information forms a genome-scale regulatory
network constructed for ASR.

Figure 2.   Regulon prediction using RECTA pipeline (red) and validation and
enrichment using literature information and gene blast (blue). All processes were
shown in rectangles and results were highlighted with corresponding background
colors. In the computational pipeline, 51 regulons with assigned motifs and operons
were analyzed sequentially through significant TFBS pairing, DEG conformation, and
TF blast. Only regulons contained DEGs (ten) and had related mapped TF (eight)
were believed to be the final predicted ASR-related regulons (five). These five
regulons were then merged into four and using the corresponding TFs to represent
their names. In the literature validation process, known ASR-related transporters were
first mapped to the MG1363 genome and resulted in 33 genes. Those genes were then
searched in 51 regulons and determine six related regulons. All regulons resulted from
both computational pipeline and literature validation were combined, along with the
information of functional modules, to determine the GRN.

Figure 3. A working model of the transcriptional gene regulatory network in response
to pH change in *L. lactis*. The mechanism is activated by the change of proton signal

in a cell. *RcfB* is assumed to be the overall activator for the rest seven regulons and
controls the ASR functional module solely. Three kinds of literature verified and
significant ASR-related regulons, *llrA*, *llrC* and *hllA*, two insufficient significant
regulon *llrD* (regulon #39) and regulon #8 (llmg_1803) predicted via our workflow
but results under 0.8 motif similarity cutoff or could not find hit, and one putative
significant regulon NHP6A control the seven functional modules which are
experimentally verified in close species MG1363. The other significant regulon *ccpA*
failed to be confirmed by any literature proved genes or transporters. Two extra
functional modules, GAD, and urea degradation show no direct connection to all
seven the regulons. One or more homology genes are found in MG1363 for all the
nine modules using BLAST. The solid arrows indicate regulation between
regulons/TFs and functional modules/genes, and the dashed arrows indicate uncertain
control processes. Besides, two ovals indicate two trans-membrane proteins, one is
confirmed as F0/F1ATPase and the other one with dash line that we still not find its
related information in the public-domain literature.

TABLE LEGEND

Table 1. 14 significant regulons that are verified and mapped to known TFs.
According to analyses, operon numbers and DEG determination (yes or no), matched
template TFs and mapped TFs were assigned for each significant regulon,
respectively, and were aligned based on regulon ID number. Five regulons containing
DEG and having the corresponding TF at the same time were bolded, as which were
being computationally verified regulons to be responsible for acid stress in MG1363.

Regulon ID	No. of operons	DEG	TF template	TF(gene) blast in MG1363
Regulon2	**82**	**Y**	**spo0A**	**_llrC_(llmg_0414)**
Regulon3	32	Y	FoxQ1	N/A
Regulon4	20	Y	SPT2	N/A
Regulon7	**49**	**Y**	**lhfB**	**_hllA_(llmg_0496)**
Regulon10	5	N	GAL80	-(llmg_0271)
Regulon12	**259**	**Y**	**CovR**	**_llrA_(llmg_0908)**
Regulon15	**19**	**Y**	**c4494**	**_ccpA_(llmg_0775)**
Regulon20	79	Y	NHP6A	N/A
Regulon28	5	Y	1Z916	N/A
Regulon31	**65**	**Y**	**ihfA**	**_hllA_(llmg_0496)**
Regulon37	10	N	CovR	_llrA_(llmg_0908)
Regulon40	7	Y	Awh	N/A
Regulon44	12	N	YBR182C	N/A
Regulon47	5	N	RHE_PF00288	_ccpA_(llmg_0775)

Table 2. Known ASR-related gene mapping from literature in response to pH change. Literature-supported ASR-related genes found in close species or other *L. lactis* strains. The template transporters and genes were first identified in published studies from the NCBI and UniProt databases. *L. lactis Il1403* was used as the organism which is very close to MG1363 if template gene existed. Only 36 templates that successfully mapped to the MG1363 genome were listed, which resulted in 33 genes. All mapped genes and corresponding templated were organized by their regulated pathways which were further used as functional modules. Mapped genes were searched in 51 regulons to build the connections between functional modules and regulons.

Template organisms			MG1363	
Organisms	Transporters	Functions/ Pathways	Mapped genes (locus tag)	Regulons
L. lactis	*ldh* *ldhB* *ldhX*	LDH	*ldh* (llmg_1120) *ldhB* (llmg_0392, llmg_0475) *ldhX* (llmg_1429)	NHP6A, *llrA*
L. lactis	*gadB* *gadC*	GAD	*gadB* (llmg_1179) *gadC* (llmg_1178)	N/A
L. lactis	*arcA* *arcB* *arcC1* *arcC2* *argF*	ADI pathway	*arcA* (llmg_2313) *arcB* (llmg_2312) *arcC1* (llmg_2310) *arcC2* (llmg_2309) *argF* (llmg_1754)	NHP6A, *llrA, llrC, hllA*
Bacteria	ureA/B/C$	Urea degradation	*pyrC* (llmg_1508)	N/A
L. lactis	atpEBFHAGDC$$	F0/F1ATPase	llmg_1952, llmg_1951, llmg_1950, llmg_1949, llmg_1948, llmg_1947, llmg_1946, llmg_1945	*llrA*, (Regulon8，llmg_1803)$$$
L. lactis	rcfB	Acid response	rcfB (llmg_2512)	(Regulon39, *llrD*)$$$
L. lactis, *E. coli* K12	*dnak* *groEL* *groES* *grpE* *clpE* *clpC* *clpP*	Chaperone, Protein repair and protease	*dnaK* (llmg_1574) *groEL2* (llmg_0411) *groES* (llmg_0410) *grpE* (llmg_1575) *clpE* (llmg_0528) *clpC* (llmg_0615) *clpP* (llmg_0638)	*llrA*
L. lactis,	*dltC, agK,* *SGP, ffh*	Envelope alterations	llmg_0878	NHP6A, *llrA*

			llmg_0374, llmg_0534,	(Regulon39,
Bacillus subtilis				
L. lactis	*recA, uvr, smn*	DNA repair	llmg_0374, llmg_0534, llmg_1718 llmg_1221	(Regulon39, *llrD*)[$$$]

771 [$]Three subunits of urease enzymes coded by ureABC operon found preserved in
772 multiple bacteria.
773 [$$]Altogether eight genes.
774 [$$$]The homolog prediction or motif research results with low homolog similarity but
775 have meaningful biological relevance.
776
SUPPLEMENTAL MATERIALS

Supplement Number 1: Figure S1 operon_distribution.pptx;

The distribution of operon numbers among 122 clusters.

Supplement Number 2: Figure S2 motif_logos_for_14_significant_regulons.pptx;

The motif logos substracted from 14 significant regulons, used for TFBS comparison.

Supplement Number 3: Dataset S1 NC_009004_operons_and_genes.xlsx;

The operon prediction results for MG1363.

Supplement Number 4: Dataset S2 122_operons_to_clusters.xlsx;

The gene clustering results for microarray "GSE47012" containing operons.

Supplement Number 5: Dataset S3 51_motif-regulons.xlsx;

The summarized motif prediction results for regulon prediction.

Supplement Number 6: Dataset S4 14_verified_regulons.xlsx;

Computational validation results for predicted regulons.

Supplement Number 7: Dataset S5 differential_expression_genes.xlsx;

86 up-regulated genes and 55 down-regulated genes.

Supplement Number 8: Dataset S6 trans-memberane.xlsx;

The prediction results for trans-membrane proteins.