Mapping diversification metrics in macroecological studies: Prospects and

challenges

Julián A. Velasco[1*] & Jesús N. Pinto-Ledezma[2,3]

[1]Museo de Zoología Alfonso L. Herrera, Facultad de Ciencias, Universidad

Nacional Autónoma de México, Mexico City, Mexico.

[2]Department of Ecology, Evolution, and Behavior, University of Minnesota, 1479

Gortner Ave, Saint Paul MN 55108, United States.

[3]Museo de Historia Natural Noel Kempff Mercado, Universidad Autónoma Gabriel

René Moreno, Av. Irala 565, C.C. 2489, Santa Cruz de la Sierra, Bolivia.

*Corresponding author: juvelas@gmail.com

ORCID: https://orcid.org/0000-0002-2183-5758

Abstract

The intersection of macroecology and macroevolution is one of the most active research areas today. Macroecological studies are increasingly using phylogenetic diversification metrics to explore the role of evolutionary processes in shaping present-day patterns of biodiversity. Evolutionary explanations of species richness gradients are key for our understanding of how diversity accumulated in a region. For instance, the present-day diversity in a region can be a result of *in situ* diversification, extinction, or colonization from other regions, or a combination of all of these processes. However, it is unknown whether these metrics capture well these diversification and dispersal processes across geography. Some metrics (e.g., mean root distance -MRD-; lineage diversification-rate -DR-; evolutionary distinctiveness -ED-) seem to provide very similar geographical patterns regardless of how they were calculated (e.g., using branch lengths or not). The lack of appropriate estimates of extinction and dispersal rates in phylogenetic trees can limit our conclusions about how species richness gradients emerged. With a review of the literature and complemented by an empirical comparison, we show that phylogenetic metrics by itself are not capturing well the speciation, extinction and dispersal processes across the geographical gradients. Furthermore, we show how new biogeographic methods can improve our inference of past events and therefore our conclusions about the evolutionary mechanisms driving regional species richness. Finally, we recommend that future studies include several approaches (e.g., spatial diversification modelling, parametric biogeographic methods) to disentangle the relative the role of

40   speciation, extinction and dispersal in the generation and maintenance of species

41   richness gradients.

43   Keywords: Phylogeny, Geography, Spatial diversification, Macroevolution, Species

44   richness, Regional assemblages

Introduction

The causes of spatial variation of biodiversity are one of the most fundamental questions in ecology, biogeography and macroecology (Brown 1995, 2014, Brown and Lomolino 1998, Hawkins et al. 2012, Fine 2015, Jablonski et al. 2017). Current studies are integrating in a single framework the ecological and evolutionary mechanisms driving regional species diversity (McGaughran 2015, Pärtel et al. 2016, Cabral et al. 2017, Leidinger and Cabral 2017). However, only three macroevolutionary processes ultimately can modify the number of species in a region: speciation, extinction and dispersal (Wiens 2011, Fine 2015, Jablonski et al. 2017) (Figure 1). These processes can be modulated by species' traits varying within clades (Paper et al. 2016, 2017, Jezkova and Wiens 2017, Moen and Wiens 2017), age of region (e.g., time-for-speciation effect; Stephens & Wiens, 2003), geographical area (Losos and Schluter 2000), or climatic conditions (Condamine et al. 2013, Lewitus and Morlon 2017).

The integration of different disciplines such as molecular phylogenetics, palaeontology, and historical biogeography have allow to infer a series of macroevolutionary processes across geography (Diniz-Filho et al. 2013, Fritz et al. 2013). It is well-known that fossil information is key to estimate with high confidence rates of speciation and extinction (Sepkoski 1998, Foote 2000, Quental and Marshall 2010, Rabosky 2010b). New methods correcting for sampling bias are able to generate improvements in the estimates of the speciation, extinction and net diversification rates (Silvestro et al. 2014, 2016).

However, the causal mechanisms that underlying the geographical diversity

gradients only can be established with greater confidence for a few taxonomic

groups with adequate fossil record, such as marine bivalves (Jablonski et al. 2006,

2017), mammals (Silvestro et al. 2014) or plants (Antonelli et al. 2015).

As fossil data is not available or incomplete for most extant groups, model-

based approaches used to estimate speciation and extinction rates in

palaeontology were adapted to study the macroevolutionary dynamics using

phylogenetic information (Nee et al. 1994, Morlon et al. 2010, Stadler 2013).

Molecular phylogenies are becoming essential to the study of diversification

dynamics across temporal and spatial scales for extant taxa (Wiens and

Donoghue 2004, Rabosky and Lovette 2008, Stadler 2013, Morlon 2014, Schluter

and Pennell 2017). Therefore, it is possible to reconstruct past diversification

process based on the branching events of a phylogeny using a set of birth-death

models (Nee et al. 1994, Nee 2006, Morlon et al. 2010, Stadler 2013, Morlon

2014). These birth-death models allow infer either a homogeneous process for an

entire clade (Nee et al. 1994, Magallón and Sanderson 2001) or a heterogeneous

process varying in time or in specific subclades of a tree (Paper et al. 2006,

Rabosky and Lovette 2008, Alfaro et al. 2009). However, these birth-death models

only account for temporal variation of the macroevolutionary processes and how

translate these processes to the geography is still a matter of debate.

Macroecological studies use two main approaches to link the estimates of

diversification with the geographical ranges of species (Hawkins et al. 2007, Algar

et al. 2009, Qian et al. 2014, Pinto-Ledezma et al. 2017, Velasco et al. 2018). The

first one uses a set of phylogenetic metrics as a proxy to capture the geographical

signature of lineage diversification dynamics (Diniz-Filho *et al.*, 2013; Fritz *et al.*,

2013; Table 1; Figure 2). These metrics provide either an estimate of a per-species

rate of diversification (e.g., mean root distance –MRD-, residual phylogenetic

diversity –rPD-, mean diversification rate –MDR-), the phylogenetic structure of

regional assemblages (e.g., phylogenetic species variability –PSV-) or the average

age of co-occurring lineages in a given area (e.g., mean age; see Table 1). Each

metric is calculated for each species in the phylogeny; therefore, we can associate

the species' values to its corresponding geographical range and generate a map

with average values for cells or regions. Although these phylogenetic metrics only

account for speciation events, macroecologists have used these maps as a proxy

to test some evolutionary-based hypothesis in macroecological research (Diniz-

Filho *et al.*, 2013; Fritz *et al.*, 2013; see Table 2 for a compendium of these

hypotheses).

The second approach used by macroecologists consists in the explicit

estimation of diversification parameters across geography (Goldberg et al. 2005,

2011, Ramiadantsoa et al. 2017). For instance, the geographic state speciation

and extinction model –GeoSSE (Goldberg et al. 2011; Table 1) allows estimating

speciation, extinction and dispersal parameters across two regions. It is possible

to disentangle the relative role of each one of these processes on the generation

and maintenance of the geographical diversity gradients (Rolland et al. 2014,

Pulido-Santacruz and Weir 2016, Pinto-Ledezma et al. 2017). In addition, a

recently developed Bayesian approach (BAMM; Rabosky 2014, Rabosky et al.

2014) allows to infer the balance of speciation and extinction in the generation of

these biodiversity gradients (Rabosky et al. 2015, Sánchez-Ramírez et al. 2015,

Morinière et al. 2016, Pinto-Ledezma et al. 2017). The BAMM approach allows

both the inference of macroevolutionary dynamics for an entire clade (i.e., a

macroevolutionary regime; Rabosky 2014) and also get estimates of per-species

diversification rates (i.e., as a phylogenetic metric; Rabosky 2016) that can be

mapped in a geographical domain. Although all these methods aim to obtain a

geographical picture of the diversification processes, it remains unexplored if they

can effectively capture these dynamics across regions.

In this paper, we conducted a review on macroecological literature to

evaluate how evolutionary and biogeographic processes contribute to shape

geographical species richness gradients. We review only those papers that make

explicit use of phylogenetic metrics and/or explicit diversification approaches

(Table 1). We divided our review in three main sections. In the first one, we

discuss how studies use phylogenetic metric to test some evolutionary-based

hypotheses underlying geographical diversity gradients and we explore some

limitations of these metrics (see also Table 2). Also, we discuss to what extent

these metrics are able to capture macroevolutionary dynamics in a spatial explicit

context. We illustrate these using three case studies (Furnariides birds, Hylid

frogs, and *Anolis* lizards; Figure 1) and explore how another approaches (e.g.,

diversification modelling and biogeographical approaches) can complement our

inferences about diversification process across geography. In the second section,

we discuss how dispersal and extinction processes are limiting these

diversification inferences and we propose some research avenues to attempt to

solve these problems. Using an explicit biogeographical approach, we test the

role of dispersal on the geographical species richness patterns of the three case

studies. Finally, in the third section, we call for the adoption of complementary

approaches (e.g., extensive simulations, parametric biogeographical methods) in

macroecological research with the aim to evaluate the relative role of speciation,

extinction and dispersal process driving geographical biodiversity gradients.

LITERATURE REVIEW

We conducted a literature search in Web of Science for studies that

explicitly have addressed questions on how speciation, extinction and dispersal

have shaped geographical species richness gradients. We selected those papers

that used either phylogenetic metrics (e.g., mean root distance -MRD-,

phylogenetic diversity -PD-, phylogenetic species variability -PSV-; diversification

rate –DR-; mean Ages; Table 1) or explicit macroevolutionary approaches (e.g.,

GeoSSE, BAMM; Goldberg *et al*., 2011; Rabosky 2014). We compiled a list of 44

papers (Table A1), but we are aware that this likely is not an exhaustive search.

The majority of papers reviewed are testing historical process shaping latitudinal

diversity gradients (LDG) in various taxa.

TESTING EVOLUTIONARY HYPOTHESIS USING PHYLOGENETIC METRICS

AND EXPLICIT DIVERSIFICATION APPROACHES

Here, we discuss how phylogenetic metrics are used to test evolutionary

hypotheses related to the generation and maintenance of geographical diversity.

Several studies used the mean root distance -MRD- metric to evaluate whether

regional assemblages are composed of "basal" or "derivate" linages. First, this

terminology should be avoided because it provides an incorrect interpretation of

the phylogenetic trees (Baum et al. 2005, Crisp and Cook 2005, Omland et al.

2008). Although this metric does not incorporate information from branch lengths

(Algar et al. 2009, Qian et al. 2015), it does provide the average number of nodes

separating each species in a given region from the root of the phylogeny (Kerr and

Currie 1999). MRD therefore provides information about the number of

cladogenetic events (splits) that have occurred through the history of co-occurring

lineages in each region (Pinto-Ledezma et al. 2017, Velasco et al. 2018). Under

this view, MRD should be interpreted as a metric of total diversification (Rabosky

2009), where high MRD values indicating regional assemblages dominated by

extensive cladogenesis and low MRD values indicating assemblages with few

cladogenetic events. A main concern with this metric concern with the fact that it

does not provide any information about what macroevolutionary dynamics have

taken place in a region. For example, it is very hard to establish whether MRD

allows to distinguish between diversity-dependent (Rabosky 2009, Rabosky and

Hurlbert 2015) or time-dependent (Wiens 2011, Harmon and Harrison 2015)

processes dominating regional diversity. Although the distinction between these

two dynamics, and its relationship with the origin and maintenance of regional

diversity, is an intense topic in the macroevolutionary literature (Rabosky 2009,

2013, Wiens 2011, Cornell 2013, Harmon and Harrison 2015, Rabosky and

Hurlbert 2015). However, more empirical and theoretical work is necessary to

establish what scenario plays a significant role in regional species richness

assembly (Rabosky 2012, Etienne et al. 2012, Valente et al. 2015, Graham et al.

2018). It might worth to establish whether local ecological process scaling up to

regional scales or emergent effects (i.e., the existence of a strong equilibrium

process) governed the build-up of regional diversity (Cornell 2013, Harmon and

Harrison 2015, Rabosky and Hurlbert 2015, Marshall and Quental 2016).

The time-for-species effect hypothesis state that the regional build-up of

species richness is directly proportional to the colonization time of its constituent

clades (Stephens and Wiens 2003b; Table 2). However, many phylogenetic

metrics used to test this hypothesis, did not incorporate any age information (Fritz

and Rahbek 2012, Qian et al. 2015). Qian et al. (2015) did some additional

predictions for the time effect hypothesis regarding the phylogenetic structure of

regional assemblages. We suggest that these predictions are not easily deduced

from the original statement of the time-for-speciation effect hypothesis (Stephens

& Wiens 2003). For instance, Qian et al. (2015, p. 7) predicted that regions with

low species richness (e.g., extra-tropical regions) should be composed of more

closely related species than regions with high species richness (e.g., tropical

regions). This assumes that regions with low species richness were colonized

recently and therefore these lineages had little time for speciation. However, it is

also plausible consider that high extinction occurred in these poor species

richness regions by marginal climatic niche conditions preventing adaptive

diversification (Wellborn and Langerhans 2015). By contrast, regions with high

species richness might also be assembled by multiple dispersals from nearby

regions becoming to be a macroevolutionary sink (Goldberg et al. 2005). In this

latter case, the species richness was not build-up by *in situ* speciation mainly but

by continued dispersal through time. To evaluate which of these scenarios is more

plausible it is necessary to adopt an approach that explicitly infer the number of

the dispersal and cladogenetic events across areas (Roy and Goldberg 2007,

Dupin et al. 2017).

Differences in species diversification are also considered as a main driver of

the geographical diversity gradient for many groups (Kennedy et al. 2014, Pinto-

Ledezma et al. 2017). This hypothesis states (Table 2) that differences in net

diversification rates between areas are the main driver of differences in regional

species richness between areas. Davies and Buckley (2011) used the phylogenetic

diversity controlled by species richness (i.e., residual PD –rPD-) to distinguish

areas with different evolutionary processes. These authors predicted that areas

where rapid speciation and low immigration events from other areas occurred, are

dominated by large adaptive radiations (e.g., large islands; Losos and Schluter

2000). By contrast, areas with slow speciation and colonized by multiple lineages

through time should have high values of residual PD.

The "out of the tropics" -OTT- hypothesis (Jablonski et al. 2006; Table 2)

states that latitudinal diversity gradient is due to that the majority of lineages

originated in the tropics and then migrated to extratropical regions. Under this

hypothesis, tropics harbour higher net diversification rates (higher speciation and

lower extinction) than extratropical regions and dispersal rates are higher from the

tropics to extratropical regions than the reverse (Jablonski et al. 2006; Table 1).

For instance, Rolland et al. (2014) used the GeoSSE model to test this hypothesis

in the generation of the latitudinal mammal diversity gradient. They found that net

diversification rates (i.e., the balance of speciation minus extinction) was higher in

tropical than in temperate regions and dispersal rates were higher from the tropics

to temperate regions than the reverse. Also, Pinto-Ledezma et al. (2017) used the

GeoSSE model to test an analogue hypothesis to OTT, as form of Out of the

Forest hypothesis (OTF), using Furnariides birds as a clade model. Their favoured

a model where open areas have higher speciation, extinction and dispersal rates

than forest habitats. All these results suggest that it is reasonable to use either

phylogenetic metrics or explicit diversification approaches (e.g., the GeoSSE

model) to evaluate a set of evolutionary-based hypotheses as a main driver of

geographical diversity gradients. However, we show here (see below) that these

approaches fail to capture the evolutionary and biogeographic processes at

spatial scales.

Are phylogenetic metrics capturing well the diversification process across

geography?

A deep understanding of evolutionary processes affecting regional species

assemblages is coming from the integration of molecular phylogenies and fossil

record (Quental and Marshall 2010, Marshall 2017). From this integration of

neontological and paleontological perspectives, it is clear that both approaches

are necessary to test evolutionary-based hypothesis in macroecological research.

Several hypotheses were proposed to explain geographical diversity patterns,

particularly the latitudinal diversity gradient –LDG- (see Table 2 for a summary and

compilation of the main hypotheses reported in the literature). Although the ideal

approach is to generate robust conclusions from multiple lines of evidence (e.g.,

fossil record, molecular phylogenies, biogeographical inference) it is clear that this

information is scarce for many taxonomic groups. Many macroecological studies

have adopted either phylogenetic metrics or explicit diversification approaches

(e.g., the GeoSSE model) to evaluate the relative contribution of speciation,

extinction and dispersal on the resulting geographical diversity gradients (Table

S1).

Phylogenetic metrics can be easily visualized in a geographical context and

several inferences about ecological (e.g., dispersal) and evolutionary (e.g.,

speciation) process can be done. As these metrics provide a *per-species level*

diversification metric for each species in a phylogeny, it is possible to associate

these values with the corresponding species' geographical range and obtain a

mean value for cells or regions in a given geographical domain (Table 1; Figure 2).

By contrast, explicit diversification approaches (e.g., GeoSSE; BAMM; fitting

models) provide a *per-lineage level diversification* metric for a given clade or a

regional assemblage (Rabosky 2016a). However, in some cases, it is possible to

generate a *per-species level diversification* metric with these approaches. For

instance, Pérez-Escobar et al. (2017) used the function GetTipsRates in

BAMMtools (Rabosky et al. 2014) to map speciation rates for Neotropical orchids.

These two approaches (phylogenetic metrics and lineage diversification)

potentially can provide complementary pictures about how macroevolutionary

dynamics have taken place in the geography. One the one hand, it is possible to

estimate diversification rates for a given clade using the number of species, its

age and a birth-death models (Magallón and Sanderson 2001, Nee 2006,

Sánchez-Reyes et al. 2017). These model-fitting approaches allow to whether

diversity- or time-dependent diversification process has taken place in a regional

assemblage (Etienne et al. 2012, Rabosky 2014, Valente et al. 2015). On the other

hand, per-species diversification rate metrics allow establishing the potential of

each individual species to generate more species (Jetz et al. 2012, Rabosky 2014,

2016a). However, these approaches imply at least a different process, which left a

different signature on the geography. Phylogenetic metrics captures a total

diversification process (Rabosky 2009), whereas lineage diversification

approaches (e.g., BAMM) can potentially provide information about an individual

diversification process (Rabosky 2013). In addition, still is not clear whether

phylogenetic metrics can provide an accurate description of the diversification

dynamics across geography.

The first step to clarify how well these phylogenetic metrics behave is to

establish a comparison within and between different taxonomic groups. To

evaluate how different phylogenetic metrics vary across geography and their

relationship with species richness, we used two empirical data sets from our own

empirical work (furnariid birds and anole lizards; (Pinto-Ledezma et al. 2017,

Velasco et al. 2018) and a data set compiled from several sources (hylid frogs;

(Wiens et al. 2006, Algar et al. 2009, Pyron 2014a). We mapped across geography

five phylogenetic metrics (Table 1, Figure 2). We selected these three data sets

because previous work analysed how evolutionary-based hypotheses affected the

present-day species richness gradient (Wiens et al. 2006, Algar et al. 2009, Pinto-

Ledezma et al. 2017, Velasco et al. 2018).

Figure 2 shows the geographical pattern of species richness and the five

phylogenetic metrics for *Anolis* lizards, hylid frogs and Furnariides birds. For all

clades, there is a higher species concentration near to the Ecuador. Higher

species concentration for hylids and Furnariides can be found in the Amazon and

the Atlantic forest and for *Anolis* lizards in Central America and the Caribe (Figure

2A-C; see also Algar et al. 2009, Pinto-Ledezma et al. 2017, and Velasco et al.

2018, for a detailed description of the geographical species richness pattern for

these clades, respectively). In terms of the geographical pattern of each

phylogenetic metric (Figure 2D-R), in most of the cases cells with higher metric

values are related to cells that contain high species richness and vice versa

(Figure 2D-R; Figure A1). However, the degree and the direction of this

relationship changes according to the phylogenetic metric used. For example,

MRD, a metric of species derivedness, shows a negative correlation with species

richness (Figure 2J-L; Figure A1). Importantly, the spatial relationships between

species richness and phylogenetic metrics found in our analysis could simply be

the result of aggregated species-level attributes within cells or assemblages

(Hawkins et al. 2017). Hence, any conclusion derived from these relationships

needs to be considered carefully. In addition, there are different levels of

correlation between phylogenetic metrics (Figure A1). For example, MDR - MA

present a high but negative correlation, and rPD - PSV and MRD - MDR present a

mid-high positive correlation (Figure A1). Although there are few studies

comparing correlations between metrics (Vellend et al. 2010, Miller et al. 2017), to

our knowledge, none previous study compares the similarity of these

diversification metrics (Table 1). However, some of these metrics sharing

mathematical assumptions, which increase the likelihood of correlation between

them. For example, for ultrametric trees, metrics as MDR could be approximated

by considering the mean root distance (i.e. MRD metric) from the tips to the root

(Freckleton et al. 2008), so further studies exploring the mathematical relation

between metrics are needed.

In order to assess if the cells/assemblages on average do not represent a

random sampling from the species pool, we applied a simple permutation test to

explore the non-randomness in each of the phylogenetic metrics. We applied a

null model where the presence-absence matrix (i.e., PAM) was randomly shuffled

1000 times, but maintaining the frequency of species occurrence and observed

richness in the cells/assemblages (Gotelli 2000). This kind of null model is

standard in studies at the community/assemblage level that use phylogenetic

information (Cavender-Bares et al. 2004, 2006). Interestingly, none of the

phylogenetic metrics deviates from the null expectation for the three clades

(Figure 3). Also, very few cells/assemblages present p-values below the 0.05

threshold, thus indicating that the cells/assemblages present random association

among species (Figure 3). These results should be supported by repeating

analyses with more clades at different spatial extents, but again, we stress that

any result obtained with the use of phylogenetic metrics need to be interpreted

carefully.

A brief comparison between phylogenetic metrics and explicit diversification

and biogeographic approaches

We compared the phylogenetic metrics enunciated in Table 1, which have

been the most used in macroecological research. We explored whether the

geographical patterns of these phylogenetic metrics in three empirical examples

coincide with the macroevolutionary dynamics inferred using explicit modelling

diversification approaches. In particular, we implemented the GeoSSE model to

estimate the three parameters (speciation, extinction, and dispersal) between two

areas in each taxonomic group (Table 3 and 4). In addition, we used the BAMM

approach to generate the per-species level diversification metric implemented in

the software BAMM 2.5.0 (Rabosky 2014). In the following section, we discuss

each metric and we compare them with the explicit diversification approaches.

Residual Phylogenetic Diversity (rPD)

In the case of furnariid birds, we show that forest areas tend to exhibit

slightly higher values of rPD in contrast with open areas (Figure 4; see also Figure

2). According to Davies and Buckley's logic, these areas exhibit slow

diversification and frequent dispersal from open areas. Pinto-Ledezma et al.

(2017) using GeoSSE and BAMM approaches indicated that open areas exhibit

higher net diversification rates than open areas (Table 3). For hylid frogs, we found

that tropical areas tend to exhibit higher rPD values than extratropical regions

(Figure 5). However, by adopting an explicit diversification approach (GeoSSE and

BAMM), we found that net diversification rates were similar in both regions (Table

3). In the case of *Anolis* lizards, the rPD values were higher in the continent than in

the island areas (Figure 6). However, using GeoSSE and BAMM, we found that

both rates were similar (Table 3). In a recent paper, Poe et al. (in press) also found

that macroevolutionary rates are similar between insular and mainland clades. All

these results suggest that rPD likely does not provide an accurate signature of the

macroevolutionary dynamic at spatial scales. In fact, it seems that rPD tends to

overestimate differences between regions when a stationary diversification

process is occurring across geography. A potential solution might be rethinking

the way in which we visualize rPD across geography in contrast with the original

meaning by Davies & Buckley (2011; see also Forest et al. 2007).

Mean root distance (MRD)

As we discussed, *MRD* captures a total diversification value portraying the

number of cladogenetic events co-occurring in a given region. In the case of

furnariid birds, we found that MRD values tend to be higher in open than forest

areas (Figure 4). Accordingly, this metric suggests that more cladogenetic events

were accumulated in open areas (i.e., more total diversification; Rabosky 2009).

Therefore, this metric, for this bird clade, is consistent with results from explicit

diversification approaches (Table 3). For hylid frogs, it seems that there are no

differences in MRD values between extratropics and tropics areas (Figure 5).

However, tropical areas have some cells with very high values. Again, MRD

provide an accurate description of the total diversification pattern in this clade

across the latitudinal gradient. In *Anolis* lizards, we found that MDR values tend to

be lower in islands in comparison with mainland areas (Figure 6). In this case,

MRD did not provide an accurate description of the evolutionary processes

occurring between the mainland and insular anole assemblages. However, there is

also a high probability that the high MRD values in the mainland are a direct reflect

of an idiosyncratic evolutionary trajectory of each one of the two clades that

radiated there (i.e., *Draconura* and *Dactyloa* clade; see Poe et al. 2017, Velasco et

al. 2018). These two clades seems to exhibit differential diversification dynamics

across geography (Velasco et al. 2018) but further research might be necessary to

evaluate these differences.

Phylogenetic species variability (PSV)

The PSV metric provides information about how related are the species in a

given regional assemblage. In hylid frogs, we found that tropical assemblages

tend to be composed of more related species than extratropical assemblages

(Figure 5). Hylid assemblages in extratropical areas are composed of multiple

lineages that dispersed from tropical areas and then diversified there. We found

higher dispersal rates from tropical to temperate regions than vice versa (Table 3

and 4). The same tendency is present in the case of furnariid birds where open

areas exhibit higher PSV values than forest areas (Figure 4) and dispersal rates

were higher from open to forest areas than the reverse (Table 3). By contrast, we

did not find any evidence for differences in PSV values between island and

mainland *Anolis* assemblages (Figure 6). In addition, the dispersal rates were very

low between these two regions (Table 3; Poe et al. 2017). All these results confirm

that the PSV metric can provide some insights about how dispersal process have

shaped regional assemblages. We find evidence that low PSV values (i.e.,

phylogenetically over-dispersed faunas) are influenced by multiple dispersals

along its evolutionary history.

Mean diversification rate (MDR)

Jetz et al. (2012) proposed MDR metric as a species-level speciation rate

metric based in the branch length along the path from the root of a tree to each

individual species. In furnariid birds, we noted that MDR was slightly higher in

open versus forest areas and the same pattern is present using the BAMM

approach (Pinto-Ledezma et al. 2017; Figure 5; Table 3). For hylid frogs,

extratropical regions tend to exhibit higher values than tropical regions (Figure 5).

MDR seems to capture well the differences in macroevolutionary diversification for

these taxa along the latitudinal diversity gradient. A similar pattern is present when

the BAMM approach is used (Table 3). We consider that both metrics (MDR vs

per-species diversification rate from BAMM) leave the same signature in the

geography. In *Anolis*, we found that insular assemblages tend to exhibit higher

MDR values than continental assemblages (Figure 6), However, there is no

difference in the macroevolutionary dynamic between these two areas for the

Anolis lizards clade (Poe et al. in press, Velasco et al. 2018).

Mean ages (MA).

The average of ages of co-occurring lineages are used to test evolutionary

hypothesis about whether a region maintains older lineages than others (e.g., a

museum) or a combination of old and recent lineages (e.g., OTT hypothesis, Table

1). Although this metric does not provide any inference of the ancestral area of the

clade, it is possible to implement an explicit biogeographic approach to test this

(see below). For example, in hylid frogs, we found that extratropical areas are

composed of older lineages than tropical regions (Figure 5). The biogeographic

parametric approach infers this same area as ancestral for the entire lineage

(Figure A2). In furnariid birds, mean ages metric revealed that older lineages have

accumulated more in forest than open areas (Figure 4). In accordance, the

ancestral area inferred with a parametric biogeographic method was the forest

area (Figure A3). In the case of the anole lizards, insular settings tend to be

composed of older lineages than continents. However, the ancestral area for the

entire anole clade is the mainland, particularly South America (Poe et al. 2017).

The mainland *Anolis* radiation is composed of two clades, one clade that

originated in South America (the *Dactyloa* clade; Poe et al. 2017) and colonized

Caribbean islands, and the other clade (the *Norops* clade; Poe et al. 2017) that

originated in the Caribbean islands and then colonized back the mainland in

Middle America and then dispersed to South America. Therefore, the

biogeographical history of the *Anolis* radiation is complex and involves multiple

dispersals between islands and mainland areas (Poe et al. 2017; Figure A4). In

general, mean ages does not provide enough information about the biogeographic

origin and maintenance of a clade. This happens because multiple dispersals and

in situ cladogenesis might erase any simplistic pattern elucidated for this metric,

as found in the case of the *Anolis* lizards.

HOW DISPERSAL AND EXTINCTION AFFECT INFERENCES OF

GEOGRAPHICAL DIVERSIFICATION GRADIENTS?

Dispersal is another key macroevolutionary process that ultimately determines the

number of a species in a region (Roy and Goldberg 2007, Eiserhardt et al. 2013,

Rolland et al. 2014, Chazot et al. 2016). However, few studies evaluated explicitly

how the direction of dispersals between region contributes to the generation of

regional differences between areas (Chown and Gaston 2000, Goldberg et al.

2005, 2011, Jablonski et al. 2006). Roy and Goldberg (2007) showed with

simulations that dispersal asymmetry between areas had a strong impact in the

regional species richness and the average age of these lineages. Accordingly,

phylogenetic metrics can be sensitive to dispersal between areas because it is

impossible to distinguish which lineages originated by *in situ* speciation or simply

due to dispersal from nearby areas. Goldberg et al. (2011) developed the GeoSSE

model to evaluate how range evolution affected diversification rates in a

phylogenetic comparative approach. The GeoSSE model only considers three

states (A: endemic species to a region; B: endemic species to another region; and

AB for widespread species) and makes a series of assumptions that can be

problematic. The first assumption of the GeoSSE model is that a time-dependent

process dominates the diversification dynamic in each region (Stephens and

Wiens 2003, Wiens 2011). This assumption conflicts with a diversity-dependent

process assumption and this debate is far from being resolved (Cornell 2013,

Harmon and Harrison 2015, Rabosky and Hurlbert 2015). The second problematic

assumption has to do with the fact that the GeoSSE model consider dispersal

rates as stable through time and lineages. In other words, the dispersal ability and

therefore the frequency of transitions between areas are constant across the

evolutionary history of a clade. There are many empirical evidence showing that

dispersal rates vary across time and space among lineages (McPeek and Holt

1992, Sanmartín et al. 2008, Robledo-Arnuncio et al. 2014).

Regardless of these major assumptions, the GeoSSE model has been

adopted to evaluate relative contributions of speciation, extinction and dispersal

to the generation of species richness gradients (e.g., Rolland et al. 2014, Pyron

2014b, Staggemeier et al. 2015, Looney et al. 2016, Morinière et al. 2016, Pulido-

Santacruz and Weir 2016, Alves et al. 2017, Hutter et al. 2017, Pinto-Ledezma et

al. 2017). In a recent study, Rabosky and Goldberg (2015) found that state-

dependent diversification models tend to inflate excessively the false discovery

rates (i.e., type I error rates). In particular, Rabosky and Goldberg (2015) found

that these models tend to find false associations between trait shifts and shifts in

macroevolutionary dynamics. Although the Rabosky and Goldberg's study was

not based on the GeoSSE model, it is clear that transitions between areas (i.e.,

dispersal events) can be falsely associated with shifts in speciation and extinction

rates across the phylogeny. Alves et al. (2017) also found that geographical

uncertainties in the assignment of species to a given area affect the parameter

estimates (i.e., speciation, extinction and dispersal rates) in the GeoSSE model.

Same authors also evaluated how incorrect assignments of bat species to tropical

or extra-tropical regions can generate erroneous conclusions about the relative

role of speciation, extinction and dispersal on a latitudinal diversity gradient. From

these studies, it is clear that dispersal is a major issue that needs to be evaluated

explicitly in macroecological studies.

Pulido-Santacruz and Weir (2016) also used the GeoSSE model to disentangle

the relative effect of speciation, extinction and dispersal on the latitudinal bird

diversity gradient. They found that extinction was prevalent across all bird clades

and therefore they suggest this as a main driver of the geographical bird diversity

gradient. Pyron (2014c), also using the GeoSSE model, found that temperate

diversity in reptiles is due to higher extinction in these areas. We consider that

extinction inferences from the GeoSSE model should be treated with caution. For

the few clades where fossil record is abundant (e.g., marine bivalves; Jablonski et

al. 2006), studies point out to conclude that extinction differences between

regions should be treated with caution due to the potential sampling bias

(Jablonski et al. 2006, 2017). In addition, studies based on extensive simulations

found that extinction inferences based only in molecular phylogenies are not

reliable (Rabosky 2010a, 2016b, Quental and Marshall 2010), although extinction

rates can be estimated relatively well using medium to large phylogenies (Beaulieu

& O'Meara 2015).

In a recent review, Sanmartín and Meseguer (2016) proposed that it is possible

to detect the extinction signature in molecular phylogenies using extensive

simulations and lineage-through-time –LTT- plots (see also Antonelli and

Sanmartín 2011). These authors also found that many birth-death models leave a

similar phylogenetic imprint, which make indistinguishable some scenarios. In

addition, extinction events can affect substantially the ancestral range estimates,

and therefore dispersal and extinction parameters in several parametric

biogeographic methods (e.g., Dispersal-Vicariance –DIVA- and Dispersal-

Extinction-Cladogenesis –DEC- models; Ronquist 1997, Ree et al. 2005).

Sanmartín and Meseguer (2016) finally proposed that the adoption of a

hierarchical Bayesian approach using continuous-time Markov Chain models will

allow a better estimation of extinction both in geography and in the phylogeny

(Sanmartín et al. 2008, Sanmartin et al. 2010).

Recently, Rabosky and Goldberg (2017) developed a semi-parametric

method (FiSSE) to correct the statistical problems found in BiSSE models by

themselves in a previous paper (Rabosky and Goldberg 2015). However, the

FiSSE method does not allow the evaluation of the contribution of dispersal on

regional species richness. In any case, the best suitable framework to estimate

relative contributions of speciation, extinction and dispersal might be the GeoSSE

model (or parametric biogeographic models; e.g., Matzke 2014; see below),

although it requires the simulation of a series of null scenarios to evaluate the

statistical power in each case (see Alves et al. 2017, Pinto-Ledezma et al. 2017 for

a few examples). For instance, Pinto-Ledezma et al. (2017) developed a

parametric bootstrapping approach simulating traits to evaluate whether empirical

inferences are different from the simulated. They simulated 100 datasets of neutral

characters along a set of empirical phylogenies and using this new information

repeated the same procedure with empirical data (see Appendix S1 in Pinto-

Ledezma et al. 2017 for details of the bootstrapping approach). This

bootstrapping procedure assumes no direct effect of the geographic character

states on the parameter estimations (Feldman et al. 2016, Pinto-Ledezma et al.

2017).

Finally, it should be clear that more research would be necessary to

establish how extinction affect estimation parameters in state-dependent

diversification approaches (e.g., the GeoSSE model). For instance, the inclusion-

exclusion of extinct species in simulated phylogenies using birth-death models

could substantially affect the geographical inferences of speciation, extinction and

dispersal parameters in the GeoSEE model. This kind of approach might provide

some lights on how to biased can be the parameter estimates with only molecular

phylogenies using the GeoSSE model or any other modeling approach.

Parametric biogeographical approaches in macroecological studies.

The use of parametric biogeographic approaches is an optimal solution to

estimate dispersals across time and space (Matzke 2014, Dupin et al. 2017).

These methods are promising in identifying the relative roles of cladogenetic and

anagenetic processes shaping regional species richness. Recently, Dupin et al.

(2017) developed a biogeographical stochastic mapping to infer the number of

dispersals, and other biogeographical events, in the evolutionary history of

Solanaceae plants across the world. This approach allows the inference from

multiple process including sympatric speciation, allopatric speciation, founder-

event speciation, range expansion (i.e., dispersal without speciation) and local

extinction (i.e., range contractions) based on a time-calibrated phylogenetic tree

and the occurrence of species in geographical regions (see also Matzke 2014 for

more detailed description of the method). These explicit biogeographical

approaches are promising in macroecological studies since they allow to test

simultaneously a set of evolutionary process during the diversification of a clade in

a region (Velasco 2018). In addition, with these new approaches it is possible to

differentiate effectively between macroevolutionary sources and sink areas

(Goldberg et al. 2005, Castroviejo-Fisher et al. 2014, Poe et al. 2017). For

instance, Poe et al. (2017) used a parametric biogeographical approach to

estimate the number of events among regions and distinguish those areas where

many cladogenetic events occurred (i.e., *in situ* speciation) and areas where

almost all its diversity was build-up from extensive colonization of other regions.

The biogeographical stochastic mapping (BSM) method developed by

Dupin et al. (2017) is promising to estimate more accurately the number of

dispersal events between regions based on a better estimation of the ancestral

area for a clade. We evaluated how dispersal rates between regions can affect

inferences drawn only from phylogenetic metrics in our three data sets. We

implemented GeoSSE and BSM approaches for each data set (Table 3 and 4). For

the case of hylid frogs, we counted the inferred number of dispersal events

between tropical and extra-tropical regions in the Americas (Table 3; see also

(Wiens et al. 2006, Algar et al. 2009). For furnariid birds, we counted the number

of dispersal events between open and forest areas (Table 3; see also Pinto-

Ledezma et al. 2017). Finally, for anole lizards, we counted the number of

dispersal events between insular and mainland areas (Table 3; see also Algar and

Losos 2011, Poe et al. 2017, Velasco et al. 2018).

Using biogeographical stochastic mapping –BSM-, we inferred the number

of dispersal events from one region to another for each one of the three

taxonomic groups examined (Table 3). The BSM approach allows us to

disentangle which dispersals were only range expansions and which dispersals

generated a speciation event (i.e., a founder-event speciation; (Barton and

Charlesworth 1984, Templeton 2008). For furnariid birds, we found that range

expansions were three times higher from forest areas to open areas than the

reverse and founder events were twice higher from forest to open areas than the

opposite (Table 3). This result suggests that differences in species richness

between forest and open areas are due by recurrent dispersal events along the

furnariid diversification history (Pinto-Ledezma et al. 2017). Pinto-Ledezma et al.

(2017) found a similar result using the GeoSSE approach, but they conducted a

parametric simulation approach to evaluate whether there was a direct effect of

the geographic location on the parameter estimates. Their results show that the

GeoSSE approach, in this case, had limited power to detect a signature of

geographic region on speciation, extinction and dispersal rates. With the

implementation of the BSM approach here, we corroborate Pinto-Ledezma et al.'s

findings with improved statistical power. In the case of hylid frogs and the

transitions between tropical and extra-tropical areas, we found that the BSM

approach inferred more dispersal events from tropical to extra-tropical regions

(Table 3 and 4). However, the number of founder events was relatively low in

comparison with range expansions (Table 3). These results suggest that few

dispersal events have occurred across the diversification of hylid frogs and

corroborate that the species richness in each region largely originated by *in situ*

speciation modulated by climatic factors (Wiens et al. 2006, Algar et al. 2009).

Finally, for *Anolis* lizards, we found that dispersal events between insular and

mainland regions were relatively low (Table 3 and 4). We did not find evidence of

any expansion range events from mainland to island or vice versa. This also

corroborates previous findings that evolutionary radiation of anole in insular and

mainland settings is due to extensive *in situ* diversification (Poe et al. in press,

2017, Algar and Losos 2011).

These results point out that the BSM approach (Dupin et al. 2017) is a

promising approach when we are interested in testing the role of anagenetic and

cladogenetic events on the resulting geographical species richness gradients.

Although parametric biogeographic approaches are still in their infancy (Sanmartín

2012, Matzke 2014, Dupin et al. 2017), these methods allow us to evaluate

macroevolutionary dynamics (i.e., speciation and extinction) in an explicit

geographical context. These methods are statistical powerful and make use of a

series of explicit geographic range evolution models (Velasco 2018).

TOWARD AN INTEGRATION OF BIOGEOGRAPHICAL AND SPECIES

DIVERSIFICATION APPROACHES IN MACROECOLOGICAL STUDIES.

Although different parametric biogeographic methods have been

developing at least for the last 20 years (Ronquist 1997, Ree et al. 2005, Landis et

al. 2013, Matzke 2014, Dupin et al. 2017), the adoption of these methods to test

evolutionary-based hypotheses underlying geographical diversity gradient has

been rare. For instance, few studies examined here tested the effect of dispersal

events in the generation of regional species richness assemblages. It should clear

that the current paradigm in biogeography makes a call for an evaluation of the

relative frequency of cladogenetic and anagenetic process during the

biogeographical history of lineages. The adoption of parametric approaches in

future macroecological studies will contribute to an improvement of the estimation

of speciation, extinction and dispersal processes as drivers of the geographical

diversity gradients. In addition, we also think that it is necessary to establish which

macroevolutionary dynamics govern regional assemblages. Phylogenetic

approaches based on fitting diversification models help to test whether regional

species richness is due to diversity dependence (i.e., ecological limits), time

dependence, or environmental factors (Rabosky and Lovette 2008, Etienne et al.

2012, Etienne and Haegeman 2012, Condamine et al. 2013). We also stress that

the adoption of many approaches providing multiple lines of evidence will help to

disentangle the evolutionary and ecological causes of biodiversity gradients.

Some recent studies have pointed toward this strategy and have begun to provide

evidence from many lines to understand how evolutionary processes underlying

species richness gradients works (Hutter et al. 2017, Pinto-Ledezma et al. 2017).

Conclusions and recommendations

The resulting geographical pattern of several phylogenetic metrics did not

provide any robust evidence of a spatially explicit diversification dynamic. As we

have shown, these resulting geographical patterns did not differ from that

generated by a simple null model. It is hard to untangle causal mechanisms (i.e.,

speciation, extinction, and dispersal) from only the geographical signature that

these metrics attempt to capture. We recommend that phylogenetic metrics

should be used only to visualize geographical patterns of total diversification (e.g.,

MRD, residual PD; MDR), phylogenetic structure (e.g., PSV), or mean ages of co-

distributed species (e.g., MA) (Table 1). We suggest that conclusions about the

role of evolutionary processes in the generation and maintenance of species

richness gradients based only in these phylogenetic metrics should be avoided

and additional approaches always should be used.

Some explicit diversification approaches (e.g., model fitting approaches;

(Etienne et al. 2012, Rabosky 2014, Valente et al. 2015) are useful to establish the

macroevolutionary dynamics operating at regional scales. Although some

approaches (e.g, the GeoSSE model) allow us to evaluate the relative role of the

ultimate process that modify the regional species diversity, its statistical power

(e.g., high Type I errors) has been challenged by simulation and empirical studies.

Furthermore, the extinction and dispersal estimates inferred by the GeoSSE model

tend to be unbiased. Parametric biogeographic approaches are becoming a

standard tool to evaluate how evolutionary processes can explain the

geographical distribution of extant taxa. These approaches are promising and

should be extensively used because allow us to estimate the relative frequency of

cladogenetic and anagenetic process shaping the regional species richness.

It is necessary that macroecological studies use a combination of explicit

diversification approaches and parameter biogeographic methods with the aim to

clarify how evolutionary process have shaped regional species richness

assemblages. As Jablonski et al. (2017) have outlined, one of the main obstacles

to generate an appropriate understanding of the causal mechanisms underlying

geographical diversity gradients has been that many studies have tested a single

hypothesis, either evolutionary or ecological, as an explanatory factor. We suggest

that ecological and evolutionary hypotheses should be tested simultaneously to

explain the relative contribution of each process to the regional diversity. As

shown by our empirical comparison of phylogenetic metrics, explicit

diversification models, and historical biogeographic methods have showed, it is

necessary to obtain evidence of different approaches to guarantee sound

conclusions about the evolutionary causes of these biodiversity gradients.

Acknowledgments

We thank Josep Padullés for reviewing the English. JAV is supported by a

Postdoctoral fellowship from DGAPA program at Facultad de Ciencias of the

UNAM. JAV thanks to Oscar Flores Villela for providing a working space at Museo

of Zoologia "Alfonso L. Herrera" and insightful discussions about historical

biogeography and evolutionary biology. JNPL is supported by a Postdoctoral

research fellowship from the Grand Challenge in Biology Postdoctoral Program,

College of Biological Sciences, University of Minnesota.

Literature cited

Alfaro, M. E. et al. 2009. Nine exceptional radiations plus high turnover explain

species diversity in jawed vertebrates. - Proc. Natl. Acad. Sci. U. S. A. 106:

13410–4.

Algar, A. C. and Losos, J. B. 2011. Evolutionary assembly of island faunas

reverses the classic island – mainland richness difference in Anolis lizards.:

1125–1137.

Algar, A. C. et al. 2009. Evolutionary constraints on regional faunas: whom, but

not how many. - Ecol. Lett.: 57–65.

Alves, D. M. C. C. et al. 2017. Geographical diversification and the effect of model

and data inadequacies: The bat diversity gradient as a case study. - Biol. J.

Linn. Soc. 121: 894–906.

Antonelli, A. and Sanmartín, I. 2011. Mass extinction, gradual cooling, or rapid

radiation? Reconstructing the spatiotemporal evolution of the ancient

728   angiosperm genus Hedyosmum (Chloranthaceae) using empirical and

729   simulated approaches. - Syst. Biol. 60: 596–615.

730 Antonelli, A. et al. 2015. An engine for global plant diversity: highest evolutionary

731   turnover and emigration in the American tropics. - Front. Genet. 6: 1–14.

732 Barton, N. H. and Charlesworth, B. 1984. Genetic Revolutions, Founder Effects,

733   and Speciation. - Annu. Rev. Ecol. Syst. 15: 133–164.

734 Baum, D. A. et al. 2005. The tree-thinking challenge. - Science (80-.). 310: 979–

735   980.

736 Brown, J. 1995. Macroecology. - University of Chicago Press.

737 Brown, J. H. 2014. Why are there so many species in the tropics? - J. Biogeogr.

738   41: 8–22.

739 Brown, J. H. and Lomolino, M. V. 1998. Biogeography. - Sinauer Associates, Inc.

740 Buckley, L. B. et al. 2010. Phylogeny, niche conservatism and the latitudinal

741   diversity gradient in mammals. - Proc. Biol. Sci. 277: 2131–8.

742 Cabral, J. S. et al. 2017. Mechanistic simulation models in macroecology and

743   biogeography: state-of-art and prospects. - Ecography (Cop.). 40: 1–14.

744 Castroviejo-Fisher, S. et al. 2014. Neotropical diversification seen through

745   glassfrogs (M Ebach, Ed.). - J. Biogeogr. 41: 66–80.

746 Cavender-Bares, J. et al. 2004. Phylogenetic Overdispersion in Floridian Oak

747   Communities. - Am. Nat. 163: 823–843.

748 Cavender-Bares, J. et al. 2006. Phylogenetic structure of floridian plant

749   communities depends on taxonomic and spatial scale. - Ecology 87: 109–

750   122.

Chazot, N. et al. 2016. Into the Andes: multiple independent colonizations drive

montane diversity in the Neotropical clearwing butterflies Godyridina. - Mol.

Ecol. 25: 5765–5784.

Chown, S. L. and Gaston, K. J. 2000. Areas cradles and museums: The latitudinal

gradient in species richness. - Trends Ecol. Evol. 15: 311–315.

Condamine, F. L. et al. 2013. Macroevolutionary perspectives to environmental

change. - Ecol. Lett. 16 Suppl 1: 72–85.

Cornell, H. V 2013. Is regional species diversity bounded or unbounded? - Biol.

Rev. Camb. Philos. Soc. 88: 140–65.

Crisp, M. D. and Cook, L. G. 2005. Do early branching lineages signify ancestral

traits? - Trends Ecol. Evol. 20: 122–128.

Davies, T. J. and Buckley, L. B. 2011. Phylogenetic diversity as a window into the

evolutionary and biogeographic histories of present-day richness gradients

for mammals. - Philos. Trans. R. Soc. Lond. B. Biol. Sci. 366: 2414–25.

Diniz-Filho, J. A. F. et al. 2013. Evolutionary macroecology. - Front. Biogeogr. 5:

1–11.

Dupin, J. et al. 2017. Bayesian estimation of the global biogeographical history of

the Solanaceae. - J. Biogeogr. 44: 887–899.

Eiserhardt, W. L. et al. 2013. Dispersal and niche evolution jointly shape the

geographic turnover of phylogenetic clades across continents. - Sci. Rep. 3:

1164.

Etienne, R. S. and Haegeman, B. 2012. A Conceptual and Statistical Framework

for Adaptive Radiations with a Key Role for Diversity Dependence. - Am. Nat.

180: E75–E89.

Etienne, R. S. et al. 2012. Diversity-dependence brings molecular phylogenies

closer to agreement with the fossil record. - Proc. Biol. Sci. 279: 1300–9.

Faith, D. P. 1992. Conservation evaluation and phylogenetic diversity. - Biol.

Conserv. 61: 1–10.

Feldman, A. et al. 2016. Body sizes and diversification rates of lizards, snakes,

amphisbaenians and the tuatara. - Glob. Ecol. Biogeogr. 25: 187–197.

Fine, P. V. A. 2015. Ecological and Evolutionary Drivers of Geographic Variation in

Species Diversity. - Annu. Rev. Ecol. Evol. Syst. 46: 369–392.

Foote, M. 2000. Origination and extinction components of taxonomic diversity:

general problems. - Paleobiology 26: 74–102.

Forest, F. et al. 2007. Preserving the evolutionary potential of floras in biodiversity

hotspots. - Nature 445: 757–760.

Fritz, S. a. and Rahbek, C. 2012. Global patterns of amphibian phylogenetic

diversity. - J. Biogeogr. 39: 1373–1382.

Fritz, S. a et al. 2013. Diversity in time and space: wanted dead and alive. - Trends

Ecol. Evol. 28: 509–16.

Goldberg, E. E. et al. 2005. Diversity, Endemism, and Age Distributions in

Macroevolutionary Sources and Sinks. - Am. Nat. 165: 623–633.

Goldberg, E. E. et al. 2011. Phylogenetic inference of reciprocal effects between

geographic range evolution and diversification. - Syst. Biol. 60: 451–65.

Gotelli, N. J. 2000. Null Model Analysis of Species Co-Occurrence Patterns. -

Ecology 81: 2606.

Graham, C. H. et al. 2018. Phylogenetic scale in ecology and evolution. - Glob.

Ecol. Biogeogr. 27: 175–187.

Harmon, L. J. and Harrison, S. 2015. Species Diversity Is Dynamic and

Unbounded at Local and Continental Scales. - Am. Nat. 185: 584–593.

Hawkins, B. a et al. 2007. Climate, niche conservatism, and the global bird

diversity gradient. - Am. Nat. 170 Suppl: S16-27.

Hawkins, B. a. et al. 2012. Different evolutionary histories underlie congruent

species richness gradients of birds and mammals. - J. Biogeogr. 39: 825–

841.

Hawkins, B. A. et al. 2017. Structural bias in aggregated species-level variables

driven by repeated species co-occurrences: a pervasive problem in

community and assemblage data. - J. Biogeogr. 44: 1199–1211.

Helmus, M. R. et al. 2007. Phylogenetic measures of biodiversity. - Am. Nat. 169:

E68-83.

Hutter, C. R. et al. 2017. Rapid Diversification and Time Explain Amphibian

Richness at Different Scales in the Tropical Andes, Earth's Most Biodiverse

Hotspot. - Am. Nat. 190: 000–000.

Jablonski, D. et al. 2006. Out of the tropics: Evolutionary dynamics of the

latitudinal diversity gradient. - Science (80-.). 314: 102–106.

Jablonski, D. et al. 2017. Shaping the Latitudinal Diversity Gradient: New

Perspectives from a Synthesis of Paleobiology and Biogeography. - Am. Nat.

189: 1–12.

Jetz, W. et al. 2012. The global diversity of birds in space and time. - Nature 491:

444–448.

Jezkova, T. and Wiens, J. J. 2017. What Explains Patterns of Diversification and

Richness among Animal Phyla? - Am. Nat. 189: 000–000.

Kennedy, J. D. et al. 2014. Into and out of the tropics: The generation of the

latitudinal gradient among New World passerine birds. - J. Biogeogr. 41:

1746–1757.

Kerr, J. T. and Currie, D. J. 1999. The relative importance of evolutionary and

environmental controls on broad-scale patterns of species richness in North

America. - Ecoscience 6: 329–337.

Landis, M. J. et al. 2013. Bayesian analysis of biogeography when the number of

areas is large. - Syst. Biol. 62: 789–804.

Latham, R. E. and Ricklefs, R. E. 1993. Global Patterns of Tree Species Richness

in Moist Forests: Energy-Diversity Theory Does Not Account for Variation in

Species Richness. - Oikos 67: 325–333.

Leidinger, L. and Cabral, J. S. 2017. Biodiversity dynamics on Islands: Explicitly

accounting for causality in mechanistic models. - Diversity 9: 30.

Lewitus, E. and Morlon, H. 2017. Detecting environment-dependent diversification

from phylogenies: a simulation study and some empirical illustrations. - Syst.

Biol. in press.

Looney, B. P. et al. 2016. Into and out of the tropics: Global diversification

patterns in a hyperdiverse clade of ectomycorrhizal fungi. - Mol. Ecol. 25:

630–647.

Losos, J. B. and Schluter, D. 2000. Analysis of an evolutionary species-area

relationship. - Nature 408: 847–50.

Magallón, S. and Sanderson, M. J. 2001. Absolute diversification rates in

angiosperm clades. - Evolution 55: 1762–80.

Marshall, C. R. 2017. Five palaeobiological laws needed to understand the

evolution of the living biota. - Nat. Ecol. Evol. 1: 165.

Marshall, C. R. and Quental, T. B. 2016. The uncertain role of diversity

dependence in species diversification and the need to incorporate time-

varying carrying capacities. - Philos. Trans. R. Soc. B Biol. Sci. 371:

20150217.

Matzke, N. J. 2014. Model selection in historical biogeography reveals that

founder-event speciation is a crucial process in island clades. - Syst. Biol. 63:

951–970.

McGaughran, A. 2015. Integrating a Population Genomics Focus into

Biogeographic and Macroecological Research. - Front. Ecol. Evol. 3: 1–5.

McPeek, M. A. and Holt, R. D. 1992. The Evolution of Dispersal in Spatially and

Temporally Varying Environments. - Am. Nat. 140: 1010–1027.

Miller, E. T. et al. 2017. Phylogenetic community structure metrics and null

models: a review with new methods and software. - Ecography (Cop.). 40:

461–477.

Moen, D. S. and Wiens, J. J. 2017. Microhabitat and Climatic Niche Change

Explain Patterns of Diversification among Frog Families. - Am. Nat. 190: 29–

44.

Morinière, J. et al. 2016. Phylogenetic niche conservatism explains an inverse

latitudinal diversity gradient in freshwater arthropods. - Sci. Rep. 6: 1–12.

Morlon, H. 2014. Phylogenetic approaches for studying diversification. - Ecol.

Lett. 17: 508–25.

Morlon, H. et al. 2010. Inferring the dynamics of diversification: a coalescent

approach. - PLoS Biol. in press.

Nee, S. 2006. Birth-Death Models in Macroevolution. - Annu. Rev. Ecol. Evol.

Syst. 37: 1–17.

Nee, S. et al. 1994. The reconstructed evolutionary process. - Philos. Trans. R.

Soc. Lond. B. Biol. Sci. 344: 305–11.

Omland, K. E. et al. 2008. Tree thinking for all biology: the problem with reading

phylogenies as ladders of progress. - BioEssays 30: 854–67.

Paper, S. et al. 2006. Likelihood Methods for Detecting Temporal Shifts in

Diversification Rates. - Evolution (N. Y). 60: 1152.

Paper, S. et al. 2016. Past, future, and present of state-dependent models of

diversification. - Am. J. Bot. 103: 792–795.

Paper, S. et al. 2017. Accelerated body size evolution during cold climatic periods

in the Cenozoic. - Proc. Natl. Acad. Sci. 114: 4183–4188.

Pärtel, M. et al. 2016. Macroecology of biodiversity: disentangling local and

regional effects. - New Phytol. 211: 404–410.

Pérez-Escobar, O. A. et al. 2017. Recent origin and rapid speciation of Neotropical

orchids in the world's richest plant biodiversity hotspot. - New Phytol. 215:

891–905.

Pinto-Ledezma, J. N. et al. 2017. The geographical diversification of Furnariides:

the role of forest versus open habitats in driving species richness gradients. -

J. Biogeogr. 44: 1683–1693.

Poe, S. et al. Comparative evolution of an archetypal adaptive radiation:

Innovation and opportunity in Anolis lizards. - Am. Nat. in press.

Poe, S. et al. 2017. A Phylogenetic, Biogeographic, and Taxonomic study of all

Extant Species of Anolis (Squamata; Iguanidae). - Syst. Biol. 0: 1–35.

Pulido-Santacruz, P. and Weir, J. T. 2016. Extinction as a driver of avian latitudinal

diversity gradients. - Evolution (N. Y). 70: 860–872.

Pyron, R. A. 2014a. Biogeographic analysis reveals ancient continental vicariance

and recent oceanic dispersal in amphibians. - Syst. Biol. 63: 779–797.

Pyron, R. A. 2014b. Biogeographic Analysis Reveals Ancient Continental

Vicariance and Recent Oceanic Dispersal in Amphibians. - Syst. Biol. 63:

779–797.

Pyron, R. A. 2014c. Temperate extinction in squamate reptiles and the roots of

latitudinal diversity gradients. - Glob. Ecol. Biogeogr. 23: 1126–1134.

Qian, H. et al. 2014. Evolutionary and ecological causes of species richness

patterns in North American angiosperm trees. - Ecography (Cop.).: 1–10.

Qian, H. et al. 2015. Evolutionary and ecological causes of species richness

patterns in North American angiosperm trees. - Ecography (Cop.). 38: 241–

250.

Quental, T. B. and Marshall, C. R. 2010. Diversity dynamics: molecular

phylogenies need the fossil record. - Trends Ecol. Evol. 25: 434–41.

Rabosky, D. 2009. Ecological limits and diversification rate: alternative paradigms

to explain the variation in species richness among clades and regions. - Ecol.

Lett. 12: 735–743.

Rabosky, D. L. 2010a. Extinction rates should not be estimated from molecular

phylogenies. - Evolution 64: 1816–24.

Rabosky, D. L. 2010b. Primary controls on species richness in higher taxa. - Syst.

Biol. 59: 634–45.

Rabosky, D. L. 2012. Testing the time-for-speciation effect in the assembly of

regional biotas. - Methods Ecol. Evol. 3: 224–233.

Rabosky, D. L. 2013. Diversity-Dependence, Ecological Speciation, and the Role

of Competition in Macroevolution. - Annu. Rev. Ecol. Evol. Syst. 44: 481–502.

Rabosky, D. L. 2014. Automatic detection of key innovations, rate shifts, and

diversity-dependence on phylogenetic trees. - PLoS One in press.

Rabosky, D. L. 2016a. Reproductive isolation and the causes of speciation rate

variation in nature. - Biol. J. Linn. Soc. 118: 13–25.

Rabosky, D. L. 2016b. Challenges in the estimation of extinction from molecular

phylogenies: A response to Beaulieu and O'Meara. - Evolution (N. Y). 70:

218–228.

Rabosky, D. L. and Lovette, I. J. 2008. Density-dependent diversification in North

American wood warblers. - Proc. Biol. Sci. 275: 2363–71.

Rabosky, D. L. and Hurlbert, A. H. 2015. Species Richness at Continental Scales

Is Dominated by Ecological Limits. - Am. Nat. 185: 572–583.

Rabosky, D. L. and Goldberg, E. E. 2015. Model Inadequacy and Mistaken

Inferences of Trait-Dependent Speciation. - Syst. Biol. 64: 340–355.

Rabosky, D. L. and Goldberg, E. E. 2017. FiSSE: A simple nonparametric test for

the effects of a binary character on lineage diversification rates. - Evolution

(N. Y). 71: 1432–1442.

Rabosky, D. L. et al. 2014. BAMMtools: An R package for the analysis of

evolutionary dynamics on phylogenetic trees (S Kembel, Ed.). - Methods Ecol.

Evol. 5: 701–707.

Rabosky, D. L. et al. 2015. Minimal effects of latitude on present-day speciation

rates in New World birds. - Proc. R. Soc. B Biol. Sci. 282: 20142889.

Ramiadantsoa, T. et al. 2017. Phylogenetic Comparative Method for Geographical

Radiation. - Ann. Zool. Fennici 54: 237–257.

Ree, R. H. et al. 2005. A likelihood framework for inferring the evolution of

geographic range on phylogenetic trees. - Evolution (N. Y). 59: 2299–311.

Robledo-Arnuncio, J. J. et al. 2014. Space, time and complexity in plant dispersal

ecology. - Mov. Ecol. 2: 1–17.

Rolland, J. et al. 2014. Faster speciation and reduced extinction in the tropics

contribute to the mammalian latitudinal diversity gradient. - PLoS Biol. 12:

e1001775.

Ronquist, F. 1997. Dispersal-vicariance analysis: a new approach to the

quantification of historical biogeography. - Syst. Bot. 46: 195–203.

Roy, K. and Goldberg, E. E. 2007. Origination, Extinction, and Dispersal:

Integrative Models for Understanding Present-Day Diversity Gradients. - Am.

Nat. 170: S71–S85.

Sánchez-Ramírez, S. et al. 2015. In and out of refugia: Historical patterns of

diversity and demography in the North American Caesar's mushroom species

complex. - Mol. Ecol. 24: 5938–5956.

Sánchez-Reyes, L. L. et al. 2017. Uncovering higher-taxon diversification

dynamics from clade age and species-richness data. - Syst. Biol. 66: 367–

378.

Sanmartin, I. et al. 2010. Bayesian island biogeography in a continental setting:

the Rand Flora case. - Biol. Lett. 6: 703–707.

Sanmartín, I. 2012. Historical Biogeography: Evolution in Time and Space. - Evol.

Educ. Outreach 5: 555–568.

Sanmartín, I. and Meseguer, A. S. 2016. Extinction in phylogenetics and

biogeography: From timetrees to patterns of biotic assemblage. - Front.

Genet. 7: 1–17.

Sanmartín, I. et al. 2008. Inferring dispersal: a Bayesian approach to phylogeny-

based island biogeography, with special reference to the Canary Islands. - J.

Biogeogr. 35: 428–449.

Schluter, D. and Pennell, M. W. 2017. Speciation gradients and the distribution of

biodiversity. - Nature 546: 48–55.

Sepkoski, J. J. 1998. Rates of speciation in the fossil record. - Philos. Trans. R.

Soc. Lond. B. Biol. Sci. 353: 315–26.

Silvestro, D. et al. 2014. Bayesian estimation of speciation and extinction from

incomplete fossil occurrence data. - Syst. Biol. 63: 349–367.

Silvestro, D. et al. 2016. Fossil biogeography: a new model to infer dispersal,

extinction and sampling from palaeontological data. - Philos. Trans. R. Soc. B

Biol. Sci. 371: 20150225.

Stadler, T. 2013. Recovering speciation and extinction dynamics based on

phylogenies. 26: 1203–1219.

Staggemeier, V. G. et al. 2015. Phylogenetic analysis in Myrcia section Aulomyrcia

and inferences on plant diversity in the Atlantic rainforest. - Ann. Bot. 115:

747–761.

Stephens, P. R. and Wiens, J. J. 2003. Explaining species richness from

continents to communities: the time-for-speciation effect in emydid turtles. -

Am. Nat. 161: 112–28.

Templeton, A. R. 2008. The reality and importance of founder speciation in

evolution. - BioEssays 30: 470–479.

Valente, L. M. et al. 2015. Equilibrium and non-equilibrium dynamics

simultaneously operate in the Galápagos islands. - Ecol. Lett. 18: 844–852.

Velasco, J. A. 2018. Are Historical Biogeographical Events Able to Promote

Biological Diversification? - In: Levente, H. (ed), Pure and Applied

Biogeography. InTechOpen, pp. 15–30.

Velasco, J. A. et al. 2018. Climatic and evolutionary factors shaping geographical

gradients of species richness in Anolis lizards. - Biol. J. Linn. Soc. xx: 1–13.

Vellend, M. et al. 2010. Measuring phylogenetic biodiversity. - Front. Meas. Biol.

Divers.: 194–207.

Wellborn, G. A. and Langerhans, R. B. 2015. Ecological opportunity and the

adaptive diversification of lineages. - Ecol. Evol. 5: 176–195.

Wiens, J. J. 2011. The Causes Of Species Richness Patterns Across Space, Time,

And Clades And The Role Of "Ecological Limits." - Q. Rev. Biol. 86: 75–96.

Wiens, J. J. and Donoghue, M. J. 2004. Historical biogeography, ecology and

species richness. - Trends Ecol. Evol. 19: 639–44.

Wiens, J. J. and Graham, C. H. 2005. Niche conservatism: Integrating evolution,

ecology, and conservation biology. - Annu. Rev. Ecol. Evol. Syst. 36: 519–

539.

Wiens, J. J. et al. 2006. Evolutionary and Ecological Causes of the Latitudinal

Diversity Gradient in Hylid Frogs: Treefrog Trees Unearth the Roots of High

Tropical Diversity. - Am. Nat. 168: 579–596.

1013

1014 Table 1: Phylogenetic metrics and explicit diversification approaches used in
1015 macroecological studies to address evolutionary questions related with the
1016 geographical diversity gradients.

Metric	Author	Description	Software / R package
MRD	Kerr and Currie 1999	MRD is calculated by counting the number of nodes separating each terminal species in a regional assemblage or cell from the tips to root of the phylogenetic tree. This metric does not need that trees be ultrametric or have branch lengths.	metricTester
PD (residual)	Faith 1992	PD is calcultated by summing all the branch lengths of species co-occurring in a regional assemblage or cell. Residual PD is obtained from an ordinary least square regression between PD and species richness.	picante, metricTester, pez
PSV	Helmus et al. 2007, Algar et al. 2009	PSV is calculated from a matrix where their diagonal elements provide the evolutionary divergence (based on the branch lengths) of each terminal species from the root to the tips of the tree, and the off-diagonal elements provide the degree of shared evolutionary history among species. Values close to zero indicates that all species in a regional assemblage or cell are very close related whereas values close to one indicate that species are not related.	picante, metricTester, pez
mean DR	Jetz et al. 2012	DR is calculated as the inverse of a measure of evolutionary isolation (Redding & Mooers 2006) which sum all the edge lengths from a species to the root of the tree. The inverse of this evolutionary isolation metric therefore capture the level of splitting rate of each species (i.e., its path to a top).	FiSSE
Mean age	Latham and Ricklefs 1993	The mean age of co-occurring species in a regional assemblage or cell simply is calculated tallying the age of each most recent common ancestor (MRCA) for each species and the averaged.	None
GeoSSE	Goldberg et al. 2011	The geographic state speciation and extinction -GeoSSE- model is a trait-dependent diversification method linking geographic occurrence with diversification rates. These method allow to infer both speciation and extinction rates as movement (dispersal) rates among two regions.	Diversitree R package
BAMM	Rabosky 2014	BAMM is a method that attempt to identify whether a phylogeny exhibit a single or various macroevolutionary regimes (i.e., different diversification dynamics). As speciation, extinction and net diversification rates are considered to be heterogeneous across the phylogeny it is possible to estimate a rate for each branch or species in the tree.	BAMM software and BAMMtools R package

1017

1018 Table 2. Description of some evolutionary hypothesis tested in macroecological studies as causal mechanisms of
1019 regional species richness.

Hypothesis	References	Description	Predictions	Metrics and/or methods used to test	Limitations
Phylogenetic niche conservatism (PNC)	Wiens and Graham 2005	Phylogenetic niche conservatism is the tendency of related species to inherit niche requirements from its the most recent common ancestors (Wiens & Graham 2005).	PNC predicts that regions where a clade originated will accumulate more species simply due to more occupation time and diversification rates tend to be similar between regions. The tropical niche conservatism hypothesis (TNC; Wiens and Donoghue 2004) is based on PNC to explain differences in species richness in tropical and temperate regions.	MRD, Mean age, GeoSSE, BAMM	1) MRD metric fails to capture spatially dynamics of the balance of speciation and extinction and it is very hard to establish whether species richness in a region is only generated by higher speciation rates. Furthermore, MRD does not capture dispersal dynamic across regions and species richness in a given region can be generated from only dispersals from nearby regions (e.g., macroevolutionary sinks; CITA). 2) Mean age provide partial is able to test the role of PNC on geographical species richness because only it is possible to establish which regions have, in average, old clades and this not reflects whether many speciation events occurred there. 3) GeoSSE is potentially the only one approach that allow to disentangle these three process but it is only limited to two regions (e.g., tropical vs. temperate). In addition, GeoSSE has been criticized due its low statistical power (see Rabosky and Goldberg 2015).
Regional diversification (RD)	Buckley et al. 2010	Differences in the balance of speciation and extinction across geography can explain differences in species richness between regions.	RD predicts that regions with striking differences in species richness are due to differences in macroevolutionary dynamics between regions.	residual PD, GeoSSE, BAMM	1) Residual PD can be used to discriminate regions with rapid and slow diversification based on the expected phylogenetic diversity given species richness (Buckley et al. 2010). However, this metric ignores the contribution to dispersal to PD in a given region or cell. 2) GeoSSE can estimate speciation, extinction and dispersal rates between regions but again is limited to two regions. 3) BAMM potentially could be used to estimate speciation rates for regional clades but this method is unable to estimate dispersal rates between regions.

Out of the tropics (OTT)	Jablonski et al. 2006	Species were generated in the tropical regions and dispersed to extratropical regions but maintain its presence in its ancestral areas	High rates of speciation are predicted in tropical regions in contrast with temperate regions. Asymmetric dispersal have occurred along the biogeographical history of a taxa from tropical to temperate areas.	MRD, Mean age, GeoSSE	These metrics are the same used to test the PNC/TNC hypothesis as we discuss above.
Time for speciation effect (TEE)	Stephens and Wiens 2003	Tropical regions accumulated more species because their clades had more time to speciate than temperate regions.	Regions recently colonized had lower species richness than regions where clades colonized very early in the history of a clade.	Mean age	1) Mean age does not provide an accurate description of which lineages colonized first a region. To test this hypothesis, it might be necessary to perform an ancestral range reconstruction of all co-occurring clades and estimate its diversification rates (i.e., total diversification for each independent colonized clade; Rabosky 2009; 2012).

1020

1021 Table 3. Parameter estimates from the GeoSSE model for three taxonomic groups
1022 (Furnariides birds, hylid frogs, and *Anolis* lizards) across two regions. Areas for each
1023 taxonomic group as follows: Furnariides birds: A: Forest; B: Open areas; Hylid frogs: A:
1024 Extra tropics; B: Tropics; *Anolis* lizards: A: Islands; B: Mainland.

Group	Rates	A	B	AB
Furnariides birds	Speciation	0.139 ± 0.020	0.223 ± 0.065	0.041 ± 0.020
	Extinction	0.040 ± 0.025	0.107 ± 0.075	-
	Dispersal	0.021 ± 0.004	0.311 ± 0.114	-
	Net diversification	0.099 ± 0.005	0.116 ± 0.01	-
Hylid frogs	Speciation	0.044 ± 0.003	0.044 ± 0.003	0.041 ± 0.025
	Extinction	0.002 ± 0.002	0.002 ± 0.002	-
	Dispersal	0.001 ± 0.001	0.035 ± 0.010	-
	Net diversification	0.042 ± 0.003	0.042 ± 0.003	-
Anolis lizards	Speciation	0.058 ± 0.003	0.058 ± 0.003	1.245 ± 1.303
	Extinction	0.001 ± 0.001	0.001 ± 0.001	-
	Dispersal	0.002 ± 0.001	0.0003 ± 0.000	-
	Net diversification	0.057 ± 0.002	0.057 ± 0.002	-

1025

1026   Table 4. Frequency of dispersal events inferred using biogeographical stochastic
1027   mapping (BSM) for three taxonomic groups (Furnariides birds, hylid frogs, and *Anolis*
1028   lizards) across two regions. Areas for each taxonomic group as follows: Furnariides
1029   birds: A: Forest; B: Open areas; Hylid frogs: A: Extra tropics; B: Tropics; *Anolis* lizards:
1030   A: Islands; B: Mainland.
1031

Event	Group	Regions	A	B
Range expansions	Furnariides birds	A	0	92.62 ± 4.39
		B	31.4 ± 4.29	0
	Hylid frogs	A	0	0.64 ± 0.78
		B	12.92 ± 0.88	0
	Anolis lizards	A	0	0
		B	0	0
Founder events	Furnariides birds	A	0	24.46 ± 2.54
		B	10.88 ± 2.22	0
	Hylid frogs	A	0	0.66 ± 0.66
		B	4.3 ± 1.42	0
	Anolis lizards	A	0	2.02 ± 0.14
		B	0.12 ± 0.33	0

1032
1033

FIGURE LEGENDS

Figure 1. Diagram illustrating how differences in speciation, extinction, and dispersal
rates between regions can generate a geographical species richness gradient. The
phylogenetic trees below illustrate how the differences in speciation and extinction
rates between two regional assemblages can shape a gradient of species richness
(degraded blue colour).

1040

Figure 2. Geographical patterns of some phylogenetic metrics used in macroecological
studies to explore evolutionary process underlying geographical diversity gradients
(see also Table 1 for a detailed explanation). Left column *Anolis* lizards; Middle column:
Hylid frogs; Right column: Furnariides birds. (A-C) observed richness patterns; (D-F)
rPD: residual phylogenetic diversity (i.e., after controlling for species richness); (G-I)
PSV: phylogenetic species variability; (J-L) MRD: mean root distance; (M-O) MDR:
mean diversification rate; (P-R) Mean ages: average ages of species.

1048

Figure 3. P-values distribution for each phylogenetic metric obtained through the null
model (see main text for details). The vertical red lines represent the empirical 0.05 cut-
off. Note that for all cases very few cells are below the 0.05 cut-off. (A-C) rPD: residual
phylogenetic diversity (i.e., after controlling for species richness); (D-F) PSV:
phylogenetic species variability; (G-I) MRD: mean root distance; (J-L) MDR: mean
diversification rate; (M-O) Mean ages: average ages of species.

1055

1056   Figure 4. Variation of phylogenetic metric values for Furnariides birds in forest and
1057   open areas. rPD: residual phylogenetic diversity (i.e., after controlling for species
1058   richness); PSV: phylogenetic species variability; MRD: mean root distance; MDR: mean
1059   diversification rate; Mean ages: average ages of species.

1060

1061   Figure 5. Variation of phylogenetic metric values for Hylid frogs in tropics and extra-
1062   tropics regions. rPD: residual phylogenetic diversity (i.e., after controlling for species
1063   richness); PSV: phylogenetic species variability; MRD: mean root distance; MDR: mean
1064   diversification rate; Mean ages: average ages of species.

1065

1066   Figure 6. Variation of phylogenetic metric values for *Anolis* lizards in continental and
1067   insular areas. rPD: residual phylogenetic diversity (i.e., after controlling for species
1068   richness); PSV: phylogenetic species variability; MRD: mean root distance; MDR: mean
1069   diversification rate; Mean ages: average ages of species.

1070

DIVERSITY GRADIENTS

**Higher speciation
Lower extinction
Dispersal (out)**

**Lower speciation
Higher extinction
Dispersal (in)**

Speciation = 0.5; Extinction = 0.05

Speciation = 0.3; Extinction = 0.15

Stochastic map
ancstates: global optim, 2 areas max. d=5; e=0; j=2.9172; LnL=−102.92

Stochastic map
ancstates: global optim, 2 areas max. d=5; e=0; j=2.9704; LnL=−899.05

Millions of years ago

Stochastic map
ancstates: global optim, 2 areas max. d=0.7012; e=0; j=2.713; LnL=−20.91