Genotypic and phenotypic diversity within the neonatal HSV-2 population

Lisa N. Akhtar[1,5], Christopher D. Bowen[2], Daniel W. Renner[2], Utsav Pandey[2], Ashley N. Della Fera[3], David W. Kimberlin[4], Mark N. Prichard[4], Richard J. Whitley[4], Matthew D. Weitzman[3,5]*, Moriah L. Szpara[2]*

[1] Department of Pediatrics, Division of Infectious Diseases, Children's Hospital of Philadelphia

[2] Department of Biochemistry and Molecular Biology, Center for Infectious Disease Dynamics, and the Huck Institutes of the Life Sciences, Pennsylvania State University

[3] Department of Pathology and Laboratory Medicine, Division of Protective Immunity and Division of Cancer Pathobiology, Children's Hospital of Philadelphia

[4] Department of Pediatrics, Division of Infectious Diseases, University of Alabama at Birmingham

[5] University of Pennsylvania Perelman School of Medicine

*Corresponding Authors
Moriah L. Szpara
Dept. of Biochemistry & Molecular Biology
The Huck Institutes of the Life Sciences
W-208 Millennium Science Complex (MSC)
Pennsylvania State University
University Park, PA 16802 USA
Phone: 814-867-0008
orcid.org/0000-0001-9859-1678
Email: moriah@psu.edu

Matthew D. Weitzman
Division of Protective Immunity
The Children's Hospital of Philadelphia
4050 Colket Translational Research Building
3501 Civic Center Blvd
Philadelphia, PA 19104-4318
Phone: 267-425-2068
orcid.org/0000-0001-9713-167X
Email: weitzmanm@email.chop.edu

Author emails:
Lisa N. Akhtar: AkhtarL@email.chop.edu
Christopher D. Bowen: cdb5004@psu.edu
Daniel W. Renner: dwr19@psu.edu
Utsav Pandey: uup104@psu.edu
Ashley N. Della Fera: ashleyycoxx@gmail.com
David W. Kimberlin: DKimberlin@peds.uab.edu
Mark N. Prichard: MPrichard@peds.uab.edu
Richard J. Whitley: RWhitley@peds.uab.edu
Matthew D. Weitzman: weitzmanm@email.chop.edu
Moriah L. Szpara: moriah@psu.edu

Keywords: Neonatal, herpes simplex virus 2, human herpesvirus 2, comparative genomics, viral spread, minority variants

Abstract

More than 14,000 neonates are infected with herpes simplex virus (HSV) annually.

Approximately half display manifestations limited to the skin, eyes, or mouth (SEM disease).

The rest develop invasive infections that spread to the central nervous system (CNS disease or

encephalitis) or systemically (disseminated disease). Invasive HSV disease is associated with

significant morbidity and mortality, but viral and host factors that predispose neonates to these

forms are unknown. To define viral diversity within the infected neonatal population, we

evaluated ten HSV-2 isolates from newborns with a range of clinical presentations. To assess

viral fitness independent of host immune factors, we measured viral growth characteristics in

cultured cells and found diverse *in vitro* phenotypes. Isolates from neonates with CNS disease

were associated with larger plaque size and enhanced spread, with isolates from cerebrospinal

fluid (CSF) exhibiting the most robust growth. We sequenced complete viral genomes of all ten

neonatal viruses, providing new insights into HSV-2 genomic diversity in this clinical setting.

We found extensive inter-host and intra-host genomic diversity throughout the viral genome,

including amino acid differences in more than 90% of the viral proteome. The genes encoding

glycoprotein G (gG, US4), gI (US7), gK (UL53), and viral proteins UL8, UL20, UL24, and US2

contained variants that were found in association with CNS isolates. Many of these viral proteins

are known to contribute to cell spread and neurovirulence in mouse models of CNS disease. This

study represents the first application of comparative pathogen genomics to neonatal HSV

disease.

Importance

Herpes simplex virus (HSV) causes invasive disease in half of infected neonates, resulting in

significant mortality and permanent cognitive morbidity. The factors that contribute to invasive

disease are not understood. This study reveals diversity among HSV isolates from infected

neonates, and makes the first associations between viral genetic variations and clinical disease

manifestations. We found that viruses isolated from newborns with encephalitis show enhanced

spread in culture. These viruses contain protein-coding variations not found in viruses causing

non-invasive disease. Many of these variations are found in proteins known to impact

neurovirulence and viral spread between cells. This work advances our understanding of HSV

diversity in the neonatal population and how it may impact disease outcome.

Introduction

Each year an estimated 10,000 neonates are infected with HSV-2, and 4,000 infected with HSV-

1, worldwide (1). Infants are typically infected at the time of birth by maternal genital shedding

of HSV, most often by mothers who are not aware of their infection (2–4). The recent increase in

genital HSV-1 incidence among women of childbearing age, particularly in developed nations,

suggests that the burden of neonatal infection will continue to rise (1, 5). While some infected

infants exhibit only superficial infection limited to the skin, eyes, or mouth (SEM disease; 45%),

about half develop invasive systemic (disseminated disease; 25%) or central nervous system

(CNS disease; 30%) infections associated with significant morbidity and mortality (6, 7).

Currently, the antiviral medication acyclovir is the standard therapy for all forms of neonatal

HSV disease. Although this intervention has reduced mortality due to invasive disease, most

survivors of invasive disease are left with permanent neurodevelopmental deficits (8, 9).

The factors that predispose a neonate to invasive HSV infection are not entirely known. Recent

studies have found that some adults and children outside of the neonatal period who experience

HSV infection of the brain have a host genetic defect within the Toll-like receptor-3 (TLR3)

pathway (10, 11). Outside of the neonatal period, HSV encephalitis is rare, as are host defects in

the TLR3-pathway. By contrast, half of HSV-infected neonates experience invasive CNS or

disseminated disease, making it less likely that host genetic defects alone could account for all of

the observed cases of invasive infection in neonates. Prior clinical data on mother-to-infant

transmission of HSV indicate that most cases of neonatal disease, including invasive forms of

disease, result from newly acquired or primary HSV infection prior to the development of

maternal antibody production (2–4, 12). This suggests a window of opportunity where the

contributions of viral genetic variation to the progression of invasive infection and disease may

be greater than in adults.

Prior studies have identified viral genetic factors that influence virulence or disease for

reoviruses, influenza virus, HIV, and others (13–18). In contrast to these RNA viruses, HSV was

presumed to have lower genetic diversity and potential for variation in virulence, due to its

relatively stable DNA genome and long co-evolutionary history with humans (19). The

assumption of limited HSV heterogeneity was supported by early studies that utilized low-

resolution restriction fragment length polymorphism (RFLP) or single-gene analyses to compare

multiple HSV isolates (20–22). However Rosenthal and colleagues used RFLP and PCR analysis

of a single locus to demonstrate that a heterogeneous HSV population can exist in an invasive

neonatal infection, and provided proof of principle that natural genetic variation can impact

neurovirulence (23, 24). More recently, advances in high-throughput sequencing (HTSeq) have

enabled a re-evaluation of herpesvirus genome-wide variation, which suggests that herpesviruses

harbor extensive diversity both between strains or individuals (inter-host variation) as well as

within a single individual (intra-host variation) (25–27). These minor genetic variants may

become clinically important if a variant within the viral population becomes the new dominant

allele or genotype as a result of a bottleneck at transmission, entry into a new body compartment,

or selective pressure such as antiviral therapy (28, 29).

Several recent examples have demonstrated the potential for new insights gained by applying

HTSeq approaches to herpesvirus infections in a clinical setting. In HTSeq studies of congenital

infection by the beta-herpesvirus human cytomegalovirus (HCMV), Renzette *et al.* found

evidence for heterogeneous viral populations both within and between hosts (30–35). The levels

of diversity observed in congenital HCMV infections far exceeded those observed in adult

infections (36–38). HTSeq-based examination of vaccine-associated rashes due to the alpha-

herpesvirus varicella zoster virus (VZV) demonstrated that adult skin vesicles contain a subset of

the viral population introduced during vaccination, and found at least 11 VZV genomic loci that

were linked to rash formation (26). Recent HTSeq comparisons of adult genital HSV-2 revealed

the first evidence of a single individual shedding two distinct strains (39), demonstrated changes

in the viral genome over time in a recently infected host (40), and provided the first evidence of

ancient recombination between HSV-1 and HSV-2 (41, 42). However, to date there has been no

evaluation of genome-wide variation in neonatal HSV isolates to determine the levels of

diversity in this population, or the potential impact(s) of viral genetic variants on disease.

Until recently, several technical barriers prevented thorough assessment of neonatal HSV

genomes. A key constraint on studies of neonatal disease has been the availability of cultured,

minimally-passaged viral samples that are both associated with clinical information and have

also been maintained in a low-passage state that is appropriate for sequencing and further

experimental studies. Historically viral culture was part of the HSV diagnostic workflow, but this

has been superseded in clinical laboratory settings by the speed and sensitivity of viral detection

by PCR (43, 44, 6). This change limits neonatal HSV sample availability for *in vitro* and animal

model studies. In addition, many previously archived neonatal HSV isolates have been passaged

extensively, allowing them to acquire mutations that enhance viral growth in culture (23, 24).

Additional challenges for HTSeq approaches to neonatal HSV include the large size of the viral

genome (~152 kb), its high G+C content (~70%), and a large number of variable-number tandem

repeats in the viral genome (> 240 mini/micro-satellite repeats and >660 homopolymers of ≥6

base pairs) (45, 27). Therefore, many studies of HSV diversity (21, 46–49) or the effect of HSV

genetic variation on disease (50–52) have relied on low-resolution restriction fragment length

polymorphism (RFLP) or single-gene PCR analyses, due to the speed and ease of analysis in

comparison to whole-genome approaches (53–58). To overcome these challenges, we combined

our expertise in HSV comparative genomics and phenotypic analysis (59, 53, 60, 58) with a

unique resource of low-passage, well-annotated neonatal specimens (8, 61, 9).

Here we analyzed a set of ten low-passage clinical HSV-2 isolates collected from neonates with

HSV infection, enrolled in one of two clinical studies that spanned three decades of patient

enrollment (1981-2008) (8, 61, 9). These samples represented a wide range of clinical

manifestations including SEM, CNS, and disseminated disease, and each sample was associated

with de-identified clinical information. We defined the level of diversity in this population using

comparative genomics and an array of cell-based phenotypic assays. We found that HSV-2

isolates displayed diverse *in vitro* phenotypes, as well as extensive inter- and intra-host diversity

distributed throughout the HSV-2 genome. Finally, we found coding variations in several HSV-2

proteins associated with CNS disease. This study represents the first-ever application of

comparative pathogen genomics to neonatal HSV disease and provides a basis for further

exploration of genotype-phenotype links in this clinically vulnerable patient population.

Results

Neonatal HSV-2 samples represent a diverse clinical population

We utilized samples collected from ten HSV-2-infected neonates enrolled by the National

Institute of Allergy and Infectious Diseases Collaborative Antiviral Study Group (CASG) for

clinical trials between 1981 and 2008 (8, 61, 9). These infants encompassed a range of clinical

disease manifestations (see **Table 1**), with about half experiencing invasive CNS disease (5

patients) or disseminated (DISS) disease with CNS involvement (2 patients), and the remainder

experiencing non-invasive SEM disease (3 patients). Extensive clinical information was

available for each patient, including long-term neurocognitive and motor outcomes (**Table 1**).

This population was also diverse with respect to sex, race, gestational age, and enrollment center

(**Table 1**; enrollment center data not shown). All samples were collected at the time of diagnosis,

prior to initiation of acyclovir therapy. Each isolate was cultured once as part of the diagnostic

process, with expansion only for the experiments shown here. Although the sample size was

constrained by the rarity of neonatal HSV infection and availability of appropriately maintained

isolates, our group is similar in size to prior HTSeq comparisons of congenital HCMV samples

(31, 32, 34, 35), and is the largest group of neonatal HSV samples ever subjected to comparative

genomic and phenotypic analysis.

Neonatal HSV-2 isolates have different fitness in culture

To determine whether the viruses isolated from this neonatal population **(Table 1)** were

intrinsically different, we assessed viral growth in culture, which provides a consistent

environment that is independent of host genetic variation. To minimize the impact of immune

pressure we selected Vero monkey kidney cells, which lack an interferon response (62, 63). Each

viral isolate was applied to a confluent monolayer of cells *in vitro*, and allowed to form plaques

for 100 hours (h) (**Figure 1A**). Average plaque size differed between isolates, with six of the ten

isolates being statistically larger (indicated in green) than the other four (indicated in black,

Figure 1B) (one-way ANOVA with Holm-Sidak's multiple comparisons test, $p < 0.05$). The

average plaque size of the previously described low passage, adult HSV-2 isolate SD90e is

shown for comparison (64). The largest plaque sizes were observed in the two viruses isolated

directly from the cerebrospinal fluid (CSF; isolates CNS11 and DISS14; **Figure 1B**); these two

isolates were not statistically different from one another in size, but were significantly larger than

any of those isolated from the skin (one-way ANOVA with Holm-Sidak's multiple comparisons

test, $p < 0.05$). Plaque size was assessed at each passage in culture and remained constant from the

time the isolates were received in our laboratory (passage 2) through their genetic and

phenotypic analysis (passage 4). The variance in plaque sizes produced by a given isolate was

not statistically different between isolates **(Figure 1B)**. The differences in average plaque size

between isolates suggested that the HSV-2 populations found in each neonatal isolate are indeed

intrinsically different.

Entry kinetics, DNA replication, protein expression, and virus production do not

account for differences in plaque size

Plaque formation is a complex endpoint that involves the ability of the virus to enter the cell,

replicate its double stranded DNA genome, produce viral proteins and assemble new virions that

spread to adjacent cells. Therefore, we explored whether the differences in plaque formation

observed in Vero cells reflected inherent differences in the ability of isolates to complete each of

these stages of the viral life cycle. For these comparisons, two large-plaque-forming isolates

(CNS11 and CNS03) were compared to two small-plaque-forming isolates (CNS12 and SEM02).

First, we compared each isolate's rate of cell entry. Virus was applied to chilled cells, followed

by warming to synchronize cell entry **(Figure 2A).** A low pH solution was applied at various

points over the first hour of cell entry to inactivate any virus that had not yet entered a cell, and

plaque formation was then allowed to proceed for 100 h. We found no difference in rates of cell

entry between these four representative viral isolates (**Figure 2B**), suggesting that large plaques

did not result from increased rates of virus entry into cells.

We next infected Vero cells at high multiplicity of infection (MOI=5) to compare the outcome of

a single round of viral replication. We found that all four isolates produced similar numbers of

genome copies (as measured by qPCR for the gB gene; see Methods for details) **(Figure 2C)**,

indicating that differences in viral DNA replication did not influence plaque size. We quantified

the production of infectious virus by counting plaque-forming-units (PFU) on Vero cell

monolayers, as well as on the highly-permissive U2OS human bone osteosarcoma epithelial cell

line (65). No differences in virus production were noted between the four isolates when

quantified on either cell type (**Figure 2D, 2˜E**). U2OS cells lack innate sensing of viral infection

through the STING pathway (66) and can even support the growth of highly-defective HSV-

isolates that lack ICP0 function (65). All isolates formed large plaques on the highly-permissive

U2OS cell monolayers (**Figure S1**), allowing us to rule out the possibility that very small foci of

infection were missed during titering of the small plaque-forming isolates on Vero cells (**Figure**

2D, E). Finally, we compared viral protein production for these isolates, and found no

differences in the expression levels of a panel of HSV-2 viral proteins at either early (6 h post

infection, hpi) or late (24 hpi) time points of a single round of high MOI infection (**Figure 2F,**

G; see legend for list of proteins). Taken together, these results suggested that large- and small-

plaque-forming neonatal HSV-2 isolates did not differ significantly in viral entry, DNA

replication, infectious virus production, or protein production over a single round of infection.

Large-plaque-forming isolates exhibit enhanced cell-to-cell spread

We next assessed the ability of representative large- and small-plaque-forming isolates to spread

from cell-to-cell. Vero cell monolayers were infected at low MOI (MOI=0.001) in order to assess

differences over multiple rounds of viral replication and spread throughout the cell monolayer

(**Figure 3A**). The contribution of indirect cell-free spread was minimized by including 0.1%

human serum in the media, and changing the media every 24 h to remove released virus and

refresh serum levels. Over a 72-hour time course, cell monolayers were assessed to measure the

extent of cell-to-cell spread, either by harvesting and titering PFU production (**Figure 3B-C**), or

by fixing infected cell monolayers and evaluating the distribution of virus by

immunofluorescence (**Figure 3D**, **E**, and **Figure S2**). Viral titers recovered from harvested cells

were similar at 2 hpi, confirming that equivalent amounts of each virus were present following

initial infection (**Figure 3B**). By 72 hpi, after multiple rounds of replication and spread, the

large-plaque-forming isolates CNS11 and CNS03 had achieved viral titers significantly greater

than those of the small-plaque-forming isolates CNS12 and SEM02 (**Figure 3B, C**; two-way

ANOVA followed by Tukey's multiple comparison test, p<0.0001 at 72h). Isolate CNS11,

which was obtained directly from the CSF, produced titers statistically greater than the other

three isolates **(Figure 3B, C)**.

We also directly evaluated cell-to-cell spread by immunostaining and quantifying the distribution

of infected cells around each infectious focus. Infected Vero cell monolayers were fixed at 24,

48, and 72 hpi, and subjected to fluorescent immunocytochemistry with an antibody directed

against total HSV (**Figure 3D** and **Figure S2**). The region of HSV-positive cells surrounding a

single initial infection was greater for the large-plaque-forming neonatal HSV-2 isolates at 24

hpi, and increased by 48 and 72 hpi. The central cytolytic clearings seen in the monolayers

infected by CNS11 or CNS03 were approximately two-fold greater than those infected by

CNS12 or SEM02, reflecting the average two-fold increase in plaque size observed in

methylene-blue-stained monolayers in **Figure 1**. However, the region of infected cells

surrounding the central cytolytic clearing for CNS11 and CNS03 was dramatically larger than

for CNS12 and SEM02, suggesting that large-plaque-forming neonatal HSV-2 isolates show

greater spread from cell-to-cell than could have been predicted by measuring plaque size alone.

To quantify this increase in the area of infected cells after a low-MOI infection, each coverslip

was imaged **(Figure S2)** and the total number of immunofluorescent pixels was quantified

(Figure 3E). By 72 hpi, the area of HSV-infected cells was statistically greater for the large-

plaque-forming isolates CNS11 and CNS03, as compared to the small-plaque-forming isolates

CNS12 and SEM02 (two-way ANOVA followed by Tukey's multiple comparison test, p<0.05 at

72h). Together these data indicated that large-plaque-forming isolates shared an enhanced ability

to spread cell-to-cell through culture, in comparison to small-plaque-forming isolates.

Comparative genomics reveals genetic diversity in neonatal HSV-2 isolates

The differences identified in cell-to-cell spread between neonatal isolates in culture indicated the

existence of intrinsic differences between these viruses. To reveal how genetic variation may

contribute to viral phenotypes in culture, and ultimately to clinical disease manifestations, we

sequenced the complete viral genome of all ten neonatal HSV-2 isolates. For each isolate, we

sequenced purified viral nucleocapsid DNA and assembled a consensus genome, which

represents the most common genotype at each nucleotide locus in the viral population. The

clinical trials utilized in this study enrolled HSV-infected infants from multiple sites across the

United States (8, 9, 61). Therefore, we first assessed the overall degree of relatedness between

these viral genomes to understand whether any similarities in viral or geographic origin might

have contributed to *in vitro* or clinical phenotype patterns. In light of the known potential for

recombination in the phylogenetic history of HSV (49, 67, 53, 42), we used a graph-based

network to investigate the phylogenetic relationship between these isolates. We found a similar

degree of divergence among all ten neonatal HSV-2 consensus genomes (**Figure 4A**). We then

compared these ten neonatal HSV-2 genomes to all available HSV-2 genomes in GenBank (all of

which are adult; see **Table S1** for HSV-2 GenBank accessions and references) to discern any

geographic or other clustering (see **Table 1**). The neonatal isolates did not segregate into any

exclusive groupings, but instead intermingled among non-neonatal isolates, with deep branches

between each isolate (**Figure 4B**). These findings were corroborated using an alternative

phylogenetic clustering algorithm as well (**Figure S3**). This suggested that similarities in viral

genetic origin were not responsible for determining the cellular or clinical outcomes of neonatal

HSV-2 infection.

Overall protein-coding diversity in neonatal HSV-2 isolates is similar to that

observed in adult HSV-2 strains

We next asked whether overt defects in any single HSV-2 protein might be associated with

clinical or *in vitro* spread phenotypes. In examining the coding potential of all ten neonatal HSV-

2 isolates, we found no protein deletions or truncations encoded by any of these viral genomes.

These comparisons revealed a total of 784 nucleotide differences in 71 genes (see **Table S2**),

resulting in 342 non-synonymous AA differences in 65 proteins. This led to an average of 1%

coding variation (range 0.0 : 2.6%). This variation was spread widely throughout the HSV-2

genome, without concentration in a particular genomic region or category of protein function

(Figure 5). We found a similar level of coding diversity in this set of neonatal isolates as in a

comparable set of 10 adult HSV-2 isolates **(Figure 5, Table S2).** We also compared these

neonatal genomes to the full set of 58 annotated adult HSV-2 genomes from GenBank (listed

Table S1). As expected from the difference in sample size, there was more overall diversity

found across 58 adult HSV-2 genomes than in 10 neonatal genomes. A comparison of the dN/dS

ratio in neonatal vs. adult HSV-2 genomes revealed a similar trend with a few outliers visible on

each axis, e.g. UL38 and US8A had a higher dN/dS ratio in neonatal than in adult isolates (see

Table S2 and **Figure S4** for full comparison). These data indicated that at the consensus level,

neonatal HSV-2 isolates display substantial inter-host coding diversity spread throughout the

genome, but do not possess strikingly more diversity or an excess of genetic drift as compared to

adult isolates.

Minority variants expand the potential coding diversity of neonatal HSV-2 isolates

We next focused our attention on differences below the consensus level in each intra-host viral

population. The amino acid (AA) variations described above exist in the consensus genomes of

each isolate. Since viral replication creates a population of genomes, we next assessed whether

minor allelic variants existed within the viral population of any neonatal HSV-2 isolate, thereby

expanding the viral genetic diversity within each host. The significant depth of coverage from

deep-sequencing of each isolate allowed us to search for minority variants at every nucleotide

position of each genome. We defined a minor variant as any nucleotide allele (single nucleotide

polymorphism, or SNP) or insertion/deletion (INDEL) with frequency below 50%, but above 2%

(a conservative limit of detection; see Methods for additional criteria). We found minor variants

in the viral genome population of all 10 neonatal HSV-2 isolates, albeit to a different degree in

each isolate (**Figure 6A** and **Table S3**). In total, there were 1,821 minor variants, distributed

across all genomic regions (**Figure S5; Table S3**). For both SNPs and INDELs, intergenic

minority variants outnumbered those in genes (genic), likely reflecting the higher selective

pressures against unfavorable mutations in coding regions. The neonatal isolate DISS29 had 8-

10-fold higher levels of minority variants than other neonatal isolates (**Figures 6A and S5**), and

these variants were often present at a higher frequency or penetrance of the minor allele than

observed in other neonatal isolates (**Figures 6B and S5**). We further examined the distribution of

minor variants that occurred in genes, and found that nearly every HSV-2 protein harbored

minority variants in at least one neonatal isolate (**Figure 6C**). Only UL3, UL11, UL35, and

UL55 were completely devoid of minority variants. Three of these genes (UL3, UL11, UL35)

were also devoid of AA variations at the consensus level (**Figure 5**). These data revealed the

breadth of potential contributions of minority variants to neonatal HSV-2 biology, which could

303   undergo selection over time or in specific niches.

Coding variations identified in neonatal HSV-2 isolates associated with CNS disease

To understand how viral genetic variants might relate to clinical disease, we assessed whether

any of the consensus level coding variations identified in our group of ten neonatal HSV samples

segregated with clinical disease features. A number of AA variations were shared by the CSF-

derived isolates CNS11 and DISS14, which formed the largest plaques *in vitro*. These included

AA variants at the level of the consensus genome in the HSV-2 genes encoding glycoprotein K

(gK, UL53 gene: V323M), glycoprotein I (gI, US7 gene: R159L and P215S), UL8 (R221S), US2

(F137L), and glycoprotein G (gG, US4 gene: R338L, S442P, and E574D) (**Figure 7**). The gI

variants exist individually in other viral isolates obtained from neonates with CNS disease

(**Table S4**). Isolates collected from infants experiencing disseminated disease with CNS

involvement also shared a variant in the HSV-2 UL20 gene (P129L) (**Figure 7**). One variant in

the HSV-2 UL24 gene (V93A) was shared only by isolates from infants with SEM disease, with

all isolates from infants with CNS disease containing a valine at this position **(Figure 7).** The

sample size of these comparisons was constrained by the overall limits of neonatal HSV-2 isolate

availability. While CNS11 and DISS14 shared several variants listed in **Figure 7**, these viral

genomes were not similar at the consensus genome level (**Figure 4**). These isolates also had non-

shared coding differences in the same genes that contained shared variants (e.g. gK, UL8, gG).

This indicates a potential for convergent evolution at these loci. Many of the coding variations

identified in this dataset involve viral proteins that are known to modulate cell-to-cell spread

(68–72) and/or contribute to neurovirulence in mouse models of CNS infection (70, 73–79)

(**Figure 7** and **Table S4**), however, their role in human disease has not yet been assessed.

Discussion

Host factors have not been identified to explain the >50% of neonates experiencing invasive

CNS or disseminated forms of HSV infection. There is growing evidence that most

herpesviruses, including HSV-1 and HSV-2, contain significant genetic variation (53, 27, 56,

80–82). The potential contributions of viral genetic variation to clinical disease in neonates

therefore warrants exploration. Here, we analyzed genetic and phenotypic diversity for HSV-2

isolated from ten neonatal patients spanning two clinical studies (8, 61, 9). Although this group is

small, the associations identified here provide the first insights into the potential impact of viral

variability on clinical outcomes in neonatal HSV disease, and serve as a starting point for further

mechanistic investigation. We found that these ten neonatal HSV-2 isolates exhibited diverse

growth characteristics in culture, with larger plaque-forming isolates observed more often in

infants with CNS as opposed to SEM disease. Furthermore, we established that enhanced viral

spread through culture was the main contributor to this large plaque formation. Using

comprehensive comparative genomics, we further demonstrated that these neonatal HSV-2

isolates contained extensive genetic diversity both within and between hosts. These data revealed

several specific viral genetic variations that were associated with cases of CNS disease, in

proteins known to contribute to cell-to-cell spread and/or neurovirulence in mouse models of

CNS disease. Further studies are required to determine the impact of these variations on HSV-2

neurovirulence and progression to CNS disease.

Genomic comparison of these neonatal isolates revealed a wide range of genetic diversity. At the

consensus genome level, which reflects the most common allele in each viral population, we

found that coding differences between strains were as numerous as between previously described

sets of adult HSV-2 isolates **(Figure 5)** (80–82). Furthermore, the specific genetic variations

associated with neonatal CNS disease in our study can also be found in genital HSV-2 isolates

which are not associated with CNS disease in adults. This suggests that the dramatic differences

in clinical manifestations following HSV-2 infection in neonates, with significantly higher rates

of invasive CNS infection, are not due to unique neonatal HSV-2 strains. It is more likely that

genetic variations occurring naturally in the adult HSV-2 population, when present in the specific

context of neonatal infection, may confer neurovirulence and progression to CNS disease.

At the level of minority variants (MV), which represent rare alleles that exist within each intra-

host viral population, we found that the DISS29 harbored 8-10 fold and CNS15 harbored 3-4

fold more MV than other neonatal virus genomes (**Figures 6 and S4**). This could be indicative

of a mixed viral population (e.g. a multi-strain infection) or of decreased polymerase fidelity (28,

27). Diversity in viral populations has also been observed in congenital HCMV infection (31, 32,

35, 34). However, this is the first time that evidence has been found for this level of viral

population diversity with HSV-2. These minor genotypes may be selected or genetically isolated

in particular niches (e.g. CSF), as observed in a comparison of VZV skin vesicles (26), or by

antiviral drug selection, as was recently demonstrated in two adults with genital HSV-2 infection

(83). All of the isolates sequenced in this study were collected at the initial time of diagnosis

(≤19 days old), prior to acyclovir treatment and prior to the development of an HSV-specific

immune response. It would therefore be compelling to examine the viral genome population

from serial patient isolates over time to identify shifts in the frequency of MV due to antiviral or

immune selection. Isolates from different body sites of the same patient could likewise be

compared to determine whether particular genotypes are enriched in different body niches.

This comparison of viral genotype to clinical phenotype revealed associations between neonatal

CNS disease and several viral protein variants that may impact neurovirulence through

modulation of cell-to-cell spread. Although the sample set in this proof of concept study is small,

we observed potential patterns that warrant exploration in a larger dataset. It is important to

acknowledge that there is limited availability of samples from neonatal infection, both due to the

rarity of these infections, and the fragility and limited body size of the infected infants. These

natural circumstances lead to minimal sample collection from infected neonates. The finding that

CNS-associated isolates in our study often exhibit enhanced spread between cells in culture,

particularly those derived directly from the CSF, suggests that one or more of these variants

could be functionally significant. Coding differences in viral proteins not known to contribute to

neurovirulence were also found to be associated with neonatal CNS disease, and represent

potential novel contributions to invasive infection. These promising results warrant exploration

in a larger study, ideally with isolates from multiple time points and/or body sites from each

infected infant. This would enable a better understanding of how overall viral genetic diversity

contributes to neuroinvasion.

Methods

Viruses

Viruses were collected from neonates enrolled in clinical studies (8, 9, 61) by the Collaborative

Antiviral Study Group (CASG) at the University of Alabama Birmingham (UAB). Samples were

collected from either the cerebrospinal fluid (CSF) or skin. Enrollment in original studies was

evenly split between males and females, and included both black and white patients. Clinical

morbidity score was determined at 12 months of life as previously defined (84, 12). Initial

collection of samples, use of samples in this study, and use of de-identified clinical information

was approved by the UAB Institutional Review Board. See Supporting Information for

additional details of this and subsequent methods.

Cell culture

395   Human lung fibroblast MRC-5 cells (ATCC®, CCL-171), African green monkey kidney Vero

396   cells (ATCC®, CCL-81), and human epithelial bone osteosarcoma U2OS cells (ATCC®, HTB-

397   96) were cultured under standard conditions. Cell lines were authenticated by ATCC prior to

398   purchase, and were confirmed to be mycoplasma free throughout experiments by periodic testing

399   (LookOut Mycoplasma, Sigma).

Virus culture

401   Viruses were cultured at the time of diagnosis and snap frozen after 1 passage. Each viral isolate

402   was then passaged 3 times on MRC-5 cells at a MOI of 0.01, with harvest at the time of

403   complete cytopathic effect (between 50-70 h). Viral stocks were titered on either Vero cells (100

404   h) or U2OS cells (48 h) under a methylcellulose overlay. Plaque size and morphology did not

405   change for any viral isolates over the course of virus stock expansion.

Plaque measurements

407   Plaques were stained with 0.5% methylene blue. Serial 4X brightfield images were collected on

408   an EVOS FL Auto Imaging System and stitched by EVOS software (University of Pennsylvania

409   Cell and Developmental Biology Microscopy Core). Plaque area was measured using ImageJ

410   software.

Genome copy number estimation by quantitative PCR for UL27

412   DNA was extracted and quantitative PCR was performed with primers specific to the viral

413   glycoprotein B gene (gB; UL27) (43). Absolute quantification was calculated based on a

414   standard curve of HSV-1 strain F nucleocapsid DNA (59).

Viral entry assay

Monolayers of Vero cells were cooled to 4°C for 30 minutes prior to infection with 100 PFU of

each viral isolate. After 1 h of viral incubation at 4°C, unbound virus was removed by washing

and cells were moved to 37°C. At 0, 10, 20, 30, 45, or 60 minutes, a low-pH citrate buffer was

applied to inactivate extracellular virus. For each condition, parallel infections were performed

without the addition of citrate solution (control). Cell monolayers were washed and allowed to

form plaques under methylcellulose for 100 h. Viral entry was quantified as the fraction of

plaques formed following citrate buffer application, where 100% is the number of plaques

formed on a monolayer not treated with citrate buffer (control).

Single-step and multi-step growth curves

425   Growth curves of HSV infection were performed in Vero cells. At 2 hpi, 0.1% human serum was

426   added to reduce cell-free spread of virus. Single-step growth curves were performed at MOI=5,

427   and multi-step growth curves at MOI=0.001, as defined by titering viral stocks on U2OS cells.

428   Every 24 h, the supernatant was removed and media containing 0.1% human serum reapplied.

Immunocytochemistry

Immunocytochemistry was performed as previously described (85). Infection was detected with

rabbit anti-HSV primary antibodies (Agilent Dako, B0114) and fluorophore-conjugated anti-

rabbit secondary antibodies (Invitrogen, A-11008). Cell nuclei were counterstained with DAPI.

5X images were collected with a Leica DM6000 wide field microscope equipped with a

Photometrics HQ2 high resolution monochrome CCD camera, and processed with LAS AF

software (UPenn Cell and Developmental Microscopy Core). 10X images were collected on an

EVOS FL Auto Imaging System and stitched using EVOS software (UPenn Cell and

Developmental Microscopy Core). Exposure and gain were optimized within each experiment

for one virus at the 72 h time point and applied identically to each image within that experiment.

Subsequent image processing (ImageJ) was applied equally to all images in a given experiment.

Immunoblotting

Immunoblotting was performed as previously described (85). Equal amounts of whole cell lysate

were separated by SDS-PAGE. Membranes were immunoblotted with antibodies raised against

total HSV (Agilent Dako, B0114); glycoproteins (g)C, gD, gE, gH, and VP5 (all gifts from Gary

Cohen); ICP8 (gift from David Knipe); and GAPDH (GeneTex, GTX100118).

Viral DNA isolation and Illumina sequencing

Viral nucleocapsid DNA for genome sequencing was prepared by infecting MRC-5 cells at an

MOI ≥5 as previously described (86, 87). Viral nucleocapsid gDNA was sheared using a Covaris

M220 sonicator/disruptor under the following conditions: 60s duration, peak power 50, 10% duty

cycle, at 4°C. Barcoded sequencing libraries were prepared using the Illumina TruSeq low-

450 throughput protocol according to manufacturer's specifications and as previously described (60,

451 88). The quality of sequencing libraries was evaluated by Qubit (Invitrogen, CA), Bioanalyzer

452 (Agilent), and qPCR (KAPA Biosystems). Paired-end sequencing (2 × 300bp length) was

453 performed on an Illumina MiSeq, according to manufacturer's recommendations (17pM input).

De novo genome assembly

455 A consensus genome was assembled for each viral isolate using a previously described Viral

456 Genome Assembly (VirGA) bioinformatics workflow (60). Annotation of new genome

457 sequences was guided by the HSV-2 reference genome (strain HG52; GenBank NC_001798)

458 based on sequence homology (89).

Comparative genomics and phylogenetic analysis

The 10 neonatal HSV-2 genomes were aligned with all annotated HSV-2 genomes available in

GenBank (see **Table S1** for full list of 58 genomes; all derived from adults) using MAFFT (90).

The genome-wide alignment used a trimmed genome format (lacking the terminal repeats) to

avoid giving undue weight to these duplicated sequences. The MAFFT alignment was used to

generate a NeighborNet phylogenetic network in SplitsTree with Uncorrected P distances (49,

91, 92), as well as a Neighbor-Joining tree (Jukes-Cantor, 1000 bootstraps) in MEGA Omega for

Windows (93). A diverse subset of ten adult HSV-2 isolates was selected for protein-level

comparisons with the ten neonatal isolates (indicated in **Table S1** with an asterisk). ClustalW2

was used to construct pairwise nucleotide alignments between whole genomes, and pairwise

amino acid alignments for each gene and protein (94). Pan-HSV2 comparisons excluded three

viral proteins for which sequences are not fully determined in most published strains, likely due

to the high G+C-content and numerous tandem repeats in these regions: ICP34.5 (RL1 –

annotated/complete in only 9 genomes from **Table S1**), ICP0 (RL2 – annotated/complete in only

8 genomes from **Table S1**), and ICP4 (RS1 – annotated/complete in only 6 genomes from **Table**

S1) (80–82). Custom Python scripts were used on these alignments to identify nucleotide and

AA differences between samples.

Minority variant detection & quantification

Minority variants (MV) were detected using VarScan v2.2.11 (mpileup2snp and mpileup2indel

commands) (95), using the following parameters to differentiate true MV from technical artifacts

(26): minimum allele frequency ≥ 0.02 (2%); base call quality ≥ 20; read depth ≥ 100;

independent reads supporting minor allele ≥ 5. MV with directional strand bias $\geq 90\%$ were

excluded. The genomic location and potential impact of each MV was assessed using SnpEff and

SnpSift (96, 97). We use the conservative cutoff of 2% for detection of minority variants both to

avoid false positive signals and to provide data that is comparable to recent examples in the

genomic analysis of herpesviruses in clinical samples (98, 28, 99).

Acknowledgements

We thank Penny Jester and the UAB clinical support team for their assistance in retrieving

archival records. The UPenn Cell and Developmental Biology Microscopy Core provided

imaging assistance, and Elise M. Peaurori provided technical assistance during early stages of

this project. We thank members of the Szpara and Weitzman labs for helpful discussions. This

research was supported by NIAID grant 1R21AI140443 (MLS and MDW), with additional

support from the following: startup funds from the Pennsylvania State University (MLS), a

CURE grant from the Pennsylvania Department of Health (MLS), a grant from the National

Institutes of Health (NS082240 to MDW), and the NICHD Pediatric Scientist Development grant

K12-HD000850 (LNA). The published NIAID CASG clinical studies of neonatal HSV infection

were funded by a contract (N01-AI-62554) with the NIAID Development and Applications

Branch and by grants from the General Clinical Research Center Program (RR-032) and the state

of Alabama.

499   Author Contributions

500   Writing – Original Draft: LNA, CB, AND, and MLS; Writing – Review & Editing: LNA, MDW,

501   and MLS; Conceptualization: LNA and MLS; Investigation: LNA, CB, AND, DWR, UP;

502   Resources: DWK, MNP, and RJW; Funding Acquisition: MLS, RJW, and MDW.

503   Declaration of Interests

504   The authors declare no competing interests.

Figure Legends

Figure 1. Neonatal HSV-2 isolates generate different sized plaques in culture.

(**A**) Representative plaques are shown after virus incubation on Vero cells for 100 h. The

previously described low-passage adult HSV-2 strain SD90e is shown for comparison (64). Scale

bar = 5mm. (**B**) Quantification of plaque area on Vero cells. Dots represent 100 individually

measured plaques and black bar represents the mean. Each green isolate (large-plaque) is

statistically larger than each black isolate (small-plaque). Black isolates are not statistically

different from one another. Additionally, each CSF-derived isolate is statistically larger than all

other isolates shown. For all statistics, p<0.05 by one-way ANOVA followed by Holm-Sidak's

multiple comparisons test.

Figure 2. Increased plaque size in culture is not determined by viral entry, DNA

replication, protein expression, or infectious virus production.

Viral growth characteristics were compared for representative neonatal isolates, including large-

plaque-formers (green) and small-plaque-formers (black). (**A-B**) Viral entry kinetics. (**A**) Viral

isolates were applied to Vero cell monolayers at 4°C for 1 h to allow virus binding, then moved

to 37°C to allow virus entry. Extracellular virus was inactivated by a low-pH buffer at the times

indicated (orange arrowheads). Cell monolayers were washed and overlaid with methylcellulose.

Plaques were scored after 100 h of incubation. (**B**) Viral entry was quantified as the fraction of

plaques formed following citrate buffer application, where 100% is the number of plaques

formed on a monolayer not treated with citrate buffer (control). These data represent three

independent experiments. Two-way ANOVA followed by Tukey's multiple comparison test was

applied. (**C-E**) Single-cycle viral replication kinetics. Vero cell monolayers were infected at

MOI=5 and incubated in the presence of 0.1% human serum. Cell monolayers were harvested at

the time points indicated. **(C)** The quantity of viral genomes present was evaluated by qPCR for

UL27. Infectious virion production (titer) was evaluated by plaque formation on U2OS **(D)** or

Vero **(E)** cells. These data represent three independent experiments. Two-way ANOVA followed

by Tukey's multiple comparison test was applied. **(F-G)** Protein production. Vero cell

monolayers were infected at MOI=5 for 6 h **(F)** or 24 h **(G)**. Whole cell lysates were subjected to

immunoblot analysis with the following antibodies: gC/gD/gE/gH, four virion glycoproteins;

VP5 (UL19), capsid protein; ICP8 (UL29), viral single-strand DNA-binding protein; HSV, viral

antibody against whole HSV-1; and GAPDH, cellular glyceraldehyde-3 phosphate

dehydrogenase as a loading control.

Figure 3. Enhanced viral cell-to-cell spread contributes to increased plaque size in

culture.

(**A**) The rate of viral spread in Vero cells was compared for representative neonatal isolates,

including large-plaque-formers (green) and small-plaque-formers (black). Vero cell monolayers

were infected at MOI=0.001 in the presence of 0.1% human serum, which was replenished every

24 h. Samples were harvested at each time point and viral titer was evaluated by plaque

formation on U2OS cells **(B)** or Vero cells **(C).** These data represent three independent

experiments. Two-way ANOVA followed by Tukey's multiple comparison test, $*p<0.0001$ at 72

h. **(D)** In parallel experiments, HSV positive cells (green) were evaluated by

immunofluorescence. Cell nuclei are counterstained with DAPI (blue). Scale bar = 200 μm.

Images are representative of three independent experiments. Images of the entire 10 mm

coverslips were then captured and stitched to create a composite image (see **Figure S2**). (**E**) The

total number of immunofluorescent (green) pixels was quantified for each coverslip. Two-way

ANOVA followed by Tukey's multiple comparison test, *$p<0.05$ at 72 h.

Figure 4. Neonatal HSV-2 genomes are genetically distinct from one another and

encompass a broad range of known HSV-2 genetic diversity.

A phylogenetic network constructed among neonatal HSV-2 genomes (**A**), or neonatal and adult

HSV-2 genomes (**B**), reveals the wide genetic distribution of these unrelated isolates. HSV-2

genomes have been previously noted to lack geographic separation into clades (41, 42, 82). The

network was created using SplitsTree4, from a MAFFT trimmed genome alignment. See **Figure**

S3 for a comparison tree constructed using a neighbor-joining (NJ) algorithm. See **Table S1** for a

complete list of accessions, geographic origins, and references for all 58 adult HSV2 strains.

Figure 5: Neonatal HSV-2 proteins contain a similar degree of consensus-level

amino acid (AA) differences as adult HSV-2 isolates.

HSV-2 proteins are grouped by function, and the percentage of variable AAs in each protein

(#AA differences divided by protein length) are plotted for the 10 neonatal isolates (dark blue),

for 10 representative adult HSV-2 isolates (light blue), and for all 58 annotated adult HSV-2

genomes in GenBank (clear outlines behind light blue boxes). All adult HSV-2 genomes used for

this comparison are listed in **Table S1**.

Figure 6: Minority variants expand the range of neonatal HSV-2 coding diversity.

(**A**) Scatter plot indicates the total number of minor variants (MV; y-axis) observed in each

neonatal isolate. MV are rare alleles that exist within each viral population, below the level of the

consensus genome. The total number of MV on the left is separated into single-nucleotide

polymorphism (SNP) vs. insertion/deletion (INDEL) variants on the right (x-axis). The genomic

location of each SNP or INDEL variant is also summarized: intergenic vs. inside of genes (genic)

for INDELS, and intergenic vs. nonsynonymous or synonymous SNPs inside of genes. (**B**) The

frequency, or penetrance, of each minority variant was examined for each isolate. Data (x-axis)

was binned in increments of 5% (e.g. 0-<5% frequency, 5-<10% frequency, and so on), and

plotted according to the number of MV observed at each frequency (y-axis). SNP and INDEL

variants were combined for this analysis. (**C**) Stacked histograms show the number of genic MV

(x-axis) located in each HSV-2 coding sequence (gene; y-axis). SNP and INDEL variants were

combined for this analysis. UL3, UL11, UL35, and UL55 lacked any minority variants and are

not included in the histogram. See **Table S3** for full list of SNP and INDEL MV position and

frequency data.

Figure 7. Coding variations identified in neonatal HSV-2 isolates in proteins known

to contribute to cell-to-cell spread and neurovirulence.

The domain structure of each HSV protein is based on published literature for both HSV-1 and -

2. Red arrows indicate protein-coding variations discussed in the text, with the BLOSUM80

score for each AA substitution listed in parenthesis beneath the highlighted difference. More

detailed information regarding cell-to-cell spread and neurovirulence can be found in **Table S4**.

References

1. Looker KJ, Magaret AS, May MT, Turner KM, Vickerman P, Newman LM, Gottlieb SL.
2017. First estimates of the global and regional incidence of neonatal herpes infection.
Lancet Glob Health 5:e300–e309.

2. Stephenson-Famy A, Gardella C. 2014. Herpes Simplex Virus Infection During Pregnancy.
Obstet Gynecol Clin North Am 41:601–614.

3. Brown ZA, Wald A, Morrow RA, Selke S, Zeh J, Corey L. 2003. Effect of serologic status
and cesarean delivery on transmission rates of herpes simplex virus from mother to infant.
Jama 289:203–209.

4. Corey L, Wald A. 2009. Maternal and Neonatal HSV Infections. N Engl J Med 361:1376–
1385.

5. Bradley H, Markowitz LE, Gibson T, McQuillan GM. 2014. Seroprevalence of herpes
simplex virus types 1 and 2--United States, 1999-2010. J Infect Dis 209:325–333.

6. James SH, Kimberlin DW. 2014. Neonatal Herpes Simplex Virus Infection. Clin Perinatol.

7. Pinninti SG, Kimberlin DW. 2014. Preventing Herpes Simplex Virus in the Newborn. Clin
Perinatol 41:945–955.

8. Kimberlin DW, Lin CY, Jacobs RF, Powell DA, Frenkel LM, Gruber WC, Rathore M,
Bradley JS, Diaz PS, Kumar M, Arvin AM, Gutierrez K, Shelton M, Weiner LB, Sleasman
JW, de Sierra TM, Soong SJ, Kiell J, Lakeman FD, Whitley RJ, National Institute of
Allergy and Infectious Diseases Collaborative Antiviral Study Group. 2001. Natural history
of neonatal herpes simplex virus infections in the acyclovir era. Pediatrics 108:223–229.

9. Kimberlin DW, Whitley RJ, Wan W, Powell DA, Storch G, Ahmed A, Palmer A, Sánchez
PJ, Jacobs RF, Bradley JS. 2011. Oral acyclovir suppression and neurodevelopment after
neonatal herpes. N Engl J Med 365:1284–1292.

10. Casanova J-L. 2015. Human genetic basis of interindividual variability in the course of
infection. Proc Natl Acad Sci 201521644.

11. Zhang S-Y, Jouanguy E, Ugolini S, Smahi A, Elain G, Romero P, Segal D, Sancho-Shimizu
V, Lorenzo L, Puel A, Picard C, Chapgier A, Plancoulaine S, Titeux M, Cognet C, von
Bernuth H, Ku C-L, Casrouge A, Zhang X-X, Barreiro L, Leonard J, Hamilton C, Lebon P,
Heron B, Vallee L, Quintana-Murci L, Hovnanian A, Rozenberg F, Vivier E, Geissmann F,
Tardieu M, Abel L, Casanova J-L. 2007. TLR3 deficiency in patients with herpes simplex
encephalitis. Science 317:1522–1527.

12. Whitley R, Arvin A, Prober C, Corey L, Burchett S, Plotkin S, Starr S, Jacobs R, Powell D,
Nahmias A, Sumaya C, Edwards K, Alford C, Caddell G, Soong S-J, the National Institute
of Allergy and Infectious Diseases Collaborative Antiviral Study Group*. 1991. Predictors
of Morbidity and Mortality in Neonates with Herpes Simplex Virus Infections. N Engl J
Med 324:450–454.

13. Fields BN, Byers K. 1983. The Genetic Basis of Viral Virulence. Philos Trans R Soc Lond B
Biol Sci 303:209–218.

14. Gonzalez-Scarano F, Beaty B, Sundin D, Janssen R, Endres MJ, Nathanson N. 1988. Genetic
determinants of the virulence and infectivity of La Crosse virus. Microb Pathog 4:1–7.

15. Tscherne DM, García-Sastre A. 2011. Virulence determinants of pandemic influenza viruses.
J Clin Invest 121:6–13.

16. Müller V, Fraser C, Herbeck JT. 2011. A Strong Case for Viral Genetic Factors in HIV
Virulence. Viruses 3:204–216.

17. Robinson CM, Jesudhasan PR, Pfeiffer JK. 2014. Bacterial Lipopolysaccharide Binding
Enhances Virion Stability and Promotes Environmental Fitness of an Enteric Virus. Cell
Host Microbe 15:36–46.

18. Yoshii K, Sunden Y, Yokozawa K, Igarashi M, Kariwa H, Holbrook MR, Takashima I.
2014. A Critical Determinant of Neurological Disease Associated with Highly Pathogenic
Tick-Borne Flavivirus in Mice. J Virol 88:5406–5420.

19. Roizman B, Whitley RJ. 2001. The nine ages of herpes simplex virus. Herpes 8:23–27.

20. Sakaoka H, Kawana T, Grillner L, Aomori T, Yamiguchi T, Saito H, Fujinaga K. 1987.
Genome Variations in Herpes Simplex Virus Type 2 Strains Isolated in Japan and Sweden.
J Gen Virol 68:2105–16.

21. Sakaoka H, Kurita K, Iida Y, Takada S, Umene K, Kim YT, Ren CS, Nahmias AJ. 1994.
Quantitative analysis of genomic polymorphism of herpes simplex virus type 1 strains from
six countries: studies of molecular evolution and molecular epidemiology of the virus. J
Gen Virol 75:513–27.

22. Bowden R, Sakaoka H, Ward R, Donnelly P. 2006. Patterns of Eurasian HSV-1 molecular
diversity and inferences of human migrations. Infect Genet Evol 6:63–74.

23. Bower JR, Mao H, Durishin C, Rozenbom E, Detwiler M, Rempinski D, Karban TL,
Rosenthal KS. 1999. Intrastrain variants of herpes simplex virus type 1 isolated from a
neonate with fatal disseminated infection differ in the ICP34.5 gene, glycoprotein
processing, and neuroinvasiveness. J Virol 73:3843–53.

24. Mao H, Rosenthal KS. 2003. Strain-Dependent Structural Variants of Herpes Simplex Virus
Type 1 ICP34.5 Determine Viral Plaque Size, Efficiency of Glycoprotein Processing, and
Viral Release and Neuroinvasive Disease Potential. J Virol 77:3409–3417.

25. Renzette N, Gibson L, Jensen JD, Kowalik TF. 2014. Human cytomegalovirus intrahost
evolution—a new avenue for understanding and controlling herpesvirus infections. Curr
Opin Virol 8:109–115.

26. Depledge DP, Kundu S, Jensen NJ, Gray ER, Jones M, Steinberg S, Gershon A, Kinchington
PR, Schmid DS, Balloux F, Nichols RA, Breuer J. 2014. Deep Sequencing of Viral
Genomes Provides Insight into the Evolution and Pathogenesis of Varicella Zoster Virus
and Its Vaccine in Humans. Mol Biol Evol 31:397–409.

27. Renner DW, Szpara ML. 2018. The impacts of genome-wide analyses on our understanding
of human herpesvirus diversity and evolution. J Virol 92:e00908-17.

28. Houldcroft CJ, Bryant JM, Depledge DP, Margetts BK, Simmonds J, Nicolaou S, Tutill HJ,
Williams R, Worth AJJ, Marks SD, Veys P, Whittaker E, Breuer J. 2016. Detection of Low
Frequency Multi-Drug Resistance and Novel Putative Maribavir Resistance in
Immunocompromised Pediatric Patients with Cytomegalovirus. Front Microbiol 7.

29. Houldcroft CJ, Beale MA, Breuer J. 2017. Clinical and biological insights from viral genome
sequencing. Nat Rev Microbiol 15:183–192.

30. Renzette N, Bhattacharjee B, Jensen JD, Gibson L, Kowalik TF. 2011. Extensive genome-
wide variability of human cytomegalovirus in congenitally infected infants. PLoS Pathog
7:1–14.

31. Renzette N, Gibson L, Bhattacharjee B, Fisher D, Schleiss MR, Jensen JD, Kowalik TF.
2013. Rapid intrahost evolution of human cytomegalovirus is shaped by demography and
positive selection. PLoS Genet 9:1–14.

32. Renzette N, Pokalyuk C, Gibson L, Bhattacharjee B, Schleiss MR, Hamprecht K, Yamamoto
AY, Mussi-Pinhata MM, Britt WJ, Jensen JD, Kowalik TF. 2015. Limits and patterns of
cytomegalovirus genomic diversity in humans. Proc Natl Acad Sci 201501880.

33. Renzette N, Kowalik TF, Jensen JD. 2016. On the relative roles of background selection and
genetic hitchhiking in shaping human cytomegalovirus genetic diversity. Mol Ecol 25:403–
413.

34. Renzette N, Pfeifer SP, Matuszewski S, Kowalik TF, Jensen JD. 2017. On the Analysis of
Intrahost and Interhost Viral Populations: Human Cytomegalovirus as a Case Study of
Pitfalls and Expectations. J Virol 91:e01976-16.

35. Pokalyuk C, Renzette N, Irwin KK, Pfeifer SP, Gibson L, Britt WJ, Yamamoto AY, Mussi-
Pinhata MM, Kowalik TF, Jensen JD. 2017. Characterizing human cytomegalovirus
reinfection in congenitally infected infants: an evolutionary perspective. Mol Ecol.

36. Sijmons S, Van Ranst M, Maes P. 2014. Genomic and Functional Characteristics of Human
Cytomegalovirus Revealed by Next-Generation Sequencing. Viruses 6:1049–1072.

37. Sijmons S, Thys K, Mbong Ngwese M, Van Damme E, Dvorak J, Van Loock M, Li G,
Tachezy R, Busson L, Aerssens J, Van Ranst M, Maes P. 2015. High-Throughput Analysis
of Human Cytomegalovirus Genome Diversity Highlights the Widespread Occurrence of
Gene-Disrupting Mutations and Pervasive Recombination. J Virol 89:7673–7695.

38. Lassalle F, Depledge DP, Reeves MB, Brown AC, Christiansen MT, Tutill HJ, Williams RJ,
Einer-Jensen K, Holdstock J, Atkinson C, Brown JR, van Loenen FB, Clark DA, Griffiths
PD, Verjans GMGM, Schutten M, Milne RSB, Balloux F, Breuer J. 2016. Islands of
linkage in an ocean of pervasive recombination reveals two-speed evolution of human
cytomegalovirus genomes. Virus Evol 2:vew017.

39. Johnston C, Magaret A, Roychoudhury P, Greninger AL, Reeves D, Schiffer J, Jerome KR,
Sather C, Diem K, Lingappa JR, Celum C, Koelle DM, Wald A. 2017. Dual-strain genital
herpes simplex virus type 2 (HSV-2) infection in the US, Peru, and 8 countries in sub-

Saharan Africa: A nested cross-sectional viral genotyping study. PLOS Med 14:e1002475.

40. Minaya MA, Jensen TL, Goll JB, Korom M, Datla SH, Belshe RB, Morrison LA. 2017.
Molecular evolution of herpes simplex virus 2 complete genomes: comparison between
primary and recurrent infections. J Virol 91:e00942–17.

41. Koelle DM, Norberg P, Fitzgibbon MP, Russell RM, Greninger AL, Huang M-L, Stensland
L, Jing L, Magaret AS, Diem K, Selke S, Xie H, Celum C, Lingappa JR, Jerome KR, Wald
A, Johnston C. 2017. Worldwide circulation of HSV-2 × HSV-1 recombinant strains. Sci
Rep 7:44084.

42. Burrel S, Boutolleau D, Ryu D, Agut H, Merkel K, Leendertz FH, Calvignac-Spencer S.
2017. Ancient Recombination Events between Human Herpes Simplex Viruses. Mol Biol
Evol 34:1713–1721.

43. Ryncarz AJ, Goddard J, Wald A, Huang M-L, Roizman B, Corey L. 1999. Development of a
high-throughput quantitative assay for detecting herpes simplex virus DNA in clinical
samples. J Clin Microbiol 37:1941–1947.

44. Wald A, Huang M-L, Carrell D, Selke S, Corey L. 2003. Polymerase chain reaction for
detection of herpes simplex virus (HSV) DNA on mucosal surfaces: comparison with HSV
isolation in cell culture. J Infect Dis 188:1345–1351.

45. Szpara ML, Tafuri YR, Parsons L, Shamim SR, Verstrepen KJ, Legendre M, Enquist LW.
2011. A wide extent of inter-strain diversity in virulent and vaccine strains of
alphaherpesviruses. PLoS Pathog 7:1–23.

46. Brown J. 2004. Effect of gene location on the evolutionary rate of amino acid substitutions in
herpes simplex virus proteins. Virology 330:209–20.

47. Bowden R, Sakaoka H, Donnelly P, Ward R. 2004. High recombination rate in herpes
simplex virus type 1 natural populations suggests significant co-infection. Infect Genet Evol
4:115–23.

48. Norberg P, Bergstrom T, Rekabdar E, Lindh M. 2004. Phylogenetic Analysis of Clinical
Herpes Simplex Virus Type 1 Isolates Identified Three Genetic Groups and Recombinant
Viruses. J Virol 78:10755–10764.

49. Norberg P, Kasubi MJ, Haarr L, Bergstrom T, Liljeqvist J-A. 2007. Divergence and
recombination of clinical herpes simplex virus type 2 isolates. J Virol 81:13158–13167.

50. Lonsdale DM, Brown SM, Lang J, Subak-Sharpe JH, Koprowski H, Warren KG. 1980.
Variations in herpes simplex virus isolated from human ganglia and a study of clonal
variation in HSV-1. Ann N Y Acad Sci 291–308.

51. Liljeqvist JA, Tunback P, Norberg P. 2009. Asymptomatically shed recombinant herpes
simplex virus type 1 strains detected in saliva. J Gen Virol 90:559–566.

52. Haarr L, Nilsen A, Knappskog PM, Langeland N. 2014. Stability of glycoprotein gene
sequences of herpes simplex virus type 2 from primary to recurrent human infection, and
diversity of the sequences among patients attending an STD clinic. BMC Infect Dis 14:63.

53. Szpara ML, Gatherer D, Ochoa A, Greenbaum B, Dolan A, Bowden RJ, Enquist LW,
Legendre M, Davison AJ. 2014. Evolution and diversity in human herpes simplex virus
genomes. J Virol 88:1209–27.

54. Parsons LR, Tafuri YR, Shreve JT, Bowen CD, Shipley MM, Enquist LW, Szpara ML.
2015. Rapid genome assembly and comparison decode intrastrain variation in human
alphaherpesviruses. mBio 6:e02213-14.

55. Kolb AW, Lee K, Larsen I, Craven M, Brandt CR. 2016. Quantitative Trait Locus Based
Virulence Determinant Mapping of the HSV-1 Genome in Murine Ocular Infection: Genes
Involved in Viral Regulatory and Innate Immune Networks Contribute to Virulence. PLOS
Pathog 12:e1005499.

56. Pfaff F, Groth M, Sauerbrei A, Zell R. 2016. Genotyping of herpes simplex virus type 1
(HSV-1) by whole genome sequencing. J Gen Virol.

57. Bowen CD, Renner DW, Shreve JT, Tafuri Y, Payne KM, Dix RD, Kinchington PR,
Gatherer D, Szpara ML. 2016. Viral forensic genomics reveals the relatedness of classic
herpes simplex virus strains KOS, KOS63, and KOS79. Virology 492:179–186.

58. Pandey U, Renner DW, Thompson RL, Szpara ML, Sawtell NM. 2017. Inferred father-to-
son transmission of herpes simplex virus results in near-perfect preservation of viral
genome identity and in vivo phenotypes. Sci Rep 7:13666.

59. Szpara ML, Parsons L, Enquist LW. 2010. Sequence variability in clinical and laboratory
isolates of herpes simplex virus 1 reveals new mutations. J Virol 84:5303–13.

60. Parsons LR, Tafuri YR, Shreve JT, Bowen CD, Shipley MM, Enquist LW, Szpara ML.
2015. Rapid genome assembly and comparison decode intrastrain variation in human
alphaherpesviruses. mBio 6:e02213-14.

61. Kimberlin DW, Lin CY, Jacobs RF, Powell DA, Corey L, Gruber WC, Rathore M, Bradley
JS, Diaz PS, Kumar M, Arvin AM, Gutierrez K, Shelton M, Weiner LB, Sleasman JW, de
Sierra TM, Weller S, Soong SJ, Kiell J, Lakeman FD, Whitley RJ, National Institute of
Allergy and Infectious Diseases Collaborative Antiviral Study Group. 2001. Safety and
efficacy of high-dose intravenous acyclovir in the management of neonatal herpes simplex
virus infections. Pediatrics 108:230–238.

62. Desmyter J, Melnick JL, Rawls WE. 1968. Defectiveness of interferon production and of
rubella virus interference in a line of African green monkey kidney cells (Vero). J Virol
2:955–961.

63. Emeny JM, Morgan MJ. 1979. Regulation of the interferon system: evidence that Vero cells
have a genetic defect in interferon production. J Gen Virol 43:247–252.

64. Colgrove R, Diaz F, Newman R, Saif S, Shea T, Young S, Henn M, Knipe DM. 2014.
Genomic sequences of a low passage herpes simplex virus 2 clinical isolate and its plaque-
purified derivative strain. Virology 450–451:140–145.

65. Yao F, Schaffer PA. 1995. An activity specified by the osteosarcoma line U2OS can
substitute functionally for ICP0, a major regulatory protein of herpes simplex virus type 1. J
Virol 69:6249–6258.

66. Deschamps T, Kalamvoki M. 2017. Impaired STING Pathway in Human Osteosarcoma
U2OS Cells Contributes to the Growth of ICP0-Null Mutant Herpes Simplex Virus. J Virol
91:e00006-17.

67. Norberg P. 2010. Divergence and genotyping of human alpha-herpesviruses: an overview.
Infect Genet Evol J Mol Epidemiol Evol Genet Infect Dis 10:14–25.

68. David AT, Saied A, Charles A, Subramanian R, Chouljenko VN, Kousoulas KG. 2012. A
Herpes Simplex Virus 1 (McKrae) Mutant Lacking the Glycoprotein K Gene Is Unable To
Infect via Neuronal Axons and Egress from Neuronal Cell Bodies. mBio 3.

69. Dingwell KS, Brunetti CR, Hendricks RL, Tang Q, Tang M, Rainbow AJ, Johnson DC.
1994. Herpes simplex virus glycoproteins E and I facilitate cell-to-cell spread in vivo and
across junctions of cultured cells. J Virol 68:834–845.

70. Weber PC, Levine M, Glorioso JC. 1987. Rapid identification of nonessential genes of
herpes simplex virus type 1 by Tn5 mutagenesis. Science 236:576–579.

71. Jacobson JG, Martin SL, Coen DM. 1989. A conserved open reading frame that overlaps the
herpes simplex virus thymidine kinase gene is important for viral growth in cell culture. J
Virol 63:1839–1843.

72. Foster TP, Melancon JM, Baines JD, Kousoulas KG. 2004. The herpes simplex virus type 1
UL20 protein modulates membrane fusion events during cytoplasmic virion morphogenesis
and virus-induced cell fusion. J Virol 78:5347–5357.

73. David AT, Baghian A, Foster TP, Chouljenko VN, Kousoulas KG. 2008. The herpes simplex
virus type 1 (HSV-1) glycoprotein K(gK) is essential for viral corneal spread and
neuroinvasiveness. Curr Eye Res 33:455–467.

74. Dingwell KS, Johnson DC. 1998. The herpes simplex virus gE-gI complex facilitates cell-to-
cell spread and binds to components of cell junctions. J Virol 72:8933–8942.

75. Meignier B, Longnecker R, Mavromara-Nazos P, Sears AE, Roizman B. 1988. Virulence of
and establishment of latency by genetically engineered deletion mutants of herpes simplex
virus 1. Virology 162:251–254.

76. Rochette P-A, Bourget A, Sanabria-Solano C, Lahmidi S, Lavallée GO, Pearson A. 2015.
Mutation of UL24 impedes the dissemination of acute herpes simplex virus 1 infection
from the cornea to neurons of trigeminal ganglia. J Gen Virol 96:2794–2805.

77. Jacobson JG, Chen SH, Cook WJ, Kramer MF, Coen DM. 1998. Importance of the herpes
simplex virus UL24 gene for productive ganglionic infection in mice. Virology 242:161–
169.

78. Leiva-Torres GA, Rochette P-A, Pearson A. 2010. Differential importance of highly
conserved residues in UL24 for herpes simplex virus 1 replication in vivo and reactivation.
J Gen Virol 91:1109–1116.

79. Liu J, Lewin AS, Tuli SS, Ghivizzani SC, Schultz GS, Bloom DC. 2008. Reduction in
severity of a herpes simplex virus type 1 murine infection by treatment with a ribozyme
targeting the UL20 gene RNA. J Virol 82:7467–7474.

80. Newman RM, Lamers SL, Weiner B, Ray SC, Colgrove RC, Diaz F, Jing L, Wang K, Saif S,
Young S, Henn M, Laeyendecker O, Tobian AAR, Cohen JI, Koelle DM, Quinn TC, Knipe
DM. 2015. Genome Sequencing and Analysis of Geographically Diverse Clinical Isolates
of Herpes Simplex Virus 2. J Virol JVI.01303-15.

81. Kolb AW, Larsen IV, Cuellar JA, Brandt CR. 2015. Genomic, Phylogenetic, and
Recombinational Characterization of Herpes Simplex Virus 2 Strains. J Virol 89:6427–
6434.

82. Johnston C, Magaret A, Roychoudhury P, Greninger AL, Cheng A, Diem K, Fitzgibbon MP,
Huang M, Selke S, Lingappa JR, Celum C, Jerome KR, Wald A, Koelle DM. 2017. Highly
conserved intragenic HSV-2 sequences: Results from next-generation sequencing of HSV-2
U L and U S regions from genital swabs collected from 3 continents. Virology 510:90–98.

83. Minaya MA, Jensen T, Goll J, Korom M, Datla SH, Belshe RB, Morrison LA. 2017.
Molecular evolution of herpes simplex virus 2 complete genomes: Comparison between
primary and recurrent infections. J Virol JVI.00942-17.

84. Whitley R, Arvin A, Prober C, Burchett S, Corey L, Powell D, Plotkin S, Starr S, Alford C,
Connor J, Jacobs R, Nahmias A, Soong S-J, the National Institute of Allergy, Infectious
Diseases Collaborative Antiviral Study Group*. 1991. A Controlled Trial Comparing
Vidarabine with Acyclovir in Neonatal Herpes Simplex Virus Infection. N Engl J Med
324:444–449.

85. Avgousti DC, Della Fera AN, Otter CJ, Herrmann C, Pancholi NJ, Weitzman MD. 2017.
Adenovirus Core Protein VII Downregulates the DNA Damage Response on the Host
Genome. J Virol 91:e01089-17.

86. Szpara ML, Tafuri YR, Enquist LW. 2011. Preparation of viral DNA from nucleocapsids. J
Vis Exp JoVE 2–7.

87. Szpara M. 2014. Isolation of herpes simplex virus nucleocapsid DNA, p. 31–41. *In*
Diefenbach, RJ, Fraefel, C (eds.), Herpes Simplex Virus. Springer New York, New York,
NY.

88. Bowen CD, Renner DW, Shreve JT, Tafuri Y, Payne KM, Dix RD, Kinchington PR,
Gatherer D, Szpara ML. 2016. Viral forensic genomics reveals the relatedness of classic
herpes simplex virus strains KOS, KOS63, and KOS79. Virology 492:179–186.

89. Dolan A, Jamieson FE, Cunningham C, Barnett BC, McGeoch DJ. 1998. The genome
sequence of herpes simplex virus type 2. J Virol 72:2010–2021.

90. Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–3066.

91. Strimmer K, Wiuf C, Moulton V. 2001. Recombination analysis using directed graphical
models. Mol Biol Evol 18:97–99.

92. Norberg P, Tyler S, Severini A, Whitley R, Liljeqvist J-A, Bergstrom T. 2011. A genome-
wide comparative evolutionary analysis of herpes simplex virus type 1 and varicella zoster
virus. PloS One 6:1–8.

93. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular
Evolutionary Genetics Analysis Version 6.0. Mol Biol Evol 30:2725–2729.

94. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin
F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. 2007. Clustal W
and Clustal X version 2.0. Bioinforma Appl Note 23:2947–2948.

95. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER,
Ding L, Wilson RK. 2012. VarScan 2: Somatic mutation and copy number alteration
discovery in cancer by exome sequencing. Genome Res 22:568–576.

96. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM.
2012. A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff. Fly (Austin) 6:80–92.

97. Cingolani P, Patel VM, Coon M, Nguyen T, Land SJ, Ruden DM, Lu X. 2012. Using
Drosophila melanogaster as a Model for Genotoxic Chemical Mutational Studies with a
New Program, SnpSift. Front Genet 3.

98. Depledge DP, Yamanishi K, Gomi Y, Gershon AA, Breuer J. 2016. Deep Sequencing of
Distinct Preparations of the Live Attenuated Varicella-Zoster Virus Vaccine Reveals a
Conserved Core of Attenuating Single-Nucleotide Polymorphisms. J Virol 90:8698–8704.

99. Hage E, Wilkie GS, Linnenweber-Held S, Dhingra A, Suárez NM, Schmidt JJ, Kay-Fedorov
PC, Mischak-Weissinger E, Heim A, Schwarz A, Schulz TF, Davison AJ, Ganzenmueller
T. 2017. Characterization of Human Cytomegalovirus Genome Diversity in
Immunocompromised Hosts by Whole-Genome Sequencing Directly From Clinical
Specimens. J Infect Dis 215:1673–1683.

Tables

Table 1: Clinical characteristics associated with HSV-2 isolates from ten patients.

Clinical Isolate*	Clinical Disease at Diagnosis	Sample Source	Morbidity Score - Mental	Morbidity Score - Motor	Patient Age at Disease Onset (days)	Gestational Age at Time of Birth (weeks)	Patient Sex & Race
CNS11	CNS	CSF	4	4	12	37	M, W
DISS14	DISS + CNS	CSF	2	2	7	39	M, W
CNS03	CNS	SKIN	4	4	17	37	M, W
CNS15	CNS	SKIN	3	4	19	36	M, W
CNS17	CNS	SKIN	4	4	17	40	M, W
DISS29	DISS + CNS	SKIN	3	3	5	38	F, B
CNS12	CNS	SKIN	4	4	16	41	F, W
SEM02	SEM	SKIN	1	1	5	38	F, W
SEM13	SEM	SKIN	4	4	11	27	F, B
SEM18	SEM	SKIN	2	2	17	37	F, W

884 * Clinical isolate order based on **Figure 1**.

887   **Table 2. Genome sequencing statistics for neonatal HSV-2 strains.**

Clinical Isolate*	Average Coverage	Raw Sequence Reads	Used for Assembly**	% viral	Depth > 100	GenBank Accession
CNS11	6,081X	3.5 million	2.8 million	79%	96%	In process^
DISS14	6,267X	3.8 million	3.0 million	77%	97%	In process^
CNS03	6,269X	3.9 million	3.1 million	78%	96%	In process^
CNS15	7,486X	6.1 million	4.4 million	73%	97%	In process^
CNS17	5,059X	3.1 million	2.3 million	73%	96%	In process^
DISS29	2,588X	1.4 million	1.1 million	78%	99%	In process^
CNS12	5,839X	3.9 million	2.8 million	73%	96%	In process^
SEM02	6,455X	4.8 million	3.7 million	77%	95%	In process^
SEM13	5,291X	3.4 million	2.4 million	72%	89%	In process^
SEM18	7,514X	16.9 million	12.2 million	72%	98%	In process^

* Clinical isolate order based on **Figure 1**.
** Numbers reflect the count of sequence read 1 of paired-end reads.
^ GenBank accessions are in process. See "Related Manuscript File" uploaded for review,
which contains all 10 GenBank flatfiles.

Figures

Figure 1. Neonatal HSV-2 isolates generate different sized plaques in culture.

(**A**) Representative plaques are shown after virus incubation on Vero cells for 100 h. The

previously described low-passage adult HSV-2 strain SD90e is shown for comparison (64). Scale

bar = 5mm. (**B**) Quantification of plaque area on Vero cells. Dots represent 100 individually

measured plaques and black bar represents the mean. Each green isolate (large-plaque) is

statistically larger than each black isolate (small-plaque). Black isolates are not statistically

different from one another. Additionally, each CSF-derived isolate is statistically larger than all

other isolates shown. For all statistics, $p < 0.05$ by one-way ANOVA followed by Holm-Sidak's

multiple comparisons test.

Figure 2. Increased plaque size in culture is not determined by viral entry, DNA replication, protein expression, or infectious virus production.

Viral growth characteristics were compared for representative neonatal isolates, including large-plaque-formers (green) and small-plaque-formers (black). **(A-B)** Viral entry kinetics. **(A)** Viral

isolates were applied to Vero cell monolayers at 4°C for 1 h to allow virus binding, then moved

to 37°C to allow virus entry. Extracellular virus was inactivated by a low-pH buffer at the times

indicated (orange arrowheads). Cell monolayers were washed and overlaid with methylcellulose.

Plaques were scored after 100 h of incubation. **(B)** Viral entry was quantified as the fraction of

plaques formed following citrate buffer application, where 100% is the number of plaques

formed on a monolayer not treated with citrate buffer (control). These data represent three

independent experiments. Two-way ANOVA followed by Tukey's multiple comparison test was

applied. **(C-E)** Single-cycle viral replication kinetics. Vero cell monolayers were infected at

MOI=5 and incubated in the presence of 0.1% human serum. Cell monolayers were harvested at

the time points indicated. **(C)** The quantity of viral genomes present was evaluated by qPCR for

UL27. Infectious virion production (titer) was evaluated by plaque formation on U2OS **(D)** or

Vero **(E)** cells. These data represent three independent experiments. Two-way ANOVA followed

by Tukey's multiple comparison test was applied. **(F-G)** Protein production. Vero cell

monolayers were infected at MOI=5 for 6 h **(F)** or 24 h **(G)**. Whole cell lysates were subjected to

immunoblot analysis with the following antibodies: gC/gD/gE/gH, four virion glycoproteins;

VP5 (UL19), capsid protein; ICP8 (UL29), viral single-strand DNA-binding protein; HSV, viral

antibody against whole HSV-1; and GAPDH, cellular glyceraldehyde-3 phosphate

dehydrogenase as a loading control.

Figure 3. Enhanced viral cell-to-cell spread contributes to increased plaque size in culture.

(**A**) The rate of viral spread in Vero cells was compared for representative neonatal isolates, including large-plaque-formers (green) and small-plaque-formers (black). Vero cell monolayers were infected at MOI=0.001 in the presence of 0.1% human serum, which was replenished every 24 h. Samples were harvested at each time point and viral titer was evaluated by plaque formation on U2OS cells (**B**) or Vero cells (**C**). These data represent three independent experiments. Two-way ANOVA followed by Tukey's multiple comparison test, *$p<0.0001$ at 72 h. (**D**) In parallel experiments, HSV positive cells (green) were evaluated by immunofluorescence. Cell nuclei are counterstained with DAPI (blue). Scale bar = 200 μm. Images are representative of three independent experiments. Images of the entire 10 mm coverslips were then captured and stitched to create a composite image (see **Figure S2**). (**E**) The total number of immunofluorescent (green) pixels was quantified for each coverslip. Two-way ANOVA followed by Tukey's multiple comparison test, *$p<0.05$ at 72 h.

Figure 4. Neonatal HSV-2 genomes are genetically distinct from one another and encompass a broad range of known HSV-2 genetic diversity.

A phylogenetic network constructed among neonatal HSV-2 genomes (**A**), or neonatal and adult HSV-2 genomes (**B**), reveals the wide genetic distribution of these unrelated isolates. HSV-2 genomes have been previously noted to lack geographic separation into clades (41, 42, 82). The network was created using SplitsTree4, from a MAFFT trimmed genome alignment. See **Figure S3** for a comparison tree constructed using a neighbor-joining (NJ) algorithm. See **Table S1** for a complete list of accessions, geographic origins, and references for all 58 adult HSV2 strains.

Figure 5: Neonatal HSV-2 proteins contain a similar degree of consensus-level amino acid (AA) differences as adult HSV-2 isolates.

HSV-2 proteins are grouped by function, and the percentage of variable AAs in each protein (#AA differences divided by protein length) are plotted for the 10 neonatal isolates (dark blue), for 10 representative adult HSV-2 isolates (light blue), and for all 58 annotated adult HSV-2 genomes in GenBank (clear outlines behind light blue boxes). All adult HSV-2 genomes used for this comparison are listed in **Table S1**.

Figure 6: Minority variants expand the range of neonatal HSV-2 coding diversity.

(**A**) Scatter plot indicates the total number of minor variants (MV; y-axis) observed in each

neonatal isolate. MV are rare alleles that exist within each viral population, below the level of the

consensus genome. The total number of MV on the left is separated into single-nucleotide

polymorphism (SNP) vs. insertion/deletion (INDEL) variants on the right (x-axis). The genomic

location of each SNP or INDEL variant is also summarized: intergenic vs. inside of genes (genic)

for INDELS, and intergenic vs. nonsynonymous or synonymous SNPs inside of genes. (**B**) The

frequency, or penetrance, of each minority variant was examined for each isolate. Data (x-axis)

was binned in increments of 5% (e.g. 0-<5% frequency, 5-<10% frequency, and so on), and

plotted according to the number of MV observed at each frequency (y-axis). SNP and INDEL

variants were combined for this analysis. (**C**) Stacked histograms show the number of genic MV

(x-axis) located in each HSV-2 coding sequence (gene; y-axis). SNP and INDEL variants were

combined for this analysis. UL3, UL11, UL35, and UL55 lacked any minority variants and are

not included in the histogram. See **Table S3** for full list of SNP and INDEL MV position and

frequency data.

Figure 7. Coding variations identified in neonatal HSV-2 isolates in proteins known to contribute to cell-to-cell spread and neurovirulence.

The domain structure of each HSV protein is based on published literature for both HSV-1 and -2. Red arrows indicate protein-coding variations discussed in the text, with the BLOSUM80 score for each AA substitution listed in parenthesis beneath the highlighted difference. More detailed information regarding cell-to-cell spread and neurovirulence can be found in **Table S4**.

987 **Supplementary Information for**

988 **Genotypic and phenotypic diversity within the neonatal HSV-2 population**

989   Lisa N. Akhtar, Christopher D. Bowen, Daniel W. Renner, Utsav Pandey[2], Ashley N. Della Fera,

990   David W. Kimberlin, Mark N. Prichard, Richard J. Whitley, Matthew D. Weitzman*,

991   Moriah L. Szpara*

992
993   ***Co-Corresponding Authors**
994   Moriah L. Szpara
995   Email: moriah@psu.edu
996
997   Matthew D. Weitzman
998   Email: weitzmanm@email.chop.edu

999
1000
1001   **This file includes:**
1002   Supplementary Text
1003   Figs. S1 to S5
1004   Tables S1, S2 and S4
1005   References for SI reference citations
1006
1007   **Other supplementary materials for this manuscript include the following:**
1008   Table S3 in separate Excel file (two sheets, listing minority variant SNPs and INDELs)
1009
1010

Supplementary Text

Virus source

Viruses analyzed were collected from neonates that were culture- or PCR-confirmed to have

HSV-2 infection, enrolled in three published clinical studies (1–3) by the Collaborative Antiviral

Study Group (CASG) at the University of Alabama Birmingham (UAB). These HSV isolates

originated from patients with SEM, CNS, or disseminated (DISS) disease with CNS involvement

(**Table 1**). Samples were collected at the time of diagnosis, prior to initiation of acyclovir

therapy, from either the cerebrospinal fluid (CSF) or skin. Viral isolates are associated with de-

identified clinical information including age, sex, race, and clinical morbidity scores, as

approved by the UAB Institutional Review Board (IRB). Enrollment in the original studies was

evenly split between males and females, and included both black and white patients. Clinical

morbidity score was determined by the CASG after 12 months of life as (1) normal; (2) mild

impairment which includes ocular sequelae (keratoconjunctivitis), speech delay, or mild motor

delay; (3) moderate impairment which includes hemiparesis, persistent seizure disorder, or

developmental delay of less than or equal to 3 months adjusted developmental age; (4) severe

impairment which includes microcephaly, spastic quadriplegia, chorioretinitis or blindness, or a

serious developmental delay of least 3 months according to the Denver Developmental

Assessment Scale (4, 5). These assessments were performed at each collaborating site. Initial

collection of samples was approved by the UAB IRB. Use of these samples in this study was

approved by the UAB IRB.

Cell culture

Cell culture

Human lung fibroblast MRC-5 cells (ATCC®, CCL-171) were cultured in minimum essential

medium Eagle (MEME, Sigma-Aldrich; M5650) supplemented with 10% fetal bovine serum

(FBS, Hyclone; SH30071.03), 2mM L-glutamine (Gibco, A2916801) and 1X penicillin-

streptomycin (Gibco; 15140122). African green monkey kidney Vero cells (ATCC®, CCL-81)

and human epithelial bone osteosarcoma U2OS cells (ATCC®, HTB-96) were cultured in

Dulbecco's modified Eagle's medium with high glucose (DMEM, Hyclone; SH30081.02)

supplemented with 10% FBS, 2mM L-glutamine, and 1X penicillin-streptomycin. Cell lines

were authenticated by ATCC prior to purchase, and were confirmed to be mycoplasma free

throughout experiments by periodic testing (LookOut Mycoplasma, Sigma).

Virus culture

Initial viral isolates (passage 1-2) were obtained from CASG samples stored at the UAB. Each

isolate had been cultured at the time of diagnosis, aliquoted after one passage, and snap frozen.

To conduct viral genome sequencing and phenotypic assays, each stock was expanded over three

serial passages by infecting monolayers of human MRC-5 cells at a MOI of 0.01. Each infection

was allowed to progress to complete cytopathic effect (CPE) before harvest (between 50-70

hours), with increasing cell volume at each passage to produce sufficient virions for subsequent

phenotypic and genomic comparisons. Viral stocks were titered on monolayers of either Vero

cells or U2OS cells, for 100 or 48 hours respectively, to allow plaques to develop. Plaque

formation was facilitated by limiting viral diffusion with a methylcellulose overlay. Plaque size

and morphology were monitored carefully and did not change for any viral isolates over the

course of virus stock expansion.

Plaque measurements

After appropriate incubation plaques were stained with 0.5% methylene blue and allowed to dry.

Serial 4X brightfield images were collected on an EVOS FL Auto Imaging System and stitched

by EVOS software to create an image of the entire well (University of Pennsylvania Cell and

Developmental Biology Microscopy Core). No processing was performed. The area of 100

plaques was measured for each viral isolate using ImageJ software.

Genome copy number estimation by quantitative PCR for UL27

1059

1060   DNA was extracted using a PureLink genomic DNA mini kit (ThermoFisher Scientific). Viral

1061   genome copy number was determined using an established assay based on real-time PCR using

1062   primers and dual-fluorescent probe specific to viral glycoprotein B gene (gB; UL27) . Samples

1063   were assayed alongside a standard curve of HSV-1 strain F nucleocapsid DNA (6), on a ViiA 7

1064   Real-Time PCR System.

Viral entry assay

1065

Monolayers of Vero cells were cooled to 4°C for 30 minutes prior to infection and 100 PFU of

each viral isolate was applied to cell monolayers at 4°C for 1 hour to allow virus binding, after

which unbound virus was washed from the cells. Cells were then moved to 37°C to allow virus

entry. At 0, 10, 20, 30, 45, or 60 minutes, a low-pH citrate buffer was applied to infected cells to

inactivate virus that had not penetrated the cellular membrane. At each time point and for each

virus, parallel infections were performed without the addition of citrate solution. These served as

controls to determine the maximum number of plaques formed. Cell monolayers were washed

and overlaid with methylcellulose. Plaques were scored after 100 hours of incubation. Viral entry

was quantified as the fraction of plaques formed following citrate buffer application, where

100% is the number of plaques formed on a monolayer not treated with citrate buffer (control).

Single-step and multi-step growth curves

1076

Virus diluted in basic growth media containing 2% FBS was applied to near confluent

monolayers of Vero cells and allowed to adsorb for 1 hour. Virus was removed and cells were

incubated at 37C for the duration of infection. At 2 hpi, media containing 0.1% human serum

was added to reduce the contribution of cell-free spread of virus. Single-step growth curves were

performed at MOI=5 as defined by titering viral stocks on U2OS cells, and monolayers harvested

at 2, 6, 12, and 24 hpi. Multi-step growth curves were performed at MOI=0.001 as defined by

titering viral stocks on U2OS cells, and monolayers were harvested at 2, 24, 48, and 72 hpi.

Every 24 hours, the supernatant was removed and media containing 0.1% human serum

reapplied. At the conclusion of each infection, cells were washed two times with PBS and

collected by scraping into an equal volume of media.

Immunocytochemistry

Multi-step growth curves were performed as described above in Vero cell monolayers grown on

glass coverslips. To terminate each infection, coverslips were washed with PBS and fixed in 4%

paraformaldehyde for 15 minutes at room temperature. Cells were permeabilized with 0.5%

Triton-X, blocked in 3% BSA, and incubated with polyclonal rabbit antibodies raised against

total HSV (Agilent Dako, B0114). After washing, cells were incubated with fluorophore-

conjugated anti-Rabbit secondary antibodies (Invitrogen, A-11008) to mark HSV infected cells,

and counterstained with DAPI to mark cell nuclei. Coverslips were initially visualized with a

Leica DM6000 wide field microscope (UPenn Cell and Developmental Microscopy Core). The

5X images were collected on a Photometrics HQ2 high resolution monochrome CCD camera,

and processed with LAS AF software. The 10X images were collected on a EVOS FL Auto

Imaging System (UPenn Cell and Developmental Microscopy Core) and stitched using EVOS

software. Exposure and gain were optimized within each experiment for one virus at the 72-hour

time point and applied identically to each image within that experiment. Any subsequent image

processing (ImageJ) was applied equally to all images in a given experiment.

Immunoblotting

1102

Whole cell lysates were prepared with 1X LDS sample buffer (NuPage) and equal amounts of

lysate were separated by SDS-PAGE. Membranes were immunoblotted with polyclonal rabbit

antibodies raised against total HSV (Agilent Dako, B0114); glycoproteins (g)C, gD, gE, gH, and

VP5 (all gifts from Gary Cohen); ICP8 (gift from David Knipe); and GAPDH (GeneTex,

GTX100118). Proteins were visualized with Pierce ECL Western Blotting Substrate

(ThermoFisher Scientific) and detected using a G:Box imaging system (Syngene).

De novo **genome assembly**

1109

A consensus genome was assembled for each viral isolate using a previously described Viral

Genome Assembly (VirGA) bioinformatics workflow (7). VirGA begins by quality-filtering the

MiSeq sequence reads and removing sequences that match the host (human) genome. Thereafter

VirGA uses a combination of SSAKE *de novo* assemblies run with differing parameters, which

are then combined into a single draft genome using Celera and GapFiller (8–10). After quality-

checking and iterative improvement of the genome assembly, annotations were transferred from

the HSV-2 reference genome (strain HG52; GenBank NC_001798) to each new genome based

on sequence homology (11).

Comparative genomics and phylogenetic analysis

1118

The genomes of all 10 neonatal HSV-2 isolates were combined with all annotated HSV-2

genomes available in GenBank (see **Table S1** for full list) and aligned using MAFFT (12). All

previously published HSV-2 genomes were derived from infected adults. The genome-wide

alignment used a trimmed genome format (lacking the terminal repeats) to avoid giving undue

weight to these duplicated sequences. The MAFFT alignment was used to generate a

NeighborNet phylogenetic network in SplitsTree with Uncorrected P distances (13–15). A

diverse subset of ten adult HSV-2 isolates was selected for protein-level comparisons with the

ten neonatal isolates; these are indicated in **Table S1** with an asterisk. GenBank genomes that

lacked open reading frame (ORF) and protein annotations were excluded from all comparisons

(16, 17). ClustalW2 was used to construct pairwise nucleotide alignments between whole

genomes and pairwise amino acid alignments for each gene and protein (18). Pan-HSV2

comparisons excluded three viral proteins for which sequences are not fully determined in most

published strains, likely due to the high G+C-content and numerous tandem repeats in these

regions: ICP34.5 (RL1 – annotated/complete in only 9 genomes from **Table S1**), ICP0 (RL2 –

annotated/complete in only 8 genomes from **Table S1**), and ICP4 (RS1 – annotated/complete in

only 6 genomes from **Table S1**) (16, 19, 20). Alignments for RL1, RL2, and RS1 for neonatal

isolates were hand-curated to optimize alignment and translation around sequencing gaps, to

avoid over-representing the degree of divergence in these three genes.

1137

1138   **Supporting Figures & Tables**

1139

1140   **Figure S1. U2OS cells support large plaque formation by all isolates.**

1141   Representative plaques are shown after neonatal viruses were allowed to incubate for 100 hours

1142   on Vero **(A)** or U2OS cells **(B)**. All neonatal isolates were capable of forming large plaques on

1143   U2OS cells, which lack innate sensing of viral infection through the STING pathway (21).

1144

1145

1146

Figure S2. Cell-to-cell spread is enhanced in certain neonatal HSV-2 isolates.

Confluent Vero cell monolayers were infected at MOI=0.001 for the time points indicated, in the

presence of 0.1% human serum. HSV positive cells (green) were evaluated at each time point by

immunofluorescence. Cell nuclei are counterstained with DAPI (blue). Serial 10X images were

obtained on an EVOS FL Auto Imaging System and stitched together to create an image of the

entire experimental coverslip. Scale bar = 1mm. These images were quantified in **Figure 3E**.

1153

1154

Figure S3. Phylogenetic clustering demonstrates that neonatal HSV-2 genomes are

genetically distinct from one another and intermingle with the previously known

range of HSV-2 genetic diversity.

A Neighbor-Joining (NJ) tree network constructed among ten neonatal and 58 adult HSV-2

genomes, revealed the wide genetic distribution of the neonatal isolates. The NJ tree (Jukes-

Cantor, 1000 bootstraps) was created in MEGA from a MAFFT trimmed genome alignment.

Bootstrap values ≥ 70 are shown here. See **Figure 4** for a network graph comparison to this tree.

Table S1 contains a complete list of accessions, geographic origins, and references for all adult

HSV2 strains.

1164

1165

1166

Figure S4. Ratio of non-synonymous to synonymous coding variations in neonatal HSV-2 versus adult HSV-2 strains.

The ratio of non-synonymous (dN) to synonymous (dS) coding variations were plotted for each HSV-2 protein. The x-axis value represents the average dN/dS ratio for each protein in 58 adult HSV-2 strains, while the y-axis value represents the average dN/dS ratio in 10 neonatal isolates. Proteins with a difference in average dN/dS ratios ≥1 in neonatal vs. adult HSV-2 genomes are labeled (green indicates a higher average dN/dS in neonatal HSV-2 genomes; red indicates a higher dN/dS in adult HSV-2 genomes). The average dN/dS ratios for all proteins are listed in **Table S2**.

1176

Figure S5. Genome-wide distribution and frequency of minority variants in

neonatal HSV-2 isolates.

For each neonatal HSV-2 isolate, the graph on the left plots spatial location in the genome (x-axis) against the frequency at which each minority variant was observed. The plot on the right summarizes the number of minor variants (y-axis height) in binned increments of 1% (x-axis). The data reveal the distinctly different distribution of minority variants in DISS29, and to a lesser extent CNS15, than in other isolates. Color code matches that used in **Figure 6**. See **Table S3** for full list of SNP and INDEL MV position and frequency data.

1185   **Table S1. Adult HSV-2 genomes (58 total) used for comparative genomic analyses.**

Virus Isolate	Country (with location, if available)	GenBank Accession #	References
H1227*	Finland	KY922721	In process
H1229*	Finland	KY922722	In process
H12212*	Finland	KY922726	In process
Uganda.2*	Uganda	KX574906	(20)
Peru.8*	Peru	KX574874	(20)
Zambia.4*	Zambia	KX574883	(20)
HG52*	Scotland	NC_001798	(22)
CtSF-R*	USA	KP334093	(19)
1192*	Wisconsin, USA	KP334095	(19)
SD90e*	Carletonville, South Africa	KF781518	(23, 24)
Peru.1	Peru	KX574860	(20, 25)
Peru.12	Peru	KX574861	(20, 25)
Zambia.3	Zambia	KX574862	(20, 25)
Zimbabwe.2	Zimbabwe	KX574863	(20, 25)
Zambia.2	Zambia	KX574864	(20, 25)
Peru.9	Peru	KX574865	(20, 25)
Peru.11	Peru	KX574866	(20, 25)
Peru.10	Peru	KX574867	(20, 25)
USA.8a	USA	KX574868	(20, 25)
USA.8b	USA	KX574869	(20, 25)
USA.1b	USA	KX574870	(20, 25)
Zimbabwe.1	Zimbabwe	KX574871	(20, 25)
Peru.7	Peru	KX574872	(20, 25)
Peru.2	Peru	KX574873	(20, 25)
Peru.6	Peru	KX574875	(20, 25)
Peru.4	Peru	KX574876	(20, 25)
Peru.3	Peru	KX574877	(20, 25)
Peru.5	Peru	KX574878	(20, 25)
South Africa.2	South Africa	KX574879	(20, 25)
South Africa.3	South Africa	KX574880	(20, 25)
South Africa.4	South Africa	KX574881	(20, 25)
USA.3a	USA	KX574882	(20, 25)
USA.3b	USA	KX574884	(20, 25)
USA.2b	USA	KX574885	(20, 25)
USA.10	USA	KX574886	(20, 25)
USA.4a	USA	KX574887	(20, 25)
USA.7a	USA	KX574888	(20, 25)
USA.6a	USA	KX574889	(20, 25)

Virus Isolate	Country (with location, if available)	GenBank Accession #	References
USA.6b	USA	KX574890	(20, 25)
USA.5a	USA	KX574891	(20, 25)
South Africa.1	South Africa	KX574892	(20, 25)
Kenya.2	Kenya	KX574893	(20, 25)
Kenya.1	Kenya	KX574894	(20, 25)
USA.7b	USA	KX574895	(20, 25)
USA.5b	USA	KX574896	(20, 25)
USA.11	USA	KX574897	(20, 25)
USA.4b	USA	KX574898	(20, 25)
Uganda.1	Uganda	KX574899	(20, 25)
USA.9	USA	KX574901	(20, 25)
Tanzania.1	Tanzania	KX574902	(20, 25)
USA.1a	USA	KX574903	(20, 25)
USA.2a	USA	KX574904	(20, 25)
Zambia.1	Zambia	KX574905	(20, 25)
2011-21400	USA	KX574908	(20, 25)
H12211	Finland	KY922725	In process
H1226	Finland	KY922720	In process
H1421	Finland	KY922723	In process
H1526	Finland	KY922724	In process

1186   *Indicates 10 adult HSV2 strains used for AA comparisons in Figure 5.
1187

Table S2. Number of nucleotide and amino acid (AA) differences observed in each set of neonatal or adult HSV2 genomes.

Gene*	ORF length in NT^	protein length (#AA)	10 Neonatal # NT diff.^	10 Neonatal # AA changes	10 Neonatal %AA diff.	10 Neonatal dN/dS ratio	10 Adult # NT diff.	10 Adult # AA changes	10 Adult %AA diff.	10 Adult dN/dS ratio	58 Adult # NT diff.	58 Adult # AA diff.	58 Adult %AA diff.	58 Adult dN/dS ratio
UL1	675	225	5	2	0.9	0.7	3	2	0.9	2.0	10	6	2.7	1.5
UL2	1,005	335	7	4	1.2	1.3	10	3	0.9	0.4	14	5	1.5	0.6
UL3	669	223	1	0	0.0	0.0	2	0	0.0	0.0	8	3	1.3	0.6
UL4	606	202	5	2	1.0	0.7	6	2	1.0	0.5	10	2	1.0	0.3
UL5	2,646	882	18	4	0.5	0.3	21	3	0.3	0.2	43	11	1.2	0.3
UL6	2,037	679	9	3	0.4	0.5	6	3	0.4	1.0	26	11	1.6	0.7
UL7	891	297	4	1	0.3	0.3	5	2	0.7	0.7	15	6	2.0	0.7
UL8	2,259	753	13	5	0.7	0.6	18	7	0.9	0.6	39	18	2.4	0.9
UL9	2,616	872	9	3	0.3	0.5	14	4	0.5	0.4	35	11	1.3	0.5
UL10	1,404	468	5	1	0.2	0.3	6	1	0.2	0.2	19	5	1.1	0.4
UL11	291	97	1	0	0.0	0.0	4	2	2.1	1.0	6	3	3.1	1.0
UL12	1,863	621	6	4	0.6	2.0	14	9	1.4	1.8	42	24	3.9	1.3
UL13	1,557	519	8	2	0.4	0.3	6	2	0.4	0.5	20	5	1.0	0.3
UL14	660	220	3	0	0.0	0.0	2	1	0.5	1.0	3	2	0.9	2.0
UL15	2,205	735	2	1	0.1	1.0	4	1	0.1	0.3	20	4	0.5	0.3
UL16	1,119	373	4	2	0.5	1.0	5	3	0.8	1.5	8	4	1.1	1.0
UL17	2,109	703	13	8	1.1	1.6	9	2	0.3	0.3	24	12	1.7	1.0
UL18	957	319	4	1	0.3	0.3	4	0	0.0	0.0	12	1	0.3	0.1
UL19	4,125	1,375	14	3	0.2	0.3	21	6	0.4	0.4	41	12	0.9	0.4
UL20	669	223	4	3	1.3	3.0	6	5	2.2	5.0	10	7	3.1	2.3
UL21	1,599	533	4	3	0.6	3.0	3	2	0.4	2.0	11	7	1.3	1.8
UL22	2,517	839	22	7	0.8	0.5	20	6	0.7	0.4	46	18	2.1	0.6
UL23	1,131	377	7	5	1.3	2.5	11	6	1.6	1.2	20	12	3.2	1.5
UL24	846	282	12	3	1.1	0.3	14	4	1.4	0.4	26	11	3.9	0.7
UL25	1,758	586	7	3	0.5	0.8	8	1	0.2	0.1	25	4	0.7	0.2
UL26	1,914	638	11	3	0.5	0.4	18	7	1.1	0.6	36	16	2.5	0.8
UL26.5	990	330	8	2	0.6	0.3	11	6	1.8	1.2	25	14	4.2	1.3

Gene*	ORF length in NT^	protein length (#AA)	10 Neonatal # NT diff.^	10 Neonatal # AA changes	10 Neonatal %AA diff.	10 Neonatal dN/dS ratio	10 Adult # NT diff.	10 Adult # AA changes	10 Adult %AA diff.	10 Adult dN/dS ratio	58 Adult # NT diff.	58 Adult # AA diff.	58 Adult %AA diff.	58 Adult dN/dS ratio
UL27	2,715	905	16	10	1.1	1.7	17	8	0.9	0.9	39	17	1.9	0.8
UL28	2,358	786	14	7	0.9	1.0	11	5	0.6	0.8	39	15	1.9	0.6
UL29	3,591	1,197	27	8	0.7	0.4	32	11	0.9	0.5	83	24	2.0	0.4
UL30	3,723	1,241	19	12	1.0	1.7	39	20	1.6	1.1	68	35	2.8	1.1
UL31	918	306	6	1	0.3	0.2	9	2	0.7	0.3	18	7	2.3	0.6
UL32	1,797	599	13	4	0.7	0.4	13	7	1.2	1.2	20	8	1.3	0.7
UL33	393	131	2	0	0.0	0.0	1	0	0.0	0.0	5	1	0.8	0.3
UL34	831	277	8	4	1.4	1.0	7	2	0.7	0.4	15	5	1.8	0.5
UL35	339	113	0	0	0.0	0.0	2	1	0.9	0.0	6	4	3.5	2.0
UL36	9,369	3,123	65	31	1.0	0.9	73	33	1.1	0.8	181	82	2.6	0.8
UL37	3,345	1,115	13	8	0.7	1.6	27	16	1.4	1.5	54	25	2.2	0.9
UL38	1,401	467	10	8	1.7	4.0	22	9	1.9	0.7	38	15	3.2	0.7
UL39	3,429	1,143	64	22	1.9	0.5	72	25	2.2	0.5	118	49	4.3	0.7
UL40	1,014	338	6	2	0.6	0.5	4	0	0.0	0.0	13	3	0.9	0.3
UL41	1,479	493	6	1	0.2	0.2	13	2	0.4	0.2	26	8	1.6	0.4
UL42	1,413	471	8	4	0.8	1.0	16	8	1.7	1.0	21	12	2.5	1.3
UL43	1,245	415	10	6	1.4	1.5	8	6	1.4	3.0	27	17	4.1	1.7
UL44	1,443	481	13	6	1.2	0.9	12	7	1.5	1.4	23	15	3.1	1.9
UL45	519	173	1	1	0.6	0.0	3	1	0.6	0.0	6	3	1.7	1.0
UL46	2,169	723	20	9	1.2	0.8	18	7	1.0	0.6	40	17	2.4	0.7
UL47	2,091	697	14	6	0.9	0.8	13	9	1.3	2.3	43	20	2.9	0.9
UL48	1,473	491	6	3	0.6	1.0	7	2	0.4	0.4	18	8	1.6	0.8
UL49	903	301	19	7	2.3	0.6	17	7	2.3	0.7	30	12	4.0	0.7
UL49A	264	88	3	1	1.1	0.5	6	2	2.3	0.5	8	2	2.3	0.3
UL50	1,110	370	8	4	1.1	1.0	7	4	1.1	1.3	22	10	2.7	0.8
UL51	735	245	6	1	0.4	0.2	1	0	0.0	0.0	11	6	2.4	1.2
UL52	3,201	1,067	22	12	1.1	1.2	31	11	1.0	0.6	54	25	2.3	0.9
UL53	1,017	339	14	8	2.4	1.3	16	10	2.9	1.7	26	14	4.1	1.2
UL54	1,539	513	5	0	0.0	0.0	7	3	0.6	0.8	24	12	2.3	1.0
UL55	561	187	3	1	0.5	0.5	2	2	1.1	0.0	9	3	1.6	0.5
UL56	708	236	6	4	1.7	2.0	8	5	2.1	1.7	19	11	4.7	1.4

Gene*	ORF length in NT^	protein length (#AA)	10 Neonatal # NT diff.^	10 Neonatal # AA changes	10 Neonatal %AA diff.	10 Neonatal dN/dS ratio	10 Adult # NT diff.	10 Adult # AA changes	10 Adult %AA diff.	10 Adult dN/dS ratio	58 Adult # NT diff.	58 Adult # AA diff.	58 Adult %AA diff.	58 Adult dN/dS ratio
US1	1,245	415	16	5	1.2	0.5	14	6	1.4	0.8	38	22	5.3	1.4
US2	876	292	7	4	1.4	1.3	4	3	1.0	3.0	15	7	2.4	0.9
US3	1,446	482	9	4	0.8	0.8	7	4	0.8	1.3	17	4	0.8	0.3
US4	2,100	700	22	13	1.9	1.4	40	25	3.6	1.7	70	45	6.4	1.8
US5	279	93	4	1	1.1	0.3	8	4	4.3	1.0	11	6	6.5	1.2
US6	1,182	394	7	2	0.5	0.4	8	2	0.5	0.3	18	5	1.3	0.4
US7	1,119	373	10	6	1.6	1.5	11	5	1.3	0.8	27	14	3.8	1.1
US8	1,638	546	12	9	1.6	3.0	9	9	1.6	0.0	28	20	3.7	2.5
US8A	441	147	4	3	2.0	3.0	2	0	0.0	0.0	7	2	1.4	0.4
US9	270	90	2	0	0.0	0.0	3	1	1.1	0.5	7	4	4.4	1.3
US10	261	87	7	1	1.1	0.2	3	2	2.3	2.0	6	2	2.3	0.5
US11	456	152	2	2	1.3	0.0	0	0	0.0	0.0	136	5	3.3	0.0
US12	909	303	6	4	1.3	2.0	6	0	0.0	0.0	11	10	3.3	10.0
RL1	786	262	17	10	3.8	1.4	♦				♦			
RL2	2,502	834	40	20	2.4	1.0	♦				♦			
RS1	3957	1319	42	26	2.0	1.6	♦				♦			

* See **Table S1** for list of strain names and genome accessions for comparisons of 10 neonatal, 10 adult, and 58 adult HSV2 genomes.

♦ Less than 10 strains from Table S1 have complete sequences of these genes; see Methods for details.

^ Abbreviations: ORF, open reading frame; NT, nucleotide; AA, amino acid; diff., difference; %, percent; dN/dS, ratio of nonsynonymous to synonymous nucleotide changes

Table S4. Viral genetic variations potentially associated with clinical groups.

HSV-2 Gene: Variation	Clinical Group (Isolates w/ Variation)	Predicted effect^	Protein Function	Neurovirulence
gK (UL53): V323M	All CSF Isolates (CNS11, DISS14)	Loss of one TM domain	Glycoprotein at virion and cell surface; interacts with gB and UL20 (26); important for cytoplasmic envelopment, viral egress, and virus-induced cell fusion	gK null virus exhibits poor neuronal spread in culture including decreased retrograde and anterograde transport (27); reduced spread within eye and to nervous system, as well as increased survival, following ocular challenge in mice (28)
gI (US7): R159L + P215S	All CSF Isolates (Both: CNS11,DISS14 R159L only: CNS03, SEM13,CNS15*,DISS29* P215S only: CNS15*,DISS29)	R159L: adds one TM domain; P215S: no change	Glycoprotein at virion and cell surface; forms heterodimer with gE to form viral Fc receptor (29); required for cell-to-cell spread of virus in epithelial and neuronal cells (30, 31);	gI null virus exhibits poor neuronal spread in culture; reduced spread in retina and to retinorecipient areas of the brain following ocular challenge in mice (31)
UL8: R221S	All CSF Isolates (CNS11, DISS14)	No predicted change	Putative primase subunit of helicase primase complex; required for unwinding of viral DNA	None identified
US2: F137L	All CSF Isolates (CNS11, DISS14)	No predicted change	Tegument protein; binds and phosphorylates TAK1 to positively regulate NFkB signaling (32)	Virus containing mutated US2 forms small plaques in culture (33); Mutant virus containing a deletion of both US2 and PK result in decreased death following intracerebral injection in mice (34)
gG (US4): R338L,S442P, E574D	All CSF Isolates (CNS11, DISS14)	No predicted change	Binds and enhances function of chemokines (35); enhances neurotrophin-dependent axonal growth in culture (36)	Virus containing mutated gG results in decreased death following intracerebral injection in mice (33)
UL24: V93A	No CNS Isolates; only found in SEM isolates (SEM02, SEM13, SEM18)	No predicted change	Membrane-associated nuclear protein; associated with dispersal of nucleolin during infection (37)	UL24 null virus forms small plaques in culture (38); decreased viral load in eye and nervous system following ocular challenge in mice **(39–41)**
UL20: P129L	All DISS Isolates w/ CNS Involved (DISS14, DISS29)	Change in position of three TM domains	Membrane-associated protein interacts with gK and gB (26); required for gK glycosylation, cell surface expression (42), and virus-induced cell fusion (43); important for viral egress (44)	UL20 null virus forms small plaques in culture (43); RNA inhibition of UL20 results in decreased rates of encephalitis following HSV footpad infection in mice (45)

* present as minority variant   ^ Prediction by SMART Analysis

References

1. Kimberlin DW, Lin CY, Jacobs RF, Powell DA, Frenkel LM, Gruber WC, Rathore M, Bradley JS, Diaz PS, Kumar M, Arvin AM, Gutierrez K, Shelton M, Weiner LB, Sleasman JW, de Sierra TM, Soong SJ, Kiell J, Lakeman FD, Whitley RJ, National Institute of Allergy and Infectious Diseases Collaborative Antiviral Study Group. 2001. Natural history of neonatal herpes simplex virus infections in the acyclovir era. Pediatrics 108:223–229.

2. Kimberlin DW, Whitley RJ, Wan W, Powell DA, Storch G, Ahmed A, Palmer A, Sánchez PJ, Jacobs RF, Bradley JS. 2011. Oral acyclovir suppression and neurodevelopment after neonatal herpes. N Engl J Med 365:1284–1292.

3. Kimberlin DW, Lin CY, Jacobs RF, Powell DA, Corey L, Gruber WC, Rathore M, Bradley JS, Diaz PS, Kumar M, Arvin AM, Gutierrez K, Shelton M, Weiner LB, Sleasman JW, de Sierra TM, Weller S, Soong SJ, Kiell J, Lakeman FD, Whitley RJ, National Institute of Allergy and Infectious Diseases Collaborative Antiviral Study Group. 2001. Safety and efficacy of high-dose intravenous acyclovir in the management of neonatal herpes simplex virus infections. Pediatrics 108:230–238.

4. Whitley R, Arvin A, Prober C, Burchett S, Corey L, Powell D, Plotkin S, Starr S, Alford C, Connor J, Jacobs R, Nahmias A, Soong S-J, the National Institute of Allergy, Infectious Diseases Collaborative Antiviral Study Group*. 1991. A Controlled Trial Comparing Vidarabine with Acyclovir in Neonatal Herpes Simplex Virus Infection. N Engl J Med 324:444–449.

5. Whitley R, Arvin A, Prober C, Corey L, Burchett S, Plotkin S, Starr S, Jacobs R, Powell D, Nahmias A, Sumaya C, Edwards K, Alford C, Caddell G, Soong S-J, the National Institute of Allergy and Infectious Diseases Collaborative Antiviral Study Group*. 1991. Predictors of Morbidity and Mortality in Neonates with Herpes Simplex Virus Infections. N Engl J Med 324:450–454.

6. Szpara ML, Parsons L, Enquist LW. 2010. Sequence variability in clinical and laboratory isolates of herpes simplex virus 1 reveals new mutations. J Virol 84:5303–13.

7. Parsons LR, Tafuri YR, Shreve JT, Bowen CD, Shipley MM, Enquist LW, Szpara ML. 2015. Rapid genome assembly and comparison decode intrastrain variation in human alphaherpesviruses. mBio 6:e02213-14.

8. Myers EW, Sutton G, Delcher AL, Dew IM, Fasulo DP, Flanigan MJ, Kravitz SA, Mobarry CM, Reinert KHJ, Remington KA, Anson EL, Bolanos RA, Chou H-H, Jordan CM, Halpern AL, Lonardi S, Beasley EM, Brandon RC, Chen L, Dunn PJ, Lai Z, Liang Y, Nusskern DR, Zhan M, Zhang Q, Zheng X, Rubin GM, Adams MD, Venter CJ. 2000. A Whole-Genome Assembly of Drosophila. Science 287:2196–2204.

9. Boetzer M, Pirovano W. 2012. Toward almost closed genomes with GapFiller. Genome Biol 13:1–9.

10. Warren RL, Sutton GG, Jones SJM, Holt R a. 2007. Assembling millions of short DNA

sequences using SSAKE. Bioinformatics 23:500–1.

11. Dolan A, Jamieson FE, Cunningham C, Barnett BC, McGeoch DJ. 1998. The genome sequence of herpes simplex virus type 2. J Virol 72:2010–2021.

12. Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–3066.

13. Strimmer K, Wiuf C, Moulton V. 2001. Recombination analysis using directed graphical models. Mol Biol Evol 18:97–99.

14. Norberg P, Kasubi MJ, Haarr L, Bergstrom T, Liljeqvist J-A. 2007. Divergence and recombination of clinical herpes simplex virus type 2 isolates. J Virol 81:13158–13167.

15. Norberg P, Tyler S, Severini A, Whitley R, Liljeqvist J-A, Bergstrom T. 2011. A genome-wide comparative evolutionary analysis of herpes simplex virus type 1 and varicella zoster virus. PloS One 6:1–8.

16. Newman RM, Lamers SL, Weiner B, Ray SC, Colgrove RC, Diaz F, Jing L, Wang K, Saif S, Young S, Henn M, Laeyendecker O, Tobian AAR, Cohen JI, Koelle DM, Quinn TC, Knipe DM. 2015. Genome Sequencing and Analysis of Geographically Diverse Clinical Isolates of Herpes Simplex Virus 2. J Virol JVI.01303-15.

17. Johnston C, Magaret A, Roychoudhury P, Greninger AL, Reeves D, Schiffer J, Jerome KR, Sather C, Diem K, Lingappa JR, Celum C, Koelle DM, Wald A. 2017. Dual-strain genital herpes simplex virus type 2 (HSV-2) infection in the US, Peru, and 8 countries in sub-Saharan Africa: A nested cross-sectional viral genotyping study. PLOS Med 14:e1002475.

18. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. 2007. Clustal W and Clustal X version 2.0. Bioinforma Appl Note 23:2947–2948.

19. Kolb AW, Larsen IV, Cuellar JA, Brandt CR. 2015. Genomic, Phylogenetic, and Recombinational Characterization of Herpes Simplex Virus 2 Strains. J Virol 89:6427–6434.

20. Johnston C, Magaret A, Roychoudhury P, Greninger AL, Cheng A, Diem K, Fitzgibbon MP, Huang M, Selke S, Lingappa JR, Celum C, Jerome KR, Wald A, Koelle DM. 2017. Highly conserved intragenic HSV-2 sequences: Results from next-generation sequencing of HSV-2 U L and U S regions from genital swabs collected from 3 continents. Virology 510:90–98.

21. Deschamps T, Kalamvoki M. 2017. Impaired STING Pathway in Human Osteosarcoma U2OS Cells Contributes to the Growth of ICP0-Null Mutant Herpes Simplex Virus. J Virol 91:e00006-17.

22. Davison AJ. 2012. Evolution of sexually transmitted and sexually transmissible human herpesviruses. Ann N Y Acad Sci 1230:E37–E49.

23. Colgrove R, Diaz F, Newman R, Saif S, Shea T, Young S, Henn M, Knipe DM. 2014. Genomic sequences of a low passage herpes simplex virus 2 clinical isolate and its plaque-purified derivative strain. Virology 450–451:140–145.

24. Lai W, Chen CY, Morse SA, Htun Y, Fehler HG, Liu H, Ballard RC. 2003. Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa. Sex Transm Infect 79:202–207.

25. Koelle DM, Norberg P, Fitzgibbon MP, Russell RM, Greninger AL, Huang M-L, Stensland L, Jing L, Magaret AS, Diem K, Selke S, Xie H, Celum C, Lingappa JR, Jerome KR, Wald A, Johnston C. 2017. Worldwide circulation of HSV-2 × HSV-1 recombinant strains. Sci Rep 7:44084.

26. Foster TP, Chouljenko VN, Kousoulas KG. 2008. Functional and physical interactions of the herpes simplex virus type 1 UL20 membrane protein with glycoprotein K. J Virol 82:6310–6323.

27. David AT, Saied A, Charles A, Subramanian R, Chouljenko VN, Kousoulas KG. 2012. A Herpes Simplex Virus 1 (McKrae) Mutant Lacking the Glycoprotein K Gene Is Unable To Infect via Neuronal Axons and Egress from Neuronal Cell Bodies. mBio 3.

28. David AT, Baghian A, Foster TP, Chouljenko VN, Kousoulas KG. 2008. The Herpes Simplex Virus Type 1 (HSV-1) Glycoprotein K(gK) is Essential for Viral Corneal Spread and Neuroinvasiveness. Curr Eye Res 33:455–467.

29. Johnson DC, Frame MC, Ligas MW, Cross AM, Stow ND. 1988. Herpes simplex virus immunoglobulin G Fc receptor activity depends on a complex of two viral glycoproteins, gE and gI. J Virol 62:1347–1354.

30. Dingwell KS, Brunetti CR, Hendricks RL, Tang Q, Tang M, Rainbow AJ, Johnson DC. 1994. Herpes simplex virus glycoproteins E and I facilitate cell-to-cell spread in vivo and across junctions of cultured cells. J Virol 68:834–845.

31. Dingwell KS, Doering LC, Johnson DC. 1995. Glycoproteins E and I facilitate neuron-to-neuron spread of herpes simplex virus. J Virol 69:7087–7098.

32. Lu X, Huang C, Zhang Y, Lin Y, Wang X, Li Q, Liu S, Tang J, Zhou L. 2017. The Us2 Gene Product of Herpes Simplex Virus 2 modulates NF-κB activation by targeting TAK1. Sci Rep 7:8396.

33. Weber PC, Levine M, Glorioso JC. 1987. Rapid identification of nonessential genes of herpes simplex virus type 1 by Tn5 mutagenesis. Science 236:576–579.

34. Meignier B, Longnecker R, Mavromara-Nazos P, Sears AE, Roizman B. 1988. Virulence of and establishment of latency by genetically engineered deletion mutants of herpes simplex virus 1. Virology 162:251–254.

35. Viejo-Borbolla A, Martinez-Martín N, Nel HJ, Rueda P, Martín R, Blanco S, Arenzana-Seisdedos F, Thelen M, Fallon PG, Alcamí A. 2012. Enhancement of chemokine function as an immunomodulatory strategy employed by human herpesviruses. PLoS Pathog 8:e1002497.

36. Cabrera JR, Viejo-Borbolla A, Martinez-Martín N, Blanco S, Wandosell F, Alcamí A. 2015. Secreted herpes simplex virus-2 glycoprotein G modifies NGF-TrkA signaling to attract free nerve endings to the site of infection. PLoS Pathog 11:e1004571.

37. Lymberopoulos MH, Pearson A. 2007. Involvement of UL24 in herpes-simplex-virus-1-induced dispersal of nucleolin. Virology 363:397–409.

38. Jacobson JG, Martin SL, Coen DM. 1989. A conserved open reading frame that overlaps the herpes simplex virus thymidine kinase gene is important for viral growth in cell culture. J Virol 63:1839–1843.

39. Rochette P-A, Bourget A, Sanabria-Solano C, Lahmidi S, Lavallée GO, Pearson A. 2015. Mutation of UL24 impedes the dissemination of acute herpes simplex virus 1 infection from the cornea to neurons of trigeminal ganglia. J Gen Virol 96:2794–2805.

40. Leiva-Torres GA, Rochette P-A, Pearson A. 2010. Differential importance of highly conserved residues in UL24 for herpes simplex virus 1 replication in vivo and reactivation. J Gen Virol 91:1109–1116.

41. Jacobson JG, Chen SH, Cook WJ, Kramer MF, Coen DM. 1998. Importance of the herpes simplex virus UL24 gene for productive ganglionic infection in mice. Virology 242:161–169.

42. Foster TP, Alvarez X, Kousoulas KG. 2003. Plasma membrane topology of syncytial domains of herpes simplex virus type 1 glycoprotein K (gK): the UL20 protein enables cell surface localization of gK but not gK-mediated cell-to-cell fusion. J Virol 77:499–510.

43. Foster TP, Melancon JM, Baines JD, Kousoulas KG. 2004. The herpes simplex virus type 1 UL20 protein modulates membrane fusion events during cytoplasmic virion morphogenesis and virus-induced cell fusion. J Virol 78:5347–5357.

44. Foster TP, Melancon JM, Olivier TL, Kousoulas KG. 2004. Herpes simplex virus type 1 glycoprotein K and the UL20 protein are interdependent for intracellular trafficking and trans-Golgi network localization. J Virol 78:13262–13277.

45. Liu J, Lewin AS, Tuli SS, Ghivizzani SC, Schultz GS, Bloom DC. 2008. Reduction in severity of a herpes simplex virus type 1 murine infection by treatment with a ribozyme targeting the UL20 gene RNA. J Virol 82:7467–7474.