Systematic Discovery of Conservation States for Single-Nucleotide Annotation of

the Human Genome

Adriana Sperlea[1,2] and Jason Ernst[1,2,3,4,5,6,*]

[1] Bioinformatics Interdepartmental Program, University of California, Los Angeles, Los Angeles, California,

90095, USA.

[2] Department of Biological Chemistry, University of California, Los Angeles, Los Angeles, California, USA.

[3] Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research at University of

California, Los Angeles, Los Angeles, California, 90095, USA.

[4] Computer Science Department, University of California, Los Angeles, Los Angeles, California, 90095,

USA.

[5] Jonsson Comprehensive Cancer Center, University of California, Los Angeles, Los Angeles, California,

90095, USA.

[6] Molecular Biology Institute, University of California, Los Angeles, Los Angeles, California, 90095, USA.

*Correspondence: Jason Ernst (jason.ernst@ucla.edu)

Abstract

Comparative genomics sequence data is an important source of information for interpreting

genomes. Genome-wide annotations based on this data have largely focused on univariate

scores or binary calls of evolutionary constraint. Here we present a complementary whole

genome annotation approach, ConsHMM, which applies a multivariate hidden Markov model to

learn *de novo* different 'conservation states' based on the combinatorial and spatial patterns of

which species align to and match a reference genome in a multiple species DNA sequence

alignment. We applied ConsHMM to a 100-way vertebrate sequence alignment to annotate the

human genome at single nucleotide resolution into 100 different conservation states. These

states have distinct enrichments for other genomic information including gene annotations,

chromatin states, and repeat families, which were used to characterize their biological

significance. Conservation states have greater or complementary predictive information than

standard constraint based measures for a variety of genome annotations. Bases in constrained

elements have distinct heritability enrichments depending on the conservation state assignment,

demonstrating their relevance to analyzing phenotypic associated variation. The conservation

states also highlight differences in the conservation patterns of bases prioritized by a number of

scores used for variant prioritization. The ConsHMM method and conservation state annotations

provide a valuable resource for interpreting genomes and genetic variation.

Introduction

The large majority of phenotype-associated variants implicated by genome-wide

association studies (GWAS) fall outside of protein coding regions.[1] Identifying the causal

variants and interpreting their biological role in these less well understood non-coding regions is

a significant challenge.[2] Large-scale mapping of epigenomic data across different cell and

tissue types has been one approach for annotating and interpreting the non-coding regions of

genomes.[3–5] Using comparative genomics data to identify regions of evolutionary constraint has

been a complementary approach for these purposes.[6–9]

In addition to providing evolutionary information, comparative genomics data has the

advantage of providing information at single-nucleotide resolution. Furthermore, it is cell type

agnostic and thus informative even when the relevant cell or tissue type has not been

experimentally profiled.[10,11] The most commonly used representations of this information are

univariate scores and binary elements of evolutionary constraint, which are called based on a

multiple species DNA sequence alignment and assumed models of evolution and selection.[8,9,12–

14] Supporting the importance of these annotations, heritability analyses have recently implicated

evolutionary constrained elements as one of the annotations most enriched for phenotype

associated variants.[15] These scores and elements have also been highly informative features to

integrative methods for prioritizing pathogenic variants.[16–19] Further improvements for predicting

pathogenic variants in coding regions have been made to the integrative scores by incorporating

features defined directly from a multiple sequence alignment.[20]

While highly useful, the representation of comparative genomics information into

univariate scores or binary elements is limited in the amount of information it can convey about

the underlying multiple sequence alignment at a specific base. This limitation has become more

pronounced given the large number of species sequenced and incorporated into multi-species

alignments such as a 100-way alignment to the human genome.[21] Approaches have been

developed to associate constrained elements, regions, or individual bases with specific

branches in a phylogenetic tree.[22–28] While also useful, such directed approaches are biased to

only representing certain types of patterns present in the alignment. An alternative approach

used for comparative genomic based annotation learned patterns of different classes of

mutations between human and orangutan[29], but this approach was only applicable at a broad

region level and only incorporated information from one non-human genome.

Analogous to the many sequenced genomes available for comparative analysis, many

different datasets are available for annotating the genome based on epigenomic data.

Approaches that define 'chromatin states' based on combinatorial and spatial patterns in these

datasets have effectively summarized the information in them to provide *de novo* genome

annotations.[4,30–32] Inspired by the success of these approaches, here we develop a method,

ConsHMM, that extends ChromHMM[31] to systematically annotate genomes into 'conservation

states' at single nucleotide resolution. The conservation states assignments are based on the

combinatorial and spatial patterns of which species align to and which match a reference

genome at each nucleotide in a multiple species DNA sequence alignment. ConsHMM takes a

relatively unbiased modeling approach that does not explicitly assume a specific phylogenetic

relationship between species. The set of conservation patterns ConsHMM can infer are thus

flexible and determined directly from the DNA sequence alignment.

We applied ConsHMM to assign a conservation state to each nucleotide of the human

genome. These states are able to capture distinct enrichments for other genomic annotations

such as gene annotations, CpG islands, repeat families, chromatin states, and genetic variation.

We demonstrate that the conservation state annotations capture additional information that is

not represented by scores or binary calls of constraint. We also show how the conservation

states enable a deeper understanding of types of bases prioritized by a number of different

scores used for variant prioritization, including those scores that integrate constraint information

with a diverse set of other genomic annotations. Overall, these conservation state annotations

are a resource for interpreting the genome and potential disease-associated variation, which

complement both existing conservation and epigenomic-based annotations.

Material and Methods

Modeling conservation states with ConsHMM

ConsHMM takes as input an *N-way* multi-species sequence alignment to a designated

reference genome. For each base in the reference genome, i, ConsHMM encodes information

from the multiple species alignment into a vector, v_i, of length N-1. An element of the vector, $v_{i,j}$,

corresponds to one of three possible observation for a non-reference species j at position i. The

three possible observations are: (1) the non-reference species aligns with a non-indel nucleotide

symbol present matching the reference nucleotide, (2) the non-reference species aligns with a

non-indel nucleotide symbol present, but does not match the reference nucleotide, or (3) the

non-reference species does not align with a non-indel nucleotide symbol present.

ConsHMM assumes that these observations are generated from a multivariate HMM

where the emission parameters are assumed to be generated by a product of independent

multinomial random variables, corresponding to each species in the alignment. Formally, the

model is defined based on a fixed number of states K, and number of species in the multiple

sequence alignment N. For each state k $(k = 1,...,K)$, non-reference species j $(j = 1,...,N-1)$ and

possible observation m (m = 1, 2, or 3 as described above), there is an emission parameter:

$p_{k,j,m}$ corresponding to the probability in state k for species j of having observation m. For each

possible observation m, let $I_m(v_{i,j}) = 1$ if $v_{i,j} = m$, and 0 otherwise. Let $b_{t,u}$ be a parameter for the

probability of transitioning from state t to state u. Let $c \in C$ denote a chromosome, where C is

the set of all chromosomes in the reference genome of the multiple species alignment, and let

L_c be the number of bases on chromosome c. Let a_k (k = 1,...,K) be a parameter for the

probability of the first base on a chromosome being in state k. Let $s_c \in S_c$ be a hidden state

sequence on chromosome c and S_c be the set of all such possible state sequences. Let c_h

denote position h on chromosome c. Let s_{c_h} denote the hidden state at position c_h for state

sequence s_c.

We learn a setting of the model parameters that aims to optimize

$$P(v|a,b,p) = \prod_{c \in C} \sum_{s_c \in S_c} a_{s_{c_1}} \left(\prod_{i=2}^{L_c} b_{s_{c_{i-1}}, s_{c_i}} \right) \prod_{h=1}^{L_c} \prod_{j=1}^{N-1} \prod_{m=1}^{3} p_{s_{c_h}, j, m}^{I_m(v_{c_h, j})}$$

Once a model is learned, each nucleotide is assigned to the state with maximum

posterior probability. To conduct the model learning and state assignments, ConsHMM calls an

extended version of the ChromHMM[31] software originally designed to solve an analogous

problem of annotating a genome into chromatin states based on combinatorial and spatial

patterns of the presence of different chromatin marks. The modeling in ConsHMM differs from

the typical use of ChromHMM in three main respects: (1) the observation for each feature

comes from a three-way multinomial distribution as opposed to a Bernoulli distribution, (2) it is

applied at single nucleotide resolution as opposed to 200-bp resolution, (3) it is applied with

more features than ChromHMM models have used in the past. (2) and (3) raise scalability

issues in terms of time and memory, which we addressed in an updated version of ChromHMM

(see below).

To apply ChromHMM in the context of three-way multinomial distributions, ConsHMM

represents the three possible observations at position i for a species j with two binary variables,

y_{ij} and z_{ij}, corresponding to aligning and matching the reference genome respectively. y_{ij} has the

value of 1 if the other species aligns to the reference with a non-indel nucleotide and 0

otherwise. z_{ij} has the value of 1 if the other species has the same nucleotide as the reference

sequence and has a value of 0 if the other species has a different nucleotide present than the

reference. In the case in which $y_{ij}=0$, there is no nucleotide to compare to the reference and that

value of the z_{ij} variable is considered missing (encoded with a '2' for ChromHMM). If the value of

an observed variable is missing, ChromHMM excludes the Bernoulli random variable

corresponding to the observation from the emission distribution calculation at that position. For

each state k and species j, ChromHMM thus learns two parameters, $f_{k,j}$ and $g_{k,j}$. $f_{k,j}$ corresponds

to the probability that at a given position in state k, species j aligns to the reference genome with

a non-indel nucleotide that is $P(y_{i,j}=1 \mid s_i=k)$. $g_{k,j}$ corresponds to the probability that at a given

position in state k, species j matches the reference genome conditioned on species j aligning

with a non-indel nucleotide that is $P(z_{i,j} = 1 \mid y_{i,j}=1$ and $s_i=k)$. This representation is equivalent to

the three-way multinomial distribution, $(p_{k,j,1}, p_{k,j,2}, p_{k,j,3})$ described above where $p_{k,j,1} = P(y_{i,j}=1,$

$z_{i,j}=1 \mid s_i = k)$, $p_{k,j,2} = P(y_{i,j}=1, z_{i,j}=0 \mid s_i = k)$, and $p_{k,j,3} = P(y_{ij}=0 \mid s_i = k)$, since $p_{k,j,1} = f_{k,j} \times g_{k,j}$, $p_{k,j,2} =$

$f_{k,j} \times (1-g_{k,j})$, and $p_{k,j,3} = 1 - f_{k,j}$.

Multiple species sequence alignment choice

Our method and software can be applied to any multiple species sequence alignment

which is available in multiple alignment format (MAF) or which can be converted into this format.

For the results presented here we applied it to the 100-way Multiz vertebrate alignment with

human (hg19) as the reference genome.[21,33]

Scaling-up ConsHMM to single base resolution with hundreds of features

Since for our application ConsHMM needs to run ChromHMM at single base resolution ('-b 1' flag) with 198 features after our binary encoding (2 for each non-human species in the 100-way alignment), we had to address scalability issues in terms of both memory and time. To address the memory issue we modified ChromHMM to support only loading in main memory input for chromosomes it is actively processing, as previously ChromHMM would only support loading all data into main memory upfront. This option can now be accessed in ChromHMM through the '-lowmem' flag. To reduce the time required we used 12-parallel processors ('-p 12' flag) and we trained on a different random subset of the human genome on each iteration of the Baum-Welch algorithm. We divided each chromosome into 200kb segments (with the exception of the last segment of each chromosome which was less than this) in order to form random subsets of the human genome. We modified ChromHMM to allow training for each iteration on a randomly selected subset of 150 of these segments ('-n 150' flag), corresponding to 30MB per iteration. We ran this for 200 iterations by adding the '-d -1' flag, which removed one of ChromHMM's default stopping criterion based on computed likelihood change on the sampled data, since the likelihood is now expected to both increase and decrease between iterations as different sequences are sampled. These new options were included in version 1.13 of ChromHMM. The unique code to ConsHMM is written in Python. The code of ConsHMM shared with ChromHMM is written in Java and included with ConsHMM.

Generating genome-wide annotations

After learning a 100-state model, we used it to segment and annotate the genome at base-pair resolution into one of the 100 conservation states. Each base in the human genome is classified into the state with the highest posterior probability. ConsHMM does this by running the MakeSegmentation command of ChromHMM. Due to computational constraints, the

segmentation could not be generated for entire chromosomes at once. Instead, we ran

MakeSegmentation on the same 200kb partitioning made for learning the model. We then

merged the resulting files together using ConsHMM's mergeSegmentation.py command with

slice size parameter set to 200,000 ('-s 200000' flag) and the number of states parameter set to

100 ('-n 100 flag').

Computing enrichments for external annotations

All overlap enrichments for external annotations were computed using the ChromHMM

OverlapEnrichment command. OverlapEnrichment computes enrichments for an external

annotation in each state assuming a uniform background distribution. Specifically the fold

enrichment of a state for an external annotation is

$$\frac{\%\ of\ external\ annotation\ bases\ falling\ in\ that\ state}{\%\ of\ genome\ falling\ in\ that\ state}$$

Positional enrichments of states relative to an anchor point from an external annotation

were computed using the ChromHMM NeighborhoodEnrichment command at single base

resolution ('-b 1' flag), single base spacing from the anchor point ('-s 1') and using the '-l' and '-r'

flags to specify the size of the region of interest around the anchor point. The '-lowmem' flag

was also used for computing the enrichments for OverlapEnrichment and

NeighborhoodEnrichment.

External data sources for enrichment analyses

The external annotations of repeat elements were obtained from the UCSC genome

browser RepeatMasker track.[21,34] We generated an annotation for whether a base overlapped

any repeat element, as well as separate annotations for bases falling in each class and family of

repeat elements. The gene annotations were obtained from GENCODE v19 for hg19[35]. CpG

Island annotations were obtained from the UCSC genome browser. Annotations of SNPs with

>=1% minor allele frequency were obtained from the commonSNP147 track from the UCSC

genome browser, which is based on dbSNP build 147. GWAS catalog variants were obtained

from the NHGRI-EBI Catalog, accessed on Dec 5, 2016.[36] For annotations of DNase I

Hypersensitive Sites (DHS) processed by the Roadmap Epigenomics Consortium, we used

Macs2 narrowPeak calls.[5] The Fetal Brain and HepG2 DHS used were of epigenome samples

E082 and E118 respectively. For the median non-exonic DHS enrichments and ranking of

states in the heritability partitioning analysis we used narrowPeak calls from the ENCODE

consortium.[3] In the cases where ENCODE provided more than one replicate for a cell or tissue

type, we used the first replicate.

PhyloP and PhastCons scores and constrained element calls were obtained from the

UCSC genome browser. Assembly gap annotations were obtained from the Gap track from the

UCSC genome browser. The context-dependent tolerance score (CDTS) used was that based

on a cohort of 7784 unrelated individuals, following the analyses in Iulio et al.[37], which focused

on this version of the score. The CDTS and variants from this cohort were both lifted from hg38

to hg19 using the liftOver tool from the UCSC genome browser.[21]

Choice of number of states

We learned models with each number of states between 2 and 100 states. We set 100

as the maximum number of states we would consider for computational tractability and

maintaining a manageable number of states for analysis. The choice of a maximum of 100 also

corresponds to the number of species used and allows for the possibility of each state to cover

1% of the genome. We analyzed the Bayesian Information Criterion (BIC) for models with each

number of states between 2 and 100, and found that the BIC generally decreases as the

number of states increases in the range considered (**Figure S1**). The BIC was calculated using

the BIC_HMM function from the HMMpa R package.[38] Analyzing the 100-state model's internal

confidence estimate of its state assignments also supported a larger number of states.

Specifically, for each state in the 100-state model we computed the average posterior

probability of that state at each base in the genome assigned to it, and confirmed consistently

high average posterior probability values in the range [0.92,1.00] with a median of 0.97 (**Figure**

S2). The posterior probabilities were computed by running the MakeSegmentation command in

ChromHMM with the '-printposterior' flag. We also investigated if additional states in models

with larger number of states were biologically relevant. Specifically, we computed enrichments

for various external annotations for models with each number of states between 2 and 100 to

determine if biologically relevant enrichments were only robustly observed in models with more

than a certain number of states. In the case of CpG islands, we observed that only models with

at least 87 states consistently obtained >15 fold enrichment and only models with at least 95

states consistently obtained >30 fold enrichment (**Figure S3**). We saw a similar pattern of

increasing enrichments for annotated transcription start sites (TSSs) for models with large

number of states. We therefore decided to analyze the largest model, 100 states, that we were

considering. We note that annotations based on chromatin states used fewer number of states,

but were also defined on fewer features at a coarser resolution and had a less uniform genome

coverage.[4,30,39]

State clustering

We clustered the states based on the correlation of vectors containing the values $f_{k,j}$ and

$f_{k,j} \times g_{k,j}$ for each species j defined above. State clustering was performed using the hclust

hierarchical clustering function from the cba R package.[40] The leaves of the resulting

hierarchical tree were ordered according to the optimal leaf ordering algorithm[41] implemented in

the order.optimal R function from the cba package. We then cut the tree such that the 8 major

groups of states were designated. The full tree is provided in **Figure S4**.

Genome segmentation using uniform transition probabilities

For analyzing the effect of the transition probabilities on the genome segmentation, we

created a separate model, which was the same model we used in the main analyses, except we

set all transition probabilities to 0.01, corresponding to each state having an equal probability of

transitioning to any state including itself. We then created a new genome segmentation by

running the MakeSegmentation command in ChromHMM with this new model. For each state,

we counted how many of the bases assigned to it in the original annotation were also assigned

to it in the annotation created with the uniform transitions and divided this number by the

number of bases in the state in the original annotation. This calculation provided a fraction from

0 to 1. We also reported the number of segments produced by each model, where a segment is

defined to be one or more consecutive bases all assigned to the same state, such that any

immediately adjacent bases are assigned to a different state or states.

Gene Ontology enrichments

For each state and each protein coding gene based on GENCODE we computed the

number of bases in that state that are within +/- 2kb of the gene's TSS. In the case of genes

with multiple annotated TSSs, we used the outermost TSS. We then created a ranking of genes

for every state by sorting the genes in descending order of this number of bases. For each state

we then created a set of 969 genes that represent the top 5% of genes in the state among the

19,397 genes we considered. We performed a Gene Ontology (GO) enrichment analysis

(ontology and annotations files from Nov. 24[th], 2016) for the top 5% genes in each state using

the STEM software in batch mode with default options and the set of all genes considered as

background.[42] STEM computed an uncorrected p-value based on the hypergeometric

distribution for each term displayed in the figures summarizing the analysis. STEM also reported

corrected p-values for testing multiple GO terms for a single state based on randomization to

three significant digits, which was less than 0.001 for all p-values mentioned in the main text.

Transcription factor binding site motif enrichments

We computed the enrichment of the conservation states within 15 bases upstream and

downstream of the center point of the POU5F1 and STAT known transcription factor-binding site

motifs.[43] The enrichment was computed relative to the background regions of the genome that

were used to identify the motifs, which excluded repeat elements, coding sequence, and 3'

untranslated regions (UTRs). The *known1* version of the motifs was used for both motifs.

Clustering of cell-type specific DNase I hypersensitive site enrichments

For the clustering of DHS analysis, we first computed the enrichments of all conservation

states for DHS for 53 samples processed by the Roadmap Epigenomics consortium[5], of which

16 were originally generated by the ENCODE project consortium.[3] We then selected the subset

of states that had a fold enrichment of at least 2 in at least one sample, leading to a subset of 21

conservation states. To more directly focus on each state's relative enrichments across

samples, we log_2 transformed each enrichment value, and then normalized the enrichments for

each state by subtracting the mean enrichment across samples and dividing by the standard

deviation. We then hierarchically clustered the states based on the correlation of their

enrichments across samples and hierarchically clustered the samples based on their

correlations across states using the pheatmap R package.[44] We also computed for each sample

the fold enrichment of DHS bases for bases in CpG islands, as the ratio between the percent of

DHS bases in CpG islands and the percent of the genome falling in CpG islands.

Precision recall analysis for recovery of gene annotations and DHS

We randomly split the 200kb genome segments used for training the model and

segmentation into two halves corresponding to training and testing data. For each target set in

the precision-recall analyses we ordered the ConsHMM states in decreasing order of their

enrichment for the target among the training set bases. We then used that ordering to iteratively

add the testing set bases in each state to form cumulative sets of bases predicted to be of the

target set and computed the precision and recall for them. For each constraint score we

computed the precision-recall curve for predicting the target set in the test data using two

methods. For the first method, we directly ordered bases in descending order of their assigned

score. For the second method, we split the sorted scores into 400 bins such that each bin

contains on average 0.25% of the genome, which was the size of the smallest state of the

ConsHMM model (0.25% of the genome in state 100). Specifically, we assigned all bases in the

genome where the score was not defined to one bin and then divided the remaining bases

uniformly among the 399 other bins based on their score. In some cases score increments were

at the boundary between two bins at their provided floating-point precision, or overlapped

multiple bins. In these cases we uniformly split the target bases assigned to that score

increment into multiple bins proportionally to the overall percentage of the score increment

falling in each bin. We then treated the 400 bins as 400 states and followed the same procedure

described for the ConsHMM states. We also computed the precision and recall of bases in each

constrained element set for predicting the target set on the testing data. For the DHS analyses,

we also separately evaluated recovery of DHS bases when restricting the analysis to non-

exonic regions. Additionally, both genome-wide and within non-exonic regions, we evaluated the

recovery of DHS bases when restricting the analysis to bases distal to a TSS, defined as more

than 2kb from a TSS.

Precision recall analysis for recovery of DHS bases aggregated across cell and tissue types

For the analysis of the recovery of DHS aggregated across cell and tissue types we

concatenated DHS from 53 cell or tissue types processed by the Roadmap Epigenomics

Consortium into one annotation in which each combination of chromosome and cell or tissue

type effectively becomes a new chromosome. We then split the concatenated data into training

and testing sets as described above. We computed the enrichments of the ConsHMM states

and scores split into bins as detailed above, but multiplying the size of each state and bin by the

number of DNase I hypersensitivity data sets. The precision and recall values for the ConsHMM

states, constraint scores considered directly, constraint scores split into bins, and constrained

element sets were then computed on the testing data.

Enrichment analysis for constrained non-exonic elements assigned to phylogenetic branches

We lifted over the constrained non-exonic elements (CNEEs) from *Lowe et al.*[22] from

hg18 coordinates to hg19, using the liftOver tool from the UCSC genome browser with default

settings.[21] These elements were previously partitioned into subsets based on the inferred

branch point of origin in a phylogenetic tree.[22] We computed the enrichments of the

conservation states for all the CNEEs and for each subset of the CNEEs separately, using the

OverlapEnrichment command from ChromHMM at single nucleotide resolution (`-b 1` flag) and

using the low memory option (`-lowmem`). We also computed analogous enrichments for

CNEEs overlapping PhastCons elements called on the same 100-way alignment that the

conservation states were annotated based on. To compute the enrichments of CNEEs for bases

in CpG islands we created an annotation consisting of a state for each CNEE subset and one

additional state for bases not assigned to any CNEE. We then ran the same OverlapEnrichment

command as above to compute enrichments of CNEE bases for non-exonic CpG islands, and

non-exonic bases in general. The reported enrichment of CpG islands is the ratio of these two

enrichments, effectively computing an enrichment relative to the non-exonic background. The

set of non-exonic bases for the enrichment analysis was generated by excluding all bases

annotated as an exon in GENCODE v19.

Heritability partitioning analysis

The heritability partitioning was performed using the LD-score regression ldsc software.

[15] We partitioned the PhastCons constrained elements into two halves based on a ranking of the

conservation states. We focused on the PhastCons constrained elements for this analysis, since

it was the only element set defined on the same alignments as our conservation states. We

focused on halves since the LD-score regression estimates can be unstable for annotations

covering too small of a percentage of the genome.[15] To determine the two halves we ranked the

conservation states in descending order of median fold-enrichment of non-exonic bases for

DHS from 123 experiments from the University of Washington ENCODE group.[3] We then

divided bases in PhastCons elements between the top 7 ranked states (1-5, 8 and 28), which

contain 51.9% of bases in PhastCons elements, and the bottom 93 states, which contain the

other 48.1% of bases in PhastCons elements. We applied ldsc to these two sets for 8 traits (age

at menarche, body mass index (BMI), coronary artery disease, educational attainment, height,

low-density lipoprotein (LDL) levels, schizophrenia and smoking behavior), all of which were

previously considered in heritability partitioning analysis.[15] We followed the procedure for

partitioning heritability as done in *Finucane et al.*[15], including using the baseline annotation set

and 500 base-pair windows around annotations to dampen the artificial inflation of heritability in

neighboring regions caused by linkage disequilibrium. The baseline annotation set contains a

range of annotations including DHS. For our analysis, we first removed the constrained element

set already included in the baseline annotation set, then added our two halves of PhastCons

elements and finally ran the ldsc software on the full set of annotations.

Enrichment analysis for variant prioritization scores

For each variant prioritization score included in the conservation state enrichment

analysis of prioritized bases, we extracted the top 1%, 5% and 10% of all the bases ranked by

each score, both genome-wide and just in non-coding regions. The non-coding regions were

defined as the intersection of where the LINSIGHT and FunSeq2 scores provided a value, as

these two scores were only defined on non-coding regions. This intersection results in a set of

bases covering 90% of the genome that excludes coding regions in addition to other regions

filtered for technical reasons by either of the two methods.[19,45] For each score we chose the

score threshold that gave us a size for the top set that was as close as possible to the target

percentage, which did not always exactly match the target percentage due to the precision of

the scores. If a score did not provide a value for a particular base being considered, then that

base was assigned to the lowest value of that score, but would still be counted when

establishing the percentage thresholds. For the scores that provided separate score values for

alternate alleles at a certain position, we used the maximum of the values for all alleles. The

state enrichments were then computed using the OverlapEnrichment command from

ChromHMM at single base resolution ('-b 1' flag) and with the low memory option ('-lowmem'

flag). For the analysis restricted to non-coding regions, we also computed the enrichment of the

states for this background region using the same command. The enrichment for each score in a

state was then divided by the enrichment of the background region for the state. For the Eigen

and Eigen-PC scores we used version 1.1, for FunSeq2 we used version 2.1.6, and for CADD

we used both v1.0 and v1.4.

Results

Annotating the human genome into conservation states

We developed an approach, ConsHMM, to annotate a genome into different

conservation states based on a multiple species DNA sequence alignment (**Figure 1A,**

Methods). We model the combinatorial patterns within the alignment of which species align to

and which match a reference genome, for which we used the human genome. Specifically, at

each nucleotide in the human genome we encode one of three possible observations for each

non-reference species in the alignment: (1) aligns with a nucleotide present that is the same as

the human reference genome, (2) aligns with a nucleotide present that is different than the

human reference genome, or (3) does not have a nucleotide present in the alignment for that

position. We further model these observations as being generated from a multivariate hidden

Markov model (HMM), which probabilistically captures both the combinatorial patterns in the

observations and their spatial context. Specifically, we assume that in each state the probability

of observing a specific combination of observations is determined by a product of independent

multinomial random variables. The parameter values of these multinomial random variables will

differ between states and are learned from the data. After the model is learned, each nucleotide

in the human genome is assigned to the state that had the maximum posterior probability of

generating the observations.

ConsHMM builds on ChromHMM[31], which has previously been applied to annotate

genomes based on epigenomic data at 200-bp resolution[30], to now annotate genomes at single

nucleotide resolution based on a multiple species DNA sequence alignment (**Methods**). We

applied ConsHMM to a 100-way Multiz vertebrate alignment with the human genome and

focused our analysis here on a model learned using 100 states in order to balance recovery of

additional biological features and model tractability (**Figures 2** and **S1-S8, Tables S1** and **S2,**

Methods). We note that HMMs have previously been used to provide local smoothing of signal

for the task of identifying constrained elements.[9,14] We verified that the HMM had a smoothing

effect in our application by comparing to a segmentation derived from a model with the same

emission parameters as our learned model, but that ignored the information in the transition

parameters (**Methods**). We saw an increase in the number of segments from 889 million to 1.06

billion when not using the transition information, though a large majority of the state

assignments to individual bases were the same (**Figure S9**). We illustrate the ConsHMM

conservation state annotations at two different loci showing that different bases that are

associated with calls of evolutionary constraint from existing approaches can have very different

underlying alignment patterns and conservation state assignments (**Figures 1B** and **S10**).

Conservation state annotations genome-wide are available online (**Web Resources**)**.**

Major groups of conservation states

We hierarchically clustered the conservation states based on their align and match

probabilities, and then cut the resulting dendrogram to reveal eight notable groups of states or

distinct individual states (**Figures 2A** and **S4, Table S3, Methods**). We named the resulting

groups based on the aligning and matching properties of major subsets of species for most

states in each group. We also summarized for each individual state the most distal species to

human that had a majority of positions aligning and the closest one that did not, and similarly for

matching (**Table S3**). The first of these subsets of states was a single state (State 1;

AM_allVert) that showed high align and match probabilities through essentially all vertebrate

species considered. The second subset showed relatively high align and match probabilities for

all mammals and some non-mammalian vertebrates (States 2-4; AM_nonMam). The third

subset showed relatively high align and match probabilities for most if not all mammals, but not

non-mammalian vertebrates (States 5-22; AM_Mam). The fourth subset showed high align

probabilities for many mammalian species, but had low align probabilities for notable species

such as mouse and rat for many of the states in the group (States 23-46; AM_SMam). The

combination of the absence of mouse and rat alignments with the presence of mammals that

are assumed to have diverged earlier is consistent with the previously observed increased

substitution rates for mouse and rat.[7] The fifth subset showed high align probabilities for many

mammalian species, but did not show high match probabilities (States 47-63; A_SMam). The

sixth subset showed high align probabilities for most primates, but not for other species (States

64-89; AM_Prim). The seventh subset showed high align probabilities for at most a subset of

primates (States 90-99; AM_SPrim). The final subset was a single state (State 100; artifact) that

showed a noteworthy pattern of high align and match probabilities for most primates and non-

mammalian vertebrates, but low probabilities for non-primate mammals, consistent with a

previous observation that inclusion of non-mammalian vertebrates can be associated with

increased presence of suspiciously aligned regions.[46]

Conservation states exhibit distinct patterns of positional enrichments relative to gene

annotations and regulatory motif instances

The conservation states showed strong and distinct positional enrichments relative to

GENCODE[35] annotated gene features including transcription start sites (TSS), transcription end

sites (TES), exon start sites, and exon end sites for both protein coding genes and

pseudogenes (**Figures 3A-D** and **S11**). Notable positional enrichments were also seen for

regulatory motifs instances (**Figure 3E** and **3F**). Relative to starts of exons of protein coding

genes seven of the states (States 1-4, 7, 28, and 54) had 13 fold or greater enrichment for some

position within 20 base pairs of exon starts, both when considering all such exons and subsets

of exons in specific coding phases (**Figures 3A** and **S11A-C**). These seven states were the only

states that had a majority of positions aligning for at least some non-mammalian vertebrates,

while still having a majority of positions aligning for all primates (**Figure 2A** and **Table S3**).

Within exons we saw the strongest enrichment for states 1-4 and 54, and among these state 1

showed the strongest enrichment, as expected given its high match probabilities through all

vertebrates (**Figures 2B, 3A-B,** and **S11A-E**). Interestingly, state 1 showed very strong

enrichment (>80 fold) in the two nucleotides immediately upstream of the exon start, with the

third upstream nucleotide also having high enrichment (46 fold) (**Figure S11C**). These three

nucleotide positions correspond to the positions of the canonical 3' splice site sequence that is

highly conserved throughout vertebrates.[47] At the ends of exons of protein coding genes

(**Figure 3B**), state 1 maintained a >40 fold enrichment for six nucleotides past the end of coding

sequence corresponding to positions of the known canonical 5' splice site sequence.[47]

Downstream of the start of protein-coding exons, the enrichment profile for state 1 showed a 3-

bp oscillation period, with a dip of enrichment at each 3rd base corresponding to codon wobble

positions. In contrast, states 3 and 54, which were both associated with high align probabilities

through many vertebrates and lower match probabilities, showed the inverse oscillation pattern

to state 1 (**Figures 3A** and **S11A-C**)**.**

Relative to TSS of protein coding genes, state 28 had the strongest enrichment reaching

a maximum enrichment 30 fold at the TSS (**Figure 3C**). State 28 was associated with moderate

align and match probabilities for almost all the species present in the alignment. Consistent with

its enrichment for TSSs state 28 also had the greatest enrichment for CpG islands (32 fold).

However, state 28 also showed a 20 fold enrichment of CpG islands >2kb away from any TSS

of protein coding genes and a 10 fold enrichment for TSS of protein coding genes >2kb away

from a CpG island, suggesting the possibility that both of these features are making a partially

independent contribution to the association, or the presence of additional unannotated TSSs

that are associated with CpG islands.[48] Relative to TES of protein coding genes we saw the

enrichment peak for state 2 at almost 12 fold (**Figure S11F**), which had high align and match

probabilities for almost all vertebrates except for fish.

Relative to pseudogene exon starts and ends, states 100 and 82, both associated with

alignability to distal vertebrates without many mammals closer to human (**Figure 2B** and **Table**

S3), had strong enrichments peaking at greater than 100 and 38 fold respectively (**Figure S11G**

and **S11H**). These two states also showed the greatest enrichment relative to TSSs of

pseudogenes peaking at 184 and 68 fold for states 100 or 82 respectively (**Figure 3D**) and for

TESs of pseudogenes peaking at 199 and 61 fold respectively (**Figure S11I**).

Relative to instances of regulatory motifs, different conservation states showed single

nucleotide enrichment variation, often associated with variation in the amount of information in

the positional-weight matrix (**Figure 3E** and **3F, Methods**).[43] For example, in the case of the

POU5F1 and STAT motifs we saw state 2 from the AM_nonMam group and state 5 from the

AM_Mam group respectively reach 1.8 fold enrichments, but have lower enrichments (1.4-1.5)

at some nucleotides with lower information content. For the STAT motif, states 55-57,

associated with high align probabilities for most mammals, but high match probabilities only for

a few primates, showed enrichments that peaked at the CG dinucleotide in the center of the

motif, consistent with their genome-wide enrichments for CG dinucleotides (**Figures 3E and**

S12).

Enrichment of conservation states for different gene classes

The previous analyses demonstrated that different conservation states have distinct

enrichments in promoter regions of genes. We next investigated whether different conservation

states also exhibit distinct enrichments for different classes of genes after controlling for the

state's relative preference for promoter regions. Specifically, for each state we determined the

5% of genes with the greatest presence of the state in its promoter region and evaluated Gene

Ontology (GO) enrichments for those genes, revealing distinct enrichment patterns (**Figures 4B**

and **S13, Methods**). For example, even among states 1-3, all of which had high alignability

through at least birds and matching through mammals, we observed substantial differences in

their gene preferences. Out of these three states, state 1 (the AM_allVert group) was the only

one enriched for nucleosomes ($p<10^{-41}$; 10.5 fold), while state 3, which had high matching only

through mammals, was the only one with a significant enrichment for a set of genes related to

sensory perception of smell ($p<10^{-300}$; 15.5 fold). State 2, which had high align and match

probabilities through all vertebrates except fish, was the state most enriched for cellular

developmental processes ($p<10^{-30}$; 1.8 fold), which did not show enrichment in state 3. We also

observed notable enrichments for states with overall lower align or match probabilities. For

example, state 89, associated with high alignability and low matching in primates as well as

some alignability and low matching in non-primate mammals, was the state most enriched for

antigen binding (p<10[-14]; 6.7 fold). This is consistent with antigen binding being associated with

many species, but fast evolving.[49]

Enrichments for repeat elements in conservation states

The conservation states showed a wide range of enrichments and depletions (from 2

fold enrichment to 133 fold depletion) for bases overlapping any repeat element (**Figures 2B**

and S8).[21,34] Of the 25 states that did not have any species outside of primates with a majority of

positions aligning, all but two had an enrichment of 1.55 or greater for repeat elements, while

the other 75 states all either had an enrichment below that or did not show enrichment (**Table**

S3). The two exceptions were state 89 and state 96, neither of which showed enrichment for

repeat elements. As noted above, state 89 is likely associated with fast evolving bases shared

with some non-primate mammals, as opposed to bases new to primates. State 96 is associated

with assembly gaps (**Figure S8**). Different repeat classes and families had distinct patterns of

enrichments for different states, even though in some cases the difference in state parameters

was subtle (**Figures 4D** and **S14**). For instance, among states in the AM_Prim group, which

primarily differed in terms of the specific combinations of primates with high align and match

probabilities, we found distinct enrichments. Notably, four different states from the group

AM_Prim, 74, 86, 76, and 77, showed maximal enrichments for the DNA, LINE, LTR, and SINE

repeat classes respectively (**Figure 4D**). State 74, which is characterized by high align and

match probabilities for all primates, had an enrichment of 5.6 fold for DNA repeats, while the

enrichment for the other three classes were between 1.0 and 1.8 fold. On the other hand, state

86, which lacked alignability of a subset of primates, had a 3.0 fold enrichment for LINE repeats,

while the enrichment for the other classes were between 0.6 and 1.6 fold. States 76 and 77 had

3.3 and 4.5 fold enrichments for LTR and SINE respectively compared to 1.1 and 2.1 fold for

SINE and LTR respectively. State 76 and state 77 both had high align probabilities through

primates up to and including squirrel monkey, with the exception that state 77 lacked alignability

to gorilla. Despite these subtle differences in the alignment probabilities, these states had

substantial differences in their repeat enrichment profiles.

Relationship of conservation states to chromatin states

We compared our conservation states to annotations of the genome based on a 25-

chromatin state model defined on 127 samples of diverse cell and tissue types using imputed

data (**Figures 4A** and **S15**).[5,39] For each conservation state we determined the median

enrichment of each chromatin state across the 127 samples. Eleven different conservations

states were maximally enriched for at least one of the 25-chromatin states. Conservation state

28 showed the greatest enrichment for any chromatin state, with a 35 fold enrichment for a

chromatin state associated with active promoters, and was maximally enriched for four other

promoter associated chromatin states. Conservation state 1 was maximally enriched for five

chromatin states all associated with transcribed and exonic regions[39] (3.8-8.7 fold), which is

consistent with this conservation state being most enriched for exons. Conservation state 2 had

the maximal enrichment for five enhancer associated chromatin states (3.1-4.7 fold), while

conservation state 5 had high enrichments for these states and also had the greatest

enrichment of any conservation state for a chromatin state primarily associated with just DNase

I hypersensitivity (2.5 fold). These and other distinct enrichments of the conservation states for

the different chromatin states highlight that conservation states are able to capture multi-

dimensional information in the genome.

Conservation states capture enrichment patterns of DNase I hypersensitive sites across cell and

tissue types

The previous analysis demonstrated that conservation states can exhibit different

enrichment patterns for different chromatin states. We next investigated whether different

conservation states also capture distinct enrichment patterns for a chromatin mark across cell

and tissue types. For this we analyzed DNase I hypersensitive sites (DHS) from 53 of the 127

samples considered above for which maps of experimentally observed DHS were available from

the Roadmap Epigenomics Consortium.[5] We focused on the 21 conservation states that

exhibited at least 2 fold enrichment in at least one sample (**Figure 4C**). We then row normalized

the enrichments in order to focus on the relative enrichment patterns across cell and tissue

types (**Methods**). Hierarchical clustering of the enrichment patterns revealed two major clusters

of states (**Figure 4C**). One of these clusters contained 14 of the 21 states and was associated

with strong enrichments for fetal related samples. Ten of the states in this cluster have

maximum enrichment for a fetal sample, while the remaining four states have maximum

enrichment for the cell type Human Umbilical Vein Endothelial Cells (HUVEC). The second

major cluster consisted of seven states, all of which were enriched for CpG islands (**Figures 2B**

and **S8**). The DHS from samples that showed the greatest enrichments in states in these

clusters also had the greatest enrichment of CpG islands (**Figure 4C, Methods**), but were

biologically diverse in terms of the type of cell or tissue and could potentially reflect technical

experimental differences.

Relationship of conservation states to constraint based annotations

We next investigated the relationship of our conservation state annotations with calls

and univariate scores of evolutionary constraint. Specifically, we considered constrained

element sets based on four methods (GERP++, SiPhy-omega, SiPhy-pi, and PhastCons) and

constraint scores based on three methods (GERP++, PhastCons, and PhyloP) publicly available

for hg19 and also defined on Multiz alignments. The PhastCons and PhyloP scores and

elements we compared to were defined on the same 100-way vertebrate alignment. The

available GERP++, SiPhy-omega, and SiPhy-pi score and elements were derived from different

versions of Multiz alignments and only considered mammals.

We consistently found conservation states 1-5 to be highly enriched (>9 fold) for all

constrained element sets (**Figures 2B** and **S16A**). These states were also among the top six

states in terms of mean score for constraint scores considered (**Figure S16B**). Consistent with

this, states 1-5 were the states that had the highest average matching probability across

mammals. Two other states exhibited at least 6 fold enrichment for at least one constrained

element set: states 54 and 100. State 100, associated with putative artifacts, showed high

enrichments for PhastCons elements (15 fold) and high average scores for PhastCons and

PhyloP. This is consistent with this state having high aligning and matching probabilities

primarily in non-mammalian vertebrates and these elements and scores being defined using

such species. State 54 was consistently enriched for all the constrained elements (4-7 fold), but

did not show high mean base-wise scores particularly for the GERP++ and PhyloP scores. This

difference of high enrichment in constrained elements but not base-wise scores is consistent

with state 54 having high alignability through most vertebrates, but low matching outside

primates. More generally, we found that constrained element calls did not have the resolution to

exhibit biologically relevant single nucleotide variation in enrichments around regulatory motifs

and exon start and ends as we saw with our conservation state annotations, with the exception

of those from PhastCons (**Figures 3** and **S17**).

The objective of our conservation state annotations is different than that of binary calls

and univariate scores of evolutionary constraint, which have a more specific and complementary

goal. However, to better understand their relative biologically relevant information we compared

their ability to recover annotated starts and ends of exons and TSS and TES of genes

separately for protein coding and pseudogenes, as there are well established genome

annotations of these features (**Figures 5A-C** and **S18**). In almost all cases the conservation

states had greater information available for recovering annotated gene features. The only

exceptions were that PhyloP scores could achieve higher precision at low recall levels for

protein coding exon starts and ends, and that SiPhy-pi elements had slightly higher precision for

TSS of protein coding genes at their one recall point.

We also compared the ability of conservation states to recover bases covered by DHS,

both genome-wide and restricted to non-exonic bases, and repeated these analyses when also

restricting to bases distal to a TSS (**Figures S19** and **S20, Methods**). When considering DHS

bases in aggregate over 53 cell and tissue types both genome-wide and restricting to non-

exonic bases, we found that at the same recall level the conservation states could identify bases

in a DHS at greater precision than all constraint scores considered and PhastCons constrained

elements. GERP++, SiPhy-pi and SiPhy-Omega elements did have higher precision at their

single recall point (**Figure S19**). Similar results were seen when just considering distal regions,

except for some of the scores in the non-exonic comparison at very low recall levels. The

relative precision at the same recall levels between conservation states and the GERP++,

SiPhy-pi and SiPhy-Omega constrained element sets did not hold for all cell types (**Figure S20**).

The increase in precision for those constrained element sets in the aggregate evaluation over

constraint scores, PhastCons elements, and ConsHMM annotations might be related to the

coarser resolution at which they were defined (**Figure S17**). We note that the information about

DHS in the conservation states was complementary to that in constrained element sets, as

evidenced by the substantial variation in DHS enrichments of bases within constrained elements

depending on their conservation state (**Figures 5D, S21** and **S22**). For example, bases in

PhastCons constrained elements falling in 35 different states were depleted for Fetal Brain DHS

in non-exonic regions, covering 10% of PhastCons bases, while bases in PhastCons elements

in 12 other states were over 5-fold enriched, covering 37% of PhastCons bases. Additionally,

we saw cases where certain states had greater enrichments for DHS for their bases not in a

constrained element compared to bases in a constrained element in other states. On the other

hand, constrained element calls offered additional information, as we observed that in most

cases, for a given conservation state, bases that were in a constrained element call had greater

enrichment for DHS than those that were not.

We also analyzed the enrichments of our conservation states for previously defined nine-subsets of PhastCons constrained non-exonic elements (CNEEs) based on a directed phylogenetic approach that assigned each element to a phylogenetic branch point of origin (**Figure S23A**).[22] This demonstrated the heterogeneous nature of some of the resulting assignments when relying on directed phylogenetic partitioning approaches. For example, bases in elements assigned to originating at the branch point of the Tetrapod clade showed a high enrichment (37 fold) for state 2, as would be expected since state 2 is associated with aligning and matching through all vertebrates except fish, but an even greater enrichment (51 fold) for state 100, associated with putative artifacts (**Figure S23A**). We also evaluated whether within non-exonic regions any subset of CNEE assigned to a specific clade exhibits enrichments comparable to enrichments seen with the conservation states for CpG islands within non-exonic regions (**Figure S23C**). The most enriched subset of CNEE bases was only 6.7 fold enriched compared to the 37.6 fold enrichment observed for state 28 in non-exonic regions, and only covered 1.9% of non-exonic CpG island bases compared to 12.8% of such bases covered in state 28. A similar pattern of enrichments was observed when considering only the CNEEs overlapping a PhastCons element called on the same alignment as the conservation states (**Figure S23B** and **S23D**). These results highlight that the conservation states are able to capture additional biologically relevant information present in the alignment that is not captured by directed phylogenetic branch assignments of constrained elements.

Bases prioritized by different variant prioritization scores have distinct conservation state enrichment patterns

In addition to scores defined based on just interspecies constraint, a variety of other scores have been proposed to prioritize variants, including some based on intra-species

constraint or integrating inter-species constraint with other genomic annotations. A number of

these scores are widely used, even though a systematic understanding of different types of

bases prioritized by various scores is generally lacking. We leveraged the conservation state

annotations to more systematically understand the bases prioritized by a variety of different

scores in terms of their underlying pattern of conservation.

Specifically, we analyzed the conservation states' genome-wide enrichments of bases

prioritized by 12-different scores (CADD (v1.4), CDTS, DANN, Eigen, Eigen-PC, FATHMM-XF,

FIRE, fitCons, GERP++, PhastCons, PhyloP, and REMM) to be in the top 1, 5, and 10% of the

genome as well as the enrichment specifically in non-coding regions for those 12-scores and

two additional ones defined only on non-coding regions, LINSIGHT and FunSeq2 (**Figures 6A,**

6B and **S24-S27**).[8,9,13,16,18,19,37,45,50–54] We observed an overall strong enrichment for bases

prioritized by most scores for a specific set of conservation states. For example, the top 1%

CADD bases showed a 77.2 fold enrichment for state 1, amounting to 46% of the top 1% CADD

bases falling in this state. This enrichment was greater than that observed for any interspecies

constraint score, despite the CADD score being defined on a diverse set of genomic

annotations, including many non-conservation based annotations. There was a general

consistency in states with higher enrichment across the various measures. For example, when

considering the top 1% bases for the genome-wide analysis, the set of states that were among

the top five most enriched by at least one of the 12 scores contained only 13 states. Nine of

these 13 states (states 1-5, 7, 28, 54, 100) were in the top five for at least three scores.

However, there were important differences for the scores in the relative enrichment among

these top states, and in several cases a single score prioritized other states.

One interesting result was the wide disagreement among the scores of the relative

importance of state 2, the most enhancer enriched state, and state 28, the most promoter

enriched state, particularly in non-coding regions. For example, when considering top 1% bases

in non-coding regions, state 2 was the second or third most enriched state for CADD, Eigen,

FATHMM-XF, GERP++, LINSIGHT, PhastCons, PhyloP, and REMM prioritized variants, with

fold enrichments in the range of 24.9-47.2. On the other hand, state 28 was not one of the top

five most enriched states for any of those scores and its enrichments ranged between 0.3-6.2.

In contrast, for CDTS, DANN, and Eigen-PC state 2 only had enrichments between 0.8-2.1,

while state 28 was the first or second most enriched state for each score, with enrichments

ranging from 7.6-18.6. Also surprising was that DANN showed a depletion for state 2, which

showed high matching through all vertebrates except fish, but enriched for states that were only

associated with subsets of primates. For example, states 92, 93, and 96 did not have an

alignment frequency greater than 0.05 for any species past Gibbon, but were among the top five

states with the greatest enrichments for DANN prioritized variants, with enrichments in the

range 4.2-5.8. None of the other 13 scores considered showed enrichment for these states. This

is despite DANN using the same overall framework as CADD except using a deep neural

network, and previously reporting to be better able to predict the simulated variants used to train

CADD[51]. We verified that this difference with CADD also held for the original version of the

CADD score that used the same features as DANN (**Figure S25**). FitCons and FunSeq2 had

more balanced and relatively lower maximum enrichments for states 2 and 28. The FIRE score

was an outlier in that the maximum enrichment it had for any conservation state was only 2.8.

States for which FIRE prioritized bases showed the greatest enrichment included states such as

77, 80, and 85, which only showed substantial alignments among primates. The FIRE score

was trained based on predicting expression quantitative trait loci (eQTL) in lymphoblastoid cell

lines, which is a very different training objective than the other scores considered. It was

previously noted that this led to background selection being the most important feature to this

score.[50]

There were also strong disagreements about the relative importance of other states

across scores. State 100, associated with likely alignment artifacts, was one such state. For

example, at the top 1% threshold for the non-coding genome analysis, the state was among the

most enriched states for FATHMM-XF, FitCons, PhastCons and PhyloP with enrichments in the

range 14.7-34.5, while other scores showed more modest enrichments or even depletions,

highlighting differences in the vulnerability of each score to likely alignment artifacts. State 54,

which associated with high aligning through most vertebrates but not matching, was another

state with wide disagreement among scores in the importance of those bases, particularly in the

genome-wide analysis. At the top 1% threshold in the genome-wide analyses, this was the third

most enriched state based on CDTS, Eigen-PC, FIRE, and fitCons, with the enrichment for

fitCons reaching 21.1. In contrast, state 54 was depleted for the top GERP++ and REMM bases.

The high enrichment of fitCons for these bases is expected, as the features it considers lack the

resolution to differentiate the third codon position from the more conserved first and second

codon positions when scoring coding regions. There were also differences in the relative

importance given to state 1. For example, when considering variants genome-wide at the top

1% threshold, nine scores had the strongest enrichment for this state, but EIGEN, EIGEN-PC,

and FIRE did not. EIGEN-PC did show the strongest enrichment for state 1 at other thresholds

and EIGEN did when restricting to non-coding genome at all thresholds. However, their

inconsistent ranking of state 1 is likely reflective of the unsupervised prioritization scheme used

by these scores. Overall, these results show that the ConsHMM state annotations provides

insights into key differences in variants prioritized by various scores by systematically and in an

unbiased way capturing biologically diverse classes of nucleotides at single nucleotide

resolution.

Enrichment of conservation states for human genetic variation

Previous analyses have found a depletion of human genetic variation in evolutionarily

constrained elements.[7] Consistent with that, the greatest depletion (3.3 fold depletion) of

common SNPs from dbSNP is in state 1, the state most enriched for constrained elements.

Interestingly, six states, A_SMam states 55-57 and AM_Prim states 87-89, had enrichments in

the range 5 to 8 fold for common SNPs. These were also the six states with greatest enrichment

of CG dinucleotides (**Figure S12**). These six states have in common that they show high align

probabilities for most primates, but low match probabilities for some of those same primates.

These states are thus associated with substantial variation both among primates and among

humans. We observed similar patterns of enrichment for variants identified from whole genome

sequencing (WGS) of a cohort of 7784 unrelated individuals[37], with the levels of state

enrichments and depletions increasing with the minor allele frequency (**Figure S28**).

777   When analyzing the enrichment of GWAS catalog variants[36] relative to the background

of common SNPs we saw opposite enrichment patterns for these states (**Figure 6C** and **6D**).

For example, relative to this background, state 1 was most enriched for GWAS catalog variants,

which is consistent with previous observations of constrained elements enriching for GWAS

variants.[7] On the other hand, states 55-57 and 87-89 showed the greatest depletion. These

results suggest that common variants are less likely to be phenotypically significant if they fall in

conservation states most enriched for common genetic variation.

Constrained element enrichment for partitioned heritability of complex traits depends on

conservation state

787   Previous analyses have suggested a strong enrichment of constrained elements and

DHS for phenotype heritability.[15,55] As we saw large differences in DHS enrichments of

constrained elements depending on the conservation state, we investigated the extent to which

constrained elements in conservation states most enriched for DHS enriched for phenotype

heritability compared to the remaining states. Specifically, we ranked the conservation states in

descending order of their median enrichment for DHS from a compendium of 123 experiments

from the ENCODE consortium, within the non-exonic portion of the state (**Figure 2B,**

Methods).[3] We then partitioned bases in PhastCons constrained elements into two almost

equal size sets based on whether they overlapped one of the top seven ranked conservation

states (states 1-5, 8, 28) or not. We then computed the heritability for these two sets for eight

phenotypes in the context of a set of baseline annotations that include DHS annotations

(**Methods**).[15] For seven of the phenotypes, we found that bases in constrained elements

overlapping the top seven states had greater enrichment than those in the remaining 93 states,

often substantially so (**Figure 6E**). These results suggest additional value in the conservation

state annotations for isolating more likely disease relevant variants.

Discussion

We presented a framework for genome annotation based on comparative genomics

sequence data. Our approach learns a set of conservation states *de novo* using a multivariate

HMM based on the combinatorial and spatial patterns of which species align and match a

reference genome in a multi-species DNA sequence alignment. We applied this approach to

annotate the human genome at single nucleotide resolution into one of 100 conservation states.

Conservation state annotations exhibited substantial enrichments for a wide range of other

genomic annotations that were not provided to the model in training, thus supporting their

biological significance. Specific conservation states exhibited strong enrichments for various

gene annotations including exons, TSS and TES of genes, while others showed strong

enrichments for specific types of repeat elements. Conservation states showed differential

enrichment patterns for various classes of genes and DHS from multiple cell types, even though

they were defined independently of any functional genomics data. Specific conservation states

exhibited enrichments for common human variants, while a different set of states exhibited

enrichments for variants identified by GWAS relative to common variation.

ConsHMM differs from other comparative genomics based annotation approaches in

several respects. One difference is that it takes an unsupervised approach that does not

explicitly use a phylogenetic tree, except to the extent to which a phylogenetic tree was used to

generate the input multi-species sequence alignment. This leads to relatively unbiased, simple

and interpretable models. However, many state patterns discovered are consistent with

expected observations from commonly assumed phylogenetic relationships of the species.

While states' parameters often decreased with divergence time from human, there were a

number of exceptions. Some of these exceptions corresponded to missing specific sub-clades

of species, particularly those with long branch lengths. For example, a number of states were

not represented by mouse and rat, while being represented by more distally diverged mammals.

Other exceptions isolated putative artifacts in the alignments that might otherwise confound

analyses, as we saw for two states heavily enriched for pseudogenes. ConsHMM also differs

from other commonly used modeling approaches in how it explicitly differentiates non-aligning

bases from aligning non-matching bases, which allowed it, for example, to identify states

particularly associated with third codon positions. Another difference between the ConsHMM

annotations and standard constraint measures is that the ConsHMM annotations are defined

directly relative to the variant present in the genome being annotated. When applying

ConsHMM to annotate the human genome, a mutation unique to human would be expected to

have a much larger effect on the ConsHMM annotations than a mutation unique to a single

other species. This would not in general be expected to be the case for constraint measures

that do not differentiate the target genome for annotation from other genomes in an alignment.

An interesting future direction would be to produce and analyze individual specific ConsHMM

annotations.

Our conservation state annotation is complementary to existing binary calls and scores

of evolutionary constraint based on phylogenetic modeling. Both locations called as constrained

and those called as non-constrained are heterogeneous in their assigned conservation state.

Our annotations thus provide additional descriptive information about the conservation patterns

at each base. In terms of information for predicting external annotations, we found that in many

cases the conservation states had greater information than constraint scores or elements.

Notably, our modeling approach identified a conservation state, state 28, associated with a

pattern of aligning and matching some mammalian and non-mammalian vertebrates, but not

with high probability for any one species. This conservation state strongly enriched for

transcription start sites and CpG islands, and was not well captured by phylogenetic modeling

approaches. For other cases, such as DHS, the relative information depended on the

constrained element set or score being compared. Importantly, we observed that DHS

information provided by the states was complementary to information in the constrained element

calls. We also used the conservation state annotations to systematically understand key

similarities and differences in the patterns of conservation in bases prioritized by a large number

of different variant prioritization scores, including scores based on integrating diverse features

(**Figure 6A** and **6B**). The conservation state annotations provide a powerful framework for

gaining such an understanding, since the corresponding conservation patterns are defined

systematically in an unbiased way, at single nucleotide resolution and capture a diverse set of

biological features. Furthermore, we observed that bases in constrained elements showed

substantially different enrichments for phenotype-associated heritability, depending on their

conservation state.

The conservation states are both inspired by, and provide complementary information to,

existing chromatin state annotation approaches. While the states from the two approaches are

based on very different data and have fundamental differences, they also exhibited substantial

cross-enrichments. In general, conservation states have the advantage of providing information

at single nucleotide resolution, which we demonstrated by showing enrichments patterns in and

around coding exons and regulatory motifs. Conservation states can also provide information

about bases in the genome even if the relevant cell type has not been experimentally profiled,

while chromatin states have the advantage of directly providing cell type specific information.

We expect many applications for the methodology and annotations we have presented

here. While we applied ConsHMM here to one multiple species alignment, a 100-way Multiz

human alignment, the methodology is general, and thus can be readily applied to alignments to

other species or alignments generated by other methods.[26] The annotations we produced serve

as a resource to directly interpret other genomic datasets or variant prioritization scores. They

could also potentially be integrated with other complementary genomic annotations in methods

that produce integrated variant prioritization scores. This work represents a step in the direction

of improving whole genome annotations, which will continue to be of increasing importance

towards understanding health and disease as the availability of whole genome sequencing data

increases.

Supplemental Data

Supplemental Data include twenty-eight figures and three tables.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgements

We thank Ewan Birney and members of the Ernst lab for useful discussions. We

acknowledge funding from US National Institutes of Health grants DP1DA044371,

R01ES024995, U01HG007912 and U01MH105578 (J.E.), and T32CA201160 (A.S.), US

National Science Foundation CAREER Award #1254200 (J.E.), a Kure-IT award and an Alfred

P. Sloan Fellowship (J.E.).

Web Resources

25-state chromatin state annotations:

http://compbio.mit.edu/roadmap

CADD score v1.0:

http://krishna.gs.washington.edu/download/CADD/v1.0/whole_genome_SNVs.tsv.gz

CADD score v1.4:

http://krishna.gs.washington.edu/download/CADD/v1.4/GRCh37/whole_genome_SNVs.t

sv.gz

CDTS score:

http://www.hli-opendata.com/noncoding/coord_CDTS_percentile_N7794unrelated.txt.gz

http://www.hli-opendata.com/noncoding/SNVusedForCDTScomputation_N7794unrelate

d_allelicFrequency0.001truncated.txt.gz

CNEEs from Lowe et al.[22]:

http://www.stanford.edu/~lowec/data/threePeriods/hg19cnee.bed.gz

ConsHMM v1.0 software and ConsHMM state annotations:

https://github.com/ernstlab/ConsHMM

DANN score:

https://cbcl.ics.uci.edu/public_data/DANN/data/

EIGEN and Eigen-PC score:

https://xioniti01.u.hpc.mssm.edu/v1.1/

ENCODE DHS:

http://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeUwDnase/

FATHMM-XF score:

http://fathmm.biocompute.org.uk/fathmm-xf/

FIRE score:

https://sites.google.com/site/fireregulatoryvariation/

fitCons score:

http://compgen.cshl.edu/fitCons/0downloads/tracks/i6/scores/

FunSeq2 score:

http://org.gersteinlab.funseq.s3-website-us-east-

1.amazonaws.com/funseq2.1.2/hg19_NCscore_funseq216.tsv.bgz

GENCODE v19:

https://www.gencodegenes.org/releases/19.html

GERP++ scores and constrained element calls:

http://mendel.stanford.edu/SidowLab/downloads/gerp/

GWAS catalog variants:

LINSIGHT score:

http://compgen.cshl.edu/~yihuang/tracks/LINSIGHT.bw

Motif instances and background:

http://compbio.mit.edu/encode-motifs/

https://www.ebi.ac.uk/gwas/

Multiz 100-way alignment to hg19 reference:

http://hgdownload.soe.ucsc.edu/goldenPath/hg19/multiz100way/

REMM score:

https://zenodo.org/record/1197579/files/ReMM.v0.3.1.tsv.gz

Roadmap Epigenomics DHS:

http://egg2.wustl.edu/roadmap/data/byFileType/peaks/consolidated/narrowPeak/

SiPhy-omega and SiPhy-pi constrained element calls (hg19 liftOver):

https://www.broadinstitute.org/mammals-models/29-mammals-project-supplementary-

info

References

1. Hindorff, L.A., Sethupathy, P., Junkins, H.A., Ramos, E.M., Mehta, J.P., Collins, F.S.,
and Manolio, T.A. (2009). Potential etiologic and functional implications of genome-wide
association loci for human diseases and traits. Proc. Natl. Acad. Sci. U. S. A. *106*,
9362–9367.

2. Ward, L.D., and Kellis, M. (2012). Interpreting non-coding variation in complex
disease genetics. Nat. Biotechnol. *30*, 1095–1106.

3. ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in
the human genome. Nature *489*, 57–74.

4. Ernst, J., Kheradpour, P., Mikkelsen, T.S., Shoresh, N., Ward, L.D., Epstein, C.B.,
Zhang, X., Wang, L., Issner, R., Coyne, M., et al. (2011). Mapping and analysis of
chromatin state dynamics in nine human cell types. Nature *473*, 43–49.

5. Roadmap Epigenomics Consortium, Kundaje, A., Meuleman, W., Ernst, J., Bilenky,
M., Yen, A., Heravi-Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J., et al. (2015).
Integrative analysis of 111 reference human epigenomes. Nature *518*, 317–330.

6. Kellis, M., Wold, B., Snyder, M.P., Bernstein, B.E., Kundaje, A., Marinov, G.K., Ward,
L.D., Birney, E., Crawford, G.E., Dekker, J., et al. (2014). Defining functional DNA
elements in the human genome. Proc. Natl. Acad. Sci. *111*, 6131–6138.

7. Lindblad-Toh, K., Garber, M., Zuk, O., Lin, M.F., Parker, B.J., Washietl, S.,
Kheradpour, P., Ernst, J., Jordan, G., Mauceli, E., et al. (2011). A high-resolution map of
human evolutionary constraint using 29 mammals. Nature *478*, 476–482.

8. Pollard, K.S., Hubisz, M.J., Rosenbloom, K.R., and Siepel, A. (2010). Detection of
nonneutral substitution rates on mammalian phylogenies. Genome Res. *20*, 110–121.

9. Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K.,
Clawson, H., Spieth, J., Hillier, L.W., Richards, S., et al. (2005). Evolutionarily
conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Res. *15*,
1034–1050.

10. Cooper, G.M., and Shendure, J. (2011). Needles in stacks of needles: finding
disease-causal variants in a wealth of genomic data. Nat. Rev. Genet. *12*, 628–640.

11. Weedon, M.N., Cebola, I., Patch, A.-M., Flanagan, S.E., De Franco, E., Caswell, R.,
Rodríguez-Seguí, S.A., Shaw-Smith, C., Cho, C.H.-H., Allen, H.L., et al. (2014).
Recessive mutations in a distal PTF1A enhancer cause isolated pancreatic agenesis.
Nat. Genet. *46*, 61–64.

12. Cooper, G.M., Stone, E.A., Asimenos, G., Green, E.D., Batzoglou, S., and Sidow, A.
(2005). Distribution and intensity of constraint in mammalian genomic sequence.
Genome Res. *15*, 901–913.

13. Davydov, E.V., Goode, D.L., Sirota, M., Cooper, G.M., Sidow, A., and Batzoglou, S.
(2010). Identifying a High Fraction of the Human Genome to be under Selective
Constraint Using GERP++. PLoS Comput. Biol. *6*, e1001025.

14. Garber, M., Guttman, M., Clamp, M., Zody, M.C., Friedman, N., and Xie, X. (2009).
Identifying novel constrained elements by exploiting biased substitution patterns.
Bioinformatics *25*, i54–i62.

15. Finucane, H.K., Bulik-Sullivan, B., Gusev, A., Trynka, G., Reshef, Y., Loh, P.-R.,
Anttila, V., Xu, H., Zang, C., Farh, K., et al. (2015). Partitioning heritability by functional
annotation using genome-wide association summary statistics. Nat. Genet. *47*, 1228–
1235.

16. Kircher, M., Witten, D.M., Jain, P., O'Roak, B.J., Cooper, G.M., and Shendure, J.
(2014). A general framework for estimating the relative pathogenicity of human genetic
variants. Nat. Genet. *46*, 310–315.

17. Ritchie, G.R.S., Dunham, I., Zeggini, E., and Flicek, P. (2014). Functional annotation
of noncoding sequence variants. Nat. Methods *11*, 294–296.

18. Ionita-Laza, I., McCallum, K., Xu, B., and Buxbaum, J.D. (2016). A spectral
approach integrating functional genomic annotations for coding and noncoding variants.
Nat. Genet. *48*, 214–220.

19. Huang, Y.-F., Gulko, B., and Siepel, A. (2017). Fast, scalable prediction of
deleterious noncoding variants from functional and population genomic data. Nat.
Genet. *49*, 618–624.

20. Jagadeesh, K.A., Wenger, A.M., Berger, M.J., Guturu, H., Stenson, P.D., Cooper,
D.N., Bernstein, J.A., and Bejerano, G. (2016). M-CAP eliminates a majority of variants
of uncertain significance in clinical exomes at high sensitivity. Nat. Genet. *48*, 1581–
1586.

21. Rosenbloom, K.R., Armstrong, J., Barber, G.P., Casper, J., Clawson, H., Diekhans,
M., Dreszer, T.R., Fujita, P.A., Guruvadoo, L., Haeussler, M., et al. (2015). The UCSC
Genome Browser database: 2015 update. Nucleic Acids Res. *43*, D670-681.

22. Lowe, C.B., Kellis, M., Siepel, A., Raney, B.J., Clamp, M., Salama, S.R., Kingsley,
D.M., Lindblad-Toh, K., and Haussler, D. (2011). Three periods of regulatory innovation
during vertebrate evolution. Science *333*, 1019–1024.

23. Siepel, A., Pollard, K.S., and Haussler, D. (2006). New Methods for Detecting
Lineage-Specific Selection. In Research in Computational Molecular Biology, (Springer,
Berlin, Heidelberg), pp. 190–205.

24. Kim, S.Y., and Pritchard, J.K. (2007). Adaptive Evolution of Conserved Noncoding
Elements in Mammals. PLOS Genet. *3*, e147.

25. Marnetto, D., Mantica, F., Molineris, I., Grassi, E., Pesando, I., and Provero, P.
(2018). Evolutionary Rewiring of Human Regulatory Networks by Waves of Genome
Expansion. Am. J. Hum. Genet. *102*, 1–12.

26. Herrero, J., Muffato, M., Beal, K., Fitzgerald, S., Gordon, L., Pignatelli, M., Vilella,
A.J., Searle, S.M.J., Amode, R., Brent, S., et al. (2016). Ensembl comparative genomics
resources. Database J. Biol. Databases Curation *2016*,.

27. Cotney, J., Leng, J., Yin, J., Reilly, S.K., DeMare, L.E., Emera, D., Ayoub, A.E.,
Rakic, P., and Noonan, J.P. (2013). The Evolution of Lineage-Specific Regulatory
Activities in the Human Embryonic Limb. Cell *154*, 185–196.

28. Villar, D., Berthelot, C., Aldridge, S., Rayner, T.F., Lukk, M., Pignatelli, M., Park,
T.J., Deaville, R., Erichsen, J.T., Jasinska, A.J., et al. (2015). Enhancer Evolution
across 20 Mammalian Species. Cell *160*, 554–566.

29. Don, P.K., Ananda, G., Chiaromonte, F., and Makova, K.D. (2013). Segmenting the
human genome based on states of neutral genetic divergence. Proc. Natl. Acad. Sci.
110, 14699–14704.

30. Ernst, J., and Kellis, M. (2010). Discovery and characterization of chromatin states
for systematic annotation of the human genome. Nat. Biotechnol. *28*, 817–825.

31. Ernst, J., and Kellis, M. (2012). ChromHMM: automating chromatin-state discovery
and characterization. Nat. Methods *9*, 215–216.

32. Hoffman, M.M., Buske, O.J., Wang, J., Weng, Z., Bilmes, J.A., and Noble, W.S.
(2012). Unsupervised pattern discovery in human chromatin structure through genomic
segmentation. Nat. Methods *9*, 473–476.

33. Blanchette, M., Kent, W.J., Riemer, C., Elnitski, L., Smit, A.F.A., Roskin, K.M.,
Baertsch, R., Rosenbloom, K., Clawson, H., Green, E.D., et al. (2004). Aligning Multiple
Genomic Sequences With the Threaded Blockset Aligner. Genome Res. *14*, 708–715.

34. Smit, A., Hubley, R., and Green, P. (2013). RepeatMasker Open-4.0.

35. Harrow, J., Frankish, A., Gonzalez, J.M., Tapanari, E., Diekhans, M., Kokocinski, F.,
Aken, B.L., Barrell, D., Zadissa, A., Searle, S., et al. (2012). GENCODE: The reference
human genome annotation for The ENCODE Project.Genome Res. *22*,1760–1774.

36. Welter, D., MacArthur, J., Morales, J., Burdett, T., Hall, P., Junkins, H., Klemm, A.,
Flicek, P., Manolio, T., Hindorff, L., et al. (2014). The NHGRI GWAS Catalog, a curated
resource of SNP-trait associations. Nucleic Acids Res. *42*, D1001-1006.

37. Iulio, J. di, Bartha, I., Wong, E.H.M., Yu, H.-C., Lavrenko, V., Yang, D., Jung, I.,
Hicks, M.A., Shah, N., Kirkness, E.F., et al. (2018). The human noncoding genome
defined by genetic diversity. Nat. Genet. *50*, 333-337.

38. Witowski, V., and Foraita, D.R. (2014). HMMpa: Analysing accelerometer data using
hidden Markov models.

39. Ernst, J., and Kellis, M. (2015). Large-scale imputation of epigenomic datasets for
systematic annotation of diverse human tissues. Nat. Biotechnol. *33*, 364–376.

40. Hahsler, M. and Buchta, C.(2017). cba: Clustering for Business Analytics.

41. Bar-Joseph, Z., Gifford, D.K., and Jaakkola, T.S. (2001). Fast optimal leaf ordering
for hierarchical clustering. Bioinforma. Oxf. Engl. *17 Suppl 1*, S22-29.

42. Ernst, J., and Bar-Joseph, Z. (2006). STEM: a tool for the analysis of short time
series gene expression data. BMC Bioinformatics *7*, 191.

43. Kheradpour, P., and Kellis, M. (2014). Systematic discovery and characterization of
regulatory motifs in ENCODE TF binding experiments. Nucleic Acids Res. *42*, 2976–
2987.

44. Kolde, R. (2015). pheatmap: Pretty Heatmaps.

45. Fu, Y., Liu, Z., Lou, S., Bedford, J., Mu, X.J., Yip, K.Y., Khurana, E., and Gerstein,
M. (2014). FunSeq2: a framework for prioritizing noncoding regulatory variants in
cancer. Genome Biol. *15*, 480.

46. Chen, X., and Tompa, M. (2010). Comparative assessment of methods for aligning
multiple genome sequences. Nat. Biotechnol. *28*, 567–572.

47. Zhang, M.Q. (1998). Statistical features of human exons and their flanking regions.
Hum. Mol. Genet. *7*, 919–932.

48. Sarda, S., Das, A., Vinson, C., and Hannenhalli, S. (2017). Distal CpG islands can
serve as alternative promoters to transcribe genes with silenced proximal promoters.
Genome Res. *27*, 553–566.

49. Litman, G.W., Anderson, M.K., and Rast, and J.P. (1999). Evolution of Antigen
Binding Receptors. Annu. Rev. Immunol. *17*, 109–147.

50. Ioannidis, N.M., Davis, J.R., DeGorter, M.K., Larson, N.B., McDonnell, S.K., French,
A.J., Battle, A.J., Hastie, T.J., Thibodeau, S.N., Montgomery, S.B., et al. (2017). FIRE:
functional inference of genetic variants that regulate gene expression. Bioinforma. Oxf.
Engl. *33*, 3895–3901.

51. Quang, D., Chen, Y., and Xie, X. (2015). DANN: a deep learning approach for
annotating the pathogenicity of genetic variants. Bioinforma. Oxf. Engl. *31*, 761–763.

52. Rogers, M.F., Shihab, H.A., Mort, M., Cooper, D.N., Gaunt, T.R., and Campbell, C.
(2018). FATHMM-XF: accurate prediction of pathogenic point mutations via extended
features. Bioinforma. Oxf. Engl. *34*, 511–513.

53. Gulko, B., Hubisz, M.J., Gronau, I., and Siepel, A. (2015). A method for calculating
probabilities of fitness consequences for point mutations across the human genome.
Nat. Genet. *47*, 276–283.

54. Smedley, D., Schubach, M., Jacobsen, J.O.B., Köhler, S., Zemojtel, T., Spielmann,
M., Jäger, M., Hochheiser, H., Washington, N.L., McMurry, J.A., et al. (2016). A Whole-

Genome Analysis Framework for Effective Identification of Pathogenic Regulatory
Variants in Mendelian Disease. Am. J. Hum. Genet. *99*, 595–606.

55. Gusev, A., Lee, S.H., Trynka, G., Finucane, H., Vilhjálmsson, B.J., Xu, H., Zang, C.,
Ripke, S., Bulik-Sullivan, B., Stahl, E., et al. (2014). Partitioning Heritability of
Regulatory and Cell-Type-Specific Variants across 11 Common Diseases. Am. J. Hum.
Genet. *95*, 535–552.

1098

1099
1100
1101
Figure Legends

Figure 1: Illustration of ConsHMM modeling approach. (A) The input to ConsHMM is a

multi-species alignment, which is illustrated for a subset of 6 species aligned to the human

sequence. At each position and for each species ConsHMM represents the information as one

of three observations: (1) aligns with a non-indel nucleotide matching the human sequence

shown in blue, (2) aligns with a non-indel nucleotide not matching the human sequence shown

in yellow, or (3) does not align with a non-indel nucleotide shown in gray. **(B)** Illustration of

conservation state assignments at a locus chr22:25,024,640-25,024,812. Only states assigned

to at least one nucleotide in the locus are shown. Below the conservation state assignments is a

color encoding of the input multiple sequence alignment according to panel (A). The major clade

of species as annotated on the UCSC genome browser[21] are labeled and ordered based on

divergence from human. Above the conservation state assignments are PhastCons constrained

elements and scores and PhlyoP constraint scores. This figure and **Figure S9** together illustrate

that positions of nucleotides that have the same status in terms of being in a constrained

element or not or have similar constraint scores can be assigned to different conservation states

depending on the patterns in the underlying multiple-species alignment.

1118

Figure 2: Conservation state emission parameters learned by ConsHMM and enrichments

for other genomic annotations. (A) Each row in the heatmap corresponds to a conservation

state. For each state and species, the left half of the heatmap gives the probability of aligning to

the human sequence, which is one minus the probability of the not aligning emission.

Analogously, the right half of the heatmap gives the probability of observing the matching

emission. Each individual column corresponds to one species with the individual names

displayed in **Figure S5.** For both halves, species are grouped by the major clades and ordered

based on the hg19.100way.nh phylogenetic tree from the UCSC genome browser, with species

that diverged more recently shown closer to the left.[21] The conservation states are ordered

based on the results of applying hierarchical clustering and optimal leaf ordering.[41] The states

are divided into eight major groups based on cutting the dendrogram of the clustering. The

groups are indicated by color bars on the left hand side and a white row between them.

Transition parameters between states of the model can be found in **Figure S6**. **(B)** The columns

of the heatmap indicate the relative enrichments of conservation states for external genomic

annotations (**Methods**). For each column, the enrichments were normalized to a [0,1] range by

subtracting the minimum value of the column and dividing by the range and colored based on

the indicated scale. Values for these enrichments and additional enrichments can be found in

Figure S8 and **Table S2** and enrichments for individual repeat classes and families can be

found in **Figure S14.**

1138

Figure 3: Conservation state positional enrichments. Plots of positional fold enrichments of

conservation states relative to **(A)** start of exons of protein coding genes in phase 0, **(B)** end of

exons of protein coding genes and **(C,D)** TSS of **(C)** protein coding, and **(D)** pseudogenes

genes. Positive values represent the number of bases downstream in the 5' to 3' direction of

transcription, while negative values represent the number of bases upstream. Enrichments

relative to gene annotations are based on a genome-wide background. The subset of states

included in panels (A)-(D) were the states that had at least a 3 fold enrichment at some position

within +/-2kb from the anchor point. **(E,F)** Also shown are positional plots relative to the central

nucleotide of a set of instances of **(E)** STAT and **(F)** POU5F1 motifs. The subset of states

included in (E), (F) are the states that had an enrichment of at least 1.5 for some position within

+/-15bp from the center nucleotide of either motif. Enrichments for motif instances were

computed relative to the portion of the genome scanned for regulatory motifs in Kheradpour and

Kellis (2014), which excludes coding, 3'UTRs, and repeat elements. Additional position

enrichment plots can be found in **Figure S11**.

1153

Figure 4: Conservation states enrichment for chromatin states, GO terms, DHS and

repeat elements. (A) Median fold enrichment of conservation states (rows) for one of 25

chromatin states from a previously defined chromatin state model defined across 127 samples

of diverse cell and tissue types (columns).[15] Only conservation states that had the maximum

value for at least one chromatin state are shown, and those values are boxed. See **Figure S15**

for the enrichments of all conservation states. **(B)** $-\log_{10}$ p-value (uncorrected) of the

conservation states (rows) for the GO term (columns) where each conservation state is

associated with its top 5% genes based on promoter regions (**Methods**). Only GO terms which

were the most enriched term for some conservation state are shown, restricted to the top 10

terms based on the significance of the enrichment. Only conservation states that had the most

significant enrichment for one of the displayed GO terms are shown, with the maximal

enrichments boxed. The full set of conservation states with additional GO terms are in **Figure**

S13. **(C)** Relative enrichments of conservation states for DHS across cell and tissue types. Only

conservation states with at least a 2 fold enrichment in one sample considered are shown.

Enrichment values were $\log_2$ transformed and then row normalized by subtracting the mean

(right heatmap) and dividing by the standard deviation. States and experiments were then

hierarchically clustered and revealed two major clusters. In the top cluster conservation states

showed the greatest enrichment for experiments in which the DHS also strongly enriched for

CpG islands (top heatmap). In the bottom cluster conservation states generally had the

strongest relative preference for a number of fetal related samples. **(D)** Enrichment of

conservation states with the maximal enrichment for LINE, SINE, LTR or DNA repeats next to

the state align probabilities for primates. These states all had low align probabilities outside of

primates, but their differences among primates corresponded to substantial differences in repeat

enrichments.

1178

Figure 5: **Relationship of conservation states with constrained elements and scores.**

Precision-recall plots for recovery of **(A)** TSS of protein coding genes, **(B)** TES of protein coding

genes, and **(C)** the start of exons of protein coding genes. Recovery based on ordering

ConsHMM conservation states for their enrichment for the target set in the training data, then

cumulatively adding the states in that ranked order and evaluating on the test data is shown with

a series of blue dots (**Methods**). The first few conservation states added are labeled with their

state number. Recovery based on ranking from highest to lowest value of constraint scores is

shown with continuous lines. Recovery based on score partitioning into 400 bins and

subsequent ordering based on enrichment for the target set in the training data, then

cumulatively adding bins in that ranked order and evaluating on the test data is shown in a

series of dots of the same color as the continuous line corresponding to the score. Recovery of

target test bases by a constrained element set is shown with a single dot for each constrained

element set. See **Figure S18-20** for plots based on additional targets. **(D)** The graph shows the

fold enrichment for Fetal Brain DHS[5] within the non-exonic portion of each conservation state,

separately for those bases in a PhastCons constrained element (pink) and bases not in such an

element (blue). Enrichments within constrained elements varied substantially depending on the

conservation state. For a given conservation state, bases in a constrained element had greater

enrichments than bases not in a constrained element, illustrating complementary information of

conservation states and constrained elements. See **Figure S21** for graphs based on different

element sets or DHS data and **Figure S22** for these enrichments plotted against the size of the

set.

1200

Figure 6: Conservation states and association with human genetic variation. (A) Fold

enrichments of bases ranked in the top 1% genome-wide by 12 variant prioritization scores.

Only states that were among the top five most enriched states for at least one score are shown.

The enrichment of the top five ranking states for each score is colored according to the ranking

and the color scale shown on right. **(B)** Enrichments of bases ranked in the top 1% of the non-

coding genome by 14 variant prioritization scores. The criteria for selecting states to display and

coloring enrichments was the same as in panel (A). Enrichments at additional thresholds and for

all states both genome-wide and for the non-coding genome are in **Figure S24-S27**. The

enrichments for CADD shown here are based on v1.4, while enrichments based on the original

version of CADD are also shown in **Figure S25-S27**. (**C**) The panel displays the $\log_2$ fold

enrichment of each state for common SNPs (pink) and GWAS catalog variants relative to

common SNPs (blue). State 1, associated with high alignability and matching through all

vertebrates, showed the greatest depletion of common SNPs and the highest enrichment for

GWAS variants relative to common SNPs. States 55-57 and 87-89 exhibited the opposite

pattern having the greatest enrichment for common SNPs and the greatest depletion of GWAS

variants relative to this background. The second most depleted state for common SNPs, which

did not show enrichment for GWAS catalog SNPs, was state 96 which captured large gaps in

the assembly (**Figure S8**). (**D**) Panel shows the representation of state emission parameters

from **Figure 2A** for the subset of states highlighted in panel (C). The states with the greatest

depletion of GWAS variants all had relatively high alignability at least through primates, but low

matching probabilities for almost all species except a few closely related primates. (**E**) Applying

the heritability partitioning enrichment method of Finucane et al.[15] on two disjoint subsets of

bases in PhastCons elements, with eight phenotypes previously analyzed with heritability

partitioning in the context of a baseline set of annotations (**Methods**).[15] One set of bases are

those in PhastCons elements that are also in one of the seven conservation states showing the

greatest enrichment for DHS in its non-exonic portion (States 1-5, 8, and 28) covering 51.9% of

PhastCons bases (pink). The other set are those bases in PhastCons elements overlapping the

remaining 93 states covering 48.1% of PhastCons bases (blue).

A

Human	T	T	T	C	C	T	G	A	C	T	T
Chimp	T	T	T	C	C	T	G	A	C	T	T
Bushbaby	T	C	T	G	C	T	T	C	C	T	T
Rat	-	C	T	T	C	T	G	A	A	T	T
Alpaca	-	-	-	C	C	T	T	G	C	A	T
Megabat	T	C	-	C	C	T	G	A	T	T	T
Parrot	-	-	-	-	-	-	-	-	-	-	-

■ Aligning and matching the human sequence

■ Aligning but not matching the human sequence

■ Not aligning to the human sequence

B

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6