Systematic Discovery of Conservation States for Single-Nucleotide Annotation of

the Human Genome

Adriana Sperlea[1,2] and Jason Ernst[1,2,3,4,5,6]

[1] Bioinformatics Interdepartmental Program, University of California, Los Angeles, Los Angeles, California,

90095, USA.

[2] Department of Biological Chemistry, University of California, Los Angeles, Los Angeles, California, USA.

[3] Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research at University of

California, Los Angeles, Los Angeles, California, 90095, USA.

[4] Computer Science Department, University of California, Los Angeles, Los Angeles, California, 90095,

USA.

[5] Jonsson Comprehensive Cancer Center, University of California, Los Angeles, Los Angeles, California,

90095, USA.

[6] Molecular Biology Institute, University of California, Los Angeles, Los Angeles, California, 90095, USA.

Correspondence: Jason Ernst (jason.ernst@ucla.edu)

Abstract

Comparative genomics sequence data is an important source of information for interpreting

genomes. Genome-wide annotations based on this data have largely focused on univariate

scores or binary calls of evolutionary constraint. Here we present a complementary whole

genome annotation approach, ConsHMM, which applies a multivariate hidden Markov model to

learn *de novo* different 'conservation states' based on the combinatorial and spatial patterns of

which species align to and match a reference genome in a multiple species DNA sequence

alignment. We applied ConsHMM to a 100-way vertebrate sequence alignment to annotate the

human genome at single nucleotide resolution into 100 different conservation states. These

states have distinct enrichments for other genomic information including gene annotations,

chromatin states, and repeat families, which were used to characterize their biological

significance. Conservation states have greater or complementary predictive information than

standard constraint based measures for a variety of genome annotations. Bases in constrained

elements have distinct heritability enrichments depending on the conservation state assignment

demonstrating their relevance to analyzing phenotypic associated variation. The conservation

states also highlight similarities and differences between constrained bases identified based on

inter and intra species approaches. The ConsHMM method and conservation state annotations

provide a valuable resource for interpreting genetic variation.

Introduction

The large majority of phenotype-associated variants implicated by genome-wide

association studies (GWAS) fall outside of protein coding regions.[1] Identifying the causal

variants and interpreting their biological role in these less well understood non-coding regions is

a significant challenge.[2] Large scale mapping of epigenomic data across different cell and tissue

types has been one approach for annotating and interpreting the non-coding regions of the

genome.[3–5] Using comparative genomics data to identify regions of evolutionary constraint has

been a complementary approach for these purposes.[6–9]

In addition to providing evolutionary information, comparative genomics data has the

advantage of providing information at single-nucleotide resolution. Furthermore, it is cell type

agnostic and thus informative even when the relevant cell or tissue type has not been

experimentally profiled.[10,11] The most commonly used representations of this information are

univariate scores and binary elements of evolutionary constraint, which are called based on a

multiple species DNA sequence alignment and assumed models of evolution and selection.[8,9,12–

14] Supporting the importance of these annotations, heritability analyses have recently implicated

evolutionary constrained elements as one of the annotations most enriched for phenotype

associated variants.[15] These scores and elements have also been highly informative features to

integrative methods for prioritizing pathogenic variants.[16–19] Further improvements for predicting

pathogenic variants in coding regions have been made to the integrative scores by incorporating

features defined directly from a multiple sequence alignment.[20]

While highly useful, the representation of comparative genomics information into

univariate scores or binary elements is limited in the amount of information it can convey about

the underlying multiple sequence alignment at a specific base. This limitation has become more

pronounced given the large number of species sequenced and incorporated into multi-species

alignments such as a 100-way alignment to the human genome.[21] Approaches have been

developed to associate constrained elements, regions, or individual bases with specific

branches in a phylogenetic tree.[22–28] While also useful, such directed approaches are biased to

only representing certain types of patterns present in the alignment. An alternative approach

used for comparative genomic based annotation learned patterns of different classes of

mutations between human and orangutan[29], but this approach was only applicable at a broad

region level and only incorporated information from one non-human genome.

Analogous to the many sequenced genomes available for comparative analysis, many

different datasets are available for annotating the genome based on epigenomic data.

Approaches that define 'chromatin states' based on combinatorial and spatial patterns in these

datasets have effectively summarized the information in them to provide *de novo* genome

annotations.[4,30–32] Inspired by the success of these approaches, here we develop a new

method, ConsHMM, that extends ChromHMM[31] to systematically annotate genomes into

'conservation states' at single nucleotide resolution. The conservation states assignments are

based on the combinatorial and spatial patterns of which species align to and which match a

reference genome at each nucleotide in a multiple species DNA sequence alignment.

ConsHMM takes a relatively unbiased modeling approach that does not explicitly assume a

specific phylogenetic relationship between species. The set of conservation patterns ConsHMM

can infer are thus flexible and determined directly from the DNA sequence alignment.

We applied ConsHMM to assign a conservation state to each nucleotide of the human

genome. These states are able to capture distinct enrichments for other genomic annotations

such as gene annotations, CpG islands, repeat families, chromatin states, and genetic variation.

These conservation state annotations are a resource for interpreting the genome and potential

disease-associated variation, which complement both existing conservation and epigenomic-

based annotations.

Material and Methods

Modeling conservation states with ConsHMM

ConsHMM takes as input an *N-way* multi-species sequence alignment to a designated

reference genome. For each base in the reference genome, *i*, ConsHMM encodes the multiple

species alignment into a vector, v_i, of length *N*-1. An element of the vector, $v_{i,j}$, corresponds to

one of three possible observation for species *j* at position *i*. The three possible observations are:

(1) the other species aligns with a non-indel nucleotide symbol present matching the reference

nucleotide, (2) the other species aligns with a non-indel nucleotide symbol present, but does not

match the reference nucleotide, or (3) the other species does not align with a non-indel

nucleotide symbol present.

ConsHMM assumes that these observations are generated from a multivariate HMM

where the emission parameters are assumed to be generated by a product of independent

multinomial random variables, corresponding to each species in the alignment. Formally, the

model is defined based on a fixed number of states *K*, and number of species in the multiple

sequence alignment *N*. For each state *k (k = 1,…,K)*, species *j* (j = 1,...,*N-1*) and possible

observation *m* (*m* = 1, 2, or 3 as described above), there is an emission parameter: $p_{k,j,m}$

corresponding to the probability in state *k* for species *j* of having observation *m*. For each

possible observation *m*, let $I_m(v_{i,j}) = 1$ if $v_{i,j} = m$, and 0 otherwise. Let $b_{t,u}$ be a parameter for the

probability of transitioning from state *t* to state *u*. Let $c \in C$ denote a chromosome, where C is

the set of all chromosomes in the reference genome of the multiple species alignment, and let

L_c be the number of bases on chromosome *c*. Let a_k (*k* = 1,...,*K*) be a parameter for the

probability of the first base on a chromosome being in state *k*. Let $s_c \in S_c$ be a hidden state

sequence on chromosome *c* and S_c be the set of all such possible state sequences. Let c_h

denote position *h* on chromosome *c*. Let s_{c_h} denote the hidden state at position c_h for state

sequence s_c.

We learn a setting of the model parameters that aims to optimize

$$P(v|a, b, p) = \prod_{c \in C} \sum_{s_c \in S_c} a_{s_{c_1}} \left(\prod_{i=2}^{L_c} b_{s_{c_{i-1}}, s_{c_i}} \right) \prod_{h=1}^{L_c} \prod_{j=1}^{N-1} \prod_{m=1}^{3} p_{s_{c_h}, j, m}^{I_m(v_{c_h, j})}$$

Once a model is learned, each nucleotide is assigned to the state with maximum

posterior probability. To conduct the model learning and state assignments, ConsHMM calls an

extended version of the ChromHMM[31] software originally designed to solve an analogous

problem of annotating the genome into chromatin states based on combinatorial and spatial

patterns of the presence of different chromatin marks. The modeling in ConsHMM differs from

the typical use of ChromHMM in three main respects: (1) the observation for each feature

comes from a three-way multinomial distribution as opposed to a Bernoulli distribution, (2) it is

applied at single nucleotide resolution opposed to 200-bp resolution, (3) it is applied with more

features than ChromHMM models have used in the past. (2) and (3) raise scalability issues in

terms of time and memory which we addressed in an updated version of ChromHMM (see

below).

To apply ChromHMM in the context of three-way multinomial distributions, ConsHMM

represents the three possible observations at position *i* for a species *j* with two binary variables,

y_{ij} and z_{ij}, corresponding to aligning and matching the reference genome respectively. y_{ij} has the

value of 1 if the other species aligns to the reference with a non-indel nucleotide and 0

otherwise. z_{ij} has the value of 1 if the other species has the same nucleotide as the reference

sequence and has a value of 0 if the other species has a different nucleotide present than the

reference. In the case in which $y_{ij}=0$, there is no nucleotide to compare to the reference and that

value of the z_{ij} variable is considered missing (encoded with a '2' for ChromHMM). If the value of

an observed variable is missing, ChromHMM excludes the Bernoulli random variable

corresponding to the observation from the emission distribution calculation at that position. For

each state k and species j, ChromHMM thus learns two parameters, $f_{k,j}$ and $g_{k,j}$. $f_{k,j}$ corresponds

to the probability that at a given position in state k, species j aligns to the reference genome with

a non-indel nucleotide that is $P(y_{i,j}=1|\ s_i=k)$. $g_{k,j}$ corresponds to the probability that at a given

position in state k, species j matches the reference genome conditioned on species j aligning

with a non-indel nucleotide that is $P(z_{i,j}=1|\ y_{i,j}=1$ and $s_i=k)$. This representation is equivalent to

the three-way multinomial distribution, $(p_{k,j,1},\ p_{k,j,2},\ p_{k,j,3})$ described above where $p_{k,j,1} = P(y_{i,j}=1,$

$z_{i,j}=1\ |\ s_i = k)$, $p_{k,j,2} = P(y_{i,j}=1,\ z_{i,j}=0\ |\ s_i = k)$, and $p_{k,j,3} = P(y_{ij}=0\ |\ s_i = k)$, since $p_{k,j,1} = f_{k,j} \times g_{k,j}$, $p_{k,j,2} =$

$f_{k,j} \times (1-g_{k,j})$, and $p_{k,j,3} = 1 - f_{k,j}$.

Multiple species sequence alignment choice

Our method and software can be applied to any multiple species sequence alignment

which is available in multiple alignment format (MAF) or which can be converted into this format.

For the results presented here we applied it to the 100-way Multiz vertebrate alignment with

human (hg19) as the reference genome.[21,33]

Scaling-up ConsHMM to single base resolution with hundreds of features

Since for our application ConsHMM needs to run ChromHMM at single base resolution ('-b 1'

flag) with 198 features after our binary encoding (2 for each non-human species in the 100-way

alignment), we had to address scalability issues in terms of both memory and time. To address

the memory issue we modified ChromHMM to support only loading in main memory input for

chromosomes it is actively processing, as previously ChromHMM would only support loading all

data into main memory upfront. This option can now be accessed in ChromHMM through the '-

lowmem' flag. To reduce the time required we used 12-parallel processors ('-p 12' flag) and we

trained on a different random subset of the human genome on each iteration of the Baum-welch

algorithm. We divided each chromosome into 200kb segments (with the exception of the last

segment of each chromosome which was less than this) in order to form a random subset of the

human genome. We modified ChromHMM to allow training for each iteration on a randomly

selected subset of 150 of these segments ('-n 150' flag) corresponding to 30MB per iteration.

We ran this for 200 iterations by adding the '-d -1' flag, which removed one of ChromHMM's

default stopping criterion based on computed likelihood change on the sampled data, since the

likelihood is now expected to both increase and decrease between iterations as different

sequences are sampled. These new options were included in version 1.13 of ChromHMM. The

unique code to ConsHMM is written in Python. The code of ConsHMM shared with ChromHMM

is written in Java and included with ConsHMM.

Generating genome-wide annotations

After learning a 100-state model, we used it to segment and annotate the genome at

base-pair resolution into one of the 100 conservation states. Each base in the human genome is

classified into the state with the highest posterior probability. ConsHMM does this by running the

MakeSegmentation command of ChromHMM. Due to computational constraints, the

segmentation could not be generated for entire chromosomes at once. Instead, we ran

MakeSegmentation on the same 200kb partitioning made for learning the model. We then

merged the resulting files together using ConsHMM's mergeSegmentation.py command with

slice size parameter set to 200,000 ('-s 200000' flag) and the number of states parameter set to

100 ('-n 100 flag').

Computing enrichments for external annotations

All overlap enrichments for external annotations were computed using the ChromHMM

OverlapEnrichment command. OverlapEnrichment computes enrichments for an external

annotation in each state assuming a uniform background distribution. Specifically the fold

enrichment of a state for an external annotation is

$$\frac{\% \ of \ external \ annotation \ bases \ falling \ in \ that \ state}{\% \ of \ genome \ falling \ in \ that \ state}$$

Positional enrichments of states relative to an anchor point from an external annotation

were computed using the ChromHMM NeighborhoodEnrichment command at single base

resolution ('-b 1' flag), single base spacing from the anchor point ('-s 1') and using the '-l' and '-r'

flags to specify the size of the region of interest around the anchor point. The '-lowmem' flag

was   also   used   for   computing   the   enrichments   for   OverlapEnrichment   and

NeighborhoodEnrichment.

External data sources for enrichment analyses

The external annotations of repeat elements were obtained from the UCSC Genome Browser

RepeatMasker track.[21,34] We generated an annotation for whether a base overlapped any repeat

element, as well as separate annotations for bases falling in each class and family of repeat

elements. The gene annotations were obtained from GENCODE v19 for hg19[35]. CpG Island

annotations were obtained from the UCSC genome browser. Annotations of SNPs with >=1%

minor allele frequency were obtained from the commonSNP147 track from the UCSC genome

browser, which is based on dbSNP build 147. GWAS catalog variants were obtained from the

NHGRI-EBI Catalog, accessed on Dec 5, 2016.[36] For annotations of DHS from the Roadmap

Epigenomics Consortium, we used Macs2 narrowPeak calls.[5] The Fetal Brain and HepG2 DHS

used were of epigenome sample E082 and E118 respectively. For the median non-exonic DHS

enrichments and ranking of states in the heritability partitioning analysis we used narrowPeak

calls from the ENCODE consortium.[3]

209   PhyloP and PhastCons scores and constrained element calls were obtained from the

UCSC genome browser. Assembly gap annotations were obtained from the Gap track from the

UCSC genome browser. The context-dependent tolerance score (CDTS) used was that based

on a cohort of 7784 unrelated individuals, following the analyses in Iulio et al.[37], which focused

on this version of the score. The CDTS and variants from this cohort were both lifted from hg38

to hg19 using liftOver tool from the UCSC Genome Browser.[21]

Choice of number of states

217   We learned models with each number of states between 2 and 100 states. We set 100

as the maximum number of states we would consider for computational tractability and

maintaining a manageable number of states for analysis. The choice of a maximum of 100 also

corresponds to the number of species used and allows for the possibility of each state to cover

1% of the genome. We analyzed the Bayesian Information Criterion (BIC) for models with each

number of states between 2 and 100, and found that the BIC generally decreases as the

number of states increases in the range considered (**Supplementary Fig. 1**). The BIC was

calculated using the BIC_HMM function from the HMMpa R package.[38] Analyzing the 100-state

model's internal confidence estimate of its state assignments also supported a larger number of

states. Specifically, for each state in the 100-state model we computed the average posterior

probability of that state at each base in the genome assigned to it, and confirmed consistently

high average posterior probability values in the range [0.92,1.00] with a median of 0.97

(**Supplementary Fig. 2**). The posterior probabilities were computed by running the

MakeSegmentation command in ChromHMM with the '-printposterior' flag. We also investigated

if additional states in models with larger number of states were biologically relevant. Specifically,

we computed enrichments for various external annotations for models with each number of

states between 2 and 100 to determine if biologically relevant enrichments were only robustly

observed in models with more than a certain number of states. In the case of CpG islands, we

observed that only models with at least 87 states consistently obtained >15 fold enrichment and

only models with at least 95 states consistently obtained >30 fold enrichment (**Supplementary**

Fig. 3). We saw a similar pattern of increasing enrichments for annotated TSSs for models with

large number of states. We therefore decided to analyze the largest model, 100 states, that we

were considering. We note that annotations based on chromatin states used fewer number of

states, but were also defined on fewer features at a coarser resolution and had a less uniform

genome coverage.[4,30,39]

State clustering

We clustered the states based on the correlation of vectors containing the values $f_{k,j}$ and

$f_{k,j} \times g_{k,j}$ for each species j defined above. State clustering was performed using the hclust

hierarchical clustering function from the cba R package.[40] The leaves of the resulting

hierarchical tree were ordered according to the optimal leaf ordering algorithm[41] implemented in

the order.optimal R function from the cba package. We then cut the tree such that the 8 major

groups of states were designated. The full tree is provided in **Supplementary Fig. 4**.

Gene Ontology enrichments

For each state and each protein coding gene based on GENCODE we computed the

number of bases in that state that are within +/- 2kb of the gene's TSS. In the case of genes

with multiple annotated TSSs, we used the outermost TSS. We then created a ranking of genes

for every state by sorting the genes in descending order of this number of bases. For each state

we then created a set of 969 genes that represent the top 5% of genes in the state among the

19,397 genes we considered. We performed a Gene Ontology enrichment analysis (ontology

and annotations files from Nov. 24[th], 2016) for the top genes in each state using the STEM

software in batch mode with default options and the set of all genes considered as

background.[42] STEM computed an uncorrected p-value based on the hypergeometric

distribution for each term displayed in the figures summarizing the analysis. STEM also reported

corrected p-values for testing multiple GO terms for a single state based on randomization to

three significant digits, which was less than 0.001 for all p-values mentioned in the main text.

Transcription factor binding site motif enrichments

We computed the enrichment of the conservation states within 15 bases upstream and

downstream of the center point of the POU5F1 and STAT known transcription factor-binding site

motifs. The enrichment was computed relative to the background regions of the genome that

were used to identify the motifs, which excluded repeat elements, coding sequence, and 3'

UTRs.[43] The *known1* version of the motifs was used for both motifs.

Precision recall analysis for recovery of gene annotations and DHS

We randomly split the 200kb genome segments used for training the model and

segmentation into two halves corresponding to training and testing data. For each target set in

the precision-recall analyses we ordered the ConsHMM states in decreasing order of their

enrichment for the target among the training set bases. We then used that ordering to iteratively

add the testing set bases in each state to form cumulative sets of bases predicted to be of the

target set and computed the precision and recall for them. For each constraint score we

computed the precision-recall curve for predicting the target set in the test data using two

methods. For the first method we directly ordered bases in descending order of their assigned

score. For the second method, we split the sorted scores into 400 bins such that each bin

contains on average 0.25% of the genome, which was the size of the smallest state of the

ConsHMM model (0.25% of the genome in state 100). Specifically, we assigned all bases in the

genome where the score was not defined to one bin and then divided the remaining bases

uniformly among the 399 other bins based on their score. In some cases score increments were

at the boundary between two bins at their provided floating-point precision, or overlapped

multiple bins. In these cases we uniformly split the target bases assigned to that score

increment into multiple bins proportionally to the overall percentage of the score increment

falling in each bin. We then treated the 400 bins as 400 states and followed the same procedure

described for the ConsHMM states. We also computed the precision and recall of bases in each

constrained element set for predicting the target set on the testing data.

Clustering of cell-type specific DNase I hypersensitive site enrichments

For the clustering of DHS analysis, we first computed the enrichments of all conservation

states for DHS for 53 samples processed by the Roadmap Epigenomics consortium[5], of which

16 were originally generated by the ENCODE project consortium.[3] We then selected the subset

of states that had a fold enrichment of at least 2 in at least one sample, leading to a subset of 21

conservation states. To more directly focus on each state's relative enrichments across

samples, we $\log_2$ transformed each enrichment value, and then normalized the enrichments for

each state by subtracting the mean enrichment across samples and dividing by the standard

deviation. We then hierarchically clustered the states based on the correlation of their

enrichments across samples and hierarchically clustered the samples based on their

correlations across states using the pheatmap R package.[44]

Heritability partitioning analysis

The heritability partitioning was performed using the LD-score regression ldsc software.[15] We partitioned the PhastCons constrained elements into two halves based on a ranking of the

conservation states. We focused on the PhastCons constrained elements for this analysis, since

it was the only element set defined on the same alignments as our conservation states. We

focused on halves since the LD-score regression estimates can be unstable for annotations

covering too small of a percentage of the genome.[15] To determine the two halves we ranked the

conservation states in descending order of median fold-enrichment of non-exonic bases for

DHS from 125 experiments from the University of Washington ENCODE group.[3] We then

divided bases in PhastCons elements between the top 7 ranked states (1-5, 8 and 28), which

contain 51.9% of bases in PhastCons elements, and the bottom 93 states, which contain the

other 48.1% of bases in PhastCons elements. We applied ldsc to these two sets for 8 traits (age

at menarche, BMI, coronary artery disease, educational attainment, height, LDL levels,

schizophrenia and smoking behavior), all of which were previously considered in heritability

partitioning analysis.[15] We followed the procedure for partitioning heritability as done in

Finucane et al.[15], including using a baseline annotation set and 500 base-pair windows around

annotations to dampen the artificial inflation of heritability in neighboring regions caused by

linkage disequilibrium. The baseline annotation set contains a range of annotations including

DHS. For our analysis, we first removed the constrained element set already included in the

baseline annotation set, then added our two halves of PhastCons elements and finally ran the

ldsc software on the full set of annotations.

Results

Annotating the human genome into conservation states

We developed a novel approach, ConsHMM, to annotate a genome into different

conservation states based on a multiple species DNA sequence alignment (**Fig. 1a, Methods**).

We model the combinatorial patterns within the alignment of which species align to and which

match a reference genome, for which we used the human genome. Specifically, at each

nucleotide in the human genome we encode one of three possible observations for each other

species in the alignment: (1) aligns with a nucleotide present that is the same as the human

reference genome, (2) aligns with a nucleotide present that is different than the human

reference genome, or (3) does not have a nucleotide present in the alignment for that position.

We further model these observations as being generated from a multivariate hidden Markov

model (HMM), which probabilistically captures both the combinatorial patterns in the

observations and their spatial context. Specifically, we assume that in each state the probability

of observing a specific combination of observations is determined by a product of independent

multinomial random variables. The parameters to the distributions of these multinomial random

variables will differ between states and are learned from the data. After the model is learned,

each nucleotide in the human genome is assigned to the state that had the maximum posterior

probability of generating the observations.

ConsHMM builds on ChromHMM[31], which has previously been applied to annotate

genomes based on epigenomic data at 200-bp resolution[30], to now annotate genomes at single

nucleotide resolution based on a multiple species DNA sequence alignment (**Methods**). We

applied ConsHMM to a 100-way Multiz vertebrate alignment with the human genome and

focused our analysis here on a model learned using 100 states in order to balance recovery of

additional biological features and model tractability (**Fig. 2**, **Supplementary Fig. 1-8**)**,**

Supplementary Tables 1,2, Methods). We illustrate the ConsHMM conservation state

annotations at two different loci showing that different bases that are associated with calls of

evolutionary constraint from existing approaches can have very different underlying alignment

patterns and conservation state assignments (**Fig. 1b, Supplementary Fig. 9**). Conservation

state annotations genome-wide are available online (**Web Resources**)**.**

 Major groups of conservation states

We hierarchically clustered the conservation states based on their align and match

probabilities, and then cut the resulting dendrogram to reveal eight notable groups of states or

distinct individual states (**Fig. 2a**, **Supplementary Fig. 4, Supplementary Table 3, Methods**).

The first of these subsets of states was a single state (State 1; AM_allVert) that showed high

align and match probabilities through essentially all vertebrate species considered. The second

subset showed relatively high align and match probabilities for all mammals and some non-

mammalian vertebrates (States 2-4; AM_nonMam). The third subset showed relatively high

align and match probabilities for most if not all mammals, but not non-mammalian vertebrates

(States 5-22; AM_Mam). The fourth subset showed high align probabilities for many mammalian

species, but had low align probabilities for notable species such as mouse and rat for many of

the states in the group (States 23-46; AM_SMam). The combination of the absence of mouse

and rat alignments with the presence of mammals that are assumed to have diverged earlier is

consistent with the previously observed increased substitution rates for mouse and rat.[7] The fifth

subset showed high align probabilities for many mammalian species, but did not show high

match probabilities (States 47-63; A_SMam). The sixth subset showed high align probabilities

for most primates, but not for other species (States 64-89; AM_Prim). The seventh subset

showed high align probabilities for at most a subset of primates (States 90-99; AM_SPrim). The

final subset was a single state (State 100; artifact) that showed a noteworthy pattern of high

align and match probabilities for most primates and non-mammalian vertebrates, but low

probabilities for non-primate mammals, consistent with a previous observation that inclusion of

non-mammalian vertebrates can be associated with increased presence of suspiciously aligned

regions.[45]

Conservation states exhibit distinct patterns of positional enrichments relative to gene

annotations and regulatory motif instances

The conservation states showed strong and distinct positional enrichments relative to

GENCODE[35] annotated gene features including transcription start sites (TSS), transcription end

sites (TES), exon start sites, and exon end sites for both protein coding genes and

pseudogenes (**Fig. 3a-d**, **Supplementary Fig. 10**). Notable positional enrichments were also

seen for regulatory motifs instances (**Fig. 3e,f**). Relative to starts of exons of protein coding

genes seven of the states (States 1-4, 7, 28, and 54) had 13 fold or greater enrichment for some

position within 20 base pairs of exon starts, both when considering all such exons and subsets

of exons in specific coding phases (**Fig. 3a, Supplementary Fig. 10a-c**). All of these states

showed alignability for some non-mammalian vertebrates in addition to most mammals. Within

exons we saw the strongest enrichment for States 1-4 and 54, and among these state 1 showed

the strongest enrichment as expected given its high match probabilities through all vertebrates

(**Figs. 2b, 3a-b, Supplementary Fig. 10a-e**). Interestingly, state 1 showed very strong

enrichment (>80 fold) in the two nucleotides immediately upstream of the exon start with the

third upstream nucleotide also having high enrichment (46 fold) (**Supplementary Fig. 10c**).

These three nucleotide positions correspond to the positions of the canonical 3' splice site

sequence that is highly conserved throughout vertebrates.[46] At the ends of exons of protein

coding genes (**Fig. 3b**), state 1 maintained a >40 fold enrichment for six nucleotides past the

end of coding sequence corresponding to positions of the known canonical 5' splice site

sequence.[46] Downstream of the start of protein-coding exons, the enrichment profile for state 1

showed a 3-bp oscillation period, with a dip of enrichment at each 3rd base corresponding to

codon wobble positions. In contrast, states 3 and 54, which were both associated with high align

probabilities through many vertebrates and lower match probabilities, showed the inverse

oscillation pattern to state 1 (**Fig. 3a, Supplementary Fig. 10a-c**)**.**

Relative to TSS of protein coding genes, state 28 had the strongest enrichment reaching

a maximum enrichment 30 fold at the TSS (**Fig. 3c**). State 28 was associated with moderate

align and match probabilities for almost all the species present in the alignment. Consistent with

its enrichment for TSSs state 28 also had the greatest enrichment for CpG islands (32 fold).

However, state 28 also showed a 20 fold enrichment of CpG islands >2kb away from any TSS

of protein coding genes and a 10 fold enrichment for TSS of protein coding genes >2kb away

from a CpG island, suggesting the possibility that both of these features are making a partially

independent contribution to the association, or the presence of additional unannotated TSSs

that are associated with CpG islands.[47] Relative to TES of protein coding gene we saw the

enrichment peak for state 2 at almost 12 fold (**Supplementary Fig. 10f**), which had high align

and match probabilities for almost all vertebrates except for fish.

Relative to pseudogene exon starts and ends, states 100 and 82, both associated with

alignability to distal vertebrates without many mammals closer to human, had strong

enrichments peaking at greater than 100 and 38 fold respectively (**Supplementary Fig. 10g,h**).

These two states also showed the greatest enrichment relative to TSSs of pseudogenes

peaking at 184 and 68 fold for states 100 or 82 respectively (**Fig. 3d**) and for TESs of

pseudogenes peaking at 199 and 61 fold respectively (**Supplementary Fig. 10i**).

Relative to instances of regulatory motifs different conservation states showed single

nucleotide enrichment variation, often associated with variation in the amount of information in

the positional-weight matrix (**Fig. 3e-f** and **Methods**).[43] For example, in the case of the POU5F1

and STAT motifs we saw state 2 from the AM_nonMam group and state 5 from the AM_Mam

group respectively reach 1.8 fold enrichments but have lower enrichments (1.4-1.5) at some

nucleotides with lower information content. For the STAT motif, states 55-57, associated with

high align probabilities for most mammals, but high match probabilities only for a few primates,

enrichments peaked at the CG dinucleotide in the center of the motif consistent with their

genome-wide enrichments for CG dinucleotides (**Fig. 3e, Supplementary Fig. 11**).

Enrichment of conservation states for different gene classes

The previous analyses demonstrated that different conservation states have distinct

enrichments in promoter regions of genes. We next investigated whether different conservation

states also exhibit distinct enrichments for different classes of genes after controlling for the

state's relative preference for promoter regions. Specifically, for each state we determined the

5% of genes with the greatest presence of the state in its promoter region and evaluated Gene

Ontology (GO) enrichments for those genes, revealing distinct enrichment patterns (**Fig. 4b,**

Supplementary Fig. 12, Methods). For example, even among states 1-3, all of which had high

alignability through at least birds and matching through mammals, we observed substantial

differences in their gene preferences. Out of these three states, state 1 (the AM_allVert group)

was the only one enriched for nucleosomes ($p<10^{-41}$; 10.5 fold), while state 3, which had high

matching only through mammals, was the only one with a significant enrichment for a set of

genes related to sensory perception of smell ($p<10^{-300}$; 15.5 fold). State 2, which had high align

and match probabilities through all vertebrates except fish, was the state most enriched for

cellular developmental processes ($p<10^{-30}$; 1.8 fold), which did not show enrichment in state 3.

We also observed notable enrichments for states with overall lower align or match probabilities.

For example, state 89, associated with high alignability and low matching in primates as well as

some alignability and low matching in non-primate mammals, was the state most enriched for

antigen binding ($p<10^{-14}$; 6.7 fold). This is consistent with antigen binding being associated with

many species, but fast evolving.[48]

Enrichments for repeat elements in conservation states

The conservation states showed a wide range of enrichments and depletions (from 2

fold enrichment to 133 fold depletion) for bases overlapping any repeat element (**Fig. 2b, Fig.**

4d, Supplementary Fig. 13).[21,34] Different states had distinct patterns of enrichments for

different repeat classes and families, even though in some cases the difference in state

parameters was subtle. For instance, among states in the AM_Prim group, which primarily

differed in terms of the specific combinations of primates with high align and match probabilities,

we found distinct enrichments. Notably, four different states from the group AM_Prim, 74, 86,

76, and 77, showed maximal enrichments for the DNA, LINE, LTR, and SINE repeat classes

respectively (**Fig. 4d**). State 74, which is characterized by high align and match probabilities for

all primates, had an enrichment of 5.6 fold for DNA repeats, while the enrichment for the other

three classes were between 1.0 and 1.8 fold. On the other hand, state 86, which lacked

alignability of a subset of primates, had a 3.0 fold enrichment for LINE repeats, while the

enrichment for the other classes were between 0.6 and 1.6 fold. States 76 and 77 had 3.3 and

4.5 fold enrichments for LTR and SINE respectively compared to 1.1 and 2.1 fold for SINE and

LTR respectively. State 76 and state 77 both had high align probabilities through primates up to

and including squirrel monkey, with the exception that State 77 lacked alignability to gorilla.

Despite these subtle differences in the alignment probabilities, these states had substantial

differences in their repeat enrichment profiles.

Relationship of conservation states to chromatin states

We compared our conservation states to annotations of the genome based on a 25-

chromatin state imputation based model defined on 127 samples of diverse cell and tissue

types[5,39] (**Fig**. **4a, Supplementary Fig. 14**). For each conservation state we determined the

median enrichment of each chromatin state across the 127 samples. Eleven different

conservations states were maximally enriched for at least one of the 25-chromatin states.

Conservation state 28 showed the greatest enrichment for any chromatin state, with a 35 fold

enrichment for a chromatin state associated with active promoters, and was maximally enriched

for four other promoter associated states. Conservation state 1 was maximally enriched for five

chromatin states all associated with transcribed and exonic regions[39] (3.8-8.7 fold), which is

consistent with this conservation state being most enriched for exons. Conservation state 2 had

the maximal enrichment for five enhancer associated states (3.1-4.7 fold), while conservation

state 5 had high enrichments for these states and also had the greatest enrichment of any

conservation state for a state primarily associated with just DNase I hypersensitivity (2.5 fold).

These and other distinct enrichments of the conservation states for the different chromatin

states highlight that conservation states are able to capture multi-dimensional information in the

genome.

Conservation states capture enrichment patterns of DNase I hypersensitive sites across cell and

tissue types

The previous analysis demonstrated that conservation states can exhibit different

enrichment patterns for different chromatin states. We next investigated whether different

conservation states also capture distinct enrichment patterns for a chromatin mark across cell

and tissue types. For this we analyzed DNase I hypersensitive sites (DHS) from 53 of the 127

samples considered above for which maps of experimentally observed DHS were available from

the Roadmap Epigenomics Consortium.[5] We focused on the 21 conservation states that

exhibited at least 2 fold enrichment in at least one sample (**Fig. 4c**). We then row normalized

the enrichments in order to focus on the relative enrichment patterns across cell and tissue

types (**Methods**). Hierarchical clustering of the enrichment patterns revealed two major clusters

of states (**Fig. 4c**). One of these clusters contained 14 of the 21 states and was associated with

strong enrichments for fetal related samples. Ten of the states in this cluster have maximum

enrichment for a fetal sample, while the remaining four states have maximum enrichment for the

cell type Human Umbilical Vein Cells (HUVEC). The second major cluster consisted of seven

states, all of which were enriched for CpG islands (**Fig. 2b**, **Supplementary Fig. 8**). The

samples that showed the greatest enrichment in states in these clusters also had the greatest

enrichment of CpG islands (**Fig. 4c**), but were biologically diverse in terms of the type of cell or

tissue and could potentially reflect technical experimental differences.

Relationship of conservation states to constraint based annotations

We next investigated the relationship of our conservation state annotations with calls

and univariate scores of evolutionary constraint. Specifically, we considered constrained

element sets based on four methods (GERP++, SiPhy-omega, SiPhy-pi, and PhastCons) and

constraint scores based on three methods (GERP++, PhastCons, and PhyloP) publicly available

for hg19 and also defined on Multiz alignments (**Fig. 2b**, **Supplementary Fig. 15**). The

PhastCons and PhyloP scores and elements we compared to were defined on the same 100-

way vertebrate alignment. The available GERP++, SiPhy-omega, and SiPhy-pi score and

elements were derived from different versions of Multiz alignments and only considered

mammals.

We consistently found conservation states 1-5 to be highly enriched (>9 fold) for all

constrained element sets (**Fig. 2b**, **Supplementary Fig. 15a**). These states were also among

the top six states in terms of mean score for constraint scores considered (**Supplementary Fig.**

15b). Consistent with this, states 1-5 were the states that had the highest average matching

probability across mammals. Two other states exhibited at least 6 fold enrichment for at least

one constrained element set: states 54 and 100. State 100, associated with putative artifacts,

showed high enrichments for PhastCons elements (15 fold) and high average scores for

PhastCons and PhyloP. This is consistent with this state having high aligning and matching

probabilities primarily in non-mammalian vertebrates and these elements and scores being

defined using such species. State 54 was consistently enriched for all the constrained elements

(4-7 fold), but did not show high mean base-wise scores. The difference of high enrichment in

constrained elements but not base-wise scores is consistent with state 54 having high

alignability through most vertebrates, but low matching outside primates. More generally, we

found that constrained element calls did not have the resolution to exhibit biologically relevant

single nucleotide variation in enrichments around regulatory motifs and exon start and ends as

we saw with our conservation state annotations, with the exception of those from PhastCons

(**Fig. 3, Supplementary Fig. 16**).

The objective of our conservation state annotations is different than that of binary calls

and univariate scores of evolutionary constraint, which have a more specific and complementary

goal. However, to better understand their relative biologically relevant information we compared

their ability to recover annotated starts and ends of exons and TSS and TES of genes

separately for protein coding and pseudogenes (**Fig. 5a-c**, **Supplementary Fig. 17**). In almost

all cases the conservation states had greater information available for recovering annotated

gene features. The only exceptions were that PhyloP scores could achieve higher precision at

low recall levels for protein coding exon starts and ends and that SiPhy-pi elements had slightly

higher precision for TSS of protein coding genes at their one recall point.

We also compared the ability of conservation states to recover bases covered by DHS

sites in specific cell types both genome-wide and restricted to non-exonic bases

(**Supplementary Fig. 18**). For these analyses we generally found that at the same recall level

the conservation states could identify bases in a DHS at greater precision than constraint

scores. Compared to constrained elements the relative ability depended both on the specific

constrained element set being compared and the specific DHS experiment. Importantly, the

information about DHS in the conservation states was complementary to that in the constrained

element sets as evidenced by large variation in DHS enrichments of bases within constrained

elements depending on their conservation state (**Fig. 5d, Supplementary Fig. 19**). For

example, the enrichment of bases in PhastCons constrained elements for Fetal Brain DHS in

non-exonic regions ranged from 0.3 to 10.1 fold depending on the conservation state.

Additionally, we saw cases where certain states had greater enrichments for DHS for their

bases not in a constrained element compared to bases in a constrained element in other states.

On the other hand, constrained element calls offered additional information as we observed that

in most cases, for a given conservation state, bases that were in a constrained element call had

greater enrichment for DHS than those that were not.

We also analyzed the enrichments of our conservation states for previously defined

nine-subsets of PhastCons constrained non-exonic elements (CNEEs) based on a directed

phylogenetic approach that assigned each element to a phylogenetic branch point of origin

(**Supplementary Fig. 20a**).[22] This demonstrated in some cases the heterogeneous nature of

the resulting assignments when relying on directed phylogenetic partitioning approaches. For

example bases in elements assigned to originating at the branch point of the Tetrapod clade

showed a 37 fold enrichment for state 2, as would be expected since state 2 is associated with

aligning and matching through all vertebrates except fish, but an even greater enrichment (51

fold) for state 100, associated with putative artifacts. A similar pattern of enrichments was

observed when considering only the CNEEs overlapping a PhastCons element called on the

same alignment as the conservation states (**Supplementary Fig. 20b**).

Enrichment of conservation states for human genetic variation

Previous analyses have found a depletion of human genetic variation in evolutionarily

constrained elements.[7] Consistent with that, the greatest depletion (3.3 fold depletion) of

common SNPs from dbSNP is in state 1, the state most enriched for constrained elements.

Interestingly, six states, A_SMam states 55-57 and AM_Prim states 87-89, had enrichments in

the range 5 to 8 fold for common SNPs. These were also the six states with greatest enrichment

of CG dinucleotides (**Supplementary Fig. 11**). These six states have in common that they show

high align probabilities for most primates, but low match probabilities for some of those same

primates. These states are thus associated with substantial variation both among primates and

among humans. We observed similar patterns of enrichment for variants identified from whole

genome sequencing (WGS) of a cohort of 7784 unrelated individuals[37], with the levels of state

enrichments and depletions increasing with the minor allele frequency (**Supplementary Fig.**

21a).

When analyzing the enrichment of GWAS catalog variants[36] relative to the background

of common SNPs we saw opposite enrichment patterns for these states (**Fig. 6a,b**). For

example, relative to this background, state 1 was most enriched for GWAS catalog variants,

which is consistent with previous observations of constrained elements enriching for GWAS

variants.[7] On the other hand, states 55-57 and 87-89 showed the greatest depletion. These

results suggest that common variants are less likely to be phenotypically significant if they fall in

conservation states most enriched for common genetic variation.

Conservation states capture similarities and differences between context-dependent tolerance

score and inter-species constraint in humans

A recent study defined a context-dependent tolerance score (CDTS) based on the local

depletion of genetic variations in the same WGS data of 7794 unrelated individuals relative to

expectation determined by the DNA sequence context.[37] Bases prioritized by the score were

reported to identify regulatory elements of the genome while having limited overlap with bases

prioritized by GERP++. We used our more detailed conservation state annotations to better

understand the relationship between bases prioritized by CDTS and those falling in GERP++

defined constrained elements (**Figure 6c, Supplementary Fig. 15a, 21b**). Bases in GERP++

elements and top 1% CDTS bases both showed the highest enrichments for state 1 (14.3 fold

and 13.3 fold respectively), which was the state strongly enriched for protein coding exons. On

the other hand states 2 and 28 highlight the differences in bases prioritized by these two sets,

with state 2 preferentially enriched for GERP++ bases relative to top 1% CDTS (12.6 fold vs. 2.4

fold) and state 28 preferentially enriched for top 1 % CDTS (9.6 fold vs. 3.0 fold). State 28 was

the state strongly associated with annotated promoters of protein coding genes (**Fig 3c**) and

CpG islands (32.1 fold) while state 2 was the state most strongly enriched for candidate

614 enhancers defined by the chromatin states (**Figure 4a**), but with limited enrichment for CpG

615 Islands (1.5 fold) (**Figure 6c**). These and other state enrichments highlight specific cross-

616 species conservation patterns that are unique to each approach to capturing constraint, as well

617 as shared.

619 *Constrained element enrichment for partitioned heritability of complex traits depends on*

620 *conservation state*

621   Previous analyses have suggested a strong enrichment of constrained elements and

622 DHS for phenotype heritability.[15,49] As we saw large differences in DHS enrichments of

623 constrained elements depending on the conservation state, we investigated the extent to which

624 constrained elements in conservation states most enriched for DHS enriched for phenotype

625 heritability compared to the remaining states. Specifically, we ranked the conservation states in

626 descending order of their median enrichment for DHS from a compendium of 125 experiments

627 from the ENCODE consortium, within the non-exonic portion of the state (**Fig. 2b, Methods**).[3]

628 We then partitioned bases in PhastCons constrained elements into two almost equal size sets

629 based on whether they overlapped one of the top seven ranked conservation states (states 1-5,

630 8, 28) or not. We then computed the heritability for these two sets for eight phenotypes in the

631 context of a set of baseline annotations that include DHS annotations (**Methods**).[15] For seven of

632 the phenotypes, we found that bases in constrained element overlapping the top seven states

633 had greater enrichment than those in the remaining 93 states, often substantially so (**Fig. 6d**).

634 These results suggest additional value in the conservation state annotations for isolating more

635 likely disease relevant variants.

637 **Discussion**

638   We presented a new framework for genome annotation based on comparative genomics

639 sequence data. Our approach learns a set of conservation states *de novo* using a multivariate

HMM based on the combinatorial and spatial patterns of which species align and match a

reference genome in a multi-species DNA sequence alignment. We applied this approach to

annotate the human genome at single nucleotide resolution into one of 100 conservation states.

Conservation state annotations exhibited substantial enrichments for a wide range of other

genomic annotations that were not provided to the model in training, thus supporting their

biological significance. Specific conservation states exhibited strong enrichments for various

gene annotations including exons and TSS and TES of genes. Conservation states showed

differential enrichment patterns for various classes of genes and DHS from multiple cell types,

even though they were defined independently of any functional genomics data. Specific

conservation states exhibited enrichments for common human variants, while a different set of

states exhibited enrichments for variants identified by GWAS relative to common variation.

ConsHMM provides a novel approach to comparative genomics based genome

annotation. ConsHMM differs from other comparative genomic based annotation approaches in

that it takes an unsupervised approach that does not explicitly use a phylogenetic tree, except to

the extent to which a phylogenetic tree was used to generate the input multi-species sequence

alignment. This leads to relatively unbiased, simple and interpretable models. However, many

state patterns discovered are consistent with expected observations from commonly assumed

phylogenetic relationships of the species. While states' parameters often decreased with

divergence time from human, there were a number of exceptions. Some of these exceptions

corresponded to missing specific sub-clades of species, particularly those with long branch

lengths. For example, a number of states were not represented by mouse and rat while being

represented by more distally diverged mammals. Other exceptions isolated putative artifacts in

the alignments that might otherwise confound analyses, as we saw for two states heavily

enriched for pseudogenes.

Our conservation state annotation is complementary to existing binary calls and scores

of evolutionary constraint based on phylogenetic modeling. Both locations called as constrained

and those called as non-constrained are heterogeneous in their assigned conservation state.

Our annotations thus provide additional descriptive information about the conservation patterns

at each base. In terms of information for predicting external annotations, we found that in many

cases the conservation states had greater information than constraint scores or elements. For

other cases, such as DHS, the relative information depended on the constrained element set or

score being compared. Importantly, we observed that DHS information provided by the states

was complementary to information in the constrained element calls. We also showed how our

conservation state annotations can clarify the relationship between traditional interspecies

constraint annotations and the recently proposed CDTS measure, as we saw states uniquely

strongly enriched in the set of bases prioritized by each approach. Additionally, we observed

that bases in constrained elements showed substantially different enrichments for phenotype-

associated heritability, depending on their conservation state.

The conservation states are both inspired by, and provide complementary information to,

existing chromatin state annotation approaches. While the states from the two approaches are

based on very different data and have fundamental differences, they also exhibited substantial

cross-enrichments. In general, conservation states have the advantage of providing information

at single nucleotide resolution, which we demonstrated by showing enrichments patterns in and

around coding exons and regulatory motifs. Conservation states can also provide information

about bases in the genome even if the relevant cell type has not been experimentally profiled,

while chromatin states have the advantage of directly providing cell type specific information.

We expect many applications for the methodology and annotations we have presented

here. While we applied ConsHMM here to one multiple species alignment, a 100-way Multiz

human alignment, the methodology is general, and thus can be readily applied to alignments to

other species or alignments generated by other methods.[26] The annotations we produced serve

as a resource to directly interpret other genomic datasets. They can also be integrated with

complementary functional genomics or comparative information in methods that aim to better

prioritize disease relevant variants.[16–19,50,51] This work represents a step in the direction of

improving whole genome annotations, which will continue to be of increasing importance

towards understanding health and disease as the availability of whole genome sequencing data

increases.

Supplemental Data

Supplemental Data include twenty-one figures and three tables.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgements

We thank Ewan Birney and members of the Ernst lab for useful discussions. We

acknowledge funding from US National Institutes of Health grants DP1DA044371,

R01ES024995, U01HG007912 and U01MH105578 (J.E.), and T32CA201160 (A.S.), US

National Science Foundation CAREER Award #1254200 (J.E.), a Kure-IT award and an Alfred

P. Sloan Fellowship (J.E.). The funding bodies had no role in the development of the method or

the analysis of the results.

Web Resources

Multiz 100-way alignment to hg19 reference:

http://hgdownload.soe.ucsc.edu/goldenPath/hg19/multiz100way/

ConsHMM v1.0 software and ConsHMM state annotations:

https://github.com/ernstlab/ConsHMM

GENCODE v19:

https://www.gencodegenes.org/releases/19.html

25-state chromatin state annotations:

http://compbio.mit.edu/roadmap

GWAS catalog variants:

https://www.ebi.ac.uk/gwas/

CDTS:

http://www.hli-opendata.com/noncoding/coord_CDTS_percentile_N7794unrelated.txt.gz

SNVs from **Supplementary Fig. 21a**:

http://www.hli-opendata.com/noncoding/SNVusedForCDTScomputation_N7794unrelate

d_allelicFrequency0.001truncated.txt.gz

Roadmap DHS:

http://egg2.wustl.edu/roadmap/data/byFileType/peaks/consolidated/narrowPeak/

ENCODE DHS:

http://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeUwDnase/

GERP++ scores and constrained element calls:

http://mendel.stanford.edu/SidowLab/downloads/gerp/

SiPhy-omega and SiPhy-pi constrained element calls (hg19 liftOver):

https://www.broadinstitute.org/mammals-models/29-mammals-project-supplementary-

info

CNEEs from Lowe et al.[22]:

http://www.stanford.edu/~lowec/data/threePeriods/hg19cnee.bed.gz

Motif instances and background:

http://compbio.mit.edu/encode-motifs/

References

1. Hindorff, L.A., Sethupathy, P., Junkins, H.A., Ramos, E.M., Mehta, J.P., Collins, F.S., and
Manolio, T.A. (2009). Potential etiologic and functional implications of genome-wide association
loci for human diseases and traits. Proc. Natl. Acad. Sci. U. S. A. *106*, 9362–9367.

2. Ward, L.D., and Kellis, M. (2012). Interpreting non-coding variation in complex disease
genetics. Nat. Biotechnol. *30*, 1095–1106.

3. ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the
human genome. Nature *489*, 57–74.

4. Ernst, J., Kheradpour, P., Mikkelsen, T.S., Shoresh, N., Ward, L.D., Epstein, C.B., Zhang, X.,
Wang, L., Issner, R., Coyne, M., et al. (2011). Mapping and analysis of chromatin state
dynamics in nine human cell types. Nature *473*, 43–49.

5. Roadmap Epigenomics Consortium, Kundaje, A., Meuleman, W., Ernst, J., Bilenky, M., Yen,
A., Heravi-Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J., et al. (2015). Integrative analysis
of 111 reference human epigenomes. Nature *518*, 317–330.

6. Kellis, M., Wold, B., Snyder, M.P., Bernstein, B.E., Kundaje, A., Marinov, G.K., Ward, L.D.,
Birney, E., Crawford, G.E., Dekker, J., et al. (2014). Defining functional DNA elements in the
human genome. Proc. Natl. Acad. Sci. *111*, 6131–6138.

7. Lindblad-Toh, K., Garber, M., Zuk, O., Lin, M.F., Parker, B.J., Washietl, S., Kheradpour, P.,
Ernst, J., Jordan, G., Mauceli, E., et al. (2011). A high-resolution map of human evolutionary
constraint using 29 mammals. Nature *478*, 476–482.

8. Pollard, K.S., Hubisz, M.J., Rosenbloom, K.R., and Siepel, A. (2010). Detection of nonneutral
substitution rates on mammalian phylogenies. Genome Res. *20*, 110–121.

9. Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K., Clawson,
H., Spieth, J., Hillier, L.W., Richards, S., et al. (2005). Evolutionarily conserved elements in
vertebrate, insect, worm, and yeast genomes. Genome Res. *15*, 1034–1050.

10. Cooper, G.M., and Shendure, J. (2011). Needles in stacks of needles: finding disease-
causal variants in a wealth of genomic data. Nat. Rev. Genet. *12*, 628–640.

11. Weedon, M.N., Cebola, I., Patch, A.-M., Flanagan, S.E., De Franco, E., Caswell, R.,
Rodríguez-Seguí, S.A., Shaw-Smith, C., Cho, C.H.-H., Allen, H.L., et al. (2014). Recessive
mutations in a distal PTF1A enhancer cause isolated pancreatic agenesis. Nat. Genet. *46*, 61–
64.

12. Cooper, G.M., Stone, E.A., Asimenos, G., Green, E.D., Batzoglou, S., and Sidow, A. (2005).
Distribution and intensity of constraint in mammalian genomic sequence. Genome Res. *15*,
901–913.

13. Davydov, E.V., Goode, D.L., Sirota, M., Cooper, G.M., Sidow, A., and Batzoglou, S. (2010).
Identifying a High Fraction of the Human Genome to be under Selective Constraint Using
GERP++. PLoS Comput. Biol. *6*, e1001025.

14. Garber, M., Guttman, M., Clamp, M., Zody, M.C., Friedman, N., and Xie, X. (2009).
Identifying novel constrained elements by exploiting biased substitution patterns. Bioinformatics
25, i54–i62.

15. Finucane, H.K., Bulik-Sullivan, B., Gusev, A., Trynka, G., Reshef, Y., Loh, P.-R., Anttila, V.,
Xu, H., Zang, C., Farh, K., et al. (2015). Partitioning heritability by functional annotation using
genome-wide association summary statistics. Nat. Genet. *47*, 1228–1235.

16. Kircher, M., Witten, D.M., Jain, P., O'Roak, B.J., Cooper, G.M., and Shendure, J. (2014). A
general framework for estimating the relative pathogenicity of human genetic variants. Nat.
Genet. *46*, 310–315.

17. Ritchie, G.R.S., Dunham, I., Zeggini, E., and Flicek, P. (2014). Functional annotation of
noncoding sequence variants. Nat. Methods *11*, 294–296.

18. Ionita-Laza, I., McCallum, K., Xu, B., and Buxbaum, J.D. (2016). A spectral approach
integrating functional genomic annotations for coding and noncoding variants. Nat. Genet. *48*,
214–220.

19. Huang, Y.-F., Gulko, B., and Siepel, A. (2017). Fast, scalable prediction of deleterious
noncoding variants from functional and population genomic data. Nat. Genet. *49*, 618–624.

20. Jagadeesh, K.A., Wenger, A.M., Berger, M.J., Guturu, H., Stenson, P.D., Cooper, D.N.,
Bernstein, J.A., and Bejerano, G. (2016). M-CAP eliminates a majority of variants of uncertain
significance in clinical exomes at high sensitivity. Nat. Genet. *48*, 1581–1586.

21. Rosenbloom, K.R., Armstrong, J., Barber, G.P., Casper, J., Clawson, H., Diekhans, M.,
Dreszer, T.R., Fujita, P.A., Guruvadoo, L., Haeussler, M., et al. (2015). The UCSC Genome
Browser database: 2015 update. Nucleic Acids Res. *43*, D670-681.

22. Lowe, C.B., Kellis, M., Siepel, A., Raney, B.J., Clamp, M., Salama, S.R., Kingsley, D.M.,
Lindblad-Toh, K., and Haussler, D. (2011). Three periods of regulatory innovation during
vertebrate evolution. Science *333*, 1019–1024.

23. Siepel, A., Pollard, K.S., and Haussler, D. (2006). New Methods for Detecting Lineage-
Specific Selection. In Research in Computational Molecular Biology, (Springer, Berlin,
Heidelberg), pp. 190–205.

24. Kim, S.Y., and Pritchard, J.K. (2007). Adaptive Evolution of Conserved Noncoding Elements
in Mammals. PLOS Genet. *3*, e147.

25. Marnetto, D., Mantica, F., Molineris, I., Grassi, E., Pesando, I., and Provero, P. (2018).
Evolutionary Rewiring of Human Regulatory Networks by Waves of Genome Expansion. Am. J.
Hum. Genet. *102*, 1–12.

26. Herrero, J., Muffato, M., Beal, K., Fitzgerald, S., Gordon, L., Pignatelli, M., Vilella, A.J.,
Searle, S.M.J., Amode, R., Brent, S., et al. (2016). Ensembl comparative genomics resources.
Database J. Biol. Databases Curation *2016*,.

27. Cotney, J., Leng, J., Yin, J., Reilly, S.K., DeMare, L.E., Emera, D., Ayoub, A.E., Rakic, P.,
and Noonan, J.P. (2013). The Evolution of Lineage-Specific Regulatory Activities in the Human
Embryonic Limb. Cell *154*, 185–196.

28. Villar, D., Berthelot, C., Aldridge, S., Rayner, T.F., Lukk, M., Pignatelli, M., Park, T.J.,
Deaville, R., Erichsen, J.T., Jasinska, A.J., et al. (2015). Enhancer Evolution across 20
Mammalian Species. Cell *160*, 554–566.

29. Don, P.K., Ananda, G., Chiaromonte, F., and Makova, K.D. (2013). Segmenting the human
genome based on states of neutral genetic divergence. Proc. Natl. Acad. Sci. *110*, 14699–
14704.

30. Ernst, J., and Kellis, M. (2010). Discovery and characterization of chromatin states for
systematic annotation of the human genome. Nat. Biotechnol. *28*, 817–825.

31. Ernst, J., and Kellis, M. (2012). ChromHMM: automating chromatin-state discovery and
characterization. Nat. Methods *9*, 215–216.

32. Hoffman, M.M., Buske, O.J., Wang, J., Weng, Z., Bilmes, J.A., and Noble, W.S. (2012).
Unsupervised pattern discovery in human chromatin structure through genomic segmentation.
Nat. Methods *9*, 473–476.

33. Blanchette, M., Kent, W.J., Riemer, C., Elnitski, L., Smit, A.F.A., Roskin, K.M., Baertsch, R.,
Rosenbloom, K., Clawson, H., Green, E.D., et al. (2004). Aligning Multiple Genomic Sequences
With the Threaded Blockset Aligner. Genome Res. *14*, 708–715.

34. Smit, A., Hubley, R., and Green, P. (2013). RepeatMasker Open-4.0.

35. Harrow, J., A, F., Jm, G., E, T., M, D., F, K., Bl, A., D, B., A, Z., S, S., et al. (2012).
GENCODE: the reference human genome annotation for The ENCODE Project. Genome Res.
22, 1760–1774.

36. Welter, D., MacArthur, J., Morales, J., Burdett, T., Hall, P., Junkins, H., Klemm, A., Flicek,
P., Manolio, T., Hindorff, L., et al. (2014). The NHGRI GWAS Catalog, a curated resource of
SNP-trait associations. Nucleic Acids Res. *42*, D1001-1006.

37. Iulio, J. di, Bartha, I., Wong, E.H.M., Yu, H.-C., Lavrenko, V., Yang, D., Jung, I., Hicks, M.A.,
Shah, N., Kirkness, E.F., et al. (2018). The human noncoding genome defined by genetic
diversity. Nat. Genet. 1.

38. Witowski, V., and Foraita, D.R. (2014). HMMpa: Analysing accelerometer data using hidden
Markov models.

39. Ernst, J., and Kellis, M. (2015). Large-scale imputation of epigenomic datasets for
systematic annotation of diverse human tissues. Nat. Biotechnol. *33*, 364–376.

40. Hahsler, C.B. and M. (2017). cba: Clustering for Business Analytics.

41. Bar-Joseph, Z., Gifford, D.K., and Jaakkola, T.S. (2001). Fast optimal leaf ordering for
hierarchical clustering. Bioinforma. Oxf. Engl. *17 Suppl 1*, S22-29.

42. Ernst, J., and Bar-Joseph, Z. (2006). STEM: a tool for the analysis of short time series gene
expression data. BMC Bioinformatics *7*, 191.

43. Kheradpour, P., and Kellis, M. (2014). Systematic discovery and characterization of
regulatory motifs in ENCODE TF binding experiments. Nucleic Acids Res. *42*, 2976–2987.

44. Kolde, R. (2015). pheatmap: Pretty Heatmaps.

45. Chen, X., and Tompa, M. (2010). Comparative assessment of methods for aligning multiple
genome sequences. Nat. Biotechnol. *28*, 567–572.

46. Zhang, M.Q. (1998). Statistical features of human exons and their flanking regions. Hum.
Mol. Genet. *7*, 919–932.

47. Sarda, S., Das, A., Vinson, C., and Hannenhalli, S. (2017). Distal CpG islands can serve as
alternative promoters to transcribe genes with silenced proximal promoters. Genome Res. *27*,
553–566.

48. Litman, G.W., Anderson, M.K., and Rast, and J.P. (1999). Evolution of Antigen Binding
Receptors. Annu. Rev. Immunol. *17*, 109–147.

49. Gusev, A., Lee, S.H., Trynka, G., Finucane, H., Vilhjálmsson, B.J., Xu, H., Zang, C., Ripke,
S., Bulik-Sullivan, B., Stahl, E., et al. (2014). Partitioning Heritability of Regulatory and Cell-
Type-Specific Variants across 11 Common Diseases. Am. J. Hum. Genet. *95*, 535–552.

50. Kichaev, G., Yang, W.-Y., Lindstrom, S., Hormozdiari, F., Eskin, E., Price, A.L., Kraft, P.,
and Pasaniuc, B. (2014). Integrating Functional Data to Prioritize Causal Variants in Statistical
Fine-Mapping Studies. PLoS Genet. *10*, e1004722.

51. Pickrell, J.K. (2014). Joint Analysis of Functional Genomic Data and Genome-wide
Association Studies of 18 Human Traits. Am. J. Hum. Genet. *94*, 559–573.

52. Telenti, A., Pierce, L.C.T., Biggs, W.H., Iulio, J. di, Wong, E.H.M., Fabani, M.M., Kirkness,
E.F., Moustafa, A., Shah, N., Xie, C., et al. (2016). Deep sequencing of 10,000 human
genomes. Proc. Natl. Acad. Sci. *113*, 11901–11906.

Figure Legends

Figure 1: Illustration of ConsHMM modeling approach. (a) The input to ConsHMM is a multi-

species alignment, which is illustrated for a subset of 6 species aligned to the human sequence.

At each position and for each species ConsHMM represents the information as one of three

observations: (1) aligns with a non-indel nucleotide matching the human sequence shown in

blue, (2) aligns with a non-indel nucleotide not matching the human sequence shown in yellow,

or (3) does not align with a non-indel nucleotide shown in gray. **(b)** Illustration of conservation

state assignments at a locus chr22:25,024,640-25,024,812. Only states assigned to at least one

nucleotide in the locus are shown. Below the conservation state assignments is a color

encoding of the input multiple sequence alignment according to panel (a). The major clade of

species as annotated on the UCSC genome browser[21] are labeled and ordered based on

divergence from human. Above the conservation state assignments are PhastCons constrained

elements and scores and PhlyoP constraint scores. The figure and **Supplementary Fig. 9**

together illustrate that positions of nucleotides that have the same status in terms of being in a

constrained element or not or have similar constraint scores can be assigned to different

conservation states depending on the patterns in the underlying multiple-species alignment.

Figure 2: Conservation state emission parameters learned by ConsHMM and enrichments

for other genomic annotations. (a) Each row in the heatmap corresponds to a conservation

state. For each state and species, the left half of the heatmap gives the probability of aligning to

the human sequence, which is one minus the probability of the not aligning emission.

Analogously, the right half of the heatmap gives the probability of observing the matching

emission. Each individual column corresponds to one species with the individual names

displayed in **Supplementary Fig. 5.** For both halves species are grouped by the major clades

and ordered based on the hg19.100way.nh phylogenetic tree from the UCSC genome browser,

with species that diverged more recently shown closer to the left.[21] The conservation states are

ordered based on the results of applying hierarchical clustering and optimal leaf ordering.[41] The

states are divided into eight major groups based on cutting the dendrogram of the clustering.

The groups are indicated by color bars on the left hand side and a white row between them.

Transition parameters between states of the model can be found in **Supplementary Fig. 6**. **(b)**

The columns of the heatmap indicate the relative enrichments of conservation states for

external genomic annotations (**Methods**). For each column, the enrichments were normalized to

a [0,1] range by subtracting the minimum value of the column and dividing by the range and

colored based on the indicated scale. Values for these enrichments and additional enrichments

can be found in **Supplementary Fig. 8** and **Supplementary Table 2** and enrichments for

individual repeat classes and families can be found in **Supplementary Fig. 13.**

Figure 3: Conservation state positional enrichments. Plots of positional fold enrichments of

conservation states relative to **(a)** start of exons of protein coding genes in phase 0, **(b)** end of

exons of protein coding genes and TSS of **(c)** protein coding, and **(d)** pseudogenes genes.

Positive values represent the number of bases downstream in the 5' to 3' direction of

transcription, while negative values represent the number of bases upstream. Enrichments

relative to gene annotations are based on a genome-wide background. The subset of states

included in panels (a)-(d) were the states that had at least a 3 fold enrichment at some position

within +/-2kb from the anchor point. Also shown are positional plots relative to the central

nucleotide of a set of instances of **(e)** STAT and **(f)** POU5F1 motifs. The subset of states

included in (e), (f) are the states that had an enrichment of at least 1.5 for some position within

+/- 15bp from the center nucleotide of either motif. Enrichments for motif instances were

computed relative to the portion of the genome scanned for regulatory motifs in Kheradpour and

Kellis (2014), which excludes coding, 3'UTR, and repeat elements. Additional position

enrichment plots can be found in **Supplementary Fig. 10**.

Figure 4: Conservation states enrichment for chromatin states, GO terms, DHS and

repeat elements. (a) Median fold enrichment of conservation states (rows) for one of 25

chromatin states from a previously defined chromatin state model defined across 127 samples

of diverse cell and tissue types (columns).[15] Only conservation states that had the maximum

value for at least one chromatin state are shown, and those values are boxed. See

Supplementary Fig. 14 for the enrichments of all conservation states. **(b)** $-\log_{10}$ p-value

(uncorrected) of the conservation states (rows) for the GO term (columns) where each

conservation state is associated with its top 5% genes based on promoter regions (**Methods**).

Only GO terms which were the most enriched term for some conservation state are shown,

restricted to the top 10 terms based on the significance of the enrichment. Only conservation

states that had the most significant enrichment for one of the displayed GO terms are shown,

with the maximal enrichments boxed. The full set of conservation states with additional GO

terms are in **Supplementary Fig. 12**. **(c)** Relative enrichments of conservation states for DHS

across cell and tissue types. Only conservation states with at least a 2 fold enrichment in one

sample considered are shown. Enrichment values were $\log_2$ transformed and then row

normalized by subtracting the mean (right heatmap) and dividing by the standard deviation.

States and experiments were then hierarchically clustered and revealed two major clusters. In

the top cluster conservation states showed the greatest enrichment for experiments in which the

DHS also strongly enriched for CpG islands (top heatmap). In the bottom cluster conservation

states generally had the strongest relative preference for a number of fetal related samples. **(d)**

Enrichment of conservation states with the maximal enrichment for LINE, SINE, LTR or DNA

repeats next to the state align probabilities for primates. These states all had low align

probabilities outside of primates, but their differences among primates corresponded to

substantial differences in repeat enrichments.

Figure 5: **Relationship of conservation states with constrained elements and scores.**

Precision-recall plots for recovery of **(a)** TSS of protein coding genes, **(b)** TES of protein coding

genes, and **(c)** the start of exons of protein coding genes. Recovery based on ordering

ConsHMM conservation states for their enrichment for the target set in the training data, then

cumulatively adding the states in that ranked order and evaluating on the test data is shown with

a series of blue dots (**Methods**). The first few conservation states added are labeled with their

state number. Recovery based on ranking from highest to lowest value of constraint scores is

shown with continuous lines. Recovery based on score partitioning into 400 bins and

subsequent ordering based on enrichment for the target set in the training data, then

cumulatively adding bins in that ranked order and evaluating on the test data is shown in a

series of dots of the same color as the continuous line corresponding to the score. Recovery of

target test bases by a constrained element set is shown with a single dot for each constrained

element set. See **Supplementary Figs. 17-18** for plots based on additional targets. **(d)** The

graph shows the fold enrichment for Fetal Brain DHS[5] within the non-exonic portion of each

conservation state, separately for those bases in a PhastCons constrained element (pink) and

bases not in such an element (blue). Enrichments within constrained elements varied

substantially depending on the conservation state. For a given conservation state, bases in a

constrained element had greater enrichments than bases not in a constrained element,

illustrating complementary information of conservation states and constrained elements. See

Supplementary Fig. 19 for graphs based on different element sets or DHS data.

Figure 6: Conservation states and association with human genetic variation. **(a)** The

panel displays the $\log_2$ fold enrichment of each state for common SNPs (pink) and GWAS

catalog variants relative to common SNPs (blue). State 1, associated with high alignability and

matching through all vertebrates, showed the greatest depletion of common SNPs and the

highest enrichment for GWAS variants relative to common SNPs. States 55-57 and 87-89

exhibited the opposite pattern having the greatest enrichment for common SNPs and the

greatest depletion of GWAS variants relative to this background. The second most depleted

state for common SNPs, which did not show enrichment for GWAS catalog SNPs, was state 96

which captured large gaps in the assembly (**Supplementary Fig. 8**). **(b)** Panel shows the

representation of state emission parameters from **Fig. 2a** for the subset of states highlighted in

panel (a). The states with the greatest depletion of GWAS variants all had relatively high

alignability at least through primates, but low matching probabilities for almost all species except

a few closely related primates. (**c**) Scatter plot of conservation state fold enrichments of top 1%

CDTS bases (x-axis) and GERP++ elements (y-axis). The states are colored by their group

assignment and the size of the point corresponds to the enrichment for CpG islands as

indicated based on the scale on the right. Those states which were at least four fold enriched in

either set are labeled. (**d**) Applying the heritability partitioning enrichment method of Finucane et

al.[15] on two disjoint subsets of bases in PhastCons elements, with eight phenotypes previously

analyzed with heritability partitioning in the context of a baseline set of annotations (**Methods**).[15]

One set of bases are those in PhastCons elements that are also in one of the seven

conservation states showing the greatest enrichment for DHS in its non-exonic portion (States

1-5, 8, and 28) covering 51.9% of PhastCons bases (pink). The other set are those bases in

PhastCons elements overlapping the remaining 93 states covering 48.1% of PhastCons bases

(blue).

a

Human	T	T	T	C	C	T	G	A	C	T	T
Chimp	T	T	T	C	C	T	G	A	C	T	T
Bushbaby	T	C	T	G	C	T	T	C	C	T	T
Rat	-	C	T	T	C	T	G	A	A	T	T
Alpaca	-	-	-	C	C	T	G	G	C	A	T
Megabat	T	C	-	C	C	T	G	A	T	T	T
Parrot	-	-	-	-	-	-	-	-	-	-	-

■ Aligning and matching the human sequence

■ Aligning but not matching the human sequence

■ Not aligning to the human sequence

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6