Evidence for rapid evolution in a grassland biodiversity experiment

Sofia J. van Moorsel*[1,6], Marc W. Schmid*[1,2,6], Niels C.A.M. Wagemaker[3], Thomas

van Gurp[4], Bernhard Schmid[1] and Philippine Vergeer[3,5,6]

*Equal contribution

[1]Department of Evolutionary Biology and Environmental Sciences, University of

Zürich, Winterthurerstrasse 190, CH-8057, Switzerland

[2]MWSchmid GmbH, Möhrlistrasse 25, CH-8006 Zürich

[3]Institute for Water and Wetland Research, Radboud University Nijmegen,

Heyendaalseweg 135, NL-6500 GL Nijmegen

[4]Deena Bioinformatics, Haverlanden 237, NL-6708 GL Wageningen

[5]Department of Environmental Sciences, Plant Ecology and Nature Conservation

group, Wageningen University, Droevendaalsesteeg 3, NL-6708 PB Wageningen

[6] Authors for correspondence:

Sofia J. van Moorsel, sofia.vanmoorsel@ieu.uzh.ch

Marc W. Schmid, contact@mwschmid.ch

Philippine Vergeer, philippine.vergeer@wur.nl

Summary

- In long-term grassland biodiversity experiments positive effects of biodiversity on plant productivity commonly increase with time. Previously it was shown that differential selection in monoculture and mixed-species grassland communities could lead to the rapid emergence of monoculture and mixture phenotypes. Underlying mechanisms for such rapid phenotypic responses are however still unclear.

- We hypothesized that in biodiversity experiments pre-adapted genotypes or epigenetic variants could be sorted out from the standing genetic or epigenetic variation.

- To test if biodiversity acted as a selective environment, we grew offspring from plants that were exposed for twelve years to a monoculture or mixture environment in a biodiversity experiment (Jena Experiment) under controlled greenhouse conditions. Using epiGBS, a genotyping-by-sequencing approach combined with bisulfite conversion to provide integrative genetic and epigenetic data, we showed that plants with a monoculture or mixture background were genetically distinct.

- Our data reveal strong genetic and epigenetic divergence within species according to selection history and suggest variation in epigenetic variation to be largely under genetic control. This pattern was consistently observed across six perennial grassland species. Our results suggest that selection of genetic variants caused the rapid emergence of monoculture and mixture types in the Jena Experiment.

Key words: biodiversity, epiGBS, epigenetic variation, genetic divergence, herbaceous plant species

Introduction

The Earth's biosphere is currently challenged by the impacts of anthropogenic environmental change and plant populations may encounter new abiotic or biotic environment due to climate-induced range shifts (Ouborg *et al.,* 2006). The unprecedented rate of environmental change raises the question whether natural communities can adapt fast enough to novel abiotic or biotic conditions. Biodiversity has been shown to buffer ecosystem towards climatic extremes and novel environmental conditions (Isbell *et al.,* 2015) and in addition it was shown that co-evolution among plants comprising a community could buffer the impact of an extreme climatic event (van Moorsel *et al.*, 2018a).

Adaptive responses of plant populations to environmental factors (e.g., Schmid 1985; Joshi *et al.*, 2001) and biotic interactions (Gervasi & Schiestl 2017) are well studied, but little effort has been devoted to studying the influence of community diversity on population structure and productivity (but see Lipowsky *et al.,* 2011; Kleynhans *et al.,* 2016). In particular, the influence of multi-species interactions for the adaptive response of a species is largely unknown, despite a growing body of evidence pointing towards the importance of species-interaction networks for the maintenance of ecosystem stability (Bastolla *et al.,* 2009). It is conceivable that the feedback between species interactions and their adaptive responses shapes community-level ecosystem functioning (van Moorsel *et al.*, 2018).

In the 1960s it was proposed that there are large differences in the time scales between ecological and evolutionary processes (Slobodkin 1961), but now it is well known that micro-evolutionary and ecological processes can occur on the same temporal scale (reviewed by Hairston *et al.,* 2005; Schoener 2011). Thus, it appears that micro-evolutionary processes may allow for an evolutionary rescue in a rapidly changing environment. However, evidence of genetic adaptation to mild stresses in natural situations over short time scales of a few generations is still scarce. Understanding how biodiversity, i.e. interactions between species, shapes this evolutionary response will be instrumental to anticipate how ecosystems may change in response to global change.

Adaptation depends on several factors. For organisms with a short generation time and asexual reproduction, such as clonal populations of bacteria, mutations and horizontal gene transfer are the main sources of genetic variation (Anderson *et al.*,

2011). However, for species with longer generation times such as perennial plants

selection more likely acts on standing genetic variation (Barrett & Schluter 2008),

resulting in a sorting-out of suitable genotypes (Fakheran *et al.,* 2010). Furthermore,

plants may adjust to a novel environment by phenotypic plasticity (Price *et al.,* 2003).

A possible indication for selection in grassland plant communities is the observed

strengthening of biodiversity effects in field biodiversity experiments (Cardinale *et*

al., 2007; Fargione *et al.,* 2007; Reich *et al.,* 2012; Meyer *et al.,* 2016). An increase in

complementarity between species resulted in an increasing positive effect of diversity

on productivity over time (e.g. Meyer *et al.,* 2016). The increased complementarity

could also have been a result of phenotypic plasticity (Ghalambour *et al.,* 2007).

However, recent common garden experiments with plant material from a grassland

biodiversity experiment (the Jena Experiment, Roscher *et al.,* 2004) gave a clear

indication for genetic divergence between monoculture and mixture types in multiple

plant species (Zuppinger-Dingley *et al.,* 2014; van Moorsel *et al.,* 2018b), suggesting

that natural selection in response to community diversity had occurred in the field. As

a consequence, mixture types showed stronger complementarity effects (Loreau &

Hector 2001) than monoculture types when grown in mixture, i.e. community

productivity was increased due to increased differences in functional traits between

species for mixture types (Zuppinger-Dingley *et al.,* 2014; van Moorsel *et al.,* 2018b).

The emergence of such monoculture and mixture types, with different growth

performance and plant functional trait variation, thus suggested that community

diversity in the field likely acted as a selective environment (Zuppinger-Dingley *et al.,*

2014; Rottstock *et al.,* 2017; van Moorsel *et al.,* 2018b). However, direct evidence for

a genetic divergence between the different populations in the field experiment is still

missing.

It should be noted that epigenetics, here defined as meiotically heritable changes in

gene expression without changes to the underlying DNA sequence (Verhoeven *et al.,*

2016), has also been proposed to play a role (Bird 2007; Bossdorf *et al.,* 2008; Tilman

& Snell-Rood 2014). In a comment accompanying the publication of Zuppinger-

Dingley *et al.* (2014), Tilman and Snell-Rood wrote: "[…] chance that the differences

between the high- and low-diversity selection groups were due to genetic divergence.

However, it is possible that epigenetic factors […] could have had a simultaneous

role" (Tilman & Snell-Rood 2014). Recent work on epigenetic recombinant inbred

lines (epiRILs) of *Arabidopsis thaliana* suggests a considerable contribution of

induced epialleles to phenotypic variation, which is independent of genetic variation

(Cortijo *et al.*, 2014; Kooke *et al.*, 2015). However, the importance of epigenetics in

natural populations and whether it contributes to adaptation remains elusive

(Quadrana & Colot 2016), in particular because it is difficult to separate epigenetic

from genetic variation. A first step to disentangle genetic from epigenetic variation

was achieved with apomictic clones of *Taraxacum officinale*, suggesting that

differences in flowering time were mediated by differences in DNA methylation

(Wilschut *et al.,* 2016). However, given the fundamental difference between apomixis

(apomeiosis, parthenogenesis, autonomous endosperm formation) and sexual

reproduction, the results from this study may not be directly transferable to non-

apomictic plant species.

Here, we tested whether community diversity could act as a selective force leading

to a divergence of phenotypes into monoculture and mixture types (Zuppinger-

Dingley *et al.*, 2014; van Moorsel *et al.,* 2018b). In particular, we tested the

hypothesis that the differentiation of plants into phenotypically distinct mixture

(exhibiting stronger biodiversity effects) and monoculture types (exhibiting weaker

biodiversity effects) within species was paralleled by genetic differentiation.

Anticipating that the differentiation might also have been be due to epigenetic

variation, we chose a method which allowed us to analyze genetic and epigenetic

variation between plants of six species, all herbaceous perennials of European

grassland.

Materials and Methods

Plant selection histories

To test whether plant types selected over eleven years in mixtures differ genetically or

epigenetically from those types selected in monocultures, we chose six species grown

in monoculture and mixture plots in the Jena Experiment (Jena, Thuringia, Germany,

51˚N, 11˚E, 135 m a.s.l., see Roscher *et al.* (2004) for experimental details). The

following species belonging to three functional groups were selected: The three small

herbs *Plantago lanceolata* L*., Prunella vulgaris* L. and *Veronica chamaedrys* L*.,* the

tall herb *Galium mollugo* L*.,* and the two legumes *Lathyrus pratensis* L. and

Onobrychis viciifolia Scop. For the experiment plants from three different selection

histories were used (Fig. 1). Plants without a selection history in the experimental

field plots of the Jena experiment were obtained from commercial seed suppliers

(Rieger Hoffmann GmbH, Blaufelden-Raboldshausen, Germany and Otto Hauenstein

Samen AG, Switzerland), who also provided the seeds for the original set up of the

Jena Experiment in 2002. Plants with a selection history in either mixture or

monoculture had been growing in the Jena Experiment since 2002. In spring 2010,

plant communities of 48 plots (12 monocultures, 12 two-species mixtures, 12 four-

species mixtures and 12 eight-species mixtures) of the Jena Experiment were

collected as cuttings, multiplied by repeated further cutting and then transplanted into

an experimental garden in Zurich, Switzerland, in the identical plant composition as

the original experimental plots for the first controlled sexual reproduction among co-

selected plants (Zuppinger-Dingley *et al.*, 2014). In spring 2011, the seedlings

produced from the seeds of the first controlled sexual reproduction in Zurich were

transplanted back into those plots of the Jena Experiment from where the parents had

originally been collected. In these newly established plots, plant communities with an

identical composition to the original communities were maintained for three years

until 2014 (van Moorsel *et al.*, 2018c).

Seed collection

To ensure a second sexual reproductive event for the collection of seed material,

entire plant communities from some of the experimental plots replanted in Jena in

2011 were excavated in March 2014 and used to established new 1-m^2 plots in the

experimental garden in Zurich. For this purpose, we excavated blocks of soil (0.5 m^2)

from plots in Jena including plant vegetation, seedlings and dormant seeds to make

sure we transferred the entire plant community. These blocks of soil were then

transferred and spread out into the plots in the experimental garden. We added a 30

cm layer of soil (1:1 mixture of garden compost and field soil, pH 7.4, commercial

name Gartenhumus, RICOTER Erdaufbereitung AG, Aarberg, Switzerland) to each

plot to make sure the plants established. As for the first sexual reproduction event in

2010, netting around each plot minimized the possibility of cross-pollination between

the same species from different selection histories. Seeds from this second controlled

sexual reproduction event were collected throughout the growing season of 2014 from

monoculture plots and 4- and 8-species mixture plots. The exact community

composition of the plots the seeds originated from is listed in Table S2. Seeds from

different mother plants were pooled together. The dry seeds were stored at 5° C for

cold stratification until germination.

Common garden experiment

Seeds were germinated in December 2014 with those from the same species being

planted within the same day. Germination was done in germination soil

("Anzuchterde", Ökohum, Herbertingen, Germany) under constant conditions in the

glasshouse without additional light. Seedlings were planted in monocultures of four

individuals and mixtures of two plus two individuals into 2-L pots filled with

agricultural soil (50% agricultural sugar beet soil, 25% perlite, 25% sand; Ricoter AG,

Aarberg, Switzerland). See van Moorsel *et al*. (2018b) for the experimental design

and the planted combinations. In total, we planted 36 monoculture assemblies and 81

mixture assemblies from mixture history, 48 monoculture assemblies and 159 mixture

assemblies from monoculture history and 33 monoculture assemblies and 100 mixture

assemblies from seedlings without a common selection history. Every species

combination was replicated, if possible, six times for each selection history (resulting

in 457 pots, referred to as "assemblies", and 1828 plants). Seedlings that died within

the first two weeks were replaced with seedlings of the same age.

The experiment was set up in six blocks with each block representing a replicate.

Every block contained ca. 80 pots and within each block, pots were placed in the

glasshouse in a randomized fashion without reference to selection history or species

assembly. Single pots always contained four plants of a single selection history.

During the experiment, plants were watered according to demand and grown at

constant temperatures (17-20°C during the day, 13-17°C during the night) with no

additional light added. The plants were not fertilized. The fungicide Fenicur (*Oleum*

foeniculi, Andermatt Biocontrol) was applied on 5 May to control powdery mildew

(*Podosphaera* spp.).

Sampling

Samples for subsequent epigenetic and genetic analysis were harvested between 18

and 28 May 2015, after twelve weeks of plant growth in the glasshouse. All four

plants were sampled in each pot. One young leaf per plant was cut from the living

plant and immediately shock-frozen in liquid nitrogen. The samples were then stored
at –80° C in Zurich before shipment to the Netherlands for further processing.
215

Genetic analysis

We measured both genetic and epigenetic variation in monoculture- and mixture-type
plants when propagated in monoculture and two-species mixed assemblies using a
novel reference-free bisulfite method (van Gurp *et al.*, 2016). This method allows
measuring DNA cytosine methylation levels and identifying single nucleotide
polymorphisms with merely one lane of sequencing per 96 samples.
222

Sample procession and library preparation

The epiGBS protocol used is a further developed protocol based on the protocol of
van Gurp *et al.* (2016). The main improvements are the used enzyme combination, a
"wobble" adapter facilitating the computational removal of PCR duplicates, and a
conversion control nucleotide that allows easier Watson / Crick identification. For 348
samples, (Csp6I/NsiI) epiGBS libraries were created and sequenced on 4 Hiseq 2500
lanes. These samples were divided over six species and three selection histories (see
Tables S1, S2).
231

DNA extraction

Plant material was disrupted by bead-beating frozen leaf tissue in a 2 mL eppendorf
tube with 2-3 mm stainless steel beads. No more than 100 mg of fresh tissue was used
per sample. DNA isolation was performed using the NucleoSpin® 8 Plant II Core Kit
(740669.5 Macherey Nagel). We followed the manufacturers protocol with the
following modifications. Cell lysis was performed using Cell lysis buffer PL1 for 30
instead of 10 min. After lysis and initial centrifugation, the lysate was carefully
pipetted to a fresh 2.5 mL tubes, avoiding the cell debris. An extra centrifugation of 5
min at 18.000 g step was performed and the lysate transferred to a 96-well rack for the
next steps. At the step were the washed columns were dried, we centrifuged 5 min at
4800 g to get rid of the last remaining wash buffer. The restriction enzymes of the first
step of the epiGBS are very sensitive to ethanol and other contamination. DNA
concentration was determined using Qubit® 2.0 Fluorometric dsDNA HS Assay Kit
(Q32851 Life technologies).

DNA digestion and adapter ligation

Per individual, 30–300 ng genomic DNA was digested overnight (17 hrs) at 37° C in a

volume of 40 µL containing 1x FD buffer (Thermo Scientific), and 2 µL of both

*Csp*6I (FD0214, Thermo Scientific) and *Nsi*I (R0127S, NEB). Following digestion,

barcoded "wobble" adapters were ligated to the fragments (Fig. S1). To minimize the

possibility of misidentifying samples as a result of sequencing or adapter synthesis

error, all pair-wise combinations of barcodes differed by a minimum of three

mutational steps. Barcode lengths were modulated from 4 bp to 6 bp to maximize the

balance of the bases at each position in the overall set. For the ligation, 4 µL of a

sample specific barcode combination of both BA and CO adapters (600 pg/µL), 6 µL

T4 DNA ligase buffer, 1 µL T4 DNA ligase (M0202M, NEB) and 5 µL of distilled

water were added to the digestion mix to a total volume of 60 µL. Ligation was

performed for 3 hours at 22° C followed by 4° C overnight.

Pooling, clean-up and nick translation

In order to assess the quality of libraries, the pooling was performed per species level

in batches of around 12 samples per pool. When pooled, the total volume of the pool

was reduced by Qiaquick PCR cleanup (28104, Qiagen) to 40 µL. The libraries were

size selected by a 0.8x Agencourt AMPure XP (A63880, Beckman coulter)

purification favouring > 200 bp DNA fragments and eluted in a total volume of 22

µL. To prevent the formation of adapter dimers, the barcoded adapters were not

phosphorylated. Therefore, after the ligation, the insert DNA fragment–adapter

connection was nicked at the 3' positions of the insert DNA fragment (Fig. S1). This

nick is repaired by Nick translation that recreates the total non-(5mC) methylated

adapter strand and during that process also "unwobbles" the adapters since the

removed nucleotides are replaced by complementing nucleotides. This nick repair

prevents the partial loss of the adapter during bisulfite treatment. The nick translation

reaction (1 hour at 15° C) was performed in a reaction of 25 µL containing 19.25 µL

of the purified library, 2.5 uL of 10 mM 5-methylcytosine dNTP Mix (D1030 Zymo

research), 2.5 uL NEBuffer 2 and 0.75 uL DNA polymerase I (M0209, NEB).

Bisulfite conversion

For bisulfite conversion of non-methylated cytosines, 20 µL of the nick-translated

library was used. Bisulfite treatment was performed using the EZ DNA Methylation-

Lightning™ Kit (Zymo Research) with the following program according to the

manufacturers protocol: 8 min at 98° C, 1 h at 54 °C followed by up to 20 h at 4° C.

epiGBS PCR

Library amplification was performed in four individual 10 µL reactions containing 1

µL ssDNA template, 5 µL KAPA HiFi HotStart Uracil+ ReadyMix

(Kapabiosystems), 3 pmol of each illumina PE PCR Primer (5'-

AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTT

CCGATCT-3' and 5'-

CAAGCAGAAGACGGCATACGAGATCGGTCTCGGCATTCCTGCTGAACCGC

TCTTCCGATCT-3'). Temperature cycling consisted of 95° C for 3 min followed by

18 cycles of 98° C for 10 s, 65° C for 15 s, 72° C for 15 s with a final extension step

at 72° C for 5 min. Replicate PCR products were pooled and quantified using a

Qubit® dsDNA HS Assay Kit (Life technologies). The quality of the Libraries was

assessed by analyzing 1 µL on a High Sensitivity DNA chip on a 2100 Bioanalyzer

system (Agilent). Libraries were considered suitable for sequencing if the majority of

DNA fragments were between 150–400 bp. When the libraries passed quality control,

they were pooled according to concentration and number of samples in the species

pool, so that each individual sample was expected to yield an equal number of clusters

on the Illumina flow cell. Before sequencing, the libraries were spiked with 10% PhiX

control to increase the complexity of the libraries.

Sequencing

Finally, Paired-End sequencing was performed on a Hiseq2500 sequencer using the

HiSeq v4 reagents and the latest version of the HiSeq Control Software (v2.2.38),

which optimizes the sequencing of low-diversity libraries

(http://res.illumina.com/documents/products/technotes/technote-hiseq-low-

diversity.pdf). As the first five cycles of a sequencing run are used to calculate the

color matrix, our barcode design achieves almost perfect balance of the first five

nucleotides when equal numbers of sequences are obtained per "A" barcode. The "B"

barcodes do not have this requirement; hence same length barcodes were used.

Sequencing reads were demultiplexed and deposited at SRA (SRA accession ID

SRP132258). The barcode files can be accessed on Zenodo (DOI

10.5281/zenodo.1167563). In total, we generated 132'884'678, 130'721'280,

81'376'294, 140'883'779, 215'473'088, and 137'868'869 short reads for *G. mollugo*,

L. pratensis, *O. viciifolia*, *P. lanceolata*, *P. vulgaris*, and *V. chamaedrys*, respectively.

Statistical analysis

De-multiplexing, *de novo* reference construction, trimming, alignment, strand-specific

variant calling, and methylation calling were done for each species as described in van

Gurp *et al.* (2016). *De novo* reference sequences were annotated with DIAMOND

(protein coding genes; NCBI non-redundant proteins as reference; version 0.8.22;

Buchfink, Xie, and Huson 2015) and RepeatMasker (transposons and repeats;

Embryophyta as reference "species"; version 4.0.6; Smit, Hubley, and Green 2013–

2015). We summarized the transposable element and repeat classes into "transposons"

comprising DNA, LTR, LINE, SINE, and RC transposon, and "repeats" including

satellite, telomeric satellite, simple, rRNA, snRNA, unknown, and unclassified

repeats. The annotation was then used to classify the genetic and epigenetic variants

into the different feature contexts (e.g., to identify whether a single nucleotide

polymorphism is located in a gene or a transposon). A summary of the reference

sequences is given in Table S3.

Genetic variation: visualization of genetic distances with single

nucleotide polymorphisms (SNPs)

The following individuals with a SNP calling rate below 30% were *a priori* removed

from the analysis of genetic variation: "pool_pla_lan_15", "pool_pla_lan_46",

"pla_lan_81", "pla_lan_82", "pla_lan_52", "pla_lan_83", "pla_lan_95",

"pla_lan_111", "pru_vul_60", "pru_vul_79", "pru_vul_80", "pru_vul_87",

"pool_pru_vul_22", "pool_pru_vul_24", "ono_vic_15", and "ver_cha_73". These

samples were well distributed across the experimental group, i.e., one or two for a

single experimental group, except for the mixture-type mixture assembly of *P.*

lanceolata for which three individuals were removed. For each species, we filtered the

genetic variation data for single nucleotide polymorphisms (SNPs) sequenced in all

individuals with a total coverage between 5 and 200. SNPs homozygous for either the

reference or the alternative allele in more than 90 % of all individuals were removed

as uninformative SNPs. To reduce the impact of false positive SNP calls, we removed

all SNPs located in contigs with more than 1 SNP per 50 basepairs because higher

rates appear unlikely and may also originate from spurious alignments to the wrong

reference contig (considering that the reference contigs represent only a minor

fraction of the entire genome, there may be many reads originating from other

locations not represented with a reference contig which are still similar enough to

(wrongly) align to the reference). This also avoids that few contigs with high SNP

rates drive the tests for genetic differentiation.

SNP allele frequencies were scaled with the function "scaleGen" from adegenet

(version 2.0.1; Jombart 2008) and genetic distances between the individuals were

visualized with t-SNE (van der Maaten and Hinton 2008, van der Maaten 2014). We

calculated 100 maps starting from different random seeds and selected the map with

the lowest final error. Individual maps were calculated in R with the package Rtsne

(version 0.13; van der Maaten and Hinton 2008, van der Maaten 2014). Parameters

for the function Rtsne were pca = FALSE, theta = 0, perplexity = 10 (except for *O.*

viciifolia for which perplexity = 5). To select the SNPs with the highest differentiation

between the populations, we calculated Jost's D (Jost 2008) with the function

"basic.stats" from hierfstat (version 0.04-22; Goudet & Jombart 2015) and only

included the top 5% in the visualization.

Genetic variation: test for genetic differentiation between populations

with single nucleotide polymorphisms (SNPs)

SNP data were processed and filtered as described before. Considering that our design

only included two factors with two and three levels (assembly in the glasshouse and

selection history respectively; and incomplete for all species except *P. vulgaris*), we

did not use a hierarchical model (with assembly nested within) to test for genetic

differentiation. Instead, we tested each factor within all levels of the other factor for

genetic differentiation. Taking *P. vulgaris* as an example, we tested for genetic

differentiation between plant histories within monoculture and mixture assemblies

(between all three histories and between monoculture and mixture types), and

between assemblies within the naïve, monoculture, and mixture-type selection

histories. For each test, we extracted the corresponding individuals and tested for

genetic differentiation with the G-statistic test (Goudet *et al.,* 1996, function

gstat.randtest implemented in the package hierfstat, version 0.04-22; Goudet &

Jombart 2015). We did not correct for multiple testing because *P*-values were derived

from 999 simulations and were thus limited to a minimal value of 0.001. Instead, we

used a significance threshold of 0.01 and provide plots showing histograms of

permuted G-statistics and the observed G-statistics. This analysis was carried out with

the (1) entire data set, (2) SNPs located within genes, and (3) SNPs located within

transposons. We chose to separately test SNPs in genes and transposons because we

expected that selection more likely acted on genes and that selection of transposons

would primarily occur due to genetic linkage to an advantageous gene. In addition, we

expected that SNP calls are more reliable within genes because many transposon

families tend to be highly repetitive.

To estimate the extent to which the genetic variation was caused by the

differentiation between populations, we calculated average (i.e., across all tested

SNPs) pairwise F_{ST} values with the function pairwise.fst from the package adegenet

(version 2.0.1; Jombart 2008). Assuming that only few loci were under selection,

many SNPs have F_{ST} values close to zero (only SNPs under selection have F_{ST} values

clearly larger than zero). To estimate the maximal divergence between the

populations, we therefore also calculated the F_{ST} of each individual SNP and extracted

the 99[th] percentile (we chose the 99[th] percentile because this is more robust than the

highest value).

Epigenetic variation: characterization of genome-wide DNA methylation
levels

402 For each species, we filtered the epigenetic variation data for cytosines sequenced in

403 at least three individuals per population (i.e., experimental group) with a total

404 coverage between 5 and 200. To provide an overview of the genome-wide DNA

405 methylation levels of the six species or each experimental group per species, we

visualized the DNA methylation levels of all cytosines averaged across all individuals with violin plots. We also visualized the average DNA methylation level within genes, transposons, repeats, and unclassified reference contigs with heat maps. Both were done either using all sequence contexts (CG, CHG, CHH) at once or separately for each sequence context.

Epigenetic variation: identification of differentially methylated cytosines (DMCs)

DNA methylation data was processed and filtered as described before. Variation in DNA methylation at each individual cytosine was then analyzed with a linear model in R with the package DSS (version 2.24.0; Park and Wu 2016), according to a design with a single factor comprising all different experimental groups (similar to the approach described for RNA-Seq, Schmid 2017 and the testing procedure in Schmid *et al.,* 2018). Specific groups were compared with linear contrasts and *P*-values for each contrast were adjusted for multiple testing to reflect false discovery rates (FDR). Taking *P. vulgaris* as an example, we compared the three plant histories across both assemblies and within each assembly to each other. Likewise, we compared the two assemblies across all plant histories and within each to each other. A cytosine was defined as differentially methylated (DMC) if the FDR was below 0.01 for any of the contrasts.

Results

Visualization of genetic distances between the plant individuals using 5% of the loci with the strongest divergence between the populations clearly separated the individuals according to their selection history in five out of six species (Fig. 2). In addition to the separation by the selection history, individuals from *L. pratensis* also clustered according to the current diversity level (i.e., assembly). However, considering that the plants were assigned randomly to the assembly treatment, we expected a genetic differentiation according to the selection history but not the assembly. We therefore tested for a significant genetic divergence between the

selection histories and the assemblies with the G-statistics test (Fig. 3, S3 and S4,

Goudet *et al.,* 1996).

Genetic differentiation was significant in all data sets ($P <= 0.001$), with two

exceptions. First, the plant histories of *P. lanceolata* did not exhibit any significant

genetic differentiation (only the test with all plant histories within monoculture

assemblies in the data set with all SNPs was significant). Second, the test including

only the monoculture and mixture types within the mixture assemblies was not

significant for *L. pratensis*. In contrast, the tests comparing the monoculture and

mixture assemblies within each of the plant histories were never significant ($P >$

0.01).

To estimate the amount of genetic variation explained by the plant histories, we

calculated average pairwise F_{ST} values (Table S4) and the 99[th] percentiles of the SNP-

wise F_{ST} values (Table 1 and Tables S5, S6). Average pairwise F_{ST} values for the

different plant histories were between 0.014 (naïve *vs.* monoculture type within the

monoculture assemblies of *L. pratensis*) and 0.1 (naïve *vs.* monoculture or mixture

type within monoculture assemblies of *P. vulgaris*). The 99[th] percentiles were

markedly higher and between 0.13 (monoculture vs. mixture types within mixture

assemblies of *G. mollugo*) and 0.67 (all plant histories within mixture assemblies of *P.*

vulgaris). Thus, overall, 1.4% to 10% of the genetic variation was explained by plant

histories. However, for individual SNPs, plant histories could explain up to 67% of

the genetic variation.

To get an overview of the DNA methylation data, we visualized overall DNA

methylation levels for each plant species, sequence context (CG, CHG, CHH), and

genomic feature context (genes, transposons, repeats, and unclassified contigs, Fig. 4).

For all species, DNA methylation was generally highest in the CG context (80.3%),

lower in the CHG context (59.7%), and lowest in CHH context (13.2%). Differences

between species were most pronounced in the CHG context in which *L. pratensis*

(77.1%) and *P. lanceolata* (85.8%) exhibited markedly higher methylation levels than

the other four species (56.6%, 52.3%, 43.6%, and 53.1% in *G. mollugo, O. viciifolia,*

P. vulgaris, and *V. chamaedrys,* respectively). Within each species and context, DNA

methylation was highest in transposons and lowest in genes (Fig. 4B). Overall, these

patterns are within the range of what was reported previously for other angiosperms

(Niederhuth *et al.,* 2016).

For an initial comparison between the experimental groups, we visualized the

overall DNA methylation levels as we did about for the genetic variation, but for each

experimental group separately (Fig. S4). Given that the overall methylation levels

appeared to be similar, we tested for significant differences in DNA methylation

levels at each individual cytosine. On average, 1.8% of all tested cytosines were

significant in at least one of the tested contrasts (FDR < 0.01, "DMCs" for

differentially methylated cytosines, Table 1 for all contexts and Tables S5, S6, and S7

for each context separately). Relative to the total number of cytosines tested,

differences between plant histories (tested within or across both assemblies) were

between 0.14% and 1.22% on average across all species and between 0.03% and

1.99% per individual species. Differences between the two assemblies (tested within

or across all plant histories) were between 0.04% and 0.15% on average across all

species and between 0.03% and 0.35% per individual species. Thus, the fraction of

differentially methylated cytosines between the plant histories was generally larger

than differences between the two assemblies.

Within the plant histories, differences between the monoculture types and the naïve

plants were between 0.20% and 1.22% within species. Differences between mixture

types and naïve plants were between 0.13% and 1.27% within species. Differences

between monoculture and mixture types were between 0.03% and 1.99% within

species. However, if compared within each species separately, there were always

more DMCs in the comparisons between plants from Jena and the naïve plants than in

the comparison between monoculture and mixture types. The sole exception was *O.*

viciifolia for which there were more differences between the monoculture and mixture

types than between these and the naïve plants.

To further characterize the differences in DNA methylation, we calculated the

average change in DNA methylation at the DMCs for each contrast, across and within

all sequence contexts (CG, CHG, and CHH) and feature types (genes, transposons,

repeats, and unclassified) and visualized these differences (Fig. 5). We could not

identify clear patterns between the different comparisons with one exception:

differences in the comparisons between plants from Jena and the naïve plants within

genes (all sequence contexts) were mostly biased towards a higher methylation in the

naïve plants. Thus, plants in the Jena Experiment lost on average DNA methylation at

DMCs within genes (*O. viciifolia* was the exception with changes close to zero).

Discussion

We aimed at measuring to what extent differential selection in monoculture and

mixed species grassland communities had influenced the genetic and epigenetic

make-up of six grassland species and whether these mechanisms could have mediated

the rapid phenotypic responses that were previously observed in plants with different

diversity backgrounds (see Zuppinger-Dingley *et al.*, 2014; van Moorsel *et al.*,

2018b). Strong genetic divergence within species in response to selection history was

found in five out of six species, suggesting that in these five species different

genotypes may have been sorted out from standing genetic variation that was present

at the start of the diversity experiment. Less likely, recombination and mutation could

also have contributed.

For *P. lanceolata,* no significant differentiation was found between monoculture

and the mixture types within mixture assemblies. This may be due to the panmictic

nature of *P. lanceolata*. Movement of pollen and seeds between plots together with

high germination and establishment rates in this species may have leveled out genetic

differentiation between plots of different species diversity. Furthermore, coverage was

low and only a small proportion of the genome was covered, resulting in only 202

SNPs in the dataset. We therefore cannot exclude the possibility that genetic variation

was present but that our epiGBS screening missed the genomic regions under

selection. For future research, we propose that increasing the sequencing coverage

should be prioritized over adding more samples.

Only in *O. viciifolia* we found more genetic differences between the monoculture

and mixture types than between these and naïve plants. This may be due to the fact

that some subplots that were transplanted from the field site in Jena to the

experimental plots contained only few individuals of *O. viciifolia*. The results indicate

that *O. viciifolia* plant material originating from monoculture and mixture selection

history could have originated from a singly ancestor, which resulted in a high degree

of relatedness within each population. In contrast, the seeds sourced from the

commercial supplier likely originated from many mother plants.

Overall, only 1.4% to 10% of genetic variation was explained by plant histories.

However, for individual SNPs, plant histories could explain up to 67% of the genetic

variation. This may indicate that only few loci were under selection (high

differentiation) and that most of the genome segregated randomly (low

differentiation). Interestingly, this would suggest that the traits underlying the

differentiation within species into monoculture and mixture types are not highly

polygenic, thus making them accessible to breeding.

Our work supports the hypothesis that perennial grassland species in field

experiments evolve as a response to their biotic environment (Zuppinger-Dingley *et*

al., 2014; van Moorsel *et al.*, 2018b; van Moorsel *et al.*, 2018c). Recent experiments

showed stronger complementarity effects (Loreau & Hector 2001) in communities

with a selection history in species mixtures as opposed to plant individuals with a

selection history in monocultures (Zuppinger-Dingley *et al.*, 2014; van Moorsel *et al.*,

2018b). It has been under debate whether epigenetic or genetic mechanisms are the

main driver of observed phenotypic variation in such scenarios (Rapp & Wendel

2005; Bird 2007; Bossdorf *et al.*, 2008; Tilman & Snell-Rood 2014). In our

experiment, we found genetic and epigenetic variation. However, genetic variation

can have a major effect on epigenetic variation not only in *cis* but also in *trans* (Dubin

et al., 2015; Kawakatsu *et al.,* 2016). However, *cis* associations can also evolve

independently of underlying genetic variation, when spontaneous epimutations are

inherited through epigenetic inheritance (Taudt *et al.,* 2016). Thus, given the observed

and the additional unobserved (only parts of the genome were sequenced) genetic

variation, it seems plausible that most of the observed epigenetic variation was caused

by the underlying genetic variation. This is in line with quantitative genetics studies in

A. thaliana in which DNA methylation differences were found to be mostly associated

with underlying genetic variation (e.g., Li *et al.,* 2014).

We did, however, also observe methylation differences between the assemblies in

absence of genetic differentiation, even though these differences were small when put

into context. A recent study using the same statistical method but a more stringent

significance threshold (FDR < 0.001 instead of FDR < 0.01) compared DNA

methylation profiles of different tissue types of the liverwort *Marchantia polymorpha*,

covering important stages of the life cycle (Schmid *et al.,* 2018). With less than 1%

DMCs, the differences between the two assemblies of this study correspond to the

difference between two individual plants in Schmid *et al.* (2018). However, around

42% of all cytosines varied significantly across the entire life cycle in that study.

Thus, differences within individuals were much more pronounced than between

individuals. Several other studies provide direct or indirect evidence for clear

differences in DNA methylation between different tissue, cell types or developmental

stages (Jullien *et al.,* 2012; Calarco *et al.,* 2012; Ibarra *et al.,* 2012; Park *et al.,* 2016;

Ingouff *et al.*, 2017; Bouyer *et al.,* 2017). Clearly, this does not exclude the possibility

that epigenetics may play a role in adaptation (see Hauser *et al.,* 2011, Kronholm &

Collins 2016 and Quadrana & Colot 2016 for examples of natural epialleles), but it

makes it less likely that the strengthening of the biodiversity effect in the populations

assessed in this study was due to epigenetic differences that were not a consequence

of the underlying genetic differentiation.

578 Conclusions

We used novel genomic tools to study the importance of epigenetics for adaptation to

novel biotic conditions in six non-model plant species differentially selected in a field

biodiversity experiment. Future research should continue to integrate molecular tools

with studies on non-model species in order to push the field forward (Richards *et al*.,

2017; Heer *et al.*, 2018). Our findings suggest that selection on standing genetic

variation is a powerful driver of evolution even in the absence of many generations of

plant growth. In addition, we propose that community diversity had the selective

power to differentiate plant populations within species into mixture and monoculture

types in only a few generations. Molecular tools and the integration of evolutionary

concepts into plant community ecology can open up new alleys of research, which

should be exploited in order to understand the community evolutive processes (Shafer

et al., 2015) that lead to the plant community compositions and structures as we see

them today.

593 Acknowledgements

We thank T. Zwimpfer, M. Furler, D. Trujillo and D. Topalovic for technical

assistance. Thanks to T. Hahl for help with the glasshouse experiment. This study was

supported by the Swiss National Science Foundation (grants number 147092 and

166457 to B. Schmid) and the University Research Priority Program Global Change

and Biodiversity of the University of Zurich. S.J.V.M. was furthermore supported by

a travel grant from the ESF Congenomics network (reference number 7167).

601 Author contributions

S.J.V.M, P.V. and B.S. planned and designed the study, S.J.V.M. carried out the pot experiment and collected plant material, C.A.M.W. performed the lab work and created the sequencing library and T.V.G. created the bioinformatics pipeline. M.W.S. analyzed the data and produced the figures. S.J.V.M and M.W.S. wrote the first draft of the manuscript. All authors contributed to revisions.

References

Anderson JT, Willis JH, Mitchell-Olds T. 2011. Evolutionary genetics of plant adaptation. Trends in Genetics 27:258–266.

Barrett R, Schluter D. 2008. Adaptation from standing genetic variation. Trends in Ecology & Evolution 23:38–44.

Bastolla U, Fortuna MA, Pascual-García A, Ferrera A, Luque B, Bascompte J. 2009. The architecture of mutualistic networks minimizes competition and increases biodiversity. Nature 458:1018–1020.

Bird A. 2007. Perceptions of epigenetics. Nature 447:396–398.

Bossdorf O, Richards CL, Pigliucci M. 2008. Epigenetics for ecologists. Ecology letters 11:106–15.

Bouyer D, Kramdi A, Kassam M, Heese M, Schnittger A, Roudier F, Colot V. 2017. DNAmethylation dynamics during early plant life. Genome Biology 18:179.

Calarco JP, Borges F, Donoghue MTA, Van Ex F, Jullien PE, Lopes R, Gardner R, Berger F, Feijó JA, Becker JD. 2012. Reprogramming of DNA methylation in pollen guides epigenetic inheritance via small RNA. Cell 151:194–205.

Cardinale BJ, Wright JP, Cadotte MW, Carroll IT, Hector A, Srivastava DS, Loreau M, Weis JJ. 2007. Impacts of plant diversity on biomass production increase through time because of species complementarity. Proceedings of the National Academy of Sciences 104:18123–18128.

Cortijo S, Wardenaar R, Colomé-Tatché M, Gilly A, Etcheverry M, Labadie K, Caillieux E, Hospital F, Aury J-M, Wincker P, *et al*. 2014. Mapping the epigenetic basis of complex traits. Science 343:1145–1148.

Dubin MJ, Zhang P, Meng D, Remigereau MS, Osborne EJ, Paolo Casale F,
Drewe P, Kahles A, Jean G, Vilhjálmsson B, *et al.,***2015.** DNA methylation
in Arabidopsis has a genetic basis and shows evidence of local adaptation.
Elife 4:e05255.

Fakheran S, Paul-Victor C, Heichinger C, Schmid B, U. Grossniklaus U,
Turnbull L. 2010. Adaptation and extinction in experimentally fragmented
landscapes. Proceedings of the National Academy of Sciences 107:19120–
19125.

Fargione J, Tilman D, Dybzinski R, Lambers JHR, Clark C, Harpole WS, Knops
JM, Reich PB, Loreau M. 2007. From selection to complementarity: shifts in
the causes of biodiversity-productivity relationships in a long-term
biodiversity experiment. Proceedings of the Royal Society B: Biological
Sciences 274:871–876.

Gervasi DDL, Schiestl FP. 2017. Real-time divergent evolution in plants driven by
pollinators. Nature communication 8:14691.

Ghalambour CK, McKay JK, Carroll SP, Reznick DN. 2007. Adaptive versus
non-adaptive phenotypic plasticity and the potential for contemporary
adaptation in new environments. Functional Ecology 21:394–407.

Goudet, J, Raymond, M, Meeüs, T, Rousset, F. 1996. Testing differentiation in
diploid populations. Genetics 144:1933-1940.

Goudet J, Jombart T. 2015. hierfstat: Estimation and Tests of Hierarchical F-
Statistics. R package version 0.04-22. https://CRAN.R-
project.org/package=hierfstat

van Gurp TP, Wagemaker NCAM, Wouters B, Vergeer P, Ouborg JNJ,
Verhoeven KJF. 2016. epiGBS: reference-free reduced representation
bisulfite sequencing. Nature Methods 13:322–324.

Hairston NG, Ellner SP, Geber MA, Yoshida T, Fox JA. 2005. Rapid evolution
and the convergence of ecological and evolutionary time. Ecology Letters
8:1114–1127.

Hauser M-T, Aufsatz W, Jonak C, Luschnig C. 2011. Transgenerational epigenetic
inheritance in plants. Biochim Biophys Acta 1809:459–468.

Heer K, Mounger J, Boquete MT, Richards CL, Opgenoorth L. 2018. The
diversifying field of plant epigenetics. New Phytologist 217:988–992.

Ibarra CA, Feng X, Schoft VK, Hsieh T-F, Usawa R, Rodrigues JA, Zemach A,
Chumak N, Machlicova A, Nishimura T, *et al.*,2012. Active DNA
demethylation in plant companion cells reinforces transposon methylation in
gametes. Science 337:1360–1364.

Ingouff M, Selles B, Michaud C, Vu TM, Berger F, Schorn AJ, Autran D, Van
Durme M, Nowack MK, Martienssen RA, *et al.*,2017. Live-cell analysis of
DNA methylation during sexual reproduction in *Arabidopsis* reveals context
and sex-specific dynamics controlled by noncanonical RdDM. Genes &
Development 31:72–83.

Isbell F, Craven D, Connolly J, Loreau M, Schmid B, Beierkuhnlein C, Bezemer
TM, Bonin C, Bruelheide H, de Luca E, *et al.*,2015. Biodiversity increases
the resistance of ecosystem productivity to climate extremes. Nature 526:574–
577.

Jombart T. 2008. Adegenet: A R Package for the Multivariate Analysis of Genetic
Markers. Bioinformatics 24: 1403–5.

Joshi J, Schmid B, Caldeira MC, Dimitrakopoulos PG, Good J, Harris R, Hector
A, Huss-Danell K, Jumpponen A, Minns A *et al.*,2001. Local adaptation
enhances performance of common plant species. Ecology Letters 4:536–544.

Jost L. 2008. G_{ST} and its relatives do not measure differentiation. Molecular Ecology
17:4015–4026.

Jullien PE, Susaki D, Yelagandula R, Higashiyama T, Berger F. 2012. DNA
methylation dynamics during sexual reproduction in *Arabidopsis thaliana*.
Current Biology 22:1825–1830.

Kawakatsu T, Huang SC, Jupe F, Sasaki E, Schmitz RJ, Urich MA, Castanon R,
Nery JR, Barragan C, He Y, *et al.*,2016. Epigenomic Diversity in a Global
Collection of Arabidopsis thaliana Accessions. Cell 166:492-505.

Kleynhans EJ, Otto SP, Reich PB, Vellend M. 2016. Adaptation to elevated CO2 in
different biodiversity contexts. Nature Communications 7:12358.

Kooke R, Johannes F, Wardenaar R, Becker F, Etcheverry M, Colot V,
Vreugdenhil D, Keurentjes, JJB. 2015. Epigenetic basis of morphological
variation and phenotypic plasticity in *Arabidopsis thaliana*. Plant Cell 27:
337–348.

Kronholm I, Collins S. 2016. Epigenetic mutations can both help and hinder
adaptive evolution. Molecular Ecology 25: 1856–1868.

Li Y, Cheng R, Spokas KA, **PalmerAA, Borevitz JO. 2014.** Genetic variation for

life history sensitivity to seasonal warming in Arabidopsis thaliana. Genetics

196: 569–577.

Lipowsky A, Schmid B, Roscher C. 2011. Selection for monoculture and mixture

genotypes in a biodiversity experiment. Basic and Applied Ecology 12:360–

371.

Loreau M, Hector A. 2001. Partitioning selection and complementarity in

biodiversity experiments. Nature 412:72–76.

de Meeûs T, Goudet J. 2007. "A step-by-step tutorial to use HierFstat to analyse

populations hierarchically structured at multiple levels." *Infection, Genetics*

and Evolution 7: 731-735.

Meyer ST, Ebeling A, Eisenhauer N, Hertzog L, Hillebrand H, Milcu A, Pompe

S, Abbas M, Bessler H, Buchmann N, *et al.,*2016. Effects of biodiversity

strengthen over time as ecosystem functioning declines at low and increases at

high biodiversity. Ecosphere 7:e01619.

van der Maaten L, Hinton G. 2008. Visualizing data using t-SNE. Journal of

Machine Learning Research 9:2579–2605.

van der Maaten L. 2014. Accelerating t-SNE using tree-based algorithms. Journal of

machine learning research 15:3221–3245.

van Moorsel SJ, Hahl T, Petchey OL, Ebeling A, Eisenhauer N, Schmid B, Wagg

C. 2018a. Evolution increases ecosystem temporal stability and recovery from

a flood in grassland communities. bioRxiv.

van Moorsel SJ, Schmid MW, Hahl T, Zuppinger-Dingley D, Schmid B. 2018b.

Selection in response to community diversity alters plant performance in and

functional traits. bioRxiv.

van Moorsel SJ, Hahl T, Wagg C, De Deyn GB, Flynn DFB, Zuppinger-Dingley

D, Schmid B. 2018c. Community evolution increases plant productivity at

low diversity. Ecology Letters 21:128–137.

Niederhuth CE, Bewick AJ, Ji L, Alabady MS, Kim KD, Li Q, Rohr NA,

Rambani A, Burke JM, Udall JA, *et al.,*2016. Widespread natural variation

of DNA methylation within angiosperms. Genome Biology 17:194.

Ouborg NJ, Vergeer P, Mix C. 2006. The rough edges of the conservation genetics

paradigm for plants. Journal of Ecology 94:1233–1248.

Park K, Kim MY, Vickers M, Park J-S, Hyun Y, Okamoto T, Zilberman D,
Fischer RL, Feng X, Choi Y, *et al.,*2016. DNA demethylation is initiated in
the central cells of *Arabidopsis* and rice. PNAS 113:15138–15143.

Park Y, Wu H. 2016. Differential methylation analysis for BS-seq data under general
experimental design. Bioinformatics 32: 1446–1453.

Price TD, Qvarnstrom A, Irwin DE. 2003. The role of phenotypic plasticity in
driving genetic evolution. Proceedings of the Royal Society B: Biological
Sciences 270:1433–1440.

Quadrana L, Colot V. 2016. Plant transgenerational epigenetics. Annual Review of
Genetics 50:467–491.

Rapp RA, Wendel JF. 2005. Epigenetics and plant evolution. New Phytologist
168:81–91.

Reich PB, Tilman D, Isbell F, Mueller K, Hobbie SE, Flynn DFB, Eisenhauer N.
2012. Impacts of biodiversity loss escalate through time as redundancy fades.
Science 336:589–592.

Richards CL, Alonso C, Becker C, Bossdorf O, Bucher E, Colomé-Tatché M,
Durka W, Engelhardt J, Gaspar B, Gogol-Döring A, *et al.,*2017.
Ecological plant epigenetics: Evidence from model and non-model species,
and the way forward. Ecology Letters 20:1576–1590.

Roscher C, Schumacher J, Baade J, Wilcke W, Gleixner G, Weisser WW,
Schmid B, Schulze E-D. 2004. The role of biodiversity for element cycling
and trophic interactions: an experimental approach in a grassland community.
Basic and Applied Ecology 5:107–121.

Rottstock T, Kummer V, Fischer M, Joshi J. 2017. Rapid transgenerational effects
in *Knautia arvensis* in response to plant community diversity. Journal of
Ecology 105:714–725.

Schmid B. 1985. Clonal Growth in Grassland Perennials: III. Genetic Variation and
Plasticity Between and Within Populations of Bellis Perennis and Prunella
Vulgaris. Journal of Ecology 73:819–830.

Schmid MW. 2017. RNA-Seq data analysis protocol: Combining in-house and
publicly available data. In: Plant germline development: Methods and
protocols. Schmidt A, editor. New York, NY: Springer New York; 2017. pp.
309–35.

Schmid MW, Giraldo-Fonseca A, Rövekamp M, Smetanin D, Bowman JL,
Grossniklaus U. 2018. Extensive epigenetic reprogramming during the life
cycle of *Marchantia polymorpha*. Genome Biology 19:9.

Schoener TW. 2011. The newest synthesis: Understanding the interplay of
evolutionary and ecological dynamics. Science 331:426–429.

Shafer ABA, Wolf JBW, Alves PC, Bergström L, Bruford MW, Brännström I,
Colling G, Dalén L, De Meester L, Ekblom R. *et al.*,2015. Genomics and the
challenging translation into conservation practice. Trends in Ecology &
Evolution 30:78–87.

Slobodkin LB. 1961. Growth and regulation of animal populations. Holt, Rinehart
and Winston, New York.

Smit AFA, Hubley R, Green P. 2013–2015. RepeatMasker Open-4.0.
http://www.repeatmasker.org/.

Taudt A, Colomé-Tatché M, Johannes F. 2016. Genetic sources of population
epigenomic variation. Nature Reviews Genetics 17: 319–332.

Tilman D, Snell-Rood EC. 2014. Ecology: Diversity breeds complementarity.
Nature 515:44–45.

Verhoeven KJF, vonHoldt BM, Sork VL. 2016. Epigenetics in ecology and
evolution: what we know and what we need to know. Molecular Ecology
25:1631–1638.

Wilschut RA, Oplaat C, Snoek LB, Kirschner J, Verhoeven KJF. 2016. Natural
epigenetic variation contributes to heritable flowering divergence in a
widespread asexual dandelion lineage. Molecular Ecology 25:1759–1768.

Zuppinger-Dingley D, Schmid B, Petermann JS, Yadav V, De Deyn GB, Flynn
DFB. 2014. Selection for niche differentiation in plant communities increases
biodiversity effects. Nature 515:108–111.

Data Accessibility Statement

Supporting information

Additional Supporting Information may be found online in the Supporting
Information tab for this article:

Fig. S1 Adapter overview.

Fig. S2 Results from the G-statistic tests given all SNPs within genes.

Fig. S3 Results from the G-statistic tests given all SNPs within transposons.

Fig. S4 DNA methylation levels in % at individual cytosines across all or within each

individual sequence context (CG, CHG, CHH) for each experimental group of each

species used.

Table S1 Sample overview.

Table S2 Community diversity and composition of the plots the seeds originated

from.

Table S3 Reference sequences generated in this study.

Table S4 Average pairwise F_{ST} values.

Table S5 99^{th} percentiles of F_{ST} values in the SNPs within genes.

Table S6 99^{th} percentiles of F_{ST} values in the SNPs within transposons.

Table S7 Number of cytosines with significant differences (FDR < 0.01) in DNA

methylation between selection-history treatments and assemblies in the CG sequence

context.

Table S8 Number of cytosines with significant differences (FDR < 0.01) in DNA

methylation between selection-history treatments and assemblies in the CHG

sequence context.

Table S9 Number of cytosines with significant differences (FDR < 0.01) in DNA

methylation between selection-history treatments and assemblies in the CHH

sequence context.

Fig. 1 (a) The origin of seeds used for the glasshouse experiment and genetic analysis.

Seedlings were planted in mixtures and monocultures in Jena in the year 2002. Two

reproduction events occurred when seeds were collected and subsequently new

seedlings were produced and planted again in the same community composition. (b)

Schematic representation of the glasshouse experiment. Monoculture assemblies and

2-species mixture assemblies were planted with either plants with mixture selection

history (green), monoculture selection history (orange) or naïve plants originating

from a commercial seed supplier (blue). Figure modified after van Moorsel *et al.*

(2018b).

Fig. 2 Genetic distance between individuals of the different populations for the six

species. (a) All loci included in the analyses. (b) Only the 5% most differentiated loci

between the experimental groups included. Green: selection history in mixture,

orange: selection history in monocultures, blue: naïve selection history. Triangles:

current assembly monoculture, circles: current assembly mixture. $D_{95\%}$ is the

threshold to choose the 5% most differentiated loci based on Jost's D (the 95%

percentile in the entire data set, Jost 2008).

Fig. 3 Results from the G-statistic tests given all SNPs. Each panel shows a histogram
of permuted test statistics (999 permutations) and indicates the observed statistics by a
black dot and a segment. Test statistics are on the x-axis, frequencies on the y-axis.
Grey boxes where data was not available (experimental group missing).

Fig. 4 (a) DNA methylation levels in percent at individual cytosines across all or

within each individual sequence context (CG, CHG, CHH) for each species used in

this study shown as violin plots. The horizontal black bars correspond to the means.

(b) Average DNA methylation levels in percent for each sequence context, genomic

feature, and species shown as a heat map.

Fig. 5 Average differences in DNA methylation at significantly differentially

methylated cytosines (DMCs; FDR < 0.01) within a given sequence (all, CG, CHG,

and CHH) and feature (all, genes, transposons, repeats, unclassified) context are

shown for all contrasts. The average differences are shown as colour gradient. The

numbers within the heat map are the average differences. The asterisk marks the rows

showing that plants in the Jena field lost on average DNA methylation at DMCs

within genes compared to naïve plants.

Table 1 99th percentile of FST values in the data set with all SNPs.

	G. mollugo	*L. pratensis*	*O. viciifolia*	*P. lanceolata*	*P. vulgaris*	*V. chamaedrys*
SH in monoculture	NA	0.10641	NA	0.1413	0.6032	0.29453
SH in mixture	0.2354	NA	0.4	NA	0.67471	NA
SH contrast mono vs. mix in monoculture	0.1579	0.1	NA	0.09282	0.25	0.2571
SH contrast mono vs. mix in mixture	0.13332	0.1462	0.3333	0.05899	0.15592	0.3333
AS for naïve history	NA	NA	NA	NA	0.13182	NA
AS for mono history	0.0755	0.0909	0.08469	0.0762	0.10173	0.14305
AS for mix history	0.08355	0.0926	NA	0.09214	0.12	0.1586

865   AS = assembly, SH = selection history. For SNPs within genes or transposons see

866   Tables S5 and S6.

 Table 2 Number of cytosines with significant differences (FDR < 0.01) in DNA

 methylation between selection-history treatments and assemblies.

	G. mollugo	*L. pratensis*	*O. viciifolia*	*P. lanceolata*	*P. vulgaris*	*V. chamaedrys*	average %
SH: mixture vs. monoculture	6365 (0.55%)	239 (0.03%)	-	582 (0.11%)	6578 (0.28%)	12966 (0.60%)	0.31%
% in genes	13.68	4.6		4.47	12.82	11.51	
% in transposons	10.65	30.54		8.25	11.28	8.18	
% in repeats	4.41	4.18		8.93	6.67	4.63	
% in unclassified contigs	71.25	60.67		78.35	69.23	75.68	
>> within monoculture assembly	2332 (0.20%)	360 (0.05%)	-	216 (0.04%)	2507 (0.11%)	6507 (0.30%)	0.14%
% in genes	12.22	4.72		6.48	10.33	10.7	
% in transposons	11.11	28.61		7.87	11.61	8.16	
% in repeats	3.82	4.72		12.04	5.78	4.84	
% in unclassified contigs	72.86	61.94		73.61	72.28	76.3	
>> within mixture assembly	4617 (0.40%)	288 (0.04%)	10353 (1.99%)	2573 (0.47%)	8875 (0.37%)	5538 (0.25%)	0.59%
% in genes	11.63	4.17	12.13	3.26	11.14	9.73	
% in transposons	10.92	34.38	15.94	11.31	14.6	8.22	
% in repeats	4.7	5.21	6.93	7.5	6.49	4.66	
% in unclassified contigs	72.75	56.25	65.01	77.92	67.76	77.39	
SH: mixture vs. naïve	-	-	-	-	28970 (1.21%)	-	(1.21%)
% in genes					9.69		
% in transposons					11.84		
% in repeats					6.73		
% in unclassified contigs					71.73		
>> within monoculture assembly	-	893 (0.13%)	-	1040 (0.19%)	18439 (0.77%)	14692 (0.68%)	0.44%
% in genes		4.26		7.69	9.75	10.22	
% in transposons		26.32		9.71	12.55	7.83	
% in repeats		4.82		5.1	6.84	4.27	
% in unclassified contigs		64.61		77.5	70.87	77.68	
>> within mixture assembly	7012 (0.61%)	-	6574 (1.27%)	-	15019 (0.63%)	-	0.84%
% in genes	11.52		12.64		9.54		
% in transposons	10.95		16.06		11.85		
% in repeats	4.08		6.77		6.78		
% in unclassified contigs	73.45		64.53		71.83		
SH: monoculture vs. naïve	-	-	-	-	29052 (1.22%)	-	(1.22%)
% in genes					9.25		
% in transposons					12.52		
% in repeats					6.83		
% in unclassified contigs					71.4		
>> within monoculture assembly	-	1388 (0.20%)	-	620 (0.11%)	15877 (0.67%)	13614 (0.63%)	0.40%
% in genes		5.04		6.29	9.01	10.13	

	C1	C2	C3	C4	C5	C6	C7
% in transposons		28.6		7.1	12.36	7.28	
% in repeats		6.63		8.06	7.26	4.73	
% in unclassified contigs		59.73		78.55	71.37	77.86	
>> within mixture assembly	5336 (0.46%)	-	5464 (1.05%)	-	18419 (0.77%)	-	0.76%
% in genes	10.91		11.79		9.98		
% in transposons	9.07		15.98		12.24		
% in repeats	3.77		6.9		6.59		
% in unclassified contigs	76.26		65.34		71.19		
AS: mixture vs. monoculture	-	-	-	-	937 (0.04%)	-	(0.04%)
% in genes					8		
% in transposons					11.31		
% in repeats					8.86		
% in unclassified contigs					71.82		
>> within naïve selection history	-	-	-	-	2988 (0.13%)	-	(0.13%)
% in genes					9.64		
% in transposons					14.63		
% in repeats					7.83		
% in unclassified contigs					67.9		
>> within monoculture selection history	948 (0.08%)	191 (0.03%)	216 (0.04%)	506 (0.09%)	2264 (0.09%)	1588 (0.07%)	0.06%
% in genes	9.07	7.85	8.8	10.87	7.91	5.79	
% in transposons	10.34	31.94	18.52	6.92	14.31	12.22	
% in repeats	5.38	3.14	3.7	11.86	7.99	6.11	
% in unclassified contigs	75.21	57.07	68.98	70.36	69.79	75.88	
>> within mixture selection history	4032 (0.35%)	708 (0.10%)	-	944 (0.17%)	1857 (0.08%)	1624 (0.07%)	0.15%
% in genes	6.92	4.38		3.6	7.7	9.3	
% in transposons	12.7	29.66		11.55	16.05	8.37	
% in repeats	6.42	6.36		9	6.41	5.11	
% in unclassified contigs	73.96	59.6		75.85	69.84	77.22	
Total (percentage DMCs of tested cytosines)	21086 (1.8%)	3291 (0.5%)	16544 (3.2%)	5122 (0.9%)	65507 (2.7%)	36076 (1.7%)	1.80%

870   AS = assembly, SH= selection history. For data on separate sequence contexts see

871   Tables S7 (CG), S8 (CHG), and S9 (CHH).