Rapid evolution allows plant communities to regain stability after simulated species loss

Sofia J. van Moorsel[1,2]*, Terhi Hahl[1], Owen L. Petchey[1], Anne Ebeling[3], Nico Eisenhauer[4,5], Bernhard Schmid[1,6] and Cameron Wagg[1,7]

[1]Department of Evolutionary Biology and Environmental Studies, University of Zürich, Winterthurerstrasse 190, CH-8057, Switzerland
[2] Department of Biology, McGill University, 1205 Dr Penfield Ave, Montreal, QC, Canada
[3]Institute of Ecology and Evolution, University of Jena, Dornburger Str. 159, 07743 Jena, Germany
[4]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany
[5]Institute of Biology, Leipzig University, Deutscher Platz 5e, 04103 Leipzig, Germany
[6]Department of Geography, University of Zürich, Winterthurerstrasse 190, CH-8057, Switzerland
[7]Agriculture and Agri-Food Canada, 850 Lincoln Road, Fredericton, New Brunswick, Canada

*To whom correspondence should be addressed.
Email: sofia.vanmoorsel@ieu.uzh.ch

E-mails of authors:
terhi.hahl@gmail.com
owen.petchey@ieu.uzh.ch
nico.eisenhauer@idiv.de
anne.ebeling@uni-jena.de
bernhard.schmid@uzh.ch
cameron.wagg@canada.ca

Author contributions:
B.S. designed research, T.H. and S.J.V.M. performed research; A.E. and N.E. maintained and coordinated the field site; S.J.V.M., C.W. and B.S. analyzed data; S.J.V.M., B.S., O.L.P. and C.W. wrote the paper.

Keywords: asynchrony | co-occurrence history | disturbance | grassland biodiversity | recovery | resilience | resistance | ecosystem stability

Abstract

Experiments simulating species loss from grassland ecosystems have shown that

losing biodiversity decreases the ability of ecosystems to buffer disturbances.

However, plant or plant-soil evolutionary processes may allow ecosystems to regain

stability and resilience over time. We explored such effects in a long-term grassland

biodiversity experiment. Low diversity communities of plants with a history of co-

occurrence (selected communities) were temporally more stable than the same

communities of plants with no such history (naïve communities). Furthermore,

selected communities showed greater recovery following a major flood, resulting in

more stable post-flood productivity. These results were consistent across soil

treatments simulating the presence or absence of co-selected microbial communities.

We suggest that plant evolution in a community context can increase ecosystem

temporal stability and resistance to disturbances. Evolution can thus in part

compensate for extreme species loss as can high plant diversity in part compensate for

the missing opportunity of evolutionary adjustments.

Biodiversity experiments simulating the loss of plant species from grassland

communities have shown that less diverse communities have reduced mean [1,2] and

increased variation of aboveground biomass [3–5]. However, it is not clear whether these

communities may regain functioning and stability over time while still being at low

diversity. The few biodiversity experiments that lasted more than 10 years showed

that functioning tended to decrease in low-diversity communities and to increase in

high-diversity communities, leading to an increased slope of the biodiversity–

productivity relationship over time [6,7]. In one of these experiments, the Jena

Experiment in Germany [7], it was shown that divergent evolutionary changes of plant

species in monocultures vs. mixtures during the first 8 years contributed to this

strengthening of the biodiversity–functioning relationship [8–10]. Feedbacks between

plants and soil organisms, however, had less explanatory power [9,11].

Ecosystem resistance, recovery and resilience that underlie stability may

depend on plant diversity [12–14]. The mechanisms by which diversity stabilizes

ecosystem productivity are based on differences between genotypes or species in their

responses to the abiotic or biotic environment [3,5,15,16]. For instance, different plant

species may exhibit high performance under different environmental conditions

(termed response diversity [17]). Asynchrony of species performances, derived from

interspecific differences in responses to environmental variation, can thus allow

diverse communities to resist disturbance or recover to maintain performance, often

referred to as insurance or portfolio effect [5,16,18,19].

The development of stability over time in the long-term biodiversity

experiments had not been analyzed so far, but using data from the Jena Experiment

[20,21], we found that combined intra- and inter-annual variation in experimental

communities decreased over time for the first 8 years (i.e. ecosystem stability,

measured as inverse of coefficient of variation of total biomass, increased over time;

Fig. 1), suggesting that temporal stability could be regained. As new communities

develop following sowing, species abundance distributions and gene frequencies can

change, and such adjustments between and within species may explain increased

stability [22]. For example, a prolonged period of co-occurrence increased "stabilizing

differences" between phylogenetically distinct annual plant species in comparison

with similar pairs of species without co-occurrence history [23]. However, it has not

been tested yet if similar effects of co-occurrence history might already appear during

the course of biodiversity experiments. This is because in such experiments,

individual plants age and take up nutrients which may also affect stability besides

effects of co-occurrence history. In addition, it has proven difficult to disentangle the

contribution of soil organisms that over time associate with the plant communities.

The interplay between mutualistic and pathogenic soil organisms (bacteria, fungi,

invertebrates) and their impact on plant communities could have (de)stabilizing

effects by changing nutrient provision and the plant's health [24,25].

To test specifically for the importance of co-occurrence history we therefore

compared communities of plants of the same age but with or without co-occurrence

history on soils with or without soil organisms that had become associated with the

plants with co-occurrence history (Fig. 2). *Selected* communities were assembled with

offspring from individuals that had co-occurred in the same plots of the Jena

Experiment over 8 years. *Naïve* communities were assembled with offspring from

seeds that were obtained from the same supplier that had provided the original seeds

for the Jena Experiment. We then asked whether (1) selected plant communities had

more stable productivity than naïve plant communities, (2) potential differences in

stability between naïve and selected communities depended on plant diversity and (3)

stability was further increased when plants were growing with their native soil

organisms. Test communities ranged in species richness from one to two (low

diversity) to four and eight species (medium diversity). We assessed the combined

intra- and inter-annual variation in the test communities across seven seasonal

biomass harvests from 2012–2015 (Fig. S1).

A natural flood event [26] after the third harvest gave us the opportunity to

additionally analyze the resistance, recovery and resilience [27–29] of test communities in

response to the flood and compare the pre- with the post-flood stability using the first

three and the last three harvests, respectively. Together, resistance and recovery

determine ecosystem resilience as we define it here, namely how ecosystem

productivity differs between pre- and post-disturbance states [30].

Finding a more stable productivity for selected communities (question (1))

would provide evidence for short-term plant evolutionary processes increasing

ecosystem stability. So far, evolutionary mechanisms underlying ecosystem stability

in biodiversity experiments have only been studied in terms of phylogenetic

relatedness that reflects evolutionary processes over long time scales with conflicting

results [31,32]. Short-term evolution leading to changes at the community level, referred

to as community evolution ([9], have almost exclusively been studied in microbial

ecosystems [33–36]. However, short-term evolutionary processes could be particularly

relevant in plant communities facing rapid global change [37,38], including increased

extreme climatic events, and for conservation and restoration approaches [39], because

plants are fixed in place and can only move by propagule dispersal.

Question (2) is based on the observation that evolution in a community context

in the Jena Experiment increased niche differentiation between species in mixtures [8].

This could potentially affect species asynchrony and average population variability,

which in turn contribute to ecosystem stability. The increasing slope of biodiversity–

productivity relationships in long-term biodiversity experiments could suggest that

biodiversity–stability relationships also increase and thus differences between plant

communities with and without co-occurrence history should be larger at higher

species richness levels. Alternatively, it could be argued that such differences

reflecting the regaining of functioning and stability after species loss should be more

pronounced in plant communities that lost more species, i.e. communities of low

diversity.

Question (3) addresses the possibility that plant–soil feedback dynamics,

including both mutualistic and pathogenic effects [40,41], contributed to the increased

stability of experimental plant communities during the 8 years of the Jena Experiment

[42]. Soil microbial communities with a shared history of the respective plant

community may be more efficient in using plant-derived substrates, providing

nutrients to plant growth and protect plants against soil-borne pathogens [25]. Moreover,

a higher soil biodiversity is likely to provide nutrients at a higher and more stable

level [e.g. 43].

Given our experimental design, question (1) relates to main effects of plant co-

occurrence history, question (2) to the interaction between co-occurrence history and

plant species richness and question (3) to main effects of soil organisms (native vs.

neutral). Potential interactions between effects of soil organisms and plant co-

occurrence history were also tested but never significant. We first present the results

for the overall temporal stability of test communities and then those for the response

to the flood event.

Results

Community and population stability and species asynchrony

We used the inverse of the combined intra- and interannual variation over six

biomass harvests from spring 2012–spring 2015 (excluding the harvest taken directly

after the flood event in summer 2013) as stability measure of test communities.

Whereas the test communities showed the usual strong increase of stability with

species richness, overall differences between co-occurrence history and especially soil

treatments were small and insignificant (Table 1). However, at low diversity, selected

communities were more stable than naïve communities, reflected by a significant co-

occurrence history x species richness interaction (Table 1; Fig. 3A). This shows that

communities of low diversity regained more stability 8 years after simulated species

loss in the Jena Experiment. Population stability decreased with species richness but

at low diversity population stability was also higher in selected communities (Table 1;

Fig. 3B). In contrast, species asynchrony increased for both selected and naïve

communities in parallel with increasing species richness (Table 1; Fig. 3C). When we

corrected community stability and species asynchrony for all model terms except co-

occurrence history (i.e. taking residuals after fitting the plot x soil treatment

interaction), selected communities were more stable than naïve communities for any

given level of asynchrony (Fig. 3D).

Responses to flood event

A naturally occurring flood in early summer 2013 had devastating effects on

the plant communities and reduced productivity in the corresponding time interval [44];

Fig. 4). Diverse test communities, especially selected communities, lost the greatest

amount of biomass, i.e. showed the least resistance to the flood. When we adjusted for

community biomass before the flood, selected communities showed higher resistance

at low diversity and lower resistance at high diversity compared with the naïve

communities (Table S1; Fig. S2A). As for temporal stability, selected communities

simulated to have lost more species (monocultures) again regained more resistance.

Communities were most resistant in non-sterilized native soil. Comparing the

two other soil treatments, the plant communities were more resistant in sterilized soil

inoculated with native soil organisms than in sterilized soil inoculated with neutral

organisms (Table 2; Fig. 4, 5A). Here, an adjustment for pre-flood biomass made soil

treatments insignificant (Table S1; Fig. S2A).

Recovery of community biomass after the flood increased with species

richness and was higher in selected than in naïve communities across all diversity

levels and soil treatments (Table 2 and S1; Fig. 5B and S2B). Selected communities

had regained more resilience than the naïve communities, particularly at low diversity, as shown by the comparison of community biomass before and after recovery. However, the effect of co-occurrence was only significant if adjusted for pre-flood community biomass (Table 2 and S1; Fig. 5C and S2C). The three soil treatments strongly differed in their resilience, which averaged out their pre-flood differences in community biomass (Table 2; Fig. 3B). The water moving across plot borders during the flood may also have transported soil organisms between different soil treatments, further diluting soil treatment effects (see Fig. 2).

We then compared the combined intra- and interannual stability over the first three harvests before the flood event (2012–2013, Fig. 6A) with the last three harvests after recovery (2014–2015, Fig. 6B). The increased stability of selected communities was mainly due to higher stability after the flood. Furthermore, the greater regain of overall stability in selected communities of low diversity was due to the time before the flood, when selected communities of high richness had even lower stability than naïve communities of high richness (Fig. 6A). After the flood, when test communities had already been developing for more than two years, there was still a consistent difference across all diversity levels with selected communities being more stable than naïve communities. This suggests that these differences may have a heritable basis or, alternatively, that initial differences between selected and naïve test communities were accentuated by species abundance shifts within them. We therefore assessed species turnover rates with the Bray-Curtis similarity measure between pre- and post-flood species compositions but found that it was not influenced by co-occurrence history or soil treatments, although it increased with species richness (Table S3; Fig. S3).

Discussion

Biodiversity experiments test how simulated species loss affects ecosystem functioning and stability and, if run over longer time periods, can address the question whether both functioning and stability can be regained over time. In the Jena Experiment [45], a typical central European grassland ecosystem of 60 plant species was reduced to experimental communities containing 16, 8, 4, 2 or 1 species, which led to decreases in aboveground biomass production and its stability over time [21]. Strengthening biodiversity–productivity relationships observed in long-term biodiversity experiments [e.g. 6,46] indicate a deterioration of monocultures and low-

diversity mixtures and regaining of functioning in more diverse mixtures [7,46]. We

previously found that, in comparison with naïve communities, selected low-diversity

mixtures regained more functioning than mixtures with higher diversity [9]. Here, we

show that selected communities from the Jena Experiment also regained stability in

comparison with naïve communities, particularly the experimental communities

reflecting more extreme species loss.

Species-poor selected communities regained temporal stability

236   Selected communities had increased temporal stability of aboveground

237   biomass production but only at the lower levels of diversity up to four species. At the

238   eight-species richness level, we did not detect any differences as all communities

239   showed the highest stability values. This mirrors earlier findings for productivity

240   where biomass was similar for naïve and selected communities at the eight-species

241   richness level [9].

242   Several mechanisms could have led to the increased stability of selected

communities. For mixtures, the changes at the community level could have been

solely caused by altered species abundance distributions [47], driven by changed

performances of species due to evolution in their particular communities (Zuppinger-

Dingley et al. 2014). However, we did not detect any changes in species abundance

distributions between selected and naïve communities. A more likely explanation

therefore is that changes in genotype frequencies within species, i.e. evolution in the

community context (Strauss et al. 2006), led to increased temporal stability, which

could also explain the increased stability of selected monocultures. For five of the 60

species from the Jena Experiment the potential for such rapid evolutionary changes

and their genetic basis, with consequential epigenetic and phenotypic changes has

been confirmed [10]. The changes in genotype frequencies within species in selected

communities could be attributable to differential mortality, growth or reproduction

among the initially sown genotypes [48], recombination during sexual reproduction or,

least likely, to mutation. Rapid evolutionary changes in systems of interacting species,

referred to as community evolution [9], have so far mainly been observed in microbial

communities [33–36,49], but are here shown to also occur in communities of perennial

plant species.

260   Temporal stability at the community level in grassland ecosystems can be

261   driven by asynchronous population dynamics of species, allowing high compensatory

262   population variation to be combined with low community-level variation over time
263   [5,18,50–52]. As shown before [e.g. 4], we found that population stability decreased with
264   increasing diversity.

265   We further observed that, for any given level of asynchrony, selected
communities were more stable than naïve communities (see Fig. 3D). The greater
community stability at the same level of asynchrony suggests that evolution in
communities reduced competitive interactions between plant species, consistent with
the findings of evolutionary niche differentiation [8] and increased facilitation [53] among
species in mixtures in the Jena Experiment. By extension, similar evolutionary
processes may have occurred between genotypes within monocultures, again
consistent with previous findings showing evolutionarily changed phenotypic
variation within monocultures after eight years of selection in the Jena Experiment [9].

274   The increased stability over the first eight years of ecosystem development in
the Jena Experiment (see Fig. 1) could also have been due to the selection of soil
communities associated with the plants [41,54–56]. Therefore, we grew the selected and
naïve plant communities in three soil treatments: native soil of the selected
communities, sterilized soil inoculated with native soil organisms or sterilized soil
inoculated with neutral soil organisms collected outside the Jena Experiment.
Previously, highly significant differences between these soil treatments regarding
microbial community composition have been demonstrated [9,11], Marc W. Schmid
pers. comm. June 2019). However, contrary to our expectation, there were no
differences in plant community temporal stability between the three soil treatments
and no interactions of these with plant co-occurrence history. Thus, we conclude that
the changes in ecosystem stability observed in the Jena Experiment were not primarily
driven by plant–soil feedbacks but more likely due to plant evolution. Still, we cannot
rule out the possibility that the greater stability of selected communities was also in
part because of more stable associations with both native and neutral soil organisms.
Indeed, plants selected for eight years in monocultures in the Jena Experiment
developed positive plant–soil feedbacks [56], which may in part have contributed to
their increased stability in the present study.

Selected communities recover better from an extreme event than naïve communities

294   Diverse communities are more stable in the face of disturbance [13], for example
when they are hit by an extreme climatic event, such as a flood as happened to our test

communities halfway through the experiment in June 2013 [44]. Considering predicted

future climate scenarios with increased frequency of extreme events [57], including

floods [58], this aspect of stability may be even more relevant than temporal stability

under unperturbed conditions [59]. Further, both temporal stability and resistance to

perturbations need to be considered because they may not always be positively

correlated [60].

There are several ways in which a community may retain stability through

perturbations. Greater resistance increases ecosystem stability by reducing the loss in

productivity [61], while greater recovery increases the amount of productivity that the

ecosystem can regain after the disturbance-induced loss [28]. Together, resistance and

recovery then determine ecosystem resilience [30]. Consequently, how plant diversity

stabilizes ecosystem productivity through mediating ecosystem resistance, recovery

and resilience has become a focal question in ecology [13,51,62,63].

In our experiment, diversity reduced ecosystem resistance, in line with

previous findings for example in micro-ecosystems with ciliates responding to

warming [60]. This was because diverse communities had more biomass before the

flood and thus could also lose more, a result found previously for community

responses to drought [12,28,64]. For example, Wang et al. (2007) showed that at the same

diversity level, communities with higher productivity were more susceptible to an

experimental drought treatment. Here, when correcting for differences in pre-flood

biomass, selected communities were more resistant than naïve communities at low

diversity. At 8-species richness, the opposite pattern was found and naïve

communities were more resistant than selected communities (see Fig. S2A). Diverse

communities made up for their reduced resistance by increased recovery, as typically

found in biodiversity experiments [28,30], but, remarkably, selected communities showed

greater recovery than naïve communities along the entire species-richness gradient.

Consequently, at low diversity selected communities regained resilience, which was

not the case at higher diversity (see Fig. 4C). Changes in genotype frequencies during

community development in the first eight years of the Jena Experiment may not only

have led to increased community productivity [9] and temporal stability (see above) but

also to increased ecosystem stability in the face of disturbance, i.e. the flood event.

Key here is the adaptation to the abiotic conditions at the Jena field site, which is a

natural floodplain, where the plant communities were exposed to soil moisture

saturation at previous milder flood events in winter 2003 and winter 2005 (personal

communication with C. Roscher). Selection thus likely favored individuals with traits that allowed them to perform well under such conditions and to recover more rapidly [65,66].

However, species interactions may also have played a role for adaptation to flooding, as selected communities were more resilient at lower compared with higher diversity. Evolutionarily increased facilitation, demonstrated for these plants in the Jena Experiment [53], may in part have caused the increased resilience. This would be in line with predictions that environmental stress might select for greater facilitative effects in plant communities (stress-gradient hypothesis; [67]. Furthermore, such effects, which could be interpreted as diffuse co-evolution [68], are more likely to occur in communities of few interacting species [9,69]. This could additionally explain why selected communities regained more temporal stability and resilience at lower than at higher diversity.

Further, we tested if the association with native soil organisms over time might have affected the stability of plant communities in response to the flood. We found that pre-flood productivity was lowest in communities grown in native soil, higher in soil with native inoculum, and highest in soil with neutral inoculum (see Fig. 4B). This could have two reasons: first, a greater density of antagonistic soil biota in native and native-inoculated soils [70], or, second, a greater pool of available soil resources resulting from the soil sterilization process in the two inoculated soils [71]. Consequently, resistance was lowest for communities growing in native soil. In contrast, it seems that native soil organisms did increase recovery and resilience of plant communities to the flood event. First, by reducing pre-flood community biomass and thus cutting losses. However, previous studies showed that plants evolving under drought can select for a different set of microbial organisms than when evolving under wet conditions [55], which offers an alternative mechanism that selected communities may have associated with a better suited native soil community over time.

After the flood, the effect of soil treatments on plant community biomass was insignificant. In part, this might have been caused by an exchange of soil organisms during the flood event. However, differences between the soil treatments in terms of microbial community composition remained highly significant [11], Marc Schmid pers. comm. June 2019), which makes it more likely that the flood had an equalizing effect due to the differential resistance, recovery and resilience among the soil treatments.

After disturbance selected communities are more stable than naïve communities

As a third aspect of stability, we compared the combined intra- and inter-

annual ecosystem stability for the three harvests before and after recovery from the

flood separately [65]. Whereas the differences between selected and naïve communities

were not significant before the flood, selected communities showed increased post-

flood stability (see Table S2 and Fig. 6). This shows first that the effect of co-

occurrence history was long-lasting, i.e. at least for two years after re-establishment of

the test communities, and second that naïve communities could not make up for the

lack of co-occurrence history from spring 2011 to summer 2013. Furthermore, the

finding suggests that genotype frequencies need time to change and potential sorting

out does not happen immediately after sowing genetically diverse species collections.

Finally, selected communities of all diversity levels showed increased post-flood

stability, but selected communities at lower diversity levels regained more overall

temporal stability. This suggests that in the longer term also the more diverse

communities will regain stability.

It is important to note that the greater post-flood stability of selected

communities in our study was not driven by compositional changes as shown by the

similar compositional turnover between selected and naïve communities. We suggest

that genetic changes within the species populations comprising the selected

communities were responsible for the increased stability [10]. Independent of the

underlying mechanism, our study demonstrates that experimental plant communities

with simulated species loss over time can regain some of the temporal stability and

capacity to buffer extreme events, and this is likely due to plant evolutionary changes

in a community context [22], with few additional effects of assembling soil

communities. Our results can be viewed from two different angles. First, the greater

level of regained stability in communities of lower diversity still can only partly make

up for the stability lost because of simulated species loss in the first place. The more

optimistic view in terms of restoration approaches for plant communities is that our

findings imply that even though naïve communities may not provide the same level of

functioning as those with a history of co-occurrence, this disadvantage can be partly

compensated by sowing or planting diverse mixtures.

Methods

Field site. This study was conducted at the Jena Experiment field site (Jena, Thuringia, Germany, 51 ˚N, 11 ˚E, 135 m a.s.l.) from 2011–2015. The Jena Experiment is a long-term biodiversity field experiment located on the banks of the Saale River. In 78 experimental field plots of different diversity levels, 60 mostly perennial species typically forming species-rich grassland ecosystems under low-intensity management are grown in a number of species combinations since 2002 [45].

Co-occurrence history. This study included eleven monocultures, twelve 2-species mixtures, twelve 4-species mixtures and twelve 8-species mixtures for a total of 47 experimental plots. The original design of the Jena Experiment [45] dictated the types of monocultures and species combinations we used. We used two co-occurrence-history treatments: communities assembled with offspring of plants that had grown together for 8 years in the 47 plots of the Jena Experiment (selected communities) and communities assembled with plants without a common history of co-occurrence in the Jena Experiment (naïve communities). The plants of naïve communities were grown from seeds obtained in 2010 from the same commercial supplier (Rieger Hofmann GmbH, in Blaufelden-Raboldshausen, Germany) who provided the seeds used for the establishment of the Jena Experiment in 2002. The supplied seeds originated from various field sites in Germany and had been cultivated by reseeding every year for up to five years in monoculture. Seeds of selected communities were produced in an experimental garden in Zurich, Switzerland, from cuttings that had been made in the Jena Experiment in 2010. The cuttings were planted in Zürich in the original species combination in plots fenced with plastic netting to reduce pollination between communities [8]. A small number of seeds were additionally collected directly in the plots of the Jena Experiment. The "selected" seeds were thus offspring of plant populations that had been sown in 2002 and grown until 2010 in plots of the Jena Experiment.

For this study we wanted to exclude potential maternal carry-over effects and therefore used for both selection histories seedlings propagated in a glasshouse. To this aim, in January 2011, the seeds of selected and naïve communities were germinated in potting soil (BF4, De Baat; Holland) in a glasshouse in Zurich. Subsequently, the seedlings were transported back to the Jena Experiment field site and transplanted into 2 x 2 m smaller plots within the original large plots (in March 2011). There were four 1 x 1 m quadrats with different soil treatments in each 2 x 2 m

plot (see next section) and each quadrat was split into two 1 x 0.5 m halves ("half-

quadrats"). We planted seedlings of selected communities into one half and seedlings

of naïve communities into the other half of each quadrat in a hexagonal pattern at a

density of 210 plants per m^2 with a 6-cm distance between individuals. We planted

the species in equal proportions, but five species were excluded from both selected

and naïve communities because they were no longer present in the original plot of the

Jena Experiment. After transplanting, the seedlings received water every second day

for six weeks. The experimental set up is shown in Fig. 2.

Soil treatments. Within each 2 x 2 m plot of the 47 plots of the Jena Experiment, we

removed the original plant cover in September 2010 and used it for the plant

propagation in the experimental garden in Zurich (see previous section).

Subsequently, we excavated the soil to a depth of 0.35 m, added a 10-cm layer of sand

to the bottom of the plots and covered it with a 0.5-mm mesh net. We separated the

borders of the plots and the quadrats by plastic frames. The excavated native soil from

each of the plots was sieved and four soil treatments were prepared. Half of the soil

(approximately 600 kg per plot) was γ-irradiated to remove the original soil biota.

Half of this sterilized soil was then inoculated with 4% (by weight) of live sugar-beet

soil and 4% of sterilized native soil of the corresponding plot ("neutral soil" obtained

by inoculation). Live sugar-beet soil was added to create a neutral soil community and

was previously collected in an agricultural sugar-beet field not associated with the

Jena Experiment, but with comparable soil properties. The second half of the

sterilized soil was inoculated with 4% (by weight) of live sugar-beet soil and 4% of

live native soil of the corresponding plot ("native soil" obtained by inoculation). The

non-sterilized part of the excavated soil was used for the second two soil treatments.

Half of this soil was filled back into one quadrat of the corresponding plot ("native

soil"). The other half of the unsterilized soil was mixed among all plots and filled into

the remaining quadrats ("mixed soil"). However, this fourth soil treatment was

abandoned after two years, which is why this treatment is not included here.

The soils were left to rest in closed bags to allow for the soil chemistry to

equalize and to encourage soil biota of the inocula to colonize the sterilized soil

before planting. The soils were then added into the quadrats in December 2010 and all

quadrats were covered with a net and a water permeable black sheet to avoid spilling

between quadrats until seedling transplantation in March 2011.

Sampling of aboveground biomass. The test communities were weeded three times

a year and the plants were cut to three cm above ground twice a year at typical

grassland harvest times (late May and August) in central Europe. Plant material from

a 50 x 20 cm area in the center of each half-quadrat was collected to measure

aboveground biomass. We sorted the biomass into species, dried it at 70° C and

weighed the dried biomass. There were four May harvests (2012–2015) and three

August harvests (2012–2014) because the experiment was terminated after the fourth

May harvest in 2015.

Natural flood event. In June 2013, the field site was flooded because of heavy rains

in central Europe [26,44]. The flood duration (maximum 25 days) and depth of water

(maximum of 40 cm) was variable among plots due to small topographical differences

among the plots in the experiment [14]. The variation in flooding severity was

distributed across the diversity gradient and within plots the quadrats and half-

quadrats experienced the same flooding severity. We tested whether flood severity [44]

influenced the diversity–stability relationship and any other of the dependent variables

(data not shown). This was not the case, which made us exclude these indices in all

analyses.

Data analysis. We first calculated the stability of community aboveground biomass

as the inverse coefficient of combined intra- and inter-annual variation (CV_{com}^{-1})

among sequential spring and summer harvests. We deliberately combined seasonal

and annual variation in the stability measure to encompass both aspects of temporal

variation in a single measure. The basic sequence for this measure was spring year n,

summer year n and spring year n+1, which had shown increasing stability during the 8

selection years in the Jena Experiment (2003/4, 2005/6, 2007/8, 2009/10; see Fig. 1).

This sequence allowed us to exclude the summer harvest 2013, which was taken

shortly after the flood event and was used for the calculation of resistance and

recovery (see below); and it increased the independence of the sequential measures

from 2003–2010. We also calculated overall temporal stability for test communities

combining the three harvests before the flood with the three after recovery (i.e.

excluding the summer harvest 2013, see Fig. S1). The stability of a single community

was thus the mean community aboveground biomass (μ_{com}) divided by its standard

deviation (σ_{com}). We also calculated average population stabilities (CV_{pop}^{-1}) and

species asynchrony ($1-\theta$, where θ is synchrony) over the same time spans as stability

[18,19]. For monocultures population stability equals community stability and

asynchrony is zero. Variation in community stability, synchrony and population

stability was analyzed with linear mixed-effects models. Stability measures were log-

transformed to improve homoscedasticity and obtain normally distributed residuals in

the analyses [72]. Fixed-effects terms were plant species richness (log scale),

community-history treatment (selected vs. naïve) and soil treatment (native,

inoculated-native or inoculated-neutral soil); plots and quadrats were used as random-

effects terms to get appropriate errors for significance tests [72]. For reasons of

consistency and to allow the use of all data in analyses with covariates we included

monocultures in the analysis of asynchrony. For graphical displays of relationships

between species richness and stability measures and asynchrony means across soil

treatments were corrected for differences between plots within species-richness levels,

which corresponds to using plots and quadrats in the mixed-model analyses. Because

co-occurrence history was a split-plot/split-quadrat treatment applied within each

quadrat it was not affected by the correction. The corrections were obtained by fitting

a model with plots and quadrats only and adding the residuals to the diversity-level

means. This correction corresponds for example to that used in metabolic ecology

when metabolic rate corrected for temperature is plotted against body size [73].

To analyze community responses to the flood event in 2013, we calculated

resistance, recovery and resilience measures [27–29]; see Fig. 4). Resistance is the

difference in community biomass between the average of the three harvests before the

flood and the community biomass directly after the flood (summer 2013), more

negative values indicating lower resistance. Recovery is the difference between the

biomass produced after recovery from the flood (averaged over the three last harvests)

and the biomass directly after the flood (summer 2013), where positive values

indicate the amount of biomass recovered. Resilience is the difference between the

average biomass of the three harvests before the flood and the average biomass of the

three harvests after recovery. Positive values indicate communities were more

productive after the flood and negative values indicate that post-flood biomass had not

returned to its pre-flood state. Variation in resistance, recovery and resilience was

analyzed with linear mixed-effects models. Fixed-effects terms were plant species

richness (log scale), co-occurrence-history treatment (selected vs. naïve) and soil

treatment (native, inoculated-native, or inoculated-neutral soil); plots and quadrats

were used as random-effects terms. Since the measures of resistance, recovery and

resilience can be dependent upon the magnitude of the pre-flood biomass [12], we

analyzed additional models, which included the average of the three harvests before

the flood as covariate (see Table S1). It should be noted that summer biomass was

generally lower than spring biomass and that because the biomass directly after the

flood was measured in summer, absolute values for resistance and recovery were

overestimated. However, because we were only interested in the comparison of

measures between treatments, and these had the same conditions, we did not consider

this overestimation in absolute values as a problem.

We also analyzed pre-flood (first three harvests) and post-flood (last three

harvests) stability in the same way as described at the beginning of this section.

Furthermore, we calculated the species compositional turnover between pre- and post-

flood conditions. Because it includes species abundances, we used the Bray-Curtis

dissimilarity between pre-flood abundances of species (averaged over the first three

harvests) and the post-flood abundances of species (averaged over the last three

harvests). The analysis was the same as for the resistance, recovery, and resilience

measures.

All analyses were conducted using the software R, version 3.2.4 [74]. Mixed

models using residual maximum likelihood (REML) were fitted using the package

ASReml for R [75] and the package 'Pascal' available at GitHub [72].

References

1. Balvanera, P. *et al.* Quantifying the evidence for biodiversity effects on ecosystem functioning and services: Biodiversity and ecosystem functioning/services. *Ecol. Lett.* **9**, 1146–1156 (2006).

2. Cardinale, B. J. *et al.* The functional role of producer diversity in ecosystems. *Am. J. Bot.* **98**, 572–592 (2011).

3. Tilman, D., Lehman, C. L. & Bristow, C. E. Diversity-Stability Relationships: Statistical Inevitability or Ecological Consequence? *Am. Nat.* **151**, 277–282 (1998).

4. Tilman, D., Reich, P. B. & Knops, J. M. H. Biodiversity and ecosystem stability in a decade-long grassland experiment. *Nature* **441**, 629–632 (2006).

5. Hector, A. *et al.* General stabilizing effects of plant diversity on grassland productivity through population asynchrony and overyielding. *Ecology* **91**, 2213–2220 (2010).

6. Reich, P. B. *et al.* Impacts of biodiversity loss escalate through time as redundancy fades. *Science* **336**, 589–592 (2012).

7. Meyer, S. T. *et al.* Effects of biodiversity strengthen over time as ecosystem functioning declines at low and increases at high biodiversity. *Ecosphere* **7**, e01619 (2016).

8. Zuppinger-Dingley, D. *et al.* Selection for niche differentiation in plant communities increases biodiversity effects. *Nature* **515**, 108–111 (2014).

9. van Moorsel, S. J. *et al.* Community evolution increases plant productivity at low diversity. *Ecol. Lett.* **21**, 128–137 (2018).

10. van Moorsel, S. J. *et al.* Evidence for rapid evolution in a grassland biodiversity experiment. *Mol. Ecol.* **in press**, (2019).

11. Schmid, M. W. *et al.* Feedbacks of plant identity and diversity on the diversity and community composition of rhizosphere microbiomes from a long-term biodiversity experiment. *Mol. Ecol.* **28**, 863–878 (2019).

12. Pfisterer, A. B. & Schmid, B. Diversity-dependent production can decrease the stability of ecosystem functioning. *Nature* **416**, 84–86 (2002).

13. Isbell, F. *et al.* Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature* **526**, 574–577 (2015).

14. Fischer, F. M. *et al.* Plant species richness and functional traits affect community
stability after a flood event. *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 20150276
(2016).

15. Schmid, B. Effects of Genetic Diversity in Experimental Stands of Solidago
Altissima -- Evidence for the Potential Role of Pathogens as Selective Agents in
Plant Populations. *J. Ecol.* **82**, 165 (1994).

16. Yachi, S. & Loreau, M. Biodiversity and ecosystem productivity in a fluctuating
environment: the insurance hypothesis. *Proc. Natl. Acad. Sci.* **96**, 1463–1468
(1999).

17. Elmqvist, T. *et al.* Response diversity, ecosystem change, and resilience. *Front.*
Ecol. Environ. **1**, 488–494 (2003).

18. de Mazancourt, C. *et al.* Predicting ecosystem stability from community
composition and biodiversity. *Ecol. Lett.* **16**, 617–625 (2013).

19. Thibaut, L. M. & Connolly, S. R. Understanding diversity-stability relationships:
towards a unified model of portfolio effects. *Ecol. Lett.* **16**, 140–150 (2013).

20. Weigelt, A. *et al.* The Jena Experiment: six years of data from a grassland
biodiversity experiment. *Ecology* **91**, 930–931 (2010).

21. Weisser, W. W. *et al.* Biodiversity effects on ecosystem functioning in a 15-year
grassland experiment: Patterns, mechanisms, and open questions. *Basic Appl.*
Ecol. **23**, 1–73 (2017).

22. Strauss, S. Y., Lau, J. A. & Carroll, S. P. Evolutionary responses of natives to
introduced species: what do introductions tell us about natural communities?
Ecol. Lett. **9**, 357–374 (2006).

23. Germain, R. M., Weir, J. T. & Gilbert, B. Species coexistence: macroevolutionary
relationships and the contingency of historical interactions. *Proc. R. Soc. B Biol.*
Sci. **283**, 20160047 (2016).

24. Eisenhauer, N. *et al.* Plant Diversity Surpasses Plant Functional Groups and Plant
Productivity as Driver of Soil Biota in the Long Term. *PLoS ONE* **6**, e16055
(2011).

25. Eisenhauer, N., Reich, P. B. & Scheu, S. Increasing plant diversity effects on
productivity with time due to delayed soil biota effects on plants. *Basic Appl.*
Ecol. **13**, 571–578 (2012).

26. Blöschl, G., Nester, T., Komma, J., Parajka, J. & Perdigão, R. A. P. The June
2013 flood in the Upper Danube Basin, and comparisons with the 2002, 1954 and
1899 floods. *Hydrol. Earth Syst. Sci.* **17**, 5197–5212 (2013).

27. Schläpfer, F. & Schmid, B. Ecosystem effects of biodiversity: a classification of
hypotheses and exploration of empirical results. *Ecol. Appl.* **9**, 893–912 (1999).

28. Van Ruijven, J. & Berendse, F. Diversity enhances community recovery, but not
resistance, after drought. *J. Ecol.* **98**, 81–86 (2010).

29. Hillebrand, H. *et al.* Decomposing multiple dimensions of stability in global
change experiments. *Ecol. Lett.* **21**, 21–30 (2018).

30. Lloret, F., Keeling, E. G. & Sala, A. Components of tree resilience: effects of
successive low-growth episodes in old ponderosa pine forests. *Oikos* **120**, 1909–
1920 (2011).

31. Cadotte, M. W., Dinnage, R. & Tilman, D. Phylogenetic diversity promotes
ecosystem stability. *Ecology* **93**, (2012).

32. Venail, P. *et al.* Species richness, but not phylogenetic diversity, influences
community biomass production and temporal stability in a re-examination of 16
grassland biodiversity studies. *Funct. Ecol.* **29**, 615–626 (2015).

33. Gravel, D. *et al.* Experimental niche evolution alters the strength of the diversity–
productivity relationship. *Nature* **469**, 89–92 (2011).

34. Lawrence, D. *et al.* Species interactions alter evolutionary responses to a novel
environment. *PLoS Biol.* **10**, e1001330 (2012).

35. Fiegna, F., Moreno-Letelier, A., Bell, T. & Barraclough, T. G. Evolution of
species interactions determines microbial community productivity in new
environments. *ISME J.* **9**, 1235–1245 (2014).

36. Fiegna, F., Scheuerl, T., Moreno-Letelier, A., Bell, T. & Barraclough, T. G.
Saturating effects of species diversity on life-history evolution in bacteria. *Proc.*
R. Soc. B Biol. Sci. **282**, 20151794 (2015).

37. Schmid, B., Birrer, A. & Lavigne, C. Genetic Variation in the Response of Plant
Populations to Elevated CO2 in a Nutrient-Poor, Calcareous Grassland. in
Körner, C., Bazzaz, F.A. (Eds.), Carbon Dioxide, Populations, and Communities
31–50 (Academic Press, San Diego, California, 1996).

38. Davis, M. B., Shaw, R. G. & Etterson, J. R. Evolutionary responses to changing
climate. *Ecology* **86**, 1704–1714 (2005).

39. Nagel, R., Durka, W., Bossdorf, O. & Bucharova, A. Rapid evolution in native
plants cultivated for ecological restoration: not a general pattern. *Plant Biol.* **21**,
551–558 (2019).

40. Bever, J. D. Soil community feedback and the coexistence of competitors:
conceptual frameworks and empirical tests. *New Phytol.* **157**, 465–473 (2003).

41. van der Putten, W. H. *et al.* Plant-soil feedbacks: the past, the present and future
challenges. *J. Ecol.* **101**, 265–276 (2013).

42. Pellkofer, S., van der Heijden, M. G. A., Schmid, B. & Wagg, C. Soil
Communities Promote Temporal Stability and Species Asynchrony in
Experimental Grassland Communities. *PLOS ONE* **11**, e0148015 (2016).

43. Eisenhauer, N. Aboveground–belowground interactions as a source of
complementarity effects in biodiversity experiments. *Plant Soil* **351**, 1–22 (2012).

44. Wright, A. J. *et al.* Flooding disturbances increase resource availability and
productivity but reduce stability in diverse plant communities. *Nat. Commun.* **6**,
6092 (2015).

45. Roscher, C. *et al.* The role of biodiversity for element cycling and trophic
interactions: an experimental approach in a grassland community. *Basic Appl.*
Ecol. **5**, 107–121 (2004).

46. Guerrero-Ramírez, N. R. *et al.* Diversity-dependent temporal divergence of
ecosystem functioning in experimental ecosystems. *Nat. Ecol. Evol.* **1**, 1639–1642
(2017).

47. Vogel, A. *et al.* A new experimental approach to test why biodiversity effects
strengthen as ecosystems age. in *Advances in Ecological Research* (Academic
Press, 2019). doi:10.1016/bs.aecr.2019.06.006

48. Barrett, R. & Schluter, D. Adaptation from standing genetic variation. *Trends*
Ecol. Evol. **23**, 38–44 (2008).

49. Zhao, L., Zhang, Q.-G. & Zhang, D.-Y. Evolution alters ecological mechanisms
of coexistence in experimental microcosms. *Funct. Ecol.* **30**, 1440–1446 (2016).

50. Isbell, F., Polley, H. W. & Wilsey, B. J. Species interaction mechanisms maintain
grassland plant species diversity. *Ecology* **90**, 1821–1830 (2009).

51. Gross, K. *et al.* Species richness and the temporal stability of biomass production:
a new analysis of recent biodiversity experiments. *Am. Nat.* **183**, 1–12 (2014).

52. Flynn, D. F. B., Schmid, B., He, J.-S., Wolfe-Bellin, K. S. & Bazzaz, F. A.

Hierarchical reliability in experimental plant assemblages. *J. Plant Ecol.* **1**, 59–65

(2008).

53. Schöb, C., Brooker, R. W. & Zuppinger-Dingley, D. Evolution of facilitation

requires diverse communities. *Nat. Ecol. Evol.* **2**, 1381–1385 (2018).

54. Lau, J. A. & Lennon, J. T. Rapid responses of soil microorganisms improve plant

fitness in novel environments. *Proc. Natl. Acad. Sci.* **109**, 14058–14062 (2012).

55. terHorst, C. P., Lennon, J. T. & Lau, J. A. The relative importance of rapid

evolution for plant-microbe interactions depends on ecological context. *Proc. R.*

Soc. B Biol. Sci. **281**, 20140028–20140028 (2014).

56. Zuppinger-Dingley, D., Flynn, D. F. B., De Deyn, G. B., Petermann, J. S. &

Schmid, B. Plant selection and soil legacy enhance long-term biodiversity effects.

Ecology **97**, 918–928 (2016).

57. Stocker, T. F. *et al.* Technical summary IPCC 2013. in *Climate Change 2013: The*

Physical Science Basis. Contribution of Working Group I to the Fifth Assessment

Report of the Intergovernmental Panel on Climate Change 33–115 (Cambridge

University Press, 2013).

58. Hirabayashi, Y. *et al.* Global flood risk under climate change. *Nat. Clim Change*

3, 816–821 (2013).

59. Donohue, I. *et al.* Navigating the complexity of ecological stability. *Ecol. Lett.* **19**,

1172–1185 (2016).

60. Pennekamp, F. *et al.* Biodiversity increases and decreases ecosystem stability.

Nature **563**, 109–112 (2018).

61. Caldeira, M. C., Hector, A., Loreau, M. & Pereira, J. S. Species richness,

temporal variability and resistance of biomass production in a Mediterranean

grassland. *Oikos* **110**, 115–123 (2005).

62. Tilman, D. & Downing, J. A. Biodiversity and stability in grasslands. *Nature* **367**,

363–365 (1994).

63. Wagg, C. *et al.* Plant diversity maintains long-term ecosystem productivity under

frequent drought by increasing short-term variation. *Ecology* **98**, 2952–2961

(2017).

64. Wang, Y., Yu, S. & Wang, J. Biomass-dependent susceptibility to drought in

experimental grassland communities. *Ecol. Lett.* **10**, 401–410 (2007).

65. Garssen, A. G., Baattrup-Pedersen, A., Voesenek, L. A. C. J., Verhoeven, J. T. A.
& Soons, M. B. Riparian plant community responses to increased flooding: a
meta-analysis. *Glob. Change Biol.* **21**, 2881–2890 (2015).

66. Wright, A. J. *et al.* Plants are less negatively affected by flooding when growing
in species-rich plant communities. *New Phytol.* **213**, 645–656 (2017).

67. Callaway, R. M. *et al.* Positive interactions among alpine plants increase with
stress. *Nature* **417**, 844 (2002).

68. Fox, L. R. Diffuse Coevolution Within Complex Communities. *Ecology* **69**, 906–
907 (1988).

69. Connell, J. H. Diversity and the Coevolution of Competitors, or the Ghost of
Competition Past. *Oikos* **35**, 131 (1980).

70. Schnitzer, S. A. *et al.* Soil microbes drive the classic plant diversity–productivity
pattern. *Ecology* **92**, 296–303 (2011).

71. Gebremikael, M. T., De Waele, J., Buchan, D., Soboksa, G. E. & De Neve, S. The
effect of varying gamma irradiation doses and soil moisture content on
nematodes, the microbial communities and mineral nitrogen. *Appl. Soil Ecol.* **92**,
1–13 (2015).

72. Schmid, B., Baruffol, M., Wang, Z. & Niklaus, P. A. A guide to analyzing
biodiversity experiments. *J. Plant Ecol.* **10**, 91–110 (2017).

73. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a
Metabolic Theory of Ecology. *Ecology* **85**, 1771–1789 (2004).

74. R Development Core Team. *R: a language and environment for statistical*
computing. R Foundation for Statistical Computing, Vienna, Austria. (2017).

75. Butler, D. *asreml: asmrel() fits the linear mixed model.* (VSNI International,
2009).

Additional information

Supplementary information is available for this paper online.

Data availability. Data for this study are available from the corresponding author and
will be made publicly available upon acceptance on the Pangaea repository.

Code availability. All R scripts are available upon request from the first author.

Acknowledgements

Thanks to Debra Zuppinger-Dingley, Dan Flynn and Varuna Yadav for the establishment of the experimental plots. Thanks to the Jena Experiment for providing infrastructure and help and to D. Trujillo and M. Furler for technical assistance. This study was supported by the Swiss National Science Foundation (grant numbers 130720, 147092 and 166457 to B.S.) and the University Research Priority Program Global Change and Biodiversity of the University of Zurich. The Jena Experiment is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, FOR1451).

Author contributions

B.S. designed research, T.H. and S.J.V.M. performed research; A.E. and N.E. maintained and coordinated the field site; S.J.V.M., C.W. and B.S. analyzed data; S.J.V.M., B.S., O.L.P. and C.W. wrote the paper.

Competing interests

The authors declare no conflicts of interest.

FIGURES & TABLES

**Figure 1 | Combined intra- and inter-annual stability of experimental
communities over the first 8 years in a grassland biodiversity experiment (Jena
Experiment; species richness levels: 1, 2, 4, 8, 16, 60).** The 8-year period was
partitioned into four 2-year periods and within each stability was calculated for spring
and summer harvests in year n and spring harvest in year n+1, corresponding to the
same sequence of three harvests used in subsequent tests communities collected from
the Jena Experiment in 2010 (selected communities) or re-established from seeds of
the original supplier (naïve communities). Based on this figure we tested whether
selected communities have higher temporal stability than naïve communities if tested
under equal conditions and at the same age and time. Thick regression line includes
three outliers outside the top margin of the plot ($P = 0.037$), thin line excludes these
outliers ($P = 0.0018$).
785

Figure 2 | Experimental set-up of test communities in the field. Seeds of cuttings made from plants that had been co-occurring for eight years in 47 plots of the Jena Experiment (selected plants) and seeds purchased from a seed supplier (naïve plants) were germinated at the same time in a glasshouse. These seedlings were then planted in the field in March 2011 according to randomized planting schemes with equal species composition and abundances. Selected (light green) and of naïve communities (dark green) were grown, in the same 47 plots from which co-selected plants had been taken, in four different soil treatments (unsterilized native or mixed soil or sterilized soil with native or neutral inoculum, see Methods). The mixed-soil treatment was not used in this paper because it was harvested early for a different experiment. Co-occurrence history (selected vs. naïve) was thus a split-split plot treatment replicated for 47 community compositions (including 11 monocultures) times three soil treatments.

800
801

Figure 3 | The biodiversity–stability relationship for selected (blue) and naïve communities (red). (A) Community stability, **(B)** mean population stability, **(C)** asynchrony, **(D)** relationship between stability and asynchrony after correction for all other model terms except co-occurrence history. Colored bands show standard errors of predictions from mixed models as presented in Table 1. For significances see Table 1 (panels A–C); the slopes in panel D are significant at $P < 0.001$ and the difference between the two parallel lines is significant at $P = 0.005$. Means across the three soil treatments of values corrected for variation within diversity levels between plots and quadrats are shown in panels A–C and individual residual values are shown in panel D.

Figure 4 | Plant community biomass before and after the extreme event of a late spring flood in June 2013. Points indicate the average community biomass for **(A)** selected (blue) and naïve communities (red) and **(B)** native soil (blue), sterilized soil with native inoculum ("inoculated", green) and sterilized soil with neutral inoculum ("neutral", orange). Resistance is the difference in biomass between the average of the three harvests before the flood and the biomass directly after the flood (label "Flood" on x-axis corresponding to summer harvest 2013). Recovery is the difference in biomass between the average of the three harvests after recovery from the flood and the biomass directly after the flood. Resilience is the difference in biomass between the average of the three harvests after recovery from the flood and the average of the three harvests before the flood. Means and standard errors calculated from raw data.

Figure 5 | Resistance, recovery, and resilience responses in terms of changes in
community biomass caused by the extreme event of a late spring flood in June
2013. (A) Biodiversity–resistance relationships, **(B)** biodiversity–recovery
relationships and **(C)** biodiversity–resilience relationships for selected (blue) and
naïve communities (red). Colored bands show standard errors of predictions from
mixed models as presented in Table 2. For significances see Table 2. Means across
the three soil treatments of values corrected for variation within diversity levels
between plots and quadrats are shown. The dashed line is drawn at 0 in each graph.
(Similar plots with values corrected for variation in pre-flood biomass as covariate are
shown in Fig. S3.)

Figure 6 | The biodiversity–stability relationship for selected (blue) and naïve communities (red). (A) The three harvests before the extreme event of a late spring flood in June 2013 and **(B)** the three harvests after recovery from the flood. Colored bands show standard errors of predictions from mixed models as presented in Table S2. $P < 0.001$ for the effect of log richness in post-flood stability and $P = 0.027$ for the effect of co-occurrence history on post-flood stability. For other test-statistics see Table S2. Means across the three soil treatments of values corrected for variation within diversity levels between plots and quadrats are shown.

847 **Table 1 | Mixed-model ANOVA results for log-transformed community stability, log-transformed mean population stability and**
848 **untransformed asynchrony.** The effects of species richness (log scale), soil treatments and co-occurrence history on the stability of community
849 and population biomass and on asynchrony across the entire experimental period from 2012–2015 were analyzed (excluding the time point
850 immediately after the extreme event of a late spring flood in June 2013). Bold italic text highlights significant effects.
851

Fixed terms	DF_{num}	Stability			Population variance			Asynchrony		
		DF_{den}	F	P	DF_{den}	F	P	DF_{den}	F	P
Log richness (R_{log})	1	***44.1***	***10.74***	***0.002***	***44.1***	***5.27***	***0.027***	***44.1***	***143.00***	***<0.001***
Soil treatment (SH)	2	87.1	0.64	0.529	87.1	1.30	0.278	87.1	0.87	0.424
Co-occurrence history (CH)	1	135.0	1.80	0.181	135.0	3.79	0.054	135.0	0.50	0.479
CH x R_{log}	2	87.2	0.05	0.954	87.2	0.01	0.992	87.2	0.38	0.685
PH x R_{log}	1	***135.0***	***4.79***	***0.030***	***135.0***	***8.38***	***0.004***	135.0	0.05	0.830
Random terms	N	$Var.\ 10^{-3}$	$SE\ 10^{-3}$		$Var.\ 10^{-3}$	$SE\ 10^{-3}$		$Var.\ 10^{-3}$	$SE\ 10^{-3}$	
Plot	46	100.1	25.9		95.6	23.5		17.9	4.6	
Plot x SH	137	15.5	10.9		13.9	7.4		-0.1	2.0	
Residual	274	92.3	11.2		58.4	7.1		20.0	2.5	

852
Note: DF_{num} = numerator degrees of freedom, DF_{den} = denominator degrees of freedom, F = variance ratio, P = probability of type-I error.

Table 2 | Mixed-model ANOVA results for resistance, recovery, and resilience of
community biomass in response to the extreme event of a late spring flood in
June 2013. The effects of species richness (log scale), soil treatments and co-
occurrence history on responses of community biomass to flooding were analyzed.
Bold italic text highlights significant effects. (Similar ANOVAs with pre-flood
biomass as covariate are shown in Table S3.)

		Resistance			Recovery			Resilience		
Fixed terms	DF_{num}	DF_{den}	F	P	DF_{den}	F	P	DF_{den}	F	P
Log richness (R_{log})	1	***44.2***	***9.41***	***0.004***	***44.1***	***15.95***	***<0.001***	44.2	1.69	0.200
Soil treatment (SH)	2	***87.3***	***14.07***	***<0.001***	87.2	0.29	0.745	***87.3***	***6.12***	***0.003***
Co-occurrence history (CH)	1	***135***	***4.19***	***0.043***	***135***	***14.50***	***<0.001***	135	3.48	0.064
SH x R_{log}	2	***87.5***	***5.95***	***0.004***	87.4	1.73	0.184	***87.5***	***6.97***	***0.002***
CH x R_{log}	1	***135***	***5.32***	***0.023***	135	0.48	0.488	135	2.65	0.106
Random terms	N	*Var.*	*SE*		*Var.*	*SE*		*Var.*	*SE*	
Plot	46	3645	1074		2234	771		6910	2238	
Plot x SH	137	775	702		-158	745		1933	1784	
Residual	274	6246	760		7851	956		15914	1937	

862 *Note*: DF_{num} = numerator degrees of freedom, DF_{den} = denominator degrees of
863 freedom, F = variance ratio, P = probability of type-I error.