Evolution increases ecosystem temporal stability and recovery from a flood in

grassland communities

Sofia J. van Moorsel[1], Terhi Hahl[1], Owen L. Petchey[1], Anne Ebeling[2], Nico

Eisenhauer[3,4], Bernhard Schmid[1] and Cameron Wagg[1*]

[1] URPP Global Change and Biodiversity and Department of Evolutionary Biology

and Environmental Studies, University of Zürich, Winterthurerstrasse 190, CH-8057,

Switzerland

[2] Institute of Ecology and Evolution, University of Jena, Dornburger Str. 159, 07743

Jena, Germany

[3] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig,

Deutscher Platz 5e, 04103 Leipzig, Germany

[4] Institute of Biology, Leipzig University, Johannisallee 21, 04103 Leipzig, Germany

[*] Correspondence should be addressed to:

C.W. cameron.wagg@ieu.uzh.ch

Keywords: community evolution | compensatory dynamics | environmental

perturbation | flood | grassland biodiversity | recovery | resilience | resistance | stability

Abstract

Understanding factors that increase ecosystem stability is critical in the face of

environmental change. Biodiversity plays a key role in buffering ecosystems against

disturbances such as extreme climatic events. The evolution of biological

communities within their local environment may also increase ecosystem stability and

resilience, but this has yet to be tested. Here, we provide evidence for such

evolutionary effects using a long-term grassland biodiversity experiment.

Communities of plants with a history of co-occurrence (co-selected communities)

were temporally more stable at low diversity than the same communities of plants

with no such history (naïve communities). Furthermore, co-selected communities

exhibited greater recovery following a major flood, resulting in more stable post-flood

productivity. These results demonstrate that community evolution can increase

ecosystem stability under normal circumstances and in response to extreme

disturbance, but also suggest that high diversity can in part compensate for

evolutionary naïvety.

Introduction

It has long been recognized that greater biodiversity can stabilize ecosystem functioning[1–5]. Such findings emphasize the importance of biodiversity for maintaining ecosystem functioning under future anticipated extreme climatic events[6,7]. The positive effect of biodiversity on maintaining ecosystem productivity over many years can be attributed to greater temporal stability and improved resistance, recovery and resilience to environmental disturbances[4,8,9]. Greater resistance increases ecosystem stability by reducing the loss in productivity[10], while greater recovery increases the amount of productivity that the ecosystem can regain after the disturbance-induced loss[11]. Together resistance and recovery determine ecosystem resilience as we define it here, namely how ecosystem productivity differs between pre- and post-disturbance states[12]. Consequently, how plant diversity stabilizes ecosystem productivity through mediating ecosystem resistance, recovery and resilience has become a focal question in ecology[3,4,9,10,13,14].

Many of the underlying mechanisms by which diversity stabilizes ecosystem productivity are based on the inherent differences among species in their niche requirements and life strategies[15–19]. For instance, different plant species may exhibit high performance under different environmental conditions (termed response diversity). Consequently, a greater plant diversity may stabilize ecosystem productivity under normally fluctuating environmental conditions and especially under environmental disturbance, because there is a higher probability that some species may perform well at any given time point. Asynchrony of species performances, derived from interspecific differences in responses to environmental variation, can thus allow more diverse ecosystems to resist more or recover faster to maintain performance, often referred to as the insurance or portfolio effect[16–18].

Species asynchrony has been conceptually and empirically demonstrated as a mechanism by which biodiversity can stabilize ecosystem productivity[16,18,20–23].

There are a number of community and population attributes associated with ecosystem stability. Ecosystem resistance, recovery and resilience that underlie stability may be dependent upon plant diversity[4,13], plant density[24] and plant functional traits[25]. However, we lack information about the importance of evolutionary processes that may be occurring over the same temporal scales across which ecosystem stability is measured[26]. So far, evolutionary mechanisms underlying the biodiversity–stability relationship have been considered in terms of phylogenetic relatedness that reflects evolutionary mechanisms over broad time scales[27,28]. It remains unclear whether evolution over short time scales can increase the stability of communities under normally fluctuating environmental conditions as well as in response to extreme climatic events by potentially improving resistance, recovery or resilience. Such evolution leading to changes at the community level is referred to as community evolution[29], but so far has almost exclusively been studied in microbial ecosystems[30–32].

Here we test the hypothesis that short-term community evolution in grassland ecosystems will affect ecosystem stability and that this effect may depend on plant diversity, which was experimentally manipulated. This hypothesis is based on community evolution increasing niche differentiation[26], and niche differentiation having the potential to affect community biomass[29], population variability, and population synchrony, all of which contribute to ecosystem stability. We measured ecosystem stability as temporal variation in primary productivity during normal environmental fluctuations. Furthermore, we measured ecosystem responses to disturbance by an extreme event, a naturally occurring major flood, as resistance,

recovery, resilience and post-disturbance stability of productivity[33]. We compared co-

selected communities with naïve communities of 1, 2, 4 or 8 plant species. Co-

selected communities were assembled with offspring from individuals that had co-

occurred in the same communities over 8 years. Naïve communities were assembled

with individuals that were obtained from the same supplier that provided the original

seeds at the beginning of the 8-year selection period of the co-selected

communities[26,29]. To take into account the importance of the local environment, plant

communities were planted in their home soil ("native") or a sterilized soil that was

inoculated with either their native soil biota ("inoculated") or with soil biota from a

different field ("neutral"). Co-selected and naïve communities were grown from

2012–2015 at the field site in Jena, Germany. Productivity was measured in May of

each year and in August of 2012–2014.

Results

Temporal stability, asynchrony and population variation

102   Community evolution significantly modified the diversity–stability

relationship and the diversity–population variation relationship (interaction plant

history x species richness in Table 1). Stability more strongly increased with diversity

in the naïve than in the co-selected communities (Fig. 1a). Conversely, population

variation increased more strongly with diversity in the co-selected than in the naïve

communities. Species asynchrony significantly increased with diversity (main effect

of species richness in Table 1). Different soil treatments did not alter the diversity–

stability relationship (Fig. 1b) and generally, the community-evolution treatments and

soil treatments did not affect stability, asynchrony and population variation (Table 1).

111   The different diversity–stability relationships between the two community-

evolution treatments were related to different asynchrony and stability relationships.

That is, asynchrony of species-level productivity fluctuations was less positively

correlated with ecosystem stability in co-selected than in naïve communities or, in

other words, community evolution significantly reduced the coupling between

species-level asynchrony and ecosystem stability (Fig. 1c). In contrast to the altered

relationship between asynchrony and stability, community evolution did not alter the

relationship between population variation and stability (Fig. 1d).

Responses to an extreme flood event

A naturally occurring flood in early summer 2013 had devastating effects on

the plant communities and reduced productivity in the corresponding time interval

(Fig. 2). Whereas diversity had no general effect on resistance to the flood, the

diversity–resistance relationships differed significantly between the two community-

evolution treatments (Table 2). This resulted from the co-selected communities

having a generally lower resistance at high diversity (Fig. 3a; Table 2). Native and

inoculated soil treatments resulted in highest resistance to the flood, in particular at

high diversity (Fig. 3b).

Ecosystem recovery following the flood was independently increased by

diversity and by community evolution, but soil treatments had no significant effects

(Fig. 3c, d; Table 2). At low (but not at high) diversity, co-selected communities were

more resilient (Fig. 3e). On the other hand, similar to temporal stability, high diversity

could compensate for the reduced resilience of naïve as compared with co-selected

communities (marginally significant interaction plant history x species richness in

Table 2). The effect of species richness on resilience was strongest on the soil that

was inoculated with native soil because of the very low resilience of the

corresponding monocultures (Fig. 3f; Table 2). Species turnover was not influenced

by community-evolution or soil treatments, but did increase with species richness

(Fig. S1; Table S1). Whereas before the flood, co-selected and naïve communities

were equally stable (Fig. 4a), the flood event significantly destabilized post-flood

productivity in the naïve compared with the co-selected communities (Fig. 4b; Table

S2).

Discussion

The potential for greater diversity to maintain a greater ecosystem functioning

over time has been well recognized[1–5]. Additionally, the positive effects of

biodiversity on ecosystem functioning have been shown to strengthen over time[34–36].

Such changes in the functioning of plant communities may reflect evolutionary

processes that occur over the same temporal scales across which ecosystem stability is

being measured. For instance, there is evidence that such temporal changes in the

functioning of communities may reflect rapid evolutionary increases in species

complementarity and community productivity[26,29,32]. Yet the importance of the

interactive effects of biodiversity and community evolution for ecosystem stability

has not been tested so far. Our study provides strong evidence that community

evolution maintained a more stable primary productivity at low diversity, which

consequently altered the diversity–stability relationship under normally fluctuating

environmental conditions. Furthermore, we found that co-selected communities had

an overall greater recovery and post-perturbation stability following a naturally

occurring extreme climatic event. In comparison to the plant diversity and

community-evolution treatments, treatments simulating co-selected vs. novel soil

microbial communities only had minor effects on ecosystem stability. Overall, our

findings demonstrate the potential importance of community-wide evolutionary

processes for maintaining ecosystem functioning and highlights the need to further

consider the integration of evolutionary processes in understanding biodiversity–

ecosystem functioning relationships.

Several mechanisms could have led to the observed differences between co-

selected and naïve communities. First, the changes at community level could have

been due to altered species abundance distributions. However, this would have had to

be related to changed performances of species due to community evolution, because

co-selected and naïve communities only differed in regard to selection history but not

with regard to initial species composition or environment. Furthermore, we could not

detect any significant changes in species abundance distributions. This leaves a

second explanation for the observed community-level effects, phenotypic changes

within species. Such changes could have a genetic or other heritable basis such as

epigenetic or maternal carry-over effects[37]. The latter are unlikely because the

communities were started from seeds rather than cuttings and the effects were

observed over a 4-year time span. We tested in a separate study using a reduced-

representation bisulfate sequencing method[38] for five of the 60 species from the Jena

Experiment if selection on the field site led to genetic or epigenetic changes and found

evidence for genetic but not for epigenetic changes[39]. These genetic changes could

have been due to differential mortality, growth or reproduction among the initially

sown genotypes[40], recombination during sexual reproduction in the field or the

experimental garden or, presumably least likely, to mutation and selection, all

occurring before the start of the present experiment.

Stability, synchrony and population variation

Numerous studies have shown that biodiversity increases ecosystem stability

due to the effects of species asynchrony in diverse communities, allowing high

compensatory population variation to be combined with low community-level

variation over time[15,17,22,41]. Here we also observed such compensatory dynamics, but

this was significantly modified by community evolution. We found that the positive

diversity–stability relationship was stronger in naïve plant communities where species

did not share a common selection history. While this positive diversity–stability

relationship in these naïve plant communities provides further support for the notion

that diversity is a key component underlying ecosystem stability, it also indicates that

the effects of plant diversity on stability may be particularly strong in newly

assembled plant communities. The weaker effect of diversity on stability in the co-

selected communities was due to greater stability at lower diversity in comparison to

the naïve plant communities. These findings suggested that the evolutionary history of

co-occurring species compensated for lower biodiversity by exhibiting a more stable

productivity and, conversely, that biodiversity in naïve communities could partly

compensate for a lack of evolutionarily increased ecosystem stability.

The modified diversity–stability relationship between the co-selected and

naïve plant communities may at least in part have been due to the dampening effect of

community evolution on the positive relationship between species asynchrony and

stability. As expected, species asynchrony increased stability overall. However, in co-

selected communities, species asynchrony was less positively associated with

stability, indicating community evolution led to a partial decoupling of asynchrony

and stability and thus allowed for higher stability at low diversity compared with

naïve communities. More specifically, in communities where species synchrony was

high, such as in less diverse communities, co-selected plant communities were generally more stable than naïve plant communities. At low diversity, species in co-selected communities compared with species in naïve communities were also generally less temporally variable in their productivity (see significant interaction plant history x species richness in Table 1). Therefore, the co-selected plant communities with low asynchrony were able to maintain greater community stability. On the other hand, the stronger coupling between asynchrony and stability allowed naïve compared with co-selected communities to have a more stable productivity at high diversity, where species fluctuated more asynchronously through time.

The reduced temporal stability at low synchrony in our co-selected plant communities (see Fig. 1c) may be due to the effect of community evolution on species competitive interactions. For instance, it has been previously shown that more diverse communities can result in the selection for characteristics by which individuals avoid competition and exhibit greater complementarity[26,29]. Such changes in species interactions can impact species dynamics and ultimately the stability of the net community productivity. This is because species competing more strongly with one another can exhibit greater asynchrony in their temporal performance since temporal variations in the environment may temporarily favor the competitive advantage of one species over another resulting in their negative temporal covariance[15,20,41]. Such destabilizing effects of evolution have been demonstrated in microalgae communities where a greater phylogenetic distance among community members results in weaker competitive interactions, that in turn reduces the competition-driven temporal asynchrony among species and the compensatory dynamics that are required to stabilize the net productivity of the communities[28].

The interactions between plants and their soil communities are well known to

influence ecosystem functioning[43] and these interactions likely change over ecological

time-scales[44–46]. We therefore anticipated that soils would play an important role in

ecosystem stability. However, we found little evidence that our soil treatments

influenced the temporal performance of plant communities under field conditions and

did not have any interactive effects with the community-evolution treatments on the

temporal performance of the plant communities. However, we did observe that the

soils influenced the pre-flood productivity of the plant communities. In particular, we

found that the pre-flood productivity was generally lower in communities grown in

native soil, which might have been due to a greater density of antagonistic soil biota

or the inoculated soils may have had a greater pool of available soil resources

resulting from the soil sterilization process[47]. The difference between inoculated and

native soils was, perhaps literally, "washed away" by the flood event. The flood event

may thus have equalized the soil properties among soil treatments and consequently

dampened any potential plant–soil interactions that have been observed elsewhere to

influence ecosystem stability[48,49].

Responses to an extreme flood event

Our study system was exposed to a naturally occurring extreme flood event

that strongly reduced the productivity of the plant communities[50]. Here we took

advantage of this to further assess the hypothesis that community evolution may

enhance ecosystem stability in response to disturbance events, due to greater

resistance, recovery or resilience. Biodiversity decreased resistance, confirming

previous findings[11,24], but the relationship was context-dependent. Community

evolution reduced resistance to flooding at high diversity (see Fig. 3a). This lower

resistance can be attributed to greater pre-flood productivity of the co-selected

communities. In this sense, co-selected communities had "more to lose" when faced

with this extreme climate event, an observation reported also in other grassland

systems in response to drought[11]. Thus, overall the reduced productivity of naïve

communities resulted in less absolute loss in productivity due to the flood and their

greater resistance. It was previously shown that selection for niche differentiation

results in higher community productivity[26]. Our results indicate that the selection-

driven increase in productivity may consequently reduce the resistance to extreme

climate events at higher levels of diversity. However, the greater productivity of co-

selected plant communities may have also allowed them to recover back to their pre-

disturbed state and maintain a more stable post-flood productivity. This is supported

by our analysis using pre-flood productivity as a covariate (see Supporting

Information, Fig. S2, Table S3). Once we corrected for the pre-flood productivity,

species richness increased resilience in both co-selected and naïve communities, and

we observed a strong interactive effect of community evolution and species richness

for ecosystem resistance (Table 2).

Biodiversity had a strong positive effect on the recovery and, in the naïve

communities also on resilience of the plant communities, paralleling numerous other

studies emphasizing the importance of biodiversity for ecosystem recovery and

resilience[4,11,12,26]. In addition, community evolution further increased recovery and

resilience (see Fig. 3c, e). The positive effect of community evolution on post-flood

recovery suggests that the local environment may have acted as a selective filter on

these plant communities. The ancestral communities of the co-selected plants were

initially sown into the field site in 2002, which is a natural floodplain where the plant

communities were exposed to environmental conditions related to soil moisture

saturation at previous milder flood events in winter 2003 and winter 2005 (personal

communication with C. Roscher) than the one in 2013 and thus selection likely

favored individuals with traits that allowed them to perform well under such

conditions and recover more rapidly[51]. It has also been shown that community

evolution can result in increased plant species complementarity in this system[26]. In

line with this, the result that community evolution in these plant communities resulted

in greater recovery may be suggestive of selection for greater facilitative effects after

the perturbation as may be anticipated under the stress-gradient hypothesis[53].

The greater recovery in the co-selected communities also resulted in a greater

resilience up to a diversity level of 4 species and was followed by an increased post-

flood stability compared with naïve communities. This means that community

evolution aided the return of the functioning of these communities to their pre-

perturbed state and their pre-perturbation temporal performance. Our results

demonstrate that community evolution can have strong effects on altering the

population- and community-level mechanisms underpinning increased stability in

ecosystem productivity, specifically mechanisms relating synchrony, recovery, and

resilience. Finally, it is important to note that the greater post-flood stability in our

study was not due to compositional changes as indicated by the similar compositional

turnover between naïve and co-selected plant communities. Thus, the increased

resistance and post-flood stability can be attributed to the greater recovery and post-

flood performance of the individual species functioning in concert. We suggest that

genetic changes within the species comprising the co-selected communities were

responsible for the increased stability. In a study comparing plant individuals within

species from different selection backgrounds in the Jena Experiment (high vs. low

diversity), we did not find evidence for epigenetic divergence, but a strong genetic

signal[39]. Likely two different evolutionary processes — co-selection between the

species within each particular community composition and "diffuse" co-selection

among all species within the community — have improved the species' abilities to

function more complementarily with other species in the community[26,29].

Conclusions

Our findings show that ecosystem stability and recovery can not only be

promoted by biodiversity but also by evolutionary processes in plant communities

over the time scales at which stability is being measured. In particular, we found that

community evolution can enhance the stability of ecosystem productivity when

diversity is low, whereas in the absence of community evolution a greater diversity

was more critical for increasing stability. The community evolution-driven increase in

ecosystem stability and recovery was likely linked to the selection on individual plant

characteristics that facilitated greater recovery in these plant communities. If

evolution in grassland communities increases ecosystem resilience and stability,

maintaining co-selected plant communities could be a crucial precaution under global

change and increased frequency of extreme climatic events[6,7]. Furthermore,

integrating evolutionary processes into the temporal changes in biodiversity–

ecosystem functioning relationships is likely to be a promising future avenue for

predicting how ecosystems may respond to climatic extremes and biodiversity loss.

Methods

Field site. This study was conducted at the Jena Experiment field site (Jena, Thuringia, Germany, 51 ˚N, 11 ˚E, 135 m a.s.l.) from 2011 to 2015. The Jena Experiment is a long-term biodiversity field experiment located on the banks of the Saale River. In 78 experimental field plots of different diversity levels, 60 central European grassland species are grown in a number of species combinations since 2002[54].

Community-evolution treatments. This study included eleven monocultures, twelve 2-species mixtures, twelve 4-species mixtures and twelve 8-species mixtures for a total of 47 experimental plots. We used two community-evolution treatments: plants with eight years of shared community selection in these experimental plots (co-selected communities) and plants without a common selection history in the Jena Experiment (naïve communities). The naïve plant seeds without a common selection history were obtained from the same commercial seed supplier (Rieger Hofmann GmbH, in Blaufelden-Raboldshausen, Germany) who provided the seeds used for the establishment of the original Jena Experiment plant communities[54]. The supplied seeds originated from various field sites in Germany and have been cultivated by reseeding every year for at least five years in monoculture. Seeds of co-selected communities were produced in an experimental garden in Zurich, Switzerland, from cuttings that had been made in the Jena Experiment. The cuttings were planted in Zürich in the original species combination in plots fenced with plastic netting to reduce pollination between communities[26]. A small number of seeds were additionally collected directly in the plots of the Jena Experiment. The "selected" seeds were thus offspring of plant populations that had been sown in 2002 and grown until 2010 in plots of the Jena Experiment.

In January 2011, the seeds of co-selected and naïve communities were

germinated in potting soil (BF4, De Baat; Holland) in a glasshouse in Zurich.

Subsequently, the seedlings were transported back to the Jena Experiment field site

and transplanted into 2 x 2 m subplots of the original plots (in March 2011). There

were four 1 x 1 m quadrats with different soil treatments in each subplot (see next

section) and each quadrat was split into two 1 x 0.5 m halves ("half-quadrats"). We

planted seedlings of co-selected communities into one half and seedlings of naïve

communities into the other half of each quadrat in a hexagonal pattern at a density of

210 plants per m^2 with a 6-cm distance between individuals. We planted the species in

equal proportions, but five species were excluded from both co-selected and naïve

communities because they were no longer present in the original plot of the Jena

Experiment. After transplanting, the seedlings received water every second day for six

weeks.

Soil treatments. Within each 2 x 2 m subplot of the 47 plots of the Jena Experiment,

we removed the original plant cover in September 2010 and used it for the plant

propagation in the experimental garden in Zurich (see previous section).

Subsequently, we excavated the soil to a depth of 0.35 m, added a 10-cm layer of sand

to the bottom of the plots and covered it with a 0.5 mm mesh net. We separated the

borders of the subplots and the quadrats by plastic frames. The excavated native soil

from each of the plots was sieved and four soil treatments were prepared. Half of the

soil (approximately 600 kg per plot) was γ-irradiated to remove the original soil biota.

Half of this sterilized soil was then inoculated with 4% (by weight) of live sugar-beet

soil and 4% of sterilized native soil of the corresponding plot ("neutral soil" obtained

by inoculation). Live sugar-beet soil was added to create a neutral soil community and

was previously collected in an agricultural sugar-beet field not associated with the

Jena Experiment, but with comparable soil properties. The second half of the

sterilized soil was inoculated with 4% (by weight) of live sugar-beet soil and 4% of

live native soil of the corresponding plot ("native soil" obtained by inoculation). The

non-sterilized part of the excavated soil was used for the second two soil treatments.

Half of this soil was filled back into one quadrat of the corresponding plot ("native

soil"). The other half of the unsterilized soil was mixed among all plots and filled into

the remaining quadrats. However, this fourth soil treatment was abandoned after two

years, which is why this treatment is not included here.

The soils were left to rest in closed bags to allow for the soil chemistry to

equalize and to encourage soil biota of the inocula to colonize the sterilized soil

before planting. The soils were then added into the quadrats in December 2010 and all

quadrats were covered with a net and a water permeable black sheet to avoid spilling

between quadrats until seedling transplantation in March 2011.

Sampling of aboveground biomass. The test communities were weeded three times

a year and the plants were cut to three cm above ground twice a year at typical

grassland harvest times (late May and August) in central Europe. Plant material from

a 50 x 20 cm area in the centre of each half-quadrat was collected to measure

aboveground biomass. We sorted the biomass into species, dried it at 70°C and

weighed the dried biomass. There were four May harvests (2012–2015) and three

August harvest (2012–2015) because the experiment was terminated after the fourth

May harvest in 2015.

Natural flood event. In June 2013, the field site was flooded due to heavy rains in

central Europe[50,55]. The flood duration (maximum 25 days) and depth of water

(maximum of 40 cm) was variable among plots and quadrats due to small

topographical differences among the plots in the experiment[25]. The variation in

flooding severity was distributed across the diversity gradient and within subplots the

quadrats and half-quadrats experienced the same flooding severity. We tested whether

flood severity[50] influenced the diversity–stability relationship and any other of our

dependent variables (data not shown). This was not the case, which made us exclude

these indices in all analyses.

Data analysis. We first calculated the stability of ecosystem functioning as the

inverse coefficient of variation (CV_{com}^{-1}) in the net community biomass over six time

points. We excluded the flooding time point to avoid any confounding effects of the

dramatic biomass loss in response to the flood. The stability of a single community is

thus the mean net community biomass (μ_{com}) divided by its standard deviation (σ_{com}).

To explain the change in stability with biodiversity and between the community-

evolution treatments we split the community-level variation across the six harvests

(CV_{com}) into its two component parts, these being the weighted population variation

(CV_{pop}) and the species synchrony (θ) that are defined elsewhere[18,22]. Variation in

these indices of stability, synchrony and population stability was analyzed with linear

mixed-effects models. Fixed-effects terms were plant species richness (log scale),

community-evolution treatment (plant history co-selected vs. naïve) and soil treatment

(native, inoculated and neutral soil); subplots and quadrats were used as random-

effects terms to get appropriate errors for significance tests[56].

We calculated the resistance, recovery and resilience (see Fig. 2) to assess the

communities' responses to the flood event. Resistance is the change in productivity

between the average of the three harvests prior to the flood and the productivity

during the flood event, more negative values indicating lower resistance. Recovery is

the difference in the biomass produced post-flood (averaged over the three post-flood

harvests) from the biomass produced during the flood event, where positive values

indicate the amount of biomass recovered. Resilience is the difference between the

pre- and post-flood biomasses where positive values indicate communities that were

more productive after than before the flood event, whereas negative values indicate

that the post-flood productivity had not returned to its pre-flood state. Variation in

resistance, recovery and resilience was analyzed with linear mixed-effects models.

Fixed-effects terms were plant species richness (log scale), community-evolution

treatment (plant history co-selected vs. naïve) and soil treatment (native, inoculated

and neutral soil); subplots and quadrats were used as random-effects terms. Since the

measures of resistance, recovery and resilience can be dependent upon the magnitude

of the pre-flood productivity[24,57], we created additional models which included the

pre-flood productivity as a covariate (see Supporting Information).

We calculated the species compositional turnover between pre- and post-flood

conditions. Because it includes species abundances, we used the Bray-Curtis

dissimilarity between pre-flood abundances of species (averaged over the three pre-

flood harvests) and the post-flood abundances of species (averaged over the three

post-flood harvests). The analysis was the same as for the resistance, recovery and

resilience measures. We also analyzed pre-flood (three harvests before the flooding

event) and post-flood (three harvests after the flooding event) stability in the same

way as described at the beginning of this section.

All analyses were conducted using the software R, version 3.2.4[58]. Mixed

models using residual maximum likelihood (REML) were fitted using the package

ASReml for R[59] and the package 'Pascal' available at GitHub[56].

References

1. Tilman, D., Reich, P. B. & Knops, J. M. H. Biodiversity and ecosystem stability

in a decade-long grassland experiment. *Nature* **441,** 629–632 (2006).

2. Proulx, R. *et al.* Diversity promotes temporal stability across levels of ecosystem

organization in experimental grasslands. *PLoS ONE* **5,** e13382 (2010).

3. Tilman, D. & Downing, J. A. Biodiversity and stability in grasslands. *Nature* **367,**

363–365 (1994).

4. Isbell, F. *et al.* Biodiversity increases the resistance of ecosystem productivity to

climate extremes. *Nature* **526,** 574–577 (2015).

5. Schläpfer, F. & Schmid, B. Ecosystem effects of biodiversity: a classification of

hypotheses and exploration of empirical results. *Ecol. Appl.* **9,** 893–912 (1999).

6. Stocker, T. F. *et al.* Technical summary IPCC 2013. in *Climate Change 2013: The*

Physical Science Basis. Contribution of Working Group I to the Fifth Assessment

Report of the Intergovernmental Panel on Climate Change 33–115 (Cambridge

University Press, 2013).

7. Rockström, J. *et al.* A safe operating space for humanity. *Nature* **461**, 472–475

(2009).

8. Oliver, T. H. *et al.* Biodiversity and resilience of ecosystem functions. *Trends*

Ecol. Evol. **30,** 673–684 (2015).

9.   Wagg, C. *et al.* Plant diversity maintains long-term ecosystem productivity under

frequent drought by increasing short-term variation. *Ecology* **98,** 2952–2961

(2017).

10. Caldeira, M. C., Hector, A., Loreau, M. & Pereira, J. S. Species richness,

temporal variability and resistance of biomass production in a Mediterranean

grassland. *Oikos* **110,** 115–123 (2005).

11. Van Ruijven, J. & Berendse, F. Diversity enhances community recovery, but not

resistance, after drought. *J. Ecol.* **98,** 81–86 (2009).

12. Lloret, F., Keeling, E. G. & Sala, A. Components of tree resilience: effects of

successive low-growth episodes in old ponderosa pine forests. *Oikos* **120,** 1909–

1920 (2011).

13. Hautier, Y. *et al.* Anthropogenic environmental changes affect ecosystem stability

via biodiversity. *Science* **348,** 336–340 (2015).

14. Gross, K. *et al.* Species richness and the temporal stability of biomass production:

a new analysis of recent biodiversity experiments. *Am. Nat.* **183,** 1–12 (2014).

15. Tilman, D., Lehman, C. L. & Bristow, C. E. Diversity-stability relationships:

statistical inevitability or ecological consequence? *Am. Nat.* **151,** 277–282 (1998).

16. Yachi, S. & Loreau, M. Biodiversity and ecosystem productivity in a fluctuating

environment: the insurance hypothesis. *Proc. Natl. Acad. Sci.* **96,** 1463–1468

(1999).

17. Hector, A. *et al.* General stabilizing effects of plant diversity on grassland

productivity through population asynchrony and overyielding. *Ecology* **91,** 2213–

2220 (2010).

18. Thibaut, L. M. & Connolly, S. R. Understanding diversity-stability relationships:

towards a unified model of portfolio effects. *Ecol. Lett.* **16,** 140–150 (2013).

19. Isbell, F. *et al.* High plant diversity is needed to maintain ecosystem services.

Nature **477,** 199–202 (2011).

20. Gonzalez, A. & Loreau, M. The causes and consequences of compensatory

dynamics in ecological communities. *Annu. Rev. Ecol. Evol. Syst.* **40,** 393–414

(2009).

21. Loreau, M., Mouquet, N. & Gonzalez, A. Biodiversity as spatial insurance in

heterogeneous landscapes. *Proc. Natl. Acad. Sci.* **100,** 12765–12770 (2003).

22. de Mazancourt, C. *et al.* Predicting ecosystem stability from community

composition and biodiversity. *Ecol. Lett.* **16,** 617–625 (2013).

23. Leary, D. J. & Petchey, O. L. Testing a biological mechanism of the insurance

hypothesis in experimental aquatic communities *J. of Anim. Ecol.* **78,** 1143–1151

(2009).

24. Pfisterer, A. B. & Schmid, B. Diversity-dependent production can decrease the

stability of ecosystem functioning. *Nature* **416,** 84–86 (2002).

25. Fischer, F. M. *et al.* Plant species richness and functional traits affect community

stability after a flood event. *Philos. Trans. R. Soc. B Biol. Sci.* **371,** 20150276

(2016).

26. Zuppinger-Dingley, D. *et al.* Selection for niche differentiation in plant

communities increases biodiversity effects. *Nature* **515,** 108–111 (2014).

27. Cadotte, M. W., Dinnage, R. & Tilman, D. Phylogenetic diversity promotes

ecosystem stability. *Ecology* **93,** S223–S233 (2012).

28. Venail, P. A., Alexandrou, M. A., Oakley, T. H. & Cardinale, B. J. Shared

ancestry influences community stability by altering competitive interactions:

evidence from a laboratory microcosm experiment using freshwater green algae.

Proc. R. Soc. B Biol. Sci. **280,** 20131548–20131548 (2013).

29. van Moorsel, S. J. *et al.* Community evolution increases plant productivity at low

diversity. *Ecol. Lett.* **21,** 128–137 (2018).

30. Fiegna, F., Moreno-Letelier, A., Bell, T. & Barraclough, T. G. Evolution of

species interactions determines microbial community productivity in new

environments. *ISME J.* **9,** 1235–1245 (2014).

31. Fiegna, F., Scheuerl, T., Moreno-Letelier, A., Bell, T. & Barraclough, T. G.

Saturating effects of species diversity on life-history evolution in bacteria. *Proc.*

R. Soc. B Biol. Sci. **282,** 20151794 (2015).

32. Lawrence, D. *et al.* Species interactions alter evolutionary responses to a novel

environment. *PLoS Biol.* **10,** e1001330 (2012).

33. Garssen, A. G., Baattrup-Pedersen, A., Voesenek, L. A. C. J., Verhoeven, J. T. A.

& Soons, M. B. Riparian plant community responses to increased flooding: a

meta-analysis. *Glob. Change Biol.* **21,** 2881–2890 (2015).

34. Cardinale, B. J. *et al.* Impacts of plant diversity on biomass production increase

through time because of species complementarity. *Proc. Natl. Acad. Sci.* **104,**

18123–18128 (2007).

35. Reich, P. B. *et al.* Impacts of biodiversity loss escalate through time as

redundancy fades. *Science* **336,** 589–592 (2012).

36. Meyer, S. T. *et al.* Effects of biodiversity strengthen over time as ecosystem

functioning declines at low and increases at high biodiversity. *Ecosphere* **7,**

e01619 (2016).

37. Verhoeven, K. J. F., vonHoldt, B. M. & Sork, V. L. Epigenetics in ecology and

evolution: what we know and what we need to know. *Mol. Ecol.* **25,** 1631–1638

(2016).

38. van Gurp, T. P. *et al.* epiGBS: reference-free reduced representation bisulfite

sequencing. *Nat. Methods* **13,** 322–324 (2016).

39. van Moorsel, S. J. Community evolution increases ecosystem functioning and

stability. (University of Zurich, 2017).

40. Barrett, R. & Schluter, D. Adaptation from standing genetic variation. *Trends*

Ecol. Evol. **23,** 38–44 (2008).

41. Isbell, F. I., Polley, H. W. & Wilsey, B. J. Biodiversity, productivity and the

temporal stability of productivity: patterns and processes. *Ecol. Lett.* **12,** 443–451

(2009).

42. Loreau, M. & de Mazancourt, C. Species synchrony and its drivers: Neutral and

nonneutral community dynamics in fluctuating environments. *Am. Nat.* **172,** E48–

E66 (2008).

43. van der Putten, W. H. *et al.* Plant-soil feedbacks: the past, the present and future

challenges. *J. Ecol.* **101,** 265–276 (2013).

44. Lau, J. A. & Lennon, J. T. Rapid responses of soil microorganisms improve plant

fitness in novel environments. *Proc. Natl. Acad. Sci.* **109,** 14058–14062 (2012).

45. terHorst, C. P., Lennon, J. T. & Lau, J. A. The relative importance of rapid

evolution for plant-microbe interactions depends on ecological context. *Proc. R.*

Soc. B Biol. Sci. **281,** 20140028–20140028 (2014).

46. Zuppinger-Dingley, D., Flynn, D. F. B., De Deyn, G. B., Petermann, J. S. &

Schmid, B. Plant selection and soil legacy enhance long-term biodiversity effects.

Ecology **97,** 918–928 (2016).

47. Gebremikael, M. T., De Waele, J., Buchan, D., Soboksa, G. E. & De Neve, S. The

effect of varying gamma irradiation doses and soil moisture content on

nematodes, the microbial communities and mineral nitrogen. *Appl. Soil Ecol.* **92,**

1–13 (2015).

48. Yang, G. *et al.* The interaction between arbuscular mycorrhizal fungi and soil

phosphorus availability influences plant community productivity and ecosystem

stability. *J. Ecol.* **102,** 1072–1082 (2014).

49. Pellkofer, S., van der Heijden, M. G. A., Schmid, B. & Wagg, C. Soil

communities promote temporal stability and species asynchrony in experimental

grassland communities. *PLoS ONE* **11,** e0148015 (2016).

50. Wright, A. J. *et al.* Flooding disturbances increase resource availability and

productivity but reduce stability in diverse plant communities. *Nat. Commun.* **6,**

6092 (2015).

51. Wright, A. J. *et al.* Plants are less negatively affected by flooding when growing

in species-rich plant communities. *New Phytol.* **213,** 645–656 (2017).

52. Loeuille, N. Influence of evolution on the stability of ecological communities:

Evolution and stability of communities. *Ecol. Lett.* **13,** 1536–1545 (2010).

53. Callaway, R. M. *et al.* Positive interactions among alpine plants increase with

stress. *Nature* **417,** 844 (2002).

54. Roscher, C. *et al.* The role of biodiversity for element cycling and trophic

interactions: an experimental approach in a grassland community. *Basic Appl.*

Ecol. **5,** 107–121 (2004).

55. Blöschl, G., Nester, T., Komma, J., Parajka, J. & Perdigão, R. A. P. The June

2013 flood in the Upper Danube Basin, and comparisons with the 2002, 1954 and

1899 floods. *Hydrol. Earth Syst. Sci.* **17,** 5197–5212 (2013).

56. Schmid, B., Baruffol, M., Wang, Z. & Niklaus, P. A. A guide to analyzing

biodiversity experiments. *J. Plant Ecol.* **10,** 91–110 (2017).

57. Wang, Y., Yu, S. & Wang, J. Biomass-dependent susceptibility to drought in experimental grassland communities. *Ecol. Lett.* **10,** 401–410 (2007).

58. R Development Core Team. *R: a language and environment for statistical computing.* (R Foundation for Statistical Computing, Vienna, Austria, 2017).

59. Butler, D. *asreml: asmrel() fits the linear mixed model.* (VSNI International, 2009).

Additional information

Supplementary information is available for this paper online.

Data availability. Data for this study are available from the corresponding author and will be made publicly available upon acceptance on the Pangaea repository.

Code availability. All R scripts are available upon request from the corresponding author C.W.

Acknowledgements

Thanks to Debra Zuppinger-Dingley, Dan Flynn and Varuna Yadav for the establishment of the experimental plots. Thanks to the Jena Experiment for providing infrastructure and help and to D. Trujillo and M. Furler for technical assistance. This study was supported by the Swiss National Science Foundation (grant numbers 130720, 147092 and 166457 to B.S.) and the University Research Priority Program Global Change and Biodiversity of the University of Zurich. The Jena Experiment is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, FOR1451).

Author contributions

B.S. designed research, T.H. and S.J.V.M. performed research; S.J.V.M., C.W. and

B.S. analyzed data; S.J.V.M., B.S. and C.W. wrote the paper with substantial

contributions of the other authors.

Competing interests

The authors declare no conflicts of interest.

 FIGURES & TABLES

Fig. 1 | The biodiversity–stability relationship in response to community-evolution and soil treatments. a, Greater plant diversity is required for greater stability, but more strongly so in naïve communities (plant history x species richness (log-transformed): $F_{1, 135.0} = 4.794$, $P = 0.030$). **b,** Different soils did not alter the overall biodiversity–stability relationship (soil treatment x species richness (log-transformed): $F_{2, 87.1} = 0.048$, $P = 0.954$). The three soil treatments were: native soil, sterilized soil inoculated with native soil (inoculated) and sterilized soil (neutral).

Means and standard errors are shown for each diversity level. **c,** Stability increased

with asynchrony strongly for both naïve and selected communities (main effect

asynchrony: $F_{1, 204.5} = 84.55$, $P = $ <0.001) but stability increased more steeply with

asynchrony in naïve plant communities than in co-selected communities (plant history

x asynchrony interaction: $F_{1, 145.8} = 3.93$, $P = 0.049$). **d,** The relationship between

stability and population variation was not altered by community evolution (plant

history x population CV: $F_{1, 160.3} = 1.348$, $P = 0.247$).

Fig. 2 | The flood event greatly reduced ecosystem productivity. Points indicate **a**,

the average productivity for co-selected and naïve plant communities and **b**, the

average productivity for the three soil treatments native soil, sterilized soil inoculated

with native soil (inoculated) and sterilized soil (neutral). Resistance is the change in

productivity between the average of the three harvests prior to the flood and the

productivity during the flood event (label "Flood" on x-axis corresponding to August

2013). Recovery is the change in productivity from this level to the average of the

three post-flood harvests. Resilience is the change from the average of the three pre-

flood harvests to the average of the three post-flood harvests.

Fig. 3 | Resistance, recovery and resilience responses to a major flood event.
Biodiversity–resistance relationships are shown in response to **a**, community-evolution and **b**, soil treatments. Biodiversity–recovery relationships are shown in response to **c**, community-evolution and **d**, soil treatments. Biodiversity–resilience relationships are shown in response to **e**, community-evolution and **f**, soil treatments. Means and standard errors are shown for each diversity level.

Fig. 4 | Influence of the community-evolution treatments on stability pre-flood vs.

post-flood. Means for each diversity and community-evolution treatment with error

bars indicating the model-estimated standard error are shown. **a,** naïve plant

communities and co-selected communities were equally stable pre-flood ($F_{1,\ 130.8}$ =

1.54, P = 0.217). **b,** co-selected plant communities were more stable than naïve plant

communities post-flood ($F_{1,\ 131.6}$ = 4.94, P = 0.028). Results are presented on a log-

scale.

Table 1 | Mixed-model ANOVA results for stability, population variance and

synchrony. The effects of species richness (log-scale), community-evolution (plant

history) and soil treatments on the stability of community productivity, population

variance and synchrony across the entire experimental period form 2012–2015. The

time point of the flooding event was excluded in the calculations of stability,

population variance and asynchrony to prevent any confounding effects of the flood.

Bold italic text highlights significant effects.

		Stability			Population variance			Asynchrony		
Fixed terms	DF_{num}	DF_{den}	F	P	DF_{den}	F	P	DF_{den}	F	P
Log richness (R_{log})	1	*44.1*	*10.74*	*0.002*	*44.1*	*5.27*	*0.027*	*44.1*	*143*	*<0.001*
Plant history (PH)	1	135	1.805	0.181	135	3.79	0.054	135	0.5	0.479
Soil treatment (SH)	2	87.1	0.641	0.529	87.1	1.3	0.278	87.1	0.87	0.424
PH x R_{log}	1	*135*	*4.794*	*0.030*	*135*	*8.38*	*0.004*	135	0.05	0.830
SH x R_{log}	2	87.2	0.048	0.954	87.2	0.01	0.992	87.2	0.38	0.685
		Var.	*SE*		*Var.*			*Var.*	*SE*	
Random terms	N	10^{-3}	10^{-3}		10^{-3}	*SE* 10^{-3}		10^{-3}	10^{-3}	
Plot	46	100.1	25.9		95.6	23.5		17.9	4.57	
Plot x SH	137	15.5	10.9		13.9	7.4		-0.05	1.96	
Residual	274	92.3	11.23		58.4	7.1		20.02	2.46	

Note: DF_{num} = numerator degrees of freedom, DF_{den} = denominator degrees of freedom, F =

variance ratio, P = probability of type-I error.

Table 2 | Mixed effect ANOVA results for resistance, recovery and resilience. The

effects of species richness (log-scale), community-evolution (plant history) and soil

treatments are shown for the changes in productivity due to flooding. Bold italic text

highlights significant effects.

		Resistance			Recovery			Resilience		
Fixed terms	DF_{num}	DF_{den}	F	P	DF_{den}	F	P	DF_{den}	F	P
Log richness (R_{log})	1	***44.2***	***9.413***	***0.004***	***44.1***	***15.95***	***<0.001***	44.2	1.69	0.200
Plant history (PH)	1	***135***	***4.19***	***0.043***	***135***	***14.5***	***<0.001***	135	3.476	0.064
Soil treatment (SH)	2	***87.3***	***14.07***	***<0.001***	87.2	0.295	0.746	***87.3***	***6.116***	***0.003***
PH x R_{log}	1	***135***	***5.323***	***0.023***	135	0.484	0.488	135	2.65	0.106
SH x R_{log}	2	***87.5***	***5.949***	***0.004***	87.4	1.728	0.184	***87.5***	***6.97***	***0.002***
Random terms	N	*Var.*	*SE*		*Var.*	*SE*		*Var.*	*SE*	
Plot	46	3645.1	1073.9		2233.7	771.4		6909.8	2237.8	
Plot x SH	137	775.2	702.4		-157.6	744.9		1933.2	1784.5	
Residual	274	6245.8	760.2		7851.01	955.6		15913.8	1937	

695 *Note*: DF_{num} = numerator degrees of freedom, DF_{den} = denominator degrees of freedom, F =

696 variance ratio, P = probability of type-I error.