Title: Cardiovascular Disease Risk Factor Responses to a Type 2 Diabetes Care Model Including Nutritional Ketosis at One Year: An Open Label, Non-Randomized, Controlled Study

Nasir H. Bhanpuri, PhD (Corresponding Author)

    Virta Health, San Francisco, CA, USA

    ████████████████

Sarah J. Hallberg, DO, MS

    Indiana University Health Arnett, Medically Supervised Weight Loss, Lafayette, IN, USA

    Virta Health, San Francisco, CA, USA

    ████████████

Paul T. Williams, PhD

    Independent Consultant, Lafayette, CA, USA

    ████████████████

Amy L. McKenzie, PhD

    Virta Health, San Francisco, CA, USA

    ████████████

Kevin D. Ballard, PhD

    Miami University, Oxford, OH, USA

    ████████████

Wayne W. Campbell, PhD

    Purdue University, West Lafayette, IN, USA

    ████████████

James P. McCarter, MD, PhD

Virta Health, San Francisco, CA, USA

Washington University School of Medicine, Department of Genetics, St. Louis, MO, USA

███████████████

Stephen D. Phinney, MD, PhD

Virta Health, San Francisco, CA, USA

███████████████

Jeff S. Volek, PhD, RD

Virta Health, San Francisco, CA, USA

The Ohio State University, Department of Human Sciences, Columbus, OH, USA

███████████████

Abstract

Background

Cardiovascular disease (CVD) is a leading cause of death among adults with type 2 diabetes

mellitus (T2D). We recently reported that glycemic control in patients with T2D can be significantly

improved through a continuous care intervention (CCI) including nutritional ketosis. The purpose of this

study was to examine CVD risk factors in this cohort.

Methods

We investigated CVD risk factors in patients with T2D who participated in a one year open label,

non-randomized, controlled study. The CCI group (n = 262) received treatment from a health coach and

medical provider. A usual care (UC) group (n = 87) was independently recruited to track customary T2D

progression. Circulating biomarkers of cholesterol metabolism and inflammation, blood pressure (BP),

carotid intima media thickness (cIMT), multi-factorial risk scores and medication use were examined.

Results

The CCI group consisted of 262 patients (baseline mean(SD): age 54(8) y, BMI 40.4(8.8)

kg/m^2). Intention-to-treat analysis (% change) revealed the following at 1-year with P values < 0.0019

indicating statistical significance after adjustment for multiple comparisons: total LDL-particles (LDL-P)

(-4.9%, P=0.02), small LDL-P (-20.8%, P=1.2x10^{-12}), LDL-P size (+1.1%, P=6.0x10^{-10}), ApoB (-1.6%,

P=0.37), ApoA1 (+9.8%, P<10^{-16}), ApoB/ApoA1 ratio (-9.5%, P=1.9x10^{-7}), triglyceride/HDL-C ratio (-

29.1%, P<10^{-16}), large VLDL-P (-38.9%, P=4.2x10^{-15}), and LDL-C (+9.9%, P=4.9x10^{-5}). Additional

effects were reductions in blood pressure, high sensitivity C-reactive protein, and white blood cell count

(all P<1x10^{-7}) while cIMT was unchanged. The 10-year atherosclerotic cardiovascular disease (ASCVD)

risk score decreased -11.9% (P=4.9x10^{-5}). Antihypertensive medication use was discontinued in 11.4 %

of CCI participants (P=5.3x10^{-5}). The UC group of 87 patients (baseline mean(SD): age 52(10)y, BMI

36.7(7.2) kg/m^2) showed no significant changes. After adjusting for baseline differences when comparing

CCI and UC groups, significant improvements for the CCI group included small LDL-P, ApoA1,

triglyceride/HDL-C ratio, HDL-C, hs-CRP, and ASCVD score. The CCI group showed a greater rise in

LDL-C.

Conclusions

A continuous care treatment including nutritional ketosis in patients with T2D improved most

biomarkers of CVD risk after one year. The increase in LDL-cholesterol appeared limited to the large

LDL subfraction. LDL particle size increased, total LDL-P and ApoB were unchanged, and inflammation

and blood pressure decreased.

Trial registration

Clinicaltrials.gov: NCT02519309. Registered 10 August 2015

<u>Keywords</u>

ketosis, carbohydrate restriction, type diabetes, cardiovascular disease, risk factor, atherogenic

dyslipidemia, inflammation, blood pressure, antihypertensive medication, continuous remote care

<u>Background</u>

Despite advances in the prevention and treatment of cardiovascular disease (CVD), it remains the leading

cause of death in adults across the world [1]. Specifically, among those with type 2 diabetes (T2D) in the

U.S., CVD accounts for 44% of mortality [2]. T2D rates have doubled over the past twenty years [3] and

CVD risk increases two to four-fold with a diagnosis of T2D [4], warranting the identification of novel

interventions to combat T2D. Intensive lifestyle interventions with dietary carbohydrate restriction [5-8],

including the recently described continuous remote care model, which helps patients with T2D sustain

nutritional ketosis [9,10], have demonstrated improved glycemic control concurrent with medication

reduction. However, the long-term sustainability and impact of these interventions on CVD risk and lipid

profiles remains a subject of debate [11,12].

Atherogenic dyslipidemia, a known risk factor for CVD [13], is highly prevalent in patients with T2D

[14] and tightly linked to high-carbohydrate diets[15]. The condition is characterized by increased

triglycerides, decreased high-density lipoprotein cholesterol concentration (HDL-C) and increased small

low-density lipoprotein particle number (small LDL-P). Evidence suggests that increased very low-

density lipoprotein particle number (VLDL-P), and in particular large VLDL-P, may be one of the key

underlying abnormalities in atherogenic dyslipidemia [14,16-18]. Elevated concentrations of small LDL

are often associated with increased total LDL particle number (LDL-P) and ApoB [19,20]. Particularly in

patients with insulin resistance and T2D, elevated LDL-P and ApoB may exist even with normal to low

LDL-C values [19,21,22]. Reliance on LDL-C for risk assessment in T2D patients may miss the impact of

atherogenic dyslipidemia and elevated LDL-P. It has been proposed that LDL-P or ApoB may be superior

to LDL-C as a predictor of CVD [22-25].

Previous studies of carbohydrate restriction of up to 1-year found a consistent decrease in triglycerides

and increase in HDL-C, while LDL-C slightly increased or decreased [15,26-28]. Although LDL-C is a

risk factor for CVD, low LDL-C may belie elevations in small LDL, LDL-P or ApoB. Conversely,

increased LDL-C with a low carbohydrate diet may primarily reflect the large LDL subfraction and may

not increase CVD risk if total LDL-P or ApoB concentrations are unchanged or decline.

Inflammation, as assessed by elevated high-sensitivity C-reactive protein (hsCRP) or white blood cell

count (WBC) [29-32], is an independent CVD risk factor and is involved in all stages of atherogenesis

[33]. Inflammation is often observed in T2D concurrent with atherogenic dyslipidemia [34] and

represents an additional CVD risk even in individuals with low to normal LDL-C [35,36]. Hypertension is

an additive risk factor in this patient population. Tighter blood pressure control has been associated with

reduction in the risk of deaths related to diabetes. This included decreased CVD, stroke and

microvascular complications [37].

For this open label, non-randomized, controlled, before-and-after study, we investigated the effects of a

continuous care intervention (CCI) on CVD risk factors. The CCI included individualized digital support

with telemedicine, health coaching, education in nutritional ketosis, biometric feedback, and an online

peer-support community. Given the multi-faceted pathophysiology of CVD, we assessed the one-year

responses in several biomarkers related to cholesterol and lipoprotein metabolism, blood pressure, and

inflammation, as well as carotid intima media thickness (cIMT) and medication use. Some results were

previously reported in relation to glycemic control [10] and are presented here as they pertain to CVD

risk.

Methods

Intervention

As previously described [9,10], we utilized a prospective, longitudinal study design with a cohort of

patients with T2D from the greater Lafayette, Indiana, USA, region who self-selected to participate in the

CCI (Clinicaltrials.gov Identifier NCT02519309). Patients in the CCI were provided access to a web-

based software application (app) for biomarker reporting and monitoring including body weight, blood

glucose and blood betahydroxybutyrate (BHB; a marker of ketosis). The remote care team consisted of a

health coach and physician or nurse practitioner who provided nutritional advice and medication

management, respectively. Patients were guided by individualized nutrition recommendations to achieve

and sustain nutritional ketosis. CCI participants self-selected to receive education via either an onsite

group setting (CCI-onsite) or via the app (CCI-web). Social support was provided via an online peer

community. This study was approved by the Franciscan Health Lafayette Institutional Review Board, and

participants provided written informed consent.

Glycemic control medication management was previously described [9,10]. For participants with a

history of hypertension, a home automatic sphygmomanometer was supplied. Participants reported their

home readings in the app daily to weekly depending on recent control and instruction from the

supervising physician. Antihypertensive prescriptions were adjusted based on home readings and

reported symptoms. Health coaches responded to patient app reported readings of systolic blood pressure

less than 110 mmHg with specific questions about symptoms of hypotension. Following resolution of

hypertension, diuretics were the first antihypertensive medications to be discontinued, followed by beta

blockers, unless the participant had a history of coronary artery disease. Angiotensin-converting-enzyme

inhibitors (ACEs) and angiotensin II receptor blockers (ARBs) were generally continued due to known

renal protection with diabetes [38,39]. Statin medications were adjusted when appropriate to maintain a

goal of LDL-P under 1000 nmol·L^{-1} or participant preference after full risk-benefit discussion.

To track T2D progression in the same geography and health system as the CCI, an independent cohort of

patients with T2D who received usual care (UC) were recruited. These patients were referred to registered

dietitians providing dietary advice according to American Diabetes Association guidelines [40].

Outcome Measures

Anthropometrics and vital signs for the CCI group were obtained at baseline, 70 days, and one year. A

stadiometer was used to assess height and used in the calculation of body mass index. A calibrated scale

in the clinic measured weight to the nearest 0.1 lb (Model 750, Detecto; Webb City, MO, USA) and

values were converted to kg. Participants were provided a cellular-connected home scale for daily weight.

Blood pressure was obtained manually by trained staff after participants rested in a seated position for

five minutes. Adverse events were reported and reviewed by the Principal Investigator and the

Institutional Review Board.

Fasting blood draws for the CCI group were collected at baseline, 70 days, and one-year follow-up

(ranging from 11 - 15 months). Blood analytes were determined via standard procedures at a Clinical

Laboratory Improvement Amendment (CLIA) accredited laboratory on the day of sample collection or

from stored serum. Serum aliquots were stored at -80°C and thawed for determination of ApoB, ApoA1,

total cholesterol, triglycerides, and direct HDL-C concentrations by FDA approved methods (Cobas c501,

Roche Diagnostics; Indianapolis, IN, USA). LDL was calculated using the Friedewald equation [41].

Lipid subfractions were quantified using clinical NMR LipoProfile® (LabCorp, Burlington NC, USA;

[42]). The LipoProfile3 algorithm used in the present investigation was used previously to relate lipid

subfractions to CVD risk [35,43,44]. The NMR-derived lipoprotein insulin resistance score (LP-IR) is

proposed to be associated with the homeostasis model assessment of insulin resistance (HOMA-IR) and

glucose disposal rate (GDR) [45]. The multifactorial 10-year atherosclerotic cardiovascular disease

(ASCVD) risk score was also computed [46].

Anthropometrics, vital signs and fasting blood draws for the UC group were obtained at baseline and one

year as described above using the same clinical facilities and laboratory and data collection methods.

Home biometrics for the UC group were not tracked and 70-day outcomes were not measured.

Carotid ultrasonography for cIMT measure was performed at baseline and one year in CCI and UC

groups to characterize atherosclerotic risk. Ultrasound technicians were trained according to protocols

that were previously tested and used to assess subclinical atherosclerosis [47,48]. The right and left

common carotid arteries were imaged one cm distal to the carotid bulb using a L12-3 multi-frequency

linear-array transducer attached to a high-resolution ultrasound system (Phillips EPIQ 5, Amsterdam,

Netherlands). Longitudinal images were captured in three imaging planes: anterior, lateral, and posterior.

Digital images were analyzed using edge-detection software (Carotid Analyzer for Research; Medical

Imaging Application, Coralville, IA) to trace the lumen-intima and intima-medial boundaries of the

artery. Analyses were performed by the same blinded investigator to obtain right and left mean arterial

diameter and mean cIMT.

Statistics

JMP software (version 5.1, SAS Institute; Cary, SC, USA) was used for all statistical analyses except

multiple imputation. Multiple imputation, conducted with Stata software (version 11, StataCorp; College

Station, TX, USA), was used to estimate means and standard errors describing the variability between

imputations. Seven hundred imputations from multivariate normal regression were run to estimate the

missing values (4% missing at baseline and 22% missing at one year). Two-sample t-tests were used to

test for significance of the differences in baseline biomarker values between groups. Two-sample t-tests

were also used to test for differences between one-year changes between groups. Paired t-tests and

analysis of covariance (ANCOVA) when adjusted for baseline covariates (sex, age, baseline BMI, and

African-American race) were used to test for significance of within-group changes. To reduce skewness

before testing for significance, triglyceride, triglyceride/HDL-C ratio and hsCRP were first log-

transformed, however aside from P values, the tables present results from the untransformed data. Percent

change in a given biomarker was calculated as the mean difference value divided by the mean baseline

value. The standard error of percent change of a given biomarker was calculated as the standard error of

the change divided by the mean baseline value. Significant changes in proportions (e.g. medication use)

were tested using McNemar's test with continuity correction in completers, and linear regression of the

changes in the dichotomous states when missing outcome data were imputed.

Throughout the manuscript, standard deviations are presented within parentheses and standard errors are

presented following "±" symbol. Nominal significance levels (P) are presented in the tables; however, a

significance level of P<0.0019 ensures simultaneous significance at P<0.05 for a Bonferroni adjustment

for the 26 variables examined. Unless otherwise noted, results presented are intention-to-treat analyses

(all starters) with missing values estimated by imputation. Some results are designated as completer

analyses (excluding participants who withdrew or lacked biomarkers at one year).

<u>Results</u>

<u>Baseline characteristics of participants</u>

The baseline characteristics of the 262 T2D participants who began the CCI are shown in Table 1. There

were no significant differences in baseline characteristics between groups self-selecting web-based (CCI-

web) and onsite education (CCI-onsite) (Supplementary Table 1) nor were there significant differences in

biomarker changes at one year between the groups (Supplementary Table 2). Therefore, results for the

two groups were combined for further analyses.

The baseline characteristics of participants with measurements at both baseline and one year were not

significantly different from dropouts and participants with missing data after correcting for multiple

comparisons (Supplementary Table 1).

An independently recruited UC group of 87 T2D participants, which provided an observational

comparison group from the same geography and health system, showed no significant differences from

the CCI group in baseline characteristics except mean body weight and BMI were higher in the CCI

versus the UC group (Table 1, P<0.001).

<u>Changes in biomarkers of CVD risk at one year</u>

Two-hundred eighteen (83%) participants remained enrolled in the CCI group at one year. One-year

changes in CVD biomarkers are detailed in Table 2 and percent changes from baseline are shown in

Figure 1. The within-CCI group changes in the following lipids and lipoproteins were all statistically

significant after adjusting for multiple comparisons (P<0.0019), reported here as mean percent difference

from baseline: ApoA1 (+9.9%), ApoB/ApoA1 ratio (-9.5%), triglycerides (-24.4%), LDL-C (+9.6%),

HDL-C (+18.1%), triglyceride/HDL-C ratio (-29.1%), large VLDL-P (-38.9%), small LDL-P (-20.8%),

LDL-particle size (+1.1%), total HDL-P (+4.9%), and large HDL-P (+23.5%). There were no significant

changes after adjusting for multiple comparisons in total LDL-P (-4.9%, p=0.02) or ApoB (-1.6%,

p=0.37).

The CCI group experienced significant reductions in systolic BP (-4.8%), diastolic BP (-4.3%), hs-CRP (-

39.3%) and WBC count (-9.1%). Significant reductions were observed in overall use of antihypertensive

medication (-11.4%) and diuretics (-9.6%) whereas changes in ACE or ARB (0.7%) and statin (-3.3%)

use were not significant. Significant reductions were observed in both multivariate metrics: 10-year

ASCVD risk (-11.9%) and LP-IR (-19.6%). There was no significant change in cIMT (averaged right and

left values). In addition, changes in cIMT were not significantly correlated with baseline LDL-P or LDL-

C, or changes in LDL-P or LDL-C (all p≥0.33).

One-year results from the UC group are provided in Table 2 and Figure 1. Within the UC group, after

adjustment for multiple comparisons there were no significant changes at one year. After adjusting for

differences in baseline characteristics and multiple comparisons, the changes observed at one year for the

following biomarkers were significantly different between the CCI and UC groups (mean ΔCCI – mean

ΔUC, where Δ is one year minus baseline): small LDL-P (-178 nmol·L^{-1}), ApoA1 (+16 mg·dL^{-1}),

triglyceride/HDL-C ratio (-3.1), LDL particle size (+0.31 nm), HDL-C (+8 mg·dL^{-1}), LDL-C (+22 mg·dL^{-}

^{1}), hs-CRP (-4.0 mg·dL^{-1}), and LP-IR (-12).

<u>Range of outcomes</u>

The distribution and range of intervention response for the CCI and UC groups were compared for LDL-

P, small LDL-P, large VLDL-P, ApoB, ApoA1, ApoB/ApoA1 ratio, and TG/HDL-C ratio

(Supplementary Figure 1). Ranges of change observed in the CCI group were within the ranges observed

in the UC group for increases in LDL-P, small LDL-P, ApoB and ApoB/ApoA1 ratio. There were two

CCI participants (2/203, 1.0%) whose change in large VLDL-P exceeded the maximum observed in the

UC group (15.2 nmol·L^{-1}). There was one CCI participant (1/185, 0.5%) whose change in ApoA1 was

less than the minimum observed in the UC group (58 mg·dL^{-1}) and one CCI participant (1/186, 0.5%)

whose change in triglyceride/HDL-C ratio was higher than the maximum observed in the UC group

(64.9).

<u>Discussion</u>

This study demonstrates that a CCI utilizing remote physician and health coach support with nutritional

ketosis beneficially altered most CVD risk factors in patients with T2D at one year. Changes included:

decreased small LDL-P, triglycerides, blood pressure and antihypertensive medication, hsCRP, and WBC

count; increased HDL-C and LDL particle size; no change in LDL-P, ApoB, and cIMT and an increase in

LDL-C. Combined with the previously reported improvements in glycemic control and reduction in

obesity [10], which reduce CVD risk [49], these results demonstrate multiple additional benefits of the

CCI with the exception of increased LDL-C.

Studies of dietary carbohydrate restriction, with a presumed increase in saturated fat intake, have shown

modest changes in LDL-C levels [15,26-28,50]. The mean 10 mg/dL change observed in the CCI group in

this study is numerically higher than the upper range of values reported by meta-analysis of lipid changes

over one year related to carbohydrate restriction (-7 mg/dl to +7 mg/dl) [51]. Higher LDL-C is related to

increased CVD risk [52,53], but also is *inversely* correlated with mortality in two large prospective

studies and a systemic review [54-56]. Additionally, there is no evidence that increasing or decreasing

LDL-C with diet interventions has any impact on mortality. LDL-C increased in the current study but

both ApoB and LDL-P, measures found to be better predictors of CVD risk, did not change significantly

[20-23,25,57]. In addition, the reduction in small LDL-P, increase in LDL size, and decrease in large

VLDL-P that occurred in the present investigation are also associated with reduced CVD risk [58-60].

While mean response of CCI participants demonstrated an improvement in most lipid biomarkers and

CVD risk factors other than LDL-C, we investigated whether a minority of participants might have

unfavorable responses to the intervention. Our results suggest that a small number of participants ($\leq$ 1%)

demonstrated changes at one year outside the range of what was observed in a usual care population

(Supplementary Figure 1). Thus, these results counter the concern that a significant portion of participants

may have an extremely adverse reaction to the CCI (due to presumed increase in saturated fat intake) as

compared to UC.

Inflammation is directly involved in all aspects of the pathogenesis of CVD [33]. High-sensitivity CRP

and WBC count are widely accepted markers of inflammation and risk factors for CVD [29-32]. In

addition to reducing cholesterol, reduction in inflammation may be a secondary mechanism of statins in

lowering CVD risk [61-63]. The present study demonstrated a 39% reduction of hsCRP and 9% reduction

in WBC count in the CCI, indicating a significant reduction in inflammation at one year. This response

may be due in part to suppression of the NLRP3 inflammasome by BHB [64].

The reduction of blood pressure with concurrent reduction in antihypertensive medication was also

significant. Blood pressure goals were recently reduced [65] and strong evidence exists that elevated

blood pressure is a primary cardiovascular risk factor [66]. An analysis of a large T2D population

suggested that antihypertensive medication may have limited effectiveness in reducing the prevalence of

hypertension in these patients [67], whereas a study of weight loss interventions showed that a decrease in

blood pressure predicted regression of carotid vessel wall volume [68]. Thus, additional lifestyle

interventions that can augment blood pressure reduction such as the CCI described here may reduce CVD

events. Additionally, the antihypertensives that were primarily decreased in the current study were shown

to increase the risk for diabetes [69]. Their removal may represent further metabolic benefit.

Carotid intima media thickness (cIMT) is a non-invasive measure of subclinical atherosclerosis that is

significantly associated with CVD morbidity and mortality [47,48,70,71]. However, a recent meta-

analysis in 3,902 patients with T2D found that cIMT progression over an average of 3.6 years did not

correlate with CVD events [70]. We found no significant change in cIMT from baseline to one year in

either the CCI or UC groups. Progression or regression of cIMT may take multiple years to manifest and

may require a larger cohort to achieve statistical significance [71]. In summary, the cIMT results from this

study provide no evidence of vascular harm or benefit from one year of nutritional ketosis in patients with

T2D.

<u>Strengths and limitations of the study</u>

Prior studies have demonstrated favorable improvements in atherogenic dyslipidemia with minimal or no

change in LDL-C and LDL-P following managed ketogenic diets in small short-term randomized trials.

This study's strengths include its larger cohort with high retention, prospective design and one-year

duration. The study was the first to assess ApoB and ApoA1 in a T2D population adhering to a ketogenic

diet. This study also has real-world application due to the outpatient setting without the use of meal

replacements or food provisions.

Limitations of this study include the lack of randomization between the CCI and UC groups. In addition,

the intervention provided to CCI participants was of greater intensity than UC. This was a single site

study and the racial composition of study participants was predominantly Caucasian. The study was not

of sufficient size and duration to determine significant differences in CVD morbidity or mortality. Since

the intervention led to concurrent weight loss and improvements in cardiovascular health, it is difficult to

conclude how much of the improvement can be attributed to weight loss versus other simultaneous

metabolic changes. Future trials could include a longer multi-site, randomized controlled trial to allow for

hard end point evaluation. Greater racial and ethnic diversity, a broader age range, and greater disease

severity could also be evaluated.

<u>Conclusions</u>

A T2D intervention combining technology-enabled continuous remote care with individualized plans

encouraging nutritional ketosis has demonstrated diabetes status improvement while improving many

CVD risk factors including atherogenic dyslipidemia, inflammation and blood pressure while decreasing

use of antihypertensive mediations. Ongoing research will determine the continued safety, sustainability,

and effectiveness of the intervention.

<u>List of abbreviations</u>

ACE: angiotensin-converting-enzyme inhibitors

ApoA1: apolipoprotein A1

ApoB: apolipoprotein B

ARB: angiotensin II receptor blockers

ASCVD risk score: 10-year atherosclerotic cardiovascular disease risk score

BHB: beta-hydroxybutyrate

BP: blood pressure

CCI: continuous care intervention

CCI-onsite: subset of CCI participants who selected to receive onsite education

CCI-web: subset of CCI participants who selected to receive web-based education

cIMT: carotid intima media thickness

CVD: cardiovascular disease

GDR: glucose disposal rate

HOMA-IR: homeostatic model assessment of insulin resistance

hsCRP: high sensitive C-reactive protein

LP-IR: lipoprotein insulin resistance score

T2D: type 2 diabetes

UC: usual care

VLDL-P: very low-density lipoprotein particle number

WBC: white blood cell

Declarations

Ethics approval and consent to participate

This study was approved by the Franciscan Health Lafayette Institutional Review Board, and participants

provided written informed consent.

Consent for publication

Not Applicable

Availability of data and material

The datasets used and/or analysed during the current study are available from the corresponding author on

reasonable request.

<u>Competing interests</u>

NHB, SJH, ALM, JPM, and SDP are employees of Virta Health Corp. and have been offered stock

options. SDP and JSV are founders of Virta Health Corp. PTW and KDB are paid consultants of Virta

Health Corp. WWC has no competing interests.

<u>Funding</u>

The study was funded by Virta Health Corp.

<u>Authors' contributions</u>

SJH, ALM, WWC, JPM, SDP and JSV conceptualized and designed the study. PTW performed the

formal analysis. SJH and ALM contributed to the investigation. NHB and SJH wrote the original draft.

NHB and PTW created the data tables and visualizations. All authors contributed to revising and editing

of the manuscript. All authors read and approved the final manuscript.

<u>Acknowledgements</u>

The authors are extremely grateful to the research participants for offering their time and energy to

participate in advancing scientific discovery. Thanks to Indiana University Health (IUH) staff, especially

Tamara Hazbun, Monica Keyes, Danielle Wharff, Patti McKee, Joni Anderson, Zachary Roberts,

Christina Selsor, and Douglas Jackson, and Virta Health staff Rachel Bolden, Sydney Rivera, and Deklin

Veenhuizen for contributions to various aspects of the study including patient care, study coordination

and data processing. Thanks to the health coaches who provided guidance to CCI participants: Brittanie

Volk, Brent Creighton, Theresa Link, Bobbie Glon, and Marcy Abner. Thanks to Roxie McKee of IUH,

Dave Gibson and Jennifer Powers of Washington University, and Teryn Sapper and staff from the Volek

Laboratory at The Ohio State University for assistance in sample analysis, storage and/or transportation

logistics. Thanks to Angela Fountain, Irinia Shalaurova, and Jim Otvos of LabCorp for guidance on

interpreting LipoProfile results. Thank you to Ronald Krauss for guidance on lipid analysis and Ethan

Weiss for critical reading of the manuscript.

References

1. Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, et al. Heart Disease and Stroke

Statistics-2017 Update: A Report From the American Heart Association. Circulation. American Heart

Association, Inc; 2017;135:e146–e603.

2. Gregg EW, Gu Q, Cheng YJ, Narayan KMV, Cowie CC. Mortality trends in men and women with

diabetes, 1971 to 2000. Ann Intern Med. 2007;147:149–55.

3. Centers for Disease Control and Prevention (CDC). National Diabetes Statistics Report, 2017. 2017 Jul

pp. 1–20.

4. Martín-Timón I, Sevillano-Collantes C, Segura-Galindo A, Del Cañizo-Gómez FJ. Type 2 diabetes and

cardiovascular disease: Have all risk factors the same strength? World J Diabetes. 2014;5:444–70.

5. Saslow LR. An Online Intervention Comparing a Very Low-Carbohydrate Ketogenic Diet and

Lifestyle Recommendations Versus a Plate Method Diet in Overweight Individuals With Type 2

Diabetes: A Randomized Controlled Trial. J Med Internet Res. 2017;19:e36.

6. Boden G, Sargrad K, Homko C, Mozzoli M, Stein TP. Effect of a low-carbohydrate diet on appetite,

blood glucose levels, and insulin resistance in obese patients with type 2 diabetes. Ann Intern Med.

2005;142:403–11.

7. Yancy WS, Foy M, Chalecki AM, Vernon MC, Westman EC. A low-carbohydrate, ketogenic diet to

treat type 2 diabetes. Nutr Metab. 2005;2:34.

8. Westman EC, Yancy WS, Mavropoulos JC, Marquart M, McDuffie JR. The effect of a low-

carbohydrate, ketogenic diet versus a low-glycemic index diet on glycemic control in type 2 diabetes

mellitus. Nutr Metab. 2008;5:36.

9. McKenzie A, Hallberg S, Creighton BC, Volk BM, Link T, Abner M, et al. A Novel Intervention

Including Individualized Nutritional Recommendations Reduces Hemoglobin A1c Level, Medication

Use, and Weight in Type 2 Diabetes. JMIR Diabetes. 2017;2:e5.

10. Hallberg SJ, McKenzie AL, Williams PT, Bhanpuri NH, Peters AL, Campbell WW, et al.

Effectiveness and Safety of a Novel Care Model for the Management of Type 2 Diabetes at 1 Year: An

Open-Label, Non-Randomized, Controlled Study. Diabetes Therapy. 2018;1–30.

11. Wood TR, Hansen R, Sigurðsson AF, Jóhannsson GF. The cardiovascular risk reduction benefits of a

low-carbohydrate diet outweigh the potential increase in LDL-cholesterol. Br J Nutr. 2016;115:1126–8.

12. Mansoor N, Vinknes KJ, Veierød MB, Retterstøl K. Effects of low-carbohydrate diets v. low-fat diets

on body weight and cardiovascular risk factors: a meta-analysis of randomised controlled trials. Br J Nutr.

2016;115:466–79.

13. Fruchart J-C, Sacks F, Hermans MP, Assmann G, Brown WV, Ceska R, et al. The Residual Risk

Reduction Initiative: a call to action to reduce residual vascular risk in patients with dyslipidemia. The

American Journal of Cardiology. 2008;102:1K–34K.

14. Arca M, Pigna G, Favoccia C. Mechanisms of diabetic dyslipidemia: relevance for atherogenesis.

Curr Vasc Pharmacol. 2012;10:684–6.

15. Volek JS, Fernandez ML, Feinman RD, Phinney SD. Dietary carbohydrate restriction induces a

unique metabolic state positively affecting atherogenic dyslipidemia, fatty acid partitioning, and

metabolic syndrome. Prog. Lipid Res. 2008;47:307–18.

16. Taskinen M-R, Borén J. New insights into the pathophysiology of dyslipidemia in type 2 diabetes.

Atherosclerosis. 2015;239:483–95.

17. Ginsberg BH. Factors Affecting Blood Glucose Monitoring: Sources of Errors in Measurement.

Journal of Diabetes Science and Technology. 2009;3:903–13.

18. Adiels M, Olofsson S-O, Taskinen M-R, Borén J. Overproduction of very low-density lipoproteins is

the hallmark of the dyslipidemia in the metabolic syndrome. Arteriosclerosis, Thrombosis, and Vascular

Biology. 2008;28:1225–36.

19. Cromwell WC, Otvos JD. Heterogeneity of low-density lipoprotein particle number in patients with

type 2 diabetes mellitus and low-density lipoprotein cholesterol <100 mg/dl. The American Journal of

Cardiology. 2006;98:1599–602.

20. Sniderman AD, St-Pierre AC, Cantin B, Dagenais GR, Després J-P, Lamarche B.

Concordance/discordance between plasma apolipoprotein B levels and the cholesterol indexes of

atherosclerotic risk. The American Journal of Cardiology. 2003;91:1173–7.

21. Otvos JD, Mora S, Shalaurova I, Greenland P, Mackey RH, Goff DC. Clinical implications of

discordance between low-density lipoprotein cholesterol and particle number. Journal of Clinical

Lipidology. 2011;5:105–13.

22. Cromwell WC, Otvos JD, Keyes MJ, Pencina MJ, Sullivan L, Vasan RS, et al. LDL Particle Number

and Risk of Future Cardiovascular Disease in the Framingham Offspring Study - Implications for LDL

Management. Journal of Clinical Lipidology. 2007;1:583–92.

23. Sniderman AD, Toth PP, Thanassoulis G, Furberg CD. An evidence-based analysis of the National

Lipid Association recommendations concerning non-HDL-C and apoB. Journal of Clinical Lipidology.

2016;10:1248–58.

24. Toth PP, Grabner M, Punekar RS, Quimbo RA, Cziraky MJ, Jacobson TA. Cardiovascular risk in

patients achieving low-density lipoprotein cholesterol and particle targets. Atherosclerosis.

2014;235:585–91.

25. Barter PJ, Ballantyne CM, Carmena R, Castro Cabezas M, Chapman MJ, Couture P, et al. Apo B

versus cholesterol in estimating cardiovascular risk and in guiding therapy: report of the thirty-person/ten-

country panel. J. Intern. Med. Blackwell Publishing Ltd; 2006. pp. 247–58.

26. Seshadri P, Iqbal N, Stern L, Williams M, Chicano KL, Daily DA, et al. A randomized study

comparing the effects of a low-carbohydrate diet and a conventional diet on lipoprotein subfractions and

C-reactive protein levels in patients with severe obesity. Am. J. Med. 2004;117:398–405.

27. Volek JS, Sharman MJ, Gómez AL, DiPasquale C, Roti M, Pumerantz A, et al. Comparison of a very

low-carbohydrate and low-fat diet on fasting lipids, LDL subclasses, insulin resistance, and postprandial

lipemic responses in overweight women. J Am Coll Nutr. 2004;23:177–84.

28. Volek JS, Phinney SD, Forsythe CE, Quann EE, Wood RJ, Puglisi MJ, et al. Carbohydrate Restriction

has a More Favorable Impact on the Metabolic Syndrome than a Low Fat Diet. Lipids. 2008;44:297–309.

29. Kannel WB, Anderson K, Wilson PW. White blood cell count and cardiovascular disease. Insights

from the Framingham Study. JAMA. 1992;267:1253–6.

30. Yarnell JW, Baker IA, Sweetnam PM, Bainton D, O'Brien JR, Whitehead PJ, et al. Fibrinogen,

viscosity, and white blood cell count are major risk factors for ischemic heart disease. The Caerphilly and

Speedwell collaborative heart disease studies. Circulation. 1991;83:836–44.

31. Folsom AR, Aleksic N, Catellier D, Juneja HS, Wu KK. C-reactive protein and incident coronary

heart disease in the Atherosclerosis Risk In Communities (ARIC) study. American Heart Journal.

2002;144:233–8.

32. Ridker PM, Buring JE, Shih J, Matias M, Hennekens CH. Prospective study of C-reactive protein and

the risk of future cardiovascular events among apparently healthy women. Circulation. 1998;98:731–3.

33. Libby P, Ridker PM, Hansson GK, Leducq Transatlantic Network on Atherothrombosis.

Inflammation in atherosclerosis: from pathophysiology to practice. Journal of the American College of

Cardiology. 2009;54:2129–38.

34. Biondi-Zoccai GGL, Abbate A, Liuzzo G, Biasucci LM. Atherothrombosis, inflammation, and

diabetes. Journal of the American College of Cardiology. 2003;41:1071–7.

35. Wong ND, Zhao Y, Quek RGW, Blumenthal RS, Budoff MJ, Cushman M, et al. Residual

atherosclerotic cardiovascular disease risk in statin-treated adults: The Multi-Ethnic Study of

Atherosclerosis. Journal of Clinical Lipidology. 2017;11:1223–33.

36. Sampson UK, Fazio S, Linton MF. Residual cardiovascular risk despite optimal LDL cholesterol

reduction with statins: the evidence, etiology, and therapeutic challenges. Curr Atheroscler Rep.

2012;14:1–10.

37. Group UPDS. Tight blood pressure control and risk of macrovascular and microvascular

complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. BMJ.

1998;317:703–13.

38. Schmieder RE, Ruilope L-M, Barnett AH. Renal protection with angiotensin receptor blockers: where

do we stand. J. Nephrol. 2011;24:569–80.

39. Jafar TH, Schmid CH, Landa M, Giatras I, Toto R, Remuzzi G, et al. Angiotensin-converting enzyme

inhibitors and progression of nondiabetic renal disease. A meta-analysis of patient-level data. Ann Intern

Med. 2001;135:73–87.

40. ADA. Standards of Medical Care in Diabetes - 2017. Diabetes Care; 2017;40.

41. Tremblay AJ, Morrissette H, Gagné J-M, Bergeron J, Gagné C, Couture P. Validation of the

Friedewald formula for the determination of low-density lipoprotein cholesterol compared with beta-

quantification in a large population. Clin. Biochem. 2004;37:785–90.

42. Jeyarajah EJ, Cromwell WC, Otvos JD. Lipoprotein particle analysis by nuclear magnetic resonance

spectroscopy. Clin. Lab. Med. 2006;26:847–70.

43. van Schalkwijk DB, de Graaf AA, Tsivtsivadze E, Parnell LD, van der Werff-van der Vat BJC, van

Ommen B, et al. Lipoprotein metabolism indicators improve cardiovascular risk prediction. Schäfer A,

editor. PLoS ONE. 2014;9:e92840.

44. May HT, Anderson JL, Winegar DA, Rollo J, Connelly MA, Otvos JD, et al. Utility of high density

lipoprotein particle concentration in predicting future major adverse cardiovascular events among patients

undergoing angiography. Clin. Biochem. 2016;49:1122–6.

45. Shalaurova I, Connelly MA, Garvey WT, Otvos JD. Lipoprotein Insulin Resistance Index: A

Lipoprotein Particle–Derived Measure of Insulin Resistance. Metabolic Syndrome and Related Disorders.

2014;12:422–9.

46. Goff DC, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB, Gibbons R, et al. 2013 ACC/AHA

guideline on the assessment of cardiovascular risk: a report of the American College of

Cardiology/American Heart Association Task Force on Practice Guidelines. Circulation. 2014. pp. S49–

73.

47. Doneen AL, Bale BF. Carotid intima-media thickness testing as an asymptomatic cardiovascular

disease identifier and method for making therapeutic decisions. Postgrad Med. 2013;125:108–23.

48. Stein JH, Korcarz CE, Hurst RT, Lonn E, Kendall CB, Mohler ER, et al. Use of Carotid Ultrasound to

Identify Subclinical Vascular Disease and Evaluate Cardiovascular Disease Risk: A Consensus Statement

from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force Endorsed

by the Society for Vascular Medicine. Journal of the American Society of Echocardiography.

2008;21:93–111.

49. UK Prospective Diabetes Study UKPDS Group. Intensive blood-glucose control with sulphonylureas

or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes

(UKPDS 33). The Lancet. 1998;352:837–53.

50. Volek JS, Sharman MJ, Forsythe CE. Modification of lipoproteins by very low-carbohydrate diets. J.

Nutr. 2005;135:1339–42.

51. Nordmann AJ, Nordmann A, Briel M, Keller U, Yancy WS, Brehm BJ, et al. Effects of Low-

Carbohydrate vs Low-Fat Diets on Weight Loss and Cardiovascular Risk Factors: A Meta-analysis of

Randomized Controlled Trials. Arch. Intern. Med. 2006;166:285–93.

52. Law MR, Wald NJ, Rudnicka AR. Quantifying effect of statins on low density lipoprotein cholesterol,

ischaemic heart disease, and stroke: systematic review and meta-analysis. BMJ. 2003;326:1423–0.

53. Giugliano RP, Pedersen TR, Park J-G, De Ferrari GM, Gaciong ZA, Ceska R, et al. Clinical efficacy

and safety of achieving very low LDL-cholesterol concentrations with the PCSK9 inhibitor evolocumab:

a prespecified secondary analysis of the FOURIER trial. Lancet. 2017;390:1962–71.

54. Zuliani G, Volpato S, Dugo M, Vigna GB, Morieri ML, Maggio M, et al. Combining LDL-C and

HDL-C to predict survival in late life: The InChianti study. Abete P, editor. PLoS ONE.

2017;12:e0185307.

55. Orozco-Beltran D, Gil-Guillen VF, Redon J, Martin-Moreno JM, Pallares-Carratala V, Navarro-Perez

J, et al. Lipid profile, cardiovascular disease and mortality in a Mediterranean high-risk population: The

ESCARVAL-RISK study. Vinciguerra M, editor. PLoS ONE. 2017;12:e0186196.

56. Ravnskov U, Diamond DM, Hama R, Hamazaki T, Hammarskjöld B, Hynes N, et al. Lack of an

association or an inverse association between low-density-lipoprotein cholesterol and mortality in the

elderly: a systematic review. BMJ Open. 2016;6:e010401.

57. Schmidt C, Bergstrom G. Apolipoprotein B and Apolipopotein A-I in Vascular Risk Prediction-A

Review. Curr Pharm Des. 2014;20:6289–98.

58. Austin MA, Breslow JL, Hennekens CH, Buring JE, Willett WC, Krauss RM. Low-density

lipoprotein subclass patterns and risk of myocardial infarction. JAMA. 1988;260:1917–21.

59. Tani S, Yagi T, Atsumi W, Kawauchi K, Matsuo R, Hirayama A. Relation between low-density

lipoprotein cholesterol/apolipoprotein B ratio and triglyceride-rich lipoproteins in patients with coronary

artery disease and type 2 diabetes mellitus: a cross-sectional study. Cardiovasc Diabetol. 2017;16:123.

60. Bertsch RA, Merchant MA. Study of the Use of Lipid Panels as a Marker of Insulin Resistance to

Determine Cardiovascular Risk. Perm J. 2015;19:4–10.

61. Ridker PM, Danielson E, Fonseca FAH, Genest J, Gotto AM Jr, Kastelein JJP, et al. Rosuvastatin to

Prevent Vascular Events in Men and Women with Elevated C-Reactive Protein. N Engl J Med.

2008;359:2195–207.

62. Albert MA, Danielson E, Rifai N, Ridker PM, PRINCE Investigators. Effect of statin therapy on C-

reactive protein levels: the pravastatin inflammation/CRP evaluation (PRINCE): a randomized trial and

cohort study. JAMA. 2001;286:64–70.

63. Asher J, Houston M. Statins and C-reactive protein levels. J Clin Hypertens (Greenwich).

2007;9:622–8.

64. Youm Y-H, Nguyen KY, Grant RW, Goldberg EL, Bodogai M, Kim D, et al. The ketone metabolite

β-hydroxybutyrate blocks NLRP3 inflammasome–mediated inflammatory disease. Nat. Med.

2015;21:263–9.

65. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison Himmelfarb C, et al. 2017

ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention,

Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American

College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. Journal

of the American College of Cardiology. 2017.

66. Ettehad D, Emdin CA, Kiran A, Anderson SG, Callender T, Emberson J, et al. Blood pressure

lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis.

Lancet. 2016;387:957–67.

67. Kuznik A, Mardekian J. Trends in utilization of lipid- and blood pressure-lowering agents and goal

attainment among the U.S. diabetic population, 1999-2008. Cardiovasc Diabetol. 2011;10:31.

68. Shai I, Spence JD, Schwarzfuchs D, Henkin Y, Parraga G, Rudich A, et al. Dietary intervention to

reverse carotid atherosclerosis. Circulation. 2010;121:1200–8.

69. Gress TW, Nieto FJ, Shahar E, Wofford MR, Brancati FL. Hypertension and antihypertensive therapy

as risk factors for type 2 diabetes mellitus. Atherosclerosis Risk in Communities Study. N. Engl. J. Med.

2000;342:905–12.

70. Lorenz MW, Price JF, Robertson C, Bots ML, Polak JF, Poppert H, et al. Carotid intima-media

thickness progression and risk of vascular events in people with diabetes: results from the PROG-IMT

collaboration. Diabetes Care. 2015;38:1921–9.

71. Riley WA. Cardiovascular risk assessment in individual patients from carotid intimal-medial

thickness measurements. Curr Atheroscler Rep. 2004;6:225–31.

Figure Legends

Figure 1. Change in biomarkers for CCI group.

Bars indicate CCI group mean percent change in biomarkers based on the intention-to-treat analysis with

missing values imputed. Percent change is computed as the change in mean values from baseline to one

year divided by the mean baseline value. *Percent change* = 100*[(*1 year value - baseline*

value)/(*baseline*)]. Negative values indicate a decrease from baseline to one year while positive values

indicate an increase. The ** symbol after the biomarker label indicates a statistically significant within

group change from baseline (P<0.0019, P adjusted for multiple comparisons). Error bars represent ± SE.

SE as Percent = 100*[(*1 year value SE*)/(*baseline*)]

Figure 2. Change in biomarkers for UC group.

Bars indicate UC group mean percent change in biomarkers based on the intention-to-treat analysis with

missing values imputed. Percent change is computed as the change in mean values from baseline to one

year divided by the mean baseline value. *Percent change* = 100*[(*1 year value - baseline*

value)/(*baseline*)]. Negative values indicate a decrease from baseline to one year while positive values

indicate an increase. (None of the within group changes were statistically significant, i.e. all P>0.0019, P

adjusted for multiple comparisons.) Error bars represent ± SE. *SE as Percent* = 100*[(*1 year value*

SE)/(*baseline*)]

Supplementary Figure 1. Distribution of changes in selected biomarkers for CCI and UC

completers

Histograms of changes at one year for CCI (blue) and UC (gray) are overlaid. Very few (≤1%) CCI

participants demonstrated changes in an undesirable direction at one year that was outside the range of

changes observed in the UC group for key lipid and lipoprotein particles. (A) Apolipoprotein B (B)

Apolipoprotein A1 (C) Apolipoprotein B/Apolipoprotein A1 ratio (D) LDL-P (E) Small LDL-P (F) Large

VLDL-P (G) Triglyceride/HDL-C ratio

	All		Completers with data	
Table 1. Baseline characteristics Continuous Care Intervention (CCI) and Usual Care (UC) groups.				
	N	**Mean (SD) or ±SE**	**N**	**Mean (SD) or ±SE**
Age (years)				
CCI-all education[1]	262	54 (8)	218	54 (8)
Usual care[1]	87	52 (10)	78	52 (10)
CCI-all vs. usual care[2]		1 ± 1		2 ± 1*
Female (%)				
CCI-all education[1]	262	66.8 ± 2.9	218	65.1 ± 3.2
Usual care[1]	87	58.6 ± 5.3	78	60.3 ± 5.5
CCI-all vs. usual care[2]		8.2 ± 6.0		4.9 ± 6.4
Smokers (%)				
CCI-all education[1]	211	3.8 ± 1.3	176	4.0 ± 1.5
Usual care[1]	87	14.9 ± 3.8	78	14.1 ± 3.9
CCI-all vs. usual care[2]		-11.2 ± 4.0†		-10.1 ± 4.2*
Weight-clinic (kg)				
CCI-all education[1]	257	116.5 (25.9)	184	115.4 (24.6)
Usual care[1]	83	105.6 (22.2)	69	106.8 (22.2)
CCI-all vs. usual care[2]		10.9 ± 2.9‡		8.6 ± 3.2†
BMI (kg·m^{-2})				
CCI-all education[1]	257	40.4 (8.8)	184	39.9 (7.9)
Usual care[1]	83	36.7 (7.3)	69	37.1 (7.6)
CCI-all vs. usual care[2]		3.7 ± 1.0‡		2.7 ± 1.1†
Hemoglobin A1c (%)				
CCI-all education[1]	262	7.60 (1.50)	204	7.49 (1.40)
Usual care[1]	87	7.64 (1.76)	72	7.74 (1.82)
CCI-all vs. usual care[2]		-0.04 ± 0.21		-0.25 ± 0.24

Systolic blood pressure (mmHg)				
CCI-all education[1]	260	132 (14)	187	133 (15)
Usual care[1]	79	130 (14)	67	129 (13)
CCI-all vs. usual care[2]		2 ± 2		4 ± 2*
Diastolic blood pressure (mmHg)				
CCI-all education[1]	260	82 (8)	187	82 (8)
Usual care[1]	79	82 (9)	67	81 (8)
CCI-all vs. usual care[2]		0 ± 1		0 ± 1
ApoB (mg·dL-1)				
CCI-all education[1]	248	105 (29)	186	103 (28)
Usual care[1]	79	107 (28)	59	106 (30)
CCI-all vs. usual care[2]		-2 ± 4		-2
ApoA1 (mg·dL-1)				
CCI-all education[1]	248	146 (28)	185	146 (29)
Usual care[1]	79	149 (22)	59	148 (21)
CCI-all vs. usual care[2]		-3 ± 3		-2 ± 3
ApoB/ApoA1 ratio				
CCI-all education[1]	248	0.74 (0.23)	185	0.73 (0.23)
Usual care[1]	79	0.73 (0.23)	59	0.73 (0.25)
CCI-all vs. usual care[2]		0.01 ± 0.03		0.00 ± 0.04
Triglycerides (mg·dL^{-1})				
CCI-all education[1]	247	197 (143)	186	201 (153)
Usual care[1]	79	283 (401)	59	297 (458)
CCI-all vs. usual care[2]		-86 ± 46*		-97 ± 61
LDL-C (mg·dL^{-1})				
CCI-all education[1]	232	103 (33)	172	100 (33)
Usual care[1]	70	102 (36)	48	100 (38)

CCI-all vs. usual care[2]		1 ± 5		0 ± 6
HDL-C (mg·dL^{-1})				
CCI-all education[1]	247	42 (13)	186	42 (14)
Usual care[1]	79	38 (11)	59	37 (11)
CCI-all vs. usual care[2]		5 ± 2†		5 ± 2†
Triglycerides/HDL-C ratio				
CCI-all education[1]	247	5.9 (7.1)	186	6.1 (7.9)
Usual care[1]	79	10.5 (23.2)	59	11.5 (26.5)
CCI-all vs. usual care[2]		-4.6 ± 2.6		-5.4 ± 3.5
Large VLDL-P (nmol·L^{-1})				
CCI-all education[1]	259	10 (8)	203	9 (8)
Usual care[1]	83	12 (12)	68	12 (13)
CCI-all vs. usual care[2]		-2 ± 1		-2 ± 2
Total LDL-P (nmol·L^{-1})				
CCI-all education[1]	259	1300 (465)	203	1296 (476)
Usual care[1]	83	1289 (511)	68	1243 (484)
CCI-all vs. usual care[2]		11 ± 63		52 ± 68
Small LDL-P (nmol·L^{-1})				
CCI-all education[1]	259	774 (377)	203	778 (378)
Usual care[1]	83	719 (322)	68	699 (326)
CCI-all vs. usual care[2]		55 ± 42		789 ± 48
LDL-particle size (nm)				
CCI-all education[1]	259	20.30 (0.55)	201	20.3 (0.55)
Usual care[1]	83	20.33 (0.56)	68	20.32 (0.55)
CCI-all vs. usual care[2]		-0.03 ± 0.07		-0.03 ± 0.08
Total HDL-P (μmol·L^{-1})				
CCI-all education[1]	259	31.3 (6.4)	203	31.7 (6.4)
Usual care[1]	83	29.9 (5.8)	68	30.2 (6.0)

CCI-all vs. usual care[2]		1.4 ± 0.8		1.5 ± 0.9
Large HDL-P (µmol·L[-1])				
CCI-all education[1]	259	4.3 (2.5)	203	4.2 (2.5)
Usual care[1]	83	3.8 (2.1)	68	3.8 (2.1)
CCI-all vs. usual care[2]		0.4 ± 0.3		0.4 ± 0.3
LP-IR score				
CCI-all education[1]	259	72 (17)	203	72 (18)
Usual care[1]	83	75 (16)	68	74 (17)
CCI-all vs. usual care[2]		-3 ± 2		-2 ± 2
C-reactive protein (mg·L[-1])				
CCI-all education[1]	249	8.5 (14.5)	193	9.0 (16.1)
Usual care[1]	85	8.9 (8.6)	70	9.1 (9.0)
CCI-all vs. usual care[2]		-0.3 ± 1.3		-0.1 ± 1.6
WBC				
CCI-all education[1]	260	7.2 (1.9)	204	7.1 (1.8)
Usual care[1]	86	8.1 (2.4)	72	8.3 (2.4)
CCI-all vs. usual care[2]		-0.9 ± 0.3†		-1.2 ± 0.3§
10-year ASCVD risk (%)				
CCI-all education[1]	198	11.1 (9.1)	135	12.1 (9.3)
Usual care[1]	72	11.8 (10.8)	55	11.4 (10.8)
CCI-all vs. usual care[2]		-0.6 ± 1.4		0.8 ± 1.6
CIMT-average (mm)				
CCI-all education[1]	236	0.681 (0.108)	144	0.692 (0.113)
Usual care[1]	84	0.681 (0.116)	68	0.680 (0.111)
CCI-all vs. usual care[2]		-0.001 ± 0.014		0.013 ± 0.016
Statin (%)				
CCI-all education[1]	262	50.0 ± 3.1	218	51.8 ± 3.4
Usual care[1]	87	58.6 ± 5.3	73	54.8 ± 5.8

CCI-all vs. usual care[2]		-8.6 ± 6.1		-3.0 ± 6.7
Any antihypertensive medication (%)				
CCI-all education[1]	262	67.2 ± 2.9	218	68.4 ± 3.2
Usual care[1]	87	52.9 ± 5.4	73	50.7 ± 5.9
CCI-all vs. usual care[2]		14.3 ± 6.1*		17.7 ± 6.7†
ACE or ARB (%)				
CCI-all education[1]	262	29.4 ± 2.8	218	28.0 ± 3.0
Usual care[1]	87	18.4 ± 4.2	73	16.4 ± 4.3
CCI-all vs. usual care[2]		11.0 ± 5.0*		11.5 ± 5.3*
Diuretics (%)				
CCI-all education[1]	262	40.8 ± 3.0	218	41.3 ± 3.3
Usual care[1]	87	29.9 ± 4.9	73	24.7 ± 5.0
CCI-all vs. usual care[2]		11.0 ± 5.8		16.6 ± 6.1†

[1]Mean and standard deviations for continuous variables, percents and standard errors for categorical variables

[2]Difference between means or percentages ± 1 standard error of the difference. Significant baseline difference between means or percentages at 0.05≤P<0.01 (*); 0.01≤P<0.001 (†); 0.001≤P<0.0001 (‡); and P<0.0001 (§).

Table 2. One-year Biomarker Change Continuous Care Intervention compared to Usual Care groups.									

		Completers					All starters (Dropouts imputed)[4]		
	N		Unadjusted		Adjusted for baseline[3]		Unadjusted		
		One Year Mean ±SE	Difference (SD) or ±SE	Signif-icance[5]	Difference ±SE	Signif-icance[5]	One Year Mean ±SE	Difference ±SE	Signif-icance[5]
ΔWeight-clinic (kg)									
CCI-all education[1]	184	101.2 ± 1.6	-14.2 (10.3)	$<10^{-16}$	-13.8 ± 0.6	$<10^{-16}$	102.7 ± 1.5	-13.8 ± 0.7	$<10^{-16}$
Usual care[1]	69	106.8 ± 2.7	0.04 (5.9)	0.95	-1.1 ± 1.1	0.29	107.3 ± 2.6	-0.2 ± 0.8	0.85
CCI-all vs. usual care[2]			-14.3 ± 1.0	$<10^{-16}$	-12.7 ± 1.3	$<10^{-16}$		-13.7 ± 1.1	$<10^{-16}$
ΔHemoglobin A1c (%)									
CCI-all education[1]	204	6.20 ± 0.07	-1.29 (1.32)	$<10^{-16}$	-1.32 ± 0.09	$<10^{-16}$	6.29 ± 0.07	-1.30 ± 0.09	$<10^{-16}$
Usual care[1]	72	7.94 ± 0.22	0.20 (1.35)	0.21	0.22 ± 0.16	0.17	7.84 ± 0.19	0.20 ± 0.15	0.18
CCI-all vs. usual care[2]			-1.49 ± 0.18	4.4×10^{-16}	-1.54 ± 0.19	4.4×10^{-16}		-1.50 ± 0.17	$<10^{-16}$
ΔSystolic blood pressure (mmHg)									
CCI-all education[1]	187	126 ± 1	-7 (16)	1.3×10^{-8}	-7 ± 1	1.6×10^{-7}	126 ± 1	-6 ± 1	1.3×10^{-8}
Usual care[1]	67	129 ± 2	0 (18)	0.91	0 ± 2	0.83	129 ± 2	-1 ± 2	0.67
CCI-all vs. usual care[2]			-7 ± 2	0.005	-6 ± 3	0.02		-5 ± 2	0.02
ΔDiastolic blood pressure (mmHg)									

CCI-all education[1]	187	78 ± 1	-4 (9)	1.4×10^{-7}	-4 ± 1	6.2×10^{-7}	79 ± 1	-4 ± 1	7.2×10^{-8}
Usual care[1]	67	81 ± 1	0 (10)	0.92	0 ± 1	0.75	81 ± 1	-1 ± 1	0.45
CCI-all vs. usual care[2]			-3 ± 1	0.01	-3 ± 1	0.03		-3 ± 1	0.06
ΔApoB (mg·dL-1)									
CCI-all education[1]	186	103 ± 2	-1 (24)	0.69	-0 ± 2	0.82	104 ± 2	-2 ± 2	0.37
Usual care[1]	59	107 ± 5	2 (37)	0.75	1 ± 4	0.9	106 ± 4	0 ± 4	0.95
CCI-all vs. usual care[2]			-2 ± 5	0.66	-1 ± 5	0.83		-2 ± 5	0.67
ΔApoA1 (mg·dL-1)									
CCI-all education[1]	185	160 ± 3	14 (24)	8.9×10^{-16}	14 ± 2	4.4×10^{-16}	160 ± 2	14 ± 2	$<10^{-16}$
Usual care[1]	59	145 ± 3	-3 (19)	0.18	-2 ± 3	0.55	147 ± 3	-2 ± 3	0.37
CCI-all vs. usual care[2]			18 ± 3	4.7×10^{-9}	16 ± 4	2.2×10^{-5}		17 ± 3	1.4×10^{-7}
ΔApoB/ApoA1									
CCI-all education[1]	185	0.67 ± 0.02	-0.06 (0.17)	1.8×10^{-6}	-0.06 ± 0.02	0.003	0.67 ± 0.02	-0.07 ± 0.01	1.9×10^{-7}
Usual care[1]	59	0.76 ± 0.04	0.03 (0.29)	0.42	0.02 ± 0.03	0.5	0.74 ± 0.03	0.02 ± 0.03	0.58
CCI-all vs. usual care[2]			-0.09 ± 0.04	0.02	-0.08 ± 0.03	0.02		-0.09 ± 0.03	0.01
ΔTriglycerides (mg·dL^{-1})									
CCI-all education[1]	186	151 ± 11	-49 (168)	5.6×10^{-5}	-50 ± 16	0.001	148 ± 12	-48 ± 13	$<10^{-16}$
Usual care[1]	59	327 ± 65	30 (301)	0.44	31 ± 29	0.27	305 ± 48	28 ± 32	0.43
CCI-all vs. usual care[2]			-80 ± 41	0.05	-81 ± 33	0.02		-76 ± 35	9.9×10^{-7}
ΔLDL-C (mg·dL^{-1})									

CCI-all education[1]	172	111 ± 3	11 (32)	7.7×10^{-6}	11 ± 3	2.6×10^{-5}	113 ± 3	10 ± 2	4.9×10^{-5}
Usual care[1]	48	90 ± 4	-11 (38)	0.05	-11 ± 5	0.03	90 ± 5	-11 ± 5	0.02
CCI-all vs. usual care[2]			22 ± 6	0.0003	22 ± 6	0.0002		21 ± 5	9.9×10^{-5}
ΔHDL-C (mg·dL^{-1})									
CCI-all education[1]	186	50 ± 1	8 (12)	$<10^{-16}$	7 ± 1	$<10^{-16}$	50 ± 1	8 ± 1	$<10^{-16}$
Usual care[1]	59	35 ± 2	-2 (9)	0.15	-1 ± 2	0.69	37 ± 2	-1 ± 1	0.41
CCI-all vs. usual care[2]			9 ± 1	1.7×10^{-10}	8 ± 2	9.9×10^{-6}		9 ± 2	1.2×10^{-8}
ΔTriglycerides/HDL-C ratio									
CCI-all education[1]	186	4.3 ± 0.6	-1.8 (9.4)	$<10^{-16}$	-1.9 ± 0.9	$<10^{-16}$	4.1 ± 0.6	-1.6 ± 0.7	$<10^{-16}$
Usual care[1]	59	12.5 ± 2.7	0.9 (16.1)	0.1	1.2 ± 1.6	0.16	11.2 ± 2.1	1.0 ± 1.7	0.24
CCI-all vs. usual care[2]			-2.8 ± 2.2	3.1×10^{-10}	-3.1 ± 1.8	5.5×10^{-7}		-2.6 ± 1.8	4.5×10^{-9}
ΔLarge VLDL-P (nmol·L^{-1})									
CCI-all education[1]	203	6 ± 1	-4 (7)	5.6×10^{-15}	-4 ± 1	1.6×10^{-14}	6 ± 1	-4 ± 1	4.2×10^{-15}
Usual care[1]	68	12 ± 2	0 (8)	0.71	0 ± 1	0.60	12 ± 1	0 ± 1	0.77
CCI-all vs. usual care[2]			-3 ± 1	0.001	-3 ± 1	0.002		3 ± 1	0.0007
ΔTotal LDL-P (nmol·L^{-1})									
CCI-all education[1]	203	1234 ± 30	-62 (375)	0.02	-57 ± 29	0.05	1235 ± 29	-64 ± 26	0.02
Usual care[1]	68	1196 ± 60	-47 (491)	0.43	-67 ± 53	0.21	1231 ± 57	-57 ± 56	0.31
CCI-all vs. usual care[2]			-15 ± 65	0.82	10 ± 62	0.87		-7 ± 62	0.91
ΔSmall LDL-P (nmol·L^{-1})									

	N								
CCI-all education[1]	203	614 ± 22	-164 (332)	2.2×10^{-12}	-161 ± 24	4.1×10^{-11}	613 ± 21	-161 ± 23	1.2×10^{-12}
Usual care[1]	68	724 ± 44	25 (370)	0.57	16 ± 45	0.71	740 ± 41	18 ± 42	0.67
CCI-all vs. usual care[2]			-189 ± 51	0.0002	-177 ± 52	0.0007		-179 ± 48	0.0002
ΔLDL-particle size (nm)									
CCI-all education[1]	201	20.53 ± 0.04	0.23 (0.54)	1.7×10^{-9}	0.23 ± 0.04	8.9×10^{-9}	20.53 ± 0.04	0.23 ± 0.04	6.0×10^{-10}
Usual care[1]	68	20.25 ± 0.07	-0.08 (0.53)	0.24	-0.08 ± 0.07	0.25	20.25 ± 0.07	-0.07 ± 0.06	0.25
CCI-all vs. usual care[2]			0.30 ± 0.07	4.4×10^{-5}	0.31 ± 0.08	0.0002		0.30 ± 0.07	3.8×10^{-15}
ΔTotal HDL-P (μmol·L⁻¹)									
CCI-all education[1]	203	33.2 ± 0.5	1.5 (4.9)	1.2×10^{-5}	1.5 ± 0.4	2.1×10^{-5}	32.8 ± 0.4	1.5 ± 0.3	5.6×10^{-6}
Usual care[1]	68	29.4 ± 0.8	-0.8 (4.7)	0.15	-0.8 ± 0.6	0.23	29.2 ± 0.7	-0.7 ± 0.6	0.23
CCI-all vs. usual care[2]			2.3 ± 0.7	0.0004	2.3 ± 0.7	0.003		2.2 ± 0.7	0.0008
ΔLarge HDL-P (μmol·L⁻¹)									
CCI-all education[1]	203	5.3 ± 0.2	1.0 (2.2)	2.5×10^{-11}	1.0 ± 0.2	4.1×10^{-11}	5.3 ± 0.2	1.0 ± 0.2	1.2×10^{-11}
Usual care[1]	68	3.9 ± 0.3	0.1 (1.6)	0.69	0.2 ± 0.3	0.44	3.9 ± 0.3	0.1 ± 0.2	0.74
CCI-all vs. usual care[2]			0.9 ± 0.3	0.0002	0.8 ± 0.3	0.01		0.9 ± 0.3	0.0004
ΔLP-IR score									
CCI-all education[1]	203	58 ± 2	-14 (18)	$<10^{-16}$	-14 ± 1	$<10^{-16}$	58 ± 1	-14 ± 1	$<10^{-16}$
Usual care[1]	68	74 ± 2	-1 (16)	0.73	-2 ± 2	0.41	75 ± 2	-1 ± 2	0.74
CCI-all vs. usual care[2]			-13 ± 2	3.8×10^{-9}	-12 ± 3	6.2×10^{-6}		-13 ± 2	6.2×10^{-9}
ΔC-reactive protein (mg·L⁻¹)									

CCI-all education[1]	193	5.7 ± 0.5	-3.3 (13.4)	$<10^{-8}$	-3.1 ± 1.0	$<10^{-16}$	5.6 ± 0.6	-3.6 ± 1.1	$<10^{-16}$
Usual care[1]	70	10.4 ± 1.8	1.3 (13.3)	0.94	0.9 ± 1.7	0.88	10.3 ± 1.6	1.3 ± 1.5	0.93
CCI-all vs. usual care[2]			-4.7 ±1.9	1.2×10^{-6}	-4.0 ± 2.0	3.0×10^{-5}		-4.9 ± 1.8	9.3×10^{-7}
ΔWBC (k·mm⁻³)									
CCI-all education[1]	204	6.5 ± 0.1	-0.7 (1.4)	2.1×10^{-11}	-0.7 ± 0.1	2.1×10^{-11}	6.6 ± 0.1	-0.7 ± 0.1	3.2×10^{-11}
Usual care[1]	72	8.3 ± 0.3	-0.1 (1.6)	0.76	-0.1 ± 0.2	0.74	8.1 ± 0.3	-0.1 ± 0.2	0.76
CCI-all vs. usual care[2]			-0.6 ± 0.2	0.003	-0.6 ± 0.2	0.004		-0.6 ± 0.2	0.003
Δ10-year ASCVD risk (%)									
CCI-all education[1]	135	10.5 ± 0.7	1.6 (5.4)	0.0004	-1.5 ± 0.6	0.01	9.6 ± 0.5	-1.3 ± 0.3	4.9×10^{-5}
Usual care[1]	55	12.7 ± 1.5	1.4 (9.3)	0.28	1.1 ± 1.0	0.27	12.9 ± 1.2	1.2 ± 0.9	0.17
CCI-all vs. usual care[2]			03.0 ± 1.3	0.03	-2.6 ± 1.2	0.03		-2.6 ± 1.0	0.008
ΔCIMT-average (mm)									
CCI-all education[1]	144	0.695 ± 0.009	0.002 (0.055)	0.63	0.003 ± 0.004	0.45	0.685 ± 0.010	0.002 ± 0.004	0.65
Usual care[1]	68	0.680 ± 0.013	0.004 (0.041)	0.37	0.002 ± 0.006	0.74	0.680 ± 0.013	0.001 ± 0.006	0.87
CCI-all vs. usual care[2]			-0.002 ± 0.007	0.74	0.001 ± 0.008	0.87		0.001 ± 0.007	0.88
ΔStatin (%)									
CCI-all education[1]	218	48.2 ± 3.4	-3.7 (34.4)	0.12	-3.6 ± 2.4	0.13	46.7 ± 3.2	-3.3 ± 2.3	0.15
Usual care[1]	73	64.4 ± 5.6	9.6 (37.9)	0.03	9.5 ± 4.3	0.03	67.4 ± 5.4	8.8 ± 4.3	0.04
CCI-all vs. usual care[2]			-13.3 ± 5.0	0.008	-13.2 ± 5.0	0.009		-12.1 ± 4.9	0.01
ΔAny antihypertensive medication (%)									

CCI-all education[1]	218	56.4 ± 3.4	-11.9 (42.3)	3.2×10^{-5}	-11.9 ± 2.9	3.6×10^{-5}	55.8 ± 3.3	-11.4 ± 2.8	5.3×10^{-5}
Usual care[1]	73	60.3 ± 5.8	9.6 (41.4)	0.05	9.6 ± 5.1	0.06	61.2 ± 5.6	8.3 ± 4.8	0.09
CCI-all vs. usual care[2]			-21.5 ± 5.6	0.0002	-21.6 ± 6.0	0.0004		-19.7 ± 5.6	0.0004
ΔACE or ARB (%)									
CCI-all education[1]	218	28.9 ± 3.1	0.9 (27.1)	0.62	1.5 ± 1.9	0.42	30.0 ± 2.9	0.6 ± 1.9	0.76
Usual care[1]	73	21.9 ± 4.9	5.5 (28.3)	0.1	3.7 ± 3.3	0.27	23.4 ± 4.7	5.0 ± 3.3	0.13
CCI-all vs. usual care[2]			-4.6 ± 3.8	0.23	-2.1 ± 3.9	0.59		-4.4 ± 3.8	0.24
ΔDiuretics (%)									
CCI-all education[1]	218	31.7 ± 3.2	-9.6 (41.3)	0.0006	-9.5 ± 2.7	0.0004	31.3 ± 3.1	-9.7 ± 2.7	0.0004
Usual care[1]	73	30.1 ± 5.4	5.5 (32.9)	0.16	5.2 ± 4.8	0.28	33.0 ± 5.3	3.2 ± 4.1	0.44
CCI-all vs. usual care[2]			-15.1 ± 4.8	0.001	-14.7 ± 5.6	0.009		-12.8 ± 4.9	0.009

[1] Means (standard deviations) are presented. Sample sizes, means, and significance levels refer to subjects with baseline and one-year measurements for *completers*, and to 349 subjects (262 intervention and 87 usual care) for *all starters*. Significance levels for *completers* refer to one-sample t-test with or without adjustment. Untransformed triglyceride and C-reactive protein values are presented, however, their statistical significances were based on their log-transformed values. All CC refers to the CCI-web and CCI-onsite combined.

[2] Mean differences ± one standard error. Significance levels refer to two-sample t-test or analysis of covariance for the differences.

[3] Adjusted for sex, age, baseline BMI, baseline insulin use (user vs. non-user), and African-American race.

[4] Imputed values based on 700 iterations from multivariate normal regression.

[5] A significance level of $P<0.0019$ ensures overall simultaneous significance of $P\leq0.05$ over the 26 variables using Bonferroni correction.

647
648

Intention-to-treat (Missing Values Imputed)

Intention-to-treat (Missing Values Imputed)