Environmental determinants of pyoverdine production, exploitation and competition in natural *Pseudomonas* communities

Elena Butaitė[1], Jos Kramer[1], Stefan Wyder[1] and Rolf Kümmerli[1]

1 Department of Plant and Microbial Biology, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

Corresponding author: Rolf Kümmerli, Department of Plant and Microbial Biology, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland; phone: +41 (0)44 63 54801; fax: +41 (0)44 63 48204; email: rolf.kuemmerli@uzh.ch

14   **Summary**

Many bacteria rely on the secretion of siderophores to scavenge iron from the environment.

16   Laboratory studies revealed that abiotic and biotic factors together determine how much

siderophores bacteria make, and whether siderophores can be exploited by non-producing

18   cheaters or be deployed by producers to inhibit competitors. Here, we explore whether these

insights apply to natural communities, by comparing the production of the siderophore

20   pyoverdine among 930 *Pseudomonas* strains from 48 soil and pond communities. We found

that pH, iron content, carbon concentration, and community diversity determine pyoverdine

22   production levels, and the extent to which strains are either stimulated or inhibited by

heterologous pyoverdines. While pyoverdine non-producers occurred in both habitats, their

24   prevalence was higher in soils. Environmental and genetic analysis suggest that non-

producers can evolve as cheaters, exploiting heterologous pyoverdine, but also due to

26   pyoverdine disuse in environments with increased iron availability. Overall, we found that

environmental factors explained between-strain variation in pyoverdine production much

28   better in soils than in ponds, presumably because high strain mixing in ponds prevents local

adaption. Our study sheds light on the complexity of natural bacterial communities, and

30   provides first insights into the multivariate nature of siderophore-based iron acquisition and

competition among environmental pseudomonads.

Introduction

Iron is a key growth-limiting factor for most bacteria. It is required for various fundamental cellular processes, such as DNA biosynthesis and respiration (Andrews *et al.*, 2003). However, iron is often limited in nature because it is either insoluble in its ferric (Fe^{+3}) form at circumneutral pH and aerobic conditions, host-bound in the context of infections, or occurs only at very low concentrations in habitats like the ocean (Andrews *et al.*, 2003; Miethke and Marahiel, 2007; Boyd and Ellwood, 2010). To overcome iron limitation, many bacteria secrete siderophores, iron-scavenging molecules that have a high affinity for ferric iron (Neilands, 1981; Hider and Kong, 2010). These molecules bind iron from natural sources and can then be taken up by cells via specific receptors. Given the importance of iron, it comes as no surprise that bacteria have evolved sophisticated mechanisms to accurately adjust the level of siderophore production to match prevailing levels of iron limitation (Andrews *et al.*, 2003; Parrow *et al.*, 2013; Frawley and Fang, 2014).

In addition to their role in provisioning producers with iron, siderophores can also have fitness consequences for other community members, including non-producers and individuals with different siderophore systems (Loper and Henkels, 1999; D'Onofrio *et al.*, 2010; Cordero *et al.*, 2012; Traxler *et al.*, 2012; Bruce, Cooper, *et al.*, 2017; Butaitė *et al.*, 2017). This is because siderophores can be shared between individuals with compatible receptors, which can select for exploitation by cheating strains that do not produce or produce less of the specific siderophore, yet capitalize on the siderophores produced by others (West *et al.*, 2006; Jiricny *et al.*, 2010). Moreover, siderophores can lock iron away from competitors with incompatible receptors, and can therefore be involved in inter-strain competition (Buyer and Leong, 1986; Joshi *et al.*, 2006; Butaitė *et al.*, 2017; Niehus *et al.*, 2017; Schiessl *et al.*, 2017; Sexton *et al.*, 2017). Although laboratory studies have uncovered many of the molecular,

ecological and evolutionary aspects of fine-tuned siderophore regulation and siderophore-

58 mediated social interactions, we know surprisingly little about the determinants of

siderophore production under natural conditions in complex multi-species communities.

Here, we address this issue by examining how environmental factors of natural soil and pond

62 habitats relate to the amount of siderophores produced by their inhabitants, and the ability of

natural isolates to affect other members of their community via their siderophores. In our

64 study, we focus on the phylogenetically diverse group of fluorescent pseudomonads, whose

members are known to produce the green-fluorescent pyoverdine as their primary siderophore

66 (Meyer *et al.*, 2008; Cornelis, 2010; Cézard *et al.*, 2015). From laboratory studies,

predominantly carried out with *Pseudomonas aeruginosa*, we know that iron concentration,

68 organic carbon composition, pH, and community diversity all influence the level of

pyoverdine production. Specifically, bacteria gradually downscale pyoverdine production in

70 response to increased iron concentration (Tiburzi *et al.*, 2008; Kümmerli *et al.*, 2009), when

iron is bound to relatively weak organic chelators (Dumas *et al.*, 2013), and at low pH where

72 solubility of iron is increased (Loper and Henkels, 1997). In addition, *P. aeruginosa*

upregulates pyoverdine production in the presence of competing strains or species (Harrison

74 *et al.*, 2008; Kümmerli *et al.*, 2009). Interestingly, pyoverdine-mediated social interactions

were found to be influenced by similar environmental factors. For example, increased iron

76 availability allows producers to down-scale pyoverdine production, which lowers their

susceptibility to exploitation by non-producers (Kümmerli *et al.*, 2009). Similarly, increased

78 carbon availability reduces the relative metabolic costs of pyoverdine production, and thus

decreases the advantage of cheaters (Brockhurst *et al.*, 2008; Sexton and Schuster, 2017).

80 Finally, community composition was found to be a key factor determining who interacts with

whom, thereby influencing the relative importance of competition versus cooperation (Griffin

82 *et al.*, 2004; Inglis *et al.*, 2016; Leinweber *et al.*, 2017).

84 To examine whether these factors influence pyoverdine production and pyoverdine-mediated

social interactions in natural communities, we isolated a total of 930 pseudomonads from 24

86 soil and 24 pond communities. We measured the pyoverdine production of all isolates in

standard iron-limited medium, and related this measure to the total iron and carbon

88 concentrations as well as the pH of their environment, and to the phylogenetic diversity of

their community. For a subset of strains, we additionally quantified the effect of pyoverdine-

90 containing supernatant on the fitness of co-occurring strains. For these experiments, we

focused on pyoverdine non-producers as supernatant-recipients because the stimulation or

92 inhibition of their growth under iron-limited conditions provides direct evidence for whether

or not they possess matching receptors for the uptake of heterologous siderophores (Butaitė *et*

94 *al.*, 2017).

96 **Results**

Pyoverdine production profiles differ between soil and pond pseudomonads

98 Prior to phenotypic screening, we sequenced the housekeeping *rpoD* gene, commonly used

for phylogenetic affiliation of pseudomonads, of all 930 isolates to confirm that they indeed

100 belong to this taxonomic group (Fig. S1). Once this was confirmed, we measured the

pyoverdine production of all isolates (using the natural fluorescence of this molecule) in iron-

102 limited casamino acids (CAA) medium supplemented with transferrin as an iron chelator. We

found that isolates greatly varied in the amount of pyoverdine they make (Fig. 1A). While

104 some strains produced no measurable amount of pyoverdine, others produced much more than

our laboratory reference strains (pyoverdine values are scaled relative to the average

106   production level of the laboratory reference strains listed in Table S1). Across all isolates,

there was a positive correlation between overall pyoverdine levels and their growth in iron-

108   limited CAA, suggesting that pyoverdine is important to overcome iron limitation (linear

mixed model, LMM: $t_{917.7} = 18.62$, $p < 0.001$; Fig. 1B).

There were several significant differences between soil and pond isolates. First, the relative

112   pyoverdine production was significantly lower for soil than for pond isolates (mean ± SE for

soil isolates: 0.571 ± 0.022; for pond isolates: 0.832 ± 0.018; LMM: $t_{21.5} = -4.36$, p < 0.001;

114   Fig. 1A). Second, there were significantly more pyoverdine non-producers (defined as

producing less than 5% of laboratory reference strains) in soil than in pond communities

116   (19.7% vs. 8.3%; Fisher's exact test: $p < 0.0001$; maximal non-producer frequency per

community was 75% and 20% in soils and ponds, respectively). Finally, the coefficient of

118   variation (CV) in relative pyoverdine production among isolates from the same community

was significantly higher in soil than in pond communities (mean CV ± SE for soil

120   communities: $85.2 \pm 8.3\%$; for pond communities: $46.2 \pm 2.4\%$; LMM: $t_{22.0} = 4.39$, $p < 0.001$;

Fig. S2).

To explore the relationship between phylogenetic diversity and pyoverdine production, we

124   constructed maximum likelihood phylogenetic trees based on partial *rpoD* gene sequences for

soil and pond isolates (Fig. S1). We found that phylogenetic diversity (normalized by the

126   number of isolates per community) was significantly lower in soil than pond communities

(Faith's phylogenetic diversity, median ± [1st quartile | 3rd quartile], for soil communities:

128     $0.09 \pm [0.07 \mid 0.11]$; for pond communities: $0.11 \pm [0.09 \mid 0.15]$; Mann-Whitney U-test: $W = 178$, $p = 0.023$).

We further examined whether there is a phylogenetic signal for pyoverdine production (i.e. whether closely related isolates show similar pyoverdine production) (Fig. S1). In line with our previous findings, based on a smaller dataset (Butaitė *et al.*, 2017), we found a very weak phylogenetic signal for soil communities (Blomberg's $K = 0.02 \pm [0.01 \mid 0.12]$), and a low but slightly increased phylogenetic signal in pond communities ($K = 0.12 \pm [0.03 \mid 0.25]$; test for differences between soil and pond communities, Wilcoxon rank sum test: $W = 186$, $p = 0.036$). These analyses confirmed that pyoverdine production levels can vary substantially even among closely related strains.

Environmental determinants of pyoverdine production

Next, we tested for a relationship between the pyoverdine production profiles described above and different abiotic (pH, total iron and carbon concentrations) and biotic (phylogenetic community diversity) variables of the environments the isolates originated from. While these environmental variables varied considerably between communities (Fig. S3, Table S2), they were often correlated within communities (Table S3). To account for the resulting collinearities, we first carried out separate principal component analyses (PCAs) for soil and pond that each incorporated the four environmental variables. PCAs break collinearities by converting the values of potentially correlated variables into a set of uncorrelated variables called principal components (PCs). These PCs can then be analysed using standard linear models. Here, we focused on the first two PCs (henceforth called SPC1 and SPC2 for soil, and PPC1 and PPC2 for pond), which explained 88.0% (soil) and 88.3% (pond) of the total variance observed (Table 1). Each PC is characterized by a set of positive and/or negative loadings, which describe how much it is influenced by each of the environmental variables fed into the PCA.

For soil communities, SPC1 was positively loaded by carbon concentration, and negatively by pH and iron concentration, indicating that SPC1 reflects a trade-off (i.e. a negative correlation) between carbon concentration on the one hand, and pH and iron concentration on the other hand. By contrast, SPC2 was solely and positively loaded by community diversity (Table 1A). When relating these PCs to the pyoverdine production profiles of the soil isolates, we found a positive correlation between relative pyoverdine production and community diversity (SPC2; $t_{18.3}$ = 3.36, p = 0.0034; Fig. 2A). In contrast, there was no association between the relative pyoverdine production and the trade-off between carbon versus pH and iron captured by SPC1 ($t_{10.6}$ = -1.93, p = 0.0809). Because there were many pyoverdine non-producers, present in 22 out of 24 soil communities, we further examined whether the likelihood of being a pyoverdine non-producer correlated with the SPCs. We found that the likelihood of being a non-producer was highest in communities characterized by relatively high levels of carbon combined with relatively low levels of pH and iron (SPC1; z = 2.85, p = 0.0044; Fig. 2B). Conversely, the likelihood of being a non-producer did not correlate with community diversity (SPC2; z = -0.72, p = 0.4745).

For pond communities, PPC1 was positively loaded by iron and carbon, and negatively by pH and community diversity, indicating that PPC1 reflects a trade-off between these groups of environmental variables (Table 1B). Conversely, PPC2 was positively loaded by pH and carbon. Note that this positive association is independent of and occurs simultaneously with the trade-off between these variables in PPC1 (Table 1B). When feeding these PCs into a linear mixed model we found a negative association between relative pyoverdine production and PPC2 ($t_{10.9}$ = -2.92, p = 0.0140), indicating that the relative pyoverdine production was lower among pond isolates from communities characterized by higher levels of carbon and pH (Fig. 2C). In contrast, PPC1 was not linked to the pyoverdine production of pond isolates

($t_{10.0} = 0.68$, $p = 0.5120$). We further examined whether the likelihood of being a pyoverdine

182   non-producer correlated with one of the PPCs, but this was neither the case for PPC1 ($z = -1.16$, $p = 0.2450$) nor for PPC2 ($z = 0.79$, $p = 0.4320$).

The genetic basis of pyoverdine non-production

The occurrence of many pyoverdine non-producers was unexpected, and prompted us to explore their genetic makeup in more detail. Our previous study (Butaitė *et al.*, 2017),

entailing whole-genome sequencing of environmental isolates, revealed that two different types of non-producers exist: those with a highly truncated pyoverdine locus, and those with a

seemingly intact, yet largely silent locus. To find out whether these types also occur here, and whether the frequency of the two types correlates with environmental variables, we screened

all 130 non-producers for the presence of the *pvdL* gene. This gene encodes an essential and conserved non-ribosomal peptide synthetase involved in pyoverdine synthesis (Ravel and

Cornelis, 2003; Smith *et al.*, 2005). The presence or absence of the *pvdL* gene hence indicates whether non-producers possess a silent yet intact locus or a truncated dysfunctional locus,

respectively.

We found that 32 out of 91 (35.2%) soil non-producers and 15 out of 39 (38.5%) pond non-producers were positive for the presence of *pvdL* (no significant difference between soil and

pond: $\chi^2 = 2.48$, $p = 0.1151$). For soil communities, the likelihood of being *pvdL*-positive was lowest in communities characterized by a combination of relatively high carbon

concentration, and a low pH and iron concentration (SPC1: $z = -2.07$, $p = 0.0386$; Fig. 2D). For pond communities, meanwhile, none of the two PPCs were associated with the likelihood

of being *pvdL*-positive (PPC1: $z = -1.54$, $p = 0.1230$; PPC2: $z = -0.83$, $p = 0.4090$).

Supernatants from pyoverdine producers affect the growth of non-producers

To estimate the extent to which pyoverdine could be involved in social interactions between strains, we compared the growth of non-producers in iron-limited medium alone versus their growth in medium supplemented with pyoverdine-containing supernatants of producers from the same community. Overall, we fed 53 non-producers (from 12 soil and pond communities each) with pyoverdine-containing supernatants from four to six different producers from the same community. This resulted in a total of 152 soil and 151 pond non-producer-supernatant combinations.

We found that the fitness effects of pyoverdine-containing supernatants on non-producers covered the entire range from almost complete growth inhibition to high stimulation (Fig. S4). While the likelihood of stimulation did not significantly differ between the two habitats (soil vs. pond: $z = 1.80$, $p = 0.072$; Fig. 3A), the likelihood of inhibition was significantly higher in pond than in soil communities ($z = 2.86$, $p = 0.004$; Fig. 3B). Note that 'stimulation' or 'inhibition' describe the respective cases where non-producers grew significantly better or worse, with pyoverdine-containing supernatants than without. Although supernatants contain other growth-modulating components in addition to pyoverdine, we have previously shown that the growth effects of this supernatant assay on non-producers are mainly caused by pyoverdine, and likely driven by the presence (stimulation) and absence (inhibition) of a matching receptor for pyoverdine uptake (Butaitė *et al.*, 2017).

Environmental determinants of pyoverdine-mediated social interactions

Next, we tested whether the likelihood of stimulation and inhibition correlate with environmental factors. To this end, we reused the first two PCs describing the relationships between our four environmental variables of interest (pH, iron and carbon concentrations,

community diversity) as explanatory variables. Because the supernatant assay involved pairs

232   of strains (i.e. the supernatant donor and the recipient), we further included the phylogenetic

relatedness between isolates, based on *rpoD* sequence similarity, into our statistical models.

For soil communities, we found that the likelihood of stimulation correlated with SPC1,

236   suggesting that non-producers were most likely to be stimulated by heterologous pyoverdine

when originating from environments featuring a combination of a high carbon concentration

238   on the one hand, and a low pH and iron concentration on the other hand ($z = 2.21$, $p = 0.0269$;

Fig. 4A). Furthermore, the likelihood of stimulation tended to increase with community

240   diversity (SPC2; $z = 1.96$, $p = 0.0502$), and increased with the relatedness between strains

(*rpoD* identity; $z = 3.44$, $p = 0.0006$; Fig. 4B). The opposite trends were generally observed

242   for the likelihood of inhibition: it correlated negatively with SPC1 ($z = -1.94$, $p = 0.0525$), and

relatedness (*rpoD* identity; $z = -2.80$, $p = 0.0052$), but not with community diversity (SPC2; z

244    $= -1.01$, $p = 0.3144$).

246   For pond communities, we observed that the likelihood of stimulation negatively correlated

with PPC1, suggesting that non-producers were most likely to be stimulated by heterologous

248   pyoverdine when originating from environments featuring low levels of carbon and iron

together with high pH and community diversity ($z = -2.13$, $p = 0.0333$; Fig. 4C). Moreover,

250   the likelihood of stimulation depended on an interaction between PPC2 and the phylogenetic

relatedness between the supernatant donor and the recipient (interaction: $z = 2.72$, $p = 0.0066$;

252   main effects: *rpoD* identity; $z = 1.31$, $p = 0.1900$; and PPC2; $z = 0.65$, $p = 0.5132$; Fig. 4D). In

particular, a relatively high phylogenetic relatedness increased the likelihood of stimulation in

254   communities featuring high levels of PPC2 (i.e. simultaneously high levels of pH and carbon),

whereas this relationship was reversed in communities featuring low levels of PPC2 (Fig.

4D). By contrast, neither the PPCs nor *rpoD* identity correlated with the likelihood of inhibition (PPC1: $z = 1.82$, $p = 0.0688$; PPC2: $z = -1.30$, $p = 0.1943$; *rpoD* identity: $z = -1.37$, $p = 0.1701$).

Discussion

The beauty of laboratory experiments in microbiology is that factors influencing bacterial physiology, behaviour, and fitness can be investigated one at the time under controlled and replicable conditions. This approach contrasts with the situation bacteria typically face in nature, where environmental conditions often fluctuate rapidly in unpredictable manners, with multiple variables simultaneously influencing bacterial behaviour and fitness. This raises the question whether the factors affecting bacterial behaviour in vitro also play a role under natural conditions. Our study tackled this question by examining whether factors shown to influence an important bacterial trait in the laboratory, the production of siderophores used for iron-scavenging, also affect this behaviour in natural communities. As a model system, we focused on the siderophore pyoverdine produced by *Pseudomonas* bacteria. Our investigations involving 930 *Pseudomonas* isolates, originating from 24 soil and 24 pond communities, yielded several novel insights. First, we found that pH, concentrations of total iron and carbon, and community diversity, all shown to be important determinants of pyoverdine production in the laboratory, are indeed correlated with the level of pyoverdine produced by natural *Pseudomonas* isolates. Second, we observed that the same environmental variables also correlated with pyoverdine-mediated social interactions, measured by the extent to which secreted pyoverdine could promote or inhibit the growth of other members of the community. Third, we showed that the way these environmental variables correlated with pyoverdine production and social interactions differed fundamentally between soil and pond communities. Finally, our data suggest that trade-offs and interactions between environmental

factors, which are typically ruled out in the laboratory, could be more predictive of bacterial

behaviour in nature than the main effect of a single factor.

We observed that isolates from soil and pond communities differ substantially in their pyoverdine production profiles and in the way their production levels seem to be influenced

by environmental factors. In particular, we found that: (a) soil isolates produced on average significantly less pyoverdine than pond isolates (Fig. 1A); (b) there were significantly more

pyoverdine non-producers in soil than in pond (Fig. 1A; Table S2); and (c) the four environmental variables examined (community diversity, pH, total iron and carbon

concentrations) were all associated with at least one aspect of pyoverdine production for soil isolates, whereas the pyoverdine production of pond isolates seemed to be much less affected

by these variables (Fig. 2). Observation (a) can relatively easily be explained by the fact that diffusivity is higher and iron concentration orders of magnitude lower in ponds compared to

soils (Fig. S3, Table S2), such that higher siderophore investments are required for successful iron scavenging. Observation (b) was more surprising, as we initially expected the opposite

pattern, if non-producers evolve as cheaters, exploiting pyoverdines produced by others for iron scavenging. This is because the more diffusive nature of ponds combined with low iron

availability should provide ample opportunities for pyoverdine exploitation. The opposite pattern observed here could suggest that there are other forces than cheating that select for

non-production (see detailed discussion in next paragraph). Observation (c), meanwhile, could be the result of ponds being more open systems, where new strains can easily be

introduced and strain mixing is conceivably high due to the more diffusive and less structured environment. Contrary to soils, where local adaptation has been observed (Bruce, West, *et al.*,

2017), increased strain mixing in ponds might even out any local variation in community composition and hamper adaptation to prevailing local environmental conditions.

We propose three different mutually non-exclusive reasons for the evolution of pyoverdine non-producers in natural habitats. First, non-producers could evolve as cheaters exploiting the

pyoverdine produced by others. In pond communities, this is a reasonable scenario because the prevailing conditions of high strain mixing, severe iron limitation and low carbon

availability have been shown to promote cheating in laboratory experiments (Brockhurst *et al.*, 2008; Kümmerli *et al.*, 2009; Sexton and Schuster, 2017). In soil communities, on the

other hand, cheaters could be favoured because this habitat sustains high cell densities, a factor previously shown to promote cheating (Ross-Gillespie *et al.*, 2009; Cordero *et al.*,

2012; Scholz and Greenberg, 2015; Weigert and Kümmerli, 2017). While the complexity of natural habitats makes it difficult to assess the relative importance of these selection

pressures, both habitats clearly have features that could select for cheating. Second, non-producers could evolve under conditions where pyoverdine is not stringently needed for iron

acquisition. The idea that disuse drives pyoverdine loss is supported by our observation that the likelihood of being a non-producer was highest under conditions with highest predicted

iron bioavailability (i.e. in soils with high carbon concentration in combination with low pH and iron concentration; Fig. 2B). Iron bioavailability is predicted to be highest under these

conditions because iron solubility increases dramatically at low pH in soil, with the total iron content playing a minor role (Petruzzelli, 1989; Rieuwerts *et al.*, 1998). Moreover, high

carbon concentrations decrease the cost of pyoverdine production (Brockhurst *et al.*, 2008; Sexton and Schuster, 2017), and can also increase the bioavailability of iron, via the metal-

complexing properties of organic compounds, such as humic and fulvic acids (Lindsay, 1991; Hassler *et al.*, 2009; Krachler *et al.*, 2015). The hypothesis that pyoverdine is lost due to

disuse is further supported by our observation that the frequency of non-producers that have permanently lost the pyoverdine locus (i.e. *pvdL*-negative isolates) peaked exactly under the

conditions of predicted highest iron bioavailability (Fig. 2D). Finally, non-producers could

also evolve in the context of heavy metal detoxification. For instance, low pH in soils is known to increase the solubility of toxic heavy metals (Petruzzelli, 1989). Pyoverdine can bind many of those metals (usually with lower affinity than iron), and thereby contribute to the detoxification of the environment (Braud *et al.*, 2009; Schalk *et al.*, 2011). Since detoxification also constitutes a public good, it is possible that non-producers evolved as cheaters in a context of heavy metal detoxification (O'Brien *et al.*, 2014; Hesse *et al.*, 2018).

We now turn to our findings from the supernatant assay, where we fed pyoverdine-containing supernatants from producers to non-producers from the same community. For soil communities, we found that the likelihood of non-producers to be stimulated by supernatants increased under conditions of higher predicted iron availability (Fig. 4A) and when relatedness among strains was high (Fig. 4B). This could indicate that in relatively iron-replete soils and among closely related strains, the selection pressure for producers to evolve cheating-resistance mechanisms is reduced. Conversely, under stringent iron limitation (i.e. high pH combined with low carbon concentration) the likelihood of pyoverdine-mediated stimulation was lower, and inhibition increased. Under these conditions, selection thus seem to favour exclusive non-exploitable pyoverdines, for which non-producers lack uptake receptors (Smith *et al.*, 2005; Lee *et al.*, 2012). This is conceivable because cheating is thought to be especially harmful for producers when iron is scarce and carbon as a building block for pyoverdine is limited (Brockhurst *et al.*, 2008; Kümmerli *et al.*, 2009; Sexton and Schuster, 2017). Taken together, the patterns discovered here suggest that the role of pyoverdine changes with increasing iron availability: from a shareable, yet exploitable public good, to an inhibitory substance, locking iron away from competitors lacking a matching uptake receptor (Buyer and Leong, 1986; Butaitė *et al.*, 2017; Niehus *et al.*, 2017; Schiessl *et al.*, 2017; Sexton *et al.*, 2017).

For pond communities, we initially proposed that the openness of aquatic systems, characterized by increased strain influx from surrounding habitats and increased strain

mixing, reduce the potential for local strain adaptation to abiotic factors. These environmental properties are likely to also affect pyoverdine-mediated social interaction patterns. For

instance, high levels of mixing could promote random interactions between pyoverdine producers and non-producers. This in turn could reduce the likelihood of non-producers to

possess a matching receptor for the uptake of a specific pyoverdine, and thus lead to the high levels of pyoverdine-mediated growth inhibition observed in our pond communities (Fig. 3B).

Nonetheless, we also found evidence for non-random interaction patterns, as the likelihood of non-producers to be stimulated by pyoverdine-containing supernatant from producers was

highest in ponds with low predicted iron availability (high pH combined with low carbon and iron concentrations, but also high community diversity; Fig. 4C). This pattern opposes the one

observed for soil non-producers (Fig. 4A), suggesting that selection for cheating and cheating resistance differ between the two habitat types. This notion is also supported by the

occurrence of complex interactions between inter-strain relatedness and the environmental variables in ponds (Fig. 4D), but not in soils.

In conclusion, our study reveals that connecting laboratory to field studies is a challenging

374   task for microbiologists. While laboratory studies are typically carried out with isogenic strains and engineered single-gene knockout mutants grown under controlled conditions, field

376   studies, like ours, face an enormous strain diversity, and a plethora of environmental factors that all simultaneously vary and influence bacterial behaviour. Despite this complexity, we

378   managed to obtain first insights into the importance of siderophore-based iron acquisition and competition in natural communities of pseudomonads. In particular, we found evidence that

380   pyoverdine production levels of natural isolates and the ability to use heterologous pyoverdine

are simultaneously influenced by multiple environmental variables, including pH, iron and

382 carbon concentrations, with patterns clearly differing between soil and pond habitats. Overall,

we advocate the need for further studies that embrace the complexity of natural systems, in

384 order to unravel how microbial behaviour and strain interactions shape community

composition and functioning.

Experimental Procedures

388 **Sampling and isolation of pseudomonads.** We sampled eight *Pseudomonas* communities at

each of three pond and three soil locations (48 communities in total). While one pond and one

390 soil location were situated on the Irchel campus of the University of Zurich (47.40° N, 8.54°

E), the other two soil and two pond habitats were located at Seleger Moor Park (47.25° N,

392 8.51° E), Switzerland. The a priori reasoning for sampling at these different locations was that

we expected ecological parameters to differ quite substantially between the habitats in a city

394 park and a more natural environment. We sampled soil cores and pond water, isolated 952

strains on a medium selective for fluorescent pseudomonads, and preserved the isolates as

396 freezer stocks as described in detail elsewhere (Butaitė *et al.*, 2017). We provided each isolate

with a unique identification code, consisting of a location ID ('s1 = Seleger Moor soil#1', 's2

398 = Seleger Moor soil#2', 's3 = Irchel soil', '1 = Seleger Moor pond#1', '2 = Seleger Moor

pond#2', '3 = Irchel pond'), a community ID (small letters 'a' to 'h for soil communities;

400 capital letters 'A' to 'H' for pond communities) and an isolate number (1 to 20).

402 **Measuring environmental parameters.** We measured the pH of soil and pond samples using

the Atago portable pH meter DPH-2 (for field measurements) and the Metrohm 744 pH meter

404 (for laboratory measurements). The pH of pond samples was first measured in the laboratory

and then later confirmed by measurements directly in the field. Prior to soil pH

406 measurements, the soil samples were suspended 1:5 (w/v) in 0.01 M $CaCl_2$ solution, shaken for 1 h, and left to sediment.

We measured the percentage of total carbon in soil and pond samples with a CHN (Leco
Truspec micro; Leco Instruments, USA) and a TOC analyser (Dimatoc 2000; Dimatec, Germany), respectively. Measurements were carried out by staff operating the specific in-
house service at the University of Zurich. Prior to the measurements, we first dried soil samples for two days at 40 °C, then grounded them to a homogenous powder and dried them
again at 40 °C overnight.

To estimate the iron concentration of the sampled soils and ponds, we used the ICP-OES methodology (Vista-MPX, Varian, USA). For soils, we first dried and homogenized the plant-
free samples (plant material was removed using a 0.6 mm stainless steel sieve). Soil samples were then digested with HCl and HNO_3 for 90 min at 120 °C, diluted in Nanopure water and
filtered. The pond samples were directly filtered using a 0.2 µm filter and then acidified with 65% HNO_3 to pH < 3. To make iron concentrations comparable between soil (µg/g) and pond
(µg/L) samples, we converted pond concentrations to µg/g by assuming that 1 ml = 1 g.

rpoD amplification and sequencing. To verify that the 952 isolates are indeed *Pseudomonas*, we PCR amplified and sequenced a part of the *rpoD* gene for 930 isolates (PCR or sequencing
failed for 22 isolates, which were excluded from further experiments). This housekeeping gene is commonly used for phylogenetic affiliation of pseudomonads (Mulet *et al.*, 2009;
Ghyselinck *et al.*, 2013). PCR mixtures were prepared, PCR reactions were carried out and the products were sequenced using PsEG30F and PsEG790R primers (Mulet *et al.*, 2009) as
described elsewhere (Butaitė *et al.*, 2017).

Community diversity and phylogenetic relatedness. A codon-aware nucleotide alignment of *rpoD* was generated using local TranslatorX v1.1 (Abascal *et al.*, 2010) with the MAFFT v7.271 (Katoh and Standley, 2013) aligner. We manually curated and trimmed the alignment at both ends resulting in a high-quality alignment of 908 sequences (including 21 reference strains; as described elsewhere (Butaitė *et al.*, 2017) over 609 nucleotides. The phylogenetic tree was inferred by RAxML v7.0.4 (Stamatakis, 2006) using the General Time Reversible (GTR) + G model with 100 bootstraps. We calculated phylogenetic diversity within communities based on the cophenetic function from the ape package in R software and normalized values by the number of tips. For some analyses, we calculated the relatedness between strains by carrying out pairwise alignments of *rpoD* sequences using the 'water' application from EMBOSS (Rice *et al.*, 2000).

pvdL amplification. We used the presence of the *pvdL* gene as a proxy for the presence of the pyoverdine biosynthesis locus in all 130 pyoverdine non-producers. *pvdL* is involved in the biosynthesis of the pyoverdine chromophore, and is the most conserved part of the locus across different pseudomonads (Ravel and Cornelis, 2003). To identify a conserved region of this gene, we aligned the *pvdL* sequences (retrieved from GenBank) of seven pseudomonads: *P. aeruginosa* PAO1, *P. chlororaphis* O6, *P. chlororaphis subsp.aureofaciens* 30-84, *P. fluorescens* A506, *P. fluorescens* SS101, *P. protegens* Pf-5, *P. syringae pv.syringae* B64. We then designed the following primers to amplify a conserved region in the isolated non-producers (pvdL_fw: CATGATGAGCAACCACCACATC, pvdL_rv: CGCTGGTCGTAGGACAGGTG; product size: 827 bp). We prepared PCR mixtures as for the *rpoD* gene amplification (Butaitė *et al.*, 2017), just this time we used a 'Hot Start' version of the Master Mix. Bacterial biomass was taken from fresh cultures grown in liquid LB

456   medium (1 µl). We used the following PCR conditions: denaturation at 94.5 °C for 5 min; 30

cycles of amplification (1 min denaturation at 94 °C, 1 min primer annealing at 57 °C, and 1

458   min primer extension at 68 °C); final elongation at 68 °C for 10 min. As a positive control, we

used 72 pyoverdine producers (pond and soil, 36 each). We considered an isolate as *pvdL*-

460   positive when the PCR yielded a product of the expected size. Among the 72 pyoverdine

producers (which should all have *pvdL*), we found seven to be negative for *pvdL*. We thus

462   estimate the rate of false-negatives to be 9.7% (i.e. the risk to falsely identify isolates as *pvdL*-

negatives).

Measurement of growth and pyoverdine production levels. To evaluate whether and to

466   what extent the isolates can produce pyoverdine, we grew them under iron-limited conditions

and measured their pyoverdine production levels. We first grew the isolates in 150 µl LB in

468   96-well plates overnight (16 - 18 h) static at room temperature. We then transferred 2 µl of

overnight cultures to 200 µl iron-limited casamino acids medium (CAA containing 5 g

470   casamino acids, 1.18 g $K_2HPO_4 \cdot 3H_2O$, 0.25 g $MgSO_4 \cdot 7H_2O$ per liter) supplemented with 25

mM HEPES buffer, 20 mM $NaHCO_3$ and 100 µg/ml of the strong iron chelator human apo-

472   transferrin in a 96-well plate. All chemicals were purchased from Sigma-Aldrich,

Switzerland.

Each plate had isolates from one community in triplicates and eight reference strains known

476   to produce pyoverdine (see Table S1 for a description of these strains). After 18 h of

incubation at room temperature, we measured growth (optical density OD at 600 nm) and

478   pyoverdine production levels (relative fluorescence units, RFU, with excitation: 400 nm and

emission: 460 nm) with an Infinite M200 Pro microplate reader (Tecan Group Ltd.,

480   Switzerland). We then scaled the growth and relative pyoverdine production values for each

isolate by dividing their OD_{600} and RFU by the average respective OD_{600} and RFU of the

482   reference strains.

484   We also carried out a control experiment to verify that it is indeed iron limitation that induces

the observed growth and pyoverdine production patterns, and not a specific effect of

486   transferrin as an iron chelator. For this, we repeated the growth assay with the synthetic iron

chelator 2,2'-dipyridyl (200 µM). The growth assay was carried out for a subset of soil (n =

488   100: 28 pyoverdine non-producers and 72 producers) and pond isolates (n = 99: 27 non-

producers and 72 producers), from 12 soil and 12 pond communities, in three replicates. The

490   assay yielded qualitative very similar results for both iron chelators (Fig. S5). Moreover, in an

earlier study we could demonstrate that the environmental isolates ($n = 315$) could use CAA

492   as a carbon source, demonstrating that the poor growth of non-producers is due to their

inability to produce pyoverdine, and not because they are unable to grow in CAA (Butaitė *et*

494   *al.*, 2017).

496   **Supernatant assay.** To quantify the effect that pyoverdine-containing supernatants from

producers has on non-producers, we set up a supernatant growth assay using a subset of

498   isolates and communities (24 communities in total: four communities per location and habitat

type). For each community, we first selected six pyoverdine producers. To be considered for

500   the supernatant assays, producers had to: (a) grow better than the corresponding non-

producers under iron-limited condition; and (b) differ in their *rpoD* sequence, and thus

502   represent phylogenetically different strains. These criteria were met for 22 communities. For

the two remaining communities (3E and s2e), we picked two producers with identical *rpoD*

504   sequence, whereby for s2e we were limited to four producers satisfying (a). For each

community, we further randomly selected up to three non-producers (upon availability). In

total, we had 26 and 27 non-producers (relative pyoverdine production < 5 % relative to the eight reference pyoverdine producers listed in Table S1) from soil and pond communities, respectively. This resulted in 152 soil non-producer-supernatant and 151 pond non-producer-supernatant combinations.

To generate pyoverdine-containing supernatants, we first grew the producers (n = 142) in 200 µl LB medium in 96-well plates overnight at 25 °C. Then, 20 µl of overnight cultures were added to 2 ml of CAA with 200 µM 2,2'-dipyridyl in 24-well plates and incubated static at 25 °C for about 22 h. We centrifuged cultures for 10 min at 3,500 rpm (Eppendorf Centrifuge 5804R) and then transferred 900 µl of supernatants to PALL AcroPrep Advance 96-well 1 ml filterplates (with 0.2 µm supor membrane), attached to an autoclaved 1.2 ml 96-well PCR plate (VWR). We centrifuged the assemblies of filter and collection plates at 2,500 rpm for 15 min. The collection plates with sterile supernatants and blank medium were sealed with Greiner SILVERseal and kept at -20 °C.

Afterwards we grew the non-producers in 200 µl of LB in 96-well plates overnight static at 25 °C. We then added 2 µl of non-producer cultures to CAA with 200 µM 2,2'-dipyridyl supplemented with 20 µl of (a) producer supernatant or (b) CAA with 200 µM 2,2'-dipyridyl that went through the same treatment as supernatants, i.e. filtering and freezing. Total culturing volume was 200 µl. Each treatment was set up in four replicates. Plates were incubated static for 17 h at 25 °C. The final OD_{600} of the cultures was measured using the Tecan microplate reader. We considered a supernatant effect as 'stimulation' or 'inhibition' when non-producers grew significantly better or worse, respectively, with a supernatant than without, based on a Wilcoxon test.

Statistical analysis. We used linear mixed-effects (LMM) and linear generalized mixed-effects (GLMM) models for statistical data analysis. Since strains isolated from the same and close communities (50 cm apart; we called neighbouring communities a 'core') might not be independent from one another, we included core and community identities as random factors into our models. Whenever appropriate, we used natural log-transformations to meet the assumption of normally distributed residuals. All statistical analyses were carried out using the R 3.1.2 statistics software (www.r-project.org).

The environmental variables were often highly correlated (Table S3), which led to high collinearities in statistical models (variance inflation factor (vif) $\gg$ 10). We thus conducted principal component analyses (PCAs) to obtain non-correlated principal components (PCs) reflecting single or combinations of different environmental variables (entered after centering and scaling to unit variance, respectively). We performed two separate PCAs for soil and pond communities, because the overall differences in pH and the total concentrations of iron and carbon (Table S2), together with differences in the signs of some of the correlations among environmental variables in the soil and pond (Table S3), also lead to high collinearity (variance inflation factor $\gg$ 10) in models deploying PCs based on the whole dataset. We thus used the PCs resulting from separate PCAs for soil and pond (Table 1) to analyse their effects on pyoverdine production and social interactions.

Acknowledgments. We dedicate this article to Prof. Helmut Brandl, who inspired and advised us on this project, but sadly passed away in 2017. We thank Roland Dünner for permission to sample at Seleger Moor park, René Husi for carbon concentration measurements, and Prof. Rainer Schulin and Björn Studer from ETH Zurich for iron concentration measurements at their facility. This work was funded by the Swiss National

556 Science Foundation (grants no. 139164 and 165835 to RK) and the Forschungskredit of the University of Zurich (FK-15-082 to EB and FK-17-111 to JK). The authors declare no

558 conflict of interest.

560 **Author contribution.** EB and RK designed the study, EB isolated the strains and performed the experiments, EB, JK, SW and RK analysed the data and wrote the paper.

References

564   Abascal, F., Zardoya, R., and Telford, M.J. (2010) TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations. *Nucleic Acids Res.* **38**: W7–

566   W13.

Andrews, S.C., Robinson, A.K., and Rodríguez-Quiñones, F. (2003) Bacterial iron

568   homeostasis. *FEMS Microbiol. Rev.* **27**: 215–237.

Boyd, P.W. and Ellwood, M.J. (2010) The biogeochemical cycle of iron in the ocean. *Nat.*

570   *Geosci.* **3**: 675–682.

Braud, A., Hoegy, F., Jezequel, K., Lebeau, T., and Schalk, I.J. (2009) New insights into the

572   metal specificity of the *Pseudomonas aeruginosa* pyoverdine-iron uptake pathway. *Environ. Microbiol.* **11**: 1079–1091.

574   Brockhurst, M.A., Buckling, A., Racey, D., and Gardner, A. (2008) Resource supply and the evolution of public-goods cooperation in bacteria. *BMC Biol.* **6**: 20.

576   Bruce, J.B., Cooper, G.A., Chabas, H., West, S.A., and Griffin, A.S. (2017) Cheating and resistance to cheating in natural populations of the bacterium *Pseudomonas fluorescens*.

578   *Evolution*.

Bruce, J.B., West, S.A., and Griffin, A.S. (2017) Bacteriocins and the assembly of natural

580   *Pseudomonas fluorescens* populations. *J. Evol. Biol.* **30**: 352–360.

Butaitė, E., Baumgartner, M., Wyder, S., and Kümmerli, R. (2017) Siderophore cheating and

582   cheating resistance shape competition for iron in soil and freshwater *Pseudomonas* communities. *Nat. Commun.* **8**: 414.

584   Buyer, J. and Leong, J. (1986) Iron transport-mediated antagonism between plant growth-promoting and plant-deleterious *Pseudomonas* strains. *J. Biol. Chem.* **261**: 791–794.

586   Cézard, C., Farvacques, N., and Sonnet, P. (2015) Chemistry and biology of pyoverdines, *Pseudomonas* primary siderophores. *Curr. Med. Chem.* **22**: 165–186.

588   Cordero, O.X., Ventouras, L.-A., DeLong, E.F., and Polz, M.F. (2012) Public good dynamics drive evolution of iron acquisition strategies in natural bacterioplankton populations.

590   *Proc. Natl. Acad. Sci. U. S. A.* **109**: 20059–64.

Cornelis, P. (2010) Iron uptake and metabolism in pseudomonads. *Appl. Microbiol.*

592   *Biotechnol.* **86**: 1637–1645.

D'Onofrio, A., Crawford, J.M., Stewart, E.J., Witt, K., Gavrish, E., Epstein, S., et al. (2010)

594   Siderophores from neighboring organisms promote the growth of uncultured bacteria. *Chem. Biol.* **17**: 254–264.

596 Dumas, Z., Ross-Gillespie, A., and Kümmerli, R. (2013) Switching between apparently
redundant iron-uptake mechanisms benefits bacteria in changeable environments. *Proc.*
598 *Biol. Sci.* **280**: 20131055.

Frawley, E.R. and Fang, F.C. (2014) The ins and outs of bacterial iron metabolism. *Mol.*
600 *Microbiol.* **93**: 609–616.

Ghyselinck, J., Coorevits, A., Van Landschoot, A., Samyn, E., Heylen, K., and De Vos, P.
602 (2013) An *rpoD* gene sequence based evaluation of cultured *Pseudomonas* diversity on
different growth media. *Microbiology* **159**: 2097–108.

604 Griffin, A.S., West, S.A., and Buckling, A. (2004) Cooperation and competition in pathogenic
bacteria. *Nature* **430**: 1024–1027.

606 Harrison, F., Paul, J., Massey, R.C., and Buckling, A. (2008) Interspecific competition and
siderophore-mediated cooperation in *Pseudomonas aeruginosa*. *ISME J.* **2**: 49–55.

608 Hassler, C.S., Havens, S.M., Bullerjahn, G.S., McKay, R.M.L., and Twissa, M.R. (2009) An
evaluation of iron bioavailability and speciation in western Lake Superior with the use of
610 combined physical, chemical, and biological assessment. *Limnol. Oceanogr.* **54**: 987–
1001.

612 Hesse, E., O'Brien, S., Tromas, N., Bayer, F., Luján, A.M., van Veen, E.M., et al. (2018)
Ecological selection of siderophore-producing microbial taxa in response to heavy metal
614 contamination. *Ecol. Lett.* **21**: 117–127.

Hider, R.C. and Kong, X. (2010) Chemistry and biology of siderophores. *Nat. Prod. Rep.* **27**:
616 637.

Inglis, R.F., Biernaskie, J.M., Gardner, A., and Kümmerli, R. (2016) Presence of a loner
618 strain maintains cooperation and diversity in well-mixed bacterial communities. *Proc. R.*
Soc. London B Biol. Sci. **283**:.

620 Jiricny, N., Diggle, S.P., West, S.A., Evans, B.A., Ballantyne, G., Ross-Gillespie, A., and
Griffin, A.S. (2010) Fitness correlates with the extent of cheating in a bacterium. *J. Evol.*
622 *Biol.* **23**: 738–747.

Joshi, F., Archana, G., and Desai, A. (2006) Siderophore cross-utilization amongst
624 rhizospheric bacteria and the role of their differential affinities for Fe3+ on growth
stimulation under iron-limited conditions. *Curr. Microbiol.* **53**: 141–147.

626 Katoh, K. and Standley, D.M. (2013) MAFFT multiple sequence alignment software version
7: improvements in performance and usability. *Mol. Biol. Evol.* **30**: 772–80.

628 Krachler, R., Krachler, R.F., Wallner, G., Hann, S., Laux, M., Cervantes Recalde, M.F., et al.
(2015) River-derived humic substances as iron chelators in seawater. *Mar. Chem.* **174**:

630   85–93.

Kümmerli, R., Jiricny, N., Clarke, L.S., West, S.A., and Griffin, A.S. (2009) Phenotypic
632   plasticity of a cooperative behaviour in bacteria. *J. Evol. Biol.* **22**: 589–598.

Lee, W., van Baalen, M., and Jansen, V.A.A. (2012) An evolutionary mechanism for diversity
634   in siderophore-producing bacteria. *Ecol. Lett.* **15**: 119–125.

Leinweber, A., Inglis, R.F., and Kümmerli, R. (2017) Cheating fosters species co-existence in
636   well-mixed bacterial communities. *ISME J.* **11**: 1179–1188.

Lindsay, W.L. (1991) Iron oxide solubilization by organic matter and its effect on iron
638   availability. *Plant Soil* **130**: 27–34.

Loper, J.E. and Henkels, M.D. (1997) Availability of iron to *Pseudomonas fluorescens* in
640   rhizosphere and bulk soil evaluated with an ice nucleation reporter gene. *Appl. Environ.*
      Microbiol. **63**: 99–105.

Loper, J.E. and Henkels, M.D. (1999) Utilization of heterologous siderophores enhances
642   levels of iron available to *Pseudomonas putida* in the rhizosphere. *Appl. Environ.*
644   *Microbiol.* **65**: 5357–63.

Meyer, J.-M., Gruffaz, C., Raharinosy, V., Bezverbnaya, I., Schäfer, M., and Budzikiewicz,
646   H. (2008) Siderotyping of fluorescent *Pseudomonas*: molecular mass determination by
      mass spectrometry as a powerful pyoverdine siderotyping method. *BioMetals* **21**: 259–
648   271.

Miethke, M. and Marahiel, M. a (2007) Siderophore-based iron acquisition and pathogen
650   control. *Microbiol. Mol. Biol. Rev.* **71**: 413–451.

Mulet, M., Bennasar, A., Lalucat, J., and García-Valdés, E. (2009) An *rpoD*-based PCR
652   procedure for the identification of *Pseudomonas* species and for their detection in
      environmental samples. *Mol. Cell. Probes* **23**: 140–7.

654 Neilands, J.B. (1981) Iron absorption and transport in microorganisms. *Annu. Rev. Nutr.* **1**:
      27–46.

656 Niehus, R., Picot, A., Oliveira, N.M., Mitri, S., and Foster, K.R. (2017) The evolution of
      siderophore production as a competitive trait. *Evolution* **71**: 1443–1455.

658 O'Brien, S., Hodgson, D.J., and Buckling, A. (2014) Social evolution of toxic metal
      bioremediation in *Pseudomonas aeruginosa*. *Proc. R. Soc. B Biol. Sci.* **281**: 20140858.

660 Parrow, N.L., Fleming, R.E., and Minnick, M.F. (2013) Sequestration and scavenging of iron
      in infection. *Infect. Immun.* **81**: 3503–14.

662 Petruzzelli, G. (1989) Recycling wastes in agriculture: heavy metal bioavailability. *Agric.*
      Ecosyst. Environ. **27**: 493–503.

664 Ravel, J. and Cornelis, P. (2003) Genomics of pyoverdine-mediated iron uptake in pseudomonads. *Trends Microbiol.* **11**: 195–200.

666 Rice, P., Longden, I., and Bleasby, A. (2000) EMBOSS: the European Molecular Biology Open Software Suite. *Trends Genet.* **16**: 276–7.

668 Rieuwerts, J.S., Thornton, I., Farago, M.E., and Ashmore, M.R. (1998) Factors influencing metal bioavailability in soils: preliminary investigations for the development of a critical 670 loads approach for metals. *Chem. Speciat. Bioavailab.* **10**: 61–75.

Ross-Gillespie, A., Gardner, A., Buckling, A., West, S.A., and Griffin, A.S. (2009) Density 672 dependence and cooperation: Theory and a test with bacteria. *Evolution* **63**: 2315–2325.

Schalk, I.J., Hannauer, M., and Braud, A. (2011) New roles for bacterial siderophores in 674 metal transport and tolerance. *Environ. Microbiol.* **13**: 2844–2854.

Schiessl, K.T., Janssen, E.M.-L., Kraemer, S.M., McNeill, K., and Ackermann, M. (2017) 676 Magnitude and mechanism of siderophore-mediated competition at low iron solubility in the *Pseudomonas aeruginosa* pyochelin system. *Front. Microbiol.* **8**: 1964.

678 Scholz, R.L. and Greenberg, E.P. (2015) Sociality in *Escherichia coli*: enterochelin is a private good at low cell density and can be shared at high cell density. *J. Bacteriol.* **197**: 680 2122–8.

Sexton, D.J., Glover, R.C., Loper, J.E., and Schuster, M. (2017) *Pseudomonas protegens* Pf-5 682 favors self-produced siderophore over free-loading in interspecies competition for iron. *Environ. Microbiol.* **19**: 3514–3525.

684 Sexton, D.J. and Schuster, M. (2017) Nutrient limitation determines the fitness of cheaters in bacterial siderophore cooperation. *Nat. Commun.* **8**: 230.

686 Smith, E.E., Sims, E.H., Spencer, D.H., Kaul, R., and Olson, M. V. (2005) Evidence for diversifying selection at the pyoverdine locus of *Pseudomonas aeruginosa*. *J. Bacteriol.* 688 **187**: 2138–2147.

Stamatakis, A. (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses 690 with thousands of taxa and mixed models. *Bioinformatics* **22**: 2688–2690.

Tiburzi, F., Imperi, F., and Visca, P. (2008) Intracellular levels and activity of PvdS, the 692 major iron starvation sigma factor of *Pseudomonas aeruginosa*. *Mol. Microbiol.* **67**: 213–227.

694 Traxler, M.F., Seyedsayamdost, M.R., Clardy, J., and Kolter, R. (2012) Interspecies modulation of bacterial development through iron competition and siderophore piracy. 696 *Mol. Microbiol.* **86**: 628–44.

Weigert, M. and Kümmerli, R. (2017) The physical boundaries of public goods cooperation

698   between surface-attached bacterial cells. *Proceedings. Biol. Sci.* **284**:.

West, S.A., Griffin, A.S., Gardner, A., and Diggle, S.P. (2006) Social evolution theory for
700   microorganisms. *Nat. Rev. Microbiol.* **4**: 597–607.

Figure 1. Environmental pseudomonads vary greatly in their level of pyoverdine production and ability to grow under iron-limited conditions. (A) Our phenotypic screen involving 930 natural pseudomonads from 24 soil and 24 pond communities (n = 462 soil and n = 468 pond) revealed that soil isolates produce significantly less pyoverdine than pond isolates, and that there are significantly more non-producers in soil than in pond communities. **(B)** There is a positive correlation between the growth (optical density measured at 600 nm) of isolates and their pyoverdine production level, indicating that pyoverdine is important to cope with iron limitation. Pyoverdine production and growth were measured in iron-limited CAA medium, and scaled relative to the values of eight laboratory reference strains (Table S1). Values represent means across three replicates for soil (yellow triangles) and pond (blue circles) isolates. Dashed lines indicate pyoverdine production (horizontal line in A, vertical line in B) and growth (horizontal line in B) of the reference strains. The box plot shows the median (bold line), the 1[st] and 3[rd] quartile (box), and the 5[th] and 95[th] percentile (whiskers).

Figure 2. Correlations between parameters of pyoverdine production and environmental variables. In soil communities (*n* = 462 isolates), relative pyoverdine production in iron-limited medium increased with community diversity (SPC2) **(A)**, whereas the likelihood of being a pyoverdine non-producer correlated with SPC1, i.e. increased with higher concentration of carbon in combination with lower pH and reduced iron concentration **(B)**. For pond communities (*n* = 468 isolates), relative pyoverdine

production correlated negatively with PPC2, i.e. decreased with higher concentrations of of carbon in combination with increased pH **(C)**. A PCR amplification of the gene *pvdL* in the 91 non-producers from soil revealed a negative correlation between the likelihood of having *pvdL* and SPC1 **(D)**. Since *pvdL* serves as an indicator for the presence of the pyoverdine biosynthesis locus, this analysis indicates that the likelihood of still having *pvdL*, although being a non-producer, increases in habitats characterized by high pH and iron in combination with low carbon. Solid lines with shaded areas show significant correlations together with the 95% confidence band for soil (yellow) and pond (blue) communities. Dashed lines depict scaled pyoverdine production of reference strains (Table S1).

Figure 3. Pyoverdine-containing supernatants from producers can both stimulate and inhibit the growth of non-producers. (A) The likelihood for non-producers to be stimulated by pyoverdine-containing supernatants from a producer did not significantly differ between pond and soil isolates. **(B)** In contrast, the likelihood for non-producers to be inhibited by producer supernatants was significantly higher for pond than for soil isolates. The supernatant assay involved 53 (soil 26 / pond 27) non-producers, and 142 (70/72) producer supernatants from a subset of communities (12/12), resulting in a total of 303 (152/151) non-producer-supernatant combinations, each replicated three to four times. 'Stimulation' or 'inhibition' corresponded to cases where non-producers grew significantly better or worse with supernatant than without, respectively. Box plots show the median (bold line), the 1st and 3rd quartile (box), and the 5th and 95th percentile (whiskers).

Figure 4. Correlations between parameters characterizing pyoverdine-mediated social interactions, environmental variables, and phylogenetic relatedness between strains. For soil non-producers, the likelihood to be stimulated by supernatants containing pyoverdine from a producer correlated positively with SPC1, i.e. it increased with increased carbon concentration, but reduced levels of pH and iron **(A)**; and phylogenetic relatedness between non-producer and producer based on *rpoD* identity **(B)**. For pond non-producers, the

likelihood to be stimulated by producer supernatants correlated negatively with PPC1, i.e. it decreased with decreased pH and community diversity combined with increased carbon and iron concentrations **(C)**. In addition, there was a significant interaction between the *rpoD*-based phylogenetic relatedness between isolates and PPC2, whereby the likelihood of stimulation significantly increased with phylogenetic relatedness for higher values of pH and carbon (high PPC2), but decreased with phylogenetic relatedness for lower values of pH and carbon (low PPC2) **(D)**. Solid lines with shaded areas show significant correlations together with the 95% confidence band for soil (yellow) and pond (blue) communities. Data shown is based on 152 non-producer-supernatant combinations for soil, and 151 non-producer-supernatant combinations for pond.

Table 1. Loadings of the abiotic and biotic environment-defining variables onto the principal components for **(A)** soil and **(B)** pond communities.

(A) Soil	SPC1	SPC2	SPC3	SPC4
community diversity	-0.1729	**0.9358**	-0.2445	-0.1862
pH	**-0.5396**	0.1584	**0.7812**	0.2711
iron	**-0.5751**	-0.1406	-0.5674	0.5723
carbon	**0.5901**	0.2819	0.090	**0.7512**
explained variance [%]	62.3	25.7	9.9	2.1

(B) Pond	PPC1	PPC2	PPC3	PPC4
community diversity	**-0.4892**	-0.3813	**-0.7797**	0.0857
pH	**-0.5161**	**0.5353**	-0.0114	**-0.6686**
iron	**0.5449**	-0.3705	-0.2391	**-0.7132**
carbon	**0.4443**	**0.6564**	-0.5786	0.1924
explained variance [%]	69.5	18.8	9.2	2.5

SPC = Soil Principal Components

PPC = Pond Principal Components

Significant loadings are given in bold