A probabilistic model of sleep structure suggests a primary role for wake probability in overall sleep in *Drosophila melanogaster*

Timothy D. Wiggin[1], Patricia R. Goodwin[1], Nathan C. Donelson[2], Chang Liu[1], and Leslie C. Griffith[1]*

[1]Department of Biology, National Center for Behavioral Genomics and Volen Center for Complex Systems, Brandeis University, Waltham, MA, 02454-9110 USA
[2]Biogen, Cambridge, MA

*Correspondence:
Leslie C. Griffith
Brandeis University
Department of Biology MS008
415 South Street,
Waltham, MA, 02454-9110, USA
Tel: 781 736 3125
griffith@brandeis.edu

Acknowledgements

We would like to thank Dr. Jané Kondev and the students in the Brandeis Quantitative Biology Research Community (QBReC) who assisted in collecting preliminary data.

Grants

This work was funded by R01 MH067284 (to LCG) and F32 NS098624 (to TDW).

ABSTRACT

In humans and flies, sleep is a highly organized behavior, occurring in discrete bouts

that are consolidated at night but have a more fragmented structure during the day.

Longer episodes correlate with transitions to deeper sleep states for both species, and

sleep fragmentation has been shown to have deleterious outcomes for a large number

of physiological processes. In mammals and birds, EEG provides an excellent measure

of sleep structure and state. In invertebrates, assessment of sleep structure has relied

on measures of episode duration, which can fail to capture important aspects of sleep

architecture. Here we model *Drosophila* sleep as a probabilistic behavior and develop

analytical tools to measure transition probabilities and the organization of sleep

episodes within a defined analysis window. We show that these tools provide insight

into the underlying mechanisms of normal sleep and into how they are altered in

pathological conditions. The probability of transition to the wake state, P(Wake) is a

major driver of normal sleep and responsive to dopaminergic tone, suggesting it is a

measure of arousal. Our method for assessing sleep fragmentation adds important

information about the organization of sleep bouts that is missing in measures of episode

duration alone. Importantly, the ability to obtain a measure of the probability structure

underlying sleep in a particular condition or genotype provides a way to compare the

state of brain sleep circuits across conditions that utilizes information from all of the

many processes that influence sleep.

INTRODUCTION

Sleep is a broadly conserved, restorative, and essential behavior. In many animals, sleep progresses through a sequence of stages with successively higher arousal thresholds (for reveiw see Eban-Rothschild and Bloch, 2012). The length of time spent continuously asleep determines the likelihood of reaching deeper sleep states, which are associated with restorative function (Scammell et al., 2017). Conditions which disrupt long episodes of sleep in mammals are associated with cognitive disruption and poor health outcomes (Baud et al., 2013; Lim et al., 2012; Qian et al., 2016; Saint Martin et al., 2016). In the fruit fly *Drosophila melanogaster* there is also evidence for deeper sleep stages that are only reached during long sleep bouts (Nitz et al., 2002; van Alphen et al., 2013; Yap et al., 2017), and sleep has been shown to be important for memory formation (Berry et al., 2015; Donlea et al., 2011). The neural mechanisms that regulate sleep episode duration and the transitions between sleep and wake are therefore of significant clinical and basic science interest.

Transitions between sleep and wake can appear sharp and switch-like. In the early 20th century, von Economo (von Economo, 1930) argued that the transition between sleep and wake was localized to a single "nervous center." In contemporary literature, localization of function is still expressed using the idea of "sleep-promoting" or "wake-promoting" cells or neurons. However, far from arising from a single center, there are many interconnected cell groups that participate in sleep transitions in the mammalian brain (Scammell et al., 2017). Even in *Drosophila*, which have much smaller brains, the localization of sleep initiation is complicated, with dozens of cellular loci that can, in specific contexts, drive sleep/wake transitions (Tomita et al., 2017). While the

idea of a single, localized sleep switch is not tenable, understanding the relationship

between the many neurons that have been shown to be able to affect sleep has been

challenging.

One obstacle to obtaining a more holistic view of the distributed sleep circuit is

that there is not a metric for comparison of the function of different brain areas or

genotypes which captures the structure of sleep or the probabilistic nature of sleep

transitions. In this paper we develop new analysis tools that will be useful for assessing

both sleep structure and the probabilistic nature of behavioral transitions. We provide a

new way to measure sleep fragmentation (the Hamming Index) that captures

information about the organization of episodes as well as their duration. We also

develop a method for obtaining a global assessment of the state of the sleep system

expressed as probabilities of behavioral transitions. This is useful due to the complex

nature of sleep circuits/genes and provides a way to compare the "state" of the system

between genotypes or treatments. Lastly, we provide evidence that state transition

probabilities can be tied to biological processes. We find that P(Wake) is strongly

influenced by dopamine, the major neurochemical involved in arousal in *Drosophila*

(Andretic et al., 2005; Kume et al., 2005).

EXPERIMENTAL PROCEDURES

Animals

Flies were raised on cornmeal-dextrose-yeast food in a 25°C incubator. The

incubator maintained a 12 hour:12 hour light:dark cycle. Flies were collected 0-2 days

post-eclosion and housed in mixed-sex, low-density vials until the beginning of the

experiment (typically 12 male and 12 female flies). All flies used in behavioral

experiments were females that were co-housed with males, and were therefore

presumably mated. For experiments in which the age of the flies is not specified, flies

were 6 – 10 days old at the beginning of data collection. Transgenic flies were as

follows: per^{01}(CS) – gift of Michael Rosbash, w^{1118} and w^{1118}; dat^{fmn} – gift of Amita

Sehgal, PBac{WH}inc^{f00285}, w^{1118} (inc^2; Bloomington Stock Center # 18307).

Sleep Measurements

Flies were briefly anesthetized with CO_2, loaded into glass tubes containing 5%

agarose/2% sucrose food, and tubes were mounted in *Drosophila* Activity Monitor

(DAM) boards (Trikinetics, Waltham, MA). Experiments were carried out at 25°C with a

12 hour:12 hour light:dark cycle, unless otherwise specified. Flies were given a day to

acclimate to the DAM system before data collection began. Sleep deprivation was

performed by securing the DAM systems to a motorized vortexer and shaking the flies

with an inter-pulse period of 10 seconds and a pulse duration of 2 seconds. Unshaken

control flies were housed in the same incubator, but not attached to the vortexer.

Generation of In Silico data

Simulation data was generated by custom programs written in Matlab

(MathWorks, Natick, MA). The Markov chain simulation accepted as input the number of

simulations, the duration (in time steps) of each simulation, and the values of P(Wake)

and P(Doze) for the *in silico* flies. The simulation was initialized in a random state with

equal probability of wake and sleep at *time* = 1. At each subsequent time step *i*, the

simulator: 1) Generated a random number between 0 – 1, 2) Checked activity at *time*(i –

1), 3) Compared the random number with the behavioral transition probability. If the fly

was active at *time*(i – 1), it was compared with P(Doze). If the fly was inactive, it is

compared with P(Wake). 4) If the random number was less than or equal to the

appropriate transition probability, the fly switches states. 5) The simulation proceeded to

the next time step. The first 100 time steps were discarded to minimize the contribution

of the random initialization state to the behavioral output. The Power-Law simulation

proceeded along similar lines, however rather than a single P(Wake) and P(Doze), the

power-law function was solved for the behavioral state duration at time = i, and the

result was compared with the random number 0 – 1. Activity was coded as 1, inactivity

was coded as 0.

Sleep Data Analysis

Both DAM system and *in silico* data were analyzed with a common set of

modular programs written in Matlab. Raw beam crossing data from the DAM was

trimmed to the appropriate experimental time period and binned to 1 minute intervals

using DAMFileScan (TriKinetics). The experimental design and fly genotype metadata

were transcribed into a text file. The Matlab program imported the DAM data and

metadata. Flies with 13 or more hours of continuous inactivity at the end of an

experiment were excluded as potentially dead. Continuous periods of inactivity lasting 5

minutes or longer were classified as sleep. Standard sleep measures were calculated

as follows: Activity while awake was the total beam breaks divided by the number

minutes with activity. Total Sleep was the count of bins classified as sleep. Mean

episode duration was the total sleep divided by the number of episodes.

The Hamming index was calculated as follows: the Hamming distance was

calculated between the experimental data and every possible fully consolidated

equivalent. A fully consolidated equivalent is a sleep record with the same total sleep as

the experimental data, but only one sleep episode. The fully consolidated templates

were generated iteratively by creating an empty variable of equal length to the

experimental window, placing an episode with duration equal to total sleep at the first

bin of that variable, and moving the episode through the duration of the variable one bin

at a time. The minimum value of Hamming distance between the experimental data and

a fully consolidated equivalent is the Hamming Index for that fly. If there is no sleep, or if

the fly exclusively sleeps during the analysis window, the Hamming Index is undefined.

Probability of wake, or P(Wake), was calculated as follows: 1) Count every bin of

inactivity except the last one. This is the denominator, the total number state transitions

from inactivity. 2) For every bin of inactivity, count the number of times the fly is active in

the subsequent bin. This is the numerator, the total number of state transitions from

inactive to active. 3) Divide the total state transitions from inactive to active by the total

state transitions from inactive. If there were no bins where the fly does not move (*i.e.*,

denominator is zero), the transition probability is undefined. Probability of doze, or

P(Doze), was calculated identically, but with activity and inactivity reversed.

Curve fitting was performed using the Matlab curve fitting toolbox using the least

absolute residual method. Fit parameters are reported ± the 95% confidence interval.

Statistical analysis was performed using the Matlab statistics toolbox, data was tested

for main effects using ANOVA, and post-hoc t-tests were corrected using Tukey's

Honestly Significant Difference Procedure (hsd).

RESULTS

Conditional Probability and Information Theory Measures of Sleep

One approach to quantifying the structure of fly sleep is to consider an animal's

behavior as switching between two states: active and inactive. Between each

observation, the fly may choose to either remain in its current state or switch. The

activity pattern of the fly reveals the overall probability of choosing each state, and the

conditional probability of switching e.g. from active to inactive. We propose to call these

conditional probabilities P(Wake), the probability that the fly will switch from the inactive

state to the active state, and P(Doze), the probability that the fly will switch from the

active state to the inactive state (see *Experimental Procedures* and Fig. 1 for detailed

explanation).

We also propose a new method for measuring sleep fragmentation that we call

the Hamming distance sleep fragmentation index, or Hamming Index. The most

commonly used method for analyzing sleep fragmentation is based on sleep episode

duration or episode count. Episode duration and number of episodes are important

features of sleep structure, but information about when the episodes occur is also

important. The Hamming Index is the minimum Hamming distance (number of

mismatches divided by sequence length, Hamming, 1950) between the behavioral

output of the fly and any fully consolidated equivalent – a sleep pattern with the same

total duration of sleep that is organized into a single episode. The Hamming index

therefore captures data about both the degree to which sleep occurs in long episodes

and the clumping of those episodes within an analysis window. Three example activity

patterns with the same total sleep duration but different sleep structures demonstrate

some properties of these measures (Fig. 1).

Circadian and Environmental Regulation of Sleep Structure

In order to assess the usefulness of our approach to quantifying fly sleep

structure, we analyzed sleep data acquired with the well-characterized *Drosophila*

Activity Monitor (DAM) system (Trikinetics, Waltham, MA). Sleep in female wildtype

(WT, *Canton Special*, $n = 60$) and circadian defective (per^{01} mutant, $n = 57$) flies was

measured for 3 days in a 12:12 light:dark cycle (LD), after which the lights were set to

off, and sleep was measured for 6 days in constant darkness (DD) (Fig. 2A-B). There is

a significant main effect of day (ZT0-12) v. night (ZT12-24) and a significant interaction

between time of day and genotype for all measures of sleep quantified (Repeated

measures ANOVA; all $p < 0.0001$). Flies in DD without a functional circadian clock

produce identical sleep during the subjective day and night: the total duration, mean

episode duration, level of fragmentation, and probability of behavioral state transition do

not differ significantly (all $p > 0.3$; Fig. 2C-G). Light alone is a potent cue for

differentiating daytime and nighttime sleep. In per^{01} flies lacking an internal circadian

clock, light causes significant decreases in total sleep and mean episode duration, and

significant increases in sleep fragmentation, P(Wake), and P(Doze) (all $p < 0.0001$; Fig.

2C-G). The circadian clock also drives significant differences between daytime and

nighttime sleep in the absence of light. WT flies in DD conditions have less total sleep,

higher sleep fragmentation, and a higher P(Wake) during subjective day than night (all p

< 0.0001; Fig. 2C-G). Mean sleep episode duration and P(Doze) are regulated by

environmental light alone (all p < 0.0001), but not by the circadian clock in DD (all p >

0.06; Fig. 2D,G). Finally, when flies have two sources of information about time of day,

both an LD cycle and circadian clock, flies sleep more with longer mean episode

duration at night (all p < 0.0001). However, sleep fragmentation as measured by

Hamming Index, is significantly higher at night than during the day (p < 0.0001).

P(Wake) is significantly lower at night in WT flies in LD (p < 0.0001), while P(Doze) is

not significantly regulated by time of day (p = 0.067).

These group comparisons lead to three overall conclusions. First, P(Wake) is

regulated more strongly by environmental light than by the circadian clock. This is

consistent with studies in which LD cycles outside the ability of the clock to entrain are

presented (e.g. Chiesa et al., 2007; Madrid et al., 1998). Second, P(Doze) is not

strongly regulated by light or by the circadian clock. This suggests that P(Wake) may be

the major regulator of total sleep and strongly influence mean episode duration. Lastly,

the regulation of sleep fragmentation appears to be more complex than the regulation of

total sleep.

In order test these group-level associations, we measured both linear and non-

linear relationships between the behavioral state transition probabilities P(Wake) and

P(Doze), and measures of sleep structure (Table 1). Across conditions and genotypes,

P(Wake) is strongly correlated with both total sleep and mean episode duration, but is

not significantly correlated with sleep fragmentation. P(Doze) is moderately, but

significantly, correlated with total sleep, mean episode duration, and sleep

fragmentation. P(Wake) and P(Doze) are not correlated with one another. While these

correlations are suggestive, they cannot unambiguously demonstrate causation,

especially due to the complexity of sleep regulation in flies (Tomita et al., 2017). An *in*

silico approach to sleep offers a more controlled arena in which to investigate the

potential causal relationships among these behavioral parameters.

In Silico Sleep Recapitulates Features of In Vivo Fly Sleep

We used an agent-based simulation paradigm to generate *in silico* sleep data.

Each *in silico* fly was initialized in a random state (either active or inactive) and with a

P(Wake) and P(Doze). At each subsequent time step, the simulation used the current

state of the fly and the behavioral transition probability to determine the new state of the

fly. WT and per^{01} fly sleep data (from Fig. 2) were used to generate empirical heat maps

of the relationship between P(Doze), P(Wake), and sleep parameters (Fig. 3A).

Corresponding *in silico* sleep data was generated for each combination of P(Doze) and

P(Wake) (range: 0-1 binned to 25 steps, $n = 100$ simulations per bin; Fig. 3B).

Comparison of the *in vivo* and *in silico* datasets shows qualitative similarity with some

notable quantitative differences, especially in mean sleep episode duration (Fig. 3A,B,

middle column). One possible explanation for this quantitative discrepancy is that our

model of state transition probability is too simple. For example, previous reports have

shown that sensory thresholds and neural activity change as a function of the duration

that the fly has been asleep (Nitz et al., 2002; van Alphen et al., 2013; Yap et al., 2017).

This is similar to the situation in humans, where transition models have to consider

sleep sub-states (Perez-Atencio et al., 2018).

In order to integrate this information into our models, we calculated P(Wake) and

P(Doze) as a function of behavioral state duration (Fig. 3C,D). We found both P(Wake)

and P(Doze) have a power-law relationship with the duration of activity state. We then

used the values for P(Wake) and P(Doze), and the best-fit power-law functions, to

generate new *in silico* data matched to LD day, LD night, DD day, and DD night in WT

flies. We then compared the deviation of our simulations from the *in vivo* WT fly data (*n*

= 100 *in silico* flies per condition; Fig. 3E). We refer to the simulations with a fixed

P(Wake) and P(Doze) as "Markov Chain simulations," because they are a special case

of a Markov chain. Both the Markov chain and power law *in silico* flies accurately

reproduce total sleep of their matched *in vivo* group (Fig. 3E, *left column*). The Markov

chain simulations systematically underestimate sleep episode duration, while the power

law simulations match the *in vivo* data (Fig. 3E, *middle column*). Both the Markov and

power law simulations produce more fragmented *in silico* sleep than shown in the *in*

vivo data (Fig. 3E, *right column*). Across groups, the majority of both the Markov chain

and power law simulations show small standardized differences from *in vivo* data (|*Z*-

score| < 2), indicating that the relatively low complexity simulations are capable of

capturing a meaningful fraction of the *in vivo* behavior.

Our simulation approach assumes that population-wide fit of behavioral state

transition probability can be used to simulate *in silico* individuals. We tested the

assumption that individual fly behavior is faithfully represented by population means

using two approaches (Fig. 4). First, we tested the hypothesis that variation of a single

fly's behavior over time is comparable to the variation in behavior between flies (Fig.

4A,B). We found instead that for both P(Wake) and P(Doze), there is significantly

greater variability between flies than within a single fly across time (two tailed *T*-test; p <

0.0001). Second, we tested the hypothesis that estimates of P(Wake) and P(Doze)

derived from each individual fly would converge toward the population estimate of these

measures. Population estimates of P(Wake) and P(Doze) are centrally located within

the distribution of individual fly derived estimates (Fig. 4C,D). Population estimates of

the power law distribution parameters of P(Wake) and P(Doze) are also relatively

faithfully reproduced by the distribution of individual fly estimates, with the notable

exception of the lower-bound parameter of P(Wake) (Fig. 4E-J). In all individual flies,

the lower bound of P(Wake) was estimated to be zero, while as a population the lower

bound was 0.5% (Fig. 4I). This discrepancy is likely due to the difficulty of getting a

good estimate of low frequency events from limited data, and would be most apparent

when estimating the frequency of long duration sleep episodes. While it is in principle

possible to estimate the single fly P(Wake) and P(Doze), fitting a power law function to

individual fly data is limited by the ability to accurately assess long-duration episode

frequency. The distribution of P(Wake) and P(Doze) and their power law fit parameters

are unimodal, allowing population-based averages to approximate a "typical" fly.

Effect of Increased Homeostatic Sleep Pressure on Behavioral State Transitions

Sleep deprivation in flies, like in other animals, leads to increased subsequent

sleep (Shaw et al., 2000). A common model for the generation of increased sleep after

release from sleep deprivation is that there is an increase in sleep "pressure" which

alters sleep drive (Fuller et al., 2006). In order to quantify how sleep pressure affects

behavioral state transitions, we performed a sleep deprivation experiment during which

WT flies were shaken periodically for 12 hours overnight (n = 120 shaken, 117 control;

Fig. 5A). Compared with a control group of flies, flies that were shaken slept less while

the shaker was on, and more the following day (Fig. 5A). We defined two time windows:

PRE, ZT0-6 the morning prior to sleep deprivation, and POST, ZT0-6 the morning

following sleep deprivation. Sleep deprived flies, but not control flies, slept significantly

more POST than PRE and had significantly less sleep fragmentation as assessed both

by Hamming Index (all p < 0.0001; Fig. 5B,C) and mean episode duration (in min, for

shaken group, PRE = 40.9 (SD 44.4), POST = 206.8 (SD 137.9), p < 0.0001, for control

group, PRE = 40.2 (SD 49.1), POST = 52.2 (SD 61.7), p = 0.2). P(Wake) was

significantly decreased and P(Doze) was significantly increased in sleep deprived flies

(all p < 0.0001), but not control flies, during the POST interval (Fig. 5D,E).

While both P(Wake) and P(Doze) are significantly different following sleep

deprivation, three lines of evidence suggest that P(Wake) is principally responsible for

the change in behavior: 1) change in total sleep during POST v. PRE is more strongly

correlated with change in P(Wake) (R = -0.97, p < 0.0001) than change in P(Doze) (R =

0.19, p = 0.06), 2) The effect size of the change in P(Wake) is larger than the effect size

of P(Doze) (Cohen's d = -1.46 and 0.81, respectively, (Cohen, 1992)) and 3)

Superimposing the observed change in P(Wake) and P(Doze) on simulated data, the

change along the P(Wake) axis is more consequential than the change along the

P(Doze) axis (Fig. 5F,G).

Examining the history-dependence of P(Wake) and P(Doze) provides additional

insight into how sleep pressure modifies behavioral transitions (Fig. 5H,I). Neither the

initial value (PRE: 1.24 ± 0.08, POST:1.30 ± 0.06) nor the long-term lower bound of

P(Wake) (PRE: 0, POST: 0) differs between PRE and POST, but the power law

exponent of decay is substantially faster in POST (PRE: 1.03 ± 0.04, POST: 1.20 ±

0.04; Fig. 5H). The shape of P(Doze) diverges between PRE and POST during

movement episodes longer than 6 minutes (Fig. 5I) but, because fewer than 20% of

wake episodes last longer than 6 minutes, this divergence does not result in decreased

overall P(Doze) (Fig. 5E). In WT flies, it appears that the primary mechanism generating

homeostatic rebound sleep is a decreased probability of waking up during a sleep

episode rather than the initiation of additional episodes. The change in P(Wake) does

not result from a generalized decrease, but from the flies more rapidly entering deep

sleep after initiating a sleep episode. It is clear from the heat maps (Fig. 5F,G),

however, that if a fly's initial position (i.e. their respective P(Wake) and P(Doze)) was in

the lowest left part of the map, the effect of changes in P(Doze) might be behaviorally

relevant. The fact that both mechanisms are engaged could be important in animals,

e.g. mutants, whose initial sleep parameters are abnormal.

Mutations Affecting Sleep Act Through Both Wake and Doze Probability

Sleep is regulated by a multitude of molecular processes, and many null and

hypomorphic mutations disrupt sleep in flies (Helfrich-Forster, 2018). We examined how

two known sleep mutants, *fumin* (*dat^{fmn}*; *n* = 39 mated female flies; Kume et al., 2005)

and *insomniac* (*inc^2*; *n* = 33; Stavropoulos and Young, 2011), affect behavioral state

transition probabilities (Fig. 6). We found that *fumin* flies show the previously

demonstrated decreased total sleep relative to control flies with the same genetic

background (*w^{1118}*; *n* = 40; p < 0.0001; Fig.6A,B). Female *insomniac* flies, in our

experimental conditions, unexpectedly have increased daytime total sleep and

unchanged nighttime sleep ($p < 0.0001$ and $p = 0.16$, respectively; Fig. 6A,B).

Interestingly, we find that *insomniac* flies are significantly hyperactive when awake ($p <$

0.0001; Fig. 6C) in addition to being hyper-somnolent. Both *fumin* and *insomniac* flies

have significantly shorter mean sleep episode duration during the day ($p < 0.012$), but

only *fumin* flies have shorter sleep episodes at night ($p < 0.0001$; Fig. 6D). Finally,

during the day *fumin* flies have significantly less fragmented sleep while *insomniac* flies

have significantly more fragmented sleep ($p < 0.0001$). There are no significant

differences in sleep fragmentation at night as assessed by Hamming Index ($p > 0.7$).

Visual examination of individual sleep rasters confirms that despite shorter mean sleep

episodes, sleep and activity in *fumin* flies is not substantially more disorganized than in

wild type (Fig. S1). In summary, *fumin* flies sleep less than WT and in shorter episodes

both during the day and night. *Insomniac* sleep only differs from WT sleep during the

day, during which they sleep more, in shorter episodes, with greater sleep

fragmentation.

Both the *fumin* and *insomniac* phenotypes can be explained by changes in

P(Wake) and P(Doze) (Fig. 6F-G). *Fumin* flies have increased P(Wake) and decreased

P(Doze) both in the day and night ($p < 0.002$), which our simulations predict would

cause the observed phenotypes (Fig. 6H,G). The *insomniac* mutation does not

significantly alter P(Wake) ($p > 0.25$), but significantly increases P(Doze) both during the

day and night ($p < 0.0001$), and shows behavior that consistent with our *in silico*

predictions (Fig. 6H,I). The power law functions of P(Wake) and P(Doze) provide

additional insight into this process (Fig. 6J,K). The increase in P(Wake) between WT

and *fumin* flies is driven by a difference between the respective power law exponents of

decay (WT: 1.16 ± 0.03, *fumin*: 0.83 ± 0.07; Fig. 6H). The increase in P(Doze) between

WT and *insomniac* flies is driven by differences between both the initial value of

P(Doze) (WT: 0.62 ± 0.03, *insomniac*: 1.06 ± 0.31) and the respective power law

exponents of decay (WT: 0.66 ± 0.02, *insomniac*: 0.47 ± 0.16). The divergent effects of

these two mutations, while complex, are well explained by a relatively simple model of

behavioral state transition probability.

DISCUSSSION

Sleep serves many physiological purposes, so it is unsurprising that it should be

responsive to many different variables. Competing or interacting switches which are

organized hierarchically can allow many factors to be integrated, but how to express the

global state of such a complex system is not clear. In this paper we model sleep in

terms of the probabilities of state transitions. We also introduce a new measure of sleep

fragmentation, the Hamming Index, that gives a quantitative assessment of the

organization of the sleep episodes in the epoch being studied.

Sleep as a probabilistic behavior has been extensively modelled in both rodents

and humans (Bianchi et al., 2012; Perez-Atencio et al., 2018; Stephenson et al., 2016;

Yang and Hursch, 1973). Previous descriptions of *Drosophila* sleep structure have

focused on describing the relative frequency of sleep episodes of different durations

(Ueno et al., 2012; Diamond, 2016). These reports each found that fly sleep follows a

power-law distribution of sleep episode duration, and that the distribution is affected by

a variety of factors, including environment and dopamine levels. Each report also

suggested a metric for quantifying sleep structure: coefficient of determination

(Diamond, 2016), and power-law fit coefficients (Ueno et al., 2012). This report both

agrees with and builds upon this previous work. We find that P(Wake) is strongly

correlated with mean episode duration (R = -0.89) and that both P(Wake) and P(Doze)

follow a power-law dependence upon duration of sleep and wake, respectively (Fig. 3).

We also find that dopamine profoundly affects P(Wake), consistent with a shift in the

distribution of episode duration (Ueno et al., 2012). By focusing not on sleep episodes

themselves, but on the state transitions that define sleep episodes, we also gain insight

into the mechanisms that generate the observed distribution of sleep episode duration.

This mechanistic advantage is most apparent in the generation of *in silico* data for a

wide range of conditional probabilities, and allows us to plot the relative positions of flies

in a conditional probability space.

Fragmentation of sleep is a critical human health issue, and the major way that

this is assessed in patient populations is by use of measures analogous to sleep bout

duration (sleep stage percentages and sleep efficacy) from polysomnography records.

These methods, however, can fail to capture fragmentation even in patients with

obstructive sleep apnea (Swihart et al., 2008). Models that extract information about

transitions and the underlying organization of episodes have been shown to be better

predictors of fragmentation and morbidity (Bianchi et al., 2010; Bianchi et al., 2012;

Swihart et al., 2008). For *Drosophila* sleep the bout length serves as the major metric

for fragmentation. Episode duration, however, can fail to identify changes in

consolidation. For example in wildtype animals in DD there is no difference between

episode duration in subjective day and night, but there are clock-dependent processes

consolidating sleep that can be detected in the Hamming Index (Fig. 2).

There are, however, cases where Hamming Index can indicate no fragmentation

even when there is a decrease mean episode duration. Total sleep in *fumin* animals,

which have high dopaminergic tone (Kume et al., 2005), is severely decreased, as is

mean episode duration. Hamming Index for this genotype is actually decreased during

the day and unchanged at night. This is likely due to the fact that there are very few

sleep episodes, which decreases the degrees of freedom for the organization of sleep in

that window. Examination of the sleep rasters of these genotypes (Fig. S1), however,

suggests that there are differences in organization that are not picked up when

considering the entire night as one analysis block. Sleep during the night is likely not

regulated by a single process- early night sleep and late night sleep show differences in

bout length and structure in both in humans (Norman et al., 2006; Yang and Hursch,

1973) and flies (Kunst et al., 2014). A sliding window analysis of Hamming Index (or of

bout length) would likely better capture nighttime sleep structure and provide a better

platform for identifying fragmentation. In most cases, however, even when considering

the night as a single analysis window, mean episode duration and Hamming Index are

in agreement and since they provide complimentary information, the use of Hamming

Index will increase confidence in the assessment of relative consolidation.

The utility of these new analysis tools is demonstrated by using them on sleep

data from wild type and clockless (*per[01]*) mutants. We find that both light and the

circadian clock can alter the structure of sleep. For flies without a functioning clock, the

structure of sleep is completely dependent on daily exposure to light. Light drives the

majority of daily sleep into the night and also reduces its fragmentation as measured by

both mean episode duration and Hamming Index. For wildtype flies, light during the day

also appears to promote nighttime sleep. Most of the difference between day and night

sleep can be attributed to changes in P(Wake). This could be explained by actions of

dopamine, which have been shown to influence the sleep function of the clock in LD

(Shang et al., 2011).

Pathological changes in sleep secondary to acute deprivation or mutations can

also be described using our metrics. Homeostatic rebound sleep in wild type animals,

like normal sleep, appears to be controlled primarily by P(Wake) and results in a

significant decrease in fragmentation as measured by the Hamming Index and mean

episode duration.

The two sleep mutants we examined appear to affect sleep via changes in

different underlying parameters. *Fumin*, which is a mutation in the dopamine reuptake

transporter, drastically increases P(Wake), in line with the idea that dopamine is a major

determinant of arousal. *Insomniac* flies on the other hand increase P(Doze) during both

the day and night. The analysis of transition probabilities suggests that while there are

similarities in these two genotypes at a macroscopic scale, the underlying molecular

and circuit processes they affect must be different.

Arousal state has a major influence on sleep amount and architecture (Ouyang et

al., 2004; Phillips et al., 2013; Scammell et al., 2017; Tagliazucchi and van Someren,

2017) and may be one of the key underlying processes that coordinate the actions of

the disparate sleep circuits. Using our new analysis tools, we show for wild type flies,

the major determinant of sleep structure is P(Wake), and we posit this is a direct

measure of arousal. P(Wake) is tied to the activity of dopaminergic neurons since *fumin*

mutants, which have an increase in dopaminergic tone (Kume et al., 2005) have a

greatly increased P(Wake). This suggests that P(Wake) may be an important parameter

to measure when assessing the global state of the distributed sleep circuit, arousal and

the involvement of dopaminergic systems in a phenotype.

LITERATURE CITED

472 Andretic, R., van Swinderen, B., and Greenspan, R.J. (2005). Dopaminergic modulation

of arousal in Drosophila. Curr Biol *15*, 1165-1175.

Baud, M.O., Magistretti, P.J., and Petit, J.M. (2013). Sustained sleep fragmentation

affects brain temperature, food intake and glucose tolerance in mice. Journal of sleep

research *22*, 3-12.

Berry, J.A., Cervantes-Sandoval, I., Chakraborty, M., and Davis, R.L. (2015). Sleep

Facilitates Memory by Blocking Dopamine Neuron-Mediated Forgetting. Cell *161*, 1656-

1667.

Bianchi, M.T., Cash, S.S., Mietus, J., Peng, C.K., and Thomas, R. (2010). Obstructive

sleep apnea alters sleep stage transition dynamics. PLoS ONE *5*, e11356.

Bianchi, M.T., Eiseman, N.A., Cash, S.S., Mietus, J., Peng, C.K., and Thomas, R.J.

(2012). Probabilistic sleep architecture models in patients with and without sleep apnea.

Journal of sleep research *21*, 330-341.

Chiesa, J.J., Diez-Noguera, A., and Cambras, T. (2007). Effects of transient and

continuous wheel running activity on the upper and lower limits of entrainment to light-

dark cycles in female hamsters. Chronobiol Int *24*, 215-234.

Cohen, J. (1992). Statistical power analysis. Current directions in psychological science.

Diamond, J.M. (2016). Goodness of fit to a mathematical model for Drosophila sleep

behavior is reduced in hyposomnolent mutants. PeerJ *4*, e1533.

Donlea, J.M., Thimgan, M.S., Suzuki, Y., Gottschalk, L., and Shaw, P.J. (2011).

Inducing sleep by remote control facilitates memory consolidation in Drosophila.

Science *332*, 1571-1576.

Eban-Rothschild, A., and Bloch, G. (2012). Social influences on circadian rhythms and

sleep in insects. Adv Genet *77*, 1-32.

Fuller, P.M., Gooley, J.J., and Saper, C.B. (2006). Neurobiology of the sleep-wake

cycle: sleep architecture, circadian regulation, and regulatory feedback. J Biol Rhythms

21, 482-493.

Hamming, R.W. (1950). "Error detecting and error correcting codes". The Bell System

Technical Journal *29*, 147-160.

Helfrich-Forster, C. (2018). Sleep in Insects. Annu Rev Entomol *63*, 69-86.

Koh, K., Evans, J.M., Hendricks, J.C., and Sehgal, A. (2006). A Drosophila model for

age-associated changes in sleep:wake cycles. Proc Natl Acad Sci U S A *103*, 13843-

13847.

Kume, K., Kume, S., Park, S.K., Hirsh, J., and Jackson, F.R. (2005). Dopamine is a

regulator of arousal in the fruit fly. J Neurosci *25*, 7377-7384.

Kunst, M., Hughes, M.E., Raccuglia, D., Felix, M., Li, M., Barnett, G., Duah, J., and

Nitabach, M.N. (2014). Calcitonin gene-related peptide neurons mediate sleep-specific

circadian output in Drosophila. Curr Biol *24*, 2652-2664.

Lim, A.S., Yu, L., Costa, M.D., Leurgans, S.E., Buchman, A.S., Bennett, D.A., and

Saper, C.B. (2012). Increased fragmentation of rest-activity patterns is associated with a

characteristic pattern of cognitive impairment in older individuals. Sleep *35*, 633-640B.

Madrid, J.A., Sanchez-Vazquez, F.J., Lax, P., Matas, P., Cuenca, E.M., and Zamora, S.

(1998). Feeding behavior and entrainment limits in the circadian system of the rat. Am J

Physiol *275*, R372-383.

Nitz, D.A., van Swinderen, B., Tononi, G., and Greenspan, R.J. (2002).

Electrophysiological Correlates of Rest and Activity in *Drosophila melanogaster*. Curr

Biol *12*, 1934-1940.

Norman, R.G., Scott, M.A., Ayappa, I., Walsleben, J.A., and Rapoport, D.M. (2006).

Sleep continuity measured by survival curve analysis. Sleep *29*, 1625-1631.

Ouyang, M., Hellman, K., Abel, T., and Thomas, S.A. (2004). Adrenergic signaling plays

a critical role in the maintenance of waking and in the regulation of REM sleep. J

Neurophysiol *92*, 2071-2082.

Perez-Atencio, L., Garcia-Aracil, N., Fernandez, E., Barrio, L.C., and Barios, J.A.

(2018). A four-state Markov model of sleep-wakefulness dynamics along light/dark cycle

in mice. PLoS ONE *13*, e0189931.

Phillips, A.J., Robinson, P.A., and Klerman, E.B. (2013). Arousal state feedback as a

potential physiological generator of the ultradian REM/NREM sleep cycle. Journal of

theoretical biology *319*, 75-87.

Qian, Y., Yi, H., Zou, J., Meng, L., Tang, X., Zhu, H., Yu, D., Zhou, H., Su, K., Guan, J.*,*

et al. (2016). Independent Association between Sleep Fragmentation and Dyslipidemia

in Patients with Obstructive Sleep Apnea. Scientific reports *6*, 26089.

Saint Martin, M., Sforza, E., Crawford-Achour, E., Pascal, L., Leinard, G., and Frederic,

R. (2016). Sleep breathing disorders and cognitive decline in healthy elderly followed for

eight years: the PROOF cohort. Annals of Physical and Rehabilitation Medicine *595*,

e99.

Scammell, T.E., Arrigoni, E., and Lipton, J.O. (2017). Neural Circuitry of Wakefulness

and Sleep. Neuron *93*, 747-765.

Shang, Y., Haynes, P., Pirez, N., Harrington, K.I., Guo, F., Pollack, J., Hong, P., Griffith,

L.C., and Rosbash, M. (2011). Imaging analysis of clock neurons reveals light buffers

the wake-promoting effect of dopamine. Nat Neurosci *14*, 889-895.

Shaw, P.J., Cirelli, C., Greenspan, R.J., and Tononi, G. (2000). Correlates of sleep and

waking in *Drosophila melanogaster*. Science *287*, 1834-1837.

Stavropoulos, N., and Young, M.W. (2011). insomniac and Cullin-3 regulate sleep and

wakefulness in Drosophila. Neuron *72*, 964-976.

Stephenson, R., Caron, A.M., and Famina, S. (2016). Significance of the zero sum

principle for circadian, homeostatic and allostatic regulation of sleep-wake state in the

rat. Physiology & behavior *167*, 35-48.

Swihart, B.J., Caffo, B., Bandeen-Roche, K., and Punjabi, N.M. (2008). Characterizing

sleep structure using the hypnogram. J Clin Sleep Med *4*, 349-355.

Tagliazucchi, E., and van Someren, E.J.W. (2017). The large-scale functional

connectivity correlates of consciousness and arousal during the healthy and

pathological human sleep cycle. Neuroimage *160*, 55-72.

Tomita, J., Ban, G., and Kume, K. (2017). Genes and neural circuits for sleep of the fruit

fly. Neuroscience research *118*, 82-91.

Ueno, T., Masuda, N., Kume, S., and Kume, K. (2012). Dopamine modulates the rest

period length without perturbation of its power law distribution in Drosophila

melanogaster. PLoS ONE *7*, e32007.

van Alphen, B., Yap, M.H., Kirszenblat, L., Kottler, B., and van Swinderen, B. (2013). A

dynamic deep sleep stage in Drosophila. J Neurosci *33*, 6917-6927.

Vienne, J., Spann, R., Guo, F., and Rosbash, M. (2016). Age-Related Reduction of

Recovery Sleep and Arousal Threshold in Drosophila. Sleep *39*, 1613-1624.

von Economo, C. (1930). Sleep as a problem of localization. Journal of Nervous and

Mental Disease *71*, 249-259.

Yang, M.C., and Hursch, C.J. (1973). The use of a semi-Markov model for describing

sleep patterns. Biometrics *29*, 667-676.

Yap, M.H.W., Grabowska, M.J., Rohrscheib, C., Jeans, R., Troup, M., Paulk, A.C., van

Alphen, B., Shaw, P.J., and van Swinderen, B. (2017). Oscillatory brain activity in

spontaneous and induced sleep stages in flies. Nature communications *8*, 1815.

Figure 1 – Conditional Probability and Information Theory Measures of Sleep.

Three exemplar sleep patterns (Observation #1-3) are shown that have the same total

sleep and median time of sleep, but different structure. White squares: wake, Black

squares: sleep. The fully consolidated equivalent of each of these observations is the

same (middle rows). During the calculation of Hamming Index, the Hamming distance to

all possible fully consolidated equivalents is calculated to find the minimum value.

Blocks that differ between each observation and the fully consolidated equivalent are

red, blocks that are the same are gray. Transitions between behavioral states are

indicated by curved arrows. Transitions from active to inactive (Doze) are above the

blocks, transitions from inactive to active (Wake) are below. The calculation for P(Wake)

is as follows: Count every measurement bin where the fly does not move except the last

one. This the total number of transitions *from* inactive. For every measurement bin

where the fly does not move, count the number where the fly moves at the next bin.

This is the number of transitions *from* inactive *to* active. Divide the transitions *from*

inactive *to* active by the total number of transitions to calculate the transition probability

of inactive to active, or P(Wake). The calculation for P(Doze) proceeds in the same way,

630   but with "active" and "inactive" switched. See *Experimental Procedures* for additional

631   details.

Figure 2 – Circadian and Environmental Regulation of Sleep Structure. (A-B)

Circadian profile of sleep in a 12:12 Light:Dark cycle (LD; A) and in constant darkness

(DD; B). Shaded regions indicate lights on and lights off times (A) or subjective day and

night (B). Lines are the population average, shaded area is the 95% confidence interval

of the mean. Individual profiles are the average of 3 days of behavior. DD sleep

measured on days 4-6 of constant darkness. (C-G) Box and whisker plots of total sleep

(C), mean sleep episode duration (D), Hamming index (E), P(Wake) (F), and P(Doze)

(G). Boxes show the 25/50/75th percentiles for each group, whiskers indicate the

minimum and maximum. *n.s.* indicates no significant difference, * $p < 0.05$, ** $p < 0.005$,

*** $p < 0.0005$.

TABLE 1 – Spearman Correlations Among Sleep Measures

	P(Wake) Rho	P(Wake) p-value	P(Doze) Rho	P(Doze) p-value
Total Sleep	-0.9804	$< 10^{-10}$	0.2221	1.2129e-06
Mean Episode Duration	-0.8869	$< 10^{-10}$	-0.1588	5.7062e-04
Hamming Index	-0.0249	0.5918	0.4665	$< 10^{-10}$
P(Doze)	-0.0769	0.0966		

Figure 3 – *In Silico* Sleep Recapitulates Features of *In Vivo* Fly Sleep. (A, B)

Heatmaps of *in vivo* (A) and *in silico* (B) sleep properties. (A) Each *In vivo* fly was

assigned to one of 25 bins from 0 to 1 based upon it's respective P(Doze) and P(Wake).

The mean value of total sleep (*left*), mean sleep episode duration (*center*), and

Hamming index (*right*) was calculated for all samples in the bin, and the bin was

assigned a color according to the legend (*bottom*). Coordinates without any samples are

displayed as white. (B) For the central value of each P(Doze) and P(Wake) bin, 100 *in*

silico flies were simulated. The mean value of each parameter of these simulations is

displayed using the same color map as the *in vivo* data. Coordinates coded white for *in*

silico Hamming index have an undefined value because no behavioral transitions were

generated for the duration of the epoch. (C, D) P(Wake) and P(Doze) plotted as a

function of the duration of inactivity (C) and activity (D), respectively. These data are

derived from the night epoch of CS flies in LD conditions. Data was fit using the

equation $a(x + 1)^{-b} + c$, where *a*, *b*, and *c* are free parameters. Black points are

experimental data, red and green lines are the power law functions that best fit the data.

(E) *Z*-scores of *in vivo* and *in silico* sleep. The population-derived values of P(Doze) and

P(Wake), and their power-law equivalents, were derived for each listed *in vivo*

experimental condition and used to simulate sleep behavior. *Z*-scores were calculated

using the sample mean and standard deviation of the corresponding *in vivo*

experimental group.

Figure 4 – Population and Individually Derived Estimates of Behavioral State

Transition Probability. (A, B) Histograms of median absolute deviation (MAD) of

P(Wake) (A) and P(Doze) (B). Intra-fly variability is calculated based on 3 day epochs

and 3 night epochs of behavior from each fly. Inter-fly variability is based on the MAD of

the population during each epoch (*n* = 60 WT flies). Low sample size of inter-fly

variability in (A) precludes concluding if the data are bimodally distributed. (C, D)

Histograms of P(Wake) and P(Doze) calculated for each fly. The population derived

estimate is shown as a dashed red line. (E-J) Histograms of the estimates of each

parameter, *a, b,* and *c*, of a power law fit using the equation $a(x + 1)^{-b} + c$ of individual

fly behavior. Data in (E-J) are from the nighttime behavior of CS flies in LD conditions.

Figure 5 - Effect of Increased Homeostatic Sleep Pressure on Behavioral State

Transitions. (A) Circadian profile of sleep over 3 days, including one night of sleep

deprivation (labeled "Shaker on"). Shaded regions indicate lights on and lights off times.

Analysis windows for PRE and POST are indicated. Mean sleep is indicated by lines,

shaded region is the 95% confidence interval. (B-E) Box and whisker plots of total sleep

(B), Hamming index (C), P(Wake) (D), and P(Doze) (E). Boxes indicate the 25/50/75[th]

percentiles for each group, whiskers indicate the minimum and maximum. (F,G)

Heatmaps of simulated sleep at each level of P(Wake) and P(Doze), reproduced from

Fig. 3. Circles indicate the coordinates of the mean P(Wake) and P(Doze) of sleep

deprivation group before sleep deprivation (white circles) and following sleep

deprivation (black circles). Dashed lines show the change along each dimension. (H,I)

Power law function fits of P(Wake) and P(Doze) during PRE and POST. PRE: blue,

POST: magenta. Open circles are the empirical data, lines are the best fit, and shaded

regions are the 95% prediction bounds for the fit. *n.s.* indicates no significant difference,

* $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.

Figure 6 - Mutations Affecting Sleep Act Through Both Wake and Doze

Probability. (A) Circadian profile of sleep in a 12:12 Light:Dark cycle. Shaded regions

indicate lights on and lights off times. Lines are the population average, shaded area is

the 95% confidence interval of the mean. Individual profiles are the average of 3 days of

behavior. (B-G) Box and whisker plots of total sleep (B), activity while awake (C), mean

sleep episode duration (D), Hamming index (E), P(Wake) (F), and P(Doze) (G). Boxes

indicate the 25/50/75th percentiles for each group, whiskers indicate the minimum and

maximum. (H,I) Heatmaps of simulated sleep at each level of P(Wake) and P(Doze),

reproduced from Fig. 2. Circles indicate the coordinates of the mean P(Wake) and

P(Doze) of the isotype control (*w^{1118}*; black), *fumin* (red), and *insomniac* (blue) flies

during daytime. (J,K) Power law function fits of P(Wake) and P(Doze) of each genotype

during daytime. *w^{1118}*: black, *fumin*: red, *insomniac*: blue. Open circles are the empirical

data, lines are the best fit, and shaded regions are the 95% prediction bounds for the fit.

The P(Wake) *fumin* fit and P(Doze) *insomniac* fit do not extend across the full range of

x-axis because fits were only evaluated in the range for which 2 empirical data points

were collected, and no flies of those genotypes produced sufficiently long episodes of

inactivity or activity, respectively. *n.s.* indicates no significant difference, * $p < 0.05$, ** p

< 0.005, *** $p < 0.0005$.

Figure S1 – Sleep Rasters of Control and *fumin* flies. 24 hours sleep rasters of ten

w[1118] flies (Top) and ten *fumin* flies (Middle and Bottom). In the top two sets of rasters,

white is wake, black is sleep. In the bottom set of rasters, white is sleep and black is

wake. The gray filled area is night.