# Neo-formation of chromosomes in bacteria

Olivier B. Poirion[1,2†] & Bénédicte Lafay[1,2,3*]

[1] Université de Lyon, F-69134 Lyon, France

[2] CNRS (French National Center for Scientific Research) UMR5005, Laboratoire Ampère, École Centrale de Lyon, 36 avenue Guy de Collongue, 69134 Écully, France

[3] CNRS (French National Center for Scientific Research) UMR5558, Laboratoire de Biométrie et Biologie Évolutive, Université Claude Bernard − Lyon 1, 43 boulevard du 11 novembre 1918, 69622 Villeurbanne cedex, France

[†] Current address: Center for Epigenomics, Department of Cellular and Molecular Medicine, University of California, San Diego, School of Medicine, 9500 Gilman Drive, La Jolla, CA 92093, USA

[*] Author for correspondence: benedicte.lafay@univ-lyon1.fr

**ABSTRACT**

Although the bacterial secondary chromosomes/megaplasmids/chromids, first noticed

about forty years ago, are commonly held to originate from stabilized plasmids, their

true nature and definition are yet to be resolved. On the premise that the integration of a

replicon within the cell cycle is key to deciphering its essential nature, we show that the

content in genes involved in the replication, partition and segregation of the replicons

and in the cell cycle discriminates the bacterial replicons into chromosomes, plasmids,

and another class of essential genomic elements that function as chromosomes. These

latter do not derive directly from plasmids. Rather, they arise from the fission of a multi-

replicon molecule corresponding to the co-integrated and rearranged ancestral

chromosome and plasmid. All essential replicons in a distributed genome are thus neo-

chromosomes. Having a distributed genome appears to extend and accelerate the

exploration of the bacterial genome evolutionary landscape, producing complex

regulation and leading to novel eco-phenotypes and species diversification.

 **INTRODUCTION**

Chromosomes are the only components of the genome that encode the necessary

information for replication and life of the cell/organism under normal growth conditions.

Their number varies across taxa, a single chromosome being the standard in bacteria

(Krawiec and Riley, 1990). Evidence accumulated over the past forty years is proving

otherwise: bacterial genomes can be distributed on more than one chromosome-like

autonomously replicating genomic element (replicon) (Casjens, 1998; diCenzo and

Finan, 2017; Mackenzie et al., 2004). The largest, primary, essential replicon (ER) in a

multipartite genome corresponds to a *bona fide* chromosome and the additional,

secondary, ERs (SERs) are expected to derive from accessory replicons (plasmids

(Lederberg, 1998)). The most popular model of SER formation posits that a plasmid

acquired by a mono-chromosome progenitor bacterium is stabilized in the genome

through the transfer from the chromosome of genes essential to the cell viability

(diCenzo and Finan, 2017; diCenzo et al., 2013; Slater et al., 2009). The existence in

SERs of plasmid-like replication and partition systems (Dubarry et al., 2006; Egan and

Waldor, 2003; Livny et al., 2007; MacLellan et al., 2004, 2006; Slater et al., 2009;

Yamaichi et al., 2007) as well as experimental results (diCenzo et al., 2014) support this

view. Yet, the duplication and maintenance processes of SERs contrast with the typical

behaviour of plasmids for which both the timing of replication initiation and the

centromere movement are random (Million-Weaver and Camps, 2014; Reyes-Lamothe

et al., 2014). Indeed, the SERs share many characteristic features with chromosomes:

enrichment in Dam methylation sites of the replication origin (Egan and Waldor, 2003;

Gerding et al., 2015), presence of initiator titration sites (Egan and Waldor, 2003;

Venkova-Canova and Chattoraj, 2011), synchronization of the replication with the cell

cycle (De Nisco et al., 2014; Deghelt et al., 2014; Egan and Waldor, 2003; Egan et al.,

2004; Fiebig et al., 2006; Frage et al., 2016; Kahng and Shapiro, 2003; Rasmussen et al.,

2007; Srivastava et al., 2006; Stokke et al., 2011), KOPS-guided FtsK translocation (Val

et al., 2008), FtsK-dependent dimer resolution system (Val et al., 2008), MatP/matS

macrodomain organisation system (Demarre et al., 2014), and similar fine-scale

segregation dynamics (Fiebig et al., 2006; Frage et al., 2016). Within a multipartite

genome, the replication of the chromosome and that of the SER(s) are initiated at

different time points (De Nisco et al., 2014; Deghelt et al., 2014; Fiebig et al., 2006;

Frage et al., 2016; Rasmussen et al., 2007; Srivastava et al., 2006; Stokke et al., 2011),

and use replicon-specific systems (Drevinek et al., 2008; Egan and Waldor, 2003;

Galardini et al., 2013; MacLellan et al., 2004, 2006; Slater et al., 2009). Yet, they are

coordinated, hence maintaining the genome stoichiometry (Deghelt et al., 2014; Egan et

al., 2004; Fiebig et al., 2006; Frage et al., 2016; Stokke et al., 2011). In the few species

where this was studied, the replication of the SER is initiated after that of the

chromosome (De Nisco et al., 2014; Deghelt et al., 2014; Fiebig et al., 2006; Frage et al.,

2016; Rasmussen et al., 2007; Srivastava, 2006; Stokke et al., 2011) under various

modalities. In the Vibrionaceae, the replication of a short region of the chromosome

licenses the SER duplication (Baek and Chattoraj, 2014; Kemter et al., 2018), and the

advancement of the SER replication and segregation triggers the divisome assembly

(Galli et al., 2016). In turn, the altering of the chromosome replication does not affect

the replication initiation control of the SER in α-proteobacterium *Ensifer/Sinorhizobium*

*meliloti* (Frage et al., 2016).

Beside the exploration of the replication/segregation mechanistic, studies of multipartite

genomes, targeting a single bacterial species or genus (diCenzo et al., 2013, 2014;

Dubarry et al., 2006; Mackenzie et al., 2004; Slater et al., 2009) or using a more

extensive set of taxa (diCenzo and Finan, 2017; Harrison et al., 2010), relied on

inadequate (replicon size, nucleotide composition, coding of core essential genes for

growth and survival (diCenzo and Finan, 2017; Harrison et al., 2010; Liu et al., 2015);

Figure 1) and/or oriented (presence of plasmid-type systems for genome maintenance

and replication initiation (Harrison et al., 2010)) criteria to characterize the SERs.

**Figure 1. Structural features of the replicons**

Boxplots of the lengths (base pairs) and numbers of genes (ORFs), protein-coding genes (CDS), pseudogenes,
ribosomal RNA genes and transfer RNA genes for the 2016 chromosomes (blue), 129 SERs (orange), and 2783
plasmids (green) included in the final dataset (4928 replicons).

While clarifying the functional and evolutionary contributions of each type of replicon

to a multipartite genome in given bacterial lineages (Galardini et al., 2013; Harrison et

al., 2010; MacLellan et al., 2004; Slater et al., 2009), these studies produced no absolute

definition of SERs (diCenzo and Finan, 2017; Harrison et al., 2010) or universal model

for their emergence (diCenzo and Finan, 2017; diCenzo et al., 2013, 2014; Galardini et

al., 2013; Harrison et al., 2010). We thus set out investigating the nature(s) and origin(s)

of these replicons using as few assumptions as possible.

**RESULTS**

**Replicon inheritance systems as diagnostic features**

We did not limit our study to a particular multipartite genome or a unique gene family.

Rather, we performed a global analysis encompassing all bacterial replicons whose

complete sequence was available in public sequence databases (Figure 2). We reasoned

that the key property discriminating the chromosomes from the plasmids is their

transmission from mother to daughter cells during the bacterial cell cycle. The functions

involved in the replication, partition and maintenance of a replicon, *i.e.*, its inheritance

systems (ISs), thence are expected to reflect the replicon degree of integration into the

host cycle.

We first faced the challenge of identifying all IS functional homologues. The inheritance

of genetic information requires functionally diverse and heterogeneous actuators

depending on the replicon type and the characteristics of the organism. Also, selecting

sequence orthologues whilst avoiding false positives (*e.g.*, sequence paralogues) can be

tricky since remote sequence homology most likely prevails among

chromosome/plasmid protein-homologue pairs.

**Figure 2. Analytical procedure**

110    Starting from an initial dataset of 5125 replicons, we identified 358,624 putative IS

111    functional homologues, overall corresponding to 1711 Pfam functional domains (Figure

112    3a), using a query set of 47,604 chromosomal and plasmidic IS-related proteins selected

113    from the KEGG and ACLAME databases (Tables 1 and 2).

114    **Table 1. ACLAME families used in the building of the query set**

| PROCESS | FAMILY | PROTEIN DESCRIPTION |
|---|---|---|
| | 32 | RepB, pi, initiator protein, RepE, RepA |
| | 76 | Rep, RepB, Rep of rolling circle initiator, RepA, RepU, OrfB, Rep2 |
| | 107 | RepC, RepCa1, RepCa2, RepCd |
| | 114 | Helicase, UrvD rep helicase, helicase super family 1, Yga2F, helicase II |
| | 118 | CdsE, CdsJ |
| | 133 | RepA, W0005, RepA1/A2 |
| | 171 | RepA, RepB, putative theta replicative protein |
| | 207 | replicative DNA helicase, DnaB, pGP1 |
| | 208 | RepA, W0013, W0041, RepFIB |
| | 224 | long form TrfA, TrfA1, TrfA2, S-TrfA, plasmid initiation protein |
| | 237 | RepA, putative RepA, truncated RepA |
| | 244 | RepA, RepB, CopB, repA1/A2, w0004 |
| | 294 | Rop regulatory protein, RNAI modulator, RNA modulator, plasmid copy number control |
| | 297 | primase activity/DNA initiation, LtrC/LtrC-like hypothetical protein, PcfD |
| | 330 | DNA repair/ DNA helicase, type III restriction enzyme, res subunit, DEAD/DEAH box helicase |
| | 377 | replicase, replication initiation, RepC, RepJ, RepE, RepL |
| | 383 | RepA, Rb100 |
| | 404 | RepA,RepB,RepW |
| Replication | 412 | Rep, RepA |
| | 423 | truncated RCR replication, RepRC, RepB, OrfA |
| | 426 | cell division control protein 6 homolog |
| | 440 | Rep 14-4, rm protein, RepA hypothetical protein |
| | 451 | RepA, host type : *Corynebacterium* |
| | 477 | Rep, RepS, RepE, host type : *Bacillus*, RepS, RepR |
| | 612 | RepL, replication initiation |
| | 775 | DNA helicase activity, RepA, putative helicase |
| | 854 | DNA helicase activity, RepC, putative initiator protein |
| | 921 | RepA |
| | 931 | DNA replication initiation, putative protein, CdsD |
| | 1005 | helicase activity, putative protein, hypothetical helicase |
| | 1055 | RNA polymerase σ factor, σ 70 family, bacteriocin uviA, sigF/V/G, tetR, host type : *Clostridium* |
| | 1095 | DNA repair/helicase, RuvB, DNA pol III γ and τ subunits, DNA pol δ subunit |
| | 1099 | putative theta replicase, RepB, Rep2 |
| | 1187 | DNA replication, RepH, RepI |
| | 1288 | RepA |
| | 1345 | DNA primase activity, DNA primase , primase CHC2 family |
| | 1398 | helicase activity, GcrE, GcrC |

| | | |
|---|---|---|
| | *1652* | DNA repair/exonuclease activity, DNA exonuclease protein, SbcCD related protein |
| | *1837* | putative replication protein |
| | *2881* | RepC-like, Pif |
| Partition | *4* | plasmid partition protein, ParA, ParA IncC protein, ParA InC1/ IncC2, SopA, virC1 |
| | *14* | RepB, RepB partitionning, KorB repressor and partitionning, ParB-like domain, YefA, YdeB, ParB, ParB-like |
| | *102* | DNA binding, partitionning protein, control protein, ParB, VirB, partition protein B |
| | *128* | DNA segregation/DNA translocase activity, cell division FtsK/ SpoIIIE, SpoI, TraB |
| | *289* | ParM family, go : translocase, hypothetical protein, rode shape protein, putative ATPase of class HSP70 |
| | *316* | microfilament motor activity, ParM family, StbA protein, stable inheritance protein, ParA |
| | *318* | ATPase, regulation of cell division, chromosome patition, GumC, ExoP related protein, EpsB, MPA1 family |
| | *427* | ATPase family, ParR family, ParB, StbB, mediator of plasmid stability |
| | *875* | DNA binding, partitionning protein family ParB/Spo0J, YPMT1.28c |
| | *876* | DNA binding, partitionning protein family ParB/Spo0J, YPMT1.29c |
| | *983* | DNA binding, ParB, CopG |
| | *1227* | DNA plasmid copy number control, CopG |
| | *2158* | RepC |
| | *2894* | DNA binding |
| Dimer resolution | *5* | serine based recombinase activity, ylb, resolvase, second invertase, TniR, ParA |
| | *10* | tyrosine-based recombinase, integrase, putative integrase, Xer, recombinase-like SAM |
| | *101* | plasmid dimer resolution, tyrosine-based recombinase, yld, SAM-like protein |
| | *170* | tyrosine-based recombinase, OrfA, hypothetical protein |
| | *589* | tyrosine based protein, Fis protein |
| | *688* | tyrosine based protein, SAM like protein, XerD |
| Maintenance | *100* | Postsegregational killing system vapBC/vag |
| | *136* | Postsegregational killing system parDE |
| | *156* | Postsegregational killing system epsilon-zeta |
| | *201* | Postsegregational killing system higBA |
| | *212* | Postsegregational killing system parDE |
| | *293* | Postsegregational killing system mazEF |
| | *319* | Postsegregational killing system relBE |
| | *326* | Postsegregational killing system mazEF |
| | *335* | Postsegregational killing system HOK/SOK |
| | *338* | Postsegregational killing system parDE |
| | *356* | Postsegregational killing system parDE |
| | *366* | Postsegregational killing system vapBC/vag |
| | *380* | Postsegregational killing system phD-doc |
| | *428* | Postsegregational killing system ccd |
| | *470* | Postsegregational killing system yacA |
| | *474* | Postsegregational killing system relBE |
| | *515* | Postsegregational killing system relBE |
| | *556* | Postsegregational killing system higBA |
| | *563* | Postsegregational killing system ccd |
| | *588* | Postsegregational killing system higBA |
| | *677* | Postsegregational killing system higBA |
| | *798* | Postsegregational killing system mazEF |
| | *916* | Postsegregational killing system relBE |
| | *1031* | Postsegregational killing system HOK/SOK |
| | *1180* | Postsegregational killing system vapXD |
| | *1308* | Postsegregational killing system HicAB |
| | *1559* | Postsegregational killing system epsilon-zeta |
| | *1927* | Postsegregational killing system mazEF |

| | | | |
|---|---|---|---|
| *3357* | Postsegregational killing system, plasmid maintenance | | |
| *4776* | Postsegregational killing system, parC | | |
| *4777* | Postsegregational killing system parDE, parD | | |
| *16584* | Postsegregational killing system vapXD | | |

**115  Table 2.  KEGG "Prokaryotic-type chromosome" orthology groups used in the building of the**
**116        query set**

| BRITE HIERARCHY | | KEGG ENTRY | NAME | DEFINITION |
|---|---|---|---|---|
| Chromosome replication | Initiation factors (bacterial) | *K02313* | DnaA | chromosomal replication initiator protein |
| | | *K02314* | DnaB | replicative DNA helicase [EC:3.6.4.12] |
| | | *K03346* | DnaB2, DnaB | replication initiation and membrane attachment protein |
| | | *K02315* | DnaC | DNA replication factor, helicase loader |
| | | *K02316* | DnaG | DNA primase [EC:2.7.7.-] |
| | | *K11144* | DnaI | primosomal protein DnaI |
| | | *K05787* | HupA | DNA-binding protein HU-alpha |
| | | *K03530* | hupB | DNA-binding protein HU-beta |
| | | *K04764* | IhfA, HimA | integration host factor subunit alpha |
| | | *K05788* | IhfB, HimD | integration host factor subunit beta |
| | | *K03111* | ssb | single-strand DNA-binding protein |
| | Terminus site-binding protein | *K10748* | Tus, Tau | DNA replication terminus site-binding protein |
| | DNA methylation enzyme | *K06223* | Dam | DNA adenine methylase [EC:2.1.1.72] |
| | Prevention of re-replication factors | *K10763* | Hda | DnaA-homolog protein |
| | | *K03645* | SeqA | negative modulator of initiation of replication |
| Chromosome partition | MukBEF complex | *K03632* | MukB | chromosome partition protein MukB |
| | | *K03804* | MukE | chromosome partition protein MukE |
| | | *K03633* | MukF | chromosome partition protein MukF |
| | Condensin-like complex | *K03529* | Smc | chromosome segregation protein |
| | | *K05896* | ScpA | segregation and condensation protein A |
| | | *K06024* | ScpB | segregation and condensation protein B |
| | Divisome proteins | *K03585* | AcrA | membrane fusion protein |
| | | *K01448* | AmiA,AmiB, AmiC | N-acetylmuramoyl-L-alanine amidase [EC:3.5.1.28] |
| | | *K13052* | DivIC, DivA | cell division protein DivIC |
| | | *K03590* | FtsA | cell division protein FtsA |
| | | *K05589* | FtsB | cell division protein FtsB |
| | | *K09812* | FtsE | cell division transport system ATP-binding protein |
| | | *K03587* | FtsI | cell division protein FtsI [EC:2.4.1.129] |
| | | *K03466* | FtsK, SpoIIIE | DNA segregation ATPase FtsK/SpoIIIE, S-DNA-T family |
| | | *K03586* | FtsL | cell division protein FtsL |
| | | *K03591* | FtsN | cell division protein FtsN |
| | | *K03589* | FtsQ | cell division protein FtsQ |
| | | *K03588* | FtsW, SpoVE | cell division protein FtsW |
| | | *K09811* | FtsX | cell division transport system permease protein |
| | | *K03531* | FtsZ | cell division protein FtsZ |
| | | *K09888* | ZapA | cell division protein ZapA |
| | | *K03528* | ZipA | cell division protein ZipA |
| | Inhibitors of FtsZ assembly | *K04074* | DivIVA | cell division initiation protein |
| | | *K06286* | EzrA | septation ring formation regulator |

| | | | |
|---|---|---|---|
| | | K03610 | MinC | septum site-determining protein MinC |
| | | K03609 | MinD | septum site-determining protein MinD |
| | | K03608 | MinE | cell division topological specificity factor |
| | | K05501 | SlmA, Ttk | TetR/AcrR family transcriptional regulator |
| | | K09772 | SepF | cell division inhibitor SepF |
| | | K13053 | SulA | cell division inhibitor, FtsZ assembly inhibitor |
| | Other chromosome partitioning proteins | K04095 | Fic | cell filamentation protein |
| | | K04094 | Gid, TrmFO | methylenetetrahydrofolate--tRNA-[uracil-5-)-methyltransferase [EC:2.1.1.74] |
| | | K03495 | GidA, MnmG, MTO1 | tRNA uridine 5-carboxymethylaminomethyl modification enzyme |
| | | K03501 | GidB, RsmG | 16S rRNA [guanine527-N7)-methyltransferase [EC:2.1.1.170] |
| | | K03569 | MreB | rod shape-determining protein MreB and related proteins |
| | | K03570 | MreC | rod shape-determining protein MreC |
| | | K03571 | MreD | rod shape-determining protein MreD |
| | | K03593 | Mrp | ATP-binding protein involved in chromosome partitioning |
| | | K03496 | ParA, Soj | chromosome partitioning protein |
| | | K03497 | ParB, Spo0J | chromosome partitioning protein, ParB family |
| | | K02621 | ParC | topoisomerase IV subunit A [EC:5.99.1.-] |
| | | K02622 | ParE | topoisomerase IV subunit B [EC:5.99.1.-] |
| | | K11686 | RacA | chromosome-anchoring protein RacA |
| | | K05837 | RodA, MrdB | rod shape determining protein RodA |
| | | K03645 | SeqA | negative modulator of initiation of replication |
| | | K03733 | XerC | integrase/recombinase XerC |
| | | K04763 | XerD | integrase/recombinase XerD |
| Nucleoid | HNS (histone-like nucleoid structuring protein) | K03746 | H-NS | DNA-binding protein H-NS |
| | | K11685 | StpA | DNA-binding protein StpA |
| | HU (heat unstable protein) | K05787 | HupA | DNA-binding protein HU-alpha |
| | | K03530 | HupB | DNA-binding protein HU-beta |
| | IHF (integration host factor) | K04764 | IhfA, HimA | integration host factor subunit alpha |
| | | K05788 | IhfB, HimD | integration host factor subunit beta |
| | Other nucleoid associated proteins | K05516 | CbpA | curved DNA-binding protein |
| | | K12961 | DiaA | chromosomal replication initiator protein |
| | | K02313 | DnaA | DnaA initiator-associating protein |
| | | K04047 | Dps | starvation-inducible DNA-binding protein |
| | | K03557 | Fis | Fis family transcriptional regulator, factor for inversion stimulation protein |
| | | K03666 | Hfq | host factor-I protein |
| | | K05596 | IciA | chromosome initiation inhibitor, LysR family transcriptional regulator |
| | | K03719 | Lrp | leucine-responsive regulatory protein, Lrp/AsnC family transcriptional regulator |
| | | K05804 | Rob | right origin-binding protein, AraC family transcriptional regulator |

**Figure 3. Properties of the IS clustering**

(a) Frequency distribution of the 358,624 putative IS protein homologues according to their number of functional domains (0 to
69) per protein (left), and occurrences of the 1711 functional Pfam domains (right). The 20 top most frequently encountered
functional domains are indicated. (b) Size distribution of the 7013 clusters, each comprising from a single to 1990 proteins. (c)
Percentage distribution of the most frequent annotation per cluster among all clusters (left) and among clusters with multiple
annotations (right). (d) Distribution of the most frequent annotation per cluster among the 917 excluded clusters (left) and the
6096 clusters retained for the analysis (right).

We then inferred 7013 homology groups using a clustering procedure and named the

clusters after the most frequent annotation found among their proteins (Figure 3b,c). Most

clusters were characterized by a single annotation whilst the remaining few (4.7%) each

harbored from 2 to 710 annotations, the most frequent annotation in a cluster generally

representing more than half of all annotations (Figure 3c). The removal of false positives

left 267,497 IS protein homologues distributed in 6096 clusters (Figure 3d) and coded by

4928 replicons out of the initial replicon dataset. Following the Genbank/RefSeq

annotations, our final dataset comprised 2016 complete genome sets corresponding to

3592 replicons (2016 chromosomes, 129 SERs, and 1447 plasmids) and 1336 plasmid

genomes (Supplementary table 1), irregularly distributed across the bacterial phylogeny

(Figure 4a). Multi-ER genomes are observed in 5.0% of all represented bacterial genera

and constitute 5.7% of the complete genomes (averaged over genera) available at the time

of study (Figure 4b). They are merely incidental (0.2% in Firmicutes) or reach up to

almost one third of the genomes (30.1% in β-Proteobacteria) depending on the lineage,

and are yet to be observed in most bacterial phyla, possibly because of the poor

representation of some lineages. Although found in ten phyla, they occur more than once

per genus in only three of them: Bacteroidetes, Proteobacteria and Spirochaetae.

**Figure 4.  Taxonomic structure of the replicon dataset**

Numbers of replicons (a) and complete genomes (b), and represented bacterial genera are shown according to datasets and host taxonomy. Surfaces represent the numbers of bacterial genera, replicons or genomes within each category. Percentages of multipartite genomes and corresponding bacterial genera are calculated for each host phylum or class (Proteobacteria).

**Exploration of the replicon diversity**

We explored the differences and similarities of the bacterial replicons with regard to

their IS usage using a data mining and machine learning approach (Methods). The 6096

retained IS clusters were used as distinct variables to ascribe each of the 4928 replicons

with a vector according to its IS usage profile. We transformed these data into bipartite

graphs depending on the number of proteins from the IS clusters coded by each replicon.

Bipartite graphs display both the vectors (replicons) and the variables (protein clusters)

together with their respective connections, and allow the interactive exploration of the

data. The majority of the replicons are interconnected (Figure 5) as testimony of the

shared evolutionary history of their IS sequences. Chromosomes and plasmids form

overall distinct groups and communities with varying degree of connectivity depending

on their functional specificities (Figure 5a) as well as on the bacterial taxonomy of their

hosts (Figure 5c). They nonetheless share many ISs, bearing witness to the continuity of

the genomic material and the extensive exchange of genetic material within bacterial

genomes. The occurrence of poorly IS cluster-connected plasmids within a group of

chromosomes did not consistently reflect a true relationship and rather resulted from

shared connections to a very small number (as low as one) of common ISs. While being

interconnected to both chromosomes and plasmids *via* numerous IS clusters, the SERs

generally stand apart from either these types of replicons and gather at the chromosome-

plasmid interface (Figure 5a,b). Their IS usage is neither chromosome-like nor plasmid-

like, suggesting that they may constitute a separate category of replicons. This is most

tangible in the case of the proteobacterial lineages where SERs occur most frequently

(top of Figure 5b).

**Figure 5.    Visualisation of the replicon IS-based relationshipss**

Gephi-generated bipartite-graphs for the whole dataset (a and b) or groups of replicons following the host taxonomy (c). Nodes correspond to the replicons (large dots) or the clusters of IS proteins (small dots). Edges linking replicons and protein clusters reflect the presence on a replicon of at least one protein of a protein cluster. Colouring according to replicon type (a and c): chromosomes (white), plasmids (grey), and SERs (blue), or according to host taxonomy (b).

All SERs in the β- and γ-Proteobacteria, and most in the α-Proteobacteria are linked to

remarkable chromosome-type IS clusters, such as AcrA, IciA, FtsE, HN-S and Lrp, as

well as to plasmid-like ParA/ParB, Rep and PSK IS clusters. A similar pattern is

observed for the SERs in actinobacterium *Nocardiopsis dassonvillei,* firmicute

*Butyrivibrio proteoclasticus*, and chloroflexi *Sphaerobacter thermophilus* and

*Thermobaculum terrenum* (Figure 5b). Interestingly, DNA primase DnaG-annotated

clusters connect the SERs present in all but one *Burkholderia* species (β-Proteobacteria)

as well as the chromosomes of all other bacteria. Since the sole exception,

*B. rhizoxinica*, possesses a SER-less reduced genome as an adaption to intracellular life,

the *Burkholderia* SERs likely originated from a single event prior to the diversification

of the genus, possibly in relation to the speciation event that gave rise to this lineage.

The second SERs harbored by only some *Burkholderia* species exhibit a higher level of

interconnection to plasmids, as do the SERs in α-proteobacterium *Sphingobium*,

cyanobacterium *Cyanothece* sp. ATCC 51142, *Deinococcus radiodurans* (Deinococcus-

Thermus) and fusobacterium *Ilyobacter polytropus*. This points to an incomplete

stabilization of the SERs into the genome that may reflect a recent, ongoing, event of

integration and/or differing selective pressures at play depending on the bacterial

lineages. At odds with these observations, some SERs group unambiguously with

chromosomes. The SERs in α-Proteobacteria *Asticcacaulis excentricus* and *Paracoccus*

*denitrificans*, Bacteroidetes *Prevotella intermedia* and *P. melaninogenica*,

acidobacterium *Chloracidobacterium thermophilum*, and cyanobacterium

*Anabaena* sp. 90 bear higher levels of interconnection to chromosomes than to plasmids

or other SERs. Indeed, the SERs in *Prevotella* spp. are hardly linked to plasmids, and

the few plasmid-like IS proteins that *C. thermophilum* SER codes (mostly Rep, Helicase

and PSK), albeit found in plasmids occurring in other phyla, are observed in none of the

Acidobacteria plasmids. An extreme situation is met in *Leptospira* spp. (Spirochaetae)

whose SERs are each linked to only three or four (out of a total of six) chromosome-like

IS clusters, always including ParA and ParB. Interestingly, the ParA cluster appears to

be specific to Spirochaetae chromosomes with the notable exception of one plasmid

found in Leptospiraceae *Turneriella parva*.

**IS-based relationships of the replicons**

We submitted the bipartite graph of the whole dataset to a community structure

detection algorithm (INFOMAP) that performs a random walk along the edges

connecting the graph vertices. We expected the replicon communities to be trapped in

high-density regions of the graph. We also performed a dimension reduction by

Principal Component Analysis followed by a hierarchical clustering procedure

(WARD). The clustering solutions (Supplementary tables 2 and 3) were meaningful

(high values reached by the stability criterion scores), and biologically relevant (efficient

separation of the chromosomes from the plasmids; high *homogeneity* values) using

either method (Table 3). In another experiment, we considered each genus as a unique

sample and averaged the variables over the replicons of the different species for each

replicon type. The aim was to control for the disparity in taxon representation of the

replicons. This dataset produced overall similar albeit slightly less stable clusters (lower

*homogeneity* values). Taxonomically homogeneous clusters of chromosomes were best

retrieved using the coupling of dimension reduction and hierarchical clustering with a

large enough number of clusters (*homogeneity* scores up to 0.93). In turn, the

community detection algorithm was more efficient in recovering the underlying

taxonomy of replicons (higher value of *completeness*), and was sole able to identify

small and scattered plasmid clusters (Supplementary tables 2 and 3).

**Table 3. Evaluation of the replicon IS-based clusterings**

| | | INDEX[a] USING IS PROTEIN SEQUENCES | | | | USING IS FUNCTIONS | |
|---|---|---|---|---|---|---|---|
| | CLUSTERING | INFOMAP | | PCA+ WARD[b] | | PCA+ WARD[b] | |
| **PROCEDURE** | Dataset[c] | $V^R$ | $\bar{V}^R_{genus}$ | $V^R$ | $\bar{V}^R_{genus}$ | $V^R_f$ | $\bar{V}^R_{f,genus}$ |
| | Parameters | 500 iterations | | $\begin{cases}k=200\\pc=30\end{cases}$ | $\begin{cases}k=200\\pc=30\end{cases}$ | $\begin{cases}k=50\\pc=4\end{cases}$ | $\begin{cases}k=20\\pc=4\end{cases}$ |
| | Number of clusters | 223 | 77 | 175 | 75 | 49 | 19 |
| | PCA explained variance | | | 57% | 58% | 87% | 85% |
| | Stability criterion ($\Delta^{Kl}$)[d] | 0.82 | 0.76 | 0.85 | 0.74 | 0.80 | 0.71 |
| **EVALUATED SEPARATION** | Chromosomes *vs.* Plasmids — *homogeneity* | 0.82 | 0.63 | 0.93 | 0.83 | 0.85 | 0.68 |
| | *completeness* | 0.15 | 0.15 | 0.25 | 0.20 | 0.30 | 0.23 |
| | *V-measure* | 0.25 | 0.24 | 0.43 | 0.32 | 0.44 | 0.35 |
| | Chromosomes *per* host phylum — *homogeneity* | 0.93 | 0.69 | 0.93 | 0.80 | 0.50 | 0.44 |
| | *completeness* | 0.60 | 0.61 | 0.35 | 0.40 | 0.27 | 0.33 |
| | *V-measure* | 0.73 | 0.65 | 0.51 | 0.53 | 0.35 | 0.38 |
| | Chromosomes *per* host class — *homogeneity* | 0.85 | 0.64 | 0.93 | 0.80 | 0.47 | 0.37 |
| | *completeness* | 0.80 | 0.82 | 0.16 | 0.58 | 0.36 | 0.41 |
| | *V-measure* | 0.82 | 0.72 | 0.66 | 0.67 | 0.41 | 0.39 |
| | Plasmids *per* host phylum — *homogeneity* | 0.88 | 0.78 | 0.06 | 0.01 | 0.02 | 0.02 |
| | *completeness* | 0.33 | 0.35 | 0.16 | 0.14 | 0.10 | 0.30 |
| | *V-measure* | 0.48 | 0.48 | 0.08 | 0.02 | 0.03 | 0.03 |
| | Plasmids *per* host class — *homogeneity* | 0.84 | 0.74 | 0.07 | 0.02 | 0.03 | 0.02 |
| | *completeness* | 0.43 | 0.51 | 0.28 | 0.36 | 0.25 | 0.28 |
| | *V-measure* | 0.57 | 0.60 | 0.12 | 0.03 | 0.05 | 0.03 |

[a] *V-measure* according to Rosenberg and Hirschberg (2007)

[b] *k*: number of input clusters; *pc*: principal components used in WARD

[c] $V^R_f$: Ensemble of all IS function-based replicon vectors ($v^r_f$); $\bar{V}^R_{f,genus}$: Ensemble of IS function-based genus-

normalized replicon vectors ($v^r_{f,genus}$)

[d] Stability criterion according to Hennig (2007)

The plasmid clusters obtained using PCA+WARD lacked taxonomical patterning and,

although highly stable, only reflected the small Euclidian distances existing among the

plasmid replicons (*e.g.*, one cluster of 2656 plasmids had a stability score of 0.975). The

clusters obtained with INFOMAP mirrored the taxonomical structure of the data,

suggesting that the taxonomic signal, expected to be associated to the chromosomes, is

preserved among the IS protein families functionally specifying the plasmids. The

presence of a majority of the SERs amongst the chromosome clusters generated by

INFOMAP confirmed the affinities between these two genomic elements and the clear

individuation of the SERs from the plasmids. However, the larger number of

chromosomal ISs often caused the PCA+WARD approach to place SERs into plasmid

clusters. The SERs in *Butyrivibrio*, *Deinococcus*, *Leptospira* and *Rhodobacter* spp.

grouped consistently with plasmids while the SERs in *Vibrionaceae* and *Brucellaceae*

formed specific clusters (Table 4). Burkholderiales and *Agrobacterium* SERs, whose

homogenous clusters tended to be unstable, exhibited a higher affinity to plasmids

overall. The SERs of *Asticcaulis*, *Paracoccus* and *Prevotella* spp. associated stably

with chromosomes using the two clustering methods (Table 4a,b) and possess IS profiles

that set them apart from both the plasmids and the other SERs.

**Table 4. IS protein cluster-based unsupervised classification of SERs**

a. INFOMAP clustering solution

| Bacterial genus | $C$ [a] | CHR% | SER% | PLD% | $wBHI$ [b] | $\overline{\Delta^c}$ [c] | $\overline{\Delta^r}$ [d] |
|---|---|---|---|---|---|---|---|
| *Agrobacterium* | 3 | 38 | 35 | 27 | 0.90 | 0.47 | 0.61 |
| *Aliivibrio* | 1 | 0 | 100 | 0 | 1.00 | 0.95 | 1.00 |
| *Anabaena* | 1 | 98 | 1 | 1 | 1.00 | 0.90 | 1.00 |
| *Asticcacaulis* | 1 | 96 | 1 | 3 | 1.00 | 0.97 | 1.00 |
| *Brucella* | 1 | 0 | 95 | 5 | 1.00 | 0.87 | 1.00 |
| *Burkholderia* | 2 | 64 | 17 | 19 | 0.99 | 0.77 | 0.99 |
| *Butyrivibrio* | 1 | 0 | 50 | 50 | 1.00 | 0.83 | 1.00 |
| *Chloracidobacterium* | 1 | 91 | <1 | 9 | 0.82 | 0.86 | 0.00 |
| *Cupriavidus* | 1 | 73 | 18 | 9 | 0.99 | 0.72 | 1.00 |
| *Cyanothece* | 1 | 0 | 6 | 94 | 0.89 | 0.61 | 0.33 |
| *Deinococcus* | 1 | 0 | 4 | 96 | 0.71 | 0.61 | 1.00 |
| *Ilyobacter* | 1 | 91 | <1 | 9 | 0.82 | 0.86 | 0.25 |
| *Leptospira* | 1 | 0 | 88 | 12 | 1.00 | 1.00 | 1.00 |

| Bacterial genus | | | | | | | |
|---|---|---|---|---|---|---|---|
| *Nocardiopsis* | 1 | 91 | <1 | 9 | 0.97 | 0.97 | 1.00 |
| *Ochrobactrum* | 1 | 0 | 95 | 5 | 1.00 | 0.87 | *n.a.n.*[e] |
| *Paracoccus* | 1 | 96 | 1 | 3 | 1.00 | 0.97 | 1.00 |
| *Photobacterium* | 1 | 96 | 1 | 3 | 0.99 | 0.79 | 1.00 |
| *Prevotella* | 1 | 96 | 2 | 2 | 0.95 | 0.92 | 1.00 |
| *Pseudoalteromonas* | 1 | 96 | 1 | 3 | 0.99 | 0.79 | 0.56 |
| *Ralstonia* | 1 | 73 | 18 | 9 | 0.99 | 0.72 | 1.00 |
| *Rhodobacter* | 1 | 0 | 40 | 60 | 1.00 | 0.71 | 1.00 |
| *Ensifer* (*Sinorhizobium*) | 2 | 0 | 2 | 98 | 0.96 | 0.65 | 0.67 |
| *Sphaerobacter* | 1 | 0 | 50 | 50 | 1.00 | 1.00 | 1.00 |
| *Sphingobium* | 2 | 77 | 1 | 22 | 0.95 | 0.90 | 0.50 |
| *Thermobaculum* | 1 | 91 | <1 | 9 | 0.82 | 0.86 | 1.00 |
| *Variovorax* | 1 | 73 | 18 | 9 | 0.99 | 0.72 | 0.90 |
| *Vibrio* | 1 | 0 | 100 | 0 | 1.00 | 0.95 | 0.89 |

[a] number of clusters containing SERs of a given bacterial genus
[b] weighted biological homogeneity index value for the phylum of the replicons in the clusters
[c] mean value of the cluster stability estimator, weighted by the cluster sizes
[d] mean value of the SER stability estimator for a given bacterial genus
[e] "*n.a.n.*", standing for "not a number", indicates that the replicon appeared in none of the bootstrap replications
performed in the clustering procedure

b. PCA+WARD clustering solution

| Bacterial genus | $C^a$ | CHR% | SER% | PLD% | $wBHI^b$ | $\overline{\Delta^c}$ [c] | $\overline{\Delta^r}$ [d] |
|---|---|---|---|---|---|---|---|
| *Agrobacterium* | 2 | 0 | 29 | 71 | 0.94 | 0.76 | 1.00 |
| *Aliivibrio* | 2 | 0 | 56 | 44 | 1.00 | 0.60 | 0.33 |
| *Anabaena* | 1 | 98 | 2 | 0 | 0.97 | 0.84 | 0.00 |
| *Asticcacaulis* | 1 | 88 | 8 | 4 | 1.00 | 0.88 | 1.00 |
| *Brucella* | 2 | 0 | 33 | 67 | 0.96 | 0.53 | 0.97 |
| *Burkholderia* | 7 | 0 | 79 | 21 | 0.97 | 0.69 | 0.84 |
| *Butyrivibrio* | 1 | <1 | 1 | 99 | 0.27 | 0.98 | 1.00 |
| *Chloracidobacterium* | 1 | <1 | 1 | 99 | 0.27 | 0.98 | 1.00 |
| *Cupriavidus* | 2 | 0 | 92 | 8 | 1.00 | 0.69 | 0.92 |
| *Cyanothece* | 1 | <1 | 1 | 99 | 0.27 | 0.98 | 1.00 |
| *Deinococcus* | 1 | <1 | 1 | 99 | 0.27 | 0.98 | 1.00 |
| *Ilyobacter* | 1 | <1 | 1 | 99 | 0.27 | 0.98 | 1.00 |
| *Leptospira* | 1 | <1 | 1 | 99 | 0.27 | 0.98 | 1.00 |
| *Nocardiopsis* | 1 | 0 | 2 | 98 | 0.58 | 0.40 | 1.00 |
| *Ochrobactrum* | 1 | 0 | 100 | 0 | 1.00 | 1.00 | 1.00 |
| *Paracoccus* | 1 | 88 | 8 | 4 | 1.00 | 0.88 | 1.00 |
| *Photobacterium* | 1 | 0 | 100 | 0 | 1.00 | 0.55 | 1.00 |
| *Prevotella* | 2 | 95 | 5 | 0 | 1.00 | 0.73 | 0.50 |
| *Pseudoalteromonas* | 2 | <1 | 1 | 99 | 0.28 | 0.82 | 0.83 |
| *Ralstonia* | 1 | 0 | 68 | 32 | 1.00 | 0.81 | 0.83 |
| *Rhodobacter* | 2 | 0 | 6 | 94 | 0.65 | 0.43 | 0.58 |
| *Ensifer* (*Sinorhizobium*) | 2 | 0 | 21 | 79 | 0.94 | 0.46 | 0.25 |
| *Sphaerobacter* | 1 | 0 | 20 | 80 | 0.93 | 0.52 | 0.50 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| *Sphingobium* | 1 | 0 | 39 | 61 | 1.00 | 0.66 | 0.83 |
| *Thermobaculum* | 1 | <1 | 1 | 99 | 0.27 | 0.98 | 1.00 |
| *Variovorax* | 1 | 0 | 67 | 33 | 1.00 | 0.48 | 0.00 |
| *Vibrio* | 2 | 0 | 56 | 44 | 1.00 | 0.60 | 0.79 |

[a] number of clusters containing SERs of a given bacterial genus
[b] weighted biological homogeneity index value for the phylum of the replicons in the clusters
[c] mean value of the cluster stability estimator, weighted by the cluster sizes
[d] mean value of the SER stability estimator for a given bacterial genus

We reached similar conclusions when performing a PCA+WARD clustering using the

117 functional annotations of the IS protein clusters rather than the IS clusters

themselves (Tables 3 and 5; Supplementary table 4).

**Table 5. Function-based unsupervised classification of SERs using PCA+WARD**

| Bacterial genus | $C^a$ | CHR% | SER% | PLD% | $wBHI^b$ | $\overline{\Delta^c}{}^c$ | $\overline{\Delta^r}{}^d$ |
|---|---|---|---|---|---|---|---|
| *Agrobacterium* | 3 | 64 | 21 | 15 | 0.86 | 0.60 | 0.81 |
| *Aliivibrio* | 1 | 0 | 70 | 30 | 1.00 | 0.70 | 0.67 |
| *Anabaena* | 1 | 77 | 4 | 19 | 0.21 | 0.64 | 1.00 |
| *Asticcacaulis* | 1 | 99 | 1 | 0 | 0.51 | 0.60 | 1.00 |
| *Brucella* | 1 | 43 | 32 | 25 | 0.75 | 0.80 | 1.00 |
| *Burkholderia* | 6 | 31 | 42 | 27 | 0.92 | 0.68 | 0.81 |
| *Butyrivibrio* | 1 | 77 | 4 | 19 | 0.21 | 0.64 | 1.00 |
| *Chloracidobacterium* | 1 | 1 | <1 | 99 | 0.29 | 0.98 | 0.00 |
| *Cupriavidus* | 1 | 5 | 95 | 0 | 1.00 | 0.66 | 0.66 |
| *Cyanothece* | 1 | 1 | <1 | 99 | 0.29 | 0.98 | 1.00 |
| *Deinococcus* | 1 | 1 | <1 | 99 | 0.29 | 0.98 | 1.00 |
| *Ilyobacter* | 1 | 77 | 4 | 19 | 0.21 | 0.64 | 1.00 |
| *Leptospira* | 1 | 1 | <1 | 99 | 0.29 | 0.98 | 1.00 |
| *Nocardiopsis* | 1 | 77 | 4 | 19 | 0.21 | 0.64 | 1.00 |
| *Ochrobactrum* | 1 | 90 | 8 | 2 | 1.00 | 0.40 | 0.73 |
| *Paracoccus* | 1 | 92 | 5 | 3 | 0.89 | 0.32 | 0.53 |
| *Photobacterium* | 1 | 0 | 70 | 30 | 1.00 | 0.70 | 0.36 |
| *Prevotella* | 2 | 86 | 3 | 11 | 0.34 | 0.62 | 0.50 |
| *Pseudoalteromonas* | 1 | 77 | 4 | 19 | 0.21 | 0.64 | 0.29 |
| *Ralstonia* | 1 | 0 | 70 | 30 | 1.00 | 0.70 | 0.22 |
| *Rhodobacter* | 2 | 27 | 32 | 41 | 0.84 | 0.73 | 0.83 |
| *Ensifer* (*Sinorhizobium*) | 2 | 25 | 48 | 27 | 0.86 | 0.76 | 0.63 |
| *Sphaerobacter* | 1 | 100 | <1 | 0 | 0.35 | 0.60 | 1.00 |
| *Sphingobium* | 2 | 62 | 17 | 21 | 0.34 | 0.64 | 0.86 |
| *Thermobaculum* | 1 | 77 | 4 | 19 | 0.21 | 0.64 | 1.00 |
| *Variovorax* | 1 | 0 | 70 | 30 | 1.00 | 0.70 | 1.00 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| *Vibrio* | 4 | 31 | 37 | 32 | 0.97 | 0.57 | 0.92 |

[a] number of clusters containing SERs of a given bacterial genus
[b] weighted biological homogeneity index value for the phylum of the replicons in the clusters
[c] mean value of the cluster stability estimator, weighted by the sizes of the clusters
[d] mean value of the SER stability estimator for a given bacterial genus

Remarkably, in this latter analysis, the chromosomes in the multipartite genomes of

*Prevotella intermedia* and *P. melaninogenica* were more similar to plasmids than to

other groups of chromosomes and to single chromosomes in other *Prevotella* species

(*P. denticola* and *P. ruminicola*).

## SER-specifying IS functions

Next, we searched which of the IS functions are specific to the SERs. We performed

several logistic regression analyses to identify over- or under-represented ISs and to

assess their respective relevance to each class of replicons. Because of their

comparatively small number, all SERs were assembled into a single group despite their

disparity. A hundred and one IS functionalities (96% of KEGG-annotated chromosome-

like functions and 72% of ACLAME-annotated plasmid-like functions) were

significantly enriched in one replicon category over the other (Table 6). The large

majority of the IS functions differentiates the chromosomes from the plasmids. The

latter are only determined by ISs corresponding to ACLAME annotations Rep, Rop and

TrfA, involved in initiation of plasmid replication, and ParA and ParB, dedicated to

plasmid partition. Some KEGG-annotated functions, *e.g.,* DnaA, DnaB or FtsZ, appear

to be more highly specific to chromosomes (higher *OR* values) than others such as

DnaC, FtsE or H-NS (lower *OR* values). Strikingly, very few functions distinguish

significantly the chromosomes from the SERs, by contrast with plasmids.

# Table 6. IS usage comparison between replicon categories

Between classes of replicons logistic regressions for each IS function. Model significance: $0 < P\_value < 0.01$: significant; $0.01 < P\_value < 0.05$: poorly significant; $0.05 < P\_value$: non significant (not shown). Odd-ratio (OR) favouring the first class: $10^0 \leq OR$, or the second class: $OR < 10^0$. IS functions biased to the same order of magnitude in chromosomes and SERs when compared to plasmids are highlighted (blue).

| | IS function | Chromosomes vs. Plasmids | | Chromosomes vs. SERs | | SERs vs. Plasmids | |
|---|---|---|---|---|---|---|---|
| | | P_value | OR | P_value | OR | P_value | OR |
| REPLICATION | CbpA | $8.20 \times 10^{-27}$ | 2,608.4 | $9.90 \times 10^{-13}$ | 22.8 | $5.60 \times 10^{-07}$ | 36.1 |
| | Dam | $6.90 \times 10^{-16}$ | 16.7 | $3.60 \times 10^{-02}$ | 2.0 | $2.40 \times 10^{-02}$ | 4.3 |
| | DiaA | $1.50 \times 10^{-15}$ | 81.9 | $1.20 \times 10^{-03}$ | 38.4 | | |
| | DnaA | $3.00 \times 10^{-44}$ | 2,118.9 | $1.10 \times 10^{-19}$ | 239.6 | $3.50 \times 10^{-03}$ | 8.3 |
| | DnaB | $1.10 \times 10^{-43}$ | 1,992.9 | $5.10 \times 10^{-19}$ | 429.4 | $8.20 \times 10^{-03}$ | 3.7 |
| | DnaB2 | $6.70 \times 10^{-03}$ | 12.6 | | | | |
| | DnaC | $6.00 \times 10^{-12}$ | 2.6 | | | $4.60 \times 10^{-02}$ | 1.5 |
| | DnaG | $2.10 \times 10^{-50}$ | 1,861.5 | $1.90 \times 10^{-21}$ | 205.3 | $2.50 \times 10^{-03}$ | 4.5 |
| | DnaI | $5.20 \times 10^{-03}$ | 18.0 | | | | |
| | Dps | $9.10 \times 10^{-21}$ | 65.3 | $3.50 \times 10^{-05}$ | 8.4 | $8.70 \times 10^{-03}$ | 6.7 |
| | Fis | $5.80 \times 10^{-07}$ | 180.9 | $3.30 \times 10^{-03}$ | 7.9 | $1.40 \times 10^{-02}$ | 25.1 |
| | Hda | $7.30 \times 10^{-07}$ | 149.1 | $5.30 \times 10^{-03}$ | 7.9 | $1.90 \times 10^{-02}$ | 18.0 |
| | Hfq | $1.40 \times 10^{-12}$ | 121.7 | $3.00 \times 10^{-04}$ | 6.9 | $8.10 \times 10^{-04}$ | 19.3 |
| | H-NS | $1.10 \times 10^{-05}$ | 2.8 | | | $3.80 \times 10^{-04}$ | 2.8 |
| | HupA | $2.70 \times 10^{-04}$ | 15.1 | | | | |
| | HupB | $1.20 \times 10^{-53}$ | 97.6 | $2.30 \times 10^{-08}$ | 6.7 | $2.40 \times 10^{-07}$ | 11.6 |
| | IciA | $7.10 \times 10^{-20}$ | 3.2 | | | $4.50 \times 10^{-07}$ | 1.8 |
| | IhfA, HimA | $1.70 \times 10^{-12}$ | 63.8 | $1.40 \times 10^{-03}$ | 10.5 | $4.90 \times 10^{-02}$ | 6.9 |
| | IhfB, HimD | $1.20 \times 10^{-14}$ | 68.4 | $4.90 \times 10^{-04}$ | 8.4 | $8.40 \times 10^{-03}$ | 9.9 |
| | Lrp | $1.60 \times 10^{-19}$ | 8.4 | | | $5.40 \times 10^{-11}$ | 8.1 |
| | Rob | $6.30 \times 10^{-19}$ | 5.3 | | | $3.40 \times 10^{-08}$ | 4.2 |
| | SeqA | $1.60 \times 10^{-03}$ | 25.9 | | | | |
| | ssb | $5.90 \times 10^{-41}$ | 298.3 | $5.00 \times 10^{-18}$ | 160.6 | | |
| PARTITION | Fic | $3.10 \times 10^{-09}$ | 10.3 | | | $8.60 \times 10^{-03}$ | 7.2 |
| | GidA, MnmG, | $5.20 \times 10^{-13}$ | 1,477.2 | $2.90 \times 10^{-08}$ | 110.6 | $4.30 \times 10^{-02}$ | 18.2 |
| | GidB, RsmG | $6.70 \times 10^{-17}$ | 6,059.9 | $2.20 \times 10^{-15}$ | 252.5 | $9.00 \times 10^{-03}$ | 32.2 |
| | MreB | $1.30 \times 10^{-21}$ | 1,598.2 | $3.90 \times 10^{-12}$ | 40.1 | $1.40 \times 10^{-05}$ | 24.1 |
| | MreC | $2.90 \times 10^{-11}$ | 1,311.2 | $1.30 \times 10^{-08}$ | 46.3 | $8.90 \times 10^{-03}$ | 32.8 |
| | MreD | $1.80 \times 10^{-08}$ | 459.2 | $6.80 \times 10^{-05}$ | 19.8 | $1.90 \times 10^{-02}$ | 18.2 |
| | Mrp | $6.60 \times 10^{-17}$ | 2,599.3 | $1.30 \times 10^{-14}$ | 35.0 | $2.50 \times 10^{-05}$ | 86.2 |
| | MukB | $2.30 \times 10^{-03}$ | 27.4 | | | $1.90 \times 10^{-02}$ | 18.2 |
| | MukE | $3.10 \times 10^{-03}$ | 21.0 | | | $1.90 \times 10^{-02}$ | 18.2 |
| | | | | | | $1.90 \times 10^{-02}$ | 18.2 |
| | ParA, Soj | $2.70 \times 10^{-38}$ | 9.9 | $9.00 \times 10^{-06}$ | 2.6 | $8.40 \times 10^{-06}$ | 3.8 |
| | ParB, Spo0J | $2.50 \times 10^{-44}$ | 13.7 | $3.00 \times 10^{-03}$ | 2.1 | $2.30 \times 10^{-06}$ | 4.1 |
| | ParC | $3.00 \times 10^{-27}$ | 4,149.3 | $3.00 \times 10^{-16}$ | 134.0 | $4.60 \times 10^{-04}$ | 12.3 |
| | ParE | $7.30 \times 10^{-26}$ | 5,842.4 | $5.70 \times 10^{-15}$ | 350.1 | $2.40 \times 10^{-04}$ | 15.8 |
| | RodA, MrdB | $2.80 \times 10^{-12}$ | 1,233.1 | $9.70 \times 10^{-10}$ | 33.0 | $2.60 \times 10^{-03}$ | 55.3 |
| | TrmFO, Gid | $1.50 \times 10^{-06}$ | 182.5 | $4.40 \times 10^{-03}$ | 8.3 | $1.90 \times 10^{-02}$ | 18.0 |
| | XerC | $1.70 \times 10^{-43}$ | 55.0 | $3.10 \times 10^{-08}$ | 8.8 | $1.80 \times 10^{-04}$ | 6.7 |
| | XerD | $1.30 \times 10^{-38}$ | 26.6 | $4.10 \times 10^{-08}$ | 3.4 | $2.50 \times 10^{-06}$ | 6.2 |
| SEGREGATION | ScpA | $1.40 \times 10^{-11}$ | 789.4 | $5.70 \times 10^{-07}$ | 42.9 | $2.10 \times 10^{-02}$ | 16.6 |
| | ScpB | $7.50 \times 10^{-32}$ | 102.5 | $1.80 \times 10^{-07}$ | 25.8 | | |
| | SepF | $1.80 \times 10^{-07}$ | 68.8 | $1.40 \times 10^{-02}$ | 12.3 | | |
| | SlmA, Ttk | $3.80 \times 10^{-09}$ | 52.3 | $1.20 \times 10^{-02}$ | 4.6 | $1.50 \times 10^{-02}$ | 7.5 |
| | Smc | $1.60 \times 10^{-08}$ | 3,090.5 | $1.40 \times 10^{-05}$ | 131.9 | | |
| | SulA | $3.30 \times 10^{-06}$ | 17.5 | | | $1.50 \times 10^{-02}$ | 10.7 |
| CELL DIVISION | AcrA | $6.60 \times 10^{-19}$ | 2.8 | $1.70 \times 10^{-02}$ | 1.1 | $5.30 \times 10^{-10}$ | 2.7 |
| | AmiA, AmiB, | $6.40 \times 10^{-36}$ | 46.4 | $2.90 \times 10^{-10}$ | 8.9 | $4.60 \times 10^{-03}$ | 3.0 |
| | DivIC, DivA | $4.90 \times 10^{-05}$ | 90.5 | $4.70 \times 10^{-02}$ | 8.1 | | |
| | DivIVA | $4.10 \times 10^{-06}$ | 128.0 | $1.10 \times 10^{-02}$ | 13.4 | | |
| | EzrA | $1.00 \times 10^{-02}$ | 13.7 | | | | |
| | FtsA | $9.50 \times 10^{-12}$ | 742.7 | $2.20 \times 10^{-08}$ | 24.6 | $2.50 \times 10^{-03}$ | 41.7 |
| | FtsB | $1.10 \times 10^{-06}$ | 167.2 | $5.40 \times 10^{-03}$ | 16.1 | | |
| | FtsE | $4.20 \times 10^{-24}$ | 2.3 | $1.30 \times 10^{-06}$ | 1.1 | $4.00 \times 10^{-11}$ | 1.9 |
| | FtsI | $9.80 \times 10^{-09}$ | 47.0 | $7.00 \times 10^{-16}$ | 3.9 | $2.20 \times 10^{-07}$ | 76.7 |
| | FtsK, SpoIIIE | $2.80 \times 10^{-37}$ | 76.9 | $2.70 \times 10^{-08}$ | 15.8 | $1.40 \times 10^{-02}$ | 4.2 |
| | FtsL | $1.20 \times 10^{-05}$ | 91.5 | $2.70 \times 10^{-02}$ | 9.8 | | |
| | FtsN | $1.60 \times 10^{-04}$ | 53.0 | | | | |
| | FtsQ | $1.70 \times 10^{-15}$ | 2,135.0 | $1.30 \times 10^{-11}$ | 99.3 | $9.00 \times 10^{-03}$ | 28.8 |
| | FtsW, SpoVE | $5.70 \times 10^{-16}$ | 4,266.4 | $4.40 \times 10^{-16}$ | 87.7 | $8.20 \times 10^{-04}$ | 55.0 |
| | FtsX | $9.30 \times 10^{-12}$ | 972.9 | $1.30 \times 10^{-08}$ | 13.8 | $4.80 \times 10^{-04}$ | 146.2 |
| | FtsZ | $3.10 \times 10^{-31}$ | 2,747.0 | $1.20 \times 10^{-19}$ | 101.6 | $9.70 \times 10^{-04}$ | 16.5 |
| | MinC | $4.40 \times 10^{-09}$ | 172.3 | $1.20 \times 10^{-02}$ | 3.0 | $5.80 \times 10^{-05}$ | 76.8 |
| | MinD | $3.10 \times 10^{-19}$ | 42.8 | $1.60 \times 10^{-04}$ | 2.3 | $5.40 \times 10^{-11}$ | 81.5 |
| | MinE | $9.00 \times 10^{-09}$ | 152.9 | $3.10 \times 10^{-02}$ | 2.6 | $5.90 \times 10^{-05}$ | 75.2 |
| | ZapA | $8.20 \times 10^{-09}$ | 602.8 | $7.40 \times 10^{-06}$ | 17.3 | $7.30 \times 10^{-03}$ | 56.1 |
| | ZipA | $7.90 \times 10^{-05}$ | 66.0 | | | | |

Left side overall label: KEGG entries (chromosome-like)

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| REPLICATION | CdsD | | | $4.40 \times 10^{-02}$ | 0.1 | | |
| | DNA helicase | $5.80 \times 10^{-21}$ | 33.6 | $2.70 \times 10^{-04}$ | 4.1 | $1.30 \times 10^{-04}$ | 9.8 |
| | Helicase-1 | $1.60 \times 10^{-27}$ | 71.1 | $1.90 \times 10^{-13}$ | 20.0 | $1.10 \times 10^{-04}$ | 4.6 |
| | DNA repair | $2.20 \times 10^{-04}$ | 34.0 | | | $5.70 \times 10^{-04}$ | 43.6 |
| | primase, LtrC | $3.10 \times 10^{-02}$ | 1.8 | | | | |
| | RepA | $5.90 \times 10^{-03}$ | 0.7 | | | | |
| | RepAEB | $1.70 \times 10^{-16}$ | 0.0 | $1.90 \times 10^{-04}$ | 0.1 | | |
| | RepC | | | | | $9.60 \times 10^{-03}$ | 2.7 |
| | RepCJE | | | $4.40 \times 10^{-02}$ | 0.1 | | |
| | RepRSE | $1.30 \times 10^{-02}$ | 0.0 | $4.90 \times 10^{-02}$ | 0.1 | | |
| | RNA polymerase | $3.20 \times 10^{-02}$ | 6.3 | | | | |
| | Rop | $3.20 \times 10^{-02}$ | 0.0 | $4.40 \times 10^{-02}$ | 0.1 | | |
| | RuvB | $1.20 \times 10^{-08}$ | 433.0 | $5.70 \times 10^{-08}$ | 17.7 | $1.40 \times 10^{-05}$ | 37.8 |
| | TrfA | $1.40 \times 10^{-02}$ | 0.3 | | | | |
| PARTITION | ATPase, TyrK, | $2.20 \times 10^{-20}$ | 19.4 | | | $8.50 \times 10^{-07}$ | 9.3 |
| | CopG | | | $2.70 \times 10^{-02}$ | 0.2 | $4.60 \times 10^{-03}$ | 23.1 |
| | DNA-binding protein | | | $4.40 \times 10^{-02}$ | 0.1 | | |
| | FtsK, SpoIIIE | $1.90 \times 10^{-07}$ | 6.0 | | | $9.90 \times 10^{-05}$ | 9.8 |
| | ParA, ParM | $1.50 \times 10^{-10}$ | 0.4 | $4.00 \times 10^{-06}$ | 0.3 | | |
| | ParB | $5.70 \times 10^{-12}$ | 0.1 | $1.40 \times 10^{-05}$ | 0.2 | | |
| | serine recombinase | $2.50 \times 10^{-06}$ | 1.4 | $1.50 \times 10^{-03}$ | 2.9 | $1.80 \times 10^{-02}$ | 0.4 |
| | tyrosine recombinase | $3.40 \times 10^{-04}$ | 3.3 | | | $7.40 \times 10^{-04}$ | 8.7 |
| | Xer-like Tyrosine | $7.60 \times 10^{-11}$ | 2.0 | | | $6.30 \times 10^{-03}$ | 1.6 |
| MAINTENANCE | Ccd (PSK) | $4.60 \times 10^{-02}$ | 3.9 | | | | |
| | HicAB (PSK) | $4.30 \times 10^{-05}$ | 25.2 | | | $4.80 \times 10^{-03}$ | 15.1 |
| | HigBA (PSK) | $3.30 \times 10^{-15}$ | 3.4 | $2.40 \times 10^{-02}$ | 1.5 | $1.20 \times 10^{-03}$ | 2.5 |
| | MazEF (PSK) | $1.20 \times 10^{-11}$ | 5.2 | $2.90 \times 10^{-02}$ | 2.6 | | |
| | ParC (PSK) | | | $4.40 \times 10^{-02}$ | 0.1 | | |
| | ParDE (PSK) | $5.50 \times 10^{-08}$ | 2.3 | | | $7.80 \times 10^{-05}$ | 3.4 |
| | PhD, Doc (PSK) | $3.20 \times 10^{-07}$ | 11.9 | | | $2.90 \times 10^{-03}$ | 8.8 |
| | plasmid maintenance | | | $4.40 \times 10^{-02}$ | 0.1 | | |
| | RelBE (PSK) | $2.70 \times 10^{-08}$ | 3.5 | | | $6.10 \times 10^{-04}$ | 4.2 |
| | VapBC/Vag (PSK) | $1.20 \times 10^{-09}$ | 3.9 | | | $1.40 \times 10^{-05}$ | 5.8 |

*ACLAME Families (plasmid-like)*

Chromosome-signature ISs are also present on the SERs, and some of them are enriched

to the same order of magnitude in both classes but not in plasmids (highlighted in

Table 6). Among these latter, helicase loader DnaC participates to the replication

initiation of the chromosome (Chodavarapu et al., 2016) whilst Walker-type ATPase

ParA/Soj interacts with ParB/Spo0J in the *parABS* chromosomal partinioning system,

and is required for proper separation of sister origins and synchronous DNA replication

(Murray and Errington, 2008). The other ISs have a regulatory role, either locally or

globally. Nucleoid-associated proteins (NAPs; Dillon and Dorman, 2010) contribute to

the replication regulation: H-NS (histone-like nucleoid structuring protein), IciA

(chromosome initiator inhibitor, LysR family transcriptional regulator), MukBEF

(condensin), and Rob/ClpB (right arm of the replication origin binding protein/curved

DNA-binding protein B, AraC family transcriptional regulator) influence both the

conformation and the functions of chromosomal DNA, replication, recombination and

repair. The NAPs also have pleiotropic regulatory roles in global regulation of gene

transcription depending on cell growth conditions (H-NS, IciA, Lrp (leucine-responsive

regulatory protein, Lrp/AsnC family transcriptional regulator), and Rob/ClpB).

Similarly, the membrane fusion protein AcrA is a growth-dependent regulator, mostly

known for its role as a peripheral scaffold mediating the interaction between AcrB and

TolC in the AcrA-AcrB-TolC Resistance-Nodule-cell Division-type efflux pump that

extrudes from the cell compounds that are toxic or have a signaling role (Du et al.,

2018). It is central to the regulation of cell homeostasis and proper development (Anes

et al., 2015; Du et al., 2018; Webber et al., 2009) as well as biofilm formation (Alav et

al., 2018). Fic (cell filamentation protein) targets the DNA gyrase B (GyrB) to regulate

the cell division and cell morphology (Lu et al., 2018) whereas SulA inhibits FtsZ

assembly, hence causing incomplete cell division and filamentation (Chen et al, 2012).

FtsE is involved in the Z-ring assembly and the initiation of constriction, and in late

stage cell separation (Meier et al, 2017).

The main divergence between SERs and chromosomes lies in the distribution patterns of

the ACLAME-annotated ISs ($OR < 10^0$ in the chromosomes *vs.* SERs comparison). Their

higher abundance on the SERs suggests a stronger link of SERs to plasmids. This

pattern may also arise from the unbalanced taxon representation in our SER dataset due

to a single bacterial lineage. For example, the presence of RepC is likely to be specific to

Rhizobiales SERs (Pinto et al., 2012).

**Identification of candidate SERs**

Since the IS profiles constitute replicon-type signatures, we searched for new putative

SERs or chromosomes among the extra-chromosomal replicons. We used the IS

functions as features to perform supervised classification analyses with various training

sets (Table 7).

**Table 7. Performance of the ERT classification procedures**

| Training Set [a] | $CV_{score}$ [b] | $\sigma_{CV_{score}}$ [c] | $OOB_{score}$ [d] | $\sigma_{OOB_{score}}$ [e] |
|---|---|---|---|---|
| $\{E_{SER}, E'_{plasmid}\}$ | 0.96 | - | 0.96 | - |
| $\{E_{SER}, E'_{plasmid}\}^{it}$ | 0.92 | 0.02 | 0.93 | 0.02 |
| $\{E_{chromosome}, E'_{plasmid}\}$ | 1.00 | - | 1.00 | - |
| $\{E_{SER}, E_{chromosome}\}^{it}$ | 0.98 | 0.00 | 0.98 | 0.01 |

[a] $E_{chromosome}$ and $E_{SER}$ are host genus-normalized sets of chromosomes or SERs, respectively (*cf.*
Table 13). $E'_{plasmid}$ is derived from the INFOMAP clustering solution, by discarding plasmids belonging to
clusters also harbouring SERs or chromosomes, and normalized according to host genus. "*it*" designates the
iterative procedure.
[b] Cross-validation score or mean of iteration cross-validation scores.
[c] Standard deviation of iteration cross-validation scores.
[d] Out-of bag estimate or mean of iteration Out-of bag estimates.
[e] Standard deviation of iteration Out-of bag estimates.

The coherence of the SER class (overall high values of the probability for a SER to be

assigned to its own class in Tables 7 and 8) confirmed that the ISs are robust genomic

markers for replicon characterization. The low SER probability scores presented by a

few SERs (Table 8) likely result from a low number of carried ISs (*e.g., Leptospira*), or

from the absence in the data of lineage-specific ISs (*e.g.*, SER idiosyncratic replication

initiator RtcB of Vibrionaceae).

**Table 8. SER probability to belong to the *SER* class**

| Genus | $\bar{P}_{SER}(SER)$ [a] |
|---|---|
| *Agrobacterium* | 0.90 |
| *Aliivibrio* | 0.87 |
| *Anabaena* | 0.94 |
| *Asticcacaulis* | 0.95 |

| | |
|---|---|
| *Brucella* | 0.92 |
| *Burkholderia* | 0.89 |
| *Butyrivibrio* | 0.83 |
| *Chloracidobacterium* | 0.88 |
| *Cupriavidus* | 0.94 |
| *Cyanothece* | 0.86 |
| *Deinococcus* | 0.78 |
| *Ensifer/Sinorhizobium* | 0.90 |
| *Ilyobacter* | 0.88 |
| *Leptospira* | 0.54 |
| *Nocardiopsis* | 0.90 |
| *Ochrobactrum* | 0.96 |
| *Paracoccus* | 0.96 |
| *Photobacterium* | 0.95 |
| *Prevotella* | 0.92 |
| *Pseudoalteromonas* | 0.91 |
| *Ralstonia* | 0.95 |
| *Rhodobacter* | 0.69 |
| *Sphaerobacter* | 0.88 |
| *Sphingobium* | 0.73 |
| *Thermobaculum* | 0.78 |
| *Variovorax* | 0.83 |
| *Vibrio* | 0.76 |

[a] SER probability, averaged per host genus, to belong to the *SER* class in the supervised classification

using $\{E_{SER}, E_{plasmid}\}^{it}$ as training set.

We detected a number of candidate SERs among the plasmids (Table 9a), most of which

are essential to the cell functioning and/or the fitness of the organism (*cf.* Box 1).

Whereas most belong to bacterial lineages known to harbour multipartite genomes,

novel taxa emerge as putative hosts to complex genomes (Rhodospirillales and, to a

lesser extent, Actinomycetales). In contrast, our analyses confirmed only one putative

SER (*Ruegeria* sp. TM1040) within the *Roseobacter* clade (Petersen et al., 2013).

Remarkably, we identified eight candidate chromosomes corresponding to two plasmids,

also identified as candidate SERs, that encode ISs hardly found in extra-chromosomal

elements (*e.g.*, DnaG, DnaB, ParC and ParE), and six SERs that part of, or all, our

analyses associate to standard chromosomes (Table 9b). Notably, *Prevotella intermedia*

SER (CP003503) shows a very high probability (> 0.98) to be a chromosome while its

annotated chromosome (CP003502), unique of its kind, falls within the plasmid class.

This approach can thus be extended to test the type of replicon for (re)annotation

purposes.

**Table 9. Identification of ERs among the extra-chromosomal replicons**

a. Candidate-SERs identified among plasmids

| Replicon | Probability[a] |
|---|---|
| *Acaryochloris marina* MBIC11017 plasmid pREB1 [CYANOBACTERIA : Chroococcales] (CP000838) | 0.578 |
| *Acaryochloris marina* MBIC11017 plasmid pREB2 [CYANOBACTERIA : Chroococcales] (CP000839) | 0.582 |
| *Agrobacterium* sp. H13-3 plasmid pAspH13-3a [α-PROTEOBACTERIA : Rhizobiales] (CP0022) | 0.565 |
| *Arthrobacter chlorophenolicus* A6 plasmid pACHL01 [ACTINOBACTERIA : Actinomycetales] (CP001342) | 0.648 |
| *Azospirillum brasilense* Sp245 plasmid AZOBR_p1 [α-PROTEOBACTERIA : Rhodospirillales] (HE577328) | 0.878 |
| *Azospirillum brasilense* Sp245 plasmid AZOBR_p2 [α-PROTEOBACTERIA : Rhodospirillales] (HE577329) | 0.591 |
| *Azospirillum brasilense* Sp245 plasmid AZOBR_p3 [α-PROTEOBACTERIA : Rhodospirillales] (HE577330) | 0.603 |
| *Azospirillum lipoferum* 4B plasmid AZO_p1e [α-PROTEOBACTERIA : Rhodospirillales] (FQ311869) | 0.722 |
| *Azospirillum lipoferum* 4B plasmid AZO_p2 [α-PROTEOBACTERIA : Rhodospirillales] (FQ311870) | 0.609 |
| *Azospirillum lipoferum* 4B plasmid AZO_p4 [α-PROTEOBACTERIA : Rhodospirillales] (FQ311872) | 0.645 |
| *Azospirillum* sp. B510 plasmid pAB510a [α-PROTEOBACTERIA : Rhodospirillales] (AP010947) | 0.732 |
| *Azospirillum* sp. B510 plasmid pAB510c [α-PROTEOBACTERIA : Rhodospirillales] (AP010949) | 0.545 |
| *Azospirillum* sp. B510 plasmid pAB510d [α-PROTEOBACTERIA : Rhodospirillales] (AP010950) | 0.530 |
| *Burkholderia phenoliruptrix* BR3459a plasmid pSYMBR3459 [β-PROTEOBACTERIA : Burkholderiales] (CP003865) | 0.663 |
| *Burkholderia phymatum* STM815 plasmid pBPHY01 [β-PROTEOBACTERIA : Burkholderiales] (CP001045) | 0.733 |
| *Burkholderia* sp. YI23 plasmid byi_1p [β-PROTEOBACTERIA : Burkholderiales] (CP003090) | 0.846 |
| *Clostridium botulinum* A3 str. Loch Maree plasmid pCLK [FIRMICUTES : Clostridiales] (CP000963) | 0.531 |
| *Clostridium botulinum* Ba4 str. 657 plasmid pCLJ [FIRMICUTES : Clostridiales] (CP001081) | 0.531 |
| *Cupriavidus metallidurans* CH34 megaplasmid [β-PROTEOBACTERIA : Burkholderiales] (CP000353) | 0.883 |
| *Cupriavidus necator* N-1 plasmid BB1p [β-PROTEOBACTERIA : Burkholderiales] (CP002879) | 0.500 |
| *Cupriavidus pinatubonensis* JMP134 megaplasmid [β-PROTEOBACTERIA : Burkholderiales] (CP000092) | 0.513 |
| *Deinococcus geothermalis* DSM 11300 plasmid1 [DEINOCOCCUS-THERMUS : Deinococcales] (CP000358) | 0.622 |
| *Deinococcus gobiensis* I-0 plasmid P1 [DEINOCOCCUS-THERMUS : Deinococcales] (CP002192) | 0.812 |
| *Ensifer/Sinorhizobium fredii* HH103 plasmid pSfHH103e [α-PROTEOBACTERIA : Rhizobiales] (HE616899) | 0.915 |
| *Ensifer/Sinorhizobium fredii* NGR234 plasmid pNGR234b [α-PROTEOBACTERIA : Rhizobiales] (CP000874) | 0.894 |
| *Ensifer/Sinorhizobium medicae* WSM419 plasmid pSMED01 [α-PROTEOBACTERIA : Rhizobiales] (CP000739) | 0.942 |
| *Ensifer/Sinorhizobium medicae* WSM419 plasmid pSMED02 [α-PROTEOBACTERIA : Rhizobiales] (CP000740) | 0.836 |
| *Ensifer/Sinorhizobium meliloti* 1021 plasmid pSymA [α-PROTEOBACTERIA : Rhizobiales] (AE006469) | 0.818 |
| *Ensifer/Sinorhizobium meliloti* 1021 plasmid pSymB [α-PROTEOBACTERIA : Rhizobiales] (AL591985) | 0.949 |
| *Ensifer/Sinorhizobium meliloti* BL2C plasmid pSINMEB01 [α-PROTEOBACTERIA : Rhizobiales] (CP002741) | 0.800 |
| *Ensifer/Sinorhizobium meliloti* BL2C plasmid pSINMEB02 [α-PROTEOBACTERIA : Rhizobiales] (CP002742) | 0.961 |
| *Ensifer/Sinorhizobium meliloti* Rm41 plasmid pSYMA [α-PROTEOBACTERIA : Rhizobiales] (HE995407) | 0.922 |
| *Ensifer/Sinorhizobium meliloti* Rm41 plasmid pSYMB [α-PROTEOBACTERIA : Rhizobiales] (HE995408) | 0.960 |
| *Ensifer/Sinorhizobium meliloti* SM11 plasmid pSmeSM11c [α-PROTEOBACTERIA : Rhizobiales] (CP001831) | 0.877 |
| *Ensifer/Sinorhizobium meliloti* SM11 plasmid pSmeSM11d [α-PROTEOBACTERIA : Rhizobiales] (CP001832) | 0.947 |
| *Methylobacterium extorquens* AM1 megaplasmid [α-PROTEOBACTERIA : Rhizobiales] (CP001511) | 0.538 |
| *Novosphingobium* sp. PP1Y plasmid Mpl [α-PROTEOBACTERIA : Sphingomonadales] (FR856861) | 0.523 |
| *Pantoea* sp. At-9b plasmid pPAT9B01 [γ-PROTEOBACTERIA : Enterobacteriales] (CP002434) | 0.527 |

| | |
|---|---|
| *Paracoccus denitrificans* PD1222 plasmid1 [α-PROTEOBACTERIA : Rhodobacterales] (CP000491) | 0.769 |
| *Ralstonia solanacearum* GMI0 plasmid pGMI0MP [β-PROTEOBACTERIA : Burkholderiales] (AL646053) | 0.861 |
| *Ralstonia solanacearum* Po82 megaplasmid [β-PROTEOBACTERIA : Burkholderiales] (CP002820) | 0.865 |
| *Ralstonia solanacearum* PSI07 megaplasmid [β-PROTEOBACTERIA : Burkholderiales] (FP885891) | 0.827 |
| *Rhizobium etli* CFN 42 plasmid p42e [α-PROTEOBACTERIA : Rhizobiales] (CP000137) | 0.700 |
| *Rhizobium etli* CFN 42 plasmid p42f [α-PROTEOBACTERIA : Rhizobiales] (CP000138) | 0.555 |
| *Rhizobium etli* CIAT 652 plasmid pA [α-PROTEOBACTERIA : Rhizobiales] (CP0010) | 0.701 |
| *Rhizobium etli* CIAT 652 plasmid pC [α-PROTEOBACTERIA : Rhizobiales] (CP001077) | 0.792 |
| *Rhizobium leguminosarum* bv. trifolii WSM1325 plasmid pR132501 [α-PROTEOBACTERIA : Rhizobiales] (CP001623) | 0.711 |
| *Rhizobium leguminosarum* bv. trifolii WSM1325 plasmid pR132502 [α-PROTEOBACTERIA : Rhizobiales] (CP001624) | 0.741 |
| *Rhizobium leguminosarum* bv. *trifolii* WSM2304 plasmid pRLG201 [α-PROTEOBACTERIA : Rhizobiales] (CP001192) | 0.777 |
| *Rhizobium leguminosarum* bv. *trifolii* WSM2304 plasmid pRLG202 [α-PROTEOBACTERIA : Rhizobiales] (CP001193) | 0.630 |
| *Rhizobium leguminosarum* bv. *viciae* 3841 plasmid pRL11 [α-PROTEOBACTERIA : Rhizobiales] (AM236085) | 0.731 |
| *Rhizobium leguminosarum* bv. *viciae* 3841 plasmid pRL12 [α-PROTEOBACTERIA : Rhizobiales] (AM236086) | 0.718 |
| *Ruegeria* sp. TM1040 megaplasmid [α-PROTEOBACTERIA : Rhodobacterales] (CP000376) | 0.667 |
| *Streptomyces cattleya* NRRL 8057 plasmid pSCA [ACTINOBACTERIA : Actinomycetales] (FQ859184) | 0.727 |
| *Streptomyces cattleya* NRRL 8057 plasmid pSCATT [ACTINOBACTERIA : Actinomycetales] (CP003229) | 0.702 |
| *Streptomyces clavuligerus* ATCC 27064 plasmid pSCL4 [ACTINOBACTERIA : Actinomycetales] (CM000914) | 0.642 |
| *Streptomyces clavuligerus* ATCC 27064 plasmid pSCL4 [ACTINOBACTERIA : Actinomycetales] (CM001019) | 0.642 |
| *Thermus thermophilus* HB8 plasmid pTT27 [DEINOCOCCUS-THERMUS : Thermales] (AP008227) | 0.500 |
| *Thermus thermophilus* JL-18 plasmid pTTJL1801 [DEINOCOCCUS-THERMUS : Thermales] (CP0033) | 0.557 |
| *Tistrella mobilis* KA081020-065 plasmid pTM2 [α-PROTEOBACTERIA : Rhodospirillales] (CP003238) | 0.578 |
| *Tistrella mobilis* KA081020-065 plasmid pTM3 [α-PROTEOBACTERIA : Rhodospirillales] (CP003239) | 0.797 |

358 b. Candidate chromosomes identified among extra-chromosomal replicons

| REPLICON | PROBABILITY [a] |
|---|---|
| *Anaeba* sp. 90 chromosome chANA02 [CYANOBACTERIA : Chroococcales] (CP003285) | 0.638 |
| *Asticcacaulis excentricus* CB 48 chromosome 2 [α-PROTEOBACTERIA : Caulobacterales] (CP002396) | 0.637 |
| *Azospirillum brasilense* Sp245 plasmid AZOBR_p1 [α-PROTEOBACTERIA : Rhodospirillales] (HE577328) | 0.774 |
| *Methylobacterium extorquens* AM1 megaplasmid [α-PROTEOBACTERIA : Rhizobiales] (CP001511) | 0.669 |
| *Nocardioides dassonvillei* DSM 43111 chromosome 2 [ACTINOBACTERIA : Actinomycetales] (CP002041) | 0.539 |
| *Paracoccus denitrificans* PD1222 chromosome2 [α-PROTEOBACTERIA : Rhodobacterales] (CP000490) | 0.778 |
| *Prevotella intermedia* 17 chromosome II [BACTEROIDETES : Bacteroidales] (CP0033) | 0.984 |
| *Prevotella melaninogenica* ATCC 845 chromosome II [BACTEROIDETES : Bacteroidales] (CP002123) | 0.698 |

359  [a] Probability for an extra-chromosomal replicon, *i.e.*, plasmid or SER, to belong to the SER (a) or Chromosome
360    (b) class according to the supervised classification procedures.

**BOX 1. CHARACTERISTICS OF CANDIDATE SERS**

According to the literature, most candidate SERs that we detected among plasmids (Table 9a) were

expected to be essential to the cell functioning and/or to the fitness of the organism.

• *Azospirillum* genomes are constituted of multiple replicons, at least one of which is expected to be

essential. The largest extra-chromosomal replicon in *A. brasilense* was proposed to be essential for

bacterial life (Wisniewski-Dyé et al*.,* 2011) since it encodes well-conserved housekeeping genes involved

in DNA replication, RNA metabolism and biosynthesis of nucleotides and cofactors, as well as in

transport and protein post-translational modifications. This replicon is unambiguously identified as a SER

by our analyses, as are additional replicons found in *A. lipoferum* and *A.* sp. B510, expected homologues

to *A. brasilense* SER (Acosta-Cruz et al*.,* 2012). In contrast, other extra-chromosomal replicons classified

as chromids by Wisniewski-Dyé et al*.* (2012) are unlikely to be true essential replicons. They were not

retrieved among our candidate SERs.

• In *Rhizobium etli* CFN42, functional interactions among sequences scattered in the different

extrachromosomal replicons are required for successful completion of life in symbiotic association

with plant roots or saprophytic growth (Brom et al., 2000). p42e (CP000137) is the only replicon other

than the chromosome that contains genes involved in the primary metabolism (Landeta et al., 2011;

Villaseñor *et al.* 2011) and evades its elimination by co-integration with other replicons including the

chromosome (Landeta et al., 2011). Furthermore, homologues to this replicon were identified in the

genomes of other *R. etli* strains as well as other *Rhizobium* species: *R. etli* CIAT652 pA,

*R. leguminosarum* bv. *viciae* 3841 pRL11, *R. leguminosarum* bv. *trifolii* WSM2304 pRLG202 and

*R. leguminosarum* bv. *trifolii* WSM1325 pR132502 (CP001075, AM236085, CP001193, and CP001624,

respectively) (Landeta et al., 2011; Villaseñor et al., 2011). These replicons were thus proposed to be

secondary chromosomes (Landeta et al., 2011).

• The genome of *Ensifer*/*Sinorhizobium meliloti* AK83 was the single multipartite-annotated

*Ensifer*/*Sinorhizobium* genomes present in our dataset. This bacterium carries two large extra-

chromosomal replicons that are involved in the establishment of the nitrogen fixation symbiosis with

legume plants. pSymA contains most of the genes involved in the nodulation and nitrogen fixation

whereas pSymB carries exopolysaccharide biosynthetic genes, also required for the establishment of the

symbiosis. Our analyses identifies candidate SERs similar to *S. meliloti* AK83 pSymA and pSymB in

other *S. meliloti* strains as well as in *S. fredii* and *S. medicae*. pSymB has been referred to as second

chromosome for carrying genes encoding essential house-keeping functions (Blanca-Ordóñez et al., 2010 ;

Galardini et al., 2011). It shows a higher level of conservation across strains and species than pSymA

(Galardini et al., 2013). pSymA, generally thought to be as stable as pSymB, greatly contribute to the

bacterial fitness in the rhizosphere (Blanca-Ordóñez et al., 2010; Galardini et al., 2013).

• The identification of *Methylobacterium extorquens* AM1 1.2 Mb megaplasmid as a SER is supported by

its presence in the genome in a predicted one copy number, by its coding a truncated *luxI* gene essential

for the operation of two chromosomally-located *luxI* genes, as well as the single *umuDC* cluster involved

in SOS DNA repair, and by the presence of a 130 kb region syntenic to a region of similar length in the

chromosome of *Methylobacterium extorquens* strain DM4 (Vuilleumier et al., 2009).

• The megaplasmid (821 kb) in *Ruegeria* sp. TM1040 carries more rRNA operons (3) than the

chromosome (1) and several unique genes (Moran et al., 2007). *Ruegeria* sp. TM1040 is the only species

in the *Roseobacter* group that possesses a SER. None of the plasmids in the other species included in our

datasets qualified as SERs according to our results in contrast to the commonly held view (Petersen et al.,

2013).

• In *Burkholderia* genus, additional ERs possess a centromere whose sequence is distinct from, but

strongly resembles that of the chromosome centromere (Dubarry et al., 2009). However, these

centromeres have a common origin and a plasmid ancestry (Passot et al., 2012). The evolution of these

replicons into SERs is best accounted for by the high level of plasticity observed in the *Burkholderia*

genomes, with extra-chromosomal replicons going through extensive exchange of genetic material among

them as well as with the chromosomes (Maida et al., 2014).

• *Acaryochloris marina* MBIC11017 pREB1 (CP000838) and pREB2 (CP000839) plasmids were

identified as candidate SERs. Both these megaplasmids code for metabolic key-proteins, and are thus

likely to contribute to the bacterium fitness (Swingley et al., 2008).

• The genomes of *Streptomyces cattleya* NRRL8057 and *S. clavuligerus* ATCC27064 harbour a linear

megaplasmid (1.8 Mb) that shows a high probability (P ≈ 0.7) to be a SER. The megaplasmid of

*S. cattleya* NRRL8057 encodes genes involved in the synthesis of various antibiotics and secondary

metabolites and is expected to be important to the life of the bacterium in its usual habitat (Barbe et al.,

2011; O'Rourke et al., 2009). In *S. clavuligerus* ATCC27064, none of the megaplasmid-encoded genes are

expected to belong to the core genome (Medema et al., 2010). However, the megaplasmid is likely to

contribute to the bacterium firness. It represents more than 20% of the coding genome and constitutes a

large reservoir of genes involved in bioactive compound production and cross-regulation with the

chromosome (Medema et al., 2010). Furthermore, *S. clavuligerus* chromosome requires the SER-encoded

*tap* gene involved in the telomere replication.

• *Butyrivibrio proteoclasticus* B316 harbours two plasmid, one of which, pCY186 plasmid (CP001813),

was identified as a candidate SERs by our analysis, albeit with a low probability (0.56). In support to this,

it carries numerous genes coding for proteins involved in replication of the chromosome (Yeoman et al.,

2011). The second plasmid in that strain, pCY360 (CP001812), also proposed to be an essential replicon

in that bacterium (Yeoman et al., 2011), presents too low a probability (P = 0.32) in our analysis to qualify

as a SER.

**DISCUSSION**

The SERs clearly stand apart from plasmids, including those that occur consistently in a

bacterial species, *e.g.*, *Lactobacillus salivarius* pMP118-like plasmids (Li et al., 2007).

The replicon size proposed as a primary classification criterion to separate the SERs

from the plasmids (diCenzo and Finan, 2017; Harrison et al., 2010) proves to be

inoperative. The IS profiles accurately identify the SERs of *Leptospira* and *Butyrivibrio*

despite their plasmid-like size, and unambiguously ascribe the chromosomes in the

reduced genomes of endosymbionts (sizes down to 139 kb) to the chromosome class.

Conversely, they assign *Rhodococcus jostii* RHA1 1.12 Mb-long pRHL1 replicon to the

plasmid class, and do not discriminate the megaplasmids (>350 kb (diCenzo and Finan,

2017)) from smaller plasmids. Plasmids may be stabilized in a bacterial population by

rapid compensatory adaptation that alleviates the fitness cost incurred by their presence

in the cell (San Millan et al., 2014; Hall et al., 2017; Stalder et al., 2017). This

phenomenon involves mutations either on the chromosome only, on the plasmid only, or

both, and does not preclude the segregational loss of the plasmid. On the contrary, SERs

code for chromosome-type IS proteins that integrate them constitutively in the species

genome and the cell cycle. The SERs thence qualify as essential replicons regardless of

their size and of the phenotypical/ecological, possibly essential, functions that they

encode and which vary across host taxa.

Yet, SERs also carry plasmid-like ISs, suggesting a role for plasmids in their formation.

The prevailing opinion assumes that SERs derive from the amelioration of

megaplasmids (diCenzo and Finan, 2017; diCenzo et al., 2013; Harrison et al., 2010;

MacLellan et al., 2004; Slater et al., 2009): a plasmid bringing novel functions for the

adaptation of its host to a new environment is stabilized into the bacterial species

genome through the transfer from the chromosome of essential genes (diCenzo and

Finan, 2017; Slater et al., 2009). However, the generalized presence of chromosome-like

ISs in the SERs of the various taxa with multipartite genomes is unlikely to derive from

the action of environment-specific and lineage-specific selective forces. In reverse, all

bacteria with similar lifestyle and exhibiting some phylogenetic relatedness may not

harbor multiple ERs (*e.g.*, α-proteobacterial nitrogen-fixing legume symbionts). Also,

the gene shuttling from chromosome to plasmid proposition fails to account for the

situation met in the multipartite genomes of *Asticaccaulis excentricus, Paracoccus*

*denitrificans* and *Prevotella* species. Their chromosome-type ISs are evenly distributed

between the chromosome and the SER whereas their homologues in the mono- or

multipartite genomes of most closely related species are primarily chromosome-coded

(see Table 10 for an example). This pattern, mirrored in their whole gene content (Naito

et al., 2016; Poirion, 2014), hints at the stemming of the two essential replicons from a

single chromosome by either a splitting event or a duplication followed by massive gene

loss. Neither mechanism informs on the presence of plasmid-type maintenance

machinery on one of the replicons. The severing of a chromosome generates a single

true replicon carrying the chromosome replication origin and an origin-less remnant,

whilst the duplication of the chromosome produces two chromosomal replicons with

identical maintenance systems. Whereas multiple copies of the chromosome are known

to cohabit constitutively in polyploid bacteria (Ohtani et al., 2010), the co-occurrence of

dissimilar chromosomes bearing identical replication initiation and partition systems is

yet to be described in bacteria.

**Table 10. IS profiles of *Paracoccus denitrificans vs. Rhodobacter sphaeroides* (Rhodobacterales)**

Chromosome-like IS functions coded only by the SER in *P. denitrificans* or *R. sphaeroides* whilst by the chromosome in other Rhodobacterales are indicated by an asterisk. Numbers corresponds to the number of homologues (*P. denitrificans* PD1222) or the pourcentage of function-coding replicons (*R. sphaeroides* and Rhodobacterales genomes).

| Group | IS FUNCTION GROUP | P.d * | chromosome1 (CP000489) | chromosome2 (CP000490) | plasmid1 (CP000491) | R.s * | chromosome % (n=12) | SER % | plasmid % (n=14) | chromosome % (n=12) | plasmid % (n=33) |
|---|---|---|---|---|---|---|---|---|---|---|---|
| REPLICATION | CbpA | | 1 | | | | 100 | | | 75 | |
| | Dam | | 1 | | | | 50 | 75 | | 25 | |
| | DnaA | * | | 1 | | | 100 | | | 92 | |
| | DnaB | * | | 1 | | | 100 | | | 100 | |
| | DnaC | | | | | | 25 | | | | 3 |
| | DnaG | * | | 1 | | | 100 | | | 100 | |
| | Dps | | 1 | | | | | | | 50 | 15 |
| | Fis | * | | 1 | | | 100 | | | 75 | |
| | Hda | | | | | | | | | 42 | |
| | Hfq | * | | 1 | | | 100 | | | 100 | |
| | H-NS | | 2 | 2 | | | 100 | | 29 | 67 | 9 |
| | HupA | | | | | | | | | 17 | |
| | HupB | | 1 | | | | 100 | | | 100 | |
| | IciA | | 1 | | | | | | 36 | 67 | |
| | IhfA, HimA | | 1 | | | | 100 | | | 100 | |
| | IhfB, HimD | * | | 1 | | * | | 100 | | 100 | |
| | Lrp | | 4 | 1 | 3 | | 100 | 25 | 50 | 100 | 9 |
| | Rob | | | | | | 25 | | | 8 | |
| | ssb | | 1 | | 3 | | 100 | 100 | | 100 | 3 |
| PARTITION | Fic | | | | | | | | 7 | 8 | 6 |
| | GidA, MnmG, MTO1 | | 1 | | | | 100 | | | 100 | |
| | GidB, RsmG | | 1 | | | | 100 | | | 100 | |
| | MreB | * | | 1 | | | 100 | | | 92 | |
| | MreC | * | | 1 | | | 100 | | | 92 | |
| | MreD | * | | 1 | | | 100 | | | 92 | |
| | Mrp | | 1 | | | | 100 | | | 100 | |
| | ParA, Soj | | 4 | 1 | | | 100 | | | 100 | 9 |
| | ParB, Spo0J | | 3 | 1 | 1 | | 100 | 75 | | 100 | 42 |
| | ParC | | 1 | 1 | | | 100 | | | 100 | |
| | ParE | | 1 | | | | 100 | | | 100 | |
| | RodA, MrdB | * | | 1 | | | 100 | | | 92 | |
| | TrmFO, Gid | * | | 1 | | | 100 | | | 100 | |
| | XerC | | 1 | | | | 100 | | | 100 | |
| | XerD | | 1 | | | | 100 | 25 | 7 | 100 | 6 |
| SEGREGATION | ScpA | | 1 | | | | 100 | | | 100 | |
| | ScpB | | 1 | | | | 100 | | | 100 | |
| | Smc | | 1 | | | | 100 | | | 100 | |
| CELL DIVISION | AcrA | | 2 | 2 | | * | | 100 | 7 | 58 | |
| | AmiA, AmiB, AmiC | | 1 | | | | 100 | | | 100 | |
| | FtsA | * | | 1 | | | 100 | | | 83 | |
| | FtsE | | 1 | | | | 100 | | 7 | 92 | |
| | FtsI | | 1 | | | | 100 | | | 92 | |
| | FtsK, SpoIIIE | | 1 | | | | 100 | | | 100 | |
| | FtsQ | * | | 1 | | | 100 | | | 92 | |
| | FtsW, SpoVE | | | 1 | | | 100 | | | 100 | |
| | FtsX | | 1 | | | | 100 | | 7 | 92 | |
| | FtsZ | * | | 1 | | | 100 | | | 100 | |
| | MinC | | | | 1 | | | | | | |
| | MinD | | | | 1 | | 75 | | | 75 | |
| | MinE | | | | 1 | | | | | | |
| | ZapA | | 1 | | | | 100 | | | 100 | |
| REPLICATION | DNA helicase | | | | | | 25 | 50 | 7 | 25 | |
| | Helicase-1 | | 1 | 1 | | | 100 | 50 | | 92 | 3 |
| | Helicase-2 | | | | | | | | | 8 | 3 |
| | DNA repair | | | | | | | | | 8 | |
| | RepA | | 2 | 2 | | | | | 43 | | 42 |
| | RepAEB | | | | 1 | | | 100 | 7 | | 15 |
| | RepC | | 1 | | | | 50 | | 43 | 8 | 24 |
| | RuvB | | 1 | | | | 100 | | 7 | 92 | |
| PARTITION | ATPase, TyrK, ExoP | | 2 | | | | 100 | 50 | 14 | 75 | 15 |
| | ParA, ParM | | | 1 | 1 | | | 100 | 100 | 8 | 88 |
| | ParB | | | | | | 75 | 100 | 93 | 25 | 64 |
| | plasmid dimer resolution | | | | | | | | 14 | | 15 |
| | Serine recombinase | | 3 | | | | 25 | | 50 | 50 | 30 |
| | Tyrosine recombinase | | | | | | 25 | | 7 | 33 | 6 |
| | Xer-like tyrosine recombinase | | | 1 | | | 25 | 50 | | 33 | 6 |
| | XerD | | | | | | 25 | | | | |
| MAINTENANCE | Ccd (PSK) | | | | | | | | 7 | | |
| | HicAB (PSK) | | | | | | | | 7 | | |
| | HigBA (PSK) | | 2 | | | | | | 14 | 17 | 6 |
| | MazEF (PSK) | | | | | | | | 7 | | 6 |
| | ParDE (PSK) | | 2 | 6 | | | 25 | 50 | 7 | 8 | 15 |
| | PhD, Doc (PSK) | | | | | | | | 7 | | |
| | RelBE (PSK) | | 1 | | | | | | | | 3 |
| | VapBC/Vag (PSK) | | 3 | 2 | | | | | 14 | 8 | 9 |

*Left hierarchy: KEGG entry (chromosome-like) covers REPLICATION, PARTITION, SEGREGATION, CELL DIVISION; ACLAME family (plasmid-like) covers REPLICATION, PARTITION, MAINTENANCE.*

We propose that the requirement for maintenance system compatibility between co-

occurring replicons is the driving force behind the presence of plasmid-type replication

initiation and maintenance systems in bacterial SERs. Indeed, genes encoding

chromosome-like replication initiators (DnaA) are hardly found on SERs. When they

are, in *Paracoccus denitrificans*, *Prevotella intermedia* and *P. melaninogenica*, the

annotated chromosome in the corresponding genome does not carry one. Similarly,

chromosomal centromeres (*parS*) are found on a single replicon within a multipartite

genome, which is the chromosome in all genomes but one. In *P. intermedia*

(GCA_000261025.1), both replication initiation and partition systems define the SER as

the *bona fide* chromosome and the annotated chromosome as an extra-chromosomal

replicon. The harmonious coexistence of different replicons in a cell requires that they

use divergent enough maintenance systems. In the advent of a chromosome fission or

duplication, the involvement of an autonomously self-replicating element different from

the chromosome is mandatory to provide one of the generated DNA molecules with a

(non-chromosomal) maintenance machinery.

'Plasmid-first' and 'chromosome-first' hypotheses can be reconciled into a unified,

general Fusion-Shuffling-Scission model of SER emergence where a chromosome and a

plasmid combine into a cointegrate (Fig. 6). Plasmids are known to merge or to integrate

chromosomes in both experimental settings (Brom et al., 2000 ; Guo et al., 2003;

Iordănescu, 1975 ; Sýkora, 1992) and the natural environment (Cervantes et al., 2011;

Naito et al., 2016; Sýkora, 1992), as are the SER and chromosome of a multipartite

genome (Val et al., 2014; Xie et al., 2017; Yamamoto et al., 2018). When integrated, the

plasmids/SERs can thus replicate with the chromosome and persist in the bacterial

lineage through several generations (Cervantes et al., 2011; Val et al., 2014; Xie et al.,

2017). The co-integrate may resolve into its original components (Guo et al., 2003; Val

et al., 2014) or give rise to novel genomic architectures (Guo et al., 2003; Cervantes et

al., 2011; Val et al., 2014). The co-integration state likely facilitates inter-replicon gene

exchanges and genome rearrangements that may lead to the translocation of large

chromosome fragments to the resolved plasmid (Guo et al., 2003; Val et al., 2014).

Multiple cell divisions, and possibly several merging-resolution rounds, could provide

time and opportunity for the plasmid-chromosome re-assortment to take place, and for

multiple essential replicons and a viable distributed genome to form ultimately. In the

novel genome, one ER retains the chromosome-like origin of replication and

centrosome, and the other the plasmidic counterparts. The novel ERs differ from the

chromosome and plasmid that gathered in the progenitor host at the onset. They thus

constitute neo-chromosomes that carry divergent maintenance machineries and can

cohabit and function in the same cell. Depending on the number of cell cycles spent as

co-integrate, the level of genome reorganization, the acquisition of genetic material and

the environmental selective pressure acting upon the host, the final essential replicons

may exhibit diverse modalities of genome integration (Figure 6).

**Figure 6. Fusion-Shuffling-Scission model of distributed genome evolution**
Origins of replication are represented by diamonds.

The Fusion-Shuffling-Scission model of genome evolution that we propose accounts for

the extreme plasticity met in distributed genomes and the eco-phenotypic flexibility of

their hosts. Indeed, having a distributed genome appears to extend and accelerate the

exploration of the genome evolutionary landscape, producing complex regulation

(diCenzo et al., 2018; Galardini et al., 2015; Jiao et al., 2018) and leading to novel eco-

phenotypes and species diversification (*e.g.*, Burkholderiaceae and Vibrionaceae).

Furthermore, this model may explain the observed separation of the replicons according

to taxonomy. Chromosomes and plasmids thus appear as extremes on a continuum of a

lineage-specific genetic material.

**MATERIALS AND METHODS**

To understand the relationships between the chromosomal and plasmidic replicons, we

focused on the distribution of Inheritance System (IS) genes for each replicon and built

networks linking the replicons given their IS functional orthologues (Fig. 2).

**Retrieval of IS functional homologues**

A sample of proteins involved in the replication and segregation of bacterial replicons

and of the bacterial cell cycle was constructed using datasets available from the

ACLAME (Leplae et al., 2010) and KEGG (Kanehisa et al., 2012) databases. Gene

ontologies for "replication", "partition", "dimer resolution", and "genome maintenance"

(Table 11) were used to select related ACLAME plasmid protein families (Table 1)

using a semi-automated procedure.

**Table 11. Gene ontologies related to plasmid ISs used to select groups of orthologous**
**proteins from the ACLAME database**

| PROCESS | ONTOLOGY | DESCRIPTION |
|---|---|---|
| | go:0006270 | DNA replication initiation |
| | phi:0000268 | plasmid vegetative DNA replication |
| | go:0003896 | DNA primase activity |
| | go:0003887 | DNA-directed DNA polymerase activity |
| Replication | go:0045020 | error-prone DNA repair |
| | go:0006260 | DNA replication |
| | phi:0000114 | DNA helicase activity |
| | go:0006281 | DNA repair |
| | phi:0000196 | plasmid copy number control |
| | go:0003677 | DNA binding |
| | 575 | plasmid partitioning protein family ParB/Spo0J |
| | go:0015616 | DNA translocase activity |
| | 576 | plasmid partitioning protein family ParM |
| | go:0000146 | microfilament motor activity |
| Partition | go:0007059 | chromosome segregation |
| | go:0015616 | DNA translocase activity |
| | go:0007059 | chromosome segregation |
| | go:0016887 | ATPase activity |
| | go:0030541 | plasmid partitioning |
| | go:0051302 | regulation of cell division |
| | phi:0000196 | plasmid copy number control |
| | phi:0000134 | site specific DNA excision |
| | phi:0000144 | serine based recombinase activity |
| | phi : 0000131 | site specific DNA recombinaison |
| Dimer resolution | phi : 0000143 | Tyrosine-based recombinase activity |
| | phi : 0000304 | plasmid dimer resolution |
| | go : 0015616 | DNA translocase activity |
| | phi:0000136 | transpositional recombination |
| | go : 0016740 | transferase activity |
| | phi : 0000262 | toxin |
| | phi:0000322 | PSK |
| | 547 | TA family parDE |
| Maintenance | 544 | TA family epsilon zeta |
| | go:0009008 | DNA methyltransferase activity |
| | phi : 0000264 | nucleoid associated protein |
| | go : 0006276 | plasmid maintenance |

KEGG orthology groups were selected following the KEGG BRITE hierarchical

classification (Table 2). Then, the proteins belonging to the relevant 92 ACLAME

protein families and 71 KEGG orthology groups (3,847 and 43,757 proteins,

respectively) were retrieved and pooled. Using this query set amounting to a total of

47,604 proteins, we performed a *blastp* search of the 6,903,452 protein sequences

available from the 5,125 complete sequences of bacterial replicons downloaded from

NCBI Reference Sequence database (RefSeq) (Pruitt et al., 2007) on 30/11/2012. We

identified 358,624 putative homologues using BLAST default parameters (Camacho et

al., 2009) and a $10^{-5}$ significance cut-off value. We chose this *E*-value threshold to

enable the capture of similarities between chromosome and plasmid proteins whilst

minimizing the production of false positives, *i.e.*, proteins in a given cluster exhibiting

small *E*-values despite not being functionally homologous. Using RefSeq ensured the

annotation consistency of the genomes included in our dataset.

**Clustering of IS functional homologues**

Using this dataset, we inferred clusters of IS functional homologues by coupling of an

*all-versus-all blastp* search using a $10^{-2}$ *E*-value threshold and a TRIBE-MCL (Enright et

al., 2002) clustering procedure. As input to TRIBE-MCL, we used the matrix of log

transformed *E*-value, $d(p_i, p_j) = -\log_{10}\left(e_{value}(p_i, p_j)\right)$, obtained from the comparisons

of all possible protein pairs. Using a granularity value of 4.0 (see below), we organized

the 358,624 IS homologues into 7013 clusters, each comprising from a single to 1990

proteins (Figure 3). We annotated IS homologues according to their best match (BLAST

hit with the lowest *E*-value) among the proteins of the query set, *i.e.*, according to one of

the 117 functions of the query set (71 from KEGG and 46 from ACLAME). Then, we

named the clusters of functional homologues using the most frequent annotation among

the proteins in the cluster. We used the number of protein annotations in a cluster to

determine the cluster quality, a single annotation being optimal. To select the best

granularity and to estimate the consistency of the clusters in terms of functional

homologues, we computed the weighted Biological Homogeneity Index (*wBHI*,

modified from the *BHI* (Datta and Datta, 2006), each cluster being weighted by its size)

and the Conservation Consistency Measure (*CCM*, similar to the *BHI* but using the

functional domains of the proteins to define the reference classes), which both take into

account the size distribution of the clusters (See next paragraph for details on index

calculation). The former gives an estimation of the overall consistency of clusters

annotations according to the protein annotations whereas the latter gives an estimation of

cluster homogeneity according to the protein domains identified beforehand. To build

the sets of functional domains, we performed an *hmmscan* (Finn et al., 2011) procedure

against the Pfam database (Finn et al., 2016) of each of the 358,624 putative IS

homologues. We annotated each protein according to the domain match(es) with *E*-value

$< 10^{-5}$ (individual *E*-value of the domain) and *c-E*-value $< 10^{-5}$ (conditional *E-* value that

measures the statistical significance of each domain). If two domains overlapped, we

only considered the domain exhibiting the smallest *E*-value. We estimated *wBHI* and

*CCM* indices for the clustering of the IS homologues and compared with values obtained

for random clusters simulated according to the cluster size distribution of the IS proteins,

irrespective of their length or function. For each of the clustering obtained for different

granularities, we constructed a random clustering following the original cluster size

distribution (assessed with a $\chi^2$ test) and composed with simulated proteins according to

the distributions of the type and number of functional domains of the data collected from

the 358,624 IS homologues. Overall, the clusters obtained using a granularity of 4.0 with

the TRIBE-MCL algorithm appeared to be homogenous in terms of proteins similarities

toward their best BLAST hits and their functional domain distributions (see below).

**Evaluation of the clustering procedures**

In order to select the best granularity and to estimate the consistency of the clusters in

terms of functional homologs, we computed the *weighted Biological Homogeneity Index*

(*wBHI*) and the *Conservation Consistency Measure* (*CCM*). The former gives an

estimate of the overall consistency of clusters annotations according to the protein

annotations whereas the latter gives an estimate of cluster homogeneity according to

protein domains identified beforehand. Although close to the *Biological Homogeneity*

*Index* (*BHI*) introduced by Datta and Datta (2006), both these indices take into account

the size distribution of the clusters.

The *BHI* was originally introduced to measure the biological homogeneity of clusters

according to reference classes to evaluate clusters obtained with microarray data (Datta

and Datta, 2006). Given a clustering $C=\{C_1,...,C_k\}$ of $k$ clusters with $n_i$ the size of the

cluster $C_{i,}$, a set of $m$ proteins $P=\{P_1,...,P_m\}$ and a set $r$ of reference classes $R$ where each

class $R_i$ could be linked to the $m$ proteins, the BHI is defined as:

$$BHI = \frac{1}{k} \sum_{i=0}^{i<k} c_i$$

609  where $c_i$ is defined as:

$$c_i = \frac{1}{(n_i(n_i-1))} \sum_{P_i,P_j \in C_i} d(P_i,P_j)$$

where $d(P_i,P_j)=1$ if $P_i$ and $P_j$ share at least one common reference class, and $d(P_i,P_j)=0$

otherwise. The reference classes here are the annotations defined according to the

protein best BLAST hit. The *BHI* is thus an easy-to-interpret measure, which value is

maximal when, for all clusters, all the proteins in a cluster share at least one annotation.

The *wBHI* is a modification of the *BHI*, where each cluster is weighted by its size $m$.

Following the previous notation scheme, the *wBHI* is defined as:

$$wBHI = \frac{1}{m} \sum_{i=0}^{i<k} 2.c_i.n_i$$

616    The *CCM* is similar to the *BHI* but the functional domains of the proteins are used to

617    define the reference classes. The distance between the proteins is here computed as the

618    Jaccard distance between the functional domain sets of the proteins. Every protein $P_i$ can

619    be described as a vector of functional domains, $D_{Pi} = \{d_1,...,d_x\}$. The Jaccard distance

620    between the two sets of domains $d_2 (P_i, P_j)$ can be defined as:

$$d_2(P_1, P_2) = 1 - \frac{|D_{P_1} \cap D_{P_2}|}{|D_{P_1} \cup D_{P_2}|}$$

621    where $D_{P_1}$ and $D_{P_2}$ are the clans or domains (when no clan could be assigned) identified

622    in $P_1$ and $P_2$ respectively. For a given cluster $C_i$, the *CCM* is calculated as:

$$CCM = \frac{1}{m} \sum_{i=0}^{i<k} 2.c'_i.n_i$$

623    where $c'_i$ is defined as:

$$c'_i = \frac{1}{(n_i(n_i-1))} \sum_{P_i, P_j \in C_i} d_2(P_i, P_j)$$

624    Clusters which proteins have similar domains result in a *CCM* value close to 0, whereas

625    a *CCM* value close to 1 indicates that the clusters hold proteins with little domain

626    overlap.

**Choice of the clustering granularity**

We tested several levels of granularity to optimize the TRIBE-MCL clustering and

obtain the most informative IS clustering in terms of functional linkage. Too low a

granularity would produce large clusters containing multiple functional families. In turn,

increasing the granularity results in the tightening of the cluster. A high granularity tends

to split clusters harboring different protein subfamilies (*e.g.,* a cluster composed of

proteins from the tyrosine recombinase superfamily) and to produce multiple clusters of

proteins belonging to a single function family according to their level of sequence

dissimilarity. Furthermore, too high a granularity would result in the formation of

numerous single protein clusters, and would dramatically increase the computation times

of the following analyses. A granularity level of 4.0 constituted a good compromise

(Figure 8). Values of *CCM* and *BHI* are slightly improved compared to granularities of

2.0 and 3.0, and the high but still workable number of clusters is expected to prevent the

formation of clusters mingling distinct protein subfamilies.

**Figure 8. Influence of granularity on the clustering**

(a) Number of clusters with more than one protein (dark diamonds) or clusters holding a single protein (pale
diamonds). (b) *BHI* (dark), *wBHI* (pale) and *CCM* (medium) scores obtained with random clusters (squares) and
normal clusters (circles), respectively.

**Assessment of the homogeneity of IS functional homologues**

The homogeneity towards the functions of the proteins in the query set relied on the

assumption that the first BLAST cut-off ($10^{-5}$ *E*-value) was stringent enough to capture

only functional homologues to the query proteins. Potential bias might nevertheless arise

from query proteins possessing a supplementary functional domain unrelated to the IS

role, or from the selection of proteins belonging to the same superfamily but differing in

function. To address these issues, we calculated the functional vectors associated to each

KEGG group or ACLAME family of the query set, as well as those for all obtained

clusters. For a protein $P_i$, we defined the associated functional vector with respect to its

set of identified domains $D_{P_i}$ and to the set of all identified domains $D=\{d_1,...,d_X\}$ as:

$$v_{P_i} = \left( n_{d_1}^{P_i}, ... , n_{d_x}^{P_i} \right)$$

655 where $n_{d_i}^{P_i}$ is the number of time $d_i$ is found in $D_{P_i}$. The functional vector associated to a given

656 cluster of proteins $C_i$ could then be defined as:

$$v_{C_i} = \left( n_{d_1}^{C_i}, ... , n_{d_x}^{C_i} \right)$$

657 where $n_{d_i}^{C_i}$ is defined as:

$$n_{d_i}^{C_i} = \frac{1}{|C_i|} \sum_{P_i \in C_i} n_{d_x}^{P_j}$$

658 For each cluster $C_0$, the cosine distance between its associated vector $v_{C_0}$ and the associated

659 vector $v_{C_a}$ of the corresponding KEGG group or ACLAME family annotations $C_a$ was then

660 computed as:

$$d_{cosine}\left(v_{C_a}, v_{C_0}\right) = 1 - \frac{\sum_{i=1}^{X} n_{d_i}^{C_0} . n_{d_i}^{C_a}}{\sqrt{\sum_{i=1}^{X} n_{d_i}^{C_0}{}^2} . \sqrt{\sum_{i=1}^{X} n_{d_i}^{C_a}{}^2}}$$

661 For each cluster $C_0$, the cosine distance between its associated vector $v_{C_0}$ and the

662 associated vector $v_{C_a}$ of the corresponding KEGG group or ACLAME family

663 annotations $C_a$ was then computed as:

$$d_{cosine}\left(v_{C_a}, v_{C_0}\right) = 1 - \frac{\sum_{i=1}^{X} n_{d_i}^{C_0} . n_{d_i}^{C_a}}{\sqrt{\sum_{i=1}^{X} n_{d_i}^{C_0}{}^2} . \sqrt{\sum_{i=1}^{X} n_{d_i}^{C_a}{}^2}}$$

The $d_{cosine}\left(v_{C_a}, v_{C_0}\right)$ values were compared with those obtained using random clusters

$C_r$ of the same size than $C_0$. For each $C_0$ and its corresponding $C_a$, 200 random clusters

and their associated distances $d_{cosine}\left(v_{C_a}, v_{C_r}\right)$, from which the corresponding empirical

distribution $D_e$ was constructed, were computed. $C_0$ is then considered as noise if

$d_{cosine}(v_{C_a}, v_{C_0}) \notin Q_{10\%}^{D_e}$ where $Q_{10\%}^{D_e}$ is the 0.1-quantile of $D_e$.

**Unsupervised analyses of the replicon space**

We represented the bacterial replicons (Supplementary Table 1) as vectors according to

their content in IS genes. The number of IS protein clusters retained for the analysis

determined the vector dimension and the number of proteins in a replicon assigned to

each cluster gave the value of each vector component. We built matrices

$P = \begin{bmatrix} p_{1,1} & \cdots & p_{1,m} \\ \vdots & \ddots & \vdots \\ p_{n,1} & \cdots & p_{n,m} \end{bmatrix}$, where $n$ is the number of replicons, $m$ the number of protein

clusters, and $p_{i,j}$ the number of proteins of the $j^{th}$ cluster encoded by a gene present on the

$i^{th}$ replicon. We constructed several datasets to explore both the replicon type and the

host taxonomy effects on the separation of the replicons in the analyses (Table 12).

**Table 12. Reference classes used in the evaluation of the replicon IS protein-based**
**unsupervised clustering solutions**

| EVALUATED SEPARATION | ENSEMBLE | NORMALIZED ENSEMBLE [a] |
|---|---|---|
| Chromosomes *vs.* Plasmids | $\left\{ R^{\{chromosome\}}, R^{\{plasmid\}} \right\}$ | $\left\{ \overline{Kl}_{genus}^{R^{\{chromosome\}}}, \overline{Kl}_{genus}^{R^{\{plasmid\}}} \right\}$ |
| Chromosomes *per* host phylum | $Kl_{phylum}^{chromosome}$ | $\overline{Kl}_{genus}^{K}\| K \in Kl_{phylum}^{chromosome}$ |
| Chromosomes *per* host class | $Kl_{class}^{chromosome}$ | $\overline{Kl}_{genus}^{K}\| K \in Kl_{class}^{chromosome}$ |
| Plasmids *per* host phylum | $Kl_{phylum}^{plasmid}$ | $\overline{Kl}_{genus}^{K}\| K \in Kl_{phylum}^{plasmid}$ |
| Plasmids *per* host class | $Kl_{class}^{plasmid}$ | $\overline{Kl}_{genus}^{K}\| K \in Kl_{class}^{plasmid}$ |

[a] Normalisation according to host genus

The taxonomic representation bias was taken into account by normalizing the data with

regard to the host genus: a consensus vector was built for each bacterial genus present in

the datasets. The value of each vector attribute was calculated as the mean of the

attribute values in the vectors of the replicons that belong to the same bacterial genus.

As a first approach, we transformed data into bipartite graphs whose vertices are the

replicons and the proteins clusters. The graphs were spatialized using the force-directed

layout algorithm ForceAtlas2 (Jacomy et al., 2014) implemented in Gephi (Bastian et al.,

2009). Bipartite graphs are a powerful way of representing the data by naturally drawing

the links between the replicons while enabling the detailed analysis of the IS cluster-

based connections of each replicon by applying forces to each node with regard to its

connecting edges. To investigate further the IS-based relationships of the replicons, we

applied the community structure detection algorithm INFOMAP (Rosvall and

Bergstrom, 2008) using the *igraph* python library (Csardi and Nepusz, 2006). We also

performed a WARD hierarchical clustering (Johnson, 1967) after a dimension reduction

of the data using a Principal Component Analysis (Hotelling, 1933). To select an

optimal number of principal components, we relied on the measurements of the cluster

stabilities using a *stability criterion* (Hennig, 2007) and retained the first 30 principal

components (57% of the total variance). For consistency purpose, the number of clusters

in the WARD analysis was chosen to match that obtained with the INFOMAP

procedure. The number of clusters used was assessed by the stability index by Fang and

Wang (2012) (Table 3). The quality of the projection and clustering results were

confirmed using the V-measure indices (Rosenberg and Hirschberg, 2007) (*homogeneity*,

*completeness*, *V-measure*) as external cluster evaluation measures (Table 3). The

*homogeneity* indicates how uniform clusters are towards a class of reference. The

*completeness* indicates whether reference classes are embedded within clusters. The *V-*

*measure* is the harmonic mean between these two indices and indicates the quality of a

clustering solution relative to the classes of reference. These three indices vary between

0 and 1, with values closest to 1 reflecting the good quality of the clustering solution.

The type of replicons (*i.e.,* plasmid or chromosome) and the taxonomic affiliation

(phylum or class) for chromosomes or plasmids were used as references classes (Table

12). Additionally, the *stability criterion* (Hennig, 2007) of individual clusters, weighted

by their size, for a given clustering result was evaluated using the bootstrapping of the

original dataset as re-sampling scheme. Individual Jaccard coefficient for each replicon

were computed as the number of times that a given replicon of a cluster in a clustering

solution is also present in the closest cluster in the resampled datasets.

**Functional characterization of the replicons and genomes**

In order to characterize the functional bias of the replicons, 117 IS functionalities (46

from ACLAME and 71 from KEGG) were considered. When equivalent in plasmids and

chromosomes, functions from ACLAME and KEGG databases were considered to be

distinct. A *n\*m* matrix $F = \begin{bmatrix} f_{1,1} & \cdots & f_{1,m} \\ \vdots & \ddots & \vdots \\ f_{n,1} & \cdots & f_{n,m} \end{bmatrix}$ with *n* the number of replicons and *m* the

number of IS functionalities, was used as input to the projection algorithms. $f_{i,j}$

represents the number of times that genes coding for proteins annotated with the $j^{th}$

function are present on the $i^{th}$ replicon. Several datasets were analysed using PCA

dimension reduction of the data followed by WARD hierarchical clustering (Table 3).

**Logistic regression analyses**

Several reference classes of replicons and complete genomes were considered for

comparison (Table 13). Ambiguous, *i.e.*, potentially adapted, plasmids belonging to

INFOMAP clusters of plasmid replicons partially composed of SERs and/or

chromosomes were removed from the plasmid class. When appropriate, the taxonomic

representation bias was taken into account by normalizing the data with regard to the

host genus as before. Logistic regressions (McCullagh and Nelder, 1989) were

performed for the 117 IS functions using the R glm package coupled to the python

binder rpy2. The computed $P_{value}$ measured the probability of a functionality to be

predictive of a given group of replicons/genomes and the *Odd-Ratio* estimated how the

functionality occurrence influenced the belonging of a replicon/genome to a given

group.

**Table 13.    Datasets used in the logistic regression analyses**

| ENSEMBLE OF REPLICONS OR GENOMES | NOTATION | DATASET | DIMENSION[a] |
|---|---|---|---|
| Genus-normalized SERs | $E_{SER}$ | $\bar{V}_{f,genus}^{R^{\{SER\}}}$ | (28, 117) |
| Genus-normalized plasmids | $E_{plasmid}$ | $\bar{V}_{f,genus}^{R^{\{plasmid\}}}$ | (262, 117) |
| Genus-normalized chromosomes | $E_{chromosome}$ | $\bar{V}_{f,genus}^{R^{\{chromosome\}}}$ | (560, 117) |

[a]  (Number of replicons, number of functions)

**Supervised classification of replicons and genomes**

In order to identify putative ill-defined SERs and chromosomes amongst plasmids, we

performed supervised classification analyses using random forest procedures (Geurts et

al., 2006). We used the IS functionalities as the set of features and the whole sets of

chromosomes, plasmids and SER as sets of samples to build four classification studies

(Table 7) and detect SER candidates (plasmids *vs.* SERs) and chromosome candidates

(chromosomes *vs.* SERs or chromosomes *vs.* plasmids). Because of the unbalanced sizes

of the training classes (SERs *vs.* chromosomes and plasmids), iterative sampling

procedures were performed using 1000 random subsets of the largest class, with a size

similar to that of the smallest class. The ensuing results were averaged to build the class

probabilities and relative importance of the variables. We also used the whole set of

plasmids when compared to SERs, to identify more robust SER candidates. The

discarding of plasmids in the iterative procedure increases the classifier sensitivity while

reducing the rate of false negatives by including more plasmid-annotated putative true

SERs, whereas it decreases the classifier precision while increasing the rate of false

positives. The ExtraTreeClassifier (a classifier similar to Random Forest) class from the

Scikit-learn python library (Pedregosa et al., 2011) was used to perform the

classifications, with *K=1000*, *max_feat=sqrt(number of variables)* and *min_split=1*. For

each run, the *feature_importances* and *estimate_proba* functions were used to compute,

respectively, the relative contribution of the input variables and the class probabilities of

replicons/genomes. The statistical probability of a replicon/genome belonging to a class

was calculated as the average predicted class of the trees in the forest. The relative

contribution of the input variables was estimated according to Breiman (2001). The

choices of the number of trees in the forest *K*, the number of variables selected for each

split *max_feat*, and the minimum number of samples required to split an internal node

*min_split* were cross-validated using a *Leave-One-Out* scheme. The performance of the

*Extremely-randomized-trees* classification procedures was assessed using a stratified 10-

fold cross-validation procedure following Han *et al.* (2012), and the out-of-bag estimate

(OOB score) (Izzenman, 2008; Pedregosa et al., 2011) computed using the *oob_score*

function of Scikit-learn python library.

**Data availability**

The data supporting the findings of this study are available within the Article and

its Supplementary Information or are available from the authors.

**REFERENCES**

Acosta-Cruz E, Wisniewski-Dyé F, Rouy Z, Barbe V, Valdés M, Mavingui P. 2012.

Insights into the 1.59-Mbp largest plasmid of *Azospirillum brasilense* CBG497.

*Archives in Microbiology*. **194**:725-736. doi: 10.1007/s00203-012-0805-2.

Alav I, Sutton JM, Rahman KM. 2018. Role of bacterial efflux pumps in biofilm

formation. *Journal of Antimicrobial Chemotherapy*. **73**:2003-2020. doi:

10.1093/jac/dky042.

Anes J, McCusker MP, Fanning S, Martins M. 2015. The ins and outs of RND efflux

pumps in Escherichia coli. *Frontiers in Microbiology*. **6**:587. doi:

10.3389/fmicb.2015.00587.

Baek JH, Chattoraj DK. 2014. Chromosome I controls chromosome II replication in

*Vibrio cholerae*. *PLOS Genetics*. **10**:e1004184. doi:

10.1371/journal.pgen.1004184.

Barbe V, Bouzon M, Mangenot S, Badet B, Poulain J, Segurens B, Vallenet D, Marlière

P, Weissenbach J. 2011. Complete genome sequence of *Streptomyces cattleya*

NRRL 8057, a producer of antibiotics and fluorometabolites. *Journal of*

*Bacteriology*. **193**:5055-5056. doi: 10.1128/JB.05583-11.

Bastian M, Heymann S, Jacomy M. 2009. Gephi: An open source software for exploring

and manipulating networks. *Third International AAAI Conference on Weblogs*

*and Social Media, San Jose Mc Enery Convention Center, May 17, 2009 - May*

*20, 2009: AAAI Publications*.

Blanco-Ordóñez H, Oliva-García JJ, Pérez-Mendoza D, Soto MJ, Olivares J, Sanjúan J,

Nogales J. 2010. pSymA-dependent mobilization of the *Sinorhizobium meliloti*

pSymB megaplasmid. *Journal of Bacteriology*. **192**:6309-6312. doi:

10.1128/JB.00549-10.

Breiman L. 2001. Random forests. *Machine Learning*. **45**:5-32. doi:
10.1023/A:1010933404324.

Brom S, García-de los Santos A, Cervantes L, Palacios R, Romero D. 2000. In
*Rhizobium etli* symbiotic plasmid transfer, nodulation competitivity and cellular
growth require interaction among different replicons. *Plasmid*. **44**:34-43. doi:
10.1006/plas.2000.1469

Camacho CJ, Coulouris G, Avagyan V, Ma N, Papdopoulos J, Bealer K, Madden TL.
2009. BLAST+: architecture and applications. *BMC Bioinformatics*. **10**:421. doi:
10.1186/1471-2105-10-421.

Casjens SR. 1998. The diverse and dynamic structure of bacterial genomes. *Annual*
*review of genetics*. **32**:339-377. doi: 10.1146/annurev.genet.32.1.339.

Cervantes L, Bustos P, Girard L, Santamaría RI, Dávila G, Vinuesa P, Romero D, Brom
S. 2011. The conjugative plasmid of a bean-nodulating *Sinorhizobium fredii*
strain is assembled from sequences of two *Rhizobium* plasmids and the
chromosome of a *Sinorhizobium* strain. *BMC microbiology*. **11**:149. doi:
10.1186/1471-2180-11-149.

Chen Y, Milam SL, Erickson HP. 2012. SulA inhibits assembly of FtsZ by a simple
sequestration mechanism. *Biochemistry*. **51**:3100–3109. doi: 10.1021/bi201669d.

Chodavarapu S, Jones AD, Feig M, Kaguni JM. 2016. DnaC traps DnaB as an open ring
and remodels the domain that binds primase. *Nucleic Acids Research*. **44**:210-
220. doi: 10.1093/nar/gkv961.

Csárdi G, Nepusz T. 2006. The igraph software package for complex network research.
*InterJournal, Complex Systems*.1695.

Datta S, Datta S. 2006. Methods for evaluating clustering algorithms for gene expression

data using a reference set of functional classes. *BMC Bioinformatics*. **7**:397. doi:

10.1186/1471-2105-7-397.

De Nisco NJ, Abo RP, Wu CM, Penterman J, Walker GC. 2014. Global analysis of cell

cycle gene expression of the legume symbiont *Sinorhizobium meliloti*.

*Proceedings of the National Academy of Sciences of the United States of*

*America*. **111**:3217-3224. doi: 10.1073/pnas.1400421111.

Deghelt M, Mullier C, Sternon JF, Francis N, Laloux G, Dotreppe D, Van der Henst C,

Jacobs-Wagner C, Letesson JJ, De Bolle X. 2014. G1-arrested newborn cells are

the predominant infectious form of the pathogen *Brucella abortus*. *Nature*

*communications*. **5**:4366. doi: 10.1038/ncomms5366.

Demarre G, Galli E, Muresan L, Paly E, David A, Possoz C, Barre F-X. 2014.

Differential management of the replication terminus regions of the two *Vibrio*

*cholerae* chromosomes during cell division. *PLOS Genetics*. **10**:e1004557. doi:

10.1371/journal.pgen.1004557.

diCenzo G, Milunovic B, Cheng J, Finan TM. 2013. The tRNAarg gene and *engA* are

essential genes on the 1.7-Mb pSymB megaplasmid of *Sinorhizobium meliloti*

and were translocated together from the chromosome in an ancestral strain.

*Journal of Bacteriology*. **195**:202-212. doi: 10.1128/JB.01758-12.

diCenzo GC, Finan TM. 2017. The divided bacterial genome: structure, function, and

evolution. *Microbiology and Molecular Biology Reviews*. **81**:e00019-00017. doi:

10.1128/MMBR.00019-17.

diCenzo GC, MacLean AM, Milunovic B, Golding GB, Finan TM. 2014. Examination

of prokaryotic nultipartite genome evolution through experimental genome

reduction. *PLOS Genetics*. **10**:e1004742. doi: 10.1371/journal.pgen.1004742.

diCenzo GC, Wellappili D, Golding GB, Finan TM. 2018. Inter-replicon gene flow

contributes to transcriptional integration in the *Sinorhizobium meliloti*

multipartite genome. *G3 (Bethesda)*. **8**:1711-1720. doi: 10.1534/g3.117.300405.

Dillon SC, Dorman CJ. 2010. Bacterial nucleoid-associated proteins, nucleoid structure

and gene expression. *Nat Rev Microbiol*. **8**:185-195. doi: 10.1038/nrmicro2261.

Drevinek P, Baldwin A, Dowson CG, Mahenthiralingam E. 2008. Diversity of the *parB*

and *repA* genes of the *Burkholderia cepacia* complex and their utility for rapid

identification of *Burkholderia cenocepacia*. *BMC microbiology*. **8**:44. doi:

10.1186/1471-2180-8-44.

Du D, Wang-Kan X, Neuberger A, van Veen HW, Pos KM, Piddock LJV, Luisi BF.

2018. Multidrug efflux pumps: structure, function and regulation. *Nature Review*

*in Microbiology*. **16**:523-539. doi: 10.1038/s41579-018-0048-6.

Dubarry N, Pasta F, Lane D. 2006. ParABS systems of the four replicons of

*Burkholderia cenocepacia*: new chromosome centromeres confer partition

specificity. *Journal of Bacteriology*. **188**:1489-1496. doi:

10.1128/JB.188.4.1489-1496.2006.

Egan ES, Lobner-Olesen A, Waldor MK. 2004. Synchronous replication initiation of the

two *Vibrio cholerae* chromosomes. *Current Biology*. **14**:R501-502. doi:

10.1016/j.cub.2004.06.036.

Egan ES, Waldor MK. 2003. Distinct replication requirements for the two *Vibrio*

*cholerae* chromosomes. *Cell*. **114**:521-530. doi: 10.1016/s0092-8674(03)00611-

1.

Enright AJ, Dongen S, Ouzounis C. 2002. An efficient algorithm for large-scale

detection of protein families. *Nucleic Acids Research*. **30**:1575-1584. doi:

10.1093/nar/30.7.1575.

Fang Y, Wang J. 2012. Selection of the number of clusters via the bootstrap method.

*Computational Statistics & Data Analysis*. **56**:468-477. doi:

10.1016/j.csda.2011.09.003.

Fiebig A, Keren K, Theriot JA. 2006. Fine-scale time-lapse analysis of the biphasic,

dynamic behaviour of the two *Vibrio cholerae* chromosomes. *Molecular*

*Microbiology*. **60**:1164-1178. doi: 10.1111/j.1365-2958.2006.05175.x.

Finn RD, Clements J, Eddy SR. 2011. HMMER web server: interactive sequence

similarity searching. *Nucleic Acids Research*. **39**:W29-37. doi:

10.1093/nar/gkr367.

Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta

M, Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The

Pfam protein families database: towards a more sustainable future. *Nucleic Acids*

*Research*. **44**:D279-285. doi: 10.1093/nar/gkv1344.

Frage B, Dohlemann J, Robledo M, Lucena D, Sobetzko P, Graumann PL, Becker A.

2016. Spatiotemporal choreography of chromosome and megaplasmids in the

*Sinorhizobium meliloti* cell cycle. *Molecular Microbiology*. **100**:808-823. doi:

10.1111/mmi.13351.

Galardini M, Mengoni A, Brilli M, Pini F, Fioravanti A, Lucas S, Lapidus A, Cheng JF,

Goodwin L, Pitluck S, Land M, Hauser L, Woyke T, Mikhailova N, Ivanova N,

Daligault H, Bruce D, Detter C, Tapia R, Han C, Teshima H, Mocali S,

Bazzicalupo M, Biondi EG. 2011. Exploring the symbiotic pangenome of the

nitrogen-fixing bacterium *Sinorhizobium meliloti*. *BMC Genomics*. **12**:235. doi:

10.1186/1471-2164-12-235.

Galardini M, Pini F, Bazzicalupo M, Biondi G, Mengoni A. 2013. Replicon-dependent

bacterial genome evolution: the case of *Sinorhizobium meliloti*. *Genome Biology*

*and Evolution*. **5**:542-558. doi: 10.1093/gbe/evt027.

Galardini M, Brilli M, Spini G, Rossi M, Roncaglia B, Bani A, Chiancianesi M, Moretto

M, Engelen K, Bacci G, Pini F, Biondi EG, Bazzicalupo M, Mengoni A. 2015.

Evolution of Intra-specific Regulatory Networks in a Multipartite Bacterial

Genome. *PLoS Computational Biology* **11**:e1004478. doi:

10.1371/journal.pcbi.1004478.

Galli E, Poidevin M, Le Bars R, Desfontaines JM, Muresan L, Paly E, Yamaichi Y,

Barre FX. 2016. Cell division licensing in the multi-chromosomal *Vibrio*

*cholerae* bacterium. *Nat Microbiology*. **1**:16094. doi:

10.1038/nmicrobiol.2016.94.

Gerding MA, Chao MC, Davis BM, Waldor MK. 2015. Molecular dissection of the

essential features of the origin of replication of the second *Vibrio cholerae*

chromosome. *MBio*. **6**:e00973. doi: 10.1128/mBio.00973-15.

Geurts P, Ernst D, Wehenkel L. 2006. Extremely randomized trees. *Machine Learning*.

**63**:3-42. doi: 10.1007/s10994-006-6226-1.

Guo FB, Ning LW, Huang J, Lin H, Zhang HX. 2010. Chromosome translocation and its

consequence in the genome of *Burkholderia cenocepacia* AU-1054. *Biochemical*

*and Bophysical Research Communications*. **403**:375-379. doi:

10.1016/j.bbrc.2010.11.039.

Guo X, Flores M, Mavingui P, Fuentes SI, Hernandez G, Davila G, Palacios R. 2003.

Natural genomic design in *Sinorhizobium meliloti*: novel genomic architectures.

*Genome Research*. **13**:1810-1817. doi: 10.1101/gr.1260903.

Hall JPJ, Brockhurst MA, Dytham C, Harrison E. 2017. The evolution of plasmid

stability: Are infectious transmission and compensatory evolution competing

evolutionary trajectories? *Plasmid*. **91**:90-95. doi:

10.1016/j.plasmid.2017.04.003.

Han J, Kamber M, Pei J. 2012. Data Mining: Concepts and Techniques, Third Edition:

Morgan kaufmann, Elsevier.

Harrison PW, Lower RP, Kim NK, Young JPW. 2010. Introducing the bacterial

'chromid': not a chromosome, not a plasmid. *Trends in Microbiology*. **18**:141-

148. doi: 10.1016/j.tim.2009.12.010.

Hennig C. 2007. Cluster-wise assessment of cluster stability. *Computational Statistics &*

*Data Analysis*. **52**:258-271. doi: 10.1016/j.csda.2006.11.025.

Hotelling H. 1933. Analysis of a complex of statistical variables into principal

components. *Journal of Educational Psychology*. **24**:417. doi:

10.1037/h0071325.

Izenman AJ. 2008. Modern multivariate statistical techniques: regression, classification,

and manifold learning: Springer-Verlag, New York Inc.

Jacomy M, Venturini T, Heymann S, Bastian M. 2014. ForceAtlas2, a continuous graph

layout algorithm for handy network visualization designed for the Gephi

software. *PLoS One*. **9**:e98679. doi: 10.1371/journal.pone.0098679.

Jiao J, Ni M, Zhang B, Zhang Z, Young JPW, Chan TF, Chen WX, Lam HM, Tian CF.

2018. Coordinated regulation of core and accessory genes in the multipartite

genome of Sinorhizobium fredii. *PLoS Genet*ics **14**:e1007428. doi:

10.1371/journal.pgen.1007428.

Johnson SC. 1967. Hierarchical clustering schemes. *Psychometrika*. **32**:241-254.

Kahng LS, Shapiro L. 2003. Polar localization of replicon origins in the multipartite

genomes of *Agrobacterium tumefaciens* and *Sinorhizobium meliloti*. *Journal of*

*Bacteriology*. **185**:3384-3391. doi: 10.1128/jb.185.11.3384-3391.2003.

Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. 2012. KEGG for integration and

interpretation of large-scale molecular data sets. *Nucleic Acids Research*.

**40**:D109-114. doi: 10.1093/nar/gkr988.

Kemter FS, Messerschmidt SJ, Schallopp N, Sobetzko P, Lang E, Bunk B, Sproer C,

Teschler JK, Yildiz FH, Overmann J, Waldminghaus T. 2018. Synchronous

termination of replication of the two chromosomes is an evolutionary selected

feature   in   Vibrionaceae.   *PLOS   Genetics*.   **14**:e1007251.   doi:

10.1371/journal.pgen.1007251.

Krawiec S, Riley M. 1990. Organization of the bacterial chromosome. *Microbiological*

*Reviews*. **54**:502-539.

Landeta C, Dávalos A, Cevallos MÁ, Geiger O, Brom S, Romero D. 2011. Plasmids

with a chromosome-like role in rhizobia. *Journal of Bacteriology*. **193**:1317-

1326. doi: 10.1128/JB.01184-10.

Lederberg J. 1998. Plasmid (1952-1997). *Plasmid*. **39**:1-9. doi: 10.1006/plas.1997.1320.

Leplae R, Lima-Mendez G, Toussaint A. 2010. ACLAME: a CLAssification of Mobile

genetic Elements, update 2010. *Nucleic Acids Research*. **38**:D57-61. doi:

10.1093/nar/gkp938.

Li Y, Canchaya C, Fang F, Raftis E, Ryan KA, van Pijkeren J-P, van Sinderen D,

O'Toole PW. 2007. Distribution of megaplasmids in *Lactobacillus salivarius* and

other   lactobacilli.   *Journal   of   Bacteriology*.   **189**:6128-6139.   doi:

10.1128/JB.00447-07.

Lioy VS, Cournac A, Marbouty M, Duigou S, Mozziconacci J, Espeli O, Boccard F,

Koszul R. 2018. Multiscale structuring of the *E. coli* chromosome by nucleoid-

associated   and   condensin   proteins.   *Cell*.   **172**:771-783   e718.   doi:

10.1016/j.cell.2017.12.027.

Liu G, Yong MY, Yurieva M, Srinivasan KG, Liu J, Lim JS, Poidinger M, Wright GD,

Zolezzi F, Choi H, Pavelka N, Rancati G. 2015. Gene essentiality is a

quantitative property linked to cellular evolvability. *Cell*. **163**:1388-1399. doi:

10.1016/j.cell.2015.10.069.

Livny J, Yamaichi Y, Waldor MK. 2007. Distribution of centromere-like parS sites in

bacteria: insights from comparative genomics. *Journal of Bacteriology*.

**189**:8693-8703. doi: 10.1128/JB.01239-07.

Lu C, Nakayasu ES, Zhang LQ, Luo ZQ. 2016. Identification of Fic-1 as an enzyme that

inhibits bacterial DNA replication by AMPylating GyrB, promoting filament

formation. *Science Signal*. **9**:ra11. doi: 10.1126/scisignal.aad0446.

Mackenzie C, Kaplan S, Choudhary M. 2004. Multiple chromosomes. *In: Miller RV and*

*Day MJ, editors. Microbial Evolution: Gene Establishment, Survival, and*

*Exchange: ASM press, Washington DC*:82-101.

Mackenzie C, Simmons AE, Kaplan S. 1999. Multiple chromosomes in bacteria: the Yin

and Yang of *trp* gene localization in *Rhodobacter sphaeroides* 2.4.1. *Genetics*.

**153**:525-538.

MacLellan SR, Sibley CD, Finan TM. 2004. Second chromosomes and megaplasmids in

bacteria. *In: Funnel BE and Phillips GJ, editors. Plasmid biology: ASM press,*

*Washington DC*:529-542.

MacLellan SR, Zaheer R, Sartor AL, MacLean AM, Finan TM. 2006. Identification of a

megaplasmid centromere reveals genetic structural diversity within the repABC

family of basic replicons. *Molecular Microbiology*. **59**:1559-1575. doi:

10.1111/j.1365-2958.2005.05040.x.

Maida I, Fondi M, Orlandini V, Emiliani G, Papaleo MC, Perrin E, Fani R. 2014.

Origin, duplication and reshuffling of plasmid genes: Insights from *Burkholderia*

*vietnamiensis* G4 genome. *Genomics*. **103**:229-238. doi:

10.1016/j.ygeno.2014.02.004.

McCullagh P, Nelder JA. 1989. Generalized linear models, Second Edition: Chapman & Hall/CRC, London. 532 p.

Medema MH, Trefzer A, Kovalchuk A, van den Berg M, Muller U, Heijne W, Wu L, Alam MT, Ronning CM, Nierman WC, Bovenberg RA, Breitling R, Takano E. 2010. The sequence of a 1.8-mb bacterial linear plasmid reveals a rich evolutionary reservoir of secondary metabolic pathways. *Genome Biology and Evolution*. **2**:212-224. doi: 10.1093/gbe/evq013.

1003 Meier EL, Daitch AK, Yao Q, Bhargava A, Jensen GJ, Goley ED. 2017. FtsEX-mediated regulation of the final stages of cell division reveals morphogenetic plasticity in *Caulobacter crescentus*. *PLoS Genetics* **13**: e1006999. doi: 10.1371/journal.pgen.1006999.

1007 Million-Weaver S, Camps M. 2014. Mechanisms of plasmid segregation: have multicopy plasmids been overlooked? *Plasmid*. **75**:27-36. doi: 10.1016/j.plasmid.2014.07.002.

1010 Murray H, Errington J. 2008. Dynamic control of the DNA replication initiation protein DnaA by Soj/ParA. *Cell*. **135**:74-84. doi: 10.1016/j.cell.2008.07.044.

1012 Naito M, Ogura Y, Itoh T, Shoji M, Okamoto M, Hayashi T, Nakayama K. 2016. The complete genome sequencing of *Prevotella intermedia* strain OMA14 and a subsequent fine-scale, intra-species genomic comparison reveal an unusual amplification of conjugative and mobile transposons and identify a novel *Prevotella*-lineage-specific repeat. *DNA Research*. **23**:11-19. doi: 10.1093/dnares/dsv032.

Ohtani N, Tomita M, Itaya M. 2010. An extreme thermophile, *Thermus thermophilus*, is a polyploid bacterium. *Journal of Bacteriology*. **192**:5499-5505. doi: 10.1128/JB.00662-10.

Passot FM, Calderon V, Fichant G, Lane D, Pasta F. 2012. Centromere binding and

evolution of chromosomal partition systems in the Burkholderiales. *Journal of*

*Bacteriology*. **194**:3426-3436. doi: 10.1128/JB.00041-12.

Pedregosa F, Weiss R, Brucher M. 2011. Scikit-learn : Machine Learning in Python.

*Journal      of      Machine      Learning      Research*.      **12**:2825-2830.

http://www.jmlr.org/papers/volume12/pedregosa11a/pedregosa11a.pdf.

Petersen J, Frank O, Göker M, Pradella S. 2013. Extrachromosomal, extraordinary and

essential--the plasmids of the *Roseobacter* clade. *Applied Microbiology and*

*Biotechnology*. **97**:2805-2815. doi: 10.1007/s00253-013-4746-8.

Pinto UM, Pappas KM, Winans SC. 2012. The ABCs of plasmid replication and

segregation.  *Nature   Review   in   Microbiology*.   **10**:755-765.   doi:

10.1038/nrmicro2882.

Pruitt KD, Tatusova T, Maglott DR. 2005. NCBI Reference Sequence (RefSeq): a

curated non-redundant sequence database of genomes, transcripts and proteins.

*Nucleic Acids Research*. **33**:D501-504. doi: 10.1093/nar/gki025.

Rasmussen T, Jensen RB, Skovgaard O. 2007. The two chromosomes of *Vibrio cholerae*

are initiated at different time points in the cell cycle. *EMBO Journal*. **26**:3124-

3131. doi: 10.1038/sj.emboj.7601747.

Reyes-Lamothe R, Tran T, Meas D, Lee L, Li AM, Sherratt DJ, Tolmasky ME. 2014.

High-copy bacterial plasmids diffuse in the nucleoid-free space, replicate

stochastically and are randomly partitioned at cell division. *Nucleic Acids*

*Research*. **42**:1042-1051. doi: 10.1093/nar/gkt918.

Rosenberg A, Hirschberg J. 2007. V-Measure: A conditional entropy-based external

cluster evaluation measure. *Proceedings of the 2007 Joint Conference on*

*Empirical Methods in Natural Language Processing and Computational Natural*

*Language Learning (EMNLP-CoNLL)* p. 410-420.
https://www.researchgate.net/publication/221012656_V-
Measure_A_Conditional_Entropy-
Based_External_Cluster_Evaluation_Measure.

Rosvall M, Bergstrom CT. 2008. Maps of random walks on complex networks reveal community structure. *Proceedings of the National Academy of Sciences of the United States of America*. **105**:1118-1123. doi: 10.1073/pnas.0706851105.

San Millan A, Peña-Miller R, Toll-Riera M, Halbert ZV, McLean aR, Cooper BS, MacLean RC. 2014. Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids. *Nature communications*. **5**:5208. doi: 10.1038/ncomms6208.

Slater SC, Goldman BS, Goodner B, Setubal JC, Farrand SK, Nester EW, Burr TJ, Banta L, Dickerman AW, Paulsen I, Otten L, Suen G, Welch R, Almeida NF, Arnold F, Burton OT, Du Z, Ewing A, Godsy E, Heisel S, Houmiel KL, Jhaveri J, Lu J, Miller NM, Norton S, Chen Q, Phoolcharoen W, Ohlin V, Ondrusek D, Pride N, Stricklin SL, Sun J, Wheeler C, Wilson L, Zhu H, Wood DW. 2009. Genome sequences of three *Agrobacterium* biovars help elucidate the evolution of multichromosome genomes in bacteria. *Journal of Bacteriology*. **191**:2501-2511. doi: 10.1128/JB.01779-08.

Srivastava P, Fekete RA, Chattoraj DK. 2006. Segregation of the replication terminus of the two *Vibrio cholerae* chromosomes. *Journal of Bacteriology*. **188**:1060-1070. doi: 10.1128/JB.188.3.1060-1070.2006.

Stalder T, Rogers LM, Renfrow C, Yano H, Smith Z, Top EM. 2017. Emerging patterns of plasmid-host coevolution that stabilize antibiotic resistance. *Scientific Reports*. **7**:4853. doi: 10.1038/s41598-017-04662-0.

Stokke C, Waldminghaus T, Skarstad K. 2011. Replication patterns and organization of

replication forks in *Vibrio cholerae*. *Microbiology*. **157**:695-708. doi:

10.1099/mic.0.045112-0.

Sýkora P. 1992. Macroevolution of plasmids: A model for plasmid speciation. *Journal

of theoretical biology*. **159**:53-65. doi: 10.1016/s0022-5193(05)80767-2.

Val M-E, Kennedy SP, El Karoui M, Bonné L, Chevalier F, Barre F-X. 2008. FtsK-

dependent dimer resolution on multiple chromosomes in the pathogen *Vibrio

cholerae*. *PLOS Genetics*. **4**:e1000201. doi: 10.1371/journal.pgen.1000201.

Val M-E, Kennedy SP, Soler-Bistué AJ, Barbe V, Bouchier C, Ducos-Galand M,

Skovgaard O, Mazel D. 2014. Fuse or die: how to survive the loss of Dam in

*Vibrio cholerae*. *Molecular Microbiology*. **91**:665-678. doi: 10.1111/mmi.12483.

Venkova-Canova T, Chattoraj DK. 2011. Transition from a plasmid to a chromosomal

mode of replication entails additional regulators. *Proceedings of the National

Academy of Sciences of the United States of America*. **108**:6199-6204. doi:

10.1073/pnas.1013244108.

Villaseñor T, Brom S, Davalos A, Lozano L, Romero D, Los Santos AG. 2011.

Housekeeping genes essential for pantothenate biosynthesis are plasmid-encoded

in *Rhizobium etli* and *Rhizobium leguminosarum*. *BMC microbiology*. **11**:66. doi:

10.1186/1471-2180-11-66.

Vuilleumier S, Chistoserdova L, Lee MC, Bringel F, Lajus A, Zhou Y, Gourion B,

Barbe V, Chang J, Cruveiller S, Dossat C, Gillett W, Gruffaz C, Haugen E,

Hourcade E, Levy R, Mangenot S, Muller E, Nadalig T, Pagni M, Penny C,

Peyraud R, Robinson DG, Roche D, Rouy Z, Saenampechek C, Salvignol G,

Vallenet D, Wu Z, Marx CJ, Vorholt JA, Olson MV, Kaul R, Weissenbach J,

Medigue C, Lidstrom ME. 2009. *Methylobacterium* genome sequences: a

reference blueprint to investigate microbial metabolism of C1 compounds from

natural and industrial sources. *PLoS One*. **4**:e5584. doi:

10.1371/journal.pone.0005584.

Webber MA, Bailey AM, Blair JM, Morgan E, Stevens MP, Hinton JC, Ivens A, Wain

J, Piddock LJ. 2009. The global consequence of disruption of the AcrAB-TolC

efflux pump in *Salmonella enterica* includes reduced expression of SPI-1 and

other attributes required to infect the host. *Journal of Bacteriology*. **191**:4276-

4285. doi: 10.1128/JB.00363-09.

Wisniewski-Dyé F, Borziak K, Khalsa-Moyers G, Alexandre G, Sukharnikov LO,

Wuichet K, Hurst GB, McDonald WH, Robertson JS, Barbe V, Calteau A, Rouy

Z, Mangenot S, Prigent-Combaret C, Normand P, Boyer M, Siguier P, Dessaux

Y, Elmerich C, Condemine G, Krishnen G, Kennedy I, Paterson AH, González

V, Mavingui P, Zhulin IB. 2011. *Azospirillum* genomes reveal transition of

bacteria from aquatic to terrestrial environments. *PLOS Genetics*. **7**:e1002430.

doi: 10.1371/journal.pgen.1002430.

Wisniewski-Dyé F, Lozano L, Acosta-Cruz E, Borland S, Drogue B, Prigent-Combaret

C, Rouy Z, Barbe V, Mendoza Herrera A, González V, Mavingui P. 2012.

Genome sequence of *Azospirillum brasilense* CBG497 and comparative analyses

of *Azospirillum* core and accessory genomes provide Insight into niche

adaptation. *Genes (Basel)*. **3**:576-602. doi: 10.3390/genes3040576.

Xie G, Johnson SL, Davenport KW, Rajavel M, Waldminghaus T, Detter JC, Chain PS,

Sozhamannan S. 2017. Exception to the rule: genomic characterization of

naturally occurring unusual *Vibrio cholerae* strains with a single chromosome.

*International Journal of Genomics*. **2017**:8724304. doi: 10.1155/2017/8724304.

Yamaichi Y, Fogel MA, McLeod SM, Hui MP, Waldor MK. 2007. Distinct centromere-

like *parS* sites on the two chromosomes of *Vibrio* spp. *Journal of Bacteriology*.

**189**:5314-5324. doi: 10.1128/JB.00416-07.

Yamamoto S, Lee K-i, Morita M, Arakawa E, Izumiya H, Ohnishi M. 2018. Single

circular chromosome identified from the genome sequence of the *Vibrio*

*cholerae* O1 bv. El Tor Ogawa strain V060002. *Genome Announcements*. **6**. doi:

10.1128/genomeA.00564-18.

Yeoman CJ, Kelly WJ, Rakonjac J, Leahy SC, Altermann E, Attwood GT. 2011. The

large episomes of *Butyrivibrio proteoclasticus* B316T have arisen through

intragenomic gene shuttling from the chromosome to smaller *Butyrivibrio*-

specific plasmids. *Plasmid*. **66**:67-78. doi: 10.1016/j.plasmid.2011.05.002.

**SUPPLEMENTARY TABLES**

Table 1. Replicon dataset

Table 2. INFOMAP IS protein-based clustering solution of the 4928 replicons

Table 3. PCA + WARD IS protein-based clustering solution of the 4928 replicons

Table 4. PCA + WARD IS function-based clustering solution of the 4928 replicons