A plant biodiversity effect resolved to a single genetic locus

Samuel E. Wuest[1,2,3*] and Pascal A. Niklaus[2]

1) Department of Plant and Microbial Biology and Zurich-Basel Plant Science Center, University of

Zurich, Zollikerstrasse 107, 8008 Zurich, Switzerland

2) Department of Evolutionary Biology and Environmental Studies and Zurich-Basel Plant Science

Center, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

3) current address: Buchenweg 5, 5436 Würenlos, Switzerland

*correspondence to: Samuel E. Wuest (samuel.wuest@ieu.uzh.ch)

Summary

Despite extensive evidence that biodiversity promotes plant community productivity, progress

towards understanding the mechanistic basis of this effect remains slow, impeding the development

of predictive ecological theory and agricultural applications. Here, we analysed non-additive

interactions between genetically divergent Arabidopsis accessions in experimental plant

communities. By combining methods from ecology and genetics, we identified a major effect locus

at which allelic differences between individuals increases above-ground productivity of

communities. In experiments with near-isogenic lines, we show that this diversity effect acts

independently of other genomic regions and can be resolved to a single region representing less

than 0.3% of the genome. Using plant-soil-feedback experiments, we also demonstrate that allelic

diversity causes genotype-specific soil legacy responses in a consecutive growing period, even after

the original community has disappeared. Our work thus suggests that positive diversity effects can

be linked to single Mendelian factors, and that a range of complex community properties can have a

simple cause. This may pave the way to a better understanding of diversity effects, and to novel

breeding strategies, focussing on phenotypic properties that manifest themselves beyond isolated

individuals, i.e. at a higher level of biological organisation.

Main Text

More than two decades of plant ecological research and the publication of hundreds of studies have

firmly established that positive biodiversity effects, in particular on community yield, are the rule

rather than exception and are often substantial[1,2]. These positive effects of biological diversity on

community functioning have been explained by larger community-level resource use promoted by

niche complementarity, facilitation, or by reduced negative density-dependent effects of enemies[3–5].

Yet our understanding of the particular driving mechanisms remains poor, for several reasons. First,

diversity effects are emergent properties that only manifest in comparisons of communities differing

in diversity[6]. Second, diversity effects, and the mechanisms that drive these, may change with

environmental conditions[7,8]. Third, while there is no doubt that functional trait differences underly

biodiversity effects[9], trait-based analyses remain to some degree phenomenological because

evolutionary forces have led to the formation of trait syndromes, i.e. to sets of highly correlated

traits that reflect fundamental trade-offs between ecological strategies[10,11]. The observed variation in

traits thus is confounded with evolutionary history[12], and it remains almost impossible to distinguish

the traits that are true drivers of biodiversity effects from traits that are merely correlated. It

therefore remains difficult to develop predictive ecological theory and apply it, for example, in

breeding and agriculture.

Most biodiversity research to date has focused on variation among species, but experimental[13],

theoretical[14], and observational[15] studies have shown that positive diversity effects on productivity

also occur at levels of organization above (e.g. landscape level) and below species (e.g. genotype

level). A substantial part of the trait variation apparent in plant communities occurs within species[16],

and increased intra-specific variation can have similar effects as inter-specific trait variation in low-

diversity systems[13,17,18]. Despite qualitative differences, there may therefore be commonalities of

trait variation within and between species with respect to effects and mechanisms, indicating that

studies at the genotype level may provide some insights into effects of species-level variation and

vice versa. A methodological advantage of intra-specific biodiversity studies is that genetic methods

can circumvent some of the problems encountered in species-level diversity studies. Specifically,

crosses between genotypes allow trait variation among individuals to be re-arranged[19] without

confounding with population structure or differentiation into ecological strategies. However, genetic

approaches are normally used to study properties of individuals rather than emergent properties of

communities. Here, we demonstrate in a case study how the genetic approach can be harnessed to

identify the genetic underpinnings of biodiversity effects.

Results

Using model plant communities (Fig. 1a), we screened ten pair-wise mixtures of genetically

divergent natural accessions of *Arabidopsis thaliana* for which mapping populations are publicly

available (Methods). Mixture communities of the two genetically divergent accessions Bayreuth

(Bay) and Shadhara (Sha) reproducibly exhibited positive net biodiversity effects, i.e. mixtures

produced a higher community-level shoot biomass than the average of their monocultures. This

depended on soil conditions, with effects that were essentially absent on peat-rich soil but grew to

an overyielding of 16% with increasing amounts of sand in the substrate (sand content × diversity:

$F_{1,160}= 4.57$, ANOVA $P < 0.05$; Fig. 1b and Supplementary Figure 1a). Analysis by additive

partitioning[6] revealed that these community-level biodiversity effects were due to complementarity

rather than selection effects (sand content × complementarity effect, $F_{1,77} = 7.21$, ANOVA $P < 0.01$;

Supplementary Figure 1b). Specifically, in our study both accessions benefited from growing in

mixed communities.

To analyse the genetic basis of the positive diversity effect in mixed Bay-Sha communities, we

performed quantitative trait locus (QTL) mapping using publicly available recombinant inbred lines

(RILs). These RILs are largely homozygous (Fig. 1c) and have been derived from a cross between

the Bay and Sha accessions, followed by multiple subsequent rounds of selfing[20]. For efficient

mapping, we capitalized on a so called competition diallel. Traditional diallel designs systematically

cross sets of parental lines realizing all possible combinations and are used in breeding to determine

the genetic basis of traits; specifically, diallel analysis partitions the trait variation of crosses into

additive contributions[21] of parental lines (general combining abilities; GCA) and cross-specific

effects (specific combining abilities; SCA), with the latter interpreted as consequences of

dominance or epistasis. By substituting individuals and crosses with communities and mixtures, the

principle of diallels can be applied to the analyses of biodiversity effects in communities[22], which

we did here (Fig. 1c, d). In this context, the distinction between maternal and paternal effects ceases

to apply, simplifying the design to a half-diallel. SCAs then quantify the deviations of mixture

yields from expectations based on additive average contributions of the two genotypes. We

combined 18 RILs and the two parental accessions Bay and Sha in a diallel, in four replicate blocks,

on sand-rich soil. We detected significant positive genotype diversity effects on above-ground

biomass production (Fig. 2a, $F_{1,189}$= 10.47, P<0.01), indicating that the traits that promote

biodiversity effects are heritable. As expected, a large proportion of the variation in SCA remained

unexplained. We therefore tested for allelic diversity effects on SCAs at 69 marker positions. Both a

marker regression technique and a standard QTL procedure revealed a major effect locus on the

lower arm of chromosome four where allelic diversity at the community level resulted in higher

SCAs (Fig. 2b and Supplementary Figure 2).

With 18 recombinant lines, mapping resolution was limited and other effect loci or genetic

interactions among loci may have gone unnoticed. We thus aimed at resolving the allelic diversity

effect further to a single Mendelian factor. For this, we isolated a family of 19 near isogenic lines

(NILs) that genetically varied only on the lower arm of chromosome 4, and in which we selected

and inferred further recombination events by molecular markers and whole-genome re-sequencing

(Fig. 3a,b, Supplementary Figure 3). With these NILs, we performed a second diallel experiment,

replicated once on peat-rich soil where we expected no diversity effects and once on sand-rich soil

where we expected positive diversity effects. Indeed, no locus was associated with positive allelic

diversity effects on above-ground biomass on peat-rich soil (Fig. 3c). In contrast, on sand-rich soil

we found a positive allelic diversity effects at a single locus (overyielding of 4.5%, Fig. 3d, $P <$

0.01), represented by a region of approx. 310 kb in size (termed locus Chr4@16.92: between 16.92

to 17.23 Mb). The overyielding of allelic mixtures of otherwise isogenic lines was transgressive:

communities that contained individuals carrying different alleles at locus Chr4@16.92 in the NIL

diallel produced more biomass than the most productive mono-allelic community ($t = 2.32$; $P =$

0.02), suggesting some form of functional complementarity between genotypes: relating to, without

reference to a specific mechanisms at play, a phenomenon of two alleles positively interacting with

each other when distributed *amongst* homozygous individuals of a community. Using structural

equation models, we tested whether these allelic diversity effects were related to observed

genotypic differences in shoot phenology or disease symptoms, but this was not the case

(Supplementary Figure 4). Interestingly, however, this analysis showed that allelic diversity already

manifested itself in increased community-level leaf cover early in the experient, when plants just

started to produce flowering bolts (Supplementary Figure 4), indicating that the effect persisted

through time. The allelic diversity effect we found, could, in principle, strictly depend on the

genetic background. To rule out this possibility, we compared monocultures and mixtures of a

second, independent pair of near-isogenic lines on both peat-rich or sand-rich soil (Supplementary

Figure 5). Again, we found a significant allelic diversity effect on above-ground biomass that depended on soil ($F_{1,164}$ = 4.17, ANOVA P=0.04 for soil x allelic diversity; 5.4% net overyielding on sand), and a significant effect of soil on the complementarity effect ($F_{1,82}$=4.8, P=0.03).

In conclusion, two consecutive diallel experiments using only 37 recombinant lines were sufficient to resolve a plant biodiversity effect to a genomic region representing ~2.5‰ of the Arabidopsis genome (containing approx. 86 genes), which emphasizes the extreme efficiency of our approach. Our work suggests that biodiversity effects between genotypes can be dissected into discrete genetic elements that have major additive contributions.

There is growing evidence that productivity responses in plant biodiversity experiments[4,5,23,24] and their increase through time[25] are related to diversity-dependent soil conditioning. To test whether allelic diversity causes effects through soil conditioning also in our study, we performed soil feedback experiments. Our objectives were twofold: first, we were interested whether there were general effects of allelic diversity on soil quality; second, we aimed to test whether allelic interactions occurred among plants that did not grow simultaneously, i.e. whether allelic effects were mediated through time by soil legacy. We assessed soil conditioning by growing indicator plants ("phytometers"[25]) on soil collected from both diallel experiments[26] when these were harvested (Fig. 1d and Supplementary Discussion). The phytometers were the two parental accessions Bay or Sha for the RIL diallel, and two near-isogenic lines in the NIL diallel.

We indeed found phytometer-specific soil legacy responses that depended on the allelic diversity of the communities that had conditioned the soils in the preceeding growing period (Fig. 4; RIL diallel: diversity at marker MSAT4.9 × phytometer; $F_{1,166}$= 6.48; ANOVA P = 0.012; NIL diallel: diversity at locus Chr4@16.92 × phytometer; $F_{1,168}$= 5.61; P = 0.02; Supplementary Table 1; Supplementary Figure 6). These phytometer-specific responses to soil legacy were independent of differences in previous community productivity and associated resource depletion (the effects

remained statistically significant and comparable in size when first adjusting for community

biomass in linear models). However, the effects differed between phytometers and experiments, i.e.

they depended on environmental or genetic context (Supplementary Discussion). This is not

surprising in light of the complex mechanisms involved. Developing a full understanding of the

biological mechanisms at play will thus require further experiments, including soil analyses.

Nonetheless, these experiments demonstrate that allelic differences at a single QTL cause

interactions between individuals within a community and also, mediated by a soil-borne factor,

through time. The latter can be perceived as an "extended phenotype" sensu Dawkins[27], the

expression of which depends on interactions between group members. We were intrigued to find

that we could, in principle, have genetically mapped the allelic diversity effect solely through its

soil legacy; in other words, by QTL mapping this extended phenotypic property of allelic mixtures

(Supplementary Figure 6e,f; Supplementary Discussion).

Discussion

Our study systematically resolves a biodiversity effect, once identified in a specific set of

interacting plants and environmental context, to between-individual allelic differences in a single

chromsomal region So far, complex emergent properties of plant communities did not necessarily

seem genetically tractable, especially since quantitative traits of individuals often are polygenic[28] if

not omnigenic in nature[29]. A single case study obviously is limited with respect to generalizations,

but we consider it possible that in many cases between-individual allelic complementarity and

resulting biodiversity effects might instead have a relatively simple genetic architecture – a feature

not uncommon for other types of biotic interactions[30]. Our genetic approach is extensible to the

study of interactions among other genotype combinations, and, with modifications, among species,

and could thus lead to fundamental new insights into the traits and genetic underpinnings of

biodiversity effects in more natural systems. Equally importantly, the genetic tractability of such

effects may allow efficient breeding of genotype mixtures that support increased yields through

some form of functional complementarity while maintaining low variation in economically relevant

traits. Biodiversity effects have received relatively little attention in breeding and conventional

agriculture, with the notable exception of crop rotation[31] and intercropping of cultivars and

species[32–34]. Instead, sustaining a growing global human population heavily depends on increasing

nutrient inputs to crop production systems[35], on breeding of single genotypes for monoculture

performance[36], and on the use of within-individual diversity effects termed heterosis[37]. Our

approach might help bypass constraints imposed on the performance of single genotypes, by

shifting breeding efforts from the individual to the system level[38].

Methods

<u>Germplasm</u>

The Shadhara and Bayreuth accessions were kindly provided by Nuno Pires (University of Zurich)

and had originally been obtained from the Nottingham Arabidopsis stock center (NASC). We

selected these two genotypes following an initial screen of ten pairs of accessions, because they

exhibited an above-average but not extreme biodiversity effect when growh together on sand-rich

soil, and because a high-quality and frequently used RIL population is publicly available[20]. It also

was interesting that the biodiversity effect vanished on peat-rich soil. The 18 RILs (representing the

"RIL-minimal set" and line 33RV191 used to generate NILs (all contained in the core-pop set of

165 lines) were ordered from the Versailles Arabidopsis stock center

(*http://publiclines.versailles.inra.fr*) and propagated in a growth chamber. A Bay×Sha RIL

(33RV191) was confirmed to be heterozygous at two PCR marker positions on chromosome 4

(Table S2). Upon selfing of this line, the two NILs 33RV191-Sha and 33RV191-Bay were isolated

(referred to as NIL-Bay or NIL-Sha hereafter), and their genomes were re-sequenced as described

below. Using the same procedure, an independent pair of near-isogenic lines was derived upon

selfing the Bay×Sha RIL 33RV77, giving rise to NILs 33RV77-Sha and 33RV77-Bay. Furthermore,

after selfing of another single heterozygous F_{10} individual of line 33RV191, we screened 160

offspring for recombination between the ShaBa5, ShaBa6 and ShaBa8 markers on chromosome

four. Upon selfing of 23 putative recombinant offspring, we isolated 19 homozygous recombinant

lines for which we confirmed a recombination event in the region by PCR. We then performed

whole-genome re-sequencing to confirm the isogenic background and to infer recombination

breakpoints for this heterogeneous inbred family (referred to as NILs throughout the text) as

described below.

<u>Soils and growth conditions</u>

Soils consisted of different mixtures of a peat and nutrient rich soil (Einheitserde ED73; pH ~5.8, N

250 mg L^{-1}; P_2O_5 300 mg L^{-1}; 75% organic matter content; Gebrüder Patzer GmbH, Sinntal-Jossa,

Germany) and finely grained quartz sand. Pot for all mixture experiments were 7×7×8 cm in size.

The experiment using the parental lines Bay and Sha was replicated on a soil quality gradient with

sand contents of 0%, 40%, 75% and 80%, which resulted in a near-linear decrease of pot-

productivity from the highest to lowest ED73 content, likely through a dilution of soil nutrients

(Fig. S1). For the rough-mapping of the diversity effect using RILs, we used a mixture of 80% sand

and 20% ED73. For the fine-mapping diallel using NILs, we used either a 80%:20% or a 20%:80%

sand:ED73 mixture.

Seeds were sown directly onto soils (approx. 10 seeds per position, 4 positions per pot, Fig. 1a).

The pots were placed in growth chambers or greenhouse compartments and covered with plastic

lids to maintain a high humidity for germination and initial seedling establishment. Additional light

was provided if necessery, achieving a photoperiod of 14–16 hours. Day-time and night-time

temperatures were maintained around 20–25 °C and 16–20 °C, respectively. Seedlings were thinned

continuously until a single healthy seedling remained per position.

Once seedlings were established, the pots were placed in a greenhouse compartment with automated

watering (every 2 days). In summer 2015, daytime temperatures were extremely high, and the first

block of the RIL diallel was therefore grown in a growth chamber with full climate control (8 h

night/16 h day; 60% humidity; 18/23°C night/day temperature). The second block was grown in the

growth chamber for a month before it was re-located to the regular greenhouse compartment.

Pots that did not contain all four originally planted individuals were discarded. Plants were

harvested 43–51 days after sowing, with the specific harvest date determined by the occurence of

approx. 5–10 dehiscent siliques on the earliest flowering genotypes within a block.

After the diallels were harvested, soil feedback trials were established by dividing the soil of a pot

into two smaller pots 5.5×5.5×6.0 cm in size. The respective phytometers (Bay or Sha for the RIL

diallel, 33RV191-Sha or 33RV191-Bay for the NIL diallel) were sown directly onto the soil. Again,

seeds were oversown and seedlings thinned continously until a single healthy individual remained.

Phytometer experiments were harvested either 36 days after sowing (peat-rich soil remaining after

the NIL diallel, harvested early because plant roots started to grow out of the pots) or 49–58 days

after sowing (sand-rich soil, each block was harvested on a single day). For all experiments, the

position of the individual pots was randomized across trays during seedling establishment, and

across watering tables after seedling establishment. Throughout the experiment, pots were re-

positioned randomly within trays and tables every 7–10 days. Pots were watered *ad libitum*, and in

case of high population densities of dark-winged fungus gnats, the systemic insectizide ActaraG

(Syngenta Agro AG) was applied according to the manufacturers recommendation. After

harvesting, plant biomass was dried at 65°C for at least three days before weighing. We determined

early rosette cover in the NIL diallel by photographing pots 27 days after sowing and estimating the

horizontally projected community-level rosette area using the Easy Leaf Area software[39]. We

further recorded the occurrence of leaf disease symptoms (wilting, blotching, or early senescence)

30 days after sowing. As a proxy for flowering time in the NIL diallel, we measured flowering bolt

height of all plants 35 days after sowing; by then, >98% of the individuals had a flowering bolt

longer than 0.5 cm). The NIL diallel was harvested 50 days after sowing.

Experimental designs

To test the soil-dependency of biodiversity effects in mixed Shadhara-Bayreuth communities, four

soil substrates varying in sand content were prepared as described above. We then grew 12 replicate

monocultures of each accession plus 24 replicate mixtures per soil type (total of 48×4= 192 pots).

The RIL diallel consisted of a half diallel replicated in four blocks. All pair-wise RIL combinations

were realized once per block except for RIL monocultures which were replicated twice. For the

follow-up soil feedback experiment, we re-used soil from only the first two blocks of the diallel. We

re-mixed the soil of each single pot after harvesting the plants, and re-distributed it into two smaller

plots that were sown with either a Shadhara or Bayreuth parental genotype that served as

phytometers.

The NIL diallel used for fine-mapping was realized in a single block that contained all pair-wise

combinations of the 19 NILs including monocultures. The subsequent soil feedback part of the

experiment was realized as described for the RIL diallel, using either 33RV191-Bay or 33RV191-

Sha genotypes as phytometers. To test if the allelic diversity effect was strictly dependent on genetic

background, we grew 21 replicate monocultures of each NIL 33RV77-Bay or 33RV77-Sha, plus 42

replicate mixtures, on either peat-rich (80% ED73, 20% sand) or sand-rich (20% ED73, 80% sand)

soil (total of 42×2×2 = 168 pots).

Genotyping and line re-sequencing

PCR-based genotyping assays (Table S2) were developed based on deletions in the Sha genome as

predicted by the Polymorph tool (http://polymorph.weigelworld.org)[40].

Barcoded libraries for genome re-sequencing were prepared using the Illumina Nextera DNA

Library Prep Kit (FC-121-1031, Illumina Inc. San Diego, CA) in combination with the Nextera

Index Kit (96 indices, FC-121-1012) and pair-end sequenced on an Illumina HiSeq 2500 (2x150 bp,

rapid run). The clustering and sequencing were performed at the Functional Genomics Center

Zurich. Sequences were aligned to the Arabidopsis genome (Col-0 genome, TAIR version 10) using

BWA[41], aligned read sorting and variant calling were performed using samtools[42]. Aligned genomic

sequences of the parental accessions Bay-0 and Sha were downloaded from the 1001 genomes

project data center (http://1001genomes.org). The VCF-file produced by the samtools software was

loaded into the R Statistical Software[43], where the subsequent analyses were performed: variant

calls were filtered (for differences in genotype calls between the Sha and Bay genomes, quality of

variant calls, population-level minimal minor allele frequency 0.2; maximum heterozygosity 0.2).

Inference of genotype calls at polymorphic sites was performed as described previously[44] and

inference of parental alleles was improved using functionality implemented in the MPR package[44].

Genotype reconstruction was then performed in R using a simple hidden Markov model as

implemented in the R package HMM, with hidden state starting probabilities (Bay, Het or Sha) all

set to 1/3, and transition probabilities from one state to itself set to 0.99998 and to the other two

states set to 0.00001 each. Emission probabilities of genotype calls given a state, e.g. Bay, were set

to 0.35, 0.25, 0.25, 0.15 for genotypes calls Bay, Het, Sha or missing, etc.

<u>Statistical analyses</u>

We analyzed data from the diallel experiments using linear mixed models summarized by analysis

of variance (ANOVA). The model terms included, in this order, the general combining abilities

(GCA) of genotypes (a factor with 20 levels in the RIL diallel and 19 levels in the NIL diallel), the

genotype diversity in the pot (GD, 1 or 2 genotypes), the allele identity in the genotype

monocultures (A, Sha or Bay), the allelic diversity in the genotype mixtures (AD, 1 [Sha/Sha or

Bay/Bay] or 2 [Sha/Bay]), and the genotype composition planted in the pot (comp). The factor GCA

was created by superimposing the model matrices for factors coding for the first and second

genotype (factors with 20 and 19 levels for RIL and NIL diallels, respectively). The significance of

GD, A, and AD were determined using F-tests with comp as error term (denominator). A and AD

were encoded in such a way that these contrasts applied only to genotype monocultures and

mixtures, respectively. Technically, this was achieved by including a third level in the factor that did

not vary in the other group. Fitting A and AD after GD therefore only explained variance in these

subsets. The diallel model was extended by additional terms and the corresponding interactions

when these applied; specifically, the RIL diallel included a block effect. The NIL diallel included

terms for soil type, and interactions of all the terms above with soil type (for example, soil×AD was

tested using soil×comp as error term). The soil feedback experiments included further interactions

with phytometer (RIL and NIL diallel), and phytometer×soil (NIL diallel). Effects of pot biomass in

the diallel (diallel biomass and diallel biomass × soil) were accounted for in these linear models, and

data were square-root transformed to obtain normally distributed residuals.

Specific combining abilities for mapping were calculated directly, within blocks, by solving the

linear model m = X GCA + SCA where X is the design matrix describing the genotype composition

of a pot. Monoculture SCAs were also determined but not used for QTL mapping of allelic diversity

within RIL mixtures. In the RIL diallel, the SCAs of each genotype composition was first calculated

per block and then aggregated over all blocks using least-square estimates. Marker regression was

performed contrasting SCAs of mono-allelic RIL mixtures ("BB" and "SS" compositions) with

mixed-allelic mixtures ("BS" compositions) using the glht-function provided by the multcomp

package[43]. QTL mapping was also performed using the R/qtl package and interval mapping

(scanone-function), with both mono-allelic compositions at a given locus re-coded as to the same

level ("mono-allelic") and compared against mixed-allelic compositions. Genome-wide significance

was assessed by resampling (n=5000).

To test the relationships of allelic diversity effects on SCAs with measured traits, we developed multip-group (sand-rich and peat-rich soil) structural equation models using lavaan (http://lavaan.ugent.be). Allelic diversity at Chr4@16.92 was included as exogenous variable; endogeneous variables were two metrics of trait variation among genotypes (which are possible indicators of complementarity), and early community-level projected leaf area (see above). Trait variation was quantified as difference among bolt length of the genotypes (square-root-transformed), and as difference in the occurence of leaf disease symptoms (a binary variable). Starting with a near-saturated model, the modelled paths were simplified in an educated way until a minimal model was found for which the model-implied and observed covariance structure among variables did not differ significantly (X^2-test).

References

1. Cardinale, B. J. *et al.* The functional role of producer diversity in ecosystems. *Am. J. Bot.* **98,** 572–592 (2011).

2. O'Connor, M. I. *et al.* A general biodiversity–function relationship is mediated by trophic level. *Oikos* **126,** 18–31 (2017).

3. Cardinale, B. J., Palmer, M. A. & Collins, S. L. Species diversity enhances ecosystem functioning through interspecific facilitation. *Nature* **415,** 426–429 (2002).

4. Schnitzer, S. A. *et al.* Soil microbes drive the classic plant diversity-productivity pattern. *Ecology* **92,** 296–303 (2011).

5. Maron, J. L., Marler, M., Klironomos, J. N. & Cleveland, C. C. Soil fungal pathogens and the relationship between plant diversity and productivity. *Ecol. Lett.* **14,** 36–41 (2011).

6. Loreau, M. & Hector, A. Partitioning selection and complementarity in biodiversity experiments. *Nature* **412,** 72–76 (2001).

7. Ratcliffe, S. *et al.* Biodiversity and ecosystem functioning relations in European forests depend on environmental context. *Ecology Letters* **20,** 1414–1426 (2017).

8. Tylianakis, J. M. *et al.* Resource heterogeneity moderates the biodiversity-function relationship in real world ecosystems. *PLoS Biol.* **6,** 947–956 (2008).

9. Flynn, D. F. B., Mirotchnick, N., Jain, M., Palmer, M. I. & Naeem, S. Functional and phylogenetic diversity as predictors of biodiversity--ecosystem-function relationships. *Ecology* **92,** 1573–1581 (2011).

10.   Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428,** 821–827 (2004).

11.   Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529,** 167–171 (2015).

12.   Cadotte, M. W., Cardinale, B. J. & Oakley, T. H. Evolutionary history and the effect of
biodiversity on plant productivity. *Proc. Natl. Acad. Sci. USA* **105,** 17012–17017 (2008).

13.   Crutsinger, G. M. *et al.* Plant genotypic diversity predicts community structure and governs
an ecosystem process. *Science* **313,** 966–968 (2006).

14.   Wang, S. & Loreau, M. Ecosystem stability in space: α, β and γ variability. *Ecol. Lett.* **17,**
891–901 (2014).

15.   Oehri, J., Schmid, B., Schaepman-Strub, G. & Niklaus, P. A. Biodiversity promotes primary
productivity and growing season lengthening at the landscape scale. *Proc. Natl. Acad. Sci.*
38, 10160–10165 (2017).

16.   Siefert, A. *et al.* A global meta-analysis of the relative extent of intraspecific trait variation in
plant communities. *Ecology Letters* **18,** 1406–1419 (2015).

17.   Crutsinger, G. M., Souza, L. & Sanders, N. J. Intraspecific diversity and dominant genotypes
resist plant invasions. *Ecol. Lett.* **11,** 16–23 (2008).

18.   Prieto, I. *et al.* Complementary effects of species and genetic diversity on productivity and
stability of sown grasslands. *Nat. Plants* **1,** 15033 (2015).

19.   Mendel, G. Versuche über Pflanzenhybriden. in *Verhandlungen des naturforschenden*
Vereines in Brünn, Band IV für das Jahr 1865 3–47 (1865).

20.   Loudet, O., Chaillou, S., Camilleri, C., Bouchez, D. & Daniel-Vedele, F. Bay-0 x Shahdara
recombinant inbred line population: A powerful tool for the genetic dissection of complex
traits in Arabidopsis. *Theor. Appl. Genet.* **104,** 1173–1184 (2002).

21.   Griffing, B. Concept of General and Specific Combining Ability in Relation to Diallel
Crossing Systems. *Aust. J. Biol. Sci.* **9,** 463–493 (1956).

22.   Griffing, B. Genetic-Analysis of Plant Mixtures. *Genetics* **122,** 943–956 (1989).

23.   Zak, D. R., Holmes, W. E., White, D. C., Peacock, A. D. & Tilman, D. Plant diversity, soil
microbial communities, and ecosystem function: Are there any links? *Ecology* **84,** 2042–
2050 (2003).

24.   Bukowski, A. R. & Petermann, J. S. Intraspecific plant-soil feedback and intraspecific
overyielding in Arabidopsis thaliana. *Ecol. Evol.* **4,** 2533–2545 (2014).

25.   Meyer, S. T. *et al.* Effects of biodiversity strengthen over time as ecosystem functioning
declines at low and increases at high biodiversity. *Ecosphere* **7,** (2016).

26.   Clements, F. E. & Goldsmith, G. W. *The Phytometer Method in Ecology.* (Carnegie
Institution of Washington, 1924).

27.   Dawkins, R. *The Extended Phenotype.* (Oxford University Press, 1982).

28. MacKay, T. F. C., Stone, E. A. & Ayroles, J. F. The genetics of quantitative traits: Challenges
and prospects. *Nature Reviews Genetics* **10,** 565–577 (2009).

29. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From
Polygenic to Omnigenic. *Cell* **169,** 1177–1186 (2017).

30. Wilfert, L. & Schmid-Hempel, P. The genetic architecture of susceptibility to parasites. *BMC*
Evol. Biol. **8,** 187 (2008).

31. Wittwer, R. A., Dorn, B., Jossi, W. & Van Der Heijden, M. G. A. Cover crops support
ecological intensification of arable cropping systems. *Sci. Rep.* **7,** 41911 (2017).

32. Finckh, M. R. *et al.* Cereal variety and species mixtures in practice, with emphasis on disease
resistance. *Agronomie* **20,** 813–837 (2000).

33. Zhu, Y. *et al.* Genetic diversity and disease control in rice. *Nature* **406,** 718–722 (2000).

34. Litrico, I. & Violle, C. Diversity in Plant Breeding: A New Conceptual Framework. *Trends in*
Plant Science **20,** 604–613 (2015).

35. Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R. & Polasky, S. Agricultural
sustainability and intensive production practices. *Nature* **418,** 671–677 (2002).

36. Edgerton, M. D. Increasing crop productivity to meet global needs for feed, food, and fuel.
Plant Physiol. **149,** 7–13 (2009).

37. Lippman, Z. B. & Zamir, D. Heterosis: revisiting the magic. *Trends in Genetics* **23,** 60–66
(2007).

38. Weiner, J., Du, Y. L., Zhang, C., Qin, X. L. & Li, F. M. Evolutionary agroecology: individual
fitness and population yield in wheat (Triticum aestivum). *Ecology* **98,** 2261–2266 (2017).

39. Easlon, H. M. & Bloom, A. J. Easy Leaf Area: Automated Digital Image Analysis for Rapid
and Accurate Measurement of Leaf Area. *Appl. Plant Sci.* **2,** apps.1400033 (2014).

40. Fitz, J. *et al.* personal communication. *http://polymorph.weigelworld.org*

41. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows – Wheeler
transform. *Bioinformatics* **25,** 1754–1760 (2009).

42. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078–
2079 (2009).

43. R Core Team. R: A Language and Environment for Statistical Computing. *R Foundation for*
Statistical Computing, Vienna, Austria ISBN 3-900051-07-0 (2017). doi:http://www.R-
project.org/

44. Xie, W. *et al.* Parent-independent genotyping for constructing an ultrahigh-density linkage
map based on population sequencing. *Proc. Natl. Acad. Sci.* **107,** 10578–10583 (2010).
423

Acknowledgements

We thank Bernhard Schmid and Ueli Grossniklaus for helpful discussions and sharing infrastructure. We thank Jordi Bacompte and Jacob Weiner for helpful comments on the manuscript. We further acknowledge Matthias Philipp for technical support, Enrica De Luca and Nicole Ponta for help with plant measurements, Matthias Furler and Dorde Topalovic for technical greenhouse support. This work was supported by an Ambizione Fellowship (PZ00P3_148223) of the Swiss National Science Foundation (to S.E.W). P.A.N acknowledges support by the University of Zurich Priority Programme "Global Change and Biodiversity". S.E.W was also financially supported by funds of the University of Zurich and the European Research Council (to Ueli Grossniklaus).

Author contributions

S.E.W. conceptualized and designed the research (with input from P.A.N.) and performed the experiments. Both authors performed the analyses and wrote the manuscript. Both authors revised and approved the final version of the manuscript.

Data availability

The datasets described in the paper and a functional annotation of the 86 genes within the fine-mapped diversity QTL are available through the Zenodo data repository (DOI:10.5281/zenodo.1254563). Sequencing data are deposited in the NCBI Short-Read Archive (accession SRP149077).

Analysis scripts are available from the authors upon request.

Competing interests

The authors declare no competing financial interests.

Figures

Figure 1 | Combining ecological concepts and genetic methods. a, Pot systems used to study diversity effects in pair-wise genotype mixtures. The inset shows a Recombinant Inbred Line (RIL) monoculture (left) and mixture (right). **b,** Net diversity effects in Bay-Sha mixtures along a peat-sand substrate gradient. Error bars denote standard errors of means (s.e.m.). n = 164 pots **c,** Outline of the diallel design and the genotypes used throughout this paper. 18 RILs and the two parental accession, or 19

near-isogenic lines (NILs) were each placed in competition with each other, allowing to assess i) effects of genotypic mixture (i.e. diagonal vs off-diagonal), or ii) effects of allelic mixture at a given locus across all genotype mixtures (i.e. comparing SS and BB vs. SB) on pot productivity. **d,** Outline of the experimental procedure used in this work. Colored labels indicate measured variables. GCA = general combining ability; SCA = specific combining ability.

Figure 2 | Allelic diversity at a major effect locus increases community productivity. a, Pot-level productivity in dependence of community type (mix = RIL mixtures vs mono = RIL monocultures), showing positive genotype mixture effects in the diallel and on sand-rich soil (values aggregated across four blocks, n = 871 pots / 210 compositions) **b,** Quantitative trait locus interval mapping of allelic diversity effects on specific combining ability (SCA). Vertical lines denote 10% and 5% genome-wide significance levels. The inset shows estimated SCA (± s.e.m.) across genotype mixtures that exhibit different allelic compositions at marker MSAT4.9 on chromosome four. LOD: logarithm of the odd.

Figure 3 | Resolving soil × allelic diversity interactions to a single Mendelian factor. a, Resequencing of near-isogenic lines (NILs) differing only on lower arm of chromosome four for fine-mapping. Shown are genotype calls at all polymorphic sites across the genome (B = homozygous for the Bay allele, het = heterozygous, S = Sha allele) in either NIL r10 (blue, top left) or NIL r96 (red, top right), as well as an overlay of the two line's genotype calls (bottom). **b,** Reconstructed genotypes across chromosome four of the 19 NILs used for fine-mapping. Each bar represents a single NIL. **c, d,** Map of allelic diversity effects across chromosome four, on either peat-rich soil (c), or sand-rich soil (d). The widths of the bars indicate the size of the regions in which no recombination events were

inferred across the whole population. Inset in (d) shows the mean ± s.e.m. of SCAs across allelic compositions at the diversity effect QTL.

Figure 4 | Allelic diversity effects persist across a generation through their soil legacy. a, Scheme of the genotype-combinations used for either the recombinant inbred lines (RILs) vs. near-isogenic lines (NILs) diallels (top), and the phytometer genotypes used in the soil-feedback phase (bottom). **b**, phytometer performance (Bay or Sha, mean ± s.e.m.) on legacy soil derived from RIL mixtures with different allelic compositions at marker MSAT4.9. **c**, phytometer performance (NIL-Bay or NIL-Sha, mean ±s.e.m.) on legacy soil derived from NIL mixtures with different allelic compositions at locus Chr4@16.92 on either sand-rich (left) or peat-rich (right) soil. Values were square-root transformed for analyses, n = 851 and 720 pots for RIL and NIL diallels, respectively.

Supplementary Tables

Supplementary Table 1 | Legacy effects of soils conditioned in RIL and NIL diallel experiments. Effects were quantified using two phytometers (factor 'phy'; Sha and Bay accessions in RIL diallel, and near-isogenic lines bearing Sha and Bay allele at putative effect locus in NIL diallel). The NIL diallel additionally was replicated on substrates differing in sand content (factor 'soil'). The term 'GCA' (general combining abilities) indicates average genotype-specific soil conditioning effects on phytometer yields. 'SCA' (specific combining abilities) captures deviations in yield from additive predictions made using GCAs. Within SCA, the following contrasts were tested: GD: Genotype diversity, i.e. whether genotype monocultures differed in feedback effects from two-genotype mixtures; A: allele-specific differences at marker MSAT4.9 (RIL diallel) and Chr4@16.92 (NIL diallel) within genotype monocultures; AD: allele-diversity effects within genotype mixtures. df and ddf indicate nominator and denominator degrees of freedom of corresponding F-tests. *** P<0.001; ** P<0.01; * P<0.05; (*) P<0.1; n.s. not significant.

Terms and contrasts	RIL diallel			NIL diallel		
	df	Denominator: ddf	Signif.	df	Denominator: ddf	Signif.
GCA	18	comp: 167	(*)	18	comp: 168	n.s.
SCA						
GD	1	comp: 167	(*)	1	comp: 168	n.s.
A	1	comp: 167	n.s.	1	comp: 168	n.s.
AD	1	comp: 167	n.s.	1	comp: 168	n.s.
Phy ₓ GCA	18	phy ₓ comp: 166	n.s.	18	phy ₓ comp: 168	*
Phy ₓ SCA						
Phy ₓ GD	1	phy ₓ comp: 166	n.s.	1	phy ₓ comp: 168	n.s.
Phy ₓA	1	phy ₓ comp: 166	n.s.	1	phy ₓ comp: 168	n.s.
Phy ₓAD	1	phy ₓ comp: 166	**	1	phy ₓ comp: 168	*
Soil ₓ GCA				18	soil ₓ comp: 150	n.s.
Soil ₓ SCA						
Soil ₓ GD				1	soil ₓ comp: 150	n.s.
Soil ₓA				1	soil ₓ comp: 150	n.s.
Soil ₓAD		does not apply		1	soil ₓ comp: 150	n.s.
Phy ₓ Soil ₓ GCA				18	phy ₓ soil ₓ comp: 146	n.s.
Phy ₓ Soil ₓ SCA						
Phy ₓ Soil ₓ GD				1	phy ₓ soil ₓ comp: 146	***
Phy ₓ Soil ₓA				1	phy ₓ soil ₓ comp: 146	n.s.
Phy ₓ Soil x AD				1	phy ₓ soil ₓ comp: 146	n.s.

507 **Supplementary Table 2: PCR markers used in this study.**

Primer	Assay	Sequence	Predicted location	Pred. fragment sizes (Sha/Bay)
SW-182	ShaBa5	ACGTATTTCGATGTATGGTCCTTG	Chr4: 16044156	550/664
SW-183		TCACGTGAATCGTATTCGTTGAAG		
SW-184	ShaBa6	CTTCTCCGCTTCAACCTCTGC	Chr4: 17709750	600/632
SW-185		AATCCAGGATTCAGAGTTGCTTTC		
SW-188	ShaBa8	TTGATTAGGGCTACGAGGATAAGG	Chr4: 16707214	408/609
SW-189		GAGTCTATTAATTATGCTTGGTGC		

 Supplementary Figures

Supplementary Figure 1 | Productivity and complementarity in Bay-Sha mixtures across a

peat-sand gradient. a, Pot-level biomass measurements of Sha and Bay monocultures or pair-wise

mixtures along a sand/peat substrate gradient. n = 164 pots **b,** Complementarity and selection

effects calculated according to the additive partitioning [1] method along the substrate gradient. Error

bars denote s.e.m.

Supplementary Figure 2 | A major effect locus driving complementarity between genotypes. a,

QTL mapping of SCA variation across allelic diversity levels using a marker regression technique

by contrasting SCAs of mono-allelic RIL mixtures (BB and SS) with bi-allelic mixtures (BS). **b**,

Pot-level productivity of each genotype composition (average of four blocks) in dependence of

allelic composition at marker MSAT4.9. $n_{composition} = 210$

Supplementary Figure 3 | Fine-mapping and persistence of allelic diversity effects in near-isogenic line. a, The comparison of the reconstruced genotype of NIL r10 (HMM Viterbi-path across all chromosomes, homozygous on lower arm of chromosome 4) in comparison to publicly available molecular marker-based genotyping data of the ancestral line from which it was derived (heterozygous on lower arm of chromosome 4) – showing a high degree of congruence between the re-constructed genotype based on whole-genome resequencing and the marker data. **b,** Schematic outline of a possible cause of the high mapping resolution achieved through the diallel design. A major advantage of the design is the joint dependency of community-level allelic diversity on recombinations *within* and *between* recombinant inbred lines, such that mapping resolution increases very quickly. **c,** Extreme example of SCA variation across genotype mixtures all containing one specific genotype (NIL r146, homozygous for the Bay-allele at locus Chr4@16.92),

 either in combinations with NILs carrying the Sha-allele (dark blue bars) or the Bay-allele (dark red

bars). Shown are the data from sand-rich soil only.

Supplementary Figure 4 | Observed above-ground trait variation does not explain

overyielding of diallelic mixtures. a, Traits measured in the NIL diallel as proxy for productivity

(early growth projected leaf cover), disease susceptibility (leaf disease at 30 days after sowing) or

phenology (bolt length at 35 days after sowing) **b,** Path diagram of multi-group structural equation

model showing direct effects of allelic diversity at locus Chr4@16.92 on SCA on sandy but not on

peat soil. Red and blue arrows show positive and negative standardized path coefficients,

respectively. n.s. = not significant. (*) = p<0.1; * = p<0.05; ** = p<0.01.

548

Supplementary Figure 5 | Soil-dependent effect of allelic diversity on overyielding and

statistical complementarity and selection effects[S1] in an independent near-isogenic

background. a, Overlay of genotype calls at all polymorphic sites across the genome (Sha and Bay =

homozygous for respective allele, het = heterozygous) of lines used for a mendelization. Top: a

comparison of the genetically independent backgrounds 33RV191-Sha (in red, this background was

used for the fine-mapping shown in Fig. 3) and line 33RV77-Sha (in blue) is shown. At the bottom, a

comparison of the near-isogenic lines 33RV77-Sha (red) vs 33RV77-Bay (blue), the two lines that were

used in the experiment shown in b and c. Purple regions depict overlap of genotype calls, vertical lines

separate the different chromosomes. **b**, Mean ±s.e. of final aboveground biomass of communities

consisting either of 33RV77-Sha plants only (BB), of mixed communities (BS), or of communities

consisting of 33RV77-Bay plants only (SS) on either peat-rich (left) or sand-rich (right) soil. **c**,

Complementarity and selection effects sensu Loreau and Hector calculated from the experiment

shown in b.

Supplementary Figure 6 | Soil feedback experiments. a, Genotype calls at all polymorphic sites

across the genome (B = homozygous for the Bay allele, H = heterozygous, S = homozygous for the

Sha allele) obtained by genome re-sequencing phytometer genotypes NIL-Bay (33RV191-Bay, top

left, red) and NIL-Sha (33RV191-Sha, top right, blue). The overlay of the genotype calls of both

lines (bottom) confirms that these lines are isogenic but for variation on lower arm of chromosome

four. The two phytometers were employed on soil derived from the NIL diallel. **b, c,** QTL mapping

by marker regression of phytometer-specific responses to soil legacy of previous generation allelic

diversity (i.e. allele-diversity × phytometer [Phy×AD] interaction in Supplementary Table 1, but in a

model without adjusting for diallel pot biomass). The mapping of diversity effects through such

influences on soil legacy could have been applied to the identification and fine-mapping of the same

major effect locus, without ever measuring biomass productivity (albeit with slightly relaxed

statistical criteria). Shown are negative $\log_{10}$-transformed P-values for each marker position in the

RIL (b) or genotype block in the NIL (c) diallels.

Supplementary Discussion

As outlined in Figures 1 and 4, and as described in the Methods, we performed two soil-feedback

experiments to test whether allelic diversity effects extend across generations. We accounted for

potential effects explainable by variation in plant productivity during the soil training phase (e.g.

nutrient draw-down or environmental correlations) in linear models (terms *diallel biomass* and

diallel biomass×soil as described in the Methods section). Significantly different soil conditioning

through allelic diversity, as assessed by phytometer performance in a next growing period, was

interpreted as an "extended phenotype"[S2] (i.e. a legacy of allelic mixture that persists through time

in the soil, even after removal of the original plant communities). These soil factor-mediated allelic

legacy effects interact with phytometer genotype, giving rise to phytometer-specific responses (term

"Phy×AD" in Supplementary Table 1).

As we emphasize, the specific mechanisms underlying both soil training and phytometer responses

await further experiments, since the response patterns (Fig. 4) do not allow for a simple mechanistic

model. One reason why it is difficult to infer specific mechanisms (e.g. specific soil factors, and

their interactions with genetic variants) is the number of variables that differed between the two

experiments because of constraints on experimental design: 1) environmental conditions (different calendar dates, different batches of soil); 2) genetic variation in the two diallel population (RILs vs NILs), potentially resulting in different effects of epistasis; and 3) phytometer genotypes (parental lines vs NILs).

In the RIL diallel, we used the two parental accession Bay and Sha as phytometers and a split-plot design to test for differential responses of these phytometer to soil conditioning. After the NIL diallel, we used two near-isogenic phytometer (33RV191-Bay and 33RV191-Sha) as phytometers in a similar test for differential responses to soil conditioning. Naively, one might expect approximately congruent responses of the phytometers carrying the same alleles at the diversity locus under question, i.e. the parental genotype Bay in the RIL diallel (Figure 4 b, yellow lines) and the NIL-Bay genotype in the NIL diallel (Figure 4 c, yellow lines) might respond similarly to the legacy of allelic diversity. However, this was not the case. For example, the Bay genotype grown on RIL diallel soil responded positively to conditioning by allelic diversity, whereas the NIL-Bay genotype in the NIL diallel responded somewhat negatively to conditioning by allelic diversity. It is noteworthy that the pattern found for Bay on RIL diallel soil is opposite to what would be expected if soil legacy was driven by a simple productivity-related depletion of resources in the conditioning phase; this suggests that allelic diversity alleviates a negative soil factor (e.g. inhibits enemies), or promotes a positive factor (e.g. mutualistic organisms).

Despite differences, the RIL and NIL soil-feedback experiments nevertheless convincingly demonstrate i) that allelic diversity effects extend across time through soil conditioning, and ii) that phytometer genotype determines the response to such conditioning (phytometer×allelic diversity interaction, both experiments, see Table S1). The fact that allelic diversity effects within a community and allele-specific legacy effects on individuals separated in time map to the same locus suggests to us that both are related to similar mechanisms.

Supplementary References

617   **Supplementary References**

619   1.   Loreau, M. & Hector, A. Partitioning selection and complementarity in biodiversity
620   experiments. *Nature* **412,** 72–76 (2001).

621   2.   Dawkins, R. *The Extended Phenotype*. (Oxford University Press, 1982).