A plant biodiversity effect resolved to a single locus

Samuel E. Wuest[1,2,*] and Pascal A. Niklaus[2]

1) Department of Plant and Microbial Biology and Zurich-Basel Plant Science Center, University of Zurich, Zollikerstrasse 107, 8008 Zurich

2) Department of Evolutionary Biology and Environmental Studies and Zurich-Basel Plant Science Center, University of Zurich, Winterthurerstrasse 190, 8057 Zurich

*correspondence to: Samuel E. Wuest (samuel.wuest@ieu.uzh.ch)

Summary

There now is pervasive evidence of positive effects of biodiversity on plant community productivity

and functioning[1,2]. Some progress has been made linking diversity effects to functional trait

variation[3], but progress towards a mechanistic understanding remains slow[4] - in part because

biodiversity effects are emergent properties of communities, and mechanisms might differ between

cases or environmental conditions[5]. Here, we analyse non-additive interactions between divergent

Arabidopsis accessions in experimental plant communities. By combining concepts and designs

from ecology and plant breeding with genetic methods, we identified a major effect locus at which

allelic diversity promotes community productivity. In further experiments with near-isogenic lines,

this diversity effect locus was resolved to a single Mendelian factor and a region representing less

than 0.3% of the genome. Using plant-soil-feedback experiments, we demonstrate that allelic

diversity causes genotype-specific soil legacy responses in the growth of a subsequent plant

generation. This suggests that asymmetric interactions of plants with soil-borne factors drive niche

complementarity, and that the effects of allelic diversity at a single locus can extend from the plant

to the system-level. Our work establishes a novel, fully reductionist approach to the study of a

complex ecological process. It may also pave the way to novel crop breeding strategies, which

focus on phenotypic properties that do not manifest at the individual-, but only at a higher level of

biological organisation.

Main Text

More than two decades of plant ecological research and the publication of hundreds of studies have

firmly established that positive biodiversity effects, in particular on community yield, are the rule

rather than exception and often substantial[1,2]. These effects have been explained by larger

community-level resource use promoted by niche complementarity, and by reduced negative

density-dependent effects of enemies[6,7]. Yet, our understanding of the particular driving

mechanisms remains poor, for several reasons. First, diversity effects are emergent properties that only manifest in comparisons of communities exhibiting different diversity levels[8]. Second, diversity effects, and the mechanisms that drive these, may change with environmental conditions[5,9]. Third, while there is no doubt that functional trait differences underly biodiversity effects[10], trait-based analyses remain to some degree phenomenological because evolutionary forces have led to the formation of trait syndromes, i.e. to sets of highly correlated traits that reflect fundamental trade-offs between ecological strategies[11,12]. The observed variation in traits thus is confounded with phylogeny[13], and it remains almost impossible to distinguish the traits that are true drivers of biodiversity effects from others that are merely correlated.

Most biodiversity research to date has focused on variation among species, but experimental[14], theoretical[15], and observational[16] studies have shown that positive diversity effects on productivity also occur at levels of organization above and below species. An important part of the trait variation apparent in plant communities occurs within species[17], and increased intra-specific variation can have similar effects as inter-specific trait variation in low-diversity systems[14,18,19]. Despite qualitative differences, there may therefore be commonalities of trait variation within and between species with respect to effects and mechanisms, indicating that studies at the genotype level may provide some insights into effects of species-level variation and vice versa.

A methodological advantage of intra-specific biodiversity studies is that genetic methods can circumvent some of the problems encountered in species-level diversity studies. Specifically, crosses between genotypes allow to re-arrange trait variation between individuals[20] without confounding with population structure or the differentiation into ecological strategies. However, genetic approaches are normally used to study properties of individuals rather than emergent properties of communities.

Here, we demonstrate in a case study how the genetic approach can be harnessed to identify the genetic underpinnings of biodiversity effects. By screening pair-wise mixtures of divergent natural

accessions of *Arabidopsis thaliana,* we identified two accessions, Bayreuth (Bay) and Shadhara

(Sha) that exhibit positive net biodiversity effects when grown together, i.e. mixtures produce a

higher community-level biomass than the average of their monocultures (Fig. 1a). This effect

depended on soil conditions, with effects that were essentially absent on peat-rich soil but grew to

an overyielding of 16% with increasing amounts of sand in the substrate (sand content × diversity:

$F_{1,160}$= 4.57, P < 0.05; Fig. 1b and Extended Data Fig. S1a). Analysis by additive

partitioning[8] revealed that these community-level biodiversity effects were due to complementarity

rather than selection effects (sand content × complementarity effect, $F_{1,77}$ = 7.21, P < 0.01; Extended

Data Fig. S1b); specifically, this indicates that both ecotypes benefited from growing in mixed

communities.

To analyse the genetic basis of the positive diversity effect in mixed Bay-Sha communities, we

performed quantitative trait locus (QTL) mapping using publicly available recombinant inbred lines

(RILs). These RILs are largely homozygous (Fig. 1c) and have been derived from a cross between

the Bay and Sha accessions, followed by multiple subsequent rounds of inbreeding[21]. For efficient

mapping, we capitalized on a so called competition diallel. Traditional diallel designs systematically

cross parental lines and are used in breeding to determine the genetic basis of traits; specifically,

diallel analysis partitions the traits of crosses into additive contributions[22] of parental lines (general

combining abilities; GCA) and cross-specific effects (specific combining abilities; SCA), with the

latter interpreted as consequences of dominance or epistasis. By substituting individuals and crosses

with communities and mixtures, the principle of diallels can be applied to the analyses of

biodiversity effects in communities[23], which we did here (Fig. 1c, d). In this context, the distinction

between maternal and paternal effects ceases to apply, simplifying the design to a half-diallel. SCAs

then quantify the deviations of mixture yields from expectations based on additive average

contributions of the two genotypes. In our study, we combined 18 RILs and the two parental

accessions Bay and Sha, in four replicate blocks, on sand-rich soil. We detected significant positive

genotype diversity effects (Fig. 2a, $F_{1,189} = 10.47$, P<0.01), indicating that the traits that promote

biodiversity effects are heritable. As expected, a large proportion of the variation in SCA remained

unexplained. We therefore tested for allelic diversity effects on SCA at 69 marker positions. Both a

marker regression technique and a standard QTL procedure revealed a major effect locus on the

lower arm of chromosome four where allelic diversity at the community level resulted in higher

SCAs (Fig. 2b and Extended Data Fig. S2).

With 18 recombinant lines, mapping resolution was limited and other effect loci or genetic

interactions among loci may have gone unnoticed. We thus aimed at resolving the allelic diversity

effect further to a single Mendelian factor. For this, we isolated a family of 19 near isogenic lines

(NILs) that genetically varied only on the lower arm of chromosome 4, and in which we selected

and inferred further recombination events by molecular markers and whole-genome resequencing

(Fig. 3a,b, Extended Data Fig. S3). With these NILs, we performed a second diallel experiment,

replicated once on peat-rich soil where we expected no diversity effects and once on sand-rich soil

where we expected positive diversity effects. Indeed, no locus was associated with positive allelic

diversity effects on peat-rich soil (Fig. 3c). In contrast, we found a positive allelic diversity effects

at a single locus (overyielding of 4.5%, Fig. 3d, P < 0.01), represented by a region of approx. 310

kb in size (termed locus Chr4@16.92 between 16.92 to 17.23 Mb). The overyielding of these allelic

mixtures was transgressive, i.e. allelic mixtures produced more biomass than the best mono-allelic

communities (t = 2.32; P = 0.02). Transgressive overyielding is generally accepted as evidence of

diversity effects that are driven by niche complementarity.

Using soils from both diallels, we next performed soil feedback experiments. We grew

phytometers[24] on soil pre-conditioned ("trained") by the growth of the specific genotype

combinations (Fig. 1d and Supplementary Discussion). These phytometers were the two parental

accessions Bay or Sha for the RIL diallel, and two near-isogenic lines in the NIL diallel. We found

significant phytometer-specific responses to soil legacy influenced by varying allelic diversity in

communities of the previous generation (Fig. 4; RIL diallel: diversity at marker MSAT4.9 ×

phytometer; $F_{1,166}$= 6.48; P = 0.012; NIL diallel: diversity at locus Chr4@16.92 × phytometer; $F_{1,168}$=

5.61; P = 0.02; Extended Data Table S1; Extended Data Fig. S4). Importantly, these phytometer-

specific responses to soil legacy were not mediated by differences in previous community

productivity and associated resource depletion (the effects remained statistically significant and

comparable in size when first adjusting for community biomass in linear models). Interestingly,

conditioning effects and phytometer responses of plants bearing the Bay allele differed in the

experiments with RILs and NILs. This indicates that genotypes with the Bay allele in both cases

strongly conditioned soils, which resulted in subsequent legacy effects. However, and not

unexpectedly given the complexity of mechanisms involved, these allele-specific effects cannot be

understood in isolation but depend on environmental conditions or genetic context (Supplementary

Discussion). Developing a full understanding of the biological mechanisms at play will thus require

further experiments, including soil analyses. Yet, our experiments demonstrate that plant genotype

interactions mediated through allelic diversity at a single locus extend their effects across

generations through their soil legacy. We were intrigued to find that we could in principle have

genetically mapped this allelic diversity effect solely through its soil legacy; in other words, by

QTL mapping an extended emergent property of allelic mixture (Extended Data Fig. S4e,f;

Supplementary Discussion).

Our study is, to the best of our knowledge, the first to systematically resolve biodiversity effects to

their genetic origin. So far, quantitative emergent properties of plant communities did not

necessarily seem genetically tractable, especially since quantitative traits of individuals often are

polygenic[25] if not omnigenic in nature[26]. The major diversity effect locus we found was resolved to

a region representing ~2.5‰ of the Arabidopsis genome (containing approx. 86 out of a total of

27'205 annotated genes) with a total of only 37 recombinant lines, which emphasizes the extreme

efficiency of diallel designs for such mapping. We consider it possible that many complementarity

and therefore biodiversity effects, both within and among species, might have a relatively simple

genetic architecture - as seen with other types of biotic interactions[27]. A single case study obviously

is limited with respect to generalizations, but our approach is extensible to the study of biodiversity

effects among other genotype combinations or even, with modifications, among species. With the

rapid progress in the field of genetics and genomics, it is to be expected that such studies will allow

for fundamental new insights into the traits and genetic underpinnings that promote biodiversity

effects in more natural systems.

Sustaining a growing global human population currently depends on increasing nutrient inputs to

crop production systems[28], on breeding[29], and on the use of within-individual diversity effects

termed heterosis[30]. Biodiversity effects have received relatively little attention in breeding and

traditional agriculture, with the notable exception of crop rotation[31] and intercropping of cultivars

and species[32–34]. Despite significant recent advances in automated, large-scale agricultural

management, a large trait variation within a crop field - as found in intercropping - is generally

considered undesirable. Our work shows that niche complementarity-mediated productivity

increases can operate by allelic diversity at a single locus and in a near-isogenic background. Given

the genetic tractability of such effects, it may thus be possible to breed mixtures of genotypes that

support increased yields through niche complementarity while maintaining low variation in

economically relevant traits. Overall, incorporating allelic diversity effects on productivity and soil

legacy into breeding programmes could help bypass constraints imposed on the performance of

single genotypes, by shifting optimization efforts from the individual to the system level[35]

Methods

Germplasm

The Shadhara and Bayreuth accessions were kindly provided by Nuno Pires (University of Zurich)

and had originally been obtained from the Nottingham Arabidopsis stock center (NASC). The 18

RILs (representing the "RIL-minimal set" and line 33RV191 used to generate NILs (all contained in

the core-pop set of 165 lines) were ordered from the Versailles Arabidopsis stock center

(*http://publiclines.versailles.inra.fr*) and propagated in a growth chamber. A Bay×Sha RIL

(33RV191) was confirmed to be heterozygous at two PCR marker positions on chromosome 4

(Table S2). Upon selfing of this line, the two NILs 33RV191-Sha and 33RV191-Bay were isolated,

and their genomes were re-sequenced as described below. Furthermore, after selfing of another

single heterozygous F_{10} individual of line 33RV191, we screened 160 offspring for recombination

between the ShaBa5, ShaBa6 and ShaBa8 markers on chromosome four. Upon selfing of 23

putative recombinant offspring, we isolated 19 homozygous recombinant lines for which we

confirmed a recombination event in the region by PCR. We then performed whole-genome re-

sequencing to confirm the isogenic background and to infer recombination breakpoints for this

heterogeneous inbred family (referred to as NILs throughout the text) as described below.

<u>Soils and growth conditions</u>

Soils consisted of different mixtures of a peat and nutrient rich soil (Einheitserde ED73; pH ~5.8, N

250 mg/l; P2O5 300 mg/l; 75% organic matter content; Gebrüder Patzer GmbH, Sinntal-Jossa,

Germany) and finely grained quartz sand, and pot size for all mixture experiments was 7×7×8 cm.

The experiment using the parental lines Bay and Sha was replicated on a soil quality gradient with

sand contents of 0%, 40%, 75% and 80%, which resulted in a near-linear decrease of pot-

productivity from the highest to lowest ED73 content (Figure S1). For the rough-mapping of the

diversity effect using RILs, we used a mixture of 80% sand and 20% ED73 mixture. For the fine-

mapping diallel using NILs, we used either a 80%:20% or a 20%:80% sand:ED73 mixture.

Seeds were sown directly onto soils (approx. 10 seeds per position, 4 positions per pot, Fig. 1a).

The pots were placed in growth chambers or greenhouse compartments and covered with plastic

lids to maintain a high humidity for germination and initial seedling establishment. Additional light

was provided if necessery, achieving a photoperiod of 14–16 hours. Day-time and night-time

temperatures were maintained around 20–25 °C and 16–20 °C, respectively. Seedlings were thinned

continuously until a single healthy seedling remained per position.

Once seedlings were established, the pots were placed in a greenhouse compartment with automated

watering (every 48 hours). In summer 2015, daytime temperatures were extremely high, and the

first block of the RIL competition diallel was therefore grown in a growth chamber with full climate

control (8hrs night/16 hrs day; 60% humidity; 18/23°C night/day temperature). The second block

was grown in the growth chamber for a month before it was re-located to the regular greenhouse

compartment.

Pots that did not contain all four originally planted individuals were discarded. Plants were

harvested 43–51 days after sowing, with the specific harvest date determined by the occurence of

approx. 5–10 dehiscent siliques on the earliest flowering genotypes within a block.

After the competition diallels were harvested, soil feedback trials were established by dividing the

soil of a pot into two smaller pots of 5.5×5.5×6.0 cm dimension and sowing the respective

phytometers (Bay or Sha for the RIL diallel, 33RV191-Sha or 33RV191-Bay for the NIL diallel)

directly onto the soil. Again, seeds were oversown and seedlings thinned continously until a single

healthy individual remained. Phytometer experiments were harvested either 36 days after sowing

(peat-rich soil remaining after the NIL diallel, harvested early because plant roots started to grow

out of the pots) or 49–58 days after sowing (sand-rich soil, each block was harvested on a single

day). For all experiments, the position of the individual pots was randomized across trays for

seedling establishment, and across watering tables after seedling establishment. Throughout the

experiment, pots were replaced randomly within trays and tables every 7–10 days. Pots were

watered *ad libitum*, and in case of high population densities of dark-winged fungus gnats, the

systemic insectizide ActaraG (Syngenta Agro AG) was applied according to the manufacturers

recommendation. After harvesting, plant biomass was dried at 65°C for at least three days before

weighing.

Experimental Designs

To test soil effects on biodiversity effects in mixed Shadhara-Bayreuth communities, four soil

substrates varying in sand content were prepared as described above. We then grew 12 replicate

monocultures of each accession plus 24 replicate mixtures per soil type (total of 48×4= 192 pots).

The RIL competition diallel consisted of a half diallel replicated in four blocks. All pair-wise RIL

combinations were realized once per block except for RIL monocultures which were replicated

twice. For the follow-up soil feedback experiment, we re-used soil from only the first two blocks of

the competition diallel. We re-mixed the soil of each single pot after harvesting the plants, and re-

distributed it into two smaller plots that were sown with either a Shadhara or Bayreuth parental

genotype that served as phytometers.

The NIL competition diallel used for fine-mapping was realized in a single block that contained all

pair-wise combinations of the 19 NILs including monocultures. The subsequent soil feedback stage

was realized as described for the RIL diallel, using either 33RV191-Bay or 33RV191-Sha genotypes

as phytometers.

Genotyping and line re-sequencing

PCR-based genotyping assays (Table S2) were developed based on deletions in the Sha genome as

predicted by the Polymorph tool (available at http://polymorph.weigelworld.org)[36].

Barcoded libraries for genome resequencing were prepared using the Illumina Nextera DNA

Library Prep Kit (FC-121-1031, Illumina Inc. San Diego, CA) in combination with the Nextera

Index Kit (96 indices, FC-121-1012) and pair-end sequenced on an Illumina HiSeq 2500 (2x150 bp,

rapid run). The clustering and sequencing were performed at the Functional Genomics Center

Zurich. Sequences were aligned to the Arabidopsis genome (Col-0 genome, TAIR version 10) using

BWA[37], aligned read sorting and variant calling were performed using samtools[38]. Aligned genomic

sequences of the parental accessions Bay-0 and Sha were downloaded from the 1001 genomes

project data center (http://1001genomes.org). The VCF-file produced by the samtools software was

loaded into the R Statistical Software [39], where the subsequent analyses were performed: firstly,

variant calls were filtered (for differences in genotype calls between the Sha and Bay genomes,

quality of variant calls, population-level minimal minor allele frequency 0.2; maximum

heterozygosity 0.2). Inference of genotype calls at polymorphic sites was performed as described

previously[40] and inference of parental alleles was improved using functionality implemented in the

MPR package[40]. Genotype reconstruction was then performed in R using a simple hidden Markov

model as implemented in the R package HMM, with hidden state starting probabilities (Bay, Het or

Sha) all set to 1/3, and transition probabilities from one state to itself set to 0.99998 and to the other

two states set to 0.00001 each. Emission probabilities of genotype calls given a state, e.g. Bay, were

set to 0.35, 0.25, 0.25, 0.15 for genotypes calls Bay, Het, Sha or missing, etc.

<u>Statistical analyses</u>

We analyzed data from the diallel experiments using linear models summarized by analysis of

variance (ANOVA). The model terms included, in this order, the general combining abilities (GCA)

of genotypes (a factor with 20 levels in the RIL diallel and 19 levels in the NIL diallel), the

genotype diversity in the pot (GD, 1 or 2 genotypes), the allele identity in the genotype

monocultures (A, Sha or Bay), the allelic diversity in the genotype mixtures (AD, 1 [Sha/Sha or

Bay/Bay] or 2 [Sha/Bay]), and the genotype composition planted in the pot (comp). The factor GCA

was created by superimposing the model matrices for factors coding for the first and second

genotype (each a factor with 20/19 levels). The significance of GD, A, and AD were determined

using F-tests with comp as error term (denominator). A and AD were encoded in such a way that

these contrasts applied only to genotype monocultures and mixtures, respectively. Technically, this

was achieved by including a third level in the factor that did not vary in the other group. Fitting A

and AD after GD therefore only explained variance in these subsets. The diallel model was

extended by additional terms and the corresponding interactions when these applied; specifically,

the RIL diallel included a block effect. The NIL diallel included terms for soil type, and interactions

of all the terms above with soil type (for example, soil×AD was tested using soil×comp as error

term). The soil feedback experiments included further interactions with phytometer (RIL and NIL

diallel), and phytometer×soil (NIL diallel). Effects of pot biomass in the competition diallel (diallel

biomass and diallel biomass × soil) were accounted for in these linear models, and data were square-

root transformed to obtain normally distributed random errors.

Specific combining abilities for mapping were calculated directly, within blocks, by solving the

linear model m = X GCA + SCA where X is the design matrix describing the genotype composition

of a pot. Monoculture SCAs were also determined but not used for QTL mapping of allelic diversity

within RIL mixtures. In the RIL diallel, the SCAs of each genotype composition was first calculated

per block and then aggregated over all blocks using least-square estimates. Marker regression was

performed using the ghlt-function provided by the multcomp package[41] and contrasting SCAs of

mono-allelic RIL mixtures ("BB" and "SS" compositions) with mixed-allelic mixtures ("BS"

compositions). QTL mapping was also performed using the R/qtl package and interval mapping

(scanone-function), with both mono-allelic compositions at a given locus re-coded as to the same

level ("mono-allelic") and compared against mixed-allelic compositions. Genome-wide significance

was assessed by resampling (5000x).

References

1.   Cardinale, B. J. *et al.* The functional role of producer diversity in ecosystems. *Am. J. Bot.* **98,**
572–592 (2011).

2.   O'Connor, M. I. *et al.* A general biodiversity–function relationship is mediated by trophic
level. *Oikos* **126,** 18–31 (2017).

3.   Cadotte, M. W. Functional traits explain ecosystem function through opposing mechanisms.
Ecology Letters **20,** 989–996 (2017).

4.   Wright, A. J., Wardle, D. A., Callaway, R. & Gaxiola, A. The Overlooked Role of Facilitation
in Biodiversity Experiments. *Trends in Ecology and Evolution* **32,** 383–390 (2017).

5.   Ratcliffe, S. *et al.* Biodiversity and ecosystem functioning relations in European forests
depend on environmental context. *Ecology Letters* **20,** 1414–1426 (2017).

6.   Schnitzer, S. A. *et al.* Soil microbes drive the classic plant diversity-productivity pattern.
Ecology **92,** 296–303 (2011).

7.   Maron, J. L., Marler, M., Klironomos, J. N. & Cleveland, C. C. Soil fungal pathogens and the
relationship between plant diversity and productivity. *Ecol. Lett.* **14,** 36–41 (2011).

8.   Loreau, M. & Hector, A. Partitioning selection and complementarity in biodiversity
experiments. *Nature* **412,** 72–76 (2001).

9.   Tylianakis, J. M. *et al.* Resource heterogeneity moderates the biodiversity-function
relationship in real world ecosystems. *PLoS Biol.* **6,** 947–956 (2008).

10. Flynn, D. F. B., Mirotchnick, N., Jain, M., Palmer, M. I. & Naeem, S. Functional and
phylogenetic diversity as predictors of biodiversity--ecosystem-function relationships.
Ecology **92,** 1573–1581 (2011).

11. Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428,** 821–827 (2004).

12. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529,** 167–171 (2015).

13. Cadotte, M. W., Cardinale, B. J. & Oakley, T. H. Evolutionary history and the effect of
biodiversity on plant productivity. *Proc. Natl. Acad. Sci. USA* **105,** 17012–17017 (2008).

14. Crutsinger, G. M. *et al.* Plant genotypic diversity predicts community structure and governs
an ecosystem process. *Science.* **313,** 966–968 (2006).

15. Wang, S. & Loreau, M. Ecosystem stability in space: α, β and γ variability. *Ecol. Lett.* **17,**
891–901 (2014).

16. Oehri, J., Schmid, B., Schaepman-Strub, G. & Niklaus, P. A. Biodiversity promotes primary
productivity and growing season lengthening at the landscape scale. *Proc. Natl. Acad. Sci.*
38, 10160–10165 (2017).

17. Siefert, A. *et al.* A global meta-analysis of the relative extent of intraspecific trait variation in
plant communities. *Ecology Letters* **18,** 1406–1419 (2015).

18. Crutsinger, G. M., Souza, L. & Sanders, N. J. Intraspecific diversity and dominant genotypes
resist plant invasions. *Ecol. Lett.* **11,** 16–23 (2008).

19. Prieto, I. *et al.* Complementary effects of species and genetic diversity on productivity and
stability of sown grasslands. *Nat. Plants* **1,** 15033 (2015).

20. Mendel, G. Versuche über Pflanzenhybriden. in *Verhandlungen des naturforschenden*
Vereines in Brünn, Band IV für das Jahr 1865 3–47 (1865).

21. Loudet, O., Chaillou, S., Camilleri, C., Bouchez, D. & Daniel-Vedele, F. Bay-0 x Shahdara
recombinant inbred line population: A powerful tool for the genetic dissection of complex
traits in Arabidopsis. *Theor. Appl. Genet.* **104,** 1173–1184 (2002).

22. Griffing, B. Concept of General and Specific Combining Ability in Relation to Diallel
Crossing Systems. *Aust. J. Biol. Sci.* **9,** 463–493 (1956).

23. Griffing, B. Genetic-Analysis of Plant Mixtures. *Genetics* **122,** 943–956 (1989).

24. Clements, F. E. & Goldsmith, G. W. *The Phytometer Method in Ecology*. (Carnegie
Institution of Washington, 1924).

25. MacKay, T. F. C., Stone, E. A. & Ayroles, J. F. The genetics of quantitative traits: Challenges
and prospects. *Nature Reviews Genetics* **10,** 565–577 (2009).

26. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From
Polygenic to Omnigenic. *Cell* **169,** 1177–1186 (2017).

27. Wilfert, L. & Schmid-Hempel, P. The genetic architecture of susceptibility to parasites. *BMC*
Evol. Biol. **8,** 187 (2008).

28. Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R. & Polasky, S. Agricultural
sustainability and intensive production practices. *Nature* **418,** 671–677 (2002).

29. Edgerton, M. D. Increasing Crop Productivity to Meet Global Needs for Feed, Food, and
Fuel. *Plant Physiol.* **149,** 7–13 (2009).

30. Lippman, Z. B. & Zamir, D. Heterosis: revisiting the magic. *Trends in Genetics* **23,** 60–66
(2007).

31. Wittwer, R. A., Dorn, B., Jossi, W. & Van Der Heijden, M. G. A. Cover crops support
ecological intensification of arable cropping systems. *Sci. Rep.* **7,** 41911 (2017).

32. Finckh, M. R. *et al.* Cereal variety and species mixtures in practice, with emphasis on disease
resistance. *Agronomie* **20,** 813–837 (2000).

33. Zhu, Y. *et al.* Genetic diversity and disease control in rice. *Nature* **406,** 718–722 (2000).

34. Litrico, I. & Violle, C. Diversity in Plant Breeding: A New Conceptual Framework. *Trends in*
Plant Science **20,** 604–613 (2015).

35. Weiner, J., Du, Y. L., Zhang, C., Qin, X. L. & Li, F. M. Evolutionary agroecology: individual
fitness and population yield in wheat (Triticum aestivum). *Ecology* **98,** 2261–2266 (2017).

36. Fitz, J. *et al.* personal communication.

37. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows − Wheeler
transform. *Bioinformatics* **25,** 1754–1760 (2009).

38. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078–
2079 (2009).

39. R Core Team. R: A Language and Environment for Statistical Computing. *R Foundation for*
Statistical Computing, Vienna, Austria ISBN 3-900051-07-0 (2017). http://www.R-
project.org/

40. Xie, W. *et al.* Parent-independent genotyping for constructing an ultrahigh-density linkage
map based on population sequencing. *Proc. Natl. Acad. Sci.* **107,** 10578–10583 (2010).

41. Hothorn, T., Bretz, F. & Westfall, P. Simultaneous inference in general parametric models.
Biometrical Journal **50,** 346–363 (2008).

Acknowledgements

We thank Bernhard Schmid and Ueli Grossniklaus for helpful discussions and generously sharing

infrastructure and lab resources. We further acknowledge Matthias Philipp for technical support,

Enrica De Luca and Nicole Ponta for help with plant measurements, Matthias Furler and Dorde

Topalovic for technical greenhouse support. This work was supported by an Ambizione Fellowship

(PZ00P3_148223) of the Swiss National Science Foundation (to S.E.W). P.A.N acknowledges

support by the University of Zurich Priority Programme "Global Change and Biodiversity". S.E.W

was also financially supported by funds of the University of Zurich and the European Research

Council (to Ueli Grossniklaus).

Author contributions

377 S.E.W. conceptualized and designed the research (with input from P.A.N.) and performed the

378 experiments. Both authors performed the analyses and wrote the manuscript. Both authors read and

379 approved the final version of the manuscript.

Data availability

382 The datasets described in the paper are available through the DataDryad repository (DOI:XXX).

383 Sequencing data are deposited in the NCBI Short-Read Archive (accession XXX).

384 Analysis scripts are available from the authors upon request.

Competing interests

387 The authors declare no competing financial interests.

Figure 1: Combining ecological concepts and genetic methods

a. Pot systems used to study diversity effects in pair-wise genotype mixtures. The inset shows a Recombinant Inbred Line (RIL) monoculture (left) and mixture (right) **b.** Net diversity effects in Bay-Sha mixtures along a peat-sand substrate gradient. Error bars denote standard errors of means (s.e.m.). **c.** Outline of the competition diallel design, the genotypes used throughout this paper. 18

RILs and the two parental accession, or 19 near-isogenic lines (NILs) are each placed in competition with each other, allowing to assess i) effects of genotypic mixture (i.e. diagonal vs off-diagonal), or ii) effects of allelic mixture at a given locus across all genotype mixtures (i.e. comparing SS and BB vs SB) on pot productivity. SCA = specific combining abilities. **d.** Outline of the experimental procedure used in this work. Colored labels indicate measured variables. GCA = general combining ability.

Figure 2: Allelic diversity at a major effect locus increases community productivity

a. Pot-level productivity in dependence of community type (mix = RIL mixtures vs mono = RIL monocultures), showing positive genotype mixture effects in the diallel and on sand-rich soil (values aggregated across four blocks). **b.** Quantitative trait locus interval mapping of allelic diversity effects on SCA. Vertical lines denote 10% and 5% genome-wide significance levels. The inset shows estimated SCA (± s.e.m.) across genotype mixtures that exhibit different allelic compositions at marker MSAT4.9 on chromosome 4. LOD: logarithm of the odd.

Figure 3: Resolving soil × allelic diversity interactions to a single Mendelian factor

a. Re-sequencing of NILs differing only on lower arm of chromosome four for fine-mapping. Shown are genotype calls at all polymorphic sites across the genome (B = homozygous for the Bay allele, het = heterozygous, S = Sha allele) in either NIL r10 (blue, top left) or NIL r96 (red, top right), as well as an overlay of the two line's genotype calls (bottom). Vertical dotted lines separate different chromosomes. **b.** Reconstructed genotypes across chromosome four of the 19 NILs used for fine-mapping. Each bar represents a single NIL. **c.** and **d.** Map of allelic diversity effects across

chromosome four, on either peat-rich soil (c.), or sand-rich soil (d.). The widths of the bars indicate

the size of the regions in which no recombination events were inferred across the whole population.

Inset in d. shows the mean ± s.e.m. SCA across allelic compositions at locus Chr4@16.92.

Figure 4: Allelic diversity effects persist across a generation through their soil legacy

a. Scheme of the genotype-combinations used for either the RIL vs. NIL diallels (top), and the

phytometer genotypes used in the soil-feedback phase (bottom). **b.** Mean phytometer (Bay or Sha,

+/- s.e.m.) performances on legacy soil derived from RIL mixtures with different allelic

compositions at marker MSAT4.9. **c.** Mean phytometer (NIL-Bay or NIL-Sha, +/- s.e.m.)

performances on legacy soil derived from NIL mixtures with different allelic compositions at locus

Chr4@16.92 on either sand-rich (left) or peat-rich (right) soil. Values were square-root transformed.

Extended Data

Table S1. Legacy effects of soils conditioned in RIL and NIL diallel experiments.

Effects were quantified using two phytometers (factor 'phy'; Sha and Bay accessions in RIL diallel,

and near-isogenic lines bearing Sha and Bay allele at putative effect locus in NIL diallel). The NIL

diallel additionally was replicated on substrates differing in sand content (factor 'soil'). The term

'GCA' (general combining abilities) indicates average genotype-specific soil conditioning effects on

phytometers. 'SCA' (specific combining abilities) captures deviations in yield from additive

predictions made using GCAs. Within SCA, the following contrasts were tested: GD: Genotype

diversity, i.e. whether genotype monocultures differed in feedback effects from two-genotype

mixtures; A: allele-specific differences at marker MSAT4.9 (RIL diallel) and Chr4@16.92 (NIL

diallel) within genotype monocultures; AD: allele-diversity effects within genotype mixtures. df and

ddf indicate nominator and denominator degrees of freedom of corresponding F-tests. *** P<0.001;

** P<0.01; * P<0.05; (*) P<0.1; n.s. not significant.

Terms and contrasts	RIL diallel			NIL diallel		
	df	Denominator: ddf	Signif.	df	Denominator: ddf	Signif.
GCA	18	comp: 167	(*)	18	comp: 168	n.s.
SCA						
GD	1	comp: 167	(*)	1	comp: 168	n.s.
A	1	comp: 167	n.s.	1	comp: 168	n.s.
AD	1	comp: 167	n.s.	1	comp: 168	n.s.
Phy × GCA	18	phy × comp: 166	n.s.	18	phy × comp: 168	*
Phy × SCA						
Phy × GD	1	phy × comp: 166	n.s.	1	phy × comp: 168	n.s.
Phy ×A	1	phy × comp: 166	n.s.	1	phy × comp: 168	n.s.
Phy ×AD	1	phy × comp: 166	**	1	phy × comp: 168	*
Soil × GCA				18	soil × comp: 150	n.s.
Soil × SCA						
Soil × GD				1	soil × comp: 150	n.s.
Soil ×A				1	soil × comp: 150	n.s.
Soil ×AD		does not apply		1	soil × comp: 150	n.s.
Phy × Soil × GCA				18	phy × soil × comp: 146	n.s.
Phy × Soil × SCA						
Phy × Soil × GD				1	phy × soil × comp: 146	***
Phy × Soil ×A				1	phy × soil × comp: 146	n.s.
Phy × Soil x AD				1	phy × soil × comp: 146	n.s.

Table S2: List of PCR markers used in this study.

Primer	Assay	Sequence	Predicted location	Pred. fragment sizes (Sha/Bay)
SW-182	ShaBa5	ACGTATTTCGATGTATGGTCCTTG	Chr4: 16044156	550/664
SW-183		TCACGTGAATCGTATTCGTTGAAG		
SW-184	ShaBa6	CTTCTCCGCTTCAACCTCTGC	Chr4: 17709750	600/632
SW-185		AATCCAGGATTCAGAGTTGCTTTC		
SW-188	ShaBa8	TTGATTAGGGCTACGAGGATAAGG	Chr4: 16707214	408/609
SW-189		GAGTCTATTAATTATGCTTGGTGC		

442 **Figure S1: Productivity and complementarity in Bay-Sha mixtures across a peat-sand**

443 **gradient**

444 **a.** Pot-level biomass measurements of Sha and Bay monocultures or pair-wise mixtures along a

445 sand/peat substrate gradient. **b.** Complementarity and selection effects calculated according to the

446 additive partitioning[8] method along the substrate gradient. Error bars denote s.e.m.

Figure S2: A major effect locus driving complementarity between genotypes

a. QTL mapping SCA variation across allelic diversity levels using a marker regression technique by contrasting SCAs of mono-allelic RIL mixtures (BB and SS) with bi-allelix mixtures (BS) **b.** Pot-level productivity of each genotype composition (averages of four blocks) in dependence of allelic composition at marker MSAT4.9.

Figure S3: Fine-mapping and persistence of allelic diversity effects in near-isogenic lines

a. The comparison of the reconstruced genotype of NIL r10 (HMM Viterbi-path across all

chromosomes, homozygous on lower arm of chromosome 4) in comparison to publicly available

molecular marker-based genotyping data of the ancestral line from which it was derived

(heterozygous on lower arm of chromosome 4) – showing a high degree of congruence between the

re-constructed genotype based on whole-genome resequencing and the marker data. **b.** Schematic

outline of a possible cause of the high mapping resolution achieved through the competition diallel

design. A major advantage of the design is the joint dependency of community-level allelic

diversity on recombinations *within* and *between* recombinant inbred lines, such that mapping

resolution increases very quickly. **c.** Extreme example of SCA variation across genotype mixtures

all containing one specific genotype (NIL r146, homozygous for the Bay-allele at locus

Chr4@16.92), either in combinations with NILs carrying the Sha-allele (dark blue bars) or the Bay-

allele (dark red bars). Shown are the data from sand-rich soil only.

Figure S4: Soil-feedback experiments

a. Genotype calls at all polymorphic sites across the genome (B = homozygous for the Bay allele, H

= heterozygous, S = homozygous for the Sha allele) obtained by genome re-sequencing phytometer

genotypes NIL-Bay (33RV191-Bay, top left, red) and NIL-Sha (33RV191-Sha, top right, blue). The

overlay of the genotype calls of both lines (bottom) confirms that these lines are isogenic but for

variation on lower arm of chromosome four. The two phytometers were employed on soil derived

from the NIL diallel. **b.** and **c.** QTL mapping by marker regression of phytometer-specific responses

to soil legacy of previous generation allelic diversity (i.e. allele-diversity × phytometer (Phy×AD)

interaction in Supplemental Table S1, but in a model without adjusting for diallel pot biomass). The

mapping of diversity effects through such influences on soil legacy could have been applied to the

identification and fine-mapping of the same major effect locus, without ever measuring biomass

productivity (albeit according to slightly relaxed statistical criteria). Shown are negative $\log_{10}$-

transformed P-values for each marker position in the RIL (b.) or genotype block in the NIL (c.)

diallels.

Supplementary Discussion

As outlined in Figures 1 and 4, and as described in the Methods, we performed two soil-feedback

experiments to test whether allelic diversity effects extend across generations after the plants were

destructively harvested. Possible effects solely explainable by variation in plant productivity during

the soil training phase (e.g. nutrient drawdown or environmental correlations) were accounted for in

the model (terms *diallel biomass* and *diallel biomass×soil* as described in the Methods section).

Significantly different soil conditioning through allelic diversity, as assessed by phytometer

performance in a next generation, can thus be seen as an extended "phenotype" (i.e. a legacy of

allelic mixture that extends throughout generations), interactions thereof with phytometer genotype

as asymetric responses (potential trade-offs) of phytometers to soil-borne factors affected by the

conditioning (term "Phy×AD" in Supplemental Table S1).

We noted, however, that the exact mechanisms underlying both soil training or phytometer

responses await further experiments, since the response patterns (Fig. 4) do not allow for a simple

mechanistic model. One problem that impedes the inference of underlying mechanisms (such as the

anticipation of specific soil factors, and their interaction with genetic variants, etc.), is the number

of variables differed between the two experiments because of constraints on experimental design: in

the RIL diallel, we used the two parental accession Bay and Sha as phytometers and a split-plot

design to test for differential responses of these phytometer to soil conditioning. After the NIL diallel, we used two near-isogenic phytometer (NIL-Bay and NIL-Sha) as phytometers in a similar test for differential responses to soil conditioning. Naively, we could expect an approximate congruence of how the different phytometers carrying the same alleles at the diversity locus responded between experiments, i.e. the parental genotype Bay in the RIL diallel (Figure 4 b, yellow lines) should exhibit approximately the same response pattern to allelic diversity legacy as the NIL-Bay genotype in the NIL diallel (Figure 4 c, yellow lines). However, for example it seems that the Bay genotype grown on RIL diallel soil responded somewhat positively to conditioning by allelic diversity (a pattern opposite to what would be expected from simple biomass-corresponding resource drawdown during the diallel experiment - and indicating that diversity led to the increased "removal" of a negatively acting soil factor, or the addition of a positively acting soil factor), whereas the NIL-Bay genotype in the NIL diallel responded somewhat negatively to conditioning by allelic diversity. Variables differing between the experiments were 1) the environment (e.g. the different dates at which the experiments were conducted, different batches of soil, greenhouse conditions, etc), 2) the genetic variation in the two diallel population (RILs vs NILs), whereby the genetic variance due to epistasis is expected to differ strongly between these two population, and 3) phytometer genotypes (parental lines vs NILs). We chose to vary the phytometer genotypes because i) we did not have NILs available for the RIL diallel, yet ii) aimed to demonstrate that asymetric soil legacy responses could arise in a near-isogenic background in the NIL diallel.

With these multiple variables differing between the two soil-feedback experiments, it is hard to explain why our naive expectation (outlined above) was not exactly met. However, and most importantly, the experiments still convincingly show that i) allelic diversity effect extend across generations through soil conditioning, and ii) phytometer genotype determines the response to such conditioning in either experiment (phytometer×allelic diversity interaction, see Table S1). Furthermore, both of these arguments apply to the NIL experiments, where the population-level

520   genetic variation (both in the training and feedback phase) was restricted to a single region in an

521   otherwise isogenic background. To us, this suggests that positive productivity effects and soil

522   conditioning are related by mechanism, but the specific details remain to be uncovered.