Competence Inhibition by the XrpA Peptide Encoded Within

the *comX* Gene of *Streptococcus mutans*

Justin Kaspar, Robert C. Shields, and Robert A. Burne[*]

Department of Oral Biology, University of Florida, Gainesville, Florida 32610

Running title: Antagonism of the ComRS-XIP circuit

Keywords: quorum sensing, intercellular signaling, transcription factor, genetic competence,

dental caries

[*]Corresponding author:

Mailing address: Department of Oral Biology, University of Florida, College of Dentistry,

P.O. Box 100424, Gainesville, FL 32610.

Phone: (352) 273-8850

Fax: (352) 273-8829

E-mail: rburne@dental.ufl.edu

SUMMARY

Streptococcus mutans displays complex regulation of natural genetic competence.

Competence development in *S. mutans* is controlled by a peptide derived from ComS (XIP);

which along with the cytosolic regulator ComR controls the expression of the alternative sigma

factor *comX*, the master regulator of competence development. Recently, a gene embedded

within the coding region of *comX* was discovered and designated *xrpA* (com<u>X</u> <u>r</u>egulatory <u>p</u>eptide

<u>A</u>). XrpA was found to be an antagonist of ComX, but the mechanism was not established. In

this study, we reveal through both genomic and proteomic techniques that XrpA is the first

describe negative regulator of ComRS systems in streptococci. Transcriptomic and promoter

activity assays in the Δ*xrpA* strain revealed an up-regulation of genes controlled by both the

ComR- and ComX-regulons. An *in vivo* protein crosslinking and *in vitro* fluorescent polarization

assays confirmed that the N-terminal region of XrpA were found to be sufficient in inhibiting

ComR-XIP complex binding to ECom-box located within the *comX* promoter. This inhibitory

activity was sufficient for decreases in P*comX* activity, transformability and ComX accumulation.

XrpA serving as a modulator of ComRS activity ultimately results in changes to subpopulation

behaviors and cell fate during competence activation.

ABBREVIATED SUMMARY

Streptococcus mutans displays complex regulation of natural genetic competence,

highlighted by a novel gene, *xrpA*, embedded within the coding region for the master regulator

ComX. We show that XrpA modulates ComRS-dependent activation of *comX* expression,

resulting in changes to sub-population behaviors, including cell lysis. XrpA is the first described

inhibitor of a ComRS system and, because it is unique to *S. mutans* it may be targetable to

prevent diseases caused by this pathogen.

INTRODUCTION

Bacterial biofilm communities coordinate behaviors in response to environmental stimuli through the use of chemical mediators that accumulate extracellularly to activate transcription of specific genes when a critical concentration is achieved, in a process termed quorum sensing (Fuqua *et al.*, 1994). In Gram-negative bacteria, diffusible acylated homoserine lactones are the principal chemical mediator that act as a proxy for cell density (Papenfort and Bassler, 2016), whereas small hydrophobic peptides fulfill a similar role in Gram-positive bacteria (Håvarstein *et al.*, 1995). More recently, quorum sensing and gene products involved in intercellular signaling have been highlighted as an area of interest for therapeutic intervention in some bacterial infections, because quorum sensing often controls the transcription of genes that contribute to virulence (Greenberg, 2003).

Genetic competence, a transient physiological state in which bacteria produce the gene products necessary for uptake of DNA from their environment was one of the first described and studied quorum sensing pathways (Tomasz, 1965). Decades later, genetic competence is still a valuable model system in molecular microbiology to unravel the complexities of signal perception, signal transduction and sub-population behaviors, and is of relevance to newer areas of research that include sociomicrobiology, interspecies antagonism and cooperativity, and microbial biogeography (Whiteley *et al.*, 2017). Genetic competence has been extensively studied in the genus *Streptococcus*, including *Streptococcus pneumoniae* (Hui *et al.*, 1995; Straume *et al.*, 2015), *Streptococcus thermophilus* (Fontaine *et al.*, 2009; Gardan *et al.*, 2013), *Streptococcus pyogenes* (Mashburn-Warren *et al.*, 2012; Wilkening *et al.*, 2015) and *Streptococcus mutans* (Li *et al.*, 2001; Son *et al.*, 2012). Interestingly, streptococci harbor two distinct peptide signaling systems that activate genetic competence: the Mitis and Anginosus groups utilize an extracellular signaling system composed of a signaling peptide termed CSP (competence stimulating peptide) and a two-component signal transduction system encoded by *comDE*. In contrast, the Bovis, Salivarius and Pyogenic streptococci employ an intercellular

signaling system that consists of the signal peptide XIP (*comX/sigX* inducing peptide) derived

from the ComS precursor, and a cytosolic Rgg-like regulator designated as ComR (Håvarstein,

2010). While these two signal systems diverge substantially in their distribution among species

and how the systems perceive and transduce their signals, stimulation of either pathway with

the cognate peptide results in the activation of transcription of an alternative sigma factor

termed ComX or SigX. ComX controls the transition into the competent state by activating the

expression of a regulon encoding gene products necessary for DNA uptake and processing.

The Mutans group of streptococci, including the human caries pathogen *S. mutans*, are

unique in that most strains encode an apparently functional ComCDE as well as ComRS

pathways. Further, either signal peptide (CSP or XIP) can trigger up-regulation of *comX*,

although different conditions, including pH, redox and growth phase, influence how effectively

each pathway is able to function (Hagen and Son, 2017). Addition of synthetic CSP (sCSP) to

growing cultures of *S. mutans* in a peptide-rich medium, such as BHI, results in activation of

transcription of genes for the biogenesis of bacteriocins via direct binding of phosphorylated

ComE to a conserved sequence in the promoter regions of these genes and operons.

Consistent with this observation, the ComCDE system of *S. mutans* appears to have evolved

from a common ancestor of the BlpCRH system of *S. pneumoniae*, which does not regulate

competence but does induce bacteriocins in the pneumococcus and some related organisms

(Johnston *et al.*, 2014). Transcription of *comX* can also be induced by CSP, but this generally

occurs in only a subset of organisms in a population, it does not involve direct binding of ComE

to the *comX* promoter, and the underlying mechanism for CSP-dependent activation of *comX* is

not well-understood (Kreth *et al.*, 2007; Hung *et al.*, 2011). The proximal regulator for direct

activation of competence is ComRS. When synthetic XIP (sXIP) is added to a peptide-free,

chemically defined medium, such as FMC or CDM, it is imported into the cytosol by the

oligopeptide permease OppA and forms a complex with ComR to activate *comX* transcription in

the entire bacterial population (Mashburn-Warren *et al.*, 2010; Son *et al.*, 2012). The XIP-ComR

complex can also activate the gene for the precursor of XIP, *comS*, creating a positive feedback

loop for amplification of the competence activation signal (Fontaine *et al.*, 2013). XIP has been

detected in culture supernates, supporting the hypothesis that XIP is a diffusible intercellular

signal (Desai *et al.*, 2012; Khan *et al.*, 2012; Wenderska *et al.*, 2012), although the mechanism

by which XIP is released into the environment can involve active transport (Chang and Federle,

2016) or cell lysis (Kaspar *et al.*, 2017), depending on the species of bacteria. Recently, it was

confirmed experimentally that XIP is able to act as a diffusible intercellular communication

molecule and that signaling can occur within biofilm populations (Shields and Burne, 2016;

Kaspar *et al.*, 2017).

While substantial progress has been made dissecting the mechanisms leading to *com*

gene activation, very little is known about regulation of the system after *comX* is induced and

late competence gene expression is active. In *S. pneumoniae*, shut-off of the ComCDE system

is regulated at multiple levels, including competition between phosphorylated and un-

phosphorylated ComE for binding sites, direct inhibition of activated ComE by the late

competence gene-encoded protein DprA, and by inhibition of ComX activity by an unknown

factor (Martin *et al.*, 2013; Mirouze *et al.*, 2013; Weng *et al.*, 2013). In streptococci that harbor

ComRS systems, the factors that regulate the Com circuit after transcriptional activation have

not been characterized in significant detail. It has been postulated that a "*comZ* gene", under the

control of ComX, exists that encodes a product that acts on the ComR-XIP complex to create a

feedback inhibition loop (Boutry *et al.*, 2013; Haustenne *et al.*, 2015). Recently, we described a

novel protein encoded within *comX* gene in an alternative (+1) reading frame that we

designated as XrpA (*comX* regulatory protein/peptide A) (Kaspar *et al.*, 2015). We described

XrpA as a novel antagonist of *comX* as loss of XrpA, either by mutating the start codon or by

introducing premature stop codons, led to an increase in transformation efficiency and

accumulation of the ComX protein, whereas overexpression of *xrpA* resulted in decreased

transformability and lower levels of ComX. However, the mechanism by which XrpA exerted its

effects was not determined. In this study, we present genetic evidence through transcriptome

profiling and biochemical evidence of protein-protein interactions that demonstrate that XrpA

affects competence development in *S. mutans* by interacting with and inhibiting ComR activity,

thus describing the first negative regulator of competence signaling that acts on the ComRS

circuit.

RESULTS

Transcriptome profiling of an XrpA-deficient strain

In our initial characterization of *xrpA*, we highlighted the unusual transcriptional

characteristics of *xrpA* and the profound influence of XrpA on the dramatically different genetic

competence phenotypes displayed by strains with polar and non-polar mutations in the *rcrR*

gene of the *rcrRPQ* operon, designated Δ*rcrR*-P and Δ*rcrR*-NP, respectively. Inactivation of

xrpA in a way that did not alter the primary sequence of ComX could convert the non-

transformable Δ*rcrR*-NP strain into the hyper-transformable state that was observed for the

Δ*rcrR*-P strain, with concomitant restoration of ComX production (Kaspar *et al.*, 2015). However,

the *rcrR* mutants displayed extreme and unusual phenotypes, so questions remain as to how

xrpA expression is regulated and what role XrpA plays in a wild-type *S. mutans* genetic

background. To begin to answer these questions, we first wanted to compare the

transcriptomes of a Δ*xrpA* strain with that of the *S. mutans* wild-type strain, UA159. For these

studies and those conducted henceforth, the Δ*xrpA* used contains a single base change at the

162^{nd} nucleotide of *comX* (*comX*::T162C), which mutates the *xrpA* start codon (ATG→ACT) and

leaves the *comX* protein coding sequence unchanged (Kaspar *et al.*, 2015).

Comparison of the transcriptome by RNA-Seq of the wild-type with the mutant lacking

XrpA when cells were grown in the chemically defined medium FMC to mid-exponential phase

revealed 56 differentially expressed genes, with 34 upregulated in Δ*xrpA* and 22 downregulated

(Figure 1A). Many of the upregulated genes were competence-related genes, including *comX*

and genes that are a part of the ComX regulon of *S. mutans* (Khan *et al.*, 2016) **(Table S2)**.

Since loss of *xrpA* caused upregulation of competence genes, we also analyzed the

transcriptome of UA159 treated with 2 μM sXIP to induce competence, with UA159 treated with

vehicle (DMSO) as a control. Cells treated with sXIP had 137 genes differentially expressed

compared to the control **(Figure 1B; Table S3)**. Several of the same genes that were the most

strongly upregulated in Δ*xrpA*, including the *comF* and *comY* operons, *drpA* and *lytF*, were also

the highest upregulated genes in the sXIP-treated cells. Interestingly, those genes that were

downregulated in the Δ*xrpA* strain differed from those downregulated by sXIP addition to

UA159. All of the downregulated genes in the Δ*xrpA* mutant were located on the TnSMu1

genomic island that encodes predicted transposases, integrases, transporter(s) and

hypothetical proteins (Waterhouse *et al.*, 2007). This genomic island was recently found to be

differentially expressed in *clpP* and *cidB* mutants, providing additional evidence for a link

between XrpA and stress responses (Chattoraj *et al.*, 2010; Kaspar *et al.*, 2015; Ahn and Rice,

2016).

Meanwhile, the upregulation of the ComX regulon in the Δ*xrpA* mutant could not be

explained by increases in expression of annotated early competence genes. However, when we

examined the region encoding *comS*, which is encoded in the intergenic region of SMU.61 and

SMU.63c in current database annotations, we found an elevated number of reads for *comS* in

the Δ*xrpA* strain compared to UA159 **(Figure 1C)**. When one considers that both *comS* and

comX are upregulated in the XrpA-deficient strain, a clearer picture emerges that XrpA most

likely exerts its influence over competence development by influencing the efficiency of ComR-

dependent activation of the *comX* and *comS* promoters, P*comX* and P*comS*.

XrpA alters comX and comS promoter activity

To determine if XrpA affects the ComR-XIP activated promoters, P*comX* and P*comS*, we

incorporated the *xrpA* start codon mutation *comX*::T162C into strains carrying GFP

transcriptional fusions to each promoter. Similarly, we transformed an *xrpA*-overexpressing

strain (184XrpA) into the GFP reporter gene fusion strains to see if increasing the amount of

XrpA produced would yield gene expression patterns that were opposite of those caused by

loss of *xrpA*. Cells were grown in chemically defined CDM, which allows for self-activation of the

ComRS system. Indeed, loss of *xrpA* resulted in both earlier and higher level expression of

P*comX* and P*comS* reporters, compared to what was observed in the wild-type genetic

background. Conversely, overexpression of *xrpA* from the strong, constitutive promoter on

pIB184 caused a decrease in GFP production from the *comX* and *comS* promoters **(Figure 2)**.

Collectively, these results validate the RNA-Seq data and provide support for the hypothesis

that XrpA negatively affects the activation of gene expression by ComR.

XrpA influences subpopulation responses to competence signals

Stimulation of genetic competence in a peptide-rich medium, such as brain-heart

infusion (BHI), by CSP results in a bimodal response where only a sub-population of cells

activate P*comX* (Son *et al.*, 2012). We reasoned that XrpA might influence the proportion of

cells that responded to sCSP in BHI. We utilized both the wild-type and Δ*xrpA* mutant carrying

P*comX* transcriptional reporter gene fusions and analyzed subpopulation behaviors using flow

cytometry three hours after sCSP addition to planktonic cultures. The percentage of cells that

were GFP-positive was more than 20% greater in the Δ*xrpA* background with addition of 100 nM

sCSP (39.7 ± 2.2 versus 61.5 ± 1.8) **(Figure 3A)** or 1000 nM sCSP (48.7 ± 1.4 compared to

73.9 ± 2.3) **(Figure 3B)**. Further, mean GFP fluorescence intensity was increased in the Δ*xrpA*

background.

Stimulation of the competence cascade by XIP in nanomolar concentrations of wild-type

S. mutans growing in a peptide-free medium results in a unimodal population response, but

addition of sXIP to cultures of *S. mutans* at concentrations higher than 1 µM can trigger cell

death in a significant fraction of the population (Wenderska *et al.*, 2012). To determine if XrpA

could influence XIP-mediated killing in cells treated with higher concentrations of sXIP, we

followed a protocol similar to the CSP experiments, but stained the cells with propidium iodide

(PI) to measure membrane integrity prior to analysis by flow cytometry. No changes were seen

in the proportions of the population that were PI-positive between the wild-type and Δ*xrpA*

background, but a clear increase in the mean GFP intensity was observed when *xrpA* was

mutated, similar to what was seen with CSP **(Figure 3C)**. However, when the *comS* gene was

removed to eliminate the positive feedback loop in the XIP signaling pathway, a distinct increase

in the proportion of PI-positive cells was seen in the Δ*xrpA* mutant population, compared to

behaviors in the wild-type genetic background (36.8 ± 0.7 versus 47.2 ± 0.9) **(Figure 3D)**.

Measurements of eDNA release from overnight cultures were used to confirm the finding that

strains that were activated for competence, but that lack *xrpA*, were more lytic than their wild-

type counterparts **(Figure 3E)**. The propensity for enhanced lysis in the absence of XrpA was

also correlated with decreased biofilm formation when either glucose or sucrose was present as

the sole carbohydrate source **(Figure 3F)**. Taken together, these data highlight that XrpA can

influence subpopulation behaviors both in terms of competence activation in environments

where peptides are present, and influence lytic behaviors associated with activation of the

competence by high, albeit physiologically relevant, concentrations of signal peptide, with the

enhanced cell death most likely being associated with more robust activation of P*comX*.

Interactions between ComR and XrpA

While genetic data from transcriptome profiling and transcriptional reporter experiments

clearly supported a role for XrpA in interference with the ComRS pathway, we wanted to test

whether these proteins could directly interact. We adapted a Strep-protein interaction (SPINE)

protocol (Herzberg *et al.*, 2007) that allows for crosslinking of proteins *in vivo*, followed by

affinity purification of target protein complexes and identification of interacting partners using

mass spectrometry (MS). For this experiment, we chose ComR as the bait by incorporating a C-

terminal Strep-tag® in front of the stop codon. The construct was also engineered in such a way

as to introduce 9 amino acids to serve as a flexible linker sequence to minimize the potential for

disrupting the native conformation of ComR; interaction of small hydrophobic peptides by Rgg-

like regulators occurs with the C-terminal domain of the proteins (Talagas *et al.*, 2016). Cultures

(500 ml) of a strain carrying the Strep-tagged ComR expressed from the strong constitutive

promoter on pIB184, and a vector-only control strain, were grown in CDM to mid-exponential

phase (OD_{600} = 0.6), at which point either 2 µM sXIP or an equivalent volume of 0.1% DMSO

control were added. After growth for an additional hour (OD_{600} = 0.8), the homobifunctional *N*-

hydroxysuccimide ester (DSP) cross-linking agent that is primary amine-reactive and contains a

thiol-cleavable bridge was added to the cells for 45 minutes at 37°C, cells were harvested, and

clarified whole cell lysate were passed over a Strep-Tactin® resin for isolation of the targeted

complex, as detailed in the methods section.

In our initial experiments, purified protein complexes were subjected to SDS-PAGE,

followed by silver staining **(Supplemental Figure 1)**. Five bands of interest that appeared in the

Strep-tagged ComR sample, but not in the vector control sample, were excised from the gel and

identified by mass spectrometry **(Table 2)**. Peptide fragments were identified that were derived

from the transcriptional regulator SgaR (SMU.289) and a putative single-stranded DNA binding

protein Ssb2 encoded by SMU.1967, as well as peptides derived from XrpA. In a second

experiment, the purified protein complexes were subjected to two-dimensional differential gel

electrophoresis (2D DIGE) **(Figure 4)**. A total of 58 spots of interest were selected and a protein

expression ratio (PER) was calculated between triplicate samples **(Supplemental Table S4)**.

Spots chosen for identification were required to have a PER >1.5 compared to the vector-only

control sample in either the sXIP treated or non-treated samples. Peptide fragments were

identified with a high degree of confidence from a putative alcohol-acetaldehyde dehydrogenase

AdhE (SMU.148), a GTP-binding protein TypA (SMU.546), the hypothetical protein encoded by

SMU.1671c and the DNA recombination and repair protein RecA (SMU.2085) **(Table 2)**. Ssb2,

which was found in the 1-D gel, was also identified. Work is currently underway to confirm

whether these identified proteins can interact with ComR and the significance of such

interaction(s) in the context of the integration of competence development with cellular

physiology and stress responses.

Importantly, peptides derived from XrpA were identified from the 2D DIGE and the

fragments detected were derived exclusively from the N-terminus of XrpA (aa 1-38), with the

fragment MFCVSKK being the peptide that was most frequently identified.

The N-terminal Region of XrpA Can Inhibit Competence Development

The results from the SPINE experiment provided evidence for an interaction between

XrpA and ComR. Based on the fragments identified by MS, we postulated that the N-terminal

domain of XrpA may be inhibitory to the function of ComR. To test this hypothesis, we

synthesized several different peptide fragments of XrpA and evaluated their inhibitory capacities

in various assays **(Table 3)**. We first tested the ability of the peptides to interfere with activation

of the P*comX* transcriptional reporter in cells growing in CDM. Of the four peptides tested that

span the entirety of the 69-aa XrpA, only XrpA-1 containing aa 5-20 of the N-terminal region of

the protein caused a marked decreased in P*comX* activity, compared to when only vehicle was

added **(Figure 5A)**. Interestingly, a higher final OD_{600} was also recorded for cultures exposed to

XrpA-1, compared to when either vehicle or any of the other three peptides were tested. To

confirm that this effect was specific for this peptide, a scrambled peptide (same aa composition,

different sequence) was synthesized and tested **(Figure 5B, Table 3)**. While some residual

inhibitory activity was evident with the scrambled peptide, compared with control and other

peptides, much of the effect was alleviated. The inhibitory effect of the XrpA-1 peptide was also

found to be dose-dependent in this experimental setup **(Figure 5C)**. A similar profile for XrpA-1

was seen using the ComR-XIP activated P*comS*::*gf*p transcriptional fusion strain (data not

shown). Notably, one of the most frequently identified peptide fragments of XrpA observed in

the SPINE experiment was not part of the sequence of XrpA-1, so we speculated that using

larger peptides might elicit a stronger inhibitory effect. Peptides N1 and N2 were synthesized

that included aa 1-18 and aa 18-38, respectively, encompassing all identified fragments from

the SPINE experiment. Indeed, these two peptides displayed a greater ability to inhibit the

induction of P*comX* activity in CDM **(Figure 5D)**. Similar to the growth profiles of the

transcriptional reporter strains seen in Figure 5A, XrpA-N1 also displayed better growth,

compared to when XrpA-N2 or vehicle was added. Together, these results confirm that the N-

terminal domain of XrpA can inhibit ComRS-dependent activation of *comX* when provided

exogenously to cells growing in a chemically defined medium.

We also tested several other competence-related phenotypes with the synthetic XrpA

peptides. In terms of transformation efficiency, a 107 ± 13-fold decrease was observed when

XrpA-1 was present, compared to the DMSO only control, and 4 ± 3 and 8 ± 5 fold decreases

were observed when XrpA-2 and XrpA-3 were present, respectively **(Figure 6A)**. In terms of

ComX accumulation, 20-25% less ComX protein was observed by western blowing in cells

treated with XrpA-1 **(Figure 6B)**. A more pronounced loss was seen upon treatment with XrpA-

N1 (30-35%, respectively). It did not appear as if ComR levels were impacted by addition of

either peptide. In fact, more ComR was present by densitometry readings in samples treated

with XrpA-2 or XrpA-3. As the addition of XrpA-1 to growing cultures resulted in higher final

OD_{600} values, we also assessed biofilm formation in the presence of the different XrpA peptides.

A significant decrease in biofilm formation was seen in both glucose- and sucrose-grown

biofilms when XrpA-1 or XrpA-N1 was present, with some decreases, when the other peptides

were added **(Figure 6C)**. An explanation for the higher OD_{600} values when XrpA-1 was present

may be related to reduced cell lysis. eDNA release was measured as described for Figure 3 and

we found significantly less eDNA accumulation in culture supernates from overnight cultures

grown in the presence of XrpA-1, but not with the other XrpA peptides, compared to DMSO-

treated controls. In all, these results support the gene fusion assays and show that the N-

terminal region of XrpA has a significant impact on competence development and biofilm-related

phenotypes when provided exogenously to *S. mutans*.

In vitro interactions of XrpA and ComR

To confirm the interaction between the N-terminal region of XrpA and ComR, we utilized

a fluorescence polarization assay where we could monitor the binding of the ComR-XIP

complex to the promoter region of *comX in vitro* using purified ComR protein and a 5' Bodipy-

labeled self-annealing stem-loop DNA probe that encompassed the ECom-box to which ComR-

XIP binds for transcriptional activation (Mashburn-Warren *et al.*, 2010; Fontaine *et al.*, 2013).

Strong, direct binding of ComR-XIP was observed to the probe in the absence of any of the

XrpA peptides with a calculated K_d of 153 ± 10 nM **(Figure 7A, Table 3)**. However, when XrpA-

1 was added to the reaction, the K_d increased to 651 ± 99 nM. Surprisingly, XrpA-2, which had

no observable effects on the phenotypes examined above, also displayed inhibitory effects,

while inclusion of XrpA-3 or XrpA-4 did not substantially alter the calculated K_d values. A similar

inhibition of ComR-XIP binding was seen when XrpA-N1 or XrpA-N2 peptides were added

individually, with XrpA-N1 having a similar effect to XrpA-1 and XrpA-N2 having a moderate

effect **(Figure 7B)**. Two different scrambled peptides, one with the same aa composition as

XrpA-1 **(Figure 7C)** and another with the same composition as XrpA-2 **(Figure 7D),** were also

tested. Similar to the transcriptional reporter assays, the XrpA-1 scrambled peptide showed

some alleviation of inhibitory properties, albeit not as substantial as XrpA-1, whereas inclusion

of the XrpA-2 scrambled peptide yielded a ComR-XIP DNA binding affinity similar to the control.

While these results suggest that N-terminal fragments of XrpA are sufficient to diminish

the ability of ComR to bind DNA, the effect could be exerted in multiple ways. For example,

XrpA could interact directly with the ComR-XIP oligomer(s) to decrease the affinity of the

complex for DNA, XrpA could compete with XIP for the SHP (XIP) binding site, or XrpA may

inhibit ComR-dependent activation of gene expression by preventing ComR-XIP from forming

higher-order oligomeric complexes. To begin to explore how XrpA influences ComR behavior,

we synthesized a fluorescein isothiocyanate (FITC)-labeled XrpA-N1 peptide and monitored its

binding to ComR by FP in the absence of DNA. Fluorescence polarization in the presence of

XrpA-N1 was increased regardless of whether sXIP was present **(Figure 7E)**, indicating that

XrpA may directly interact with ComR and that this interaction can occur even if XIP does not

occupy the SHP-binding pocket. Further, we tested the specificity of XrpA interaction with ComR

by doing fluorescent polarization experiments with two other *S. mutans* purified proteins that do

not participate directly in the regulation of competence, CcpA and SppA (SMU.508) (Abranches

et al., 2008), and neither showed any ability to interact with fluorescent XrpA-N1 **(Supplemental**

Figure 2). Finally, we conducted a cold competition fluorescence polarization assay and found

poorer binding for our peptide probe to ComR as increasing concentrations of unlabeled XrpA-

N1 peptide were added **(Figure 7F)**. Taken together, these experiments verify that XrpA

interacts directly with ComR, with the outcome being that this interaction antagonizes activation

of gene expression by the ComR-XIP complex and competence development.

DISCUSSION

The study of bacterial cell-cell communication has provided valuable insights into bacterial processes that are critical for growth, essential for the activities of complex microbial ecosystems, and impactful of human health and diseases. These communication pathways have served as tractable model systems to dissect the intricacies of specialized secretory systems for signal molecules and bacteriocins, the mechanisms for signal transduction through two-component systems and cytosolic transcriptional regulators, the hierarchical control of regulons, and how multiple sensory inputs are integrated into key physiological outcomes and manifestation of virulence. As research has continued with these systems, the apparently straightforward paradigms once described for control of quorum sensing and intercellular communication have been assimilated into increasingly complex and diverse models for cellular reprogramming. Examples of such complexity can be found with negative feedback associated with the late competence gene *dprA* of *Streptococcus pneumoniae* (Weng *et al.*, 2013), identification of the novel protein Kre that controls the bimodal regulation of ComK in *Bacillus subtilis* (Gamba *et al.*, 2015), and the recent discovery of a short, leaderless, intercellular peptide signal in Group A *Streptococcus* (Do *et al.*, 2017) that regulates protease expression. Similar advances in understanding of regulatory systems have been realized using *S. mutans* as a model organism (Lemos *et al.*, 2013). One important characteristic that distinguishes *S. mutans* from other streptococci is that it encodes the ComRS signaling systems and integrates the ComCDE bacteriocin production pathway with competence through at least two different signal peptides XIP and CSP, respectively. Here we demonstrate additional complexities in the competence pathway of *S. mutans*, which is intimately intertwined with stress tolerance (Kaspar *et al.*, 2016), by providing experimental evidence that a novel negative feedback system involving the unusual XrpA peptide is a regulator of the ComRS pathway and, consequently, of activation of late competence genes and lytic behaviors.

Using genetic and biochemical approaches, we confirmed that XrpA serves as a

negative regulator of competence development in *S. mutans* by inhibiting activation of the

targets of the ComR-XIP complex, apparently through a direct interaction with ComR that can

be demonstrated *in vivo* by cross-linking and *in vitro* using purified constituents. We first

described XrpA as an antagonist of ComX (Kaspar *et al.*, 2015) that, based on its unusual

genomic location within the *comX* gene and the inverse relationship of *xrpA*-specific mRNA

abundance to the full-length *comX* transcript and ComX protein levels, might function as an anti-

sigma factor, as opposed to acting on early competence genes (ComDE or ComRS systems).

Instead, using transcriptome profiling and transcriptional reporter assays a picture began to

emerge that XrpA influenced ComRS-dependent activation of the *comX* promoter. XrpA

functioning as an autogenous negative regulator of its own expression by blocking ComR-

dependent activation of *comX* may provide the cells with an opportunity to fine-tune ComS and

ComX production, particularly in response to environmental inputs, such as redox. A similar type

of regulation has been described for the *E. coli* RNA polymerase-binding protein DksA, which

along with the co-factor (p)ppGpp promotes a negative feedback loop on the *dksA* promoter to

keep DksA protein levels constant in different environmental conditions (Chandrangsu *et al.*,

2011). Not only is it intriguing in evolutionary terms that the XrpA negative feedback system

evolved within the *comX* coding region, but also there are potentially important physiological

ramifications of the existence of this regulatory circuit. The exact mechanism by which the *xrpA*

mRNA is translated has not been established, but we presently favor a model by which

ribosomal slippage occurs during *comX* translation at or near the *xrpA* start codon, allowing for

production of XrpA, with the efficiency of translational initiation at the *xrpA* start codon and the

stability of the 5' region of the *comX* mRNA being factors that govern the ratio of ComX to XrpA.

Confirmation of such a model is the subject of ongoing research. Nevertheless, if we accept the

premise that *xrpA* translation is not as efficient as that of *comX* when the *comX* promoter is

activated, then it can be envisioned that the negative feedback loop created by XrpA acting on

ComR modulates transcriptional initiation at the *comX* promoter; with the feedback loop

contributing to fine tuning ComX levels in response to cellular physiology and environment

and/or serving as a primary pathway to turn off the competence circuit. The former would be

consistent with the observation that signal perception, induction of *comX* and progression to the

competent state are all exquisitely sensitive to key environmental inputs that include pH (Guo *et*

al., 2014; Son *et al.*, 2015), oxidative stressors (De Furio *et al.*, 2017) and carbohydrate source

and availability (Moye *et al.*, 2016). The latter model would be consistent with the fact that, while

the MecA-Clp pathway can serve as a mechanism to shut off competence through degradation

of ComX, the cells also need a way to shut down the ComRS circuit so as not to produce

nascent ComX during competence while committing to turning off the competence regulon.

Our model for XrpA acting at the level of ComR-dependent activation of *comS* and *comX*

is supported by the transcriptional reporter data in which overexpression of *xrpA* lead to

decreased P*comX* activity (Figure 2). Recently it was suggested that an antagonist termed

"ComZ" must be present within ComRS-positive streptococci that can shut down ComRS

activity in a similar manner to DprA of the ComDE systems (Mirouze *et al.*, 2013; Haustenne *et*

al., 2015). We do not suspect that XrpA is the aforementioned ComZ, as the loss of *xrpA* only

exhibits stronger activation of ComRS-dependent promoters, but does not increase the duration

of activation, as is evident in Figure 2. Studies into ComX stability over the course of

competence activation revealed that ComX protein accumulates faster at early time points after

induction with signal peptides (10-30 minutes) in an *xrpA*-negative strain, consistent with the

data on promoter activity, but ComX protein dissipates at a similar rate in the presence or

absence of *xrpA* (data not shown). Thus, we conclude that XrpA modulates the strength of

ComRS signaling, but other factors must be present, perhaps working in concert with XrpA, to

shut off competence in *S. mutans*, further highlighting the complexity of these systems. Perhaps

other protein candidates identified in our SPINE experiment, such as *ssb2*, fulfill this role; an

area that remains to be investigated.

In *S. mutans*, competence activation has been linked to cell lysis in a process similar to

the fratricide that was first described for *S. pneumoniae.* Fratricide appears to be essential for

efficient gene transfer between bacteria in biofilm communities and to be mediated in *S. mutans*

through the activities of encoded cell wall hydrolases that are a part of the ComX regulon (Wei

and Håvarstein, 2012; Khan *et al.*, 2016), and possibly by intracellular bacteriocins (Perry *et al.*,

2009). As seen in Figure 3, the inhibition of ComRS activity by XrpA has the ability to change

the proportion of cells exhibiting responses to signal inputs in a population, both in terms of

competence activation in complex medium and cell lysis, as measured by propidium iodide

staining of XIP-treated cultures and production of eDNA, which in *S. mutans* is heavily

dependent on cell lysis (Liao *et al.*, 2014). We propose a model **(Figure 8)** in which *xrpA*

influences the decision pathway by which cells choose between a) viability and potential to

uptake DNA as a result of competence activation or b) fratricide and cell death, the latter

providing a source of genetic material and eDNA for incorporation into the extracellular matrix

during biofilm formation (Liao *et al.*, 2014). We posit that this decision network relies on the

strength and duration of P*comX* activation, although we cannot yet rule out the post-

transcriptional regulatory factors (RNA binding proteins, RNAses, or riboswitching) govern the

ratio of XrpA to ComX. Notwithstanding, when XrpA accumulates and/or is active, the strength

of ComRS activation is moderated, providing an even balance between cell viability and cell

death. In the absence of XrpA, ComRS-dependent activation causes ComX over-accumulation,

which favors lytic behavior, which can be observed in Figure 3 and is supported by the fact that

higher concentrations of exogenously supplied XIP induce cell death (Wenderska *et al.*, 2012).

Previously, we reported that overexpression of *xrpA* results in a growth defect in the presence of

oxygen, linking *xrpA* expression and production to oxidative stress tolerance (Kaspar *et al.*,

2015). One aspect not studied here is how *xrpA* might sense an oxidative environment and

integrate that response into competence activation through modulation of ComRS activity. The

C-terminal portion of XrpA, which does not appear to interact with ComR, is very hydrophobic

and may be membrane-associate (Kaspar *et al.*, 2015). Additionally, XrpA is unusual in that it

contains seven cysteine residues distributed fairly evenly across the protein that could

participate in disulfide bond formation, between XrpA proteins/peptides to influence XrpA

availability or to form covalent interactions with binding partners. ComR contains three cysteine

residues distributed evenly over its length, which could allow for covalent coupling to XrpA or

sub-fragments of XrpA. It is our working hypothesis that XrpA integrates the oxidative state of

the environment into fine-tuning the strength of the ComRS signal, tempering activation in one

condition over the other. In the case of early biofilm formation, oxygen is readily available and

could be sensed by environmental inputs such as XrpA, leading to its inactivation. In this

scenario, high ComRS activation and ComX accumulation could shift a larger population of the

cells into a lytic mode during competence activation, releasing eDNA to facilitate the formation

of a protective extracellular matrix, thereby conveying increased fitness or persistence of *S.*

mutans over other health-associated commensal streptococci. As oxygen levels and redox

potential decrease with biofilm maturation, the need for strong competence activation could be

diminished, shifting the cells into a more stable growth mode with a smaller proportion of cells

undergoing lysis. It is noteworthy also that bacteriocins are among the most highly up-regulated

genes when *S. mutans* is growing in air, compared with the transcriptome of anaerobically

growing cells (Ahn *et al.*, 2007). We are currently exploring these ideas and how the

competence regulon is integrated with biofilm development and its role in competition with

commensal streptococci. It is critical to note that ComX and XrpA are highly conserved in all

sequenced clinical isolates of *S. mutans* (Kaspar *et al.*, 2015), suggesting evolutionary pressure

to keep these pathways intact. Our working hypothesis is this evolutionary pressure arises from

the need to compete with commensal streptococci that can antagonize the growth of *S. mutans*

through a variety of mechanisms (Bowen *et al.*, 2017).

We have previously noted that *xrpA* appears to be unique to *S. mutans* (Kaspar *et al.*,

2015). Indeed, a tblastn search for identification of ORFs with similar sequences bacteria

showed that only *Streptococcus troglodytae*, a recently sequenced oral isolate from

chimpanzees that is most closely related to *S. mutans* (Okamoto *et al.*, 2013), contains an intact

xrpA coding sequence embedded within the *comX* coding region. *Streptococcus dysgalactiae*

subsp. *equisimilis* (SDSE) also contains a similar *xrpA* coding sequence, but the *xrpA* protein

coding sequence is disrupted by premature stop codons in the sequenced isolate of this

streptococcus. Thus, it appears that only *S. mutans* and extremely closely related organisms

are the only ComRS-containing streptococci that encode an XrpA-like inhibitor; there is no

evidence that XrpA is present in *S. rattus*, *S. sobrinus. S. cricetus*, *S. downeii* or other mutans

streptococci. We cannot, however, exclude that there are proteins or peptides in ComRS-

containing streptococci that play a role that is similar or identical to that of XrpA in *S. mutans.*

The unique nature of XrpA in *S. mutans* is also notable in the context that the *S. mutans* ComR

has strict recognition for its cognate XIP peptide (Shanker *et al.*, 2016), whereas ComR proteins

from Bovis and Pyogenic streptococci are more promiscuous in the XIP peptides that they are

able to recognize to enhance ComR DNA binding capacity. It is then a logical conclusion that

the Mutans group has further separated from the Bovis and Pyogenic group in terms of ComR-

XIP regulation. A logical extrapolation of these observations is to ask whether XrpA could serve

as a novel anti-caries target or therapeutic. It is important to note that other oral health-

associated commensal streptococci, such as *Streptococcus mitis*, *Streptococcus gordonii* and

Streptococcus sanguinis are all part of the Mitis group of streptococci that lack ComRS signaling

systems and rely on ComDE for competence activation (Håvarstein, 2010), so targeting XrpA

should not disrupt a healthy oral biofilm. It is encouraging that small synthetized portions of

XrpA have an effect on competence activation, as shown in Figure 6. It is critical to point out

that while theses synthetized peptides were provided exogenously in this study, we do not

suspect at this time that XrpA is actively released to the extracellular environment. As previously

discussed, our current working hypothesis is that XrpA is able to accumulate in response to

external environmental cues, such as oxidative stress, and function as a sensor inside the cells,

and at this time have no reason to believe that XrpA has an extracellular lifecycle. Our working

model must also account for the fact that XrpA fragments can elicit effects when provided

exogenously. Therefore, we are testing the hypothesis that XrpA peptides, which may be

released into the extracellular space through lysis, can be actively internalized, perhaps after

processing, to elicit their effects on ComR – similar to what has been proposed for XIP (Kaspar

et al., 2017). Mass spectrometry studies are currently ongoing to localize XrpA and other *S.*

mutans encoded peptides that affect competence (Ahn *et al.*, 2014).

In summary, this work provide additional novel insights into the complex regulatory

nature of bacterial cell-cell signaling systems, providing the organisms with multiple check

points throughout the circuit to either amplify or diminish the response to signal inputs based on

key environmental and/or physiologic cues. Future work will be focused on how environmental

inputs can influence XrpA/ComR interaction and activities, and the resulting consequences in

terms of biofilm ecology. Development of these model systems should shed further light on

microbial interactions and the importance of cell-cell signaling systems at the very earliest

stages of colonization and biofilm development. It is also interesting to ponder if the recently

discovered peptides such as XrpA, the *rcrQ*-associated peptides of *S. mutans* (Ahn *et al.*, 2014)

and the leaderless SHP of GAS (Do *et al.*, 2017) play important roles in the virulence potential

of these organisms if multiple other peptides with profound impacts on cellular behaviors are

currently hidden in the genomes of Gram-positive bacteria.

EXPERIMENTAL PROCEDURES

Bacterial Strains and Growth Conditions.

S. mutans wild-type strain UA159 and its derivatives **(Table 1)** were grown in either brain heart

infusion (BHI - Difco), FMC (Terleckyj *et al.*, 1975; Terleckyj and Shockman, 1975) or CDM

(Chang *et al.*, 2011) medium. The medium was supplemented with 10 µg ml^{-1} erythromycin, 1

mg ml^{-1} of kanamycin or 1 mg ml^{-1} spectinomycin when needed. Unless otherwise noted,

cultures were grown overnight in BHI medium with the appropriate antibiotics at 37°C in a 5%

CO_2 aerobic atmosphere. The next day, cultures were harvested by centrifugation, washed

twice in 1 mL of phosphate-buffered saline (PBS), and resuspended in PBS to remove all traces

of BHI. Cells were then diluted in the desired medium before beginning each experiment.

Synthetic XIP (sXIP, aa sequence = GLDWWSL), corresponding to residues 11-17 of ComS,

was synthesized and purified to 96% homogeneity by NeoBioSci (Cambridge, MA). The

lyophilized sXIP was reconstituted with 99.7% dimethyl sulfoxide (DMSO) to a final

concentration of 2 mM and stored in 100 µL aliquots at -20°C. Selected XrpA peptide

sequences and fluorescently-labeled derivatives were synthesized, purified and confirmed by

mass spectrometry by Biomatik USA (Wilmington, DE). XrpA peptides were also reconstituted

with DMSO to a final concentration of 1 mM and stored at -20°C.

Construction of Bacterial Strains.

Mutant strains of *S. mutans*, including inactivation of the *xrpA* start codon (Δ*xrpA*) were created

using a PCR ligation mutagenesis approach as previously described (Lau *et al.*, 2002; Kaspar *et*

al., 2015). Overexpression of genes (*xrpA*, *comR*) was achieved by amplifying the structural

genes of interest from *S. mutans* UA159 and cloning into the expression plasmid pIB184 using

the EcoRI and BamHI restriction sites (Biswas *et al.*, 2008). For *in vivo* protein-protein

interaction experiments, a Strep-tag sequencing (WSHPQFEK) was first inserted in front of the

stop codon on the pIB184-ComR overexpressing plasmid using the Q5® Site Directed

Mutagenesis Kit (New England Biolabs, Beverly, Mass.) and following the provided protocol.

After selection of the appropriate construct by sequencing, a [G4S]$_2$ Linker sequence

(ggtggaggaggctctggtggaggcggtagc) was then inserted between the *comR* and Strep-tag

sequence using the same kit and protocol. Transformants were confirmed by PCR and

sequencing after selection on BHI agar with appropriate antibiotics.   Plasmid DNA was isolated

from *E. coli* using QIAGEN (Chatsworth, Calif.) miniprep columns, and restriction and DNA-

modifying enzymes were obtained from New England Biolabs. PCRs were carried out with 100

ng of chromosomal DNA by using *Taq* DNA polymerase, and PCR products were purified with

the QIAquick kit (QIAGEN).

Transcriptome Profiling via RNA-Seq.

Selected strains of *S. mutans* to be analyzed by RNA-sequencing (UA159, Δ*xrpA*) were grown

in FMC medium to mid-exponential log phase of OD$_{600}$ = 0.5 before harvesting. For *S. mutans*

UA159 treated with 2 μM sXIP, sXIP was added at OD$_{600}$ = 0.2. RNA extraction, rRNA removal,

library construction and read analysis was conducted as previously described elsewhere (Zeng

et al., 2013; Kaspar *et al.*, 2015). Briefly, 10 μg of high-quality total RNA was processed using

the MICROB*Express*^{TM} Bacterial mRNA Enrichment Kit (Ambion of Life Technologies, Grand

Island, NY), twice, before ethanol precipitation and resuspension in 25 μL of nuclease-free

water. The quality of enriched mRNA samples was analyzed using an Agilent Bioanalyzer

(Agilent Technologies, Santa Clara, CA). cDNA libraries were generated from the enriched

mRNA samples using the TruSeq Illumina kit (Illumina, San Diego, CA), following instructions

from the supplier. Deep sequencing was performed at the University of Florida ICBR facilities

(Gainesville, FL). Approximately 20 million short-reads were obtained for each sample. After

removing adapter sequences from each short-read and trimming of the 3'-ends by quality

scores (Schmieder and Edwards, 2011), the resulting sequences were mapped onto the

reference genome of strain UA159 (GenBank accession no. AE014133) using the short-read

aligner. Mapped short-read alignments were then converted into readable formats using

SAMTOOLS (Li *et al.*, 2009). For viewing of the mapped reads aligned to the genome, .bam

files were uploaded into the Integrative Genomics Viewer (IGV – version 2.3.55) (Robinson *et*

al., 2011). A ".csv" file containing raw read counts for each replicate (3) was then uploaded to

Degust (http://degust.erc.monash.edu/) and edgeR analysis performed to determine Log2 fold

change and a false discovery rate (FDR). The P-value was obtained by taking the –log10 of the

FDR. The data files used in this study are available from NCBI-GEO (Gene Expression

Omnibus) under accession no. GSE110167.

Measurements of Promoter Activity via GFP Fluorescence.

For measurements of GFP fluorescence, cultures were inoculated from washed overnight

cultures in CDM medium at a 1:50 dilution. Inoculated medium (175 µL) was added to each well

along with a 50 µL mineral oil overlay in a Costar™ 96 well assay plate (black plate with clear

bottom; Corning Incorporated) and incubated at 37°C. At intervals of 30 minutes for a total of 18

hours, OD_{600} along with GFP fluorescence (excitation 485/20 nm, emission 528/20 nm) was

measured with a Synergy 2 multimode microplate reader (BioTek). Relative expression was

calculated by subtracting the background fluorescence of UA159 (mean from six replicates)

from raw fluorescence units of the reporter strains and then dividing by $OD_{600.}$

Flow cytometry.

Bacterial cultures were grown to $OD_{600} = 0.6$ before being harvested, washed and resuspended

in PBS before being run through a FACSCalibur™ (BD Biosciences) flow cytometer. For sCSP

experiments, cultures were grown in BHI after a 1:20 dilution from overnight culture while

cultures were grown in FMC for sXIP experiments at the same initial dilution. Both peptides

were added to the growing cultures at $OD_{600} = 0.2$. sXIP treated cells were stained with 5 µg

mL^{-1} propidium iodide (PI) for 10 minutes in the dark for analysis of membrane-compromised

cells. Cells were then sonicated in a water bath sonicator for 3 intervals of 30 seconds in 5 mL

polystyrene round-bottom tubes to achieve primarily single cells for analysis. Forward and side

scatter signals were set stringently to allow sorting of single cells. In total, 5×10^4 cells were

counted from each event, at a maximum rate of 2×10^3 cells per second, and each experiment

was performed in triplicate. Detection of GFP fluorescence was through a 530 nm (± 30 nm)

bandpass filter, and PI was detected using a 670-nm long pass filter. Data were acquired for

unstained cells and single-color positive controls so that data collection parameters could be

properly set. The data were collected using Cell Quest Pro (BD Biosciences) and analyzed with

FCS Express 4 (De Novo Software). Gating for quadrant analysis was selected by using a dot

density plot with forward and side scatter, with gates set to capture the densest section of the

plot. Graphing and statistical analyses were performed using Prism (GraphPad Software). *x*-

and *y*-axis data represent logarithmic scales of fluorescent intensity (arbitrary units).

Measurements of eDNA Release.

Overnight cultures of selected *S. mutans* strains, grown in CDM medium with addition of a final

concentration of 10 µM of synthetic XrpA peptides when noted, where measured for a final

OD_{600} and then harvested by centrifugation for their supernatant fraction. 5 mL of the resulting

supernatant was then run through QIAGEN (Chatsworth, Calif.) PCR purification columns to

capture eDNA present. The eDNA was then eluted off the column with 600 µL water, and 594

µL of this elution was mixed with 5 µL of 50 µM Sytox Green (Invitrogen) to a final concentration

of 0.5 µM. After vortexing the solution and incubation for 15 minutes in the dark at room

temperature, 200 µL of the stained samples were transferred into a Costar™ 96 well assay plate

(black plate with clear bottom; Corning Incorporated). Fluorescence (excitation 485/20 nm,

emission 528/20 nm) was measured with a Synergy 2 multimode microplate reader (BioTek)

and the resulting data was then normalized for the measured final OD_{600} nm resulting in a final

arbitrary eDNA release measurement. The data represents 3 independent biological replicates

with 3 technical replicates each. Statistical significance was determined by the Student's T-Test.

Biofilm Assays.

Selected *S. mutans* strains were grown from overnight cultures to mid-exponential phase after a

1:20 dilution in BHI broth at 37ºC in a 5% CO_2 atmosphere. The mid-exponential phase cells

were then diluted 1:100 into CDM medium with either 20 mM glucose or CDM medium

containing 15 mM glucose and 2.5 mM sucrose as a carbohydrate source. 200 µL of this dilution

was loaded into 96 well polystyrene microtiter plates and incubated in a 5% CO_2 atmosphere at

37ºC for 48 h. A final concentration of 10 µM of synthetic XrpA peptides were added when

needed. After, the medium was decanted, and the plates were washed twice with 200 µL of

sterile water to remove planktonic and loosely bound cells. The adherent bacteria were stained

with 60 µL of 0.1% crystal violet for 15 min. After rinsing twice with 200 µL of water, the bound

dye was extracted from the stained biofilm using 200 µL of ethanol:acetone (8:2) solution, twice.

The extracted dye was diluted into 1.6 mL of ethanol:acetone solution. Biofilm formation was

quantified by measuring the absorbance of the solution at OD $_{575}$ nm. The data represents 3

independent biological replicates with 4 technical replicates each. Statistical significance was

determined by the Student's T-Test.

SPINE for ComR Interactions.

A Strep-tag protein interaction experiment (SPINE) was derived from a previously published

protocol (Herzberg *et al.*, 2007). Briefly, a strain harboring a C-terminal Strep-tagged ComR,

along with the vector only control, was grown in 500 mL of CDM medium after a 1:20 dilution

from overnight cultures to an $OD_{600} = 0.6$. At this time, either 2 µM of sXIP or DMSO (vehicle,

0.1% final concentration) was added to cultures and were grown an additional hour at 37ºC in a

5% CO_2 atmosphere. After, cells were pelleted by centrifugation, washed and resuspened in 50

mM HEPES (pH 8) with a final concentration of 2.5 mM of protein crosslinker solution added

(DSP, Thermo Scientific). The cells were incubated at 37ºC for 45 minutes, at which point 50

mM Tris (pH 7.5) was added to stop the crosslinking reaction. After, cells were pelleted, washed

in Buffer W, and lysed via bead beating in Buffer W. The crosslinked Strep-tagged ComR

complex was then purified from the lysate using Strep-tactin® resin (iba) in a chromatography

column following the manufacture's protocol. Purification was verified by running the selected

fractions and elutions on 16.5% Tris-Tricine gels (BioRad) followed by silver staining and/or

western blot **(Supplemental Figure 3)**. Protein concentrations of the elutions were determined

using the bicinchoninic acid assay (BCA; Thermo Scientific). Complex-containing elutions were

then combined and precipitated by the TCA/Acetone precipitation method. Precipitant was sent

to Applied Biomics (Hayward, CA) for 2D DIGE Protein Expression Profiling which included 2D

gel electrophoresis, determination of protein expression ratios between samples, spot picking

and identification by LC-MS/MS.

Transformation Assays.

Overnight cultures were diluted 1:20 in 200 µL of FMC medium in polystyrene microtiter plates

in the presence or absence of 10 µM of synthetic XrpA peptides. The cells were grown to OD_{600}

= 0.15 in a 5% CO_2 atmosphere. When desired, 0.5 µM of sXIP was added, cells were

incubated for 10 min and 0.5 µg of purified plasmid pIB184, which harbors a erythromycin

resistance (Erm^R) gene, was added to the culture. After 2.5 h incubation at 37°C, transformants

and total CFU were enumerated by plating appropriate dilutions on BHI agar plates with and

without the addition of 1 mg mL^{-1} erythromycin, respectively. CFU were counted after 48 h of

incubation, and transformation efficiency was expressed as the percentage of transformants

among the total viable cells. Fold change was then calculated from the UA159 control with

DMSO (vehicle) addition. Statistical significance was determined by the Student's T-Test

Western blot.

Overnight cultures of *S. mutans* were diluted 1:50 into 35 mL of FMC medium and harvested by

centrifugation when the cultures reached an $OD_{600} = 0.5$. When desired, 2 µM sXIP was added

when the cultures reached an OD_{600} value of 0.2 along with 10 µM of selected XrpA peptides.

Cell pellets collected by centrifugation were washed once with buffer A (0.5 M sucrose; 10 mM

Tris-HCl, pH 6.8; 10 mM $MgSO_4$) containing 10 µg mL^{-1} of phenylmethanesulfoynl fluoride

(PMSF) (ICN Biomedicals Inc.) and resuspended in 0.5 mL Tris-buffered saline (50 mM Tris-

HCl, pH 7.5; 150 mM NaCl). Cells were lysed using a Mini Bead Beater (Biospec Products) in

the presence of 1 volume of glass beads (avg. diam. 0.1 mm) for 30 s intervals, three times,

with incubation on ice between homogenizations. Lysates were then centrifuged at 3,000 × *g* for

10 minutes at 4°C. Protein concentrations of the resulting supernates were determined using

the bicinchoninic acid assay (BCA; Thermo Scientific) with purified bovine serum albumin as the

standard. Ten microgram aliquots of proteins were mixed with 5X SDS sample buffer (200 mM

Tris-HCl, pH 6.8; 10% [v/v] SDS; 20% [v/v]; 10% [v/v] β-merchaptoethanol; 0.02% [v/v]

bromophenol blue), loaded on a 12.5% polyacrylamide gel with a 5% stacking gel and

separated by electrophoresis at 150 V for 45 minutes. Proteins were transferred to Immobilon-P

polyvinylidene difluroride (PVDF) membranes (Millipore) using a Trans-Blot Turbo transfer

system and a protocol provided by the supplier (BioRad). The membranes were treated with

either primary polyclonal anti-ComX, anti-ComR or anti-ManL (loading control) antisera at a

1:1000 dilution and a secondary peroxidase-labeled, goat anti-rabbit IgG antibody (1:5000

dilution; Kierkegaard & Perry Laboratories, USA). Detection was performed using a SuperSignal

West Pico Chemiluminescent Substrate kit (Thermo Scientific) and visualized with a FluorChem

8900 imaging system (Alpha Innotech, USA).

Purification of Recombinant ComR.

The *S. mutans* UA159 *comR* gene was amplified using primers AAAGAATCCTATGTTAAAAGA

and CACCCTAGGAGACCCATCAAA and was cloned into the BamHI and AvrII sites of the pET

45b expression vector downstream of the 6x His-tag and separated by an enterokinase

cleavage site. The resulting vector was transformed into *E. coli* DE3 cells (New England

Biolabs). To induce expression of *comR*, 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) was

added to 1 L of growing culture in LB medium once the OD_{600} reached 0.6. The culture was

grown for an additional 4 h at 37°C before the cells were pelleted by centrifugation and frozen

overnight at -20°C. The next day, the cells were lysed after suspension into B-PER (Thermo

Scientific) with addition of HALT protease inhibitor (Thermo Scientific) and cellular debris

removed by centrifugation for 20 minutes at 13,000 x g. The His-ComR was column purified

using Ni-NTA resin (QIAGEN) and eluted with 250 mM imidazole **(Supplemental Figure 4)**. To

remove the 6x His-tag, 1 mL of a 2-3 mg mL^{-1} purified His-ComR sample was dialyzed in

EKMax Buffer overnight (50 mM Tris-HCL pH 8, 1 mM $CaCl_2$, 0.1% Tween-20) with 50 U of

EKMax enterokinase (Thermo Scientific) then added and incubated overnight at 4°C. Finally,

the cleaved ComR sample was added to a dialysis cassette to exchange the EKMax buffer with

PBS pH 7.4 overnight. Final protein concentration was determined using the bicinchoninic acid

assay (BCA; Thermo Scientific) with purified bovine serum albumin as the standard. Digestion

was confirmed by SDS-PAGE and Coomassie Blue staining **(Supplemental Figure 5)**.

Fluorescence Polarization

A 5' Bodipy-labeled self-annealing, stem-loop DNA probe with sequence encompassing the

ECom-box which ComR binds to within the P*comX* promoter (5'-BODIPY FL-X -

ATGGGACATTTATGTCCTGTCCCCCACAGGACATAAATGTCCCAT - 3'), was synthesized

(Thermo Fisher) and kept at a constant concentration of 1 μM in all reactions. Purified ComR

protein was serially diluted, ranging from 5 to 2000 nM, and mixed with 10 μM sXIP and 10 μM

of selected synthetic XrpA peptides unless otherwise noted, in reaction buffer to a final volume

of 250 μL (PBS pH 7.4, 10 mM βME, 1 mM EDTA, 0.1 mg mL^{-1} BSA, 20% glycerol (v/v), 0.01%

Triton X-100 and 0.05 mg mL^{-1} salmon sperm DNA). Reactions were transferred to a Corning®

96-well, half-area, black polystyrene plate prior to incubation at 37°C for 30 minutes.

Polarization values were measured using a BioTek Synergy 2 plate reader (excitation 485 nm,

emission 528 nm), and the resulting millipolarization values were plotted for each protein

concentration tested to assess protein/peptide interactions. For fluorescent peptides

experiments, a synthetic fluorescein isothiocyanate (FITC)-labeled XrpA-N1 peptide (FITC-AHX-

MIQNCISILRHFLITLK) was used (Biomatik USA) with reaction buffer PBS pH 7.4, 10 mM βME,

0.1 mg mL^{-1} BSA, 20% glycerol (v/v), and 0.01% Triton X-100. Graphing and linear regression

analyses to determine kD values were performed using Prism (GraphPad Software).

ACKNOWLEDGEMENTS

Research reported in this publication was supported by the National Institute of Dental and

Craniofacial Research of the National Institutes of Health under Award Numbers R01 DE13239,

R01 DE023339, and T90 DE21990.

AUTHOR CONTRIBUTIONS

JK, RCS and RAB contributed to the conception and design of the study; JK and RCS

performed the experiments, acquired and analyzed the data, JK and RAB interpreted the data;

and JK and RAB wrote the manuscript.

REFERENCES

Abranches, J., Nascimento, M.M., Zeng, L., Browngardt, C.M., Wen, Z.T., Rivera, M.F., and

Burne, R.A. (2008) CcpA regulates central metabolism and virulence gene expression in

Streptococcus mutans. *J Bacteriol* **190**: 2340–9

Ahn, S.-J., Kaspar, J., Kim, J.N., Seaton, K., and Burne, R. a (2014) Discovery of novel peptides

regulating competence development in *Streptococcus mutans*. *J Bacteriol* **196**: 3735–3745.

Ahn, S.-J., and Rice, K.C. (2016) Understanding the *Streptococcus mutans* Cid/Lrg System

through CidB Function. *Appl Environ Microbiol* **82**: 6189–6203.

Ahn, S.-J., Wen, Z.T., and Burne, R.A. (2007) Effects of oxygen on virulence traits of

Streptococcus mutans. *J Bacteriol* **189**: 8519–27.

Biswas, I., Jha, J.K., and Fromm, N. (2008) Shuttle expression plasmids for genetic studies in

Streptococcus mutans. *Microbiology* **154**: 2275–82.

Boutry, C., Delplace, B., Clippe, A., Fontaine, L., and Hols, P. (2013) SOS response activation

and competence development are antagonistic mechanisms in *Streptococcus thermophilus*. *J

Bacteriol* **195**: 696–707.

Bowen, W.H., Burne, R.A., Wu, H., and Koo, H. (2017) Oral Biofilms: Pathogens, Matrix, and

Polymicrobial Interactions in Microenvironments. *Trends Microbiol*

https://doi.org/10.1016/j.tim.2017.09.008.

Chandrangsu, P., Lemke, J.J., and Gourse, R.L. (2011) The *dksA* promoter is negatively

feedback regulated by DksA and ppGpp. *Mol Microbiol* **80**: 1337–48.

Chang, J.C., and Federle, M.J. (2016) PptAB Exports Rgg Quorum-Sensing Peptides in

Streptococcus. PLoS One **11**: e0168461.

Chang, J.C., LaSarre, B., Jimenez, J.C., Aggarwal, C., and Federle, M.J. (2011) Two Group A

Streptococcal Peptide Pheromones Act through Opposing Rgg Regulators to Control Biofilm

Development. *PLoS Pathog* **7**: e1002190.

Chattoraj, P., Banerjee, A., Biswas, S., and Biswas, I. (2010) ClpP of *Streptococcus mutans*

differentially regulates expression of genomic islands, mutacin production, and antibiotic

tolerance. *J Bacteriol* **192**: 1312–23.

Desai, K., Mashburn-Warren, L., Federle, M.J., and Morrison, D.A. (2012) Development of

competence for genetic transformation of *Streptococcus mutans* in a chemically defined

medium. *J Bacteriol* **194**: 3774–80.

Do, H., Makthal, N., VanderWal, A.R., Rettel, M., Savitski, M.M., Peschek, N., *et al.* (2017)

Leaderless secreted peptide signaling molecule alters global gene expression and increases

virulence of a human bacterial pathogen. *Proc Natl Acad Sci U S A* **114**: E8498–E8507.

Fontaine, L., Boutry, C., Frahan, M.H. de, Delplace, B., Fremaux, C., Horvath, P., *et al.* (2009) A

Novel Pheromone Quorum-Sensing System Controls the Development of Natural Competence

in *Streptococcus thermophilus* and *Streptococcus salivarius. J Bacteriol* **192**: 1444–1454.

Fontaine, L., Goffin, P., Dubout, H., Delplace, B., Baulard, A., Lecat-Guillet, N., *et al.* (2013)

Mechanism of competence activation by the ComRS signalling system in streptococci. *Mol*

Microbiol **87**: 1113–32.

Fuqua, W.C., Winans, S.C., and Greenberg, E.P. (1994) Quorum sensing in bacteria: the LuxR-

LuxI family of cell density-responsive transcriptional regulators. *J Bacteriol* **176**: 269–75.

Furio, M. De, Ahn, S.J., Burne, R.A., and Hagen, S.J. (2017) Oxidative Stressors Modify the

Response of *Streptococcus mutans* to Its Competence Signal Peptides. *Appl Environ Microbiol*

83: e01345-17.

Gamba, P., Jonker, M.J., and Hamoen, L.W. (2015) A Novel Feedback Loop That Controls

Bimodal Expression of Genetic Competence. *PLoS Genet* **11**: e1005047.

Gardan, R., Besset, C., Gitton, C., Guillot, A., Fontaine, L., Hols, P., and Monnet, V. (2013)

Extracellular life cycle of ComS, the competence-stimulating peptide of *Streptococcus*

thermophilus. J Bacteriol **195**: 1845–55.

Greenberg, E.P. (2003) Bacterial communication and group behavior. *J Clin Invest* **112**: 1288–

90.

Guo, Q., Ahn, S.-J., Kaspar, J., Zhou, X., and Burne, R.A. (2014) Growth phase and pH

influence peptide signaling for competence development in *Streptococcus mutans. J Bacteriol*

196: 227–36.

Hagen, S.J., and Son, M. (2017) Origins of heterogeneity in *Streptococcus mutans* competence:

interpreting an environment-sensitive signaling pathway. *Phys Biol* **14**: 15001.

Haustenne, L., Bastin, G., Hols, P., and Fontaine, L. (2015) Modeling of the ComRS Signaling

Pathway Reveals the Limiting Factors Controlling Competence in *Streptococcus thermophilus*.

Front Microbiol **6**.

Håvarstein, L.S. (2010) Increasing competence in the genus *Streptococcus. Mol Microbiol* **78**:

541–4.

Håvarstein, L.S., Coomaraswamy, G., and Morrison, D.A. (1995) An unmodified

heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus*

pneumoniae. Proc Natl Acad Sci U S A **92**: 11140–4.

Herzberg, C., Weidinger, L.A.F., Dörrbecker, B., Hübner, S., Stülke, J., and Commichau, F.M.

(2007) SPINE: A method for the rapid detection and analysis of protein–protein interactionsin

vivo. *Proteomics* **7**: 4032–4035.

Hui, F.M., Zhou, L., and Morrison, D.A. (1995) Competence for genetic transformation in

Streptococcus pneumoniae: organization of a regulatory locus with homology to two lactococcin

A secretion genes. *Gene* **153**: 25–31.

Hung, D.C.I., Downey, J.S., Ayala, E.A., Kreth, J., Mair, R., Senadheera, D.B., *et al.* (2011)

Characterization of DNA binding sites of the ComE response regulator from *Streptococcus*

mutans. J Bacteriol **193**: 3642–52.

Johnston, C., Martin, B., Fichant, G., Polard, P., and Claverys, J.-P. (2014) Bacterial

transformation: distribution, shared mechanisms and divergent control. *Nat Rev Microbiol* **12**:

181–96.

Kaspar, J., Ahn, S.-J., Palmer, S.R., Choi, S.C., Stanhope, M.J., and Burne, R.A. (2015) A

Unique ORF within the *comX* gene of *Streptococcus mutans* Regulates Genetic Competence

and Oxidative Stress Tolerance. *Mol Microbiol* **96**: 463–482.

Kaspar, J., Kim, J.N., Ahn, S.-J., and Burne, R.A. (2016) An Essential Role for (p)ppGpp in the

Integration of Stress Tolerance, Peptide Signaling, and Competence Development in

Streptococcus mutans. Front Microbiol **7**: 1162.

Kaspar, J., Underhill, S.A.M., Shields, R.C., Reyes, A., Rosenzweig, S., Hagen, S.J., and

Burne, R.A. (2017) Intercellular communication via the *comX*-Inducing Peptide (XIP) of

Streptococcus mutans. J Bacteriol JB.00404-17.

Khan, R., Rukke, H. V., Høvik, H., Åmdal, H.A., Chen, T., Morrison, D.A., and Petersen, F.C.

(2016) Comprehensive Transcriptome Profiles of *Streptococcus mutans* UA159 Map Core

Streptococcal Competence Genes. *mSystems* **1**: e00038-15.

Khan, R., Rukke, H. V, Ricomini Filho, A.P., Fimland, G., Arntzen, M.Ø., Thiede, B., and

Petersen, F.C. (2012) Extracellular identification of a processed type II ComR/ComS

pheromone of *Streptococcus mutans. J Bacteriol* **194**: 3781–8.

Kreth, J., Hung, D.C.I., Merritt, J., Perry, J., Zhu, L., Goodman, S.D., *et al.* (2007) The response

regulator ComE in *Streptococcus mutans* functions both as a transcription activator of mutacin

production and repressor of CSP biosynthesis. *Microbiology* **153**: 1799–807.

Lau, P.C.., Sung, C.K., Lee, J.H., Morrison, D.A., and Cvitkovitch, D.G. (2002) PCR ligation

mutagenesis in transformable streptococci: application and efficiency. *J Microbiol Methods* **49**:

193–205.

LeBlanc, D.J., Lee, L.N., and Abu-Al-Jaibat, A. (1992) Molecular, genetic, and functional

analysis of the basic replicon of pVA380-1, a plasmid of oral streptococcal origin. *Plasmid* **28**:

130–145.

Lemos, J.A., Quivey, R.G., Koo, H., and Abranches, J. (2013) *Streptococcus mutans*: a new

Gram-positive paradigm? *Microbiology* **159**: 436–45.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., *et al.* (2009) The Sequence

Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–9.

Li, Y.H., Lau, P.C., Lee, J.H., Ellen, R.P., and Cvitkovitch, D.G. (2001) Natural genetic

transformation of *Streptococcus mutans* growing in biofilms. *J Bacteriol* **183**: 897–908.

Liao, S., Klein, M.I., Heim, K.P., Fan, Y., Bitoun, J.P., Ahn, S.-J., *et al.* (2014) *Streptococcus*

mutans extracellular DNA is upregulated during growth in biofilms, actively released via

membrane vesicles, and influenced by components of the protein secretion machinery. *J*

Bacteriol **196**: 2355–66.

Martin, B., Soulet, A.-L., Mirouze, N., Prudhomme, M., Mortier-Barrière, I., Granadel, C., *et al.*

(2013) ComE/ComE~P interplay dictates activation or extinction status of pneumococcal X-state

(competence). *Mol Microbiol* **87**: 394–411.

Mashburn-Warren, L., Morrison, D.A., and Federle, M.J. (2010) A novel double-tryptophan

peptide pheromone controls competence in *Streptococcus* spp. via an Rgg regulator. *Mol*

Microbiol **78**: 589–606.

Mashburn-Warren, L., Morrison, D.A., and Federle, M.J. (2012) The cryptic competence

pathway in *Streptococcus pyogenes* is controlled by a peptide pheromone. *J Bacteriol* **194**:

4589–600.

Mirouze, N., Bergé, M.A., Soulet, A.-L., Mortier-Barrière, I., Quentin, Y., Fichant, G., *et al.*

(2013) Direct involvement of DprA, the transformation-dedicated RecA loader, in the shut-off of

pneumococcal competence. *Proc Natl Acad Sci U S A* **110**: E1035-44.

Moye, Z.D., Son, M., Rosa-Alberty, A.E., Zeng, L., Ahn, S.-J., Hagen, S.J., and Burne, R.A.

(2016) Effects of Carbohydrate Source on Genetic Competence in *Streptococcus mutans*. *Appl*

Environ Microbiol AEM.01205-16.

Okamoto, M., Imai, S., Miyanohara, M., Saito, W., Momoi, Y., Abo, T., *et al.* (2013)

Streptococcus troglodytae sp. nov., from the chimpanzee oral cavity. *Int J Syst Evol Microbiol*

63: 418–422.

Papenfort, K., and Bassler, B.L. (2016) Quorum sensing signal–response systems in Gram-

negative bacteria. *Nat Rev Microbiol* **14**: 576–588.

Perry, J.A., Jones, M.B., Peterson, S.N., Cvitkovitch, D.G., and Lévesque, C.M. (2009) Peptide

alarmone signalling triggers an auto-active bacteriocin necessary for genetic competence. *Mol*

Microbiol **72**: 905–17.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and

Mesirov, J.P. (2011) Integrative genomics viewer. *Nat Biotechnol* **29**: 24–6.

Schmieder, R., and Edwards, R. (2011) Quality control and preprocessing of metagenomic

datasets. *Bioinformatics* **27**: 863–4.

Shanker, E., Morrison, D.A., Talagas, A., Nessler, S., Federle, M.J., and Prehna, G. (2016)

Pheromone Recognition and Selectivity by ComR Proteins among *Streptococcus* Species.

PLOS Pathog **12**: e1005979.

Shields, R.C., and Burne, R.A. (2016) Growth of *Streptococcus mutans* in Biofilms Alters

Peptide Signaling at the Sub-population Level. *Front Microbiol* **7**: 1075.

Son, M., Ahn, S.-J., Guo, Q., Burne, R.A., and Hagen, S.J. (2012) Microfluidic study of

competence regulation in *Streptococcus mutans*: environmental inputs modulate bimodal and

unimodal expression of *comX*. *Mol Microbiol* **86**: 258–72.

Son, M., Ghoreishi, D., Ahn, S.-J., Burne, R.A., and Hagen, S.J. (2015) Sharply Tuned pH

Response of Genetic Competence Regulation in *Streptococcus mutans*: a Microfluidic Study of

the Environmental Sensitivity of *comX*. *Appl Environ Microbiol* **81**: 5622–31.

Straume, D., Stamsås, G.A., and Håvarstein, L.S. (2015) Natural transformation and genome

evolution in *Streptococcus pneumoniae. Infect Genet Evol* **33**: 371–80.

Talagas, A., Fontaine, L., Ledesma-Garca, L., Mignolet, J., Li de la Sierra-Gallay, I., Lazar, N.,

et al. (2016) Structural Insights into Streptococcal Competence Regulation by the Cell-to-Cell

Communication System ComRS. *PLOS Pathog* **12**: e1005980.

Terleckyj, B., and Shockman, G.D. (1975) Amino acid requirements of *Streptococcus mutans*

and other oral streptococci. *Infect Immun* **11**: 656–664.

Terleckyj, B., Willett, N.P., and Shockman, G.D. (1975) Growth of several cariogenic strains of

oral streptococci in a chemically defined medium. *Infect Immun* **11**: 649–55.

Tomasz, A. (1965) Control of the Competent State in Pneumococcus by a Hormone-Like Cell

Product: An Example for a New Type of Regulatory Mechanism in Bacteria. *Nature* **208**: 155–

159.

Waterhouse, J.C., Swan, D.C., and Russell, R.R.B. (2007) Comparative genome hybridization

of *Streptococcus mutans* strains. *Oral Microbiol Immunol* **22**: 103–110.

Wei, H., and Håvarstein, L.S. (2012) Fratricide is essential for efficient gene transfer between

pneumococci in biofilms. *Appl Environ Microbiol* **78**: 5897–905.

Wenderska, I.B., Lukenda, N., Cordova, M., Magarvey, N., Cvitkovitch, D.G., and Senadheera,

D.B. (2012) A novel function for the competence inducing peptide, XIP, as a cell death effector

of *Streptococcus mutans*. *FEMS Microbiol Lett* **336**: 104–12.

Weng, L., Piotrowski, A., and Morrison, D.A. (2013) Exit from Competence for Genetic

Transformation in *Streptococcus pneumoniae* Is Regulated at Multiple Levels. *PLoS One* **8**:

e64197.

Whiteley, M., Diggle, S.P., and Greenberg, E.P. (2017) Progress in and promise of bacterial

quorum sensing research. *Nature* **551**: 313–320.

Wilkening, R. V., Chang, J.C., and Federle, M.J. (2016) PepO, a CovRS-controlled

endopeptidase, disrupts *Streptococcus pyogenes* quorum sensing. *Mol Microbiol* **99:** 71-87.

Zeng, L., Choi, S.C., Danko, C.G., Siepel, A., Stanhope, M.J., and Burne, R.A. (2013) Gene

regulation by CcpA and catabolite repression explored by RNA-Seq in *Streptococcus mutans*.

PLoS One **8**: e60465.

TABLES

TABLE 1. List of strains

Strain or Plasmid	Relevant Characteristics	Source or Reference
S. mutans Strains		
UA159	Wild-type	ATCC 700610
Δ*xrpA*	Inactivation of *xrpA* start codon; *comX*::T162C	(Kaspar *et al.*, 2015)
184XrpA/UA159	UA159 harboring overexpression of *xrpA* on pIB184, Em^R	(Kaspar *et al.*, 2015)
Δ*comR*	*comR* (SMU.61) :: Em^R	(Kaspar *et al.*, 2016)
Δ*comX*	*comX* (SMU.1997) :: Em^R	(Kaspar *et al.*, 2015)
PcomX::gfp/UA159	UA159 harboring *gfp* fluorescent reporter of P*comX*	(Son *et al.*, 2012)
PcomX::gfp/Δ*xrpA*	Δ*xrpA* harboring *gfp* fluorescent reporter of P*comX*	This study
PcomX::gfp/184XrpA	184XrpA/UA159 harboring *gfp* fluorescent reporter of P*comX*	This study
PcomS::gfp/UA159	UA159 harboring *gfp* fluorescent reporter of P*comS*	(Kaspar *et al.*, 2017)
PcomS::gfp/Δ*xrpA*	Δ*xrpA* harboring *gfp* fluorescent reporter of P*comS*	This study
PcomS::gfp/184XrpA	184XrpA/UA159 harboring *gfp* fluorescent reporter of P*comS*	This study
PcomX::gfp/Δ*comS*	Δ*comS* (*comS* :: Em^R) harboring *gfp* fluorescent reporter of P*comX*	(Son *et al.*, 2012)
PcomX::gfp/Δ*comS*Δ*xrpA*	ΔcomS in *comX*::T162C harboring gfp fluorescent reporter of P*comX*	This study
184ComR-Strep/UA159	UA159 harboring overexpression of *comR* with addition of (G4S)$_2$ linker and Strep-tag on pIB184, Em^R	This study
Plasmids		
pDL278	Escherichia coli - Streptococcus shuttle vector, Sp^R	(LeBlanc *et al.*, 1992)
pIB184	Shuttle expression plasmid with the constitutive P23 promoter, Em^R	(Biswas *et al.*, 2008)

991 * Em, erythromycin; Sp, spectinomycin.

993 TABLE 2. Results of 2D DIGE Spot Identification by LC-MS/MS
994

SMU Number	Gene/Protein Designation	Protein Description	Spot / Gel Slice Number*	Protein Score C.I. %	Total Ior Score C.I.
SMU.15	*ftsH*	cell division protein	S10	0	73
SMU.99	*fbaA*	fructose-bisphosphate aldolase	S36	100	100
SMU.148	*adhE*	bifunctional acetaldhyde-CoA / alcohol dehydrogenase	S3, S4	100	100
SMU.289	*sgaR*	putuative transcriptional regulator	G4	95	99
SMU.359	*fusA*	elongation factor G	S6	100	100
SMU.360	*gapC*	glyceraldehyde-3-phosphate dehydrogenase	S29, S30	100	100
SMU.546	*typA*	GTP-binding protein	S6	100	100
SMU.640c		transcriptional regulator	S47	41	0
SMU.714	EF-Tu	translation elongation factor	S14, S15	100	100
SMU.1077	*pgm*	phosphomannomutase	S11	100	100
SMU.1120		putative sugar ABC transporter	S36	0	96
SMU.1122	*cdd*	putative cytidine deaminase	S44	0	63
SMU.1178c		Amino acid ABC transporter	S20	0	43
SMU.1190	*pykF*	pyruvate kinase	S14, S15	100	100
SMU.1208c		hypothetical protein	S44, S47	0	50
SMU.1247	*eno*	enolase	S18	100	100
SMU.1367c		methylase	S19, S21	0	93
SMU.1535		glycogen phosphorylase	S24	0	44
SMU.1572	*murZ*	UDP-N-acetylglucosamine-1-carboxyvinyl transferase	S51	0	44
SMU.1599	celR	transcriptional regulator	S18	0	58
SMU.1671c		hypothetical protein	S20, S21	0	82
SMU.1791c		putative nucleotidyltransferase	S5	0	88
SMU.1954	*groEL*	protein chaperonin	S11	100	100
SMU.1967	*ssb2*	single-stranded DNA-binding protein	S56, S57, G3	100	100
SMU.2085	*recA*	DNA recombination and repair	S18, S19, S20, S21	100	100
SMU.2154c		putative peptidase	S1	0	92
#	*xrpA*	Potential inhibitor of competence	S1, S14, S15, S36, G5	0	82

995 *Spot (S) and gel slice (G) numbers correspond to Figures 4 and Supplemental Figure S1,
996 respectively
997 # Not annotated
998

999 TABLE 3. Synthesized XrpA Peptides and Determined ComR Kd from Fluorescent Polarization
1000

Peptide	Start Position	End Position	Sequence	Length	ComR Kd (nM)*
					153 ± 10
1	5	20	CISILRHVFLITLKMF	16	651 ± 99
2	22	34	VSKKVRNVVLIEC	13	198 ± 17
3	35	46	LMKKSVRLNTVC	12	169 ± 12
4	56	69	IFYFVIVCLHINKV	14	165 ± 13
N1	1	18	MIQNCISILRHVFLITLK	18	783 ± 204
N2	18	38	MFCVSKKVRNVVLIECLMKK	20	400 ± 49
S1			LFKFTCVRLILIISHM	16	248 ± 38
S2			KKNVIECVSVRVL	13	147 ± 11

*ComR Kd determined from fluorescent polarization experiment shown in Figure 7A-D

FIGURE LEGENDS

Figure 1. Transcriptome Analysis of ΔxrpA and UA159 with sXIP addition. Volcano plots of (A)

ΔxrpA and (B) UA159 + 2 μM sXIP from RNA-Seq results compared to UA159 of three

independent replicates grown in FMC medium to $OD_{600} = 0.5$. Log2 fold change and false

discovery rates (FDR) converted to –log10 P-values were calculated from Degust using edgeR

analysis. Genes of interest that were $\geq$1.5 log2 fold change and had a $\geq$4 –log10 P-value were

highlighted either in red (upregulated) or blue (downregulated) and are listed in Tables S2 and

S3. (C) Visual representation of read counts accumulated in either the *comS, comX* or *comY*

coding sequences in either the UA159 (blue bars) or *ΔxrpA* (orange bars) genetic backgrounds.

Mapped short read alignments were converted in ".bam" files and visualized with the IGV

genome browser.

1013

Figure 2. Effect of XrpA on activation of comX *and* comS *promoters.* Transcriptional activation

assays using a fused-*gfp* reporter for (A) P*comX* and (B) P*comS* in wild-type (UA159 – green

circles), *comX*::T162C (*ΔxrpA* – blue squares) or *xrpA* overexpression (184XrpA – orange

diamonds) genetic backgrounds. Black lines represent growth (OD_{600}, right axis) of each of the

respective reporter strains during the assay. Each data point shown is the average of three

independent biological replicates with four measured technical replicates.

1020

Figure 3. XrpA changes subpopulation behaviors. Histogram of cell counts from flow cytometry

analysis of the P*comX*::*gfp* reporter strain in UA159 (black lines) or *ΔxrpA* (red lines) grown in

BHI with addition of either (A) 100 nM or (B) 1000 nM sCSP. For experiments with FMC and

addition of 2 μM sXIP, reporter strains in either (C) UA159 or (D) *ΔcomS* background were

stained with propidium iodide before analysis. A total of 50,000 cells were counted in three

independent replicates for each experiment. (E) eDNA release of selected strains from three

independent overnight cultures grown in CDM media. eDNA release was calculated by taking

the arbitrary fluorescent units and dividing by the recorded OD_{600} at the time of harvest. (F)

Change in biofilm biomass compared to UA159 using either CDM media with 20 mM glucose

(blue bars) as a sole carbohydrate source or with 15 mM glucose and 2.5 mM sucrose (orange

bars) as the carbohydrate source after 48 hours of growth. Data represent the average of three

independent biological replicates with four technical replicates each. Statistical analysis was

calculated by the Student's *t*-test; * P < 0.05.

1034

1035   *Figure 4. 2D DIGE of SPINE Experiment.* Individual 2D gels of (A) pIB184, (B) ComR-Strep

1036   without sXIP addition and (C) ComR-Strep with 2 µM sXIP elutions obtained during SPINE

1037   experiment. (D) Combined image of each individual 2D gel with pIB184 in blue, ComR-Strep

1038   without sXIP in green and ComR-Strep with addition in red. Selected spots of interest are circled

1039   and labeled. Y-axis is labeled with molecular weights and X-axis with pH ranges.

1040

Figure 5. PcomX activities with addition of various synthetic XrpA peptides. Transcriptional

activation assays using a fused P*comX*::*gfp* reporter strain in CDM medium with addition of 10

µM of various synthesized XrpA peptides (Table 3). (A) Addition of XrpA peptides 1-4, (B)

comparison between XrpA-1 and a scrambled version of XpA-1, (C) Dose-dependent inhibition

of P*comX* activity by various concentrations of XrpA-1, (D) comparison between N-terminal

specific XrpA-N1 and XrpA-N2 peptides. Colored lines represent relative P*comX* expression

(arbitrary fluorescent units divided by OD_{600}, left axis). Black lines represent growth (OD_{600}, right

axis) of each of the respective reporter strains during the assay. Each data point shown is the

average of three independent biological replicates with four measured technical replicates.

1050

Figure 6. XrpA changes subpopulation behaviors. (A) Fold change in transformation efficiency

with 10 μM XrpA peptides compared to DMSO-only control (vehicle; =1) in FMC medium with

0.5 μM sXIP addition. Data represents the average of three independent replicates. (B) Western

blot using 10 μg of whole cell lysates of UA159 with addition of 10 μM XrpA peptides in FMC

medium with 0.5 μM sXIP addition. Cells were grown to $OD_{600\ nm}$ = 0.5 before harvesting.

Primary antisera, raised against the corresponding protein, were used for detection. ManL (the

EIIAB domain of the glucose PTS permease) served as a loading control. Numbers under each

respective blot represent densitometry readings with the DMSO control set to 100%. (C)

Change in biofilm biomass of UA159 with addition of 10 μM XrpA peptides using either CDM

media with 20 mM glucose (blue bars) as a sole carbohydrate source or with 15 mM glucose

and 2.5 mM sucrose (orange bars) as the carbohydrate source after 48 hours of growth. Data

represents the average of three independent replicates with four technical replicates each. (D)

eDNA release of UA159 with addition of 10 μM XrpA peptides from three independent overnight

cultures grown in CDM media. eDNA release was calculated by taking the arbitrary fluorescent

units and dividing by the recorded OD_{600} at the time of harvest. All statistical analysis for this

figure was calculated by the student's T-Test, * $P < 0.001$.

1067

Figure 7. Fluorescent Polarization experiments confirm XrpA-ComR interaction. Fluorescent

polarization (FP) curves of increasing concentrations of purified ComR binding to 10 nM of

P*comX* dsDNA probe in the presence of 10 μM sXIP and 10 μM of various XrpA peptides (Table

3). Control (black lines) represents binding in the absence of XrpA peptides. (A) addition of XrpA

peptides 1-4, (B) addition of XrpA peptides N1 and N2, (C) comparison between XrpA-1 and a

scrambled version of XrpA-1 and (D) comparison between XrpA-2 and a scrambled version of

XrpA-2. (E). Binding of 10 nM FITC-labeled XrpA-N1 peptide to increasing concentrations of

purified ComR, with and without 10 μM sXIP addition. No DNA probe is present in this

experiment. (F) Increasing concentrations of synthetic unlabeled XrpA-N1 peptide were

assessed for their ability to compete with FITC-labeled XrpA-N1 for their binding to increasing

concentrations of purified ComR. No DNA probe is present in this experiment. Data shown are

averaged from three independent experiments. Kd values for figures A-D are shown in Table 3.

1080

Figure 8. Model for XrpA Modulation of ComRS Signaling. Current model for the role of XrpA in

inhibition of ComRS activity and in cell fate. When present and active, XrpA (yellow) interacts

with ComR (blue) independently of whether ComR is in a complex with XIP (orange), resulting

in diminished affinity of ComR for its target in P*comX*. XrpA inhibition of ComR prevents the

ComR-XIP complex from over-amplifying the competence activation signal, thereby maintaining

a balance within the population of cells that induce competence and internalize DNA with a

group of cells that undergo lysis providing DNA as a nutrient source, a source for genetic

diversification, and/or a source of eDNA that contributes to extracellular matrix formation. One

potential mechanism by which XrpA curtails proficient ComR-XIP activation of P*comX* is through

inefficient dimer formation between ComR-XIP complexes. When XrpA is absent from the

circuit, as in the case of the *xrpA* mutant (Δ*xrpA*; *comX*::T162C), over-amplification of ComRS

signaling occurs. This leads to increases in the accumulation of the sigma factor ComX (green);

which in turn results in an increase in the subpopulation of cells that undergo cell lysis.

1094

UA159 (WT)

XIP

XrpA

ComR

P*comX*

comR

comX

xrpA

ComX

Competence — Lysis

Cell Fate

△**xrpA (*comX*::T162C)**

P*comX*

comR

comX

xrpA

ComX

ComX

Competence — Lysis

Cell Fate