1 Transcriptional Activation of Elephant Shark Mineralocorticoid Receptor by Corticosteroids,
2 Progestins and Spironolactone
3

4 Yoshinao Katsu[1,2] [*], Satomi Kohno[3], Kaori Oka[1], Xiaozhi Lin[1], Sumika Otake[1], Nisha E.
5 Pillai[4], Wataru Takagi[5], Susumu Hyodo[5], Byrappa Venkatesh[4], Michael E. Baker[6, *]
6
[1]Graduate School of Life Science, Hokkaido University, Sapporo, Japan; [2]Department of
Biological Sciences, Hokkaido University, Sapporo, Japan; [3]Department of Obstetrics and
Gynecology, Medical University of South Carolina, and Marine Biomedicine and
Environmental Science Center, Hollings Marine Laboratory, Charleston, SC, USA;
[4]Comparative Genomics Laboratory, Institute of Molecular and Cell Biology, Agency for
Science, Technology and Research (A*STAR), Singapore; [5]Laboratory of Physiology,
Atmosphere and Ocean Research Institute, University of Tokyo, Chiba, Japan; [6]Division of
Nephrology, Department of Medicine, University of California, San Diego, CA, USA
15
Correspondence:
ykatsu@sci.hokudai.ac.jp
mbaker@ucsd.edu

Summary. We report the first analysis of corticosteroid and progesterone activation of
full-length mineralocorticoid receptor (MR) from elephant shark (*Callorhinchus milii*), a
cartilaginous fish belonging to the oldest group of jawed vertebrates.   Aldosterone, cortisol,
11-deoxycorticosterone, corticosterone, 11-deoxcortisol, progesterone and
19-norprogesterone had half-maximal responses (EC50s) below 1 nM for elephant shark MR,
and hence, these steroids, including progesterone, are potential physiological
mineralocorticoids.   Aldosterone, the physiological mineralocorticoid in humans and other
terrestrial vertebrates, is not found in cartilaginous or ray-finned fishes.   Progesterone is a
precursor for corticosteroids that activate elephant shark MR, suggesting that progesterone
was an ancestral ligand for the MR.   Progesterone does not activate human, amphibian or
alligator MRs indicating that progesterone activation of the MR was lost during the transition
to terrestrial vertebrates.   RNA-seq analysis finds strong MR expression in elephant shark
ovary and testis, in which progesterone-activated MR may have novel functions.

Introduction

The mineralocorticoid receptor (MR) belongs to the nuclear receptor family, a large and diverse group of transcription factors that also includes receptors for glucocorticoids (GR), progestins (PR) androgens (AR) and estrogens (ER) (*1, 2*). Sequence analysis revealed that the MR and GR are closely related (*3*); phylogenetic analysis indicates that MR and GR evolved from a corticosteroid receptor (CR) that was present in jawless vertebrates, such as lamprey and hagfish (*4-7*). A distinct mineralocorticoid receptor (MR) first appears in cartilaginous fishes (Chondrichthyes), the oldest group of extant jawed vertebrates (gnathostomes) that diverged from bony vertebrates about 450 million years and are a crucial group in understanding the origin and evolution of jawed vertebrate morphology and physiology (*8, 9*). Like mammals, cartilaginous fishes contain the full complement of adrenal and sex steroid receptors: AR, ER, GR, MR and PR (*1, 2, 4, 10*).

Aldosterone (Aldo) is the physiological activator of transcription of human MR in epithelial tissues, such as the kidney distal collecting tubules and the colon, in which the MR regulates electrolyte homeostasis (*6, 11-14*). The MR also is found in brain, heart, aorta, lung, liver, spleen, adipose tissue, testis, breast and ovary (*12-18*). The physiological function of the MR in these tissues is still being elucidated.

The MR and other steroid receptors have a characteristic modular structure consisting of an N-terminal domain (NTD) (domains A and B), a central DNA-binding domain (DBD) (domain C), a hinge domain (D) and a C-terminal ligand-binding domain (LBD) (domain E) (*2, 4, 19-21*) (Figure 1). The E domain alone is competent to bind steroids (*4, 19, 22-24*). Interactions between the NTD (A/B domains) and the LBD and coactivators are important regulators of transcriptional activation of mammalian MRs (*25-32*) and fish MRs (*31-33*). Moreover, GAL-DBD-hinge-LBD constructs of zebrafish MR have different responses to progestins and some corticosteroids than do GAL-DBD-hinge-LBD constructs of human, chicken, alligator and Xenopus MRs (*32*).

Figure 1. Comparison of domains in elephant shark MR with vertebrate MRs. MRs from elephant shark (shark), zebrafish, coelacanth, Xenopus (frog), chicken and human are compared. The functional A/B domain to E domains are schematically represented with the numbers of amino acid residues and the percentage of amino acid identity is depicted. GenBank accession numbers: elephant shark MR (XP_007902220), zebrafish MR (NP_001093873), coelacanth MR (XP_014348128), Xenopus (NP_001084074), chicken (ACO37437), human MR (NP_000892).

The timing of the evolution of this difference between full-length and truncated MRs in ray-finned fishes and terrestrial vertebrates, as well as expression of the MR in non-epithelial tissues is not known. Also unresolved is the identity of the ancestral mineralocorticoid in cartilaginous fishes and the current mineralocorticoid in ray-finned fish because aldosterone (Aldo), the physiological mineralocorticoid in terrestrial vertebrates, is not found in cartilaginous fishes or ray-finned fishes. Aldo first appears in lobe-finned fish (*34*), the forerunners of terrestrial vertebrates (*35*). Thus, the identity of the physiological mineralocorticoid in cartilaginous fishes and ray-finned fishes is not established, although cortisol and 11-deoxycorticosterone have been proposed (*36-41*).

Complicating the identity of the physiological mineralocorticoid in cartilaginous and

ray-finned fishes is evidence that progesterone (Prog) and 19-norprogesterone (19norProg),

along with spironolactone (Spiron) (Figure 2), are transcriptional activators of several

ray-finned fish MRs (*21, 33, 40*), including zebrafish MR (*42*), and of chicken MR (*32*).   In

contrast, these steroids are antagonists for human MR (*22, 24, 43*), alligator MR and Xenopus

MR (*32*).   Ray-finned fish MRs and chicken MR differ in their response to Prog, 19norProg

and Spiron, raising the question of whether the response to progestins and Spiron evolved in

ray-finned fish before or after the divergence of ray-finned fish from the lobe-finned fish

lineage that led to tetrapods.

Transcriptional activation by corticosteroids and other 3-ketosteroids of a full-length

cartilaginous fish MR, which is the physiological MR, has not been investigated.   Only

truncated skate MR (*44*), consisting of the GAL4 DBD fused to the D domain and E domain

of the MR (MR-LBD) has been studied for its response to corticosteroids.   Carroll et al.

found that aldosterone (Aldo), corticosterone, 11-deoxycorticosterone, and cortisol (Figure 2)

are transcriptional activators of truncated skate MR.

Figure 2. Structures of steroids that are ligands for the MR. Aldo, 11-deoxycorticosterone and S are physiological mineralocorticoids in terrestrial vertebrates (*4, 6, 12, 45*). S is both a mineralocorticoid and a glucocorticoid in lamprey (*4, 46*) and a glucocorticoid in ray-finned fish (*47*). Cortisol is a physiological glucocorticoid in terrestrial vertebrates and ray-finned fish (*4, 10, 48-50*). corticosterone is a glucocorticoid in rats and mice (*4*). Aldo, 11-deoxycorticosterone, cortisol, corticosterone and Prog have a similar high affinity for human MR (*3, 51-53*). Prog, 19norProg, 17OH-Prog and Spiron are antagonists for human MR (*21, 43, 51*) and rat MR (*54, 55*). Prog, 19norProg, and Spiron are agonists for fish MRs (*21, 33, 40*). 19norProg is a weak agonist for rat MR (*56, 57*).

Elephant shark MR is an attractive receptor to study early events in the evolution of

mechanisms for regulating MR transcription because genomic analyses reveal that elephant

shark genes are evolving slowly (*9*), making their genes windows into the past.   Thus, we

decided to investigate transcriptional activation by Aldo, 11-deoxycorticosterone,

corticosterone, 11-deoxycortisol, cortisol, Prog, 19norProg, 17-hydroxyprogesterone

(17OH-Prog) and Spiron of full-length and truncated elephant shark MR.   Interestingly, all

3-ketosteroids, including progesterone, had EC50s of 1 nM or lower for full-length elephant

shark MR.   Transcriptional activation by Prog, 19norProg and Spiron of truncated elephant

shark MR resembled that of zebrafish MR and not chicken MR, indicating that progestin

activation of MR is an ancestral response, conserved in ray-finned fish, lost in Xenopus,

alligator and human MRs, and distinct from activation of chicken MR, which arose

independently.   We also performed RNA-seq analysis of elephant shark MR and find

widespread expression of MR in various elephant shark organs (gill, kidney, heart, intestine,

liver, spleen, brain), with strong MR expression in ovary and testis, which are likely targets

for Prog, suggesting a role for Prog-MR complexes in elephant shark reproduction, as well as

in some other functions in other MR-containing organs.   Indeed, our data suggest that Prog

or a related progestin may have been one of the ancestral mineralocorticoids.   Elephant

shark RNA-seq analysis also indicates that expression of human MR in diverse organs, such

as brain, heart, liver, spleen, ovary and testis, evolved in an ancestral cartilaginous fish.

Results

Functional domains of elephant shark MR and other vertebrate MRs.

In Figure 1, we compare the functional domains of elephant shark MR to selected

vertebrate MRs.   Elephant shark MR and human MR have 92% and 67% identity in DBD

and LBD, respectively.   Interestingly, elephant shark MR has similar conservation to the

DBD (91-92%) and LBD (64-69%) of other MRs.   The A, B and D domains of elephant

shark MR and other MRs are much less conserved.

Transcriptional activation of full-length and truncated elephant shark MR by

corticosteroids, progestins and spironolactone.

We screened a panel of steroids at 0.1 nM and 1 nM for transcriptional activation of

full-length and truncated elephant shark MR.   Aldo, cortisol, corticosterone,

11-deoxycorticosterone and 11-deoxycortisol were strong activators of full-length elephant

shark MR (Figure 3A) indicating that elephant shark MR has broad specificity for

corticosteroids.   Interestingly, at these low concentrations, all progestins and Spiron are

transcriptional activators of full-length elephant shark MR, with 19-norProg having the

strongest activity and 17OHProg having the weakest activity (Figure 3A).

In parallel experiments, truncated elephant shark MR, lacking the A/B domain and

containing a GAL4-DBD instead of the MR DBD, retained a strong response to all

corticosteroids and 19nor Prog (Figure 3B).   Prog and Spiron had significant, but reduced

activity, while 17OH-Prog had little activity for truncated elephant shark MR.

Figure 3. Transcriptional activation of elephant shark MR by 3-ketosteroids.   Full length and

truncated elephant shark MR were expressed in HEK293 cells with an MMTV-luciferase reporter.

A. Full length elephant shark MR.   Cells were treated with 0.1 nM or 1.0 nM Aldo, cortisol,

corticosterone, 11-deoxycorticosterone, 11-deoxycortisol, Prog, 19norProg (19NP), 17OH-Prog,

Spiron or vehicle alone (DMSO).

B. Truncated elephant shark MR.   Cells were treated with 0.1 nM or 1.0 nM Aldo, cortisol,

corticosterone, 11-deoxycorticosterone, 11-deoxycortisol, Prog, 19norProg (19NP), 17OH-Prog or

Spiron or vehicle alone (DMSO).   Results are expressed as means $\pm$ SEM, n=3.   Y-axis indicates

fold-activation compared to the activity of control vector with vehicle (DMSO) alone as 1.

EC50 values for steroid activation of elephant shark MR

Concentration-dependence of transcriptional activation of full length elephant shark

MR by corticosteroids (Aldo, cortisol, corticosterone, 11-deoxycorticosterone,

11-deoxycortisol) is shown in Figure 4A and by progestins (Prog, 19norProg, 17OH-Prog)

and Spiron in Figure 4B.   The corresponding concentration-dependent curves for truncated

elephant shark MR are shown in Figures 4C and 4D, respectively.   Table 1 summarizes the

EC50s of corticosteroids for full-length and truncated elephant shark MR.   Table 1 also

contains, for comparison, previously determined EC50s of corticosteroids for full-length and

truncated human, chicken, alligator, Xenopus and zebrafish MRs (*32*) and skate MR (*44*).

Figure 4. Concentration-dependent transcriptional activation of full-length and truncated

elephant shark MR (shark MR) by 3-ketosteroids.   Full length and truncated elephant shark MR

(shark MR) were expressed in HEK293 cells with an MMTV-luciferase reporter.   Full-length

elephant shark MR (shark MR) (A) and (C) and truncated elephant shark MR (shark MR) (B) and (D).

(A), (B) Aldo, cortisol, corticosterone, 11deoxycorticosterone or 11-deoxycortisol.

(C), (D) Aldo, Prog, 19norProg (19nor-P), 17OH-Prog (17OH-P) or Spiron.   Cells transfected with

elephant shark MR were treated with increasing concentrations of cortisol, corticosterone, Aldo,

11deoxycorticosterone, 11-deoxycortisol, Prog, 19norProg, 17OH-Prog, Spiron or vehicle alone

(DMSO).   Y-axis indicates fold-activation compared to the activity of control vector with vehicle

(DMSO) alone as 1.

Table 1. Corticosteroid activation of full-length MR and GAL4-DBD-MR-LBD

MR	Aldo	B	F	DOC	S
	EC50 (M)	EC50 (M)	EC50 (M)	EC50 (M)	EC50 (M)
Elephant shark Full	1.1×10^{-10}	1.7×10^{-10}	4.6×10^{-10}	6.3×10^{-11}	1.1×10^{-10}
	100%	101%	114%	83%	83%
Elephant shark LBD	3.7×10^{-11}	9.9×10^{-11}	1.9×10^{-10}	2.4×10^{-11}	6.8×10^{-11}
	100%	90%	79%	81%	77%
[1]**Skate LBD**	7×10^{-11}	1×10^{-10}	1×10^{-9}	3×10^{-11}	2.2×10^{-8}
[2]**Human Full**	2.7×10^{-10}	1.2×10^{-9}	5.5×10^{-9}	4.2×10^{-10}	3.6×10^{-9}
	100%	119%	133%	74%	42%
[2]**Human LBD**	2.8×10^{-10}	5.9×10^{-10}	$3. \times 10^{-9}$	1.8×10^{-9}	#-
	100%	95%	74%	44%	*8%
[2]**Chicken Full**	6.2×10^{-11}	5.1×10^{-11}	2.8×10^{-10}	3.4×10^{-11}	6.7×10^{-10}
	100%	109%	128%	110%	112%
[2]**Chicken LBD**	1.3×10^{-10}	1.6×10^{-10}	6.9×10^{-10}	1.7×10^{-10}	4.7×10^{-9}
	100%	92%	75%	92%	36%
[2]**Alligator-Full**	2.8×10^{-10}	3.6×10^{-10}	6.9×10^{-9}	2.3×10^{-10}	2.7×10^{-9}
	100%	138%	176%	85%	45%
[2]**Alligator LBD**	3.5×10^{-10}	3.8×10^{-10}	2.3×10^{-9}	5.2×10^{-10}	#-
	100%	88%	68%	51%	*8%
[2]**Xenopus Full**	4.6×10^{-10}	6.2×10^{-10}	1.1×10^{-8}	7.6×10^{-10}	9.1×10^{-9}
	100%	105%	126%	59%	31%
[2]**Xenopus-LBD**	1.5×10^{-9}	1.9×10^{-9}	1.2×10^{-8}	#-	#-
	100%	74%	37%	*10%	*6%
[2]**Zebrafish Full**	8.2×10^{-11}	3.0×10^{-10}	4.4×10^{-10}	6.3×10^{-11}	4.0×10^{-10}
	100%	112%	123%	103%	94%
[2]**Zebrafish LBD**	2.7×10^{-11}	1.5×10^{-10}	3.1×10^{-10}	1.0×10^{-10}	9.1×10^{-10}
	100%	96%	77%	99%	67%

[1] (*44*)

[2] (*32*) doi:doi: https://doi.org/10.1101/151233)

Curve did not saturate.

(%) Relative induction to Aldosterone induced activation.

* Relative induction at 1 µM compared to Aldo.

Previously we reported that progestins and Spiron are transcriptional activators of

full-length chicken and zebrafish MRs and truncated zebrafish MR (*32*).   However, EC50

values of progestins and Spiron for these MR were not determined.   We have remedied this

omission and report their EC50 values in Table 2 and Figure 5, for comparison with

full-length and truncated elephant shark MR.

Figure 5. Concentration-dependent transcriptional activation of full-length and truncated

chicken and zebrafish MR by progestins and Spironolactone.   HEK293 cells transfected with

chicken and zebrafish MR were treated with increasing concentrations of Prog, 19norProg,

17OH-Prog or Spiron.   Full-length chicken MR (A) and zebrafish MR (B) and truncated chicken MR

(C) and zebrafish MR (D).   Y-axis indicates fold-activation compared to the activity of control vector

with vehicle (DMSO) alone as 1.

Table 2. EC50 values for progestin and spironolactone activation of full-length and

GAL4-DBD-MR-LBD constructs of elephant shark, zebrafish and chicken MR

MR	Aldo	Prog	17OH-Prog	19norProg	Spiron
Elephant shark Full	1.1×10^{-10}	2.7×10^{-10}	1.4×10^{-9}	4.3×10^{-11}	5.5×10^{-10}
	100%	43%	25%	84%	45%
Elephant shark LBD	3.7×10^{-11}	4.8×10^{-10}	2.9×10^{-9}	1.8×10^{-11}	4.2×10^{-10}
	100%	40%	26%	98%	53%
[2]Zebrafish Full	8.2×10^{-11}	2.4×10^{-9}	1.8×10^{-8}	9.4×10^{-10}	3.8×10^{-9}
	100%	77%	44%	83%	54%
[2]Zebrafish LBD	2.7×10^{-11}	9.8×10^{-8}	#-	6.4×10^{-8}	#-
	100%	122%	*24%	122%	*73%
[2]Chicken Full	6.2×10^{-11}	7.1×10^{-10}	2.9×10^{-8}	6.8×10^{-10}	5.1×10^{-9}
	100%	62%	15%	68%	30%
[2]Chicken LBD	1.3×10^{-10}	#-	#-	#-	#-
	100%	*21%	-	*29%	-

[2] (*32*) doi:doi: https://doi.org/10.1101/151233)# Curve did not saturate.

(%) Relative induction to Aldosterone induced activation.

* Relative induction at 1 µM compared to Aldo.

RNA-seq analysis of elephant shark MR

We examined expression of level of elephant shark MR gene in 10 tissues based on

RNA-seq data (Figure 6). The MR gene was found to express widely in all tissues,

including gills and kidney, two traditional mineralocorticoid-responsive tissues.

Interestingly there was considerably higher expression in the ovary and testis, the two

reproductive tissues analyzed.

Figure 6. Expression level of MR gene in 10 tissues of elephant shark estimated based on

RNA-seq data. Transcript abundances are shown in terms of normalized counts called Fragments

per kilobase of exon per million fragments mapped (FPKM) (*58*). FPKM is estimated by normalizing

gene length followed by normalizing for sequencing depth.

RNA-seq analysis of human MR

   RNA-seq analysis of human MR (*59*) (Figure 7) reveals that the MR is expressed in

kidney, colon, brain, heart, liver, ovary, spleen and testis.  This diverse pattern of expression

is similar to that of elephant shark MR.

Figure 7. Expression level of human MR based on RNA-seq data. Transcript abundances are

shown in terms of normalized counts called Reads Per Kilobase of transcript per Million mapped

reads (RPKM) (*59*).

Discussion

Cartilaginous fish, including elephant sharks, occupy a key position in the evolution

of vertebrates as an out-group to ray-finned fish, the largest group of extant vertebrates, and

the lobe-finned fish, which are the forerunners of terrestrial vertebrates.   Importantly, the

elephant shark genome is evolving slowly (*9*), making it attractive for studying ancestral

proteins, including the MR, which first appears as a distinct MR ortholog in cartilaginous fish

(*5, 6, 10, 44*).

This is the first report of transcriptional activation of a full-length cartilaginous fish

MR by a panel of five corticosteroids and Prog, 19norProg, 17OH-Prog and Spiron.

Full-length elephant shark MR has EC50s below 1 nM for five corticosteroids, (Figure 4,

Table 1) including Aldo, a steroid that has not been found in cartilaginous fish.   Prog,

19norProg and Spiron also have sub-nM EC50s for full-length elephant shark MR.   In

addition to their low EC50s, all of these corticosteroids and 19norProg have similar fold

activation of transcription of full-length MR (Figure 4).   Thus, several corticosteroids, as

well as 19norProg and Prog, are potential physiological mineralocorticoids for elephant shark

MR.

Compared to full-length elephant shark MR, the EC50s of all five corticosteroids and

19norProg are lower for its truncated MR, while the EC50 for Spiron is slightly lower, and

the EC50s for Prog and 17OH-Prog are about 2-fold higher (Table 1, Table 2).   Compared to

fold activation by Aldo, the responses to 11-deoxycorticosterone, S, Prog and 17OH-Prog are

similar for full-length and truncated elephant shark MR.   However, the relative responses to

corticosterone and cortisol of truncated elephant shark MR are 10% and 35% lower,

respectively, (Figure 4, Table 1) and for 19norProg and Spiron, 14% and 8% higher

respectively, than for full-length elephant shark MR (Figure 4, Table 2).   Regarding

truncated skate MR, most of the EC50s of corticosteroids (*44*) are similar to that for elephant

shark MR (Table 1).   The exception is 11-deoxycortisol, which has over 200-fold higher

EC50 for skate MR than for elephant shark MR.

Corticosteroid and progestin activation of full-length and truncated elephant shark,

terrestrial vertebrate and zebrafish MRs

Comparison of transcriptional activation by corticosteroids and progestins of

full-length and truncated elephant shark MR with full-length and truncated human, chicken,

alligator, Xenopus and zebrafish MRs (Table 1) provides insights into the evolution of

regulation of steroid-mediated transcription of these MRs by allosteric interactions between

the NTD/DBD and LBD.   Like elephant shark MR, truncated terrestrial vertebrate MRs and

zebrafish MR have similar EC50s for Aldo, corticosterone and cortisol as their full-length

counterparts (Table 1).   However, the response of truncated human, alligator and Xenopus

MRs to 11-deoxycortisol did not saturate at 1 µM, preventing us from determining the EC50

and explaining the low fold activation by S (Table 1).   Interestingly, truncated chicken and

zebrafish MRs, which are activated by progestins, have nM EC50s for 11-deoxycortisol,

although fold activation by 11-deoxycortisol is lower than that of Aldo.   The similar activity

of truncated human, alligator and Xenopus MRs to Aldo, corticosterone and cortisol indicates

that allosteric interactions between the NTD and LBD are steroid-selective.   Overall, it

appears that corticosteroid activation, especially by 11-deoxycortisol, of elephant shark MR is

less sensitive to the loss of NTD/DBD than are human, chicken, alligator and Xenopus MRs.

Progestin activation of full-length and truncated chicken and zebrafish MRs

272   Transcriptional activation by progestins of chicken and zebrafish MRs differs from

that of elephant shark MR, especially for their truncated MRs.   Although Prog, 19norProg

and Spiron have nM EC50s for transcriptional activation of full-length chicken and zebrafish

MRs, at 1 µM Prog 19norProg and Spiron transcriptional activation of truncated chicken MR

did not saturate, and at 1 µM Spiron transcriptional activation of zebrafish MR did not

saturate (Figure 5, Table 2).   Fold activation compared to Aldo by all progestins and Spiron

is substantially lower for truncated chicken MR and for Spiron activation of zebrafish MR.

However, Prog and 19norProg have EC50s of 98 nM and 64 nM, respectively, for truncated

zebrafish MR.   In this respect, truncated zebrafish MR is closer to truncated elephant shark

MR than is truncated chicken MR.   These data suggest that transcriptional activation by

progestins of chicken MR appears to have evolved independently in terrestrial vertebrates.

Progestins may be mineralocorticoids in cartilaginous fishes

284   Prog is a precursor for the synthesis of the other corticosteroids (*5, 60*).   Two

parsimonious metabolites of progesterone are 11-deoxycorticosterone

(21-hydroxyprogesterone) and 19norProg (Figure 2), which have EC50s, 0.063 nM and 0.043

nM, respectively, for elephant shark MR.   These are the two lowest EC50s among the tested

steroids.   Moreover, 19norProg evokes a stronger response from elephant shark MR than

Aldo (Figure 4A, B).   C19 demethylase activity has been found in mammalian kidney (*56*).

If C19 demethylase is present in elephant shark, then 19norProg needs to be considered as a

potential physiological mineralocorticoid.

292   The strong response to 19norProg is interesting because *in vivo* studies in rats

revealed that 19norProg is at least a 100-fold weaker MR agonist than Aldo (*56*), while in

transfected cells, at 1 nM, 19norProg is an antagonist (*21, 43*).   Unexpectedly, compared to

Aldo, 19norAldo has less than 1% binding affinity for rat MR (*61*) indicating that the

mechanism by which C19 demethylation of Prog increases this steroid's transcriptional

activity for elephant shark MR is likely to be complex.

We propose that transcriptional activation of elephant shark MR by 19norProg, as

well as by Prog and Spiron, can be explained by a mechanism based on Geller et al.'s (*43*)

discovery that S810L human MR mutant is transcriptionally activated by 1 nM Prog,

19norProg, and Spiron, unlike wild-type human MR, in which these steroids are MR

antagonists.   Geller et al. used a 3D model of S810L MR and transcriptional analysis of a

series of mutations at Ser-810 (helix 5) and Ala-773 (helix 3) to propose that a van der Waals

contact between Leu-810 and Ala-773 was sufficient for transcriptional activation of S810L

MR by progestins.   In Figure 3C of Geller et al. (*43*), the human S810M mutant was

activated by 19norProg.   Elephant shark MR and skate MR contain a methionine at this

position and an alanine corresponding to Ala-773 (Figure 8).   Based on Geller et al.'s model,

we propose that transcriptional activation of elephant shark MR by 19norProg is due to a van

der Waals contact between Met-782 (helix 5) and Ala-745 (helix 3), which stabilizes the A

ring of 19norProg, promoting transcriptional activation.

Figure 8. Alignment of elephant shark MR to Serine-810 and Alanine-773 in helices 3-5 in human MR.

Elephant shark MR and skate MR contain a methionine corresponding to human Ser-810 and an alanine

corresponding to Ala-773.   Lamprey CR and hagfish CR also contain a corresponding methionine, as well as a

cysteine corresponding to Ala-773.   Human Ser-810 and Ala-773 are conserved in coelacanths, terrestrial

vertebrate and ray-finned fish MRs.   Amino acids that are identical to amino acids in human MR are denoted

by (-).

Our results indicate that progestin activation of the MR is an ancient response.   The

timing of evolution of Prog as an MR antagonist is not known.   One necessary, but not

sufficient event, is the evolution in helix 5 of a serine corresponding to serine-810 in human

MR.　This occurs in coelacanth MR, which contains Ser-816, corresponding to human MR

Ser-810 (Figure 8) and to elephant shark MR Met-782.　This mutation in coelacanth MR

would be expected to eliminate the proposed van der Waals contact between helix 5 and helix

3, leading to antagonist activity of Prog in some terrestrial vertebrate MRs (*32*).　However, a

serine corresponding to Ser-810 in human MR also is present in chicken MR and ray-finned

fish MRs (Figure 8), indicating another mechanism is involved in Prog activation of the MR

in these vertebrates.

Role for elephant shark MR in reproductive physiology

The evidence from RNA-seq analysis of high MR expression in ovary and testis

(Figure 6) suggests that a Prog-MR complex is important in reproductive responses in

elephant shark.　Of course, Prog also acts as a reproductive steroid in ovary and testis via

transcriptonal activation of the PR (*62, 63*).　Based on evidence that Prog activates the MR

in several ray-finned fishes (*21, 33, 40, 42*), a Prog-MR complex also may activate

transcription in reproductive tissues and other organs in ray-finned fish, as well as

cartilaginous fish.

RNA-seq analysis also finds MR expression in elephant shark gills and kidneys

(Figure 6), two classical targets for MR regulation of electrolyte transport (*6*).　RNA-seq

analysis also identifies MR expression in elephant shark heart and brain, two other organs in

which corticosteroids have important physiological actions via the MR (*15-17, 64-66*).

Indeed, RNA seq analysis of elephant shark MR indicates that expression the MR in diverse

organs was conserved during the descent from cartilaginous fishes to humans.

Materials and Methods

Chemical reagents

Aldo, cortisol, corticosterone, 11-deoxycorticosterone, 11-deoxycortisol, Prog,

19norProg, 17OH-Prog and Spiron were purchased from Sigma-Aldrich.　For reporter gene

assays, all hormones were dissolved in dimethylsulfoxide (DMSO); the final DMSO

concentration in the culture medium did not exceed 0.1%.

Construction of plasmid vectors

The full-coding regions from elephant shark MR were amplified by PCR with KOD

DNA polymerase.　The PCR products were gel-purified and ligated into pcDNA3.1 vector

(Invitrogen) for the full-coding region or pBIND vector for D-E domains {Oka, 2013 #48}.

RNA-sequence analysis

The RNA-seq reads from following tissues of elephant shark were down-loaded

from the Sequence Read Archive database of NCBI (accession number SRA054255): brain,

gills, heart, intestine, kidney, liver, muscle, ovary, spleen, and testis.   For each tissue,

sequences were assembled *de novo* using Trinity, version r2013-08-14 {Grabherr, 2011

#396}.

Gene Expression Analyses

362   To determine the expression level of MR genes, we performed abundance

estimation of transcripts from the afore mentioned 10 tissues. Trinity transcripts from all ten

tissues and full-length cDNA sequence of the MR gene were combined together and clustered

using CD-HITv4.6.1 at 100% identity {Li, 2006 #394}. RNA-seq reads from each of the ten

tissues was independently aligned to the clustered transcript sequences and abundance of MR

gene transcripts was estimated by RSEMv1.2.25 (*58*) which uses bowtiev2.2.6 for aligning

(*67*). Transcript abundances were measured in terms of normalized counts called Fragments

per kilobase of exon per million fragments mapped (FPKM) (*58*). FPKM is estimated by

normalizing the gene length followed by normalizing for sequencing depth.

Transactivation Assay and Statistical Methods

372   Transfection and reporter assays were carried out in HEK293 cells, as described

previously (*68, 69*).   All experiments were performed in triplicate.   The values shown are

mean ± SEM from three separate experiments, and dose-response data and EC50 were

analyzed using GraphPad Prism.   Comparisons between two groups were performed using

t-test, and all multi-group comparisons were performed using one-way ANOVA followed by

Bonferroni test. $P < 0.05$ was considered statistically significant.

Acknowledgments: K.O. was supported by the Japan Society for the Promotion of Science

(JSPS) Research Fellowships for Young Scientists.   This work was supported in part by

Grants-in-Aid for Scientific Research 23570067 and 26440159 (YK) from the Ministry of

Education, Culture, Sports, Science and Technology of Japan.   M.E.B. was supported by

Research fund #3096.   BV was supported by the Biomedical Research Council of A*STAR,

Singapore.

Author contributions: Y.K., S.K., K.O., X.L., S.O. and N.E.P. carried out the research.

Y.K. and S.K. analyzed data.   W.T. and S.H. aided in the collection of animals.   NEP

assembled and analyzed RNAseq data.   Y.K. and M.E.B. conceived and designed the

experiments.   Y.K., M.E.B. and B.V. wrote the paper.   All authors gave final approval for

publication.

Declaration of interests: We have no competing interests.

References

1. M. E. Baker, D. R. Nelson, R. A. Studer, Origin of the response to adrenal and sex
steroids: Roles of promiscuity and co-evolution of enzymes and steroid receptors. *J*
Steroid Biochem Mol Biol **151**, 12-24 (2015).

2. R. M. Evans, The steroid and thyroid hormone receptor superfamily. *Science* **240**,
889-895 (1988).

3. J. L. Arriza, C. Weinberger, G. Cerelli, T. M. Glaser, B. L. Handelin, D. E. Housman,
R. M. Evans, Cloning of human mineralocorticoid receptor complementary DNA:
structural and functional kinship with the glucocorticoid receptor. *Science* **237**,
268-275 (1987).

4. M. E. Baker, J. W. Funder, S. R. Kattoula, Evolution of hormone selectivity in
glucocorticoid and mineralocorticoid receptors. *J Steroid Biochem Mol Biol* **137**,
57-70 (2013).

5. M. E. Baker, Y. Katsu, 30 YEARS OF THE MINERALOCORTICOID RECEPTOR:
Evolution of the mineralocorticoid receptor: sequence, structure and function. *J*
Endocrinol **234**, T1-T16 (2017).

6. B. C. Rossier, M. E. Baker, R. A. Studer, Epithelial sodium transport and its control
by aldosterone: the story of our internal environment revisited. *Physiological reviews*
95, 297-340 (2015).

7. J. W. Thornton, Evolution of vertebrate steroid receptors from an ancestral estrogen
receptor by ligand exploitation and serial genome expansions. *Proc Natl Acad Sci U S*
A **98**, 5671-5676 (2001).

8. J. G. Inoue, M. Miya, K. Lam, B. H. Tay, J. A. Danks, J. Bell, T. I. Walker, B.
Venkatesh, Evolutionary origin and phylogeny of the modern holocephalans
(Chondrichthyes: Chimaeriformes): a mitogenomic perspective. *Molecular biology*
and evolution **27**, 2576-2586 (2010).

9. B. Venkatesh, A. P. Lee, V. Ravi, A. K. Maurya, M. M. Lian, J. B. Swann, Y. Ohta, M.
F. Flajnik, Y. Sutoh, M. Kasahara, S. Hoon, V. Gangu, S. W. Roy, M. Irimia, V. Korzh,
I. Kondrychyn, Z. W. Lim, B. H. Tay, S. Tohari, K. W. Kong, S. Ho, B.
Lorente-Galdos, J. Quilez, T. Marques-Bonet, B. J. Raney, P. W. Ingham, A. Tay, L. W.
Hillier, P. Minx, T. Boehm, R. K. Wilson, S. Brenner, W. C. Warren, Elephant shark
genome provides unique insights into gnathostome evolution. *Nature* **505**, 174-179
(2014).

10. J. T. Bridgham, S. M. Carroll, J. W. Thornton, Evolution of hormone-receptor
complexity by molecular exploitation. *Science* **312**, 97-101 (2006).

11.   J. W. Funder, Aldosterone and mineralocorticoid receptors: a personal reflection.
Molecular and cellular endocrinology **350**, 146-150 (2012).

12.   U. A. Hawkins, E. P. Gomez-Sanchez, C. M. Gomez-Sanchez, C. E. Gomez-Sanchez,
The ubiquitous mineralocorticoid receptor: clinical implications. *Curr Hypertens Rep*
14, 573-580 (2012).

13.   L. Martinerie, M. Munier, D. Le Menuet, G. Meduri, S. Viengchareun, M. Lombes,
The mineralocorticoid signaling pathway throughout development: expression,
regulation and pathophysiological implications. *Biochimie* **95**, 148-157 (2013).

14.   F. Jaisser, N. Farman, Emerging Roles of the Mineralocorticoid Receptor in
Pathology: Toward New Paradigms in Clinical Pharmacology. *Pharmacological*
reviews **68**, 49-75 (2016).

15.   E. P. Gomez-Sanchez, Mineralocorticoid receptors in the brain and cardiovascular
regulation: minority rule? *Trends Endocrinol Metab* **22**, 179-187 (2011).

16.   J. J. DuPont, I. Z. Jaffe, 30 YEARS OF THE MINERALOCORTICOID RECEPTOR:
The role of the mineralocorticoid receptor in the vasculature. *J Endocrinol* **234**,
T67-T82 (2017).

17.   M. Joels, E. R. de Kloet, 30 YEARS OF THE MINERALOCORTICOID
RECEPTOR: The brain mineralocorticoid receptor: a saga in three episodes. *J*
Endocrinol **234**, T49-T66 (2017).

18.   T. J. Cole, M. J. Young, 30 YEARS OF THE MINERALOCORTICOID RECEPTOR:
Mineralocorticoid receptor null mice: informing cell-type-specific roles. *J Endocrinol*
234, T83-T92 (2017).

19.   P. Huang, V. Chandra, F. Rastinejad, Structural overview of the nuclear receptor
superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**, 247-272
(2010).

20.   F. Rastinejad, P. Huang, V. Chandra, S. Khorasanizadeh, Understanding nuclear
receptor form and function using structural biology. *Journal of molecular*
endocrinology **51**, T1-T21 (2013).

21.   A. Sugimoto, K. Oka, R. Sato, S. Adachi, M. E. Baker, Y. Katsu, Corticosteroid and
progesterone transactivation of mineralocorticoid receptors from Amur sturgeon and
tropical gar. *Biochem J* **473**, 3655-3665 (2016).

22.   J. Fagart, J. Huyet, G. M. Pinon, M. Rochel, C. Mayer, M. E. Rafestin-Oblin, Crystal
structure of a mutant mineralocorticoid receptor responsible for hypertension. *Nature*
structural & molecular biology **12**, 554-555 (2005).

23.   Y. Li, K. Suino, J. Daugherty, H. E. Xu, Structural and biochemical mechanisms for

the specificity of hormone binding and coactivator assembly by mineralocorticoid
receptor. *Mol Cell* **19**, 367-380 (2005).

24. R. K. Bledsoe, K. P. Madauss, J. A. Holt, C. J. Apolito, M. H. Lambert, K. H. Pearce,
T. B. Stanley, E. L. Stewart, R. P. Trump, T. M. Willson, S. P. Williams, A
ligand-mediated hydrogen bond network required for the activation of the
mineralocorticoid receptor. *The Journal of biological chemistry* **280**, 31283-31293
(2005).

25. R. Rupprecht, J. L. Arriza, D. Spengler, J. M. Reul, R. M. Evans, F. Holsboer, K.
Damm, Transactivation and synergistic properties of the mineralocorticoid receptor:
relationship to the glucocorticoid receptor. *Molecular endocrinology (Baltimore, Md)*
7, 597-603 (1993).

26. F. M. Rogerson, P. J. Fuller, Interdomain interactions in the mineralocorticoid receptor.
Molecular and cellular endocrinology **200**, 45-55 (2003).

27. R. Kumar, G. Litwack, Structural and functional relationships of the steroid hormone
receptors' N-terminal transactivation domain. *Steroids* **74**, 877-883 (2009).

28. J. B. Pippal, Y. Yao, F. M. Rogerson, P. J. Fuller, Structural and functional
characterization of the interdomain interaction in the mineralocorticoid receptor. *Mol*
Endocrinol **23**, 1360-1370 (2009).

29. K. Fischer, S. M. Kelly, K. Watt, N. C. Price, I. J. McEwan, Conformation of the
mineralocorticoid receptor N-terminal domain: evidence for induced and stable
structure. *Mol Endocrinol* **24**, 1935-1948 (2010).

30. N. Faresse, Post-translational modifications of the mineralocorticoid receptor: How to
dress the receptor according to the circumstances? *J Steroid Biochem Mol Biol* **143**,
334-342 (2014).

31. P. J. Fuller, Novel interactions of the mineralocorticoid receptor. *Molecular and*
cellular endocrinology **408**, 33-37 (2015).

32. Y. Katsu, K. Oka, M. E. Baker, Evolution of steroid specificity in human, chicken,
alligator, frog and zebrafish mineralocorticoid receptors: Allosteric interactions affect
steroid specificity. *bioRxiv*,   (2018).

33. J. B. Pippal, C. M. Cheung, Y. Z. Yao, F. E. Brennan, P. J. Fuller, Characterization of
the zebrafish (Danio rerio) mineralocorticoid receptor. *Molecular and cellular*
endocrinology **332**, 58-66 (2011).

34. J. M. P. Joss, D. E. Arnoldreed, R. J. Balment, The Steroidogenic Response to
Angiotensin-Ii in the Australian Lungfish, Neoceratodus-Forsteri. *J Comp Physiol B*
164, 378-382 (1994).

35.   C. T. Amemiya, J. Alfoldi, A. P. Lee, S. Fan, H. Philippe, I. Maccallum, I. Braasch, T.
Manousaki, I. Schneider, N. Rohner, C. Organ, D. Chalopin, J. J. Smith, M. Robinson,
R. A. Dorrington, M. Gerdol, B. Aken, M. A. Biscotti, M. Barucca, D. Baurain, A. M.
Berlin, G. L. Blatch, F. Buonocore, T. Burmester, M. S. Campbell, A. Canapa, J. P.
Cannon, A. Christoffels, G. De Moro, A. L. Edkins, L. Fan, A. M. Fausto, N. Feiner,
M. Forconi, J. Gamieldien, S. Gnerre, A. Gnirke, J. V. Goldstone, W. Haerty, M. E.
Hahn, U. Hesse, S. Hoffmann, J. Johnson, S. I. Karchner, S. Kuraku, M. Lara, J. Z.
Levin, G. W. Litman, E. Mauceli, T. Miyake, M. G. Mueller, D. R. Nelson, A. Nitsche,
E. Olmo, T. Ota, A. Pallavicini, S. Panji, B. Picone, C. P. Ponting, S. J. Prohaska, D.
Przybylski, N. R. Saha, V. Ravi, F. J. Ribeiro, T. Sauka-Spengler, G. Scapigliati, S. M.
Searle, T. Sharpe, O. Simakov, P. F. Stadler, J. J. Stegeman, K. Sumiyama, D. Tabbaa,
H. Tafer, J. Turner-Maier, P. van Heusden, S. White, L. Williams, M. Yandell, H.
Brinkmann, J. N. Volff, C. J. Tabin, N. Shubin, M. Schartl, D. B. Jaffe, J. H.
Postlethwait, B. Venkatesh, F. Di Palma, E. S. Lander, A. Meyer, K. Lindblad-Toh,
The African coelacanth genome provides insights into tetrapod evolution. *Nature* **496**,
311-316 (2013).
36.   M. E. Baker, Evolution of glucocorticoid and mineralocorticoid responses: go fish.
Endocrinology **144**, 4223-4225 (2003).
37.   M. E. Baker, C. Chandsawangbhuwana, N. Ollikainen, Structural analysis of the
evolution of steroid specificity in the mineralocorticoid and glucocorticoid receptors.
BMC Evol Biol **7**, 24 (2007).
38.   N. R. Bury, A. Sturm, Evolution of the corticosteroid receptor signalling pathway in
fish. *Gen Comp Endocrinol* **153**, 47-56 (2007).
39.   T. Sakamoto, C. Mori, S. Minami, H. Takahashi, T. Abe, D. Ojima, M. Ogoshi, H.
Sakamoto, Corticosteroids stimulate the amphibious behavior in mudskipper:
potential role of mineralocorticoid receptors in teleost fish. *Physiol Behav* **104**,
923-928 (2011).
40.   A. Sturm, N. Bury, L. Dengreville, J. Fagart, G. Flouriot, M. E. Rafestin-Oblin, P.
Prunet, 11-deoxycorticosterone is a potent agonist of the rainbow trout
(Oncorhynchus mykiss) mineralocorticoid receptor. *Endocrinology* **146**, 47-55
(2005).
41.   H. Takahashi, T. Sakamoto, The role of 'mineralocorticoids' in teleost fish: relative
importance of glucocorticoid signaling in the osmoregulation and 'central' actions of
mineralocorticoid receptor. *Gen Comp Endocrinol* **181**, 223-228 (2013).
42.   Y. Katsu, M. E. Baker, Progesterone activation of zebrafish mineralocorticoid receptor

may influence growth of some transplanted tumors. *Proc Natl Acad Sci U S A* **115**, E2908-E2909 (2018).

43.   D. S. Geller, A. Farhi, N. Pinkerton, M. Fradley, M. Moritz, A. Spitzer, G. Meinke, F. T. Tsai, P. B. Sigler, R. P. Lifton, Activating mineralocorticoid receptor mutation in hypertension exacerbated by pregnancy. *Science* **289**, 119-123 (2000).

44.   S. M. Carroll, J. T. Bridgham, J. W. Thornton, Evolution of hormone signaling in elasmobranchs by exploitation of promiscuous receptors. *Molecular biology and evolution* **25**, 2643-2652 (2008).

45.   P. J. Fuller, Y. Yao, J. Yang, M. J. Young, Mechanisms of ligand specificity of the mineralocorticoid receptor. *J Endocrinol* **213**, 15-24 (2012).

46.   D. A. Close, S. S. Yun, S. D. McCormick, A. J. Wildbill, W. Li, 11-deoxycortisol is a corticosteroid hormone in the lamprey. *Proc Natl Acad Sci U S A* **107**, 13942-13947 (2010).

47.   Y. Li, A. Sturm, P. Cunningham, N. R. Bury, Evidence for a divergence in function between two glucocorticoid receptors from a basal teleost. *BMC Evol Biol* **12**, 137 (2012).

48.   A. Sturm, L. Colliar, M. J. Leaver, N. R. Bury, Molecular determinants of hormone sensitivity in rainbow trout glucocorticoid receptors 1 and 2. *Molecular and cellular endocrinology* **333**, 181-189 (2011).

49.   A. S. Arterbery, D. J. Fergus, E. A. Fogarty, J. Mayberry, D. L. Deitcher, W. Lee Kraus, A. H. Bass, Evolution of ligand specificity in vertebrate corticosteroid receptors. *BMC Evol Biol* **11**, 14 (2011).

50.   N. R. Bury, The evolution, structure and function of the ray finned fish (Actinopterygii) glucocorticoid receptors. *Gen Comp Endocrinol*,   (2016).

51.   R. Rupprecht, J. M. Reul, B. van Steensel, D. Spengler, M. Soder, B. Berning, F. Holsboer, K. Damm, Pharmacological and functional characterization of human mineralocorticoid and glucocorticoid receptor ligands. *Eur J Pharmacol* **247**, 145-154 (1993).

52.   C. Hellal-Levy, B. Couette, J. Fagart, A. Souque, C. Gomez-Sanchez, M. Rafestin-Oblin, Specific hydroxylations determine selective corticosteroid recognition by human glucocorticoid and mineralocorticoid receptors. *FEBS Lett* **464**, 9-13 (1999).

53.   O. Mani, L. G. Nashev, C. Livelo, M. E. Baker, A. Odermatt, Role of Pro-637 and Gln-642 in human glucocorticoid receptors and Ser-843 and Leu-848 in mineralocorticoid receptors in their differential responses to cortisol and aldosterone.

J Steroid Biochem Mol Biol **159**, 31-40 (2016).

54.   C. M. Kagawa, Blocking urinary electrolyte effects of desoxycorticosterone with

progesterone in rats. *Proc Soc Exp Biol Med* **99**, 705-707 (1958).

55.   G. Wambach, J. R. Higgins, Antimineralocorticoid action of progesterone in the rat:

correlation of the effect on electrolyte excretion and interaction with renal

mineralocorticoid receptors. *Endocrinology* **102**, 1686-1693 (1978).

56.   J. W. Funder, W. R. Adam, 19-Nor progesterone is a mineralocorticoid agonist.

Endocrinology **109**, 313-315 (1981).

57.   C. E. Hall, C. E. Gomez-Sanchez, S. Hungerford, E. P. Gomez-Sanchez,

Mineralocorticoid and hypertensive effects of 19-nor-progesterone. *Endocrinology*

109, 1168-1175 (1981).

58.   B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

59.   L. Fagerberg, B. M. Hallstrom, P. Oksvold, C. Kampf, D. Djureinovic, J. Odeberg, M.

Habuka, S. Tahmasebpoor, A. Danielsson, K. Edlund, A. Asplund, E. Sjostedt, E.

Lundberg, C. A. Szigyarto, M. Skogs, J. O. Takanen, H. Berling, H. Tegel, J. Mulder,

P. Nilsson, J. M. Schwenk, C. Lindskog, F. Danielsson, A. Mardinoglu, A. Sivertsson,

K. von Feilitzen, M. Forsberg, M. Zwahlen, I. Olsson, S. Navani, M. Huss, J. Nielsen,

F. Ponten, M. Uhlen, Analysis of the human tissue-specific expression by

genome-wide integration of transcriptomics and antibody-based proteomics. *Mol Cell*

Proteomics **13**, 397-406 (2014).

60.   A. H. Payne, D. B. Hales, Overview of steroidogenic enzymes in the pathway from

cholesterol to active steroid hormones. *Endocr Rev* **25**, 947-970 (2004).

61.   K. N. Wynne, J. Mercer, J. R. Stockigt, J. W. Funder, 19-nor analogs of adrenal

steroids: mineralocorticoid and glucocorticoid receptor activity. *Endocrinology* **107**,

1278-1280 (1980).

62.   S. Aquila, F. De Amicis, Steroid receptors and their ligands: effects on male gamete

functions. *Exp Cell Res* **328**, 303-313 (2014).

63.   O. M. Conneely, B. Mulac-Jericevic, F. DeMayo, J. P. Lydon, B. W. O'Malley,

Reproductive functions of progesterone receptors. *Recent Prog Horm Res* **57**, 339-355

(2002).

64.   E. R. de Kloet, C. Otte, R. Kumsta, L. Kok, M. H. Hillegers, H. Hasselmann, D.

Kliegel, M. Joels, Stress and Depression: a Crucial Role of the Mineralocorticoid

Receptor. *J Neuroendocrinol* **28**, (2016).

65.   J. W. Funder, Aldosterone and mineralocorticoid receptors in the cardiovascular

603   system. *Prog Cardiovasc Dis* **52**, 393-400 (2010).

604 66. M. J. Young, A. J. Rickard, Mineralocorticoid receptors in the heart: lessons from

605   cell-selective transgenic animals. *J Endocrinol* **224**, R1-13 (2015).

606 67. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat*

607   *Methods* **9**, 357-359 (2012).

608 68. K. Oka, S. Kohno, H. Urushitani, L. J. Guillette, Jr., Y. Ohta, T. Iguchi, Y. Katsu,

609   Molecular cloning and characterization of the corticoid receptors from the American

610   alligator. *Molecular and cellular endocrinology* **365**, 153-161 (2013).

611 69. K. Oka, A. Hoang, D. Okada, T. Iguchi, M. E. Baker, Y. Katsu, Allosteric role of the

612   amino-terminal A/B domain on corticosteroid transactivation of gar and human

613   glucocorticoid receptors. *J Steroid Biochem Mol Biol* **154**, 112-119 (2015).