Epidemiological Study of Autism Subgroups Using Autism Treatment Evaluation Checklist (ATEC) Score

Shreyas Mahapatra[1], Edward Khokhlovich, Samantha Martinez[1], Benjamin Kannel, Stephen M Edelson[2], Andrey Vyshedskiy[1,3*]

[1]Boston University, Boston, USA
[2]Autism Research Institute, San Diego, CA, USA
[3]ImagiRation LLC, Boston, MA, USA

Keywords: autism, ASD, psychological evaluations, ATEC, Autism Treatment Evaluation Checklist, ATEC norms

*Corresponding author: Andrey Vyshedskiy, Ph.D., Boston University, Boston, USA, Tel: +1 (617) 433-7724; E-mail: vysha@bu.edu

Abstract

Here we report the results of the subgroup analyses of the observational cohort of children whose parents completed the Autism Treatment Evaluation Checklist (ATEC) over the period of several years. A linear mixed effects model was used to evaluate longitudinal changes in ATEC scores within different patient subgroups. All groups decreased their mean ATEC score over time indicating improvement of symptoms, however there were significant differences between the groups. Younger children improved more than the older children. There was no difference in improvement between females vs. males. The model showed a trend towards significance for different autism severity groups. One surprising finding was that children from developed English-speaking countries improved less than children from the rest of the world.

Introduction

The regular assessment of temporal change in symptoms of children with Autism Spectrum Disorder (ASD) participating in a clinical trial has been a long-standing challenge. A common hurdle in these efforts is the availability of trained technicians needed to conduct rigorous and consistent assessment of children at multiple time points. If parents could administer regular psychometric evaluations of their children, then the cost of clinical trials will be reduced, enabling longer clinical trials with the larger number of participants. The Autism Treatment Evaluation Checklist (ATEC) was developed to provide a free and easily accessible method for caregivers to track the changes of ASD symptoms over time (Rimland & Edelson, 1999). Importantly, ATEC was not designed for diagnostic purposes, but only to measure changes in ASD severity, making it useful in monitoring behaviors over time as well as tracking the efficacy of a treatment. Various studies have sought to confirm the validity of ATEC (Al Backer, 2016; Geier, Kern, & Geier, 2013; Jarusiewicz, 2002), yet none to date have assessed longitudinal changes in participants' ATEC scores with respect to age, sex, and ASD severity. This study analyzed data reported by participants using the ATEC over a four-year time span (2013 to 2017). Assessing these data permits insights into the effects of age, sex, country of origin, and ASD severity on the longitudinal changes in ATEC score with all of these factors (save for sex) showing statistically significant differences affecting ATEC score dynamics. These findings identify specific variables capable of altering the developmental trajectory of children with ASD and indicate possible avenues of future investigation of causal relationships related to changes in ASD severity.

ATEC is comprised of four subscales: 1) Speech/Language/Communication, 2) Sociability, 3) Sensory/Cognitive Awareness, and 4) Health/Physical/Behavior. These four subscales are used to calculate the total score that ranges from 0 to 179. A lower score indicates less severe symptoms of ASD

and a higher score correlates with more severe symptoms of ASD (Rimland & Edelson, 1999). The subscales provide survey takers with the information about specific areas of behaviors which may change over time.

Previous studies have aimed to evaluate the validity and reliability of the ATEC in tracking participant's ASD severity. One trial conducted by Magiati et al., aimed to comprehensively assess ATEC's ability to longitudinally measure changes in participant performance (Magiati, Moss, Yates, Charman, & Howlin, 2011). This study utilized ATEC to monitor the progress of 22 schoolchildren over a five-year period. ATEC score was compared to age-specific cognitive, language, and behavioral metrics such as the Wechsler Preschool and Primary Scale of Intelligence. The researchers noted ATEC's high level of internal consistency as well as a highly correlative relationship with other standardized assessments used to measure the same capacities in children with ASD (Magiati et al., 2011). Charman et al. utilized ATEC amongst other measures to test the feasibility of tracking the longitudinal changes in children using caregiver-administered questionnaires and noted differential effects across subscales of ATEC, possibly driven by development-focused vs. symptom-focused subscales that are conflated in the ATEC total score (Charman, Howlin, Berry, & Prince, 2004). Another study assessing the ability of dietary intervention to affect ASD symptoms also utilized ATEC as a primary measure (Klaveness, Bigam, & Reichelt, 2013), concluding that it has "high general reliability" coupled with an ease of access. Whitehouse et al. used ATEC as a primary outcome measure for a randomized controlled trial of their iPad-based intervention for ASD named TOBY (Whitehouse et al., 2017). This trial was conducted over a 6-month time frame, with outcome assessments at the 3-month and 6-month time points. Although the study did not demonstrate significant ATEC score differences amongst test groups, the researchers reaffirmed their use of ATEC, noting its "internal consistency and adequate predictive validity" (Whitehouse et al., 2017). These studies support the viability of ATEC as a tool for longitudinal

measurement of ASD severity which can be a vital instrument in tracking symptom changes during a trial.

The current observational study was initiated nearly two decades ago when one of the authors (Stephen M. Edelson of Autism Research Institute) distributed ATEC questionnaire to parents of children with ASD. Initially, ATEC evaluations were distributed as hard copy. In 2013 the online version of ATEC was developed. The participant responses to the online version of ATEC are presented in this manuscript.

Methods

ATEC Evaluation Structure

The ATEC is a caregiver-administered questionnaire designed to measure changes in severity of ASD in response to treatment. A total score and four subscale scores are reported. Questions in the first three subscales are scored using a 0-2 scale. The fourth subscale, Health/Physical/Behavior, is scored using a 0–3 point scale. ATEC can be accessed online or in hard-copy format.

The first subscale, Speech/Language/Communication, contains 14 items and its score ranges from 0 to 28 points. The Sociability subscale contains 20 items within 0 to 40 score range. The third subscale, Sensory/Cognitive awareness, has 18 items and scores range from 0 to 36. Finally, the Health/Physical/Behavior subscale contains 25 items. The scores from each subscale are combined in order to calculate a Total Score, which ranges from 0 to 179 points. A lower score indicates a lower severity of ASD symptoms.

Collection of Evaluations

ATEC responses originated from participants voluntarily and freely completing online ATEC evaluations from 2013 to 2017.

Evaluations of ATEC score changes over time

In order to study how ATEC scores change overtime and whether those changes vary within different ASD subgroups, we first introduced a concept of a "Visit" by dividing the two-year-long observation interval into 3-month periods. All evaluations were mapped into 3-month-long bins with the first evaluation placed in the first bin. When more than one evaluation was completed within a bin, their results were averaged to calculate a single number representing this 3-month interval. We then hypothesized that there is an interaction between a Visit and a given subgroup category (age, sex, ASD severity, and country of origin). Statistically, we modeled that by applying Linear Mixed Effect (LME) model with repeated measures, where we introduced an interaction term testing the hypothesis, which, in turn, enabled generation of pairwise differences between modeled subgroups at different visits. Participant specific variability was accounted for by introducing random effect into the model.

Pairwise differences were computed by applying "LS Means " and "contrast" functions to a generated LME model. For each ATEC score and a given subgroup, the following output is generated:

1.   General ANOVA summary of the model itself, including p-values for each covariate and the interaction term among them

2.   LS Means computed for a given category at each visit (with 95% Confidence Interval)

3.   Pairwise differences between categories at different visits with p-values adjusted for multiple comparisons testing using Tukey method.

Participants

Participants were selected based on the following criteria:

1) Completeness: Participants who did not provide a DOB were excluded. As participants' dates of birth (DOB) were utilized to determine age, the availability of DOB was a vital factor.

2) Consistency: Participants must have filled out at least three questionnaires within 2 years and the interval between the first and the last evaluation was one year or longer.

3) Maximum age: Participants older than twelve years of age were excluded from this study.

4) Minimal ATEC severity: Participants with initial ATEC scores of less than 20 were excluded.

After excluding participants that did not meet these criteria, there were 2331 total participants.

Age groups

Participants were grouped based on age, calculated from the date of birth at the time of the first completed evaluation. The three age groups were: 1.2-3 years of age (YOA), 3.1-6 YOA, 6.1-12 YOA, Table 1.

Autism Severity Measurements

The initial ATEC total score was used as proxy for ASD severity. Participants were organized into three groups: mild (initial ATEC total score 20-49), moderate (initial ATEC total score 50-79), and severe (initial ATEC total score > 80), Table 2.

Country groups

Participants were split into two groups based their country. The developed English-speaking nations included participants from the United States, Canada, United Kingdom, Ireland, Australia, and New Zealand. Participants from other countries were grouped together as "the rest of the world" (Table 3).

Sex groups

83% of the 2331 participants were males (Table 4).

Results

In order to evaluate temporal dynamics of ATEC scores and their distribution within different subgroups, the concept of a "Visit" was developed by dividing the two-year-long observation interval into 3-month periods. All completed evaluations were accordingly mapped into 3-month-long bins, with the first evaluation placed in the first bin. In instances where more than one evaluation was completed by a participant within a bin, these evaluations were averaged to calculate a single value for this 3-month interval.

The least squared means (LS Means), as well as the dynamics of score changes (LS Means differences) over time (between visits) for each group are presented in the supplementary materials. The fitting of LME model allowed us not only to assess the temporal dynamics of the scores, but also to evaluate the "tightness" of each individual mean value by generating 95% confidence interval. We have observed high degree of data consistency, similar to what was reported by Magiati et. al. The differences between participant subgroups at different visits, and differences between the first and the last visit per subgroup are discussed in greater detail below.

Effect of sex on longitudinal change of ATEC scores

The interaction term between Sex group and Visits was not statistically significant (at the $\alpha = 0.05$ level of significance) for either the ATEC total score or any of subscale scores (Table 5).

Longitudinal change of ATEC scores as a function of age

The significance of interaction term between Age group and Visits shows that the dynamics of ATEC total score as well as scores in the Communication, Sociability, and Sensory subscales varies within different Age groups (Table 6). Table 7 shows LS Means for age groups at the initial and the last visits. Reduction in Total ATEC score (showing the degree of improvement) was inversely related to age (Table 7). Over the two years 1-3 YOA group improved by 28.18 units (SE=1.14, $p < .0001$), 3-6 YOA group improved by 18.69 units (SE=0.76, $p < .0001$), and 6-12 YOA group improved by 13.07 units (SE=0.98, $p < .0001$).

For age group comparisons, three pairwise comparisons were made, Table 8. Neither pairwise comparison reached statistical significance at Visit 1, but all three pairwise comparisons in Total ATEC score yielded statistically significant differences in LS Means at Visit 8 with younger children improving more than the older children (Table 8). The difference in Total ATEC score for the 1-3 YOA group relative to the 3-6 YOA group was 2.11 units (SE=0.77, $p = .49$) at Visit 1 and -7.38 units (SE=1.33, $p < .0001$) at Visit 8. Total ATEC score for the 1-3 YOA group relative to the 6-12 YOA group was 2.41 units (SE=0.87, $p = .47$) at Visit 1 and -12.71 units (SE=1.45, $p < .0001$) at Visit 8. Total ATEC score for the 3-6 YOA group relative to the 6-12 YOA group was 0.29 units (SE=0.72, $p = 1.00$) at Visit 1 and -5.33 units (SE=1.19, $p < .002$) at Visit 8. These observations were recapitulated in the Communication subscale (Table 8). For the Sociability and Sensory subscales, only the 1-3 YOA group vs. 3-6 YOA group and 1-3 YOA group vs. 6-12 YOA group yielded a statistically significant decrease in score at Visit 8 (Table 8).

The interaction term between Age Group and Visits was not statistically significant for the Physical subscale and thus age-specific means were not calculated.

Country effects on ATEC scores

Surprisingly, a comparison of developed English-speaking nations (the United States, Canada, United Kingdom, Ireland, Australia, and New Zealand) to the rest of the world demonstrated greater improvements in ATEC total score and the Sociability subscale in the non-English-speaking nations. The significance of interaction term between Country group and Visits shows that the dynamics of ATEC total score and the Communication, Sociability, and Physical subscales varies in different Country groups (Table 9). Table 10 shows LS Means for Country groups at the initial and the last visits. Reduction in Total ATEC score was greater in non-English-speaking nations group (Table 10). Over the period of two years the participants in the English-speaking nations group improved by 16.45 units (SE=0.84, p < 0.0001), and non-English-speaking nations group improved by 21.01 units (SE=0.70, p < 0.0001).

The difference in ATEC total score for the English-Speaking Counties group relative to the rest of the world was 1.03 units (SE=0.61, p = 0.95) at Visit 1 and -3.50 units (SE=1.05, p = 0.0641) at Visit 8, Table 11. This trend toward statistical significance at Visit 8 indicates that children in the English-speaking countries improve their symptoms to a smaller degree than children in the non-English speaking nations. Dissection of the Total ATEC score into subscales indicates that children in the non-English speaking nations manifested greater improvements of their symptoms in all subscales with the Sociability subscale having the greatest contribution to the difference between the groups. The difference in the Sociability subscale score for the English-Speaking Counties group relative to the rest of the world was 0.23 units (SE=0.19, p = 0.9975) at Visit 1 and -1.25 units (SE=0.33, p = 0.0145) at Visit 8, Table 11.

Change of ATEC scores as a function of ASD severity

The significance of interaction term between ASD severity group and Visits shows that the dynamics of ATEC total score and of individual scores within all subscales differs between severity

groups (Table 12). Table 13 shows the LS Mean calculations for the three severity groups (mild, moderate, severe) at Visit 1 and Visit 8. Reduction in Total ATEC score was directly related to severity (Table 13). Over the two years, the mild group improved by 9.20 units (SE=0.94, p < 0.0001), the moderate group by 20.124 units (SE=0.76, p < 0.0001), and the severe group by 29.55 units (SE=1.09, p < 0.0001).

In comparing the difference in LS Mean for ATEC total score at Visit 1, all three pairwise comparisons between severity groups yielded statistically significant differences (Table 14). This is in contrast to Visit 8, at which point none of the comparisons reached statistical significance (Table 14). The results for all four subscales mirrored those of ATEC total score, showing no statistically significant differences between severity groups at Visit 8 (Table 14). This may simply be an artifact of the definition of ASD severity, which is based solely on a child's initial Total ATEC score independent of child's age. Therefore, we added a different approach to definition of ASD severity groups.

According to ATEC norms, ATEC total scores in young children decrease exponentially with age, with a time constant of approximately 3.3 years regardless of the initial ATEC total score (Mahapatra et al., 2018). Thus, participants could be divided into three approximately equal groups using exponents decaying with a time constant of 3.3 years. The exact parameters of exponents were determined using best-fit trendlines to ATEC norms (Mahapatra et al., 2018), Table 15.

Group differences were reassessed using the severity group definition based on both the initial ATEC total score and age as specified in Table 15. The significance of interaction term between ASD severity group and Visits shows that the dynamics of ATEC total score and individual subscale scores varies between different severity groups (Table 16). Table 17 shows the LS Mean calculations for the three severity groups at Visit 1 and Visit 8. Reduction in Total ATEC score was directly related to severity (Table 17). Over the two years the mild group improved by 16.72 units (SE=0.90, p < 0.0001), the

moderate group improved by 21.85 units (SE=0.90, p < 0.0001), and the severe group improved by 20.49 units (SE=0.90, p < .0001).

In comparing the difference in LS Mean for ATEC total score at Visit 1, all three pairwise comparisons between severity groups yielded statistically significant differences (Table 18). This is in contrast to Visit 8, at which point none of the comparisons reached statistical significance, although the difference between moderate and severe groups showed a trend toward significance (p=0.0786) (Table 18). For the Communication subscale all pairwise group differences were statistically significant, confirming the advantage of severity group assignment based on both initial ATEC total score and age (Table 18). For the Sociability subscale only the moderate group vs. severe group yielded a statistically significant difference in score at Visit 8 (Table 18). There were no statistically significant differences between severity groups at Visit 8 in the Sensory subscale or the Physical subscale.

Discussion

Design considerations for an ASD early-intervention clinical trial must take into account (1) the trial duration, (2) number of participants, and (3) the quality of participant assessment. A short clinical trial of an early therapeutic intervention in 2 to 3 year old children can easily miss a target, as an improvement of symptoms may not emerge until children reach the school age. Small numbers of participants can easily skew the data as ASD is known to be a highly heterogeneous disorder. Longer clinical trials, with the greater number of participants, provide a better test for any intervention. Increasing the trial duration and the number of trial participants, however, raises the demand for regular assessment of participants by trained psychometric technicians. Furthermore, to attain the larger number of trial participants, clinical trials must accept participants across a large geographical region. The logistical issues associated with such an endeavor come at immense cost. As a result, large numbers of ASD clinical trials working under a limited budget suffer from short duration and low

participant number, often compromising the trial objectives (e.g., (Drew et al., 2002; Whitehouse et al., 2017).

ATEC was in part designed to circumvent these problems. If caregivers could serve as psychometric technicians and conduct regular evaluations of their children, the cost of clinical trials will be substantially reduced while simultaneously allowing for longer trial duration. There is an understanding in the psychological community that parents cannot be trusted with an evaluation of their own children. In fact, parents often yield to wishful thinking and overestimate their children's abilities on a single assessment. However, the pattern of changes can be generated by measuring the score dynamics over multiple assessments. When a single parent completes the same evaluation every three months over multiple years, changes in the score become meaningful. This study attempts to characterize the typical changes in ATEC score over time as a function of children age, sex, ASD severity, and country of origin in a large and diverse group of participants.

Sex does not affect ATEC Score

The prevalence of ASD is strongly male-biased, affecting 4 times as many males as females. Accordingly, we were interested in differences in the rate of improvement between participants of different sexes. No significant differences in improvement of ATEC total score were observed. This suggests that although for reasons not fully understood there is a greater incidence of ASD in males, the rate of improvement of ASD symptoms remains similar in males and females.

Effect of Age on ATEC Score

The participants' age was a significant modulating factor in determining the rate of their improvement. Younger children demonstrated greater improvement in Total ATEC score. This phenomenon was recapitulated across subscales, with differences between the 1-3 YOA group and 3-6 YOA group as well as between the 1-3 YOA group and 6-12 YOA group reaching statistical significance

for the Communication, Sociability, and Sensory subscales (Table 8). This finding is consistent with other ATEC longitudinal studies: in all of them the younger children showed greater improvement in Total ATEC score compared to the older children (compare Magiati et al. to Charman et. al. and Charman et al. to Whitehouse et al./TOBY, Table 19).

The magnitude of the annual decrease of the ATEC score was also found to be roughly similar to other reports across the studied age range. For the younger children the reduction of ATEC score seen in this study is in between those of Whitehouse et al./TOBY trial and Charman et al, Table 19. For the older children, the reduction of ATEC seen in this study is somewhat similar to that reported by Charman et al., Table 19.

The small differences between the studies can be attributed to differences in study design. In particular, the current study (1) had significantly more participants, (2) was based on greater number of ATEC evaluations, and (3) was conducted over the longer period of time than all the others discussed herein.

Effect of ASD Severity on ATEC Score

In comparing the difference in LS Mean for ATEC total score at Visit 1, all three pairwise comparisons between severity groups yielded statistically significant differences (Table 14). This is in contrast to Visit 8, at which point none of the comparisons reached statistical significance (Table 14). The results for all four subscales mirrored those of ATEC total score, showing no statistically significant differences between severity groups at Visit 8 (Table 14). This may simply be an artifact of the definition of ASD severity, which is based on a child's initial Total ATEC score. This method groups children with the same initial Total ATEC score together independent of age. Thus, children who score 80 on their initial evaluation at the age 10 are grouped together with children who score 80 on their initial evaluation at the age 2. According to ATEC norms (Mahapatra et al., 2018), these children will score 70 and 25 respectively at the age of 12, and therefore clearly belong to different severity groups. This

inconsistency in definition of ASD severity solely based on the initial Total ATEC score independent of age may explain the observation that none of the group comparisons reached statistical significance at Visit 8.

The definition of ASD severity groups based on two parameters – the initial Total ATEC score and age – yielded somewhat superior results compared to defining ASD severity based solely on the initial Total ATEC score. While both definition methods showed no statistically significant differences between severity groups at Visit 8 in ATEC total score (Tables 14 and 18), the former method showed statistically significant pairwise differences between all the groups at Visit 8 for the Communication subscale .

Role of Country of Origin

Conventional wisdom may suggest that the increased access to resources, including government-provided therapy for ASD, should lead to greater improvements. English-speaking nations (the United States, Canada, the United Kingdom, Ireland, Australia, and New Zealand) lead the world in government spending on therapy for children with ASD and therefore would be expected to produce superior outcomes of ASD therapy. Surprisingly, a comparison of English-speaking nations to the rest of the world demonstrated greater improvements in Total ATEC score as well as in each subscale within the non-English speaking nations (Table 11).

This observation runs contrary to conventional thought and underscores the consensus that there is a potential for improving the treatment of children with ASD in the developed world. While it is difficult to speculate on the reason for this disparity between developed English-speaking countries and the rest of the world, we note that child treatment is more often outsourced in the developed countries compared to more traditional societies where mother is more likely to stay at home and personally take care of a sick child.

Limitations

Participant selection presents a novel challenge in a study focused on caregiver-administered assessments. In the selection of participants for inclusion in this study, a baseline of ASD diagnosis could not have been established as child's diagnosis is not part of ATEC questionnaire. Thus, it is not impossible that some of the participants did not have ASD diagnosis altogether. E.g., parents of a neurotypical toddler worried for any reason about an ASD diagnosis could have decided to monitor toddler's development with ATEC evaluations and thus inadvertently added their normally developing child to the ATEC collection. As neurotypical children develop faster, the presence of neurotypical children in the dataset would have artificially increased the magnitude of annual changes of ATEC scores, predominantly for younger participants.

We note though that it is unlikely that there were many neurotypical participants in our database. First, ATEC is virtually unknown outside the autism community. Second, there is a little incentive for the parents of neurotypical children to complete multiple exhaustive ATEC questionnaires (unless one of the children was previously diagnosed with ASD). Third, to further limit the contribution from neurotypical children, we excluded participants with an initial ATEC total score of 20 or less, that may have represented the neurotypical population (7% of all participants). Despite this effort, the reported data may over-approximate the magnitude of annual changes of ATEC scores, especially in the younger participants.

As noted by other groups (Whitehouse et al., 2017; Charman et al., 2004), the use of ATEC as a primary outcome measure has some inherent drawbacks. While the ATEC is capable of delineating incremental differences in ASD severity amongst participants, the variety of measures amongst its subscales fails to differentiate developmental-specific changes from symptom-specific ones. This aspect of the ATEC may introduce a confounding variable when participants are at different developmental stages and follow unique developmental trajectories during a study. To mitigate these effects, trial

designs must accurately separate participants based on developmental stage. This is most often accomplished by using age as a proxy for developmental stage.

Conclusions

This manuscript attempts to characterize the typical changes in ATEC score over time as a function of children age, sex, ASD severity, and country of origin in a large and diverse group of participants. In doing so, it lends support to the efficacy of caregiver-driven psychometric observation, which when applied at scale, may be a viable alternative to using licensed technicians to assess the children.

Acknowledgments

We wish to thank Dr. Petr Ilyinskii for productive discussion and scrupulous editing of this manuscript.

Compliance with Ethical Standards

Using the Department of Health and Human Services regulations found at 45 CFR 46.101(b)(4), the Chesapeake Institutional Review Board (IRB) determined that this research project is exempt from IRB oversight.

Author contributions

Andrey Vyshedskiy, Edward Khokhlovich and Stephen M. Edelson designed the study. Stephen M. Edelson acquired the data. Edward Khokhlovich, Samantha Martinez, Benjamin Kannel, and Andrey

Vyshedskiy analyzed the data. Shreyas Mahapatra, Edward Khokhlovich, Stephen M. Edelson, and Andrey Vyshedskiy wrote the paper.

Conflicts of Interest

The authors declare no conflict of interest.

References

Al Backer, N. B. (2016). Correlation between Autism Treatment Evaluation Checklist (ATEC) and Childhood Autism Rating Scale (CARS) in the evaluation of autism spectrum disorder. *Sudanese Journal of Paediatrics*, *16*(1), 17.

Charman, T., Howlin, P., Berry, B., & Prince, E. (2004). Measuring developmental progress of children with autism spectrum disorder on school entry using parent report. *Autism*, *8*(1), 89–100.

Drew, A., Baird, G., Baron-Cohen, S., Cox, A., Slonims, V., Wheelwright, S., … Charman, T. (2002). A pilot randomised control trial of a parent training intervention for pre-school children with autism. *European Child & Adolescent Psychiatry*, *11*(6), 266–272.

Geier, D. A., Kern, J. K., & Geier, M. R. (2013). A comparison of the Autism Treatment Evaluation Checklist (ATEC) and the Childhood Autism Rating Scale (CARS) for the quantitative evaluation of autism. *Journal of Mental Health Research in Intellectual Disabilities*, *6*(4), 255–267.

Jarusiewicz, B. (2002). Efficacy of Neurofeedback for Children in the Autistic Spectrum: A Pilot Study. *Journal of Neurotherapy*, *6*(4), 39–49. https://doi.org/10.1300/J184v06n04_05

Klaveness, J., Bigam, J., & Reichelt, K. L. (2013). The varied rate of response to dietary intervention in autistic children. *Open Journal of Psychiatry*, *3*(02), 56.

Magiati, I., Moss, J., Yates, R., Charman, T., & Howlin, P. (2011). Is the Autism Treatment Evaluation

      Checklist a useful tool for monitoring progress in children with autism spectrum disorders?

      Journal of Intellectual Disability Research, *55*(3), 302–312. https://doi.org/10.1111/j.1365-

      2788.2010.01359.x

Mahapatra, S., Vyshedksy, D., Martinez, S., Kannel, B., Braverman, J., Edelson, S. M., & Vyshedskiy,

      Andrey. (2018). Autism Treatment Evaluation Checklist (ATEC) norms: a "growth chart" for ATEC

      score changes as a function of age. *Children, Accepted for publication*. https://doi.org/10.3390

Rimland, B., & Edelson, S. M. (1999). Autism Research Institute. *Autism Treatment Evaluation Checklist*

      (ATEC).

Whitehouse, A. J., Granich, J., Alvares, G., Busacca, M., Cooper, M. N., Dass, A., … others. (2017). A

      randomised controlled trial of an iPad-based application to complement early behavioural

      intervention in Autism Spectrum Disorder. *Journal of Child Psychology and Psychiatry*. Retrieved

      from http://onlinelibrary.wiley.com/doi/10.1111/jcpp.12752/full

Table 1

Characteristics and baseline measures for age groups.

Age (abbreviation used in the paper)	Participants in each age group (total)	Participants in each age group (%)	Age at baseline (mean ± SD)	Initial ATEC Total score (mean ± SD)
1.2-3 YOA (1-3)	516	22%	2.44±0.44	71±22
3.1-6 YOA (3-6)	1176	50%	4.33±0.85	62±23
6.1-12 YOA (6-12)	639	28%	8.20±1.62	62±24

Table 2

Characteristics and baseline measures for ASD severity groups

Autism Severity	Initial ATEC Total Score	Participants in each severity group (total)	Participants in each severity group (%)	Age at baseline (mean ± SD)	Initial ATEC Total score (mean ± SD)
Mild	20-49	686	29%	5.34±2.30	39±8
Moderate	50-79	1074	46%	4.82±2.34	64±8
Severe	80 ≤	571	25%	4.84±2.45	95±14

Table 3

Characteristics and baseline measures for the English-speaking countries and the rest of the world groups.

Group	Participants in each group (total)	Participants in each group (%)	Age at baseline (mean ± SD)	Initial ATEC Total score (mean ± SD)
English-speaking countries	956	41%	5.43±2.45	63±23
Rest of the world	1367	59%	4.66±2.26	65±23

Table 4

Characteristics and baseline measures for sex groups

Group	Participants in each group (total)	Participants in each group (%)	Age at baseline (mean ± SD)	Initial ATEC Total score (mean ± SD)
Males	1922	83%	4.97±2.36	64±23
Females	404	17%	5.00±2.41	65±24

Table 5. Effect of sex on ATEC scores

Score Type	p-Value
Total Score	0.7368
Subscale 1: Communication	0.8868
Subscale 2: Sociability	0.8474
Subscale 3: Sensory	0.7531
Subscale 4: Physical	0.6310

Table 6. Effect of age on ATEC scores

Score Type	p-Value
Total Score	<.0001
Subscale 1: Communication	<.0001
Subscale 2: Sociability	<.0001
Subscale 3: Sensory	<.0001
Subscale 4: Physical	0.1125

Table 7: LS Means for various age groups. Data are presented as: LS Mean (SE; 95% CI). The difference between Visit 8 and Visit 1 is presented as LS Mean (SE; P-value).

	Visit 1			Visit 8			Visit 8 – Visit 1		
	1-3 YOA	3-6 YOA	6-12 YOA	1-3 YOA	3-6 YOA	6-12 YOA	1-3 YOA	3-6 YOA	6-12 YOA
ATEC Total	65.89 (0.68; 64.55-67.23)	63.78 (0.47; 62.86-64.70)	63.48 (0.61; 62.29-64.67)	37.71 (1.13; 35.50-39.92)	45.09 (0.76; 43.60-46.58)	50.42 (0.96; 48.54-52.30)	-28.18 (1.14; <0.0001)	-18.69 (0.76; <0.0001)	-13.07 (0.98; <0.0001)
Subscale 1 Communication	16.18 (0.17; 15.86-16.50)	15.72 (0.11; 15.51-15.94)	15.45 (0.15; 15.16-15.74)	7.11 (0.26; 6.60-7.63)	11.28 (0.18; 10.94-11.62)	13.51 (0.22; 13.07, 13.95)	-9.07 (0.26; <0.0001)	-4.44 (0.17; <0.0001)	-1.94 (0.22; <0.0001)
Subscale 2 Sociability	14.08 (0.21; 13.67-14.48)	13.44 (0.14; 13.16-13.72)	13.58 (0.18; (13.22-13.94)	7.32 (0.35; 6.62-8.01)	9.37 (0.24; 8.90-9.84)	10.46 (0.30; 9.87-11.06)	-6.76 (0.37; <0.0001)	-4.07 (0.25; <0.0001)	-3.11 (0.32; <0.0001)
Subscale 3 Sensory	15.22 (0.20; 14.84-15.61)	14.69 (0.13; 14.43-14.96)	14.70 (0.17; 14.36; 15.04)	8.52 (0.33; 7.88-9.16)	10.47 (0.22; 10.04-10.91)	11.21 (0.28; 10.66-11.75)	-6.70 (0.34; <0.0001)	-4.22 (0.22; <0.0001)	-3.50 (0.29; <0.0001)
Subscale 4 Physical	Not Significant (N/S)	N/S	N/S	N/S	N/S	N/S	N/S	N/S	N/S

Table 8: LS Mean differences between Age Groups. Data are presented as: LS Mean difference (SE; P-Value)

	Visit 1			Visit 8		
	1-3 vs. 3-6	1-3 vs. 6-12	3-6 vs. 6-12	1-3 vs. 3-6	1-3 vs. 6-12	3-6 vs. 6-12
Total Score	2.11 (0.77, 0.4946)	2.41 (0.87, 0.4717)	0.29 (0.72, 1.0000)	-7.38 (1.33, <0.0001)	-12.71 (1.45, <0.0001)	-5.33 (1.19, 0.0020)
Subscale 1: Communication	0.46 (0.19, 0.7252)	0.73 (0.22, 0.1213)	0.27 (0.17, 0.9969)	-4.17 (0.31, <0.0001)	-6.40 (0.34, <0.0001)	-2.23 (0.28, <0.0001)
Subscale 2: Sociability	0.64 (0.24, 0.5252)	0.50 (0.27, 0.9748)	-0.14 (0.22, 1.0000)	-2.05 (0.42, 0.0003)	-3.14 (0.46, <0.0001)	-1.09 (0.38, 0.3622
Subscale 3: Sensory	0.53 (0.22, 0.7615)	0.52 (0.25, 0.9240)	-0.01 (0.21, 1.0000)	-1.95 (0.39, 0.0001)	-2.69 (0.42, <0.0001)	-0.73 (0.35, 0.9118)
Subscale 4: Physical	N/S	N/S	N/S	N/S	N/S	N/S

Table 9. Effects of a participants' country of origin (English-Speaking Countries group and the Non-English-Speaking Countries group) on ATEC scores

Score Type	p-Value
Total Score	<.0001
Subscale 1: Communication	<.0001
Subscale 2: Sociability	<.0001
Subscale 3: Sensory	0.0622
Subscale 4: Physical	0.0264

Table 10: LS Means of English-Speaking Countries group and the Non-English-Speaking Countries group. Data are presented as LS Mean (SE; 95% CI). The difference between Visit 8 and Visit 1 is presented as LS Mean (SE; P-value).

	Visit 1		Visit 8		Visit 8 – Visit 1	
	English-Speaking Countries	Non-English-Speaking Countries	English-Speaking Countries	Non-English-Speaking Countries	English-Speaking Countries	Non-English-Speaking Countries
ATEC Total	63.41 (0.52; 62.40-64.42)	64.44 (0.45; 63.55-65.32)	46.93 (0.83; 45.31-48.55)	43.43 (0.71; 42.04-44.82)	-16.48 (0.84; <0.0001)	-21.01 (0.70; <0.0001)
Subscale 1 Communication	15.32 (0.12; 15.07-15.56)	15.91 (0.11; 15.69-16.12)	11.30 (0.20; 10.92-11.69)	10.67 (0.17; 10.33-11.00)	-4.01 (0.20; <0.0001)	-5.24 (0.17; <0.0001)
Subscale 2 Sociability	13.40 (0.16; 13.09-13.70)	13.63 (0.14; 13.36-13.90)	9.89 (0.26; 9.38-10.40)	8.64 (0.22; 8.21-9.07)	-3.51 (0.27; <0.0001)	-4.99 (0.23; <0.0001)
Subscale 3 Sensory	N/S	N/S	N/S	N/S	N/S	N/S
Subscale 4 Physical	20.60 (0.23; 20.15-21.05)	20.30 (0.20; 19.91-20.69)	15.50 (0.38; 14.76-6.24)	14.30 (0.32; 13.66-14.93)	-3.51 (0.27; <0.0001)	-5.10 (0.39; <0.0001)

Table 11: LS Mean differences between the English-Speaking Countries group and the Non-English-Speaking Countries group. Data are presented as LS Mean difference (SE; P-Value)

	Visit 1	Visit 8
	English-Speaking Countries vs. Non-English-Speaking Countries	English-Speaking Countries vs. Non-English-Speaking Countries
Total Score	1.03 (0.61, 0.9528)	-3.50 (1.05, 0.0641)
Subscale 1: Communication	0.59 (0.15, 0.0087)	-0.64 (0.25, 0.4409)
Subscale 2: Sociability	0.23 (0.19, 0.9975)	-1.25 (0.33, 0.0145)
Subscale 3: Sensory	N/S	N/S
Subscale 4: Physical	-0.30 (0.28, 0.9994)	-1.21 (0.48, 0.4669)

Table 12. Effects of initial ASD severity on ATEC scores

Score Type	p-Value
Total Score	<.0001
Subscale 1: Communication	<.0001
Subscale 2: Sociability	<.0001
Subscale 3: Sensory	<.0001
Subscale 4: Physical	<.0001

Table 13: LS Means for various severity groups. Data are presented as: LS Mean (SE; 95% CI). The difference between Visit 8 and Visit 1 is presented as LS Mean (SE; P-value).

	Visit 1			Visit 8			Visit 8 – Visit 1		
	Mild	Moderate	Severe	Mild	Moderate	Severe	Mild	Moderate	Severe
ATEC Total	56.97 (0.81; 55.38-58.56)	64.03 (0.48; 63.08-64.97)	72.49 (0.96; 70.61-74.37)	47.77 (1.09; 45.64-49.90)	43.78 (0.76; 42.29-45.28)	42.93 (1.28; 40.42-45.45)	-9.20 (0.94; <0.0001)	-20.24 (0.76; <0.0001)	-29.55 (1.09; <0.0001)
Subscale 1: Communication	14.94 (0.16; 14.64-15.25)	15.75 (0.12; 15.51-15.98)	16.28 (0.17; 15.95-16.61)	11.41 (0.24; 10.95-11.87)	10.65 (0.19; 10.28-11.02)	10.74 (0.27; 10.21-11.26)	-3.53 (0.23; <0.0001)	-5.10 (0.19; <0.0001)	5.54 (0.27; <0.0001)
Subscale 2: Sociability	11.05 (0.20; 10.67-11.44)	13.51 (0.15; 13.22-13.80)	16.51 (0.22; 16.07-16.94)	9.62 (0.30; 9.03-10.22)	8.76 (0.24; 8.29-9.23)	9.12 (0.35; 8.42-9.81)	-1.43 (0.30; 0.006)	-4.75 (0.25; <0.0001)	-7.39 (0.35; <0.0001)
Subscale 3: Sensory	13.05 (0.19; 12.68-13.43)	14.80 (0.14; 14.52-15.07)	16.56 (0.21; 16.14-16.97)	10.78 (0.29; 10.22-11.34)	9.98 (0.22; 9.54-10.42)	9.61 (0.33; 8.96-10.26)	-2.28 (0.28; <0.0001)	-4.81 (0.23; <0.0001)	-6.94 (0.32; <0.0001)
Subscale 4: Physical	17.74 (0.28; 17.19-18.29)	20.10 (0.22; 19.68-20.53)	24.28 (0.32; 23.66-24.91)	15.64 (0.44; 14.78-16.51)	14.38 (0.35; 13.69-15.06)	14.39 (0.51; 13.38-15.40)	-2.10 (0.44; 0.0005)	-5.73 (0.36; <0.0001)	-9.89 (0.51; <0.0001)

Table 14: LS Mean differences between severity groups. Data are presented as: LS Mean difference (SE; P-Value)

	Visit 1			Visit 8		
	Mild vs. Moderate	Mild vs. Severe	Moderate vs. Severe	Mild vs. Moderate	Mild vs. Severe	Moderate vs. Severe
Total Score	-7.06 (0.90, <0.0001)	-15.52 (1.52, <0.0001)	-8.46 (1.03, <0.0001)	3.98 (1.30, 0.2533)	4.83 (1.88, 0.6243)	0.85 (1.46, 1.0000)
Subscale 1: Communication	-0.81 (0.18, 0.0028)	-1.34 (0.23, <0.0001)	-0.53 (0.19, 0.4508)	0.76 (0.29, 0.5940)	0.68 (0.36, 0.9723)	-0.09 (0.32, 1.0000)
Subscale 2: Sociability	-2.46 (0.23, <0.0001)	-5.46 (0.31, <0.0001)	-3.00 (0.25, <0.0001)	0.86 (0.38, 0.8251)	0.51 (0.48, 1.0000)	-0.35 (0.42, 1.0000)
Subscale 3: Sensory	-1.74 (0.22, <0.0001)	-3.50 (0.30, <0.0001)	-1.76 (0.24, <0.0001)	0.79 (0.36, 0.8557)	1.17 (0.45, 0.6095)	0.37 (0.39, 1.0000)
Subscale 4: Physical	-2.36 (0.33, <0.0001)	-6.54 (0.44, <0.0001)	4.18 (0.37, <0.0001)	1.27 (0.55, 0.8037)	1.25 (0.69, 0.9803)	-0.01 (0.61, 1.0000)

Table 15.			
Severity group definition based on initial ATEC total score and age			
Severity group	**Definition based on initial ATEC total score and age**	**Number of participants**	**% of participants in each group**
Mild	20<initial ATEC total score ≤17+119*exp(-age/3.3)	758	33
Moderate	17+119*exp(-age/3.3) <initial ATEC total score ≤27+189*exp(-age/3.3)	816	35
Severe	27+189*exp(-age/3.3) <initial ATEC total score	757	32

Table 16. Effects of severity groups on ATEC scores. Severity group was defined based on initial ATEC total score and age.

Score Type	p-Value
Total Score	<.0001
Subscale 1: Communication	<.0001
Subscale 2: Sociability	<.0001
Subscale 3: Sensory	<.0001
Subscale 4: Physical	<.0001

Table 17: LS Means for various severity groups defined based on initial ATEC total score and age. Data are presented as LS Mean (SE; 95% CI). The difference between Visit 8 and Visit 1 is presented as LS Mean (SE; P-value).

	Visit 1			Visit 8			Visit 8 – Visit 1		
	Mild	Moderate	Severe	Mild	Moderate	Severe	Mild	Moderate	Severe
ATEC Total	57.99 (0.90; 56.23-59.75)	63.17 (0.75; 61.70-64.65)	66.46 (0.82; 64.86-68.05)	41.27 (1.13; 39.06-43.48)	41.32 (1.03; 39.29-43.34)	45.97 (1.07; 43.88-48.07)	-16.72 (0.90; <0.0001)	-21.85 (0.90; <0.0001)	-20.49 (0.90; <0.0001)
Subscale 1: Communication	15.48 (0.20; 15.09-15.87)	15.58 (0.18; 15.22-15.94)	15.07 (0.18; 14.72-15.42)	9.30 (0.26; 8.79-9.80)	10.65 (0.25;10.18-11.14)	12.03 (0.24; 11.55-12.51)	-6.18 (0.21; <0.0001)	-4.93 (0.21; <0.0001)	-3.04 (0.21; <0.0001)
Subscale 2: Sociability	11.43 (0.25; 10.95-11.92)	13.46 (0.22; 13.02-13.89)	14.80 (0.23; 14.36-15.25)	8.32 (0.33; 7.68-8.97)	8.21 (0.32; 7.59-8,83)	9.71 (0.32; 9.09-10.34)	-3.11 (0.29; <0.0001)	-5.24 (0.29; <0.0001)	-5.09 (0.29; <0.0001)
Subscale 3: Sensory	13.14 (0.24; 12.67-13.61)	14.47 (0.21; 14.05-14.89)	15.27 (0.22; 14.83-15.70)	9.21 (0.31; 8.60-9.83)	9.27 (0.30; 8.69-9.86)	10.37 (0.30; 9.78-10.96)	-3.92 (0.27; <0.0001)	-5.20 (0.27; <0.0001)	-4.90 (0.27; <0.0001)
Subscale 4: Physical	17.40 (0.36; 16.70-18.10)	19.89 (0.33; 19.24-20.53)	22.56 (0.34; 21.90-23.22)	13.85 (0.48; 12.91-14.79)	13.22 (0.46; 12.31-14.13)	14.95 (0.47; 14.04-15.87)	-3.55 (0.42; <0.0001)	-6.67 (0.42; <0.0001)	-7.60 (0.42; <0.0001)

Table 18: LS Mean differences between severity groups defined based on initial ATEC total score and age. Data are presented as LS Mean difference (SE; P-Value)

	Visit 1			Visit 8		
	Mild vs. Moderate	Mild vs. Severe	Moderate vs. Severe	Mild vs. Moderate	Mild vs. Severe	Moderate vs. Severe
Total Score	-5.18 (0.83; <0.0001)	-8.47 (1.16; <0.0001)	-3.28 (0.89; 0.0437)	-0.05 (1.29; 1.0000)	-4.70 (1.51; 0.2269)	-4.65 (1.33; 0.0786)
Subscale 1: Communication	-0.11 (0.18; 1.0000)	0.41 (0.22; 0.9757)	0.51 (0.19; 0.5653)	-1.36 (0.30; 0.0012)	-2.73 (0.32; <0.0001)	-1.38 (0.30; 0.0014)
Subscale 2: Sociability	-2.02 (0.24; <0.0001)	-3.37 (0.30; <0.0001)	-1.35 (0.25; <0.00001)	0.11 (0.39; 1.0000)	-1.39 (0.43; 0.1643)	-1.50 (0.40; 0.0374)
Subscale 3: Sensory	-1.34 (0.23; <0.0001)	-2.13 (0.29; <0.0001)	-0.79 (0.24; 0.1361)	-0.06 (0.37; 1.0000)	-1.15 (0.41; 0.4045)	-1.10 (0.38; 0.3602)
Subscale 4: Physical	-2.49 (0.34; <0.0001)	-5.16 (0.43; <0.0001)	-2.67 (0.37; <0.0001)	0.63 (0.57; 1.0000)	-1.11 (0.62; 0.9852)	-1.74 (0.59; 0.3323)

Table 19.

Comparison of the annualized decrease of ATEC score across multiple studies.

	Number of Participants	Age at baseline (years)	Initial Total ATEC score	Number of ATEC Evaluations Completed by each Participant	Study duration (years)	Annualized change in score: LS Mean (SE) or mean±				
						Total Score	Subscale 1	Subscale 2	Subscale 3	Sul
Current Study: 1.2-5YOA	**516**	**2.4±0.5**	**71±22**	**5.9±3.7**	**2**	**-14.1 (0.6)**	**-4.5 (0.1)**	**-3.4 (0.2)**	**-3.4 (0.2)**	**N/**
Whitehouse et al./ TOBY[A]	36	3.3±0.7	71±23	3	0.5	-25.6±49.6	-8.6 ±19.0	6.7±15.0	-4.8±19.4	-5.:
Current Study: 3.1-5YOA	**1176**	**4.3±0.9**	**62±23**	**5.4±3.5**	**2**	**-9.4 (0.5)**	**-2.2 (0.1)**	**-2.0 (0.1)**	**-2.1 (0.1)**	**N/**
Charman et. al.	57	4.7±0.7	81±18	2	1	-8.5±6.9	-3.9±4.1	-1.3±5.4	-2.2±5.1	-1.:
Magiati et. al.	22	5.6±0.6	53±21	2	4.8	1.3±3.6	-0.5±0.6	0.7±1.3	0.3±1.1	0.9
Current Study: 6.1-12YOA	**639**	**8.2±1.6**	**62±24**	**5.1±2.9**	**2**	**-6.5 (0.5)**	**-1.0 (0.1)**	**-1.6 (0.2)**	**-1.8 (0.2)**	**N/**

Includes both test and control groups as there was no difference in ATEC score change between the groups.