Biochemical Characterization of RecBCD Enzyme from An Antarctic *Pseudomonas* Species and

Identification of Its Cognate Chi (χ) Sequence

Theetha L. Pavankumar*[1,2,#], Anurag Kumar Sinha*[1,3] and Malay K. Ray[1,#]

[1]CSIR-Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad-500007, INDIA

[2] Department of Microbiology and Molecular Genetics, Briggs Hall, One Shields Ave, University

of California, Davis, CA, USA.

*These authors have contributed equally to this work

Present Address:

[3] Department of Biology, University of Copenhagen, Ole Maaløes Vej 5, Copenhagen, Denmark.

Address for correspondence:

Theetha L Pavankumar

Department of Microbiology and Molecular Genetics,

One Shields Ave, University of California, Davis, CA-95616, USA

Telephone: +1 530-754-9702

Fax: +1 530-754-8973

E-mail: pavan@ucdavis.edu

Malay K. Ray

CSIR-Centre for Cellular and Molecular Biology

Uppal Road, Hyderabad 500 007, INDIA

Telephone: +91 40 2719 2512

Fax :   +91 40 2716 0591

E-mail : malay.ccmb@gmail.com

ABSTRACT

Pseudomonas syringae Lz4W RecBCD enzyme, RecBCD^{Ps}, is a trimeric protein complex

comprised of RecC, RecB, and RecD subunits. RecBCD enzyme is essential for *P. syringae*

growth at low temperature, and it protects cells from low temperature induced replication arrest.

In this study, we show that the RecBCD^{Ps} enzyme displays distinct biochemical behaviors. Unlike

E. coli RecBCD enzyme, the RecD subunit is indispensable for RecBCD^{Ps} function. The RecD

motor activity is essential for the Chi-like fragments production in *P. syringae*, highlighting a

distinct role for *P. syringae* RecD subunit in DNA repair and recombination process. Further, the

ssDNA-dependent endonuclease activity is notably absent in RecBCD^{Ps} enzyme. Here, we

demonstrate that the RecBCD^{Ps} enzyme recognizes a unique octameric DNA sequence, 5'-

GCTGGCGC-3′ (Chi^{Ps}) that attenuates nuclease activity of the enzyme when it enters dsDNA

from the 3′-end. We propose that the reduced translocation activities manifested by motor-

defective mutants cause cold sensitivity in *P. syrinage*; emphasizing the importance of DNA

processing and recombination functions in rescuing low temperature induced replication fork

arrest.

Keywords:

RecBCD enzyme, *Pseudomonas,* Chi sequence, cold adaptation, DNA damage, DNA repair and

recombination, DNA helicase and nuclease.

Abbreviations:

ATP, Adenosine triphosphate; DSB, double-strand break; 'Chi, Crossover hotspot instigator; Ni-

NTA, Nitrio tri-acetic acid; TLC, thin layer chromatography; MMC, mitomycin C; UV light, Ultra

violet; ABM, Antarctic bacterial medium; LB, Luria-Bertani medium.

INTRODUCTION

The RecBCD enzyme-mediated homologous recombination is a DNA repair pathway that ensures genome integrity by faithful repair of broken DNA in *E. coli* and many Gram-negative bacteria. The heterotrimeric RecBCD enzyme complex, comprised of RecB, RecC, and RecD subunits, is essential for double-strand breaks (DSBs) repair via homologous recombination and protects host cells from foreign DNAs and invading phages (1-4). DSBs are generated in cells by various exogenous and endogenous factors including the running of replication forks into preexisting lesions (5). RecBCD enzyme is a highly processive helicase and nuclease, which unwinds and degrades DNA strands asymmetrically from a blunt or nearly blunt dsDNA (2). Initially, RecBCD enzyme degrades 3′-ended DNA preferentially over the 5′-end of the DNA until it encounters a regulatory DNA sequence called 'Chi' (Crossover hotspot instigator; 5′-GCTGGTGG-3′ in *E. coli*) (2, 6, 7). Chi (χ) recognition switches RecBCD enzyme's polarity of DNA degradation. It attenuates 3′→5′ nuclease activity but upregulates 5′→3′ nuclease activity resulting in the production of 3'-ended ssDNA tail (8). The RecBCD enzyme loads RecA onto the 3'-terminal ssDNA (9) facilitating the homologous pairing to form Holliday junction. The RuvAB/C complex further resolve these recombination DNA intermediates to promote DNA repair process via homologous recombination (10). The behavior of RecBCD enzyme was also studied using single molecule techniques (11). Single molecule analyses of *E. coli.* RecBCD enzyme revealed that it translocates on DNA at a much higher rate, before Chi and pauses at Chi (12). The Chi recognition switches lead motor subunit of the RecBCD enzyme, from fast to slow motor (RecD to RecB), resulting in the reduction of translocation rate by one-half after Chi (13).

Previously, we have shown that the RecBCD enzyme of Antarctic *Pseudomonas syringae* Lz4W (RecBCD[Ps]) is essential for the growth at low temperature (14). Growth at low temperature induces frequent replication arrest causing fork reversal mediated DNA damage in *P. syringae* Lz4W (15). The RecBCD[Ps] enzyme thus rescues cells from replication-arrest dependent DNA

damage enabling *P. syringae* Lz4W to grow at low temperature (15). Unlike in *E. coli*, the RecD

subunit of *P. syringae* Lz4W is an indispensable subunit of RecBCD complex (16). The genetic

analysis of ATP binding and nuclease defective mutants of RecBCD[Ps] enzyme indicated that the

ATP driven motor activities of both RecD and RecB motor subunits are critical for growth at low

temperature, whereas the nuclease activity of holoenzyme is dispensable (14).

In this study, we performed a biochemical analysis of wild-type and mutant RecBCD[Ps]

enzymes to understand their biochemical role in protecting *P. syringae* Lz4W cells from low

temperature induced DNA damage. Here, we report that the ATP dependent DNA unwinding, not

the DNA degradation activity of RecBCD enzyme is critical for growth at low temperature.

Besides, inactivation of ATPase activity of RecD and RecB motor subunits has impaired the DNA

unwinding activity of RecBCD enzyme leading to cold-sensitive phenotype *in vivo*. Furthermore,

we have identified the *P. syringae* Chi sequence (5′-GCTGGCGC-3′), which attenuates the

RecBCD[Ps] nuclease activity *in vitro*.

RESULTS

Purification of RecBCD complex and its mutant variants from *P. syringae* Lz4W

The *P. syringae recCBD* genes were overexpressed on pGL10 derived plasmids in Δ*recCBD* (LCBD) strain to obtain RecB, RecC (N-terminal His-tagged) and, RecD proteins in an equimolar ratio (13). The wild-type and mutant RecBCD enzymes (RecB^{K28Q}CD, RecBCD^{K229Q}, and RecB^{D1118A}CD) were purified by two-step purification protocol as shown in the flowchart (Fig. 1A) using Ni-NTA affinity column chromatography and size exclusion chromatography. The purified fractions of RecBCD and mutants contained all the three protein subunits (RecB, RecC, and RecD) in a stoichiometric ratio (Fig. 1B). Silver nitrate staining of gels further confirmed the purity of protein fractions. However, silver staining of SDS-PAGE separated proteins showed an additional protein band of ~60 kDa size (Fig. 1C). The Mass spectrometric analysis of this protein band indicated that it belongs to HSP-60 family of chaperonin, GroEL. The appearance of GroEL as a contaminant during purification of RecBCD protein is also evidenced in *E. coli (17)*. Nonetheless, the presence of RecB, RecC, and RecD subunits was further confirmed by Western analysis using the polyclonal antibodies specific to these proteins (Fig. S1).

ATP hydrolysis activity of the wild-type and the mutant RecBCD enzymes

ATP hydrolyzing activity of wild-type and mutant RecBCD enzymes was measured by thin layer chromatography (TLC) as described in Materials and methods. The wild-type RecBCD enzyme displayed DNA stimulated ATPase activity in the presence of linear pBR322 double-stranded DNA (dsDNA). At 22ºC, RecBCD hydrolyzed ATP with the maximum velocity (V_{max}) of 236.5 μmol ATP per μmol enzyme s^{-1} and a K_m of 57.8 μM for ATP (Fig. 2A and Table 1). The mutant RecB^{D1118A}CD enzyme (nuclease defective mutant) showed ATPase activity similar to the wild-type enzyme. However, the ATP-binding defective mutant enzymes such as RecB^{K28Q}CD (mutation in the consensus ATP binding site of RecB) and RecBCD^{K229Q} (mutation in the consensus ATP binding site of RecD) showed a 10-fold decrease in ATP hydrolyzing activities (Table 1).

ATP hydrolyzing activities of RecBCD enzymes were measured at three different

temperatures (37, 22 and 4°C) (Fig. 2B). Interestingly, The RecBCD enzyme showed the highest

ATPase activity at 37°C compared to 22°C and 4°C. At 22°C, the optimum temperature for *P.*

syringae Lz4W growth, the wild-type RecBCD and the nuclease-deficient RecB^{D1118A}CD enzymes

displayed ~40% lower ATPase activity compared to 37°C. At 4°C, the activities were further

reduced to ~50% of their respective ATPase activity observed at 22°C. The ATP hydrolyzing

defective mutants showed ~10-fold decrease in ATP hydrolyzing activity compared to wild-type

RecBCD enzyme, at all the temperatures tested (Table 1 and Fig. 2C). It indicates that the

mutations were previously shown to prevent *P. syringae* growth at low temperature severely

curtailed ATP hydrolyzing activity.

DNA unwinding and degradation activities of the RecBCD enzyme at different

concentration of Mg^{++} and ATP

The DNA unwinding and degradation activity of *E. coli* RecBCD enzyme (RecBCD^{Ec}) are

free [Mg^{2+}] ion dependent. An increase in free [Mg^{2+}] increases nucleolytic cleavage by RecBCD

enzyme (18). To understand the DNA unwinding and degradation behavior of *P. syringae*

RecBCD enzyme (RecBCD^{Ps}), we performed experiments at various concentrations of

magnesium and ATP as described in Materials and methods. In the first set of reactions, the molar

concentration of magnesium was kept constant (2 mM) and the ATP concentration was varied (0-

10 mM); and in the second set of reactions, the ATP concentration was kept constant (2 mM) and

magnesium concentration was varied (0-10 mM). The results indicate that the DNA unwinding

and degradation properties of RecBCD^{Ps} enzyme are also modulated by the ratio of Mg^{2+} and

ATP. When the molar concentration of Mg^{2+} is lesser than ATP, RecBCD^{Ps} enzyme unwinds the

dsDNA substrate, and the degradation of DNA is not much pronounced (Fig. 3A, lane 4-6).

However, when the molar concentration of Mg^{++} exceeds the ATP concentration, RecBCD^{Ps}

degrades unwound DNA more vigorously (Fig. 3B, lane 4-10). The ratio of [Mg^{2+}]: [ATP] thus

affects the kinetics of DNA unwinding and degradation properties of RecBCD enzyme. We also

observed that an increase in ATP concentration more than three folds over Mg^{++} inhibits the DNA

unwinding activity of RecBCD^{Ps} enzyme (Fig. 3A, lane 7-10). The inhibition of DNA unwinding

activity is possibly due to sequestration of a Mg^{2+} ion by ATP leading to the depletion of free

magnesium in the reaction. Subsequently, the DNA unwinding and degradation experiments were

performed under specified reaction conditions. The limiting magnesium reaction condition (5 mM

ATP, 2 mM Mg^{++}) was chosen to observe the DNA unwinding activity and, the excess magnesium

condition (2 mM ATP, 6 mM Mg^{++}) to observe the DNA unwinding and degradation activities of

RecBCD enzyme. Interestingly, under excess magnesium conditions, we observed three shorter

discrete DNA fragments (Fig. 3B, Lane 5-10). These DNA fragments production by RecBCD

enzyme is possibly due to Chi-like sequence on a DNA substrate and is discussed later in the

results section.

Calcium has been shown to inhibit the nuclease activity of *E. coli* RecBCD enzyme (19).

We also studied the effects of calcium on *P. syringae* RecBCD by increasing the Ca^{++} ion

concentration in a reaction mixture that contained fixed amounts of ATP and Mg^{2+} (2 and 6 mM

respectively) (Supplementary Fig. S2). We found that similar to RecBCD^{Ec}, the high concentration

of calcium inhibits both helicase and nuclease activity of RecBCD^{Ps} enzyme.

Temperature-dependent DNA unwinding and nuclease activity of *P. syringae* RecBCD

enzyme

ATPase RecBCD mutants affect *P. syringae* Lz4W growth in a temperature-dependent

manner. Hence, we sought to assess the effects of temperature on the DNA unwinding and

degradation properties of RecBCD^{Ps} enzyme. We measured the enzyme activities at 22°C, and

4°C, using NdeI linearized [5′-^{32}P] labeled pBR322 plasmid as a substrate. Under the limiting

magnesium reaction condition (5 mM ATP, 2 mM Mg^{2+}), the DNA unwinding activity (i.e.,

production of full-length ssDNA) of RecBCD^{Ps} enzyme was detected only at 22°C and not at 4°C

(Fig. 4A). However, under excess magnesium reaction condition (2 mM ATP, 6 mM Mg^{++}) the

DNA degradation activity of RecBCD^{Ps} enzyme was observed at both 22°C and 4°C (Fig. 4B).

We calculated the rate of DNA unwinding by measuring the decreased band intensities of

5′-[γ-^{32}P]-labeled dsDNA substrates at different temperatures. The wild-type RecBCD^{Ps} unwound

the DNA at the rate of 35.1 ± 1.6 bp/sec at 22°C when ATP and Mg^{++} ratio was 5:2 (limiting

magnesium condition) (Table 2). However, the enzyme could unwind the DNA even much faster,

when the ATP and Mg^{++} ratio was changed to 2:6 (excess magnesium condition). Under the latter

condition, wild-type RecBCD^{Ps} enzyme unwound (also degraded) the DNA at the rate of 101.5 ±

3.3 bp/sec (Table 2). Notably, under the limiting magnesium reaction condition at low temperature

(4°C), the RecBCD enzyme failed to produce detectable unwound DNA products (Fig. 4A).

However, under excess magnesium conditions, RecBCD enzyme could unwind and degrade the

DNA at the rate of 55.8 ± 6.9 bp/s (Table 2, Fig. 4B), which is about 54% of the activity observed

at 22°C. The apparent unwinding rates obtained under excess magnesium conditions, based on

the disappearance of ^{32}P-end-labeled DNA substrate could be an over-estimation. It is possible

that some enzyme molecules could just nucleolytically remove the end-labeled nucleotide, but

couldn't fully unwind the dsDNA substrate.

DNA unwinding and nuclease activities of mutant RecB^{K28Q}CD, RecBCD^{K229Q}, and

RecB^{D1118A}CD enzymes

We have shown that *P. syringae* cells carrying *recB^{K28Q}CD* or *recBCD^{K229Q}* mutants are

sensitive to cold temperature, UV irradiation and Mitomycin C (MMC) (14). These two mutant

enzymes also display very weak ATPase activity. To understand the impact of these mutations

on the DNA unwinding and degradation properties of RecBCD complex, we analyzed RecB^{K28Q}CD

and RecBCD^{K229Q} enzymes *in vitro* at 22°C and 4°C. At 22°C, under limiting and excess

magnesium conditions, RecB^{K28Q}CD with the ATPase-defective RecB subunit displayed weak

helicase activity (Fig. 5A). The rate of DNA unwinding was 1.5 ± 0.7 bp/sec and 12.9 ± 3.4 bp/sec

at 22°C, under limiting and excess magnesium conditions respectively (Fig.5A and Table 2). At

4°C, on the other hand, RecB^{K28Q}CD showed no detectable DNA unwinding activity under the

excess magnesium condition (Fig. 5A and Table 2). These results suggest that RecB^{K28Q}CD is a

poor helicase-nuclease enzyme.

Compared to the RecB^{K28Q}CD enzyme, RecBCD^{K229Q} enzyme with the ATPase-defective

motor RecD subunit displayed higher DNA unwinding and nuclease activities. At 22°C, the

RecBCD^{K229Q} enzyme with the ATPase-defective RecD subunit displayed substantial DNA

unwinding activity, 27.2 ± 6.1 bp/sec and 92.6 ± 14.1 bp/sec, under limiting and excess

magnesium conditions respectively (Fig. 5B and Table 2). However, at 4°C under excess

magnesium conditions, the mutant enzyme unwound the DNA at the rate of 17.3 ± 9.6 bp/sec (Fig

5B and Table 2). The combined DNA unwinding and degradation activity of the RecBCD^{K229Q}

enzyme (under excess magnesium) were ~90% and ~30% of the wild-type RecBCD^{Ps} activity at

22°C and 4°C respectively. Interestingly, the mutant RecBCD^{K229Q} enzyme failed to produce

discrete DNA fragments (Fig. 5B).

We also tested DNA unwinding and degradation by the nuclease defective RecB^{D1118A}CD

enzyme. (Fig. 5C). The mutation in the nuclease catalytic site of RecB subunit (RecB^{D1118A}CD)

led to a complete loss of *in vivo* nuclease activity in RecBCD^{Ps} complex, without affecting the

recombination proficiency and cold adaptation of bacteria (14, 15). At 22°C, under limiting

magnesium reaction condition, RecB^{D1118A}CD enzyme produced single-stranded pBR322 DNA at

the rate 30.4 ± 1.7 bp/sec, which is 85% of the rate observed with the wild-type enzyme. Under

excess magnesium condition, the rate of DNA unwinding increased to 109.7 ± 17.5 bp/sec, which

is similar to wild-type. At 4°C, this enzyme could unwind the pBR322 DNA at the rate of 41.0 ±

8.1 bp/sec, which is about 75% of wild-type enzyme (Table 2). More importantly, under both

limiting and excess magnesium conditions, this enzyme as expected, produced only the full-length

ssDNA of pBR322 and did not degrade the DNA (Fig. 5C). From these data, it is clear that

RecB^{D1118A}CD enzyme is nuclease deficient *in vitro*, but its DNA unwinding activity is comparable

to wild-type enzyme at both 22 and 4°C (Fig. 5C and Table 2).

P. syringae RecBCD enzyme does not have endonuclease activity

The ATP-dependent endonuclease activity of wild-type RecBCD enzyme of *P. syringae*

was examined using a circular M13 ssDNA as substrate as previously described for RecBCD^{Ec}

(20-22). We tested the endonuclease activity under three different conditions of ATP and

magnesium concentrations as described in Supplementary Fig S4. Under all the three conditions,

wild-type RecBCD and mutant proteins failed to degrade M13 phage ssDNA (Fig S3). It indicates

that unlike RecBCD^{Ec}, the RecBCD^{Ps} does not exhibit endonucleolytic activity under the

conditions tested.

A specific DNA sequence on pBR322 plasmid modulates nuclease activity of *P. syringae*

RecBCD.

We noticed that, under the excess magnesium reaction conditions, RecBCD^{Ps} produced

a discrete DNA fragments shorter than the full-length unwound DNA substrate (Fig 3B, Lanes 5-

10), and the shortest DNA fragment showed a higher intensity than the other ones (Fig 3B). This

interesting observation led us to hypothesize that the *P. syringae* RecBCD enzyme recognizes a

specific DNA sequence. This specific DNA sequence could potentially be a Chi (χ) like sequence;

which alters the nuclease activity of RecBCD^{Ps} complex, allowing 3'-ended ssDNA to escape from

DNA degradation as observed earlier (2, 8, 23, 24). To further confirm that these DNA fragments

are indeed single-stranded, we performed degradation assays in the presence of Exonuclease-I

(ExoI), which specifically degrades ssDNA by its 3′→5′ exonuclease activity (25). The RecBCD^{Ps}-

generated short DNA fragments disappeared in the presence of ExoI (Fig S4A) suggesting that

they are indeed ssDNA, similar to Chi-specific DNA fragments produced by *E*. *coli* RecBCD

enzyme after Chi-recognition (8, 23, 24, 26).

Using genetic experiments, we have previously shown that RecBCD[Ps] enzyme does not

recognize *E. coli* Chi sequence (5'-GCTGGTGG-3') (14, 16). We performed biochemical

experiments using pBR322[χ+3F3H] (a pBR322 derivative, containing three tandem *E. coli* χ

sequences) (27) and pBR322 (lacking *E. coli* χ sequence) as a DNA substrate. Interestingly,

ssDNA fragments produced by RecBCD[Ps] enzyme were identical in size with both the substrates,

confirming that RecBCD[Ps] does not recognize *E. coli* Chi sequence but recognizes an unknown

DNA sequence of pBR322 plasmid (Fig S4B). To locate putative Chi sequence of RecBCD[Ps]

enzyme, we amplified 3.6 kb segment of pBR322 using primer OROPI and OROPII

(Supplementary Table ST2), which excludes rop region of the plasmid (Fig. S5A). Amplified

fragments were 5'-labeled, and the assays were performed in the presence of excess magnesium.

Apparently, all the three shorter ssDNA fragments were also observed with the 3.6 kb DNA

substrate, indicating the presence of putative Chi sequence within 3.6 kb of the plasmid

(Supplementary Fig. S5C).

Chi dependent protection of pBR322 DNA fragments are strand specific

E. coli RecBCD enzyme recognizes Chi sequence in a specific orientation, 5'-

GCTGGTGG-3', when enzyme enters through 3′-end (28) and the 3'-ended ssDNA is being

protected from RecBCD nuclease activity after Chi recognition (8, 29, 30). Here, we examined

which strand of the linearized pBR322 is being protected to produce these discrete DNA bands

(Fig. 3B). Hence, we performed DNA degradation assay with RecBCD[Ps], in which only one DNA

strand was labeled at a time. We individually labeled the OROPI and OROPII primers with γ^{32}P

at 5′-end and amplified the 3.6 kbp region of pBR322 plasmid with one unlabeled and another

^{32}P-labeled primer or with both labeled primers. In our assay, we always used sub-saturating

concentration of RecBCD[Ps] enzyme compared to DNA substrate, so that RecBCD[Ps] enzyme

enters through either one or the other end, but not through both the ends. DNA degradation

assays using these DNA substrates produced all the three bands when top strand (OROPII end)

was alone labeled, or when the both strands were end-labeled (Fig. 6A). We did not observe any ssDNA band when the bottom strand (OROPI end) was labeled (Fig. 6A), suggesting that RecBCD^{Ps} recognizes a Chi-like sequence in a specific orientation and has a polarity for Chi sequence recognition. These results also suggest that all the Chi-like sequences are on the top strand.

We then performed DNA degradation assays using two other DNA substrates amplified from the pBR322 plasmid. A PCR amplified 2.8 kbp DNA fragment (using OROPII and pBRS1R primers) and a 2.5 kbp PCR amplified fragment (using OROPII and pBRB1R primers set (Fig. S6A). The DNA band of the lowest size (~2.4 kb) and the highest intensity (the prominent DNA band) was the common ssDNA fragment produced by RecBCD^{Ps} in all the three DNA substrates (Supplementary Fig.S6B). This suggests that one of the Chi-like sequences has the strongest RecBCD-inhibitory activity and this site (we designate it as Chi^{Ps}) is located proximal to the 2.5 kbp substrate end (as the protected ssDNA was about ~ 2.4 kb). Two upper DNA bands with less intensity could be due to variants of Chi^{Ps}, which might have a weak inhibitory effect on RecBCD^{Ps} nuclease activity (see below).

Identification of Chi^{Ps} as an 8-mer (5'-GCTGGCGC-3') sequence that modulates *P. syringae* RecBCD nuclease activity and protects DNA from further degradation

To identify the precise location of Chi^{Ps} site in pBR322 DNA substrate, we generated several internally deleted constructs of pBR322 plasmid by site-directed segment-deletion as described in Materials and methods. The 3.6 kbp DNA region of pBR322 plasmid with deleted region/s were PCR amplified using OROPI and OROPII primers. DNA degradation assays were then performed with RecBCD^{Ps} enzyme using the DNA fragments with internal deletion as substrates (Supplementary Fig. S7).

The localization of Chi^{Ps} on the plasmid was first based on two hypotheses: the sequence might be GC rich and should be located close to the right end of 2.5 kbp DNA fragment. From the pBR322 nucleotide sequence analysis, it appeared that a GC-rich region spanning the region

between 273-450 nucleotides within *tet^R* gene of pBR322 might contain the Chi^Ps. Accordingly,

3.6 kbp DNA containing two deletions (Δ273-381 and Δ400-450) were initially tested by the Chi-

protection assays.

The 3.6 kbp DNA deleted for 273-381 region produced the Chi-specific high-intensity

fragment (Supplementary Fig. S7), while 400-450 nucleotides deletion failed to produce it (Fig.

6B). This indicated that the Chi^Ps sequence is located within 400-450 nucleotide region of the

pBR322 plasmid. The three additional deletions within the 400-450 region of the plasmid segment

were made; Δ401-419, Δ421-439, and Δ441-449 (Fig. 6C, D). DNA degradation assays with the

fragments in which these sequences were deleted revealed that Chi is located within the 421-439

nucleotides segment, as these deletions did not produce a protected prominent DNA fragment

(Fig. 6C). we further made two more deletion constructs (Δ421-429 and Δ431-439) within the 421-

439 nucleotides region, and tested for Chi-specific fragment production. Interestingly, deletion of

base-pair from 431-439 bp abolished the prominent DNA band (Fig. 6D) indicating that the deleted

region between 431-439 bp, 5′-GCTGGCGCC-3′ contains the putative Chi sequence of *P.*

syringae. The schematic representation of all the constructs with deleted nucleotide regions and,

the presence or absence of protected prominent DNA band (putative Chi sequence) are also

shown in Fig. 6E.

The identification of putative Chi^Ps sequence further prompted us to investigate the reason

for producing an apparent three distinct DNA fragments by RecBCD^Ps enzyme (Fig. 3B). We

speculated that pBR322 DNA substrate might have sequences that are similar but not identical

to Chi^Ps sequence, and might act like Chi^Ps sequence with weaker recognition property under the

in vitro assay conditions. The 9-mer (5′-GCTGGCGCC-3′) sequence appears only at one place

on the pBR322 substrate. Considering the *E. coli* Chi sequence is an octamer, we first looked in

to 5′-GCTGGCGC -3′ (first 8 nucleotides of the 9-mer from the 5′-end), and 5′-CTGGCGCC-3′

(eliminating first G of the 5′-end) as possible Chi-like sequences. These two octamer sequences

appear at a single location on the pBR322 substrate. However, among the 7-mer combinations

we looked into, the first 7 nucleotide sequence, 5′-GCTGGGC -3′, was located in three reasonable

positions on pBR322 substrate which could potentially produce discrete DNA bands as observed

in DNA degradation assays. Hence, we further focused on 5′-GCTGGCGC -3′ sequence as a

putative Chi[Ps] sequence. We found that the putative Chi[Ps] sequence (5′-GCTGGCGC -3′) is

located at 431-438 bp position of pBR322 and, the similar 7-mer sequences at three different

locations. The first one at 964-971 position (5′-GCTGGCG**T**-3′); the second one at 1493-1500

position (5′-GCTGGCG**G**-3′) and the third one at 2525-2532 position (5′-GCTGGCG**T**-3′) (Fig S8).

All the three-bands show 7 bases identical to the 8 bp Chi[Ps] sequence (5′-GCTGGCGC**-**3′), and

occur in the same orientation (5′→3′) as the Chi[Ps]. The RecBCD[Ps] enzyme might recognize 8 mer

sequences (5′-GCTGGCGC-3′) as well as the 7+1 mer sequences (5′-GCTGGCG+**T/G**-3′)

resulting in multiple DNA bands. Although it is expected to observe four discrete DNA bands, we

observed only three DNA bands. The fourth-expected DNA band with a size of ~230 bp (when

RecBCD[Ps] enzyme enters from 3'-side of the NdeI linearized pBR322 substrate as depicted in

Fig. S8) was not apparent in in-vitro experiments performed on agarose gels. Possibly, it is the

last Chi-like sequence to recognize by RecBCD enzyme and also being the shortest one with ~

230 bp (Fig. S8).

To better define the Chi[Ps] sequence, we mutated the **T** to **C** at the 971[st] position of pBR322,

which creates a sequence identical to the putative 8-mer Chi[Ps] sequence and performed the Chi-

protection assay with RecBCD[Ps] enzyme. Interestingly, the intensity of the Chi-protected second

DNA fragment (after T to C mutation) appeared stronger, and the intensity was similar to the

prominent DNA band (Fig. 7A), which establish the strong recognition of octamer sequence (5′-

GCTGGCGC-3′) as Chi[Ps].

Cloning of Chi[Ps] in pBKS plasmid and confirmation of ectopic Chi[Ps] activity

To further confirm that the octamer sequence (5′-GCTGGCGC-3′) is indeed the Chi

sequence in *P. syringae* and can function as an ectopic Chi[Ps] sequence, we cloned the

octanucleotide sequence into pBKS which is devoid of Chi[Ps] sequence (Table ST1). RecBCD[Ps]

reactions under excess magnesium were performed with XbaI digested linear pBKS (Chi[0])

plasmid and the Chi[Ps] containing pBKS (pBKS-Chi[Ps]) plasmid. As expected, no Chi-protected DNA

bands were observed using the linear pBKS (Chi[0]) DNA as substrate. In contrast, the expected

Chi specific DNA fragment was observed in the case of XbaI digested pBKS-Chi plasmid (Fig.

7B). These experiments confirm that RecBCD[Ps] recognizes an 8-mer sequence (5′-GCTGGCGC-

3′) as Chi[Ps] sequence during resection of double-stranded DNA ends, and thus Chi[Ps] attenuates

the nuclease activity of RecBCD enzyme and promotes DNA recombination and repair.

DISCUSSION

We have earlier shown that RecBCD protein complex is essential for cold adaptation in

Antarctic *P. syringae* Lz4W (14-16). In this study, we have analyzed the biochemical properties

of the wild-type (RecBCD^{Ps}) and three mutant enzymes (RecB^{K28Q}CD, RecBCD^{K229Q}, and

RecB^{D1118A}CD) to analyze the role of RecBCD during growth at low temperature. We also report

for the first time, *Pseudomonas* octameric Chi-sequence (Chi^{Ps}), (5′-GCTGGCGC-3′), and its

ability to attenuate the nuclease activity of *P. syringae* RecBCD enzyme *in vitro*.

Role of ATPase activity of RecB and RecD subunits

The analyses of wild-type and mutant enzymes of *P. syringae* have revealed that mutation

in the critical ATP binding sites of RecB (RecB^{K28Q}) and RecD (RecD^{K229Q}) motor subunit reduces

the ATPase activity of trimeric RecBCD complex by ~10 fold. The mutation in the nuclease active-

site does not affect the ATPase activity. *P. syringae* cells carrying the alleles of RecB^{K28Q} or

RecD^{K229Q} are sensitive to cold temperature, UV irradiation and MMC (14). The ~ 10-fold reduction

of ATPase activity observed *in-vitro* supports the idea that ATP hydrolysis by RecB and RecD

subunits in RecBCD^{Ps} holoenzyme is essential for *P. syringae* survival at low temperature.

DNA unwinding and degradation activities of RecBCD and mutant enzymes

Our data suggest that the DNA unwinding and degradation properties of RecBCD enzyme

depend on ATP and Mg^{++} concentrations, and on the temperature *in vitro*. Under limiting

magnesium reaction condition, at 22°C, the RecB^{K28Q}CD, RecBCD^{K229Q} and RecB^{D1118A}CD

enzymes have retained about 4%, 77% and 85% of the wild-type DNA unwinding activity

respectively, while at 4°C, we could not detect the DNA unwinding activity of both wild-type and

mutant RecBCD^{Ps} enzymes under the limiting magnesium reaction condition. In contrast, when

magnesium was in excess over ATP, the DNA unwinding and degradation activity of the RecBCD

enzymes could be measured 22°C and 4°C). These results suggest that the excess of magnesium

over ATP is favorable for the helicase and nuclease activities of RecBCD^{Ps} enzyme.

The RecB^{K28Q}CD enzyme is a poor helicase and with no detectable nuclease activity *in*

vitro even at 22°C. However, at 4°C, RecB^{K28Q}CD enzyme failed to unwind and degrade the DNA

under both limiting and excess magnesium reaction conditions. Hence, low-temperature growth

sensitivity of the *P. syringae recB^{K28Q}CD* mutant can be attributed to the lack of RecBCD

unwinding and/degradation activities at 4°C.

The RecBCD^{K229Q} enzyme with defective RecD ATPase unwinds and degrades the

dsDNA, albeit at the reduced rate. The DNA unwinding/degradation rate of RecBCD^{K229Q} mutant

enzyme is reduced to one-third of the wild-type enzyme at 4°C. Therefore, we propose that its

inability to support the DNA repair process and growth at low temperature is possibly due to its

decreased DNA unwinding/degradation activity, particularly at low temperature. Interestingly, the

RecBCD^{K229Q} enzyme shows lack of discrete DNA fragments (putative Chi-specific fragments)

production. However, the *E. coli* RecD ATPase defective RecBCD enzyme produces the Chi-

specific fragments *in vitro* and confers DNA repair proficiency *in vivo* (31). This converse

observation suggests that the RecD subunit of RecBCD^{Ps} enzyme has a distinct role in DNA repair

and recombination function in *P. syringae*. Despite both RecBCD^{K229Q} and RecB^{K28Q}CD mutant

enzymes being similarly defective in ATP hydrolyzing activity, their DNA unwinding activity was

largely varied. This possibly be due to the selective motor dependency of the enzyme complex,

similar to observation made in *E. coli* RecBCD enzyme (31), in which, the RecB motor is the

absolute requirement for Chi recognition and the motor activity of RecBCD complex.

The RecB^{D1118A}CD enzyme is a processive helicase with no detectable nuclease activity

in vitro, at both 22°C and 4°C. Interestingly, *P. syringae* cells carrying *recB^{D1118A}* allele are capable

of growing at low temperature, suggesting nuclease activity is not essential for its growth at low

temperature (14).

RecF pathway role in RecBCD nuclease defective strain and its importance in *P.

syringae growth at low temperature**

In *E. coli*, the RecF pathway along with RecJ (5'→3' ssDNA specific nuclease) is known

to work with nuclease-defective, RecA loading-deficient RecBCD (32). The role of RecBCD

nuclease defective - RecF hybrid pathway in *P. syringae* is also evidenced. Deletion of *recF* gene

alone (in WT cells) is cold resistant (15). However, deletion of the *recF* gene in *recB^{D1118A}* mutant

renders cold sensitivity (unpublished observation, Apuratha T Pandiyan and Malay K Ray). The

over-expression of RecJ in pRecB$^{\Delta nuc}$CD (RecB nuclease domain deleted) expressing *P. syringae*

recBCD null strain alleviated the slow growth phenotype of *P. syringae* cells at low temperature

(14), suggesting RecJ role in nuclease-defective RecBCD cells. These observations indicate

RecFOR-RecJ role in RecB nuclease-defective *P. syringae* strain.

recA deleted *P. syringae* cells grow slowly at low temperature (15). This suggests that

RecBCD enzyme alone, in the absence of RecA, can rescue low temperature-induced replication

fork arrest possibly by suppressing chromosomal lesions via DNA degradation of reversed

replication fork (33). Interestingly, the combination of a *recA* deletion and a nuclease defective

RecBCD mutation (*recB^{D1118A}CD*) causes cold sensitivity (15), suggesting a direct role of RecA in

rescuing low temperature induced replication fork arrest in a RecB nuclease defective strain.

Therefore, we propose that, in RecB nuclease defective strain, the RecF pathway enables RecA

mediated DNA repair and thus, protects cells from low temperature induced DNA damage.

Identification and characterization of Novel octameric Chi^{Ps} sequence

One of the novel findings in this study is the identification of *P. syringae* Lz4W Chi-

sequence (Chi^{Ps}), 5′-GCTGGCGC-3′. This sequence has not been identified so far in any

Pseudomonas species. Most of the identified bacterial Chi (χ) sequences are GC rich sequences

(Table 3) and the number of nucleotides in χ sites vary from 5-mer (in *Bacillus subtilis*) to 8-mer

(in *E. coli*, *L. lactis* and *H. influenza*) (34). A study using cell lysate from Pseudomonads indicated

that *Pseudomonas* species do not recognize *E. coli* χ sequence (24). We have confirmed this

observation earlier by genetic experiments (14), and in the present study, we have biochemically

identified the Chi[Ps] (5′-GCTGGCGC-3′) sequence; which is identical up to 5 bases from the 5′-

end to the *E. coli* Chi (Chi[Ec]) (5′-GCTGGTGG-3′) sequence. 7-mer (5′-GCTGGCG-3′) sequence,

with change in the last 8[th] nucleotide, are partially recognized. The mutation of 7-mer sequence

to make a perfect 8-mer Chi[Ps] sequence enables it to be strongly recognized by RecBCD[Ps].

Interestingly, appearance of three ssDNA fragments indicates that RecBCD[Ps] enzyme can

sometimes bypass Chi[Ps] and recognize the next putative Chi-like sequence. Similar observations

were made in *E. coli,* where the probability of Chi recognition by *E. coli* RecBCD enzyme and

nicking the DNA is about 30-40% (28, 35).

436   Interestingly, the difference between *E. coli* and *P. syringae* RecBCD enzymes are

confined to the last three nucleotides (5′- GCTGG<u>TGG</u> -3′ vs 5′- GCTGG<u>CGC</u> -3′). The recent

study on the molecular determinants responsible for the Chi recognition by RecBCD enzyme has

revealed the importance of RecC channel in Chi recognition. Among the 35 amino acid residues

of RecC channel examined, the Q38, T40, Q44, L64 W70 D133, L134, D136, D137, R142, R186

and D705 residues of *E. coli* RecC subunit have shown to affect the Chi recognition property of

RecBCD[Ec] enzyme (36). Surprisingly, all these residues are well conserved in the *P. syringae*

RecC subunit. Therefore, the RecC amino acids responsible for recognition of the last three

nucleotides of are still elusive. Further analysis of *E. coli* and *P. syringae* RecC subunits could

shed more insight on the molecular determinants responsible for the recognition of last three

nucleotides of Chi sequence in *E. coli*.

447   *E. coli* contains 1,008 Chi sequences (37). They are four- to eightfold more frequent than

expected by chance and appear on average once every 4.5 kbp. 75% of Chi sites are skewed

towards the replicative leading strand in *E. coli* (37) keeping with their function in stimulating

double-strand break repair upon replication fork collapse. These observations suggest a role for

the RecBCD enzyme as a repair factor functioning towards re-establishment of DNA replication

fork in case of collapse. However, this skewed nature is not applicable for *B. subtilis*, *S. aureus*

and *H. influenzae*, where the skew is statistically insignificant, and the Chi sequence (of the

respective species) is distributed all over the genome (34). The search for Chi[Ps] sequence (5′-

GCTGGCGC-3′) in the draft genome sequence of *P. syringae* Lz4W (4.98 Mb in 42 contigs,

Accession no. <u>**AOGS01000000**</u>) revealed that it contains 1541 Chi[Ps] sequences (5′-

GCTGGCGC-3′) and 4564 of 7-mer Chi[Ps] sequences (5′-GCTGGCG-3′). The Chi[Ps] sequence

appears once in every ~3 kb and is overrepresented compared to other random octamer

sequences. No skewed Chi[Ps] sequence distribution was observed in *P. syringae* Lz4W genome

(A. Pandiyan and M. K. Ray, unpublished observation). Analysis of the closely related *P.*

fluorescens Pf0-1 genome (**Accession no. <u>NC_007492.2</u>**) (38) revealed that it contains 2241

Chi[Ps] sequences and the Chi[Ps] sequence appears once in every 3 kb and, is almost equally

distributed on both strands (1119 Chi-Ps vs. 1122 complementary Chi-Ps sequences). Thus the

pattern of Chi-distribution is not universal, and although Chi[Ps] is over-represented in

Pseudomonas genome, the orientation bias is not observed.

Biochemical properties of RecBCD enzyme and its role in *P. syringae* growth at low

temperature

This study has revealed the biochemical properties of *Pseudomonas* RecBCD enzyme.

The biochemical properties of RecBCD[Ps] enzyme, compared to the RecBCD[Ec], are particularly

associated with the RecD subunit. In *E. coli,* RecD is dispensable for DNA repair process. The

recD null *E. coli* strain is hyper-recombinogenic (39) and RecBC[Ec] enzyme (without RecD)

unwinds dsDNA and loads RecA constitutively in a Chi-independent manner (30). Also, the *E. coli*

RecBCD enzyme with a mutation in the ATP binding site of RecD subunit produces Chi-specific

fragments and cells expressing the mutant enzyme are UV resistant (31). In contrast, *P. syringae*,

RecD is essential for the RecBCD's function (14) and ATP hydrolyzing activity of RecD motor is

an absolute requirement for Chi[Ps] fragments production. Also, cells expressing RecD ATPase

mutant enzyme are cold sensitive, UV and MMC sensitive (13). Importantly, Chi-like octameric

sequence (5′-GCTGGCGC-3′) attenuates nuclease activity of RecBCD*[Ps]* enzyme producing Chi[Ps]

containing ssDNA fragments and thus, can act as a Chi sequence for RecBCD enzyme of

Pseudomonas species.

Based on our results we propose a model (Figure 8) which explains the role of RecBCD

and collaborative DNA repair pathways in rescuing the replication fork arrests in *P. syringae* Lz4W

at low temperature. In this model, RecBCD enzyme can rescue replication fork from the arrest by

linear chromosomal DNA degradation in a *recA*-independent manner. When nuclease activity is

compromised, the nuclease-defective RecBCD enzyme acts with the *recF* pathway to ensure

DNA repair by homologous recombination. In this model, we also propose that motor activity of

RecBCD enzyme is essential for rescuing *P. syringae* cells from low temperature induced

replication fork arrest.

MATERIALS AND METHODS

Bacterial strains, plasmids and growth conditions

The bacterial strains and plasmids used in this study are listed in Tables ST1. The psychrophilic *P. syringae* Lz4W was isolated from a soil sample of Schirmacher Oasis, Antarctica (40) and routinely grown at 22 or 4°C (for high and low temperatures respectively) in Antarctic bacterial medium (ABM) composed of 5 g peptone and 2.0 g yeast extract per liter, as described earlier (16). *E. coli* strains were cultured at 37°C in Luria–Bertani (LB) medium, which contained 10 g tryptone, 5 g yeast extract and 10 g NaCl per liter. For solid media, 15 g bacto-agar (Hi-Media) per liter was added to ABM or LB. When necessary, LB medium was supplemented with ampicillin (100 µg ml^{-1}), kanamycin (50 µg ml^{-1}), gentamicin (15 µg ml^{-1}) or tetracycline (20 µg ml^{-1}) for *E. coli*. For *P. syringae*, the ABM was supplemented with tetracycline (20 µg ml^{-1}), kanamycin (50 µg ml^{-1}) as needed.

pBR322 plasmid DNA (4.3 Kb) and χ+-3F3H dsDNA (a pBR322 derivative, containing two sets of three tandem repeats of χ sequences (27)) were purified using a Qiagen midi kit. Plasmids were linearized with NdeI restriction endonuclease, dephosphorylated with SAP. The dephosphorylated linear dsDNA was 5′- labeled with T4-PNK and [γ-^{32}P] ATP as per the manufacturer guidelines. DNA concentrations were determined by absorbance at 260 nm using molar extinction co-efficient of 6500 M^{-1} cm^{-1} (in nucleotides). All restriction enzymes, DNA ligase T4 Polynucleotide Kinase (T4 PNK), Shrimp alkaline phosphatase (SAP) and *E. coli* SSB were purchased from New England Biolabs (MA, USA). Accuprime Pfx DNA polymerase was purchased from Novagen (WI, USA).

Antibodies and Western analysis

Production of anti-RecB, anti-RecC and anti-RecD antibodies has been described (14). For Western analysis, proteins were separated by SDS-PAGE, transferred onto Hybond C membrane (Amersham Biosciences), and probed with appropriate antibodies. The

immunoreactive protein bands were detected by alkaline phosphatase-conjugated anti-rabbit goat

antibodies (Bangalore Genie, India). For quantification, the blots were scanned with a HP scanjet

and band intensities were measured using Image J software (rsbweb.nih.gov/ij/).

Over expression and purification of recombinant proteins

The LCBD ($\Delta recBCD$) strain harboring pGHCBD, pGHCB^{K28Q}D pGHCBD^{K229Q}

pGHCB^{D1118A}D plasmids (14) were initially grown in 10 ml ABM broth containing kanamycin (50

μg/ml) for 24 hrs at 22°C. 1% of above culture was inoculated into a 2-liter conical flask containing

500 ml ABM broth with kanamycin (50 μg/ml). The culture was then incubated at 22°C with

aeration for 24 hrs. Later, the bacterial cells were harvested by centrifugation at 4°C, 6000 rpm

for 10 min. The bacterial cell pellets were stored at -80°C. The cells pellet was removed as and

when required for the purification of proteins.

All the recombinant proteins in this study were expressed with His-tag on N-terminus of

RecC and purified by nickel-nitrilotriacetic acid (Ni-NTA) affinity chromatography as described in

the manufacturer's protocol (Qiagen, New Delhi, India). In brief, cell lysate of overexpressed

strains containing His-tagged proteins were prepared by dissolving the cell pellet in 10 ml lysis

buffer (50 mM NaH$_2$PO$_4$ (pH 7.4), 300 mM NaCl, 10 mM imidazole and 10% glycerol) and lysed

by mild sonication. The sonicated cell lysate was then centrifuged with 14,000 rpm at 4°C, for 30

min to remove insoluble cellular debris. The supernatant was passed through a pre-equilibrated

column containing 1 ml slurry of Ni-NTA agarose beads and column was allowed to bind Ni-NTA

agarose beads with His-tagged proteins. Further, column was washed with 4-5 volumes of wash

buffer (50 mM NaH$_2$PO$_4$ (pH 7.4), 300 mM NaCl, 20 mM imidazole, and 10% glycerol). Finally,

the bound proteins were eluted with 1-2 ml elution buffer (50 mM NaH$_2$PO$_4$ (pH 7.4), / 300 mM

NaCl / 300 mM imidazole / 10% glycerol).

Gel filtration technique (size-exclusion column chromatography) was employed for the

purification of His-tagged RecBCD complex and the mutant protein complexes. For this,

Superose-12 gel filtration column (Pharmacia Fine Chemicals) was used in the fast protein liquid

chromatography (FPLC) (Pharmacia Fine Chemicals). Initially, the column was pre-equilibrated

with buffer contained 20 mM Tris HCl (pH 7.5), 0.1 mM EDTA, 150 mM NaCl, 0.1 mM PMSF and

10% glycerol. Later, 0.5 ml of Ni-NTA purified protein solution was injected to the column and

allowed to pass through the column at a flow rate of 0.4 ml/min. Optical density at 280 nm was

recorded for the eluted protein fractions and the fractions were collected in separate

microcentifuge tubes. The protein fractions were then analyzed on SDS-PAGE stained with

coomassie brilliant blue or silver nitrate. The gel filtration protein fractions of interest were further

membrane dialyzed in 50% glycerol containing gel filtration buffer. RecBCD enzyme

concentrations were determined by measuring OD_{280} and using molar extinction co-efficient 4.7 X

10^5 M^{-1} cm^{-1} as determined by ExPASy – ProtParam tool (https://web.expasy.org/protparam).

Thin-layer chromatography based assay

ATPase activity of RecBCD and mutant proteins was assayed at different temperatures

by thin layer chromatography (TLC) on polyethylene-imine (PEI)-cellulose plates (E-Merck,

Germany). The assay was performed as described earlier (41, 42). Assays were carried out at

indicated temperatures in a reaction volume of 20 μl containing 25 mM Tris acetate (pH 7.5), 1

mM Mg acetate, 1 mM DTT, 100mM NdeI linearized pBR322-dsDNA and 200 μM ATP as a

substrate with 2 nM RecBCD and mutant enzymes. One μl of 100 times diluted 10 mCi ml^{-1} stock

of [γ-^{32}P] ATP (specific activity 3000 Ci $mmol^{-1}$) was used as a tracer in each reaction to measure

the rate of ATP hydrolysis. Following 0, 1, 2, 3, 5 10 minutes of reaction, 0.5 μl aliquots of the

samples were spotted on TLC plate, air-dried and were allowed to develop in a mobile phase

containing 0.5 M formic acid and 0.5 M lithium chloride for 15 minutes. The TLC plate was dried

and exposed to the Phosphor imaging plate for 4-6 hrs. The Imaging plates were scanned in a

Phosphor Imager, and the amounts of ^{32}Pi and [γ-^{32}P] ATP were quantified using Image gauge

software (Fuji-3000). Further, data were analyzed using GraphPad Prism 4.0 software.

DNA unwinding assay

Plasmid DNAs were linearized with appropriate restriction enzymes in the presence of shrimp alkaline phosphatase and 5`-end-labeled by T4 polynucleotide kinase and [γ-^{32}P] ATP. Subsequent purification of labeled DNA was accomplished by passage through a MicroSpin S-200 HR column (Amersham biosciences-GE healthcare, Buckinghamshire, UK). The reaction mixtures contained 25 mM Tris acetate (pH 7.5), 2 mM magnesium acetate (as indicated), 1 mM DTT, 10 μM (nucleotides) linear pBR322 dsDNA P^{32}-labeled at 5′- end, 5 mM ATP, 2 μM *E. coli* SSB protein and 0.5 nM RecBCD^{Ps} or mutant enzymes. DNA unwinding reactions were started with the addition of either enzyme or ATP, after pre-incubation of all other components at 22 or 4°C for 5 min. Assays were stopped at the indicated times by addition of proteinase K to a final concentration of 0.5 mg/ml, which was dissolved in sample loading buffer (125 mM EDTA, 40% glycerol, 2.5% SDS, 0.25% bromophenol blue, and 0.25% xylene cyanol). After 5-min incubation with proteinase K at room temperature, the reaction products were run on a 1% (w/v) agarose gel in a 1X TBE (45 mM Tris borate (pH 8.3) and 1 mM EDTA) buffer at 25-30 constant volts for 15 hrs. Agarose gels were dried, exposed to phosphor imaging plates and quantified using Phosphor Imager (Fuji-3000). Further, data were analyzed using image gauge software.

The DNA unwinding rates of were measured by using the following formula

$$\text{nM bp s}^{-1}(\text{nM helicase})^{-1} = \text{slope X (C/ [100\%]) X } t_s \text{ X (1/E)}$$

Where C is the concentration of linear dsDNA substrate in base-pairs in nM (i.e., 5000 nM), t_s is the time in seconds, and E is the enzyme concentration in nM.

DNA degradation assay

The assays were performed as described above in DNA unwinding assay, except that the reaction mixtures contained 6 mM magnesium acetate and 2 mM ATP.

Single-stranded DNA endonuclease assay

The endonuclease activity of RecBCD enzyme on ssDNA was examined using a circular

M13 ssDNA substrate as described previously (20, 22). In brief, endonuclease activity was tested

in 3 different buffer conditions. The first reaction mixture contained 50 mM MOPS (pH=7.5), 1 mM

ATP, 10 mM $MgCl_2$, 4.16 nM circular M13 ssDNA with 0.5 nM RecBCD. The second reaction

mixture contained 25 mM Tris-acetate, 1 mM ATP, 8 mM Mg-acetate,1 mM DTT, 4.16 nM M13

ssDNA with 0.5 nM RecBCD. The third reaction mixture contained 25 mM Tris-acetate, 2 mM

ATP, 6 mM Mg-acetate, 1 mM DTT, 4.16 nM M13 ssDNA with 0.5 nM RecBCD. After the reaction,

samples were removed at the indicated times; quenched with 120 mM EDTA, 40% (v/v) glycerol,

and 0.125% bromphenol blue; and loaded on a 0.8% agarose gel in 1X TBE (90 mM Tris borate,

2 mM EDTA). The gel was run at 4 V/cm for 3 h and stained with ethidium bromide (0.5 mg/ml).

The bands were visualized by exposure to UV light.

Site directed deletion of regions from plasmid pBR322

Different internal regions of plasmid pBR322 were deleted using site directed deletion. In short,

primers were designed consists of sequences flanking the region to be deleted. PCR was

performed to amplify whole Plasmid DNA using Accuprime Pfx DNA polymerase (Invitrogen).

After PCR, reaction mixture was subjected for the overnight DpnI digestion. 5 µl of reaction

mixture was then transformed into DH5α ultra-competent cells. All selected regions, which had to

be deleted was in tetracycline resistance gene of the plasmid. Therefore, for primary screening

only those colonies were selected which were Amp[R] and Tet[S]. Deletion was further confirmed by

PCR and sequencing. All primer list and corresponding deleted regions are shown in Table ST2.

ACCESSION NUMBERS

Accession no. AOGS01000000, the draft genome sequence of *P. syringae* Lz4W and,

Accession no. NC_007492.2, the genome sequence *Pseudomonas fluorescens* Pf0-1.

SUPPLIMENTARY MATERIALS

Supplementary material related to this manuscript is attached.

ACKNOWLEDGEMENTS

Research in M.K.R. laboratory is supported by the Council of Scientific and Industrial Research

(CSIR), Government of India. A part of the work was supported by a grant to M.K.R. from

Department of Science and Technology (DST), Government of India. T.L.P and A.K.S.

acknowledge CSIR, India for research fellowships.

We also thank Prof. Stephen C Kowalczykowski, Prof. Benedicte Michel and Dr. Naofumi Handa

for their critical reading and comments on the manuscript.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest with the contents of this article.

AUTHOR CONTRIBUTIONS

Theetha L. Pavankumar & Anurag Kumar Sinha both have contributed equally to this work.

T.L.P., A.K.S. and M.K.R designed the experiments. T.L.P. and A.K.S. conducted the experiment.

T.L.P., A.K.S. and M.K.R analyzed the results and wrote the manuscript.

 FIGURE LEGENDS

FIGURE 1. Purification of His-tagged RecBCD complex and mutant enzymes by Ni-NTA

agarose column chromatography. (A) Steps involved in purification of wild-type RecBCD and

mutant proteins. **(B)** SDS-PAGE analysis of purified RecBCD protein fractions of wild-type and

mutants. Purified protein fractions were stained either with coomassie brilliant blue (B) or with

silver nitrate (C). Three protein bands of expected size corresponding to RecB, RecC and RecD

are visible on the gel. A low molecular protein (~60 kDa) observed on silver nitrate strained gel

was identified as GroEL, a HSP-60 family chaperonin.

FIGURE 2. ATP hydrolysis of wild-type and mutant RecBCD enzymes at different

temperatures. The assays were carried out by TLC method as described in Methods. **(A)** A graph

showing the concentration dependent ATP hydrolysis by RecBCD (WT) and mutant enzymes at

$22°C$. The inset in B shows a blow-up of the same data of mutant enzymes using an expanded y-

scale. **(B)** Representative of TLC plates showing the ATP hydrolysis by wild-type and mutant

RecBCD enzymes at $37°$, $22°$ and $4°C$. **(C)** ATPase activities at $37°$, $22°$ and $4°C$ are shown in

histogram with error bars. The ATP hydrolysis data presented in the histogram represents the

results obtained from three independent experiments.

FIGURE 3. Effects of magnesium and ATP on the unwinding and degradation by wild-type

RecBCD enzyme of *P. syringae*. The DNA unwinding and degradation assays were performed

with [5'-^{32}P] labeled *Nde*I linearized pBR322 plasmid DNA in the presence of different

concentrations of magnesium and ATP, as described in Methods; **(A)** the reaction mixture

contained fixed amount of Mg-acetate (2 mM) and the amount ATP was varied (0 to 10 mM) as

indicated. **(B)** The reaction mixture contained the fixed amount of ATP (2 mM) and concentration

of Mg-acetate was varied (0 to 10 mM) as indicated. The reactions were performed for 5 min and

analyzed on a 1% agarose gel containing 1X TBE buffer at a 25-30 Volts for 15 hrs. Agarose gels

were dried, exposed to phosphor imaging plates and quantified using Phosphor Imager (Fuji-

3000). The data were analyzed using image gauge software. The lanes C_1 and C_2 contain [5'-^{32}P]

labeled dsDNA and the heat denatured 5'-^{32}P labeled dsDNA respectively as a control. A [5'-^{32}P]

labeled discrete DNA bands of smaller than full-length ssDNA of pBR322 were also noticed in the

lanes 5-10 of panel B.

FIGURE 4. DNA unwinding and degradation of *Nde*I digested linear dsDNA of pBR322 by

RecBCD^{Ps} (WT) enzyme at 22° and 4°C. The DNA unwinding and degradation at 22° and 4°C

were carried out as described in Methods. **(A)** The DNA unwinding reactions contained 5 mM

ATP and 2 mM Mg^{++} (limiting magnesium condition) **(B)** The DNA degradation reactions contained

2 mM ATP and 6 mM Mg^{++} (excess magnesium condition). The reactions were initiated by adding

ATP, and stopped at the indicated times by adding stop-buffer. The lanes C_1 and C_2 contain [5'-

^{32}P] labeled *Nde*I linearised double-stranded and the heat-denatured ssDNA of pBR322 as

control. The discrete ssDNA fragments of pBR322 produced by the nuclease activity of RecBCD

enzyme are also indicated.

FIGURE 5. DNA unwinding and degradation of *Nde*I linearized dsDNA of pBR322 by mutant

RecBCD^{Ps} enzymes at 22° and 4°C. (A) DNA unwinding and degradation by RecB^{K28Q}CD

enzyme. The DNA unwinding and degradation at 22° and 4°C were carried out as described in

Methods. Note that this enzyme shows DNA unwinding (ssDNA production), but no detectable

DNA degradation at 22°C, and none were detectable at 4°C. **(B) DNA unwinding and**

degradation by RecBCD^{K229Q} enzyme. The RecBCD^{K229Q} enzyme has apparently retained both

the DNA unwinding and degradation properties. But, interestingly, the discrete DNA bands are

absent **(C) DNA unwinding by nuclease-deficient RecB^{D1118A}CD enzyme under limiting and**

excess magnesium conditions at 22 and 4°C. RecB^{D1118A}CD enzyme is unable to degrade

DNA at both 22 and 4°C and notably, the DNA unwinding seems to be faster in excess-

magnesium reaction condition (2 mM ATP:6 mM Mg^{++}) compared to limiting magnesium reaction

condition (5 mM ATP:2 mM Mg^{++}). Each reaction mixtures contained 0.5 nM of enzyme and 10

μM (nucleotides) linear [5'-^{32}P] labeled pBR322 dsDNA. The lanes C_1 and C_2 contain [5'-^{32}P]

labeled *Nde*I linearized double-stranded and the heat-denatured ssDNA of pBR322 respectively.

FIGURE 6. Identification of Chi sequence using PCR amplified pBR322 fragment (3.6 kb)

containing internal deletions as a DNA substrate. (A) Top panel; Schematic representation of

NdeI linearized pBR322 plasmid DNA. The locations of OROPI and OPROPII primers used in

PCR amplification of 3.6 kb fragments are indicated. Bottom panel; DNA degradation reactions

performed using the PCR amplified 3.6 kb DNA as substrate, having either bottom strand labeled

(left panel) or top strand labeled (middle panel) or both strand labeled (right panel). The lane C

contains [5'-^{32}P] labeled heat-denatured ssDNA as a control. Notably, the top strand labeling of

DNA substrate resulted in appearance of discrete DNA fragments. **(B)** Deletion of 400-450 bp of

pBR322 (pBR322$_{(Δ400-450)}$) resulted in disappearance of DNA fragments. But, it is clearly visible

when intact 3.6 kb pBR322 was used as a substrate, suggesting the presence of putative Chi^{Ps}

sequence in this region. **(C)** Further deletion of 401-419; 421-439 bo regions; and, **(D)** 421-429;

431-439; 441-449 bp regions of pBR322 shows that the DNA fragments are indeed from the 431-

439 bp region of the pBR322 DNA sequence. **(E)** A schematic representation of 3.6 kb region of

pBR322 and deleted regions within, are shown in the left panel. Right panel shows the presence

(**+**) or absence (**-**) of intense protected band (Chi-like fragments), when these constructs are used

as assay substrates.

FIGURE 7. (A) DNA degradation of modified fragment of pBR322 by RecBCD^Ps enzyme.

Apart from one 8-mer Chi^Ps sequence at 431th position of pBR322, there are two 7-mer similar

sequences present in the plasmid pBR322. One such similar sequence at 964^th position (5′

GCTGGCG**T** 3′) was mutated to make it an 8-mer sequence (5′ GCTGGCG**C** 3′). This substrate

(pBR322^T971C) that has two Chi sequences in the same orientation when used as a substrate,

yielded two DNA fragments with high intensity. **(B) The DNA degradation pattern of *Xba*I**

digested linear dsDNA of pBKS plasmid by RecBCD^Ps enzyme. Chi^Ps are inserted into

pBluescript vector (pBKS) by site directed insertion. *Xba*I linearized pBKS vector or pBKS with

Chi^Ps was 5'-end labeled with ^32P has been used for assays. The discrete ssDNA of expected size

was produced when pBKS (Chi^Ps) was used as a substrate, whereas it is absent, when native

pBKS was used.

FIGURE 8. Role of RecBCD dependent DNA repair pathway in rescue of low temperature

induced replication forks arrest. (i) A chromosome replicating bi-directionally, (ii) encounters

low temperature-induced chromosomal lesion or blockage, (iii), causing replication fork arrest

and fork reversal (RFR). RFR is suppressed by linearized chromosomal DNA degradation by

RecBCD enzyme and resetting of replication fork. (iv) RFR is stabilized further by RuvAB

complex and, (v) further, resolved by RuvC leading to chromosomal linearization. Linearized

chromosome is processed by either RecBCD alone or in conjunction (when RecBCD is

nuclease defective) with recFOR-recJ hybrid pathway. The defective motors activity of RecBCD

enzyme leads to chromosomal fragmentation and cell death at the low temperature.

REFERENCES

1.   Cromie GA, Connelly JC, Leach DR. Recombination at double-strand breaks and DNA
ends: conserved mechanisms from phage to humans. Mol Cell. 2001;8(6):1163-74.
2.   Dillingham MS, Kowalczykowski SC. RecBCD enzyme and the repair of double-
stranded DNA breaks. Microbiol Mol Biol Rev. 2008;72(4):642-71, Table of Contents.
3.   Kowalczykowski SC, Dixon DA, Eggleston AK, Lauder SD, Rehrauer WM.
Biochemistry of homologous recombination in Escherichia coli. Microbiol Rev.
1994;58(3):401-65.
4.   Michel B, Boubakri H, Baharoglu Z, LeMasson M, Lestini R. Recombination proteins
and rescue of arrested replication forks. DNA Repair (Amst). 2007;6(7):967-80.
5.   Kuzminov A. Single-strand interruptions in replicating chromosomes cause double-
strand breaks. Proc Natl Acad Sci U S A. 2001;98(15):8241-6.
6.   Taylor AF, Smith GR. Strand specificity of nicking of DNA at Chi sites by RecBCD
enzyme. Modulation by ATP and magnesium levels. J Biol Chem. 1995;270(41):24459-67.
7.   Stahl FW, Stahl MM. Recombination pathway specificity of Chi. Genetics.
1977;86(4):715-25.
8.   Anderson DG, Kowalczykowski SC. The recombination hot spot chi is a regulatory
element that switches the polarity of DNA degradation by the RecBCD enzyme. Genes Dev.
1997;11(5):571-81.
9.   Anderson DG, Kowalczykowski SC. The translocating RecBCD enzyme stimulates
recombination by directing RecA protein onto ssDNA in a chi-regulated manner. Cell.
1997;90(1):77-86.
10.   Hiom K, West SC. Branch migration during homologous recombination: assembly of
a RuvAB-Holliday junction complex in vitro. Cell. 1995;80(5):787-93.
11.   Pavankumar TL, Exell JC, Kowalczykowski SC. Direct Fluorescent Imaging of
Translocation and Unwinding by Individual DNA Helicases. Methods Enzymol. 2016;581:1-
32.
12.   Spies M, Bianco PR, Dillingham MS, Handa N, Baskin RJ, Kowalczykowski SC. A
molecular throttle: the recombination hotspot chi controls DNA translocation by the
RecBCD helicase. Cell. 2003;114(5):647-54.
13.   Spies M, Amitani I, Baskin RJ, Kowalczykowski SC. RecBCD enzyme switches lead
motor subunits in response to chi recognition. Cell. 2007;131(4):694-705.
14.   Pavankumar TL, Sinha AK, Ray MK. All three subunits of RecBCD enzyme are
essential for DNA repair and low-temperature growth in the Antarctic Pseudomonas
syringae Lz4W. PLoS One. 2010;5(2):e9412.
15.   Sinha AK, Pavankumar TL, Kamisetty S, Mittal P, Ray MK. Replication arrest is a
major threat to growth at low temperature in Antarctic Pseudomonas syringae Lz4W. Mol
Microbiol. 2013;89(4):792-810.
16.   Regha K, Satapathy AK, Ray MK. RecD plays an essential function during growth at
low temperature in the antarctic bacterium Pseudomonas syringae Lz4W. Genetics.
2005;170(4):1473-84.
17.   Handa N, Bianco PR, Baskin RJ, Kowalczykowski SC. Direct visualization of RecBCD
movement reveals cotranslocation of the RecD motor after chi recognition. Mol Cell.
2005;17(5):745-50.

18. Dixon DA, Kowalczykowski SC. Role of the Escherichia coli recombination hotspot,
chi, in RecABCD-dependent homologous pairing. J Biol Chem. 1995;270(27):16360-70.
19. Rosamond J, Telander KM, Linn S. Modulation of the action of the recBC enzyme of
Escherichia coli K-12 by Ca2+. J Biol Chem. 1979;254(17):8646-52.
20. Eggleston AK, Kowalczykowski SC. Biochemical characterization of a mutant recBCD
enzyme, the recB2109CD enzyme, which lacks chi-specific, but not non-specific, nuclease
activity. J Mol Biol. 1993;231(3):605-20.
21. Sun JZ, Julin DA, Hu JS. The nuclease domain of the Escherichia coli RecBCD enzyme
catalyzes degradation of linear and circular single-stranded and double-stranded DNA.
Biochemistry. 2006;45(1):131-40.
22. Yu M, Souaya J, Julin DA. The 30-kDa C-terminal domain of the RecB protein is
critical for the nuclease activity, but not the helicase activity, of the RecBCD enzyme from
Escherichia coli. Proc Natl Acad Sci U S A. 1998;95(3):981-6.
23. Dixon DA, Kowalczykowski SC. The recombination hotspot chi is a regulatory
sequence that acts by attenuating the nuclease activity of the E. coli RecBCD enzyme. Cell.
1993;73(1):87-96.
24. McKittrick NH, Smith GR. Activation of Chi recombinational hotspots by RecBCD-like
enzymes from enteric bacteria. J Mol Biol. 1989;210(3):485-95.
25. Lehman IR, Nussbaum AL. The Deoxyribonucleases of Escherichia Coli. V. On the
Specificity of Exonuclease I (Phosphodiesterase). J Biol Chem. 1964;239:2628-36.
26. Dixon DA, Churchill JJ, Kowalczykowski SC. Reversible inactivation of the
Escherichia coli RecBCD enzyme by the recombination hotspot chi in vitro: evidence for
functional inactivation or loss of the RecD subunit. Proc Natl Acad Sci U S A.
1994;91(8):2980-4.
27. Anderson DG, Churchill JJ, Kowalczykowski SC. A single mutation, RecB(D1080A,)
eliminates RecA protein loading but not Chi recognition by RecBCD enzyme. J Biol Chem.
1999;274(38):27139-44.
28. Taylor AF, Schultz DW, Ponticelli AS, Smith GR. RecBC enzyme nicking at Chi sites
during DNA unwinding: location and orientation-dependence of the cutting. Cell.
1985;41(1):153-63.
29. Amundsen SK, Taylor AF, Smith GR. The RecD subunit of the Escherichia coli
RecBCD enzyme inhibits RecA loading, homologous recombination, and DNA repair. Proc
Natl Acad Sci U S A. 2000;97(13):7399-404.
30. Churchill JJ, Anderson DG, Kowalczykowski SC. The RecBC enzyme loads RecA
protein onto ssDNA asymmetrically and independently of chi, resulting in constitutive
recombination activation. Genes Dev. 1999;13(7):901-11.
31. Spies M, Dillingham MS, Kowalczykowski SC. Translocation by the RecB motor is an
absolute requirement for {chi}-recognition and RecA protein loading by RecBCD enzyme. J
Biol Chem. 2005;280(44):37078-87.
32. Ivancic-Bace I, Peharec P, Moslavac S, Skrobot N, Salaj-Smic E, Brcic-Kostic K.
RecFOR function is required for DNA repair and recombination in a RecA loading-deficient
recB mutant of Escherichia coli. Genetics. 2003;163(2):485-94.
33. Miranda A, Kuzminov A. Chromosomal lesion suppression and removal in
Escherichia coli via linear DNA degradation. Genetics. 2003;163(4):1255-71.
34. El Karoui M, Biaudet V, Schbath S, Gruss A. Characteristics of Chi distribution on
different bacterial genomes. Res Microbiol. 1999;150(9-10):579-87.

35.   Arnold DA, Bianco PR, Kowalczykowski SC. The reduced levels of chi recognition
exhibited by the RecBC1004D enzyme reflect its recombination defect in vivo. J Biol Chem.
1998;273(26):16476-86.
36.   Handa N, Yang L, Dillingham MS, Kobayashi I, Wigley DB, Kowalczykowski SC.
Molecular determinants responsible for recognition of the single-stranded DNA regulatory
sequence, chi, by RecBCD enzyme. Proc Natl Acad Sci U S A. 2012;109(23):8901-6.
37.   Arakawa K, Uno R, Nakayama Y, Tomita M. Validating the significance of genomic
properties of Chi sites from the distribution of all octamers in Escherichia coli. Gene.
2007;392(1-2):239-46.
38.   Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FS. Enhanced
annotations and features for comparing thousands of Pseudomonas genomes in the
Pseudomonas genome database. Nucleic Acids Res. 2016;44(D1):D646-53.
39.   Lovett ST, Luisi-DeLuca C, Kolodner RD. The genetic dependence of recombination
in recD mutants of Escherichia coli. Genetics. 1988;120(1):37-45.
40.   Shivaji S, Rao NS, Saisree L, Sheth V, Reddy GS, Bhargava PM. Isolation and
identification of Pseudomonas spp. from Schirmacher Oasis, Antarctica. Appl Environ
Microbiol. 1989;55(3):767-70.
41.   Korangy F, Julin DA. Alteration by site-directed mutagenesis of the conserved lysine
residue in the ATP-binding consensus sequence of the RecD subunit of the Escherichia coli
RecBCD enzyme. J Biol Chem. 1992;267(3):1727-32.
42.   Kornberg A, Scott JF, Bertsch LL. ATP utilization by rep protein in the catalytic
separation of DNA strands at a replicating fork. J Biol Chem. 1978;253(9):3298-304.
43.   Smith GR, Comb M, Schultz DW, Daniels DL, Blattner FR. Nucleotide sequence of the
chi recombinational hot spot chi +D in bacteriophage lambda. J Virol. 1981;37(1):336-42.
44.   Chedin F, Noirot P, Biaudet V, Ehrlich SD. A five-nucleotide sequence protects DNA
from exonucleolytic degradation by AddAB, the RecBCD analogue of Bacillus subtilis. Mol
Microbiol. 1998;29(6):1369-77.
45.   Halpern D, Chiapello H, Schbath S, Robin S, Hennequet-Antier C, Gruss A, et al.
Identification of DNA motifs implicated in maintenance of bacterial core genomes by
predictive modeling. PLoS Genet. 2007;3(9):1614-21.
849

850 **TABLE 1. ATPase activity of wild type and mutant RecBCD enzymes**

Enzyme	37°C	22°C	4°C	$^a K_m$	$^a k_{cat}$
RecBCD (WT)	366.7 ± 30.9	236.5 ± 9.4	119.0 ± 19.6	57.8 ± 10.1	61.3
RecB^{D1118A}CD	346.0 ± 33.8	209.8 ± 13.9	112.4 ± 23.3	59.5 ± 8.5	55.3
RecB^{K28Q}CD	33.3 ± 0.9	16.2 ± 3.0	9.7 ± 1.5	58.3 ± 11.0	6.6
RecBCD^{K229Q}	30.5 ± 1.4	26.9 ± 2.2	9.7 ± 1.7	49.4 ± 5.8	6.1

851 ATPase activities are expressed here in units of µmol ATP hydrolyzed per second per µmol

852 RecBCD enzymes. ^{a} K_m and k_{cat} values are calculated from ATPase activity at 22 °C.

854 **TABLE 2. DNA unwinding and degradation activities of the wild type and mutant RecBCD**

855 **enzymes at 22 and 4°C**

Enzyme	Rate of DNA unwinding at 22°C (bp s^{-1})	Rate of DNA unwinding at 4°C (bp s^{-1})	Rate of DNA unwinding [a] or degradation at 22°C (bp s^{-1})	Rate of DNA unwinding [a] or degradation at 4°C (bp s^{-1})
	ATP > Mg^{++} (5:2 mM)	ATP > Mg^{++} (5:2 mM)	ATP < Mg^{++} (2:6 mM)	ATP < Mg^{++} (2:6 mM)
RecBCD^{Ps}	35.1 ± 1.6	ND [b]	101.5 ± 3.3	55.8 ± 6.9
RecB^{D1118A}CD	30.4 ± 1.7	ND	109.7 ± 17.5	41.0 ± 8.1
RecB^{K28Q}CD	1.5 ± 0.76	ND	12.9 ± 3.4	ND
RecBCD^{K229Q}	27.2 ± 6.1	ND	92.6 ± 14.1	17.3 ± 9.6

856 The values shown for the DNA unwinding/degradation rates with mean ± S.E. from two

857 independent experiments. ^{a} Rate of DNA unwinding under excess magnesium reaction condition.

858 ^{b} ND - not determined

860 **TABLE 3. Chi sequences identified in different bacteria.**

Bacteria	Chi sequence	Recombination machinery
Escherichia coli	5′ GCTGGTGG 3′	RecBCD (43)
Bacillus subtilis	5′ AGCGG 3′	AddAB (44)
Lactococcus lactis	5′ GCGCGTG 3′	RexAB (34)
Streptococcus pneumoniae	5′ GCGCGTG 3′	RexAB (45)
Staphylococcus aureus	5′ GAAGCGG 3′	- (45)
Streptococcus agalactiae	5′ GCGCGTG 3′	- (45)
Streptococcus thermophilus	5′ GCGCGTG 3′	- (45)
***Pseudomonas syringae* Lz4W**	**5′ GCTGGCGC 3′**	RecBCD (14)

861
862

A

Overexpression of RecBCD
and mutant proteins in
LCBD strain (ΔrecCBD::Tet)

↓

Cell lysis by sonication

↓

Ni-NTA- affinity
column purification

↓

Superose-12
column gel filtration

B

RecB^{His}-CD (WT) RecB^{K28Q}His-CD RecB^{D1118A} His-CD RecB^{His}-CD^{K229Q}

RecB
^{His}-RecC
RecD

C

RecB^{His}-CD (WT) RecB^{K28Q}His-CD RecB^{D1118A} His-CD RecB^{His}-CD^{K229Q}

RecB
^{His}-RecC
RecD
GroEL

Figure 1

A

B

C

Figure 2

A

ATP	-	-	0	1	2	3	4	5	6	7	8	10 mM
Mg²⁺	-	-	2	2	2	2	2	2	2	2	2	2 mM
Lane #	C₁	C₂	1	2	3	4	5	6	7	8	9	10

ssDNA

B

Mg²⁺	-	-	0	1	2	3	4	5	6	7	8	10 mM
ATP	-	-	2	2	2	2	2	2	2	2	2	2 mM
Lane #	C₁	C₂	1	2	3	4	5	6	7	8	9	10

ssDNA

Discrete
DNA
fragments

Figure 3

A

Limiting magnesium condition
ATP (5 mM) : Mg^{++} (2 mM)

ssDNA

B

Excess magnesium condition
ATP (2 mM) : Mg^{++} (6 mM)

ssDNA

Figure 4

Figure 5

Figure 6

A

dsDNA (3650 bp)

+
ssDNA

prominent
DNA
fragments

B

dsDNA

+
ssDNA

DNA
fragment

Figure 7

RecB^{D1118A}CD (helicase), RecJ-RecFOR hybrid pathway

RecBCD pathway

(i) Replicating chromosome

Low temperature induced DNA damage and replication arrest

(ii)

(iii) RF reversal (RFR)

Linear chromosomal DNA degradation by RecBCD enzyme and Replication fork resetting

(iv) RecBCD dependent degradation of RFR

(v) HJ resolution and linearization of chromosome

Defective motor activity of RecBCD enzyme

(vi) **Chromosome fragmentation Cell death and cold sensitivity**

RecBCD   RuvAB   RuvC

Figure 8