Constriction rate modulation can drive cell size control and homeostasis in *C.

*crescentus***

Ambroise Lambert[1]*, Aster Vanhecke[1]*, Anna Archetti[1], Seamus Holden[1,2], Felix

Schaber[1], Zachary Pincus[3,4], Michael T. Laub[5,6], Erin Goley[7], Suliana Manley[1]

1 Institute of Physics, École Polytechnique Fédérale de Lausanne (EPFL),

1015 Lausanne, Switzerland

2 Centre for Bacterial Cell Biology, Institute for Cell and Molecular Biosciences,

Newcastle University, Newcastle upon Tyne NE2 4AX, UK

3 Department of Genetics, Washington University in St Louis, St. Louis, MO, USA

4 Department of Developmental Biology, Washington University in St Louis, St.

Louis, MO, USA

5 Department of Biology, Massachusetts Institute of Technology, Cambridge,

MA, USA

6 Howard Hughes Medical Institute, Massachusetts Institute of Technology,

Cambridge, MA, USA

7 Department of Biological Chemistry, Johns Hopkins University School of

Medicine, Baltimore, MD, USA.

Author contributions: *These authors contributed equally.

Lead contact: suliana.manley@epfl.ch

<u>Abstract</u>

Rod-shaped bacteria typically grow first via sporadic and dispersed elongation along

their lateral walls, then via a combination of zonal elongation and constriction at the

division site to form the poles of daughter cells. Although constriction comprises up to

half of the cell cycle, its impact on cell size control and homeostasis has rarely been

considered. To reveal the roles of cell elongation and constriction in bacterial size

regulation during cell division, we captured the shape dynamics of *Caulobacter*

crescentus with time-lapse structured illumination microscopy and used molecular

markers as cell-cycle landmarks. We perturbed constriction rate using a

hyperconstriction mutant or fosfomycin inhibition. We report that constriction rate

contributes to both size control and homeostasis, by determining elongation during

constriction, and by compensating for variation in pre-constriction elongation on a

single-cell basis.

Introduction

Cell size regulation is observed nearly universally among prokaryotes [1], allowing them to both control their size at birth and to homeostatically maintain it over multiple generations [2]. Cell size control and homeostasis are critical for survival: once too small, cells lack the volume required to host the essential machinery of life [3] or initiate chromosome segregation [4], whereas cells that are too large may suffer limitations in nutrient uptake [5] and distribution [6] because of their reliance on diffusive transport.

Size regulation is the outcome of multiple sequential processes that take place during the cell division cycle. Progression through the cell cycle is marked by several key processes, including chromosome replication, segregation, and division into two daughter cells. These processes occur once per cell cycle in bacteria such as *Caulobacter crescentus* [7], in contrast to rapidly proliferating organisms such as *Escherichia coli* [8] and *Bacillus subtilis* in which cells often have multi-fork replication and which can, following nutrient up-shifts, initiate replication multiple times in a single cell cycle. In *C. crescentus*, differentiation from a swarmer to a stalked cell and the initiation of chromosome replication and segregation mark the transition from cell cycle phase G1 to S. The completion of replication marks the end of S phase. Once DNA segregation is completed, cells finish cytokinesis to form sibling stalked and swarmer cells during G2/M [9].

From the perspective of the cell wall, individual *C. crescentus* cells elongate exponentially throughout the cell cycle, as is typical for rod-shaped bacteria. Their growth is divided into a stage of pure elongation as peptidoglycan is inserted sporadically along the lateral walls, followed by a stage of mixed elongation and constriction in G2/M phase during which peptidoglycan is inserted at mid-cell to build two new poles [10,11]. Each stage governed by largely independent molecular machines. In *B. subtilis*, when cells are on average longer at the onset of constriction, they are also on average longer at division [12,13]. This suggests a simple model for cell size control, by modifying the cell length at which the divisome, the multi-protein complex which guides division, begins to generate constriction. Another possibility is

that the rate of constriction is modulated; this was shown to be the case for MatP, which

coordinates chromosome segregation and pole construction in *E. coli* [14].

Several checkpoints have been proposed to couple bacterial cell size homeostasis to

cell cycle phases. Shortly after entry into S phase, chromosome segregation must initiate

before the cytokinetic Z-ring can assemble at mid-cell, coordinated by the gradient-forming

FtsZ inhibitor MipZ [15]. In *E. coli*, the initiation of chromosome replication requires a fixed

volume per origin of replication, leading to a "sizer" [16]. Chromosome replication in G1/S may

take a constant duration, referred to as a "timer" [8]. Putative molecular mechanisms have

generally relied on the accumulation of proteins above a threshold, such as an "initiator"

triggering replication [17], or excess peptidoglycan (PG) cell wall precursors triggering

constriction [18]. A model of the latter case predicts a constant addition of volume per cell

cycle, or "adder". Indeed, an adder has been observed for *C. crescentus* under a wide range

of growth conditions [19]. Deviations from a pure adder toward a mixed relative timer and adder

have also been reported for stalked cells, observed over many generations and a range of

different temperatures [20]. Any model incorporating a sizer or adder will allow smaller cells to

increase while larger cells decrease in size over generations until both converge to a size set

by the constant of addition [21]. Thus, both provide a clear means for a population to achieve

size homeostasis.

Remarkably, although constriction makes up a significant portion of the cell cycle in

many bacteria [22], its impact on cell size control and homeostasis has rarely been considered.

Intriguingly, budding yeast may use constriction rate to modulate their size in response to

changes in growth conditions [23]. However, a single-cell study of the contribution of the

constriction stage in bacteria has been challenging, in part due to the diffraction-limited size of

the constriction site, and the need for corroboration by divisome markers to unambiguously

identify constriction onset. Furthermore, direct measurement of the instantaneous constriction

rate has not been possible.

Here, we investigated whether, and how cells adjust their constriction rate to achieve

cell size control and homeostasis. We used structured illumination microscopy (SIM) [24] to

resolve the constriction site diameter and measure the size of synchronized *C. crescentus* cells

as they progressed through their cell cycle. We show that perturbing the constriction rate

changes cell size, independent of elongation rate. Furthermore, we found that within a

population the onset of constriction and its rate are coordinated: cells that elongate more than

average before constriction undergo a more rapid constriction, leading to less elongation

during constriction, and vice-versa. This compensation leads to a higher fidelity adder than

permitted by onset control alone, allowing *C. crescentus* to better maintain its size in the face

of biological noise.

Results

Perturbing constriction rate changes cell length

To test the role of constriction, we perturbed its rate pharmacologically and genetically.

Fosfomycin inhibits the peptidoglycan (PG) synthesis enzyme MurA [25], which slows PG

synthesis and therefore the constriction rate. Additionally, the divisome includes cell wall

remodeling enzymes including the late-arriving FtsW and FtsI. Several point mutants of the

trans-glycosylase FtsW [26] and its cognate transpeptidase FtsI [27], referred to as FtsW**I*,

resulted in a gain-of-function phenotype in *C. crescentus* [28]. It was hypothesized that these

mutations maintain the enzymes in their active state, and thereby would increase the

constriction rate [28].

We resolved cell shape dynamics during the cell cycle by performing dual-color imaging

of the inner membrane and divisome proteins (FtsZ-GFP, FtsW-GFP) with time-lapse SIM

(Figure 1A, Movies S1-S3) on a synchronized population of cells. We used automated image

analysis to quantify cell shape parameters during the cell cycle (Figure 1B, Figure S1,

Methods). The overall cell length relative to the wild-type strain was shorter for FtsW**I* and

longer for fosfomycin-treated cells, consistent with previous studies [18,28] (Figures 2A, S2A,

B).

115
116
Figure 1: Experimental strategy and constriction-related models for modulation of cell size.
(A) Time-lapse SIM images: inner membrane (mCherry-MTS2, red), FtsZ (FtsZ-GFP, green). Shown
are example wild-type (WT), FtsW**I* mutant, and fosfomycin-treated cells through constriction, until
separation. (B) Analysis of cell shape parameters using sDaDa (see Supplementary methods and Figure
S1): the central line (black) is used to measure length (L); the width (W) is extracted from each
perpendicular segment; the cell contour defines cell shape (red line). (C) Constriction rate or onset
control mechanisms for length. Cells are born at time 0 with length at birth L_B, and elongate
exponentially. T_C and L_C are the time and length at constriction onset. T_G and L_G are the time and length
at the end of the cell cycle. Magenta parts of the cell contour represent lateral elongation, and cyan parts
represent septal elongation. Scale bars: 500 nm. Bicolor bars indicate the stage: pre-constriction
(magenta) and post-constriction (cyan).
128

Could elongation before the onset of constriction (Figure 1C, onset modulation) set the

differences in final length between FtsW**I* mutant, fosfomycin-treated and WT cells? The

appearance of a measurable constriction in SIM data corresponded well with the arrival of

FtsW (Figure S1C), and allowed us to separate elongation before and after constriction onset.

Differences in elongation before constriction for all conditions (Figure 2B) were insufficient to

account for the observed differences in final length (Figure 2A). Thus, we examined shape

changes during constriction (Figure 1C). Individual cells continued to elongate exponentially

with the same apparent rate, even as they changed from pure elongation to mid-cell remodeling

and constriction (Figures 2 C, E). However, the constriction rate was increased for the FtsW**I*

mutant and decreased for fosfomycin-treated cells when compared to WT (Figure 2C), leading

to differences in overall cell elongation during constriction (Figure 2D). Thus, we have

demonstrated that constriction rate modulation can be a mechanism for cell size control,

independent from onset modulation [12,13].

142
Figure 2. Differences in constriction rate yield different cell sizes and pole shapes.
Single-cell distributions of (A) length at division; (B) elongation before constriction; (C) mean constriction
rate; (D) elongation during constriction. (A-D), black bars represent the median of the population.
Number of cells N: WT: N=208, FtsW**I*: N=212, FOM: N=220. Significance: **: p<0.005, *: p<0.05,
n.s.: not significant. (E) Kymographs of representative cells, displaying cell diameter along the cell's
length (vertical axis), versus growth time post-synchrony (horizontal axis) , red indicates large diameter,
blue small diameter. The middle of the cell is indicated by the black horizontal line. Bicolor bars indicate
the stage: pre-constriction (magenta) and post-constriction (cyan). (F) Pole shape analysis. The
curvature is the reciprocal of the radius (Rc) of a circle tangent to the curve at a given point, here taken
to be the pole. Each cell contour represents a representative single cell from each condition; the
distribution of curvatures is plotted above (median value, black bar). (G) The pole region was extracted
from each contour (>6000 cells per condition) and analysed using principal component analysis
(Celltool, [46]). Shape mode 1 mostly accounts for variation in the length of the pole; shape mode 2
mostly accounts for variation in the bluntness of the pole independent of length. The distributions of each
shape mode is plotted, with examples of corresponding shapes. (see also Figure S2)
158

We found that individual cells continued to elongate at the same rate during

constriction, although different perturbations modulated their constriction rate. Thus, faster

constriction as in the case of FtsW**I* implies cells should have shorter, blunter poles, while

slower constriction as in the case of fosfomycin treatment implies they should have longer,

sharper poles. Indeed, kymographs show a more extended gradient in cell width at the poles

of fosfomycin-treated cells (Figure 2E). In contrast, FtsW**I* cells show a steeper gradient at

the poles. This was confirmed quantitatively by measuring the radius of curvature at the poles

(Figure 2F). Furthermore, a population-wide analysis of pole shape demonstrated that over

95% of the total shape variance is accounted for with two principle shape modes, which

primarily capture variation in the length and bluntness of the poles (Figure 2G). FtsW**I*,

fosfomycin-treated, and WT cells were all distinct along each of these shape axes. We also

observed differences in the width of the Z-ring, which appears laterally extended in the

fosfomycin case (Figure S2G). This may result from changes in length at constriction onset,

since the region of lowest MipZ concentration will be more extended in longer cells [15].

Elongation before and during constriction compensate one another

To better decipher the relative role of the constriction rate in cell size regulation we

further analyzed its contribution to cell size homeostasis. Our experiments were designed to

precisely measure the relative contributions to total elongation, and not to distinguish between

different general models of homeostasis, which would require measurement over thousands

of generations. We found that cells elongated with a distinct mean value for each condition

(Figure 3), and that the more individual cells elongated before, the less they elongated during

constriction across all conditions tested (Figure 3, Figure S3). Indeed, the total elongation was

independent of the relative time cells spent in elongation and constriction phases, with the

exception of fosfomycin-treated cells (Figure 3), generally consistent with an "adder."

Consequently, the variance in the total elongation was lower than the variances in elongation

before and during constriction would have independently suggested. This was true for all

populations, including under perturbed conditions (Figure S3A). These results demonstrate

compensation, or over-compensation in the case of fosfomycin (Figure 3, Figure S3B),

between elongation before and during constriction resulting in a higher fidelity homeostasis for

total elongation (Figure S3A-C).

Figure 3. Compensation of elongation before and during constriction contributes to cell size homeostasis.
Total elongation (gray) and elongation before constriction (color) for individual Wild Type, FtsW**I* and fosfomycin (FOM)-treated cells, as a function of normalized onset time (T_C/T_G). Lines represent the 20 cells moving average; the shaded zones represent the moving standard deviation. Extreme outliers, more than 2 standard deviations from the mean, were omitted for the calculation of the moving average (see also Figure S3).

What could be the mechanism for this compensation? Elongation rate and constriction

rate together determine elongation during constriction. Compensation could occur if cells which

elongate less before onset subsequently elongate more rapidly or constrict more slowly.

However, elongation rate during constriction did not negatively correlate with elongation before

constriction (Figure S4A). To better understand constriction dynamics, we examined single cell

waist widths as a function of relative duration of constriction (Figure 4A). Cells which elongated

more before constriction also spent relatively less time constricting, indicating a higher overall

constriction rate. The converse was true for cells which elongated less before constriction,

confirming that constriction rate was behind the compensatory effect. Single cells constricted

with increasing rate until division; thus, we defined two rates, corresponding to early and late

constriction (Figure S4B).   Interestingly, early instantaneous constriction rate correlated

positively with elongation before constriction, but late rate did not (Figure 4B, C). Hence, early

constriction rate changes at the single cell level to adjust elongation during and compensate

elongation before constriction.

While molecular mechanisms have been proposed for ensuring homeostasis, the

identity of the underlying regulatory factors remains controversial. A previous model estimated

PG precursor excess amount as a function of cell cycle [18]. Each cell is assumed to be born

with negligible excess, and generates an increasing excess of PG precursors. PG precursors

are synthesized in the cell volume, at a volume-dependent rate, while being depleted as they

become integrated into the cell wall (Supplementary Note 1). Using this model and

experimentally measured cell contours to estimate the changes in surface area (ΔA) and

volume (ΔV), we calculated the excess precursor area (A_{excess}) at the onset of constriction (T_c)

for individual cells at onset of constriction:

220
$$A_{excess}(T_c) = \langle \frac{\Delta A}{\Delta V} \rangle_{cell\ cycle} \Delta V(T_c) - \Delta A(T_c)$$

221   Here, $\langle \rangle_{cell\ cycle}$ refers to the value averaged over the cell cycle. Since it took on average 30

222   minutes to set up each experiment, we underestimated the volume and area at birth, leading

223   to an offset toward negative estimated precursor excess (Figure 4D). However, we expect the

224   trends to be insensitive to this shift.

225
Figure 4. Early constriction rates compensate for elongation before onset.
(A) Normalized waist width as a function of normalized time during constriction, colormap represents
the elongation before constriction. The measurable constriction was divided into early (0.9-0.6) and
late (0.6-0.3) stages. **(B)** Testing correlation between early constriction rate and elongation before
constriction in both WT and mutant strains, WT: r=0.45, p-value<0.01 mutant: r=0.24, p-value<0.01
(Pearson correlation coefficient). (C) Testing correlation between late constriction rate and elongation
before constriction, WT: r=0.05, p-value>0.48, mutant: r=0.02, p-value>0.8 (Pearson correlation
coefficient). (D) Testing correlation between estimated PG precursor excess and early constriction
rate, WT: r=0.33, p-value<0.01, mutant: r=0.29, p-value<0.01 (Pearson correlation coefficient).
N>=200 for each strain. (E) Schematic of size regulation in *C. crescentus* with mixed modulation of
constriction onset and rate. Magenta parts of the cell contour represent lateral elongation, and cyan
parts represent septal elongation. Later onset leads to higher PG precursor excess, which drives more
rapid initial constriction (dashed trajectories), and vice-versa (see also Figure S4).

To explain our data, we speculate on a parsimonious model in which PG precursor

excess also sets the rate of PG remodeling at the constriction site, and therefore the rate of

constriction: the higher the excess the shorter the constriction duration. Indeed, we observed

a negative correlation between rate of constriction and estimated excess PG precursor for WT

and FtsW**I* cells (Figure 4D, Figure S4). Furthermore, as the new cell poles are built, the

excess PG precursor should diminish, leading to a decreased creation of area per time. This

is indeed what we observe (supplementary note 2), consistent with models of constriction rate

in *E. coli* [14]. Fosfomycin inhibits PG synthesis, so we can no longer use the same

mathematical expression to estimate precursor excess, since the activity of fosfomycin would

introduce an extra depletion term. Interestingly, within our model this should lead to a slower

constriction, consistent with our observations (Figure 2C). Although we have posed the

regulatory factor to be PG precursors, this remains controversial because there is only indirect

evidence for their role. Any "X-factor" regulatory molecule for constriction rate following the

functional relationship described for surface area and volume would fit within the model we

suggest.

<u>Discussion</u>

Salmonella, E. coli and *B. subtilis* coordinate their cell size with nutrient availability,

perhaps to allow sufficient room for multi-fork replication [4,29,30]. *C. crescentus* shows no

such adaptation [19,31] , and how its size is modulated has remained a mystery. Our findings

show a clear contribution to cell size control from growth during the final constriction stage of

cell cycle. Modulation of constriction dynamics changes the overall length of cells, in a manner

which has implications for cell shape. In the hypothetical case of extremely rapid constriction,

the cell length would be set almost entirely by the growth during the pure elongation stage,

leading to short cells with blunt poles. By modulating constriction onset and rate together

(Figure 4E), cells may arrive at a variety of pole shapes, an unexpected control mechanism for

bacterial cell shape.

Intriguingly, the cell wall itself can have differential properties at the division site. In *B.*

subtilis, the division septum contains an enrichment of pentapeptides compared to the rest of

the cell envelope [32], perhaps due to a change in the crosslinking or PG composition. In *E.*

coli, glycan strands lacking stem peptides are enriched at the septum, allowing proteins

containing the PG binding (SPOR) domain to be recruited [33]. In *C. crescentus,* the hydrolase

DipM is recruited to the division site by its PG binding LysM domains, suggesting a distinct PG

chemistry [32–34] . Consistently, we observed a reduction in staining by wheat germ agglutinin

(WGA) at midcell for later stages of the cell (Figure S4I) [37]. Since cells were fixed prior to

staining, this implies a difference in the chemistry of the division site, but confounds the

interpretation of WGA staining in living cells [20]. We expect that in future studies it will be

important to use fluorescent cell-cycle markers in conjunction with fluorescent D-amino acids

[11], which together can identify cell-cycle timing and modes of growth. It would be interesting

to investigate whether the rate of constriction also affects cell wall chemistry at the division

site.

Different factors have been demonstrated to be important for determining constriction

dynamics. Before cells can build a septal wall, chromosomes must be partitioned; accordingly,

machinery which coordinates the two such as MatP in *E. coli* can also modulate constriction

rate [14]. Similarly, dynamically treadmilling FtsZ filaments act as a scaffold to direct cell wall

remodelers to the division site, and modulate their rate [38,39]. Nevertheless, the activity of

PG remodeling enzymes involved in constriction might depend on PG precursor amounts, and

thus may act as a part of a responsive machine. Since, for individual cells, elongation proceeds

exponentially with a single rate constant even as PG precursor excess is predicted to increase

over cell cycle, the elongation machinery is presumably relatively insensitive to changes in PG

precursor amounts. By coupling the elongation machinery to the PG precursor-sensitive

constriction machinery, the cell may have arrived at a simple means of compensating for

fluctuations in elongation during different phases of the cell cycle. This compensation still has

its limitations as we observed in the case of FtsW**I* (Figure S3B); in the case of large

elongation before onset, cells must still elongate by a minimum amount during Z-ring

maturation (Figure S2H) and constriction. In the future, it will be interesting to identify the

molecular partners responsible for constriction rate modulation and PG sensing, and to

experimentally investigate the mechanism behind compensation of elongation.

Acknowledgements

The authors would like to thank Jan Skotheim and Devon Chandler-Brown for critical reading of the manuscript, Manley lab members and Collier lab members for helpful discussions, and the Thanbichler lab for plasmid gifts.

Author contributions

A.L., A.V., S.M., E.G., M.T.L., S.H. conceived, designed and/or performed experiments, A.L., A.V., A.A., S.M., F. S., S.H., Z.P. created analysis tools and analyzed the data, and all authors contributed to the writing of the manuscript.

Declaration of interests

The authors declare no competing interests.

References

1. Koch, A.L. (1996). What size should a bacterium be? A question of scale. Annu. Rev. Microbiol. *50*, 317–348.

2. Wang, P., Robert, L., Pelletier, J., Dang, W.L., Taddei, F., Wright, A., and Jun, S. (2010). Robust growth of escherichia coli. Curr. Biol. *20*, 1099–1103.

3. National Research Council (US) Steering Group for the Workshop on Size Limits of Very Small Microorganisms (1999). Size Limits of Very Small Microorganisms: Proceedings of a Workshop (Washington (DC): National Academies Press (US)) Available at: http://www.ncbi.nlm.nih.gov/books/NBK224756/ [Accessed August 4, 2017].

4. Donachie, W.D., and Begg, K.J. (1989). Cell length, nucleoid separation, and cell division of rod-shaped and spherical cells of Escherichia coli. J. Bacteriol. *171*, 4633–4639.

5. Beveridge, T.J. (1988). The bacterial surface: General considerations towards design and function. Can. J. Microbiol. *34*, 363–372.

6. Schulz, H.N., and Jørgensen, B.B. (2001). Big bacteria. Annu. Rev. Microbiol. *55*, 105–137.

7. Marczynski, G.T. (1999). Chromosome Methylation and Measurement of Faithful, Once and Only Once per Cell Cycle Chromosome Replication inCaulobacter crescentus. J. Bacteriol. *181*, 1984–1993.

8. Cooper, S., and Helmstetter, C.E. (1968). Chromosome replication and the division cycle of Escherichia coli Br. J. Mol. Biol. *31*, 519–540.

9. Skerker, J.M., and Laub, M.T. (2004). Cell-cycle progression and the generation of asymmetry in Caulobacter crescentus. Nat. Rev. Microbiol. *2*, 325–337.

10. Aaron, M., Charbon, G., Lam, H., Schwarz, H., Vollmer, W., and Jacobs-Wagner, C. (2007). The
tubulin homologue FtsZ contributes to cell elongation by guiding cell wall precursor synthesis in
Caulobacter crescentus. Mol. Microbiol. *64*, 938–952.

11. Kuru, E., Hughes, H.V., Brown, P.J., Hall, E., Tekkam, S., Cava, F., De, P., Brun, Y.V., and
Vannieuwenhze, M.S. (2012). In situ probing of newly synthesized peptidoglycan in live bacteria
with fluorescent D-amino acids. Angew. Chem. - Int. Ed. *51*, 12519–12523.

12. Taheri-Araghi, S., Bradde, S., Sauls, J.T., Hill, N.S., Levin, P.A., Paulsson, J., Vergassola, M., and
Jun, S. (2015). Cell-Size Control and Homeostasis in Bacteria. Curr. Biol. *25*, 385–391.

13. Weart, R.B., Lee, A.H., Chien, A.-C., Haeusser, D.P., Hill, N.S., and Levin, P.A. (2007). A metabolic
sensor governing cell size in bacteria. Cell *130*, 335–347.

14. Coltharp, C., Buss, J., Plumer, T.M., and Xiao, J. (2016). Defining the rate-limiting processes of
bacterial cytokinesis. Proc. Natl. Acad. Sci. U. S. A. *113*, E1044–E1053.

15. Thanbichler, M., and Shapiro, L. (2006). MipZ, a Spatial Regulator Coordinating Chromosome
Segregation with Cell Division in Caulobacter. Cell *126*, 147–162.

16. Wallden, M., Fange, D., Lundius, E.G., Baltekin, Ö., and Elf, J. (2016). The Synchronization of
Replication and Division Cycles in Individual E. coli Cells. Cell *166*, 729–739.

17. Sompayrac, L., and Maaløe, O. (1973). Autorepressor Model for Control of DNA Replication.
Nature. New Biol. *241*, 133.

18. Harris, L.K., and Theriot, J.A. (2016). Relative rates of surface and volume synthesis set bacterial
cell size. Cell *165*, 1479–1492.

19. Campos, M., Surovtsev, I.V., Kato, S., Paintdakhi, A., Beltran, B., Ebmeier, S.E., and Jacobs-
Wagner, C. (2014). A constant size extension drives bacterial cell size homeostasis. Cell *159*,
1433–1446.

20. Banerjee, S., Lo, K., Daddysman, M.K., Selewa, A., Kuntz, T., Dinner, A.R., and Scherer, N.F.
(2017). Biphasic growth dynamics control cell division in Caulobacter crescentus. Nat. Microbiol.
2, 17116.

21. Jun, S., and Taheri-Araghi, S. (2015). Cell-size maintenance: universal strategy revealed. Trends
Microbiol. *23*, 4–6.

22. den Blauen, T., Hamoen, L.W., and Levin, P.A. (2017). The divisome at 25: the road ahead. Curr.
Opin. Microbiol. *36*, 85–94.

23. Leitao, R.M., and Kellogg, D.R. (2017). The duration of mitosis and daughter cell size are
modulated by nutrients in budding yeast. J Cell Biol, jcb.201609114.

24. Gustafsson, M.G.L. (2000). Surpassing the lateral resolution limit by a factor of two using
structured illumination microscopy. J. Microsc. *198*, 82–87.

25. Kahan, F.M., Kahan, J.S., Cassidy, P.J., and Kropp, H. (1974). THE MECHANISM OF ACTION OF
FOSFOMYCIN (PHOSPHONOMYCIN). Ann. N. Y. Acad. Sci. *235*, 364–386.

26. Meeske, A.J., Riley, E.P., Robins, W.P., Uehara, T., Mekalanos, J.J., Kahne, D., Walker, S., Kruse,
A.C., Bernhardt, T.G., and Rudner, D.Z. (2016). SEDS proteins are a widespread family of bacterial
cell wall polymerases. Nature *537*, 634–638.

27. Adam, M., Fraipont, C., Rhazi, N., Nguyen-Distèche, M., Lakaye, B., Frère, J.M., Devreese, B.,
Beeumen, J.V., Heijenoort, Y. van, Heijenoort, J. van, *et al.* (1997). The bimodular G57-V577
polypeptide chain of the class B penicillin-binding protein 3 of Escherichia coli catalyzes peptide
bond formation from thiolesters and does not catalyze glycan chain polymerization from the lipid
II intermediate. J. Bacteriol. *179*, 6005–6009.

28. Modell, J.W., Kambara, T.K., Perchuk, B.S., and Laub, M.T. (2014). A DNA Damage-Induced, SOS-
Independent Checkpoint Regulates Cell Division in Caulobacter crescentus. PLoS Biol. *12*.

29. Sargent, M.G. (1975). Control of cell length in Bacillus subtilis. J. Bacteriol. *123*, 7–19.

30. Schaechter, M., Maaloe, O., and Kjeldgaard, N.O. (1958). Dependency on medium and
temperature of cell size and chemical composition during balanced grown of Salmonella
typhimurium. J. Gen. Microbiol. *19*, 592–606.

31. Beaufay, F., Coppine, J., Mayard, A., Laloux, G., Bolle, X.D., and Hallez, R. (2015). A NAD-
dependent glutamate dehydrogenase coordinates metabolism with cell division in Caulobacter
crescentus. EMBO J. *34*, 1786–1800.

32. Morales Angeles, D., Liu, Y., Hartman, A.M., Borisova, M., de Sousa Borges, A., de Kok, N.,
Beilharz, K., Veening, J.-W., Mayer, C., Hirsch, A.K.H., *et al.* (2017). Pentapeptide-rich
peptidoglycan at the Bacillus subtilis cell-division site. Mol. Microbiol. *104*, 319–333.

33. Yahashiri, A., Jorgenson, M.A., and Weiss, D.S. (2015). Bacterial SPOR domains are recruited to
septal peptidoglycan by binding to glycan strands that lack stem peptides. Proc. Natl. Acad. Sci.
U. S. A. *112*, 11347–11352.

34. Goley, E.D., Comolli, L.R., Fero, K.E., Downing, K.H., and Shapiro, L. (2010). DipM links
peptidoglycan remodeling to outer membrane organization in Caulobacter. Mol. Microbiol. *77*,
56–73.

35. Möll, A., Schlimpert, S., Briegel, A., Jensen, G.J., and Thanbichler, M. (2010). DipM, a new factor
required for peptidoglycan remodelling during cell division in Caulobacter crescentus. Mol.
Microbiol. *77*, 90–107.

36. Poggio, S., Takacs, C.N., Vollmer, W., and Jacobs-Wagner, C. (2010). A protein critical for cell
constriction in the Gram-negative bacterium Caulobacter crescentus localizes at the division site
through its peptidoglycan-binding LysM domains. Mol. Microbiol. *77*, 74–89.

37. Douglass, K.M., Sieben, C., Archetti, A., Lambert, A., and Manley, S. (2016). Super-resolution
imaging of multiple cells by optimized flat-field epi-illumination. Nat. Photonics *10*, 705–708.

38. Bisson-Filho, A.W., Hsu, Y.-P., Squyres, G.R., Kuru, E., Wu, F., Jukes, C., Sun, Y., Dekker, C., Holden,
S., VanNieuwenhze, M.S., *et al.* (2017). Treadmilling by FtsZ filaments drives peptidoglycan
synthesis and bacterial cell division. Science *355*, 739–743.

39. Yang, X., Lyu, Z., Miguel, A., McQuillen, R., Huang, K.C., and Xiao, J. (2017). GTPase activity-
coupled treadmilling of the bacterial tubulin FtsZ organizes septal cell wall synthesis. Science
355, 744–747.

40. Goley, E.D., Yeh, Y.-C., Hong, S.-H., Fero, M.J., Abeliuk, E., McAdams, H.H., and Shapiro, L. (2011).
Assembly of the Caulobacter cell division machine. Mol. Microbiol. *80*, 1680–1698.

41. Holden, S.J., Pengo, T., Meibom, K.L., Fernandez, C.F., Collier, J., and Manley, S. (2014). High
throughput 3D super-resolution microscopy reveals Caulobacter crescentus in vivo Z-ring
organization. Proc. Natl. Acad. Sci. *111*, 4566–4571.

42. Schrader, J.M., and Shapiro, L. (2015). Synchronization of Caulobacter Crescentus for
Investigation of the Bacterial Cell Cycle. JoVE J. Vis. Exp., e52633–e52633.

43. Evinger, M., and Agabian, N. (1977). Envelope-associated nucleoid from Caulobacter crescentus
stalked and swarmer cells. J. Bacteriol. *132*, 294–301.

44. Thanbichler, M., Iniesta, A.A., and Shapiro, L. (2007). A comprehensive set of plasmids for
vanillate- and xylose-inducible gene expression in Caulobacter crescentus. Nucleic Acids Res. *35*,
e137.

45. Szeto, T.H., Rowland, S.L., Habrukowich, C.L., and King, G.F. (2003). The MinD Membrane
Targeting Sequence Is a Transplantable Lipid-binding Helix. J. Biol. Chem. *278*, 40050–40056.

46. Pincus, Z., and Theriot, J.A. (2007). Comparison of quantitative methods for cell-shape analysis. J.
Microsc. *227*, 140–156.

47. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image
analysis. Nat. Methods *9*, 671–675.

48. Sliusarenko, O., Heinritz, J., Emonet, T., and Jacobs-Wagner, C. (2011). High-throughput, subpixel
precision analysis of bacterial morphogenesis and intracellular spatio-temporal dynamics. Mol.
Microbiol. *80*, 612–627.

STAR Methods

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Suliana Manley (suliana.manley@epfl.ch).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Bacterial strains and growth conditions

Strains and plasmids

The strains, plasmids, oligonucleotides, restriction sites and modes of constructions used for this study

are summarized in key resources table. The WT and mutant strains were electroporated with the P_{van}

mCherry-MTS2 plasmid to yield the P_{van} mCherry-MTS2 strain. These strains were then electroporated

with the P_{xyl} FtsZ-GFP, P_{xyl} FtsW-GFP or P_{xyl} FtsW**-GFP plasmid to yield the respective dual color

strains.

Growth conditions

Liquid cultures were grown overnight at 28°C with 15 mL of M2G minimal media under mechanical

agitation (180 rpm). Each specific inducer for every different condition is described below. Fosfomycin

perturbation was achieved with a subminimal inhibitory concentration of 12.5 µg/ml added one hour prior

synchronization. To induce the expression of mCherry-MTS2 from the P_{van} promoter, 0.5 mM of vanillate

was added to the culture before overnight growth. For the expression of FtsW-eGFP, xylose was added

to reach a final concentration of 0.3 % (mass per volume) 2 hours before synchrony as optimized

previously [40]. P_{xyl} FtsZ-eGFP was induced overnight at 0.003 % xylose in M2G as optimized in a

previous study [41].

METHOD DETAILS

Sample preparation

Cells were synchronized at 4°C by Percoll density gradient [42] when they reached mid-exponential

phase (OD_{660}=0.3-0.5). A silicone gasket (Grace Biolabs, 103280) was placed on a rectangular cover

slide, and filled with 1% M2G agarose (Ultra PureTM Agarose, Sigma) containing fosfomycin, xylose

and vanillate at the appropriate concentrations when needed. Vanillate was present at 0.5 mM in all

experiments, xylose was present at 0.003% for induction of FtsZ-eGFP, but absent for the FtsW-eGFP

experiments. Fosfomycin was added to a final concentration of 10 µg/ml for drug perturbation

experiments. A cover slide was placed on top of the silicone gasket before solidification of the agarose

to achieve a flat agarose pad. After 5 min, the top cover slide was removed, and a 1 µL drop of a

synchronized cells suspension was placed on the pad. A small piece of agarose (~1 mm) was cut out

on two opposing sides to ensure aerobic conditions during imaging. After absorption of the droplet, the

pad was sealed with a plasma-cleaned #1.5 round coverslip with a diameter of 25 mm.

Image acquisition

Microscope set up

SIM microscopy was performed on the 3D NSIM Nikon microscope, with a CFI Apochromat TIRF

objective (100 x, NA 1.49, Nikon). The microscope was equipped with 400 mW, 561 nm and 480mW,

488 nm lasers (Coherent Sapphire) and a back-illuminated EMCCD camera (iXon 3, Andor Technology)

with a 512x512 pixel CCD sensor.

Acquisition settings

Dual color imaging of the cells was performed at 28°C using the 488 nm and 561 nm lasers for the

divisome protein-eGFP channel and the mCherry-MTS2 channel respectively. The camera was

operated with a readout speed of 1 MHz and a dynamic range of 16 bit to have the maximum pixel

readout speed at the highest dynamic range. The preamplifier gain and the electron multiplication gain

were set to 1 and 200 respectively to maximize the signal to noise at the chosen dynamic range. All raw

SIM images were acquired with a camera acquisition time of 200 ms and 100 ms (5 fps and 10 fps) for

the 561 and 488 channels. The laser power for both channels was 4 W/cm2. These settings yielded a

good balance between image quality and photo-bleaching.

All the raw images were acquired in 3D SIM image mode to ensure the highest signal to noise

ratio and lateral resolution. Fifteen images were captured of each 30.7×30.7 µm field of view, five phase-

shifted images per angle at each of three interference pattern angles. A full raw dual color image stack

was acquired in 17s.

Live-cell fluorescence microscopy over the cell cycle was achieved by performing time-lapse imaging.

3D SIM snapshots were captured at 5 min or 10 min (for fosfomycin-treated cells) time intervals to follow

dynamics while minimizing photo-bleaching of the sample during the image acquisition. Multiple fields

of view were imaged sequentially at each time point, allowing following up to 200 cells per experiment.

Super-resolved SIM images were reconstructed by the Nikon NIS-Elements software.

QUANTIFICATION AND STATISTICAL ANALYSIS

Analysis of cell shape dynamics

The super-resolved SIM images were processed via a custom-made software package called sDADA

(Shape Dynamics Automated Data Analysis). sDADA generates scatter plots, histograms and violin

plots in order to study key parameters controlling the cell size and homeostasis, such as: elongation

rate, constriction duration, length at birth, onset, onset time. sDADA extracts these parameters from the

analysis of the cell shape dynamics thanks to semi-interactive modules for image segmentation, edge

detection, cell filtering, cell tracking and statistical analysis (Supplementary Methods). The MATLAB-

based software package is available together with its documentation upon request.

Parameter definition

We assumed as time zero (T_0) the time at which the suspension of synchronized bacteria is added to

the agarose pad. This occurred approximately 20-40 minutes before starting time-lapse SIM acquisition.

T_C, T_Z, and T_W refer to the constriction onset time measured with different approaches. T_Z is the time of

the FtsZ assembly, which we assumed to occur when the fluorescence intensity of FtsZ-eGFP at mid-

cell was three times higher than elsewhere. T_W is the FtsW arrival time measured as the moment at

which the FtsW signal appeared stable at midcell (Figure S1C).

T_C is defined as the time at which the constriction invagination depth is equal to a predetermined

normalized waist width threshold. To find the optimal threshold, we tested different thresholds in the

reasonable range from 80% up to 99% of the maximum diameter, with step sizes of 1 %. Since FtsW

arrival time is an alternative readout of the constriction start the Tc values computed from the waist

diameter versus time should strongly and robustly correlate with the Tw values. For each threshold, we

computed the Pearson's correlation coefficient for the scatter plot of the Tc values versus the Tw values

of all the cells. We found that a threshold of 92% had the best correlation coefficient and minimal least

square error for Tc versus Tw.

Generation time (T_G) and final length (L_G) are the time and the length at which the cell divide.

Constriction duration τ is the difference between T_G and T_C, or when specified, T_W. The length at birth,

L_B and the elongation rate k are extracted from the exponential fitting of the elongation: $L(t) = L_B \cdot$

e^{kt}.The length at the constriction onset, L_C, L_Z or L_W, were measured as the lengths at T_C, T_Z and T_W

respectively. The total elongation is the difference between L_G and L_B. Elongation before constriction

and during constriction are defined respectively as the difference between L_G and L_C, and between L_C

and L_B. When specified, L_W can be used instead of L_C.

Statistical tests

For each parameter defined in the section above, the statistical significance of observed differences

between strains or conditions was tested. We compared the means of the repeats using Mood's median

test (Supplementary Methods). The correlation between variables was analyzed using Pearson's

correlation coefficient in the presence of a linear trend or the Spearman's correlation coefficient for non-

linear relationship. The experiments were performed in two independent replicates per condition with a

minimum of 100 cells analyzed per replicate, and 200 minimum per condition.

DATA AND SOFTWARE AVAILABILITY

All data and software used to support the results of this manuscript are available from the Lead Contact

upon reasonable request. Original data is available on Zenodo, DOI: 10.5281/zenodo.1172042 and

10.5281/zenodo.1172072

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains		
NA1000, synchronizable derivative of wild-type CB15	[43]	CB15N
CB15N *ftsW(F145L,A246T); ftsI(I45V)*	[28]	ML2159
WT *Pxyl::ftsZ-GFP Pvan::MTS2-CHY (GmR/KmR)*	This paper	CB15NVL1
ML2159 *Pxyl::ftsZ-GFP Pvan::MTS2-CHY (GmR/KmR)*	This paper	CB15NVL2
WT *Pxyl::ftsW-GFP Pvan::MTS2-CHY (GmR/KmR)*	This paper	CB15NVL3
ML2159 *Pxyl::ftsW**-GFP Pvan::MTS2-CHY (GmR/KmR)*	This paper	CB15NVL4
Oligonucleotides		
5'-AACTTGGTACCTCTAGAGGAAGATCTTTCATCGAGGAG-3'	This paper	MTS2_KpnI-F
5'-AACTTGAATTC AAGCTTCTAGGATCCACCGCCG-3'	This paper	MTS2_EcoRI-R
5'-CTCGAGCTCCGATGGCCTCCAACGCG-3'	This paper	oEG035
5'-GTTCGAATTCTCTCAGGCGTCCGCGCGACC-3'	This paper	oEG036
5'-AACTTGGTACCTCTAGAGGAAGATCTTTCATCGAGGAG-3'	This paper	MTS2_KpnI-F
Plasmids		
Pxyl::VENN-2 FtsW (KmR)	[40]	pEG105
Pxyl::VENN-2 FtsW** (KmR)	This paper	pEG1224
Pxyl:: GFPN-2(KmR)	[44]	pXGFPN-2
Pxyl:: GFPC-2(KmR)	[44]	pXGFPC-2
Pxyl::VENN-2(KmR)	[44]	PXVENN-2
E. coli MTS2 fused to GFP pBAD33 derived plasmid *Para-GFP::EcMTS::EcMTS (CmR)*	[45]	pSLR92
Pvan:: (KmR)	[44]	pVCHYC-4
Pxyl::ftsZ-dendra2 (KmR) used to create pVL1	[41]	pX-Ftsz-Dendra2
Pxyl::ftsZ-GFP (KmR)	This study	pVL1
Pvan::MTS2-CHY(GmR)	This study	pVL2
Pxyl::GFPN-2 FtsW** (KmR)	This study	pEG1308
Pxyl::VENN-2 FtsW (KmR)	[40]	pEG105

Pxyl::VENN-2 FtsW** (KmR)	This study	pEG1224
used to create pEG1305,pEG1308 (KmR)	This study	pXGFPN-2
Software and Algorithms		
sDaDa	This paper	See Supplementary Note
Celltool	[46]	http://zplab.wustl.edu/celltool/
ImageJ	[47]	https://imagej.nih.gov/ij/
MicrobeTracker	[48]	http://microbetracker.org/
MATLAB	The MathWorks, Natick, MA	https://ch.mathworks.com/products/matlab.html
Original data	This study	10.5281/zenodo.1172042 and 10.5281/zenodo.1172072

530
531