Title

Automatic and feature-specific (anticipatory) prediction-related neural activity in the human auditory system

Authors

Gianpaolo Demarchi*[1], Gaëtan Sanchez*[1,2] and Nathan Weisz[1]

* shared first authorship

Abstract

Prior experience shapes sensory perception by enabling the formation of expectations with regards to the occurrence of upcoming sensory events. Especially in the visual modality, an increasing number of studies show that prediction-related neural signals carry feature-specific information about the stimulus. This is less established in the auditory modality, in particular without bottom-up signals driving neural activity. We studied whether auditory predictions are sharply tuned to even carry tonotopic specific information. For this purpose, we conducted a Magnetoencephalography (MEG) experiment in which participants passively listened to sound sequences that varied in their regularity (i.e. entropy). Sound presentations were temporally predictable (3 Hz rate), but were occasionally omitted. Training classifiers on the random (high entropy) sound sequence and applying them to all conditions in a time-generalized manner, allowed us to assess whether and how carrier frequency specific information in the MEG signal is modulated according to the entropy level. We show that especially in an ordered (most predictable) sensory context neural activity during the anticipatory and omission periods contains carrier-frequency specific information.

22 Overall our results illustrate in the human auditory system that prediction-related neural
23 activity can be tuned in a tonotopically specific manner.

Introduction

Our capacity to predict incoming sensory inputs based on past experiences is fundamental to adapt our behavior in complex environments. A core enabling process is the identification of statistical regularities in sensory input, which does not require any voluntary allocation of processing resources (e.g. selective attention) and occurs more or less automatically in healthy brains [1]. Analogous to other sensory modalities [2,3], auditory cortical information processing takes place in hierarchically organized streams along putative ventral and dorsal pathways [4]. These streams reciprocally connect different portions of auditory cortex with frontal and parietal regions [4,5]. This hierarchical anatomical architecture yields auditory cortical processing regions sensitive to top-down modulations, thereby enabling modulatory effects of predictions. Building upon this integrated feedforward and top-down architecture, cortical and subcortical regions seem to be involved towards auditory prediction-error generation mechanisms [6,7]. A relevant question in this context is to what extent prediction-related top-down modulations (pre-)activate the same feature-specific neural ensembles as established for genuine sensory stimulation.

Such fine-tuning of neural activity would be suggested by frameworks that propose the existence of internal generative models [8–11], inferring causal structure of sensory events in our environment and the sensory consequences of our actions. A relevant process of validating and optimizing these internal models is the prediction of incoming stimulus events, by influencing activity of corresponding neural ensembles in respective sensory areas. Deviations from these predictions putatively lead to (prediction) error signals, which are passed on in a bottom-up manner to adapt the internal model, thereby continuously improving predictions [9] (for an alternative predictive coding architecture see [12]). According to this line of reasoning, predicted input should lead to weaker neural activation than input that was not predicted, which has been illustrated previously in the visual [13] and the auditory

modality [14]. Support for the idea that predictions engage neurons specifically tuned to (expected) stimulus features has been more challenging to address and has come mainly from the visual modality (for review see [15]). In an fMRI study Smith and Muckli showed that early visual cortical regions (V1 and V2), which process occluded parts of a scene, carry sufficient information to decode above chance different visual scenes [16]. Intriguingly, activity patterns in the occlusion condition are generalized to a non-occlusion control condition, implying context-related top-down feedback or input via lateral connections to modulate visual cortex in a feature specific manner. In line with these results, it has been shown that mental preplay of a visual stimulus sequence is accompanied by V1 activity that resembles activity patterns driven in a feedforward manner by the real sequence [1]. Beyond more or less automatically generated predictions, explicit attentional focus to specific visual stimulus categories also goes along with similar feature-specific modifications in early and higher visual cortices even in the absence of visual stimulation [17]. Actually it has been proposed that expectations increase baseline activity of sensory neurons tuned to a specific stimulus [18,19]. Moreover, another study using magnetoencephalography (MEG) and multivariate decoding analyses, revealed how expectation can induce a preactivation of stimulus template in visual sensory cortex suggesting a mechanism for anticipatory predictive perception [20]. Overall, for the visual modality, these studies underline that top-down processes lead to sharper tuning of neural activity to contain more information about the predicted and / or attended stimulus (feature).

Studies as to whether predictions in the auditory domain (pre-)activate specific sensory representations in a sharply tuned manner are scarce especially in humans (for animal works see e.g. [21,22]). Sharpened tuning curves of neurons in A1 during selective auditory attention have been established in animal experiments [23], even though this does not necessarily generalize to automatically formed predictions. A line of evidence could be drawn from research in marmoset monkeys, in which a reduction of auditory activity is seen

during vocalization ([24]; for suppression of neural activity to movement-related sounds in rats

in rats see [25]). This effect is abolished when fed back vocal utterances are pitch shifted [26]

thereby violating predictions. Such an action-based dynamic (i.e. adaptive) sensory filter

likely involves motor cortical inputs to the auditory cortex that selectively suppresses

predictable acoustic consequences of movement [27]. Interestingly, even inner speech may be

sufficient to produce reduced neural activity, but only when the presented sounds matched

those internally verbalized [28]. Using invasive recordings in a small set of human epilepsy

patients, it was shown that masked speech is restored by specific activity patterns in bilateral

auditory cortices [29], an effect reminiscent of a report in the visual modality ([1]; for other studies

investigating similar auditory continuity illusion phenomena see [30–32]). Albeit being feature

specific, this "filling-in" type of activity pattern observed during phoneme restoration cannot

clarify conclusively whether this mechanism requires top-down input. In principle these

results could also be largely generated via bottom-up thalamocortical input driving feature

relevant neural ensembles via lateral or feedforward connections. To resolve this issue,

putative feature-specific predictions needs to be also shown without confounding

feedforward input (i.e. during silence).

It has been recently shown, with a high resolution fMRI experiment, that predictive

responses to omissions follow a tonotopic organization in the auditory cortex[33]. But following

the notion that predictive processes are also proactive in an anticipatory sense, the exact

timing of the effects provides important evidence on whether predictions in the auditory

system come along with feature-specific preactivations of relevant neural ensembles[15]. To

this end higher temporal resolution techniques (e.g., EEG/MEG) are needed.

The goal of the present MEG study was to investigate in healthy human participants

whether predictions in the auditory modality are exerted in a carrier-frequency (i.e. tonotopic)

specific manner and, more importantly, whether those predictions are accompanied by

anticipatory effects. For this purpose, we merged an omission paradigm with a regularity

modulation paradigm (for overview see [34]; see **Figure 1** for details). So-called omission responses occur when an expected tone is replaced by silence. Frequently this response has been investigated in the context of Mismatch Negativity (MMN; [35]) paradigms, which undoubtedly have been the most common approach of studying the processing of statistical regularities in human auditory processing [36–39]. This evoked response occurs upon a deviance from a "standard" stimulus sequence, that is, a sequence characterized by a regularity violation regarding stimulation order. For omission responses (e.g. [40]), this order is usually established in a temporal sense, that is, allowing precise predictions when a tone will occur ([41]; for a study using a repetition suppression design see [42]). The neural responses during these silent periods are of outstanding interest since they cannot be explained by any feedforward propagation of activity elicited by a physical stimulus. Thus, omission of an acoustic stimulation will lead to a neural response, as long as this omission violates a regular sequence of acoustic stimuli, i.e. when it occurs unexpectedly. Previous works have identified auditory cortical contributions to the omission response (e.g. [42]). Interestingly, and underlining the importance of a top-down input driving the omission response, a recent DCM study by Chennu et al. [43] illustrates that it can be best explained when assuming top-down driving inputs into higher order cortical areas (e.g. frontal cortex). While establishing temporal predictions via a constant stimulation rate, we varied the regularity of the sound sequence by parametrically modulating its entropy level (see e.g. [44,45]). Using different carrier frequencies, sound sequences varied between random (high entropy; transition probabilities from one sound to all others at chance level) and ordered (low entropy; transition probability from one sound to another one above chance). Our reasoning was that omission-related neural responses should contain carrier-frequency specific information that is modulated by the entropy level of the contextual sound sequence. Using a time generalization decoding approach [46], we find strong evidence that particularly during the low entropy (i.e. highly ordered) sequence, neural activity prior to and during the omission period contains

127   carrier-frequency specific information similar to activity observed during real sound

128   presentation. Our work reveals how even in passive listening situations, the auditory system

129   extracts the statistical regularities of the sound input, continuously casting feature-specific

130   (anticipatory) predictions as (pre-)activations of carrier-frequency specific neural ensembles.

Figure 1: Experimental design. A) Transition matrices used to generate sound sequences according to the different conditions (random (RD), midminus (MM), midplus (MP) and ordered (OR)). B) Schematic examples of different sound sequences generated across time. 10% of sound stimuli were randomly replaced by omission trials (absence of sound) in each context.

Results

Single-trial neural activity contains carrier frequency information

A crucial first step and the foundation to address the questions of whether carrier-frequency specific neural activity patterns are modulated by predictions even in anticipatory or omission periods, is to establish that we can actually decode carrier-frequencies during actual sound presentation. To address this issue, we used the single trial MEG time-series data from the random (high entropy) condition and trained a classifier (LDA) to distinguish the four different carrier frequencies. Robust above chance (*p* < .05, Bonferroni corrected, grey line) classification accuracy was observed commencing ~30 ms following stimulus onset and peaking (~35%) at around 100 ms to gradually decline until 350 ms. Interestingly however, carrier-frequency specific information remained above chance at least until 700 ms post-stimulus onset (i.e. the entire period tested) meaning that this information was contained in the neural data when new sounds were presented. The pattern was overall similar for the other entropy conditions (not shown), however in this study we focus on the data from the random sequence, since this decoding was the basis (i.e. training data set) for all upcoming analysis using a time-generalization approach, in which carrier-frequency had to be decoded.

To identify the brain regions that provided informative activity, we adapted a previously reported approach [47], which projects the classifier weights obtained for the decoding of carrier frequency from sensor to source space (**Figure 2B**). Since later analysis using the time-generalization approach pointed to a differential entropy-level effect for early (50-125 ms; W1) and late (125-333 ms; W2) periods of the training time period (described below and in **Figure 3**), the projected weights are displayed separately for these periods. For both time periods it is evident that bilateral auditory cortical regions contribute informative activity, albeit with a right hemispheric dominance. While this appears similar for both

time-windows, informative activity also spreads to non-auditory cortical regions such as

frontal areas during the later (W2) period.

Overall, the analyses so far show that carrier frequency specific information can be

robustly decoded from MEG, with informative activity originating (as expected) mainly from

auditory cortex. Interestingly and going beyond what can be shown by conventional evoked

response analysis, carrier-frequency specific information is temporally persistent, potentially

reflecting a memory trace of the sound that is (re-)activated when new information arrives.

Figure 2: Decoding carrier frequencies from random sound sequences. A) Robust increase of
decoding accuracy is obtained rapidly, peaking ~100 ms after which it slowly wanes. Note however
that significant decoding accuracy is observed even after 700 ms, i.e. likely representing a memory
trace that is (re-)activated even when new tones (with other carrier frequencies) are processed. B)
Source projection of classifier weights for an early (W1) and late (W2) reveals informative activity to
mainly originate from auditory cortices, with a right hemispheric dominance. During the later (W2)
period informative activity spreads to also encompass e.g. frontal regions.

Modulations of entropy lead to clear anticipatory prediction effects

An anticipatory effect of predictions should be seen as relative increases of

carrier-frequency specific information prior to the onset of the expected sound with

increasing regularity of the sound sequence. To tackle this issue it is important to avoid

potential carry-over effects of decoding, which e.g would be present when training and

testing on ordered sound sequences. We circumvented this problem by training our classifier

only on data from the random sound sequence (see **Figure 2A**) and testing it for

carrier-frequency specific information in all other conditions using time generalization.

Applying this approach to pre- and post-sound (**Figure 3A**) or omissions (**Figure 3B**)

respectively, non-parametric cluster permutation test yields a clear prestimulus effect in both

cases indicating a linear increase of decoding accuracy with decreasing entropy level

(pre-sound: $p_{cluster}$ < 0.001; pre-omissions: $p_{cluster}$ < 0.001). In both cases, these anticipatory

effects appear to involve features that are relevant at later training time periods (~125-333

ms, W2; for informative activity in source space see also **Figure 2B**). The time courses of

averaged accuracy for this training time interval (only shown for pre-omissions in bottom

panel of **Figure 3B**) visualizes this effect with a clear relative increase of decoding accuracy

prior to expected sound onset in particular for the ordered sequence. This analysis clearly

underlines that predictions that evolve during a regular sound sequence contain

carrier-frequency specific information that are preactivated in an anticipatory manner, akin to

the prestimulus stimulus templates reported in the visual modality [20].

Figure 3: Regression analysis (using entropy level as independent variable; red colors indicate increased decoding accuracy for more regular sequences) for pre- and post-stimulus decoding using time-generalization of classifier trained on random sound sequences (used to avoid potential carry-over effects as soon as regularity is added to sound sequence). Upper panels display t-values thresholded at uncorrected $p < .05$. The areas framed in black are clusters significant at $p_{cluster} < .05$. The lower panel shows the decoding accuracy for individual conditions averaged for training times between the dashed lines. A) Display of effects pre- and post-sound, showing a clear anticipation effect and a late effect commencing after ~400 ms. The latter effect is more clearly visualized in the lower panel. Interestingly, different train times appear to dominate the anticipation and post-stimulus effects. B) Display of effects pre- and post-omission, showing a single continuous positive cluster. However, the actual t-values suggest temporally distinct maxima within this cluster underlining the dynamics around this event. Analogous to sounds a clear anticipation effect can be observed, driven by increased pre-omission decoding accuracy for events embedded in regular sequences (see lower panel). A similar increase can be seen immediately following the onset of the omission which cannot be observed following actual sound onset. Interestingly this increase is long lasting with further peaks emerging approximately at 330 ms and 580 ms.

Entropy-dependent classification accuracy of sounds

According to some predictive processing frameworks [15], predicted sounds should lead to a reduced activation as compared to cases in which the sound was unpredicted. In case the activation stems mainly from carrier-frequency specific neural ensembles, a decrease of decoding accuracy with increasing regularity (i.e. low entropy) could be expected. Using a time generalization approach, we applied the classifier trained on the carrier-frequencies from the random sound sequence (**Figure 2**) to the post-sound periods of the individual entropy conditions. Our regression approach yielded a negative relationship

between decoding accuracy and regularity at approximately 100-200 ms post-sound onset,

however this effect did not survive multiple comparison testing ($p_{cluster}$ = 0.12). Also on a

descriptive level, this effect appeared not to be strictly linear, meaning that our study does

not provide strong evidence that predictions reduce carrier-frequency specific information of

expected tones. Interestingly, a significant positive association ($p_{cluster}$ < 0.001) at a later

interval time period emerged, beginning around 370 ms post-sound onset and lasting at

least until 700 ms (**Figure 3A**). While - analogous to the aforementioned anticipation effect

(and the subsequently described omission effect) - decoding accuracy increased the more

regular the sound sequence was, this effect was most strongly driven by classifier

information stemming from earlier training time intervals (50-125 ms, W1; see **Figure 2B**).

This effect is in line with the previously described temporally extended effect for the decoding

of carrier frequency from random sound sequences and suggests that carrier-frequency

specific information is more strongly reactivated by subsequent sounds when embedded in a

more regular sound sequence.

Entropy-dependent classification accuracy of sound omissions

Our sound sequences were designed such that the onset of a sound could be

precisely predicted (following the invariant 3 Hz rhythm), but the predictability of the

carrier-frequencies was parametrically modulated according to the entropy level. Following

the illustration of anticipation related prediction effects, a core question of our study was

whether we could identify carrier-frequency specific information following an expected but

omitted sound onset. As depicted in **Figure 3B,** showing the outcome of our regression

analysis, this is clearly the case. Nonparametric cluster permutation test yielded a single

cluster that also comprises the anticipation effect described above ($p_{cluster}$ < 0.001), however -

also in contrast to the analysis locked to sounds (**Figure 3A**) - a post-omission onset

increase is clearly identifiable. Interestingly, also the post-omission effect was long lasting reaching far beyond the short interval of the omitted sound. The lower panel of **Figure 3B** shows the decoding accuracy averaged over the 125-333 ms training time interval (W1; see **Figure 2B**), illustrating the enhanced decoding accuracy especially for the ordered sequence. Even though the entropy-driven effect is clearly continuous, local peaks at ~90, 330 and 580 ms following the time of the (omitted) stimulation onsets can be identified at a descriptive level, thus tracking the stimulation rate of the sound sequence. This part shows that carrier-frequency specific information about an anticipated sound is enduring, not only encompassing prestimulus periods, but temporally persists when the prediction of a sound presentation is violated.

Interindividual neural representations of statistical regularities and their correlation with feature-specific predictions

Our previous analyses establish a clear relationship between (anticipatory) predictions of sounds and the carrier-frequency specific information contained in the neural signal. This was derived by a regression analysis across conditions using the entropy level as independent variable. An interesting follow-up question, is whether interindividual variability to derive the statistical regularity from the sound sequences would be correlated with carrier frequency specific information for predicted sounds.

To pursue this question we first tested whether information pertaining to the level of statistical regularity of the sequence is contained in the single trial level signal. Using all magnetometers (i.e. discarding the spatial pattern) and a temporal decoding approach showed that the entropy level of the condition in which the sound was embedded into could be decoded above chance from virtually any time point (**Figure 4A)**. Given the block-wise

presentation of the conditions, this temporally extended effect ($p < .05$, Bonferroni corrected,

green line) is not surprising. On top of this effect, a transient increase of decoding accuracy

following ~50-200 ms stimulus onset can be observed. In order to identify potential neural

generators that drive the described effect, the trained classifier weights were projected in the

source space (see inset of **Figure 4A**) analogous to the approach described above. Since

sensor level analysis suggested a temporally stable (in the sense of almost always

significant) neural pattern, the entire 0-330 ms time period was used for this purpose. The

peak informative activity was strongly right lateralized to temporal as well as parietal regions.

Based on this analysis, we can state that information about the regularity of the sound

sequence is contained also at the single-trial level and that temporal and parietal regions

may play an important role in this process. We applied the classifier trained on the sound

presentations to the same omission periods, in order to uncover whether the statistical

pattern information is also present following an unexpected omission (**Figure 4A**).

Interestingly, in this case a decrease of decoding accuracy was observed commencing ~120

ms after omission onset and lasting for ~200 ms. During this brief time period the entropy

level could not be decoded above chance. This effect illustrates that an unexpected omission

transiently interrupts the processing of the statistical regularity of the sound sequence.

To test whether the interindividual variability in neurally representing the entropy level

is associated with carrier-frequency specific predictions, we correlated average entropy

decoding accuracy in a 0-330 ms time-window following sound onset with time-generalized

decoding accuracy for carrier-frequency around sound or omission onset separately for the

early (W1) and late (W2) training time-windows. The analysis (**Figure 4B**) shows for early

training time-window (W1) a negative relationship ($p_{cluster} = 0.02$), meaning that participants

whose neural activity suggested a stronger representation of the level of statistical regularity

preactivate carrier-frequency specific neural patterns to a lesser extent. It should be noted

that the overall entropy-related effect was driven more by the later training-time window

(W2), for which no correlation was observed in the present analysis. Also, the correlation effect was not found when locking the analysis to the omission onset, which could either suggest a spurious finding or the fact of suboptimal power for this effect given the far lower amount of trials for the omission-centered time-generalized decoding. On a descriptive (uncorrected) level positive correlations can be seen following sound onset that are sequentially pronounced for early (W1) and late (W2) training-time windows ($p_{cluster}$ = 0.062 and $p_{cluster}$ = 0.096, respectively). Following the omission onset a late positive correlation ($p_{cluster}$ = 0.033) was observed ~500-600 ms for the late training time-window (W2), meaning that the carrier-frequency specific pattern of the omitted sound was reactivated more during the ordered sequence for participants who showed stronger encoding of the statistical regularity. Altogether, this analysis demonstrates that the brain has a continuous representation of the magnitude of regularity of the sequence and that it is modulated by the presence or (unexpected) absence of a sound. Furthermore, the results suggests that the interindividual variability in this more global representation of the input regularity could influence the exertion of carrier-frequency specific neural patterns preceding and following sound.

Figure 4: Decoding of entropy level of sound sequence and correlation main prediction effects gained from time-generalization analysis (see Figure 3). A) A classifier was trained to decode the entropy level from MEG activity elicited by sounds and tested around relevant events, i.e. sound or omission onset. Robust increases of decoding accuracy can be observed following sound onsets. Right temporal and parietal regions appear to contribute most informative activity (small inset). While overall decoding accuracy is above chance level mostly throughout the entire period, this pattern breaks down briefly following an omission. B) Average entropy level decoding following sound onset (0-330 ms) was taken and (Spearman) correlated with the time-generalized decoding accuracy of the low entropy condition. A significant negative correlation especially with early training time-window patterns (W1) can be seen in the anticipation period towards a sound, that was however not observed prior to omissions. Following the onset of omissions nonparametric cluster permutation testing pointed to a late positive correlation with the late activation patterns (W2).

Discussion

In this study, we investigate neural activity during passive listening to auditory tone sequences by manipulating respective entropy levels and thereby the predictability of an upcoming sound. We used MVPA applied to MEG data to first show that neural responses contain sufficient information to decode the carrier-frequency of tones. Using classifiers trained on random sound sequences in a condition- and time-generalized manner our main result is that carrier-frequency specific information increases the more regular (i.e.

predictable) the sound sequence becomes, especially in the anticipatory and post-omission

periods. This study provides strong support that prediction-related processes in the human

auditory system are sharply tuned to contain tonotopically specific information. While the

finding of sharp tuning of neural activity is not surprising, given in particular invasive

recordings from the animal auditory cortex (e.g. during vocalizations, see [26]; or shift of tuning

curves following explicit manipulations of attention to specific tone frequencies, see [23,48], our

work is a critical extension of previous human studies for which a tonotopically tuned effects

of predictions has not been shown so far. Critically, given that omission responses have

been considered as pure prediction signals [34,49], our work illustrates that sharp tuning via

predictions does not require bottom-up thalamocortical drive.

Carrier-frequency specific information following sound onset is contained in single-trial MEG

data

To pursue our main research question, we relied on MVPA applied to MEG data [46,50].

Prior to addressing whether (anticipatory) prediction-related neural activity contains

carrier-frequency specific information, an important sanity check was first to establish the

decoding performance when a sound was presented in the random sequence. A priori, this

is not a trivial undertaking given that the small spatial extent of the auditory cortex ([51] likely

produces highly correlated topographical patterns for different pure tones and the fact that

mapping tonotopic organization using noninvasive electrophysiological tools has had mixed

success (for critical overview see e.g. [52]). Considering this challenging background it is

remarkable that all participants showed a stable pattern with marked post-stimulus onset

decoding increases after ~50 ms. While a peak is reached around 100 ms post-sound onset

after which decoding accuracy slowly declines, it remains above chance for at least 700 ms.

This observation is remarkable given the passive setting for the participant (i.e. no task

involved with regards to sound input) and the very transient nature of evoked responses to

sounds that are commonly used in cognitive neuroscience. Our analysis shows that neural

activity patterns containing carrier-frequency specific information remain present for an

extended time putatively representing a memory trace of the sound that is available when

new acoustic information impinges on the auditory system. This capacity is of key

importance for forming associations across events, thereby enabling the encoding of the

statistical regularity of the input stream [53,54].

The previous result underlines that noninvasive electrophysiological methods such as

MEG can be used to decode low-level auditory features such as the carrier-frequency of

tones. This corroborates and extends findings from the visual modality for which successful

decoding of low-level stimulus features such as contrast edge orientation have been

demonstrated previously [55]. However, the analysis leading to this conclusion included all

sensors and was therefore spatially agnostic. We used an approach introduced by Marti and

Dehaene [47] to describe informative activity on the source level. Based on the subsequent

entropy-related effects identified in the time-generalization approach, we focussed on an

earlier (W1, 50-125 ms) and a later time-window (W2, 125-333 ms). While informative

activity was observed bilaterally especially in auditory regions along the Superior Temporal

Gyrus in both hemispheres, the pattern was stronger on the right side. Furthermore, on a

descriptive level, informative activity was spread more frontally in the later time-window,

implying an involvement of hierarchically downstream regions. Overall this analysis suggests

that carrier-frequency specific information mainly originates from auditory cortical regions,

but that regions not classically considered as auditory processing regions may contain

feature-specific information as well [5].

This analysis was of not only relevant as a sanity check, but also because the trained

classifiers were used and time-generalized to all entropy levels. While this approach yielded

highly significant effects (see also below for discussion), decoding accuracy was not high in

absolute terms especially for the anticipatory and omission periods. However, it should be

noted that we refrained from a widespread practice of subaveraging trials [50,56], which boosts

classification accuracies significantly. When compared to cognitive neuroscientific M/EEG

studies that perform decoding on the genuine single trials and a focus on group level effects

(rather than feature-optimizing on individual level as in BCI applications), the strength of our

effects are comparable (e.g. [47,57]).

(Anticipatory) Predictions are formed in spectrally sharply-tuned manner

Using an MVPA approach with time-generalization allowed us to assess whether

carrier-frequency related neural activity prior to or during sound / omission is systematically

modulated by the entropy level. When training and testing within regular (i.e. low entropy)

sound sequences carry-over effects could be artificially introduced that would be erroneously

interpreted as anticipation effects: in these conditions the preceding sound (with its

carrier-frequency) already contains information about the upcoming sound. To circumvent

this problem, we consistently used a classifier trained on sounds from the random sequence

- i.e. where neural activity following a sound is not predictive of an upcoming sound - and

applied it to all conditions. Using a regression analysis, we could derive in a time-generalized

manner to what extent carrier-frequency specific decoding accuracy of the (omitted) sound

was modulated by the entropy of the sound sequence in which the event was embedded.

The act of predicting usually contains a notion of pre-activating relevant neural ensembles, a

pattern that has been previously illustrated in the visual modality (e.g. [1,20]). For the omission

response, this was put forward by Bendixen et al. [49], even though the reported evoked

response effects cannot be directly seen as signatures of preactivation. Chouiter et al.[58] use

also a decoding approach, and show an effect of frequency/pitch after expected but omitted

sounds, but doesn't look for an anticipatory effect. In line with this preactivation view, our

main observation was that carrier-frequency specific information increased with increased

regularity of the tone sequence already in the pre-sound/omission period, clearly showing

sharp tuning of neural activity in anticipation of the expected sound. This effect was in particular pronounced for later training time periods (W2; see **Figure 3**) which contained informative activity also beyond auditory regions (e.g. frontal cortex; see **Figure 2B**). This finding critically extends the aforementioned previous research done in the visual modality, clearly establishing feature-specific tuning of anticipatory prediction processes in the auditory system. Our finding supports and enhances as well a recent high field fMRI study for the auditory domain[33], where the lower temporal resolution of the technique could not permit the separation of pure prediction and and preactivation effects.

According to most predictive coding accounts expected stimuli should lead to attenuated neural responses, which has been confirmed for auditory processing using evoked M/EEG (e.g. [14]) or BOLD responses (e.g. [13]). Thus a reduction of carrier-frequency specific information could also be expected when sounds were embedded in a more ordered sound sequence. While such an association was descriptively observed in early training- (W1) and testing-time (< 200 ms) intervals, it was statistically not significant and the individual conditions displayed in **Figure 3A** also suggest such a relationship not to be strictly linear. This observation may be reconciled when dissociating the strength of neural responses (e.g. averaged evoked M/EEG or BOLD response) from the feature specific information in the sigal, as has been described the visual modality: here reduced neural responses in the visual cortex have been reported, while at the same time representational information is enhanced [19]. An enhancement of representational - i.e. carrier-frequency specific - information was observed in our study at late testing-time intervals, following ~500 ms after sound onset. This effect was broad in terms of the training-time window, however it was largest for early intervals (W1; see **Figure 3A**). The late onset suggests this effect to be a consequence of the subsequent sound presentation, i.e. a reactivation of the carrier-frequency specific information within more ordered sound sequences, which would be crucial in establishing and maintaining neural representations of the statistical regularity of

the sound sequence. While this effect derived from time-generalization analysis is not

identical to the temporal decoding result described above, it is fully in line with it in a sense

that feature specific information is temporally persistent - even in this passive setting - far

beyond the typical time-windows examined in cognitive neuroscience.

Next to the anticipatory and post-sound period, we were also interested in the

strength of carrier-frequency specific information following an omission. Since no

feedforward activity along the ascending auditory pathway can account for omission-related

activity, they have been considered to reflect pure prediction responses [59]. Suggestive

evidence comes from an EEG experiment ([36], but see also [37]) in which sounds following

button presses with a fix delay could either be random or of a single identity. The authors

show evoked omission responses to be not only sensitive to timing, but also to the predicted

identity of the stimulus. However, from the evoked response it is not possible to infer which

feature specific information the signal carries (see comment above). Our result significantly

extends this research by illustrating carrier-frequency specific information to increase

following omission onset the more regular the sound sequence is. Descriptively this occurs

rapidly following omission onset with a peak ~100 ms, however further peaks can be

observed approximately following the stimulation rate up to at least 600 ms. The late

reactivations of the carrier-frequency specific information of the missing sound is in line with

the temporally persisting effects described above, pointing to an enhanced association of

past und upcoming stimulus information in the ordered sequence. However, in contrast to

the sound centered analysis, the post-omission entropy-related effects are mainly driven by

the late training time-window (W2) analogous to the anticipatory effect. Thus a speculation

based on this temporal effect could be that whereas reactivation of carrier-frequency specific

information following sounds by further incoming sounds in an ordered sequence engage

hierarchically upstream regions in the auditory system, omission-related carrier-frequency

specific activity engages downstream areas including some conventionally considered

non-auditory (e.g. frontal cortex).

Interindividual variability of entropy encoding influences carrier-frequency related prediction

effects

The overall individual ability to represent statistical regularities could have profound

implications for various behavioural domains [54]. While the main analysis pertained to

decoding of carrier-frequency specific (low-level) information, we also addressed whether a

representation of a more abstract feature such as the sequence's entropy level could also be

decoded from the noninvasive data. Functionally, extracting regularities requires an

integration over a longer time period and previous MEG works focussing on evoked

responses have identified in particular slow (DC) shifts to reflect transitions from random to

regular sound [60]. This fits with our result showing that the entropy level of a sound sequence

can be decoded above chance at virtually any time point, implying an ongoing (slow)

process tracking regularities that is transiently increased following the presentation of a

sound. Our across-participant correlation approach is suggestive that indeed the individual's

disposition to correctly represent the level of regularity is linked to pre- and

post-sound/omission engagement of carrier-frequency specific neural activity patterns. While

some open questions remain (e.g. the prestimulus discrepancy between sound and omission

correlation patterns), we consider this line of research as very promising.

Taken together, the successful decoding of low- and high-level auditory information

underlines the significant potential of applying MVPA tools to noninvasive

electrophysiological data to address research questions in auditory cognitive neuroscience

that would be difficult to pursue using conventional approaches. In particular our approach

may be a reliable and easy avenue to parametrize an individual's ability to represent

statistical regularities, without the need to invoke behavioural responses that may be

influenced by multiple non-specific factors. This could be especially valuable when studying

different groups for which conventional paradigms as relying on overt behavior may be

problematic such as children or various patient groups (e.g. disorders of consciousness).

Conclusion

Predictive processes should (pre-)engage feature-specific neural assemblies in a

top-down manner. However only little direct evidence exist for this notion in the human

auditory system [29,33]. We significantly advance this state by introducing a hybrid regularity

modulation and omission paradigm, in which expectations of upcoming carrier-frequency of

tones were controlled in a parametric manner. Using MVPA, our results unequivocally show

an increase of carrier-frequency specific information during anticipatory as well as (silent)

omission periods the more regular the sequence becomes. Our findings and outlined

approach holds significant potential to address in-depth further questions surrounding the

role of internal models in auditory perception.

Methods and Materials

Participants

A total of 34 volunteers (16 females) took part in the experiment, giving written

informed consent. At the time of the experiment, the average age was 26.6 ± 5.6 SD years.

All participants reported no previous neurological or psychiatric disorder, and reported

normal or corrected-to-normal vision. One subject was discarded from further analysis, since

in a first screening it has been assessed that she has been exposed to a wrong entropy

sequence (twice MP and no OR). The experimental protocol was approved by the ethics

committee of the University of Salzburg and has been carried out in accordance with the Declaration of Helsinki.

Stimuli and experimental procedure

Before entering the Magnetoencephalography (MEG) cabin, five head position indicator (HPI) coils were applied on the scalp. Anatomical landmarks (nasion and left/right pre-auricular points), the HPI locations, and around 300 headshape points were sampled using a Polhemus FASTTRAK digitizer. After a 5 min resting state session (not reported in this study), the actual experimental paradigm started. The subjects watched a movie ("Cirque du Soleil: Worlds Away") while passively listening to tone sequences. Auditory stimuli were presented binaurally using MEG-compatible pneumatic in-ear headphones (SOUNDPixx, VPixx technologies, Canada). This particular movie was chosen for the absence of speech and dialogue, and the soundtrack was substituted with the sound stimulation sequences. These sequences were composed of four different pure (sinusoidal) tones, ranging from 200 to 2000 Hz, logarithmically spaced (that is: 200 Hz, 431 Hz, 928 Hz, 2000 Hz) each lasting 100 ms (5 ms linear fade in / out). Tones were presented at a rate of 3 Hz. Overall the participants were exposed to four blocks, each containing 4000 stimuli, with every block lasting about 22 mins. Each block was balanced with respect to the number of presentations per tone frequency. Within the block, 10% of the stimuli were omitted, thus yielding 400 omission trials (100 per omitted sound frequency). While within each block, the overall amount of trials per sound frequency was set to be equal, blocks differed in the order of the tones, which were parametrically modulated in their entropy level using different transition matrices [61]. In more detail, the random condition (RD; see **Figure 1**) was characterized by equal transition probability from one sound to another, thereby preventing any possibility of accurately predicting an upcoming stimulus (high entropy). In the ordered

condition (OR), presentation of one sound was followed with high (75%) probability by another sound (low entropy). Furthermore, two intermediate conditions were included (midminus and midplus, labelled MM and MP respectively [61]). The probability on the diagonal was set to be equiprobable (25%) across all entropy conditions, thereby controlling for the influence of self-repetitions. The experiment was programmed in MATLAB 9.1 (The MathWorks, Natick, Massachusetts, U.S.A) using the open source Psychophysics Toolbox [62].

MEG data acquisition and preprocessing

The brain magnetic signal was recorded at 1000 Hz (hardware filters: 0.1 - 330 Hz) in a standard passive magnetically shielded room (AK3b, Vacuumschmelze, Germany) using a whole head MEG (Elekta Neuromag Triux, Elekta Oy, Finland). Signals were capture by 102 magnetometers and 204 orthogonally placed planar gradiometers at 102 different positions. We use a signal space separation algorithm implemented in the Maxfilter program (version 2.2.15) provided by the MEG manufacturer to remove external noise from the MEG signal (mainly 16.6Hz, and 50Hz plus harmonics) and realign data to a common standard head position (-trans default Maxfilter parameter) across different blocks based on the measured head position at the beginning of each block [63].

Data analysis was done using the Fieldtrip toolbox [64] (git version 20170919) and in-house built scripts. First, a high-pass filter at 0.1 Hz (6th order zero-phase Butterworth filter) was applied to the continuous data. Subsequently, for Independent Component Analysis, continuous data were chunked in 10s blocks and down-sampled at 256 Hz. The resulting components were manually scrutinized to identify eye blinks, eye movements, heartbeat and 16 and ⅔ train power supply artifacts. Finally, the continuous data were segmented from 1000ms before to 1000 ms after target stimulation onset and the artifactual components projected out (3.0 ± 1.5 SD components removed on average per each

547   subject). The resulting data epochs were down-sampled to 100 Hz for the decoding analysis.

548   Finally, the epoched data was 30 Hz lowpass-filtered (6th order zero-phase Butterworth filter)

549   prior to further analysis. Following these preprocessing steps, no trials therefore were

550   rejected [65].

Multivariate Pattern Analysis (MVPA)

We used multivariate pattern analysis as implemented in the MVPA-Light

(https://github.com/treder/MVPA-Light, commit 003a7c) package [66,67], forked and modified in

order to extract the classifier weights (https://github.com/gdemarchi/MVPA-Light/tree/devel)

[47]. MVPA decoding was performed on single trial sensor-level (102 magnetometers) data

using a time-generalization [46] analysis. Overall, three decoding approaches were taken:

1. *Entropy-level decoding*: To investigate how brain activity is modulated by the different

entropy levels, we kept only trials either with sound presentation (removing omission

trials), or all the omissions (discarding the sounds). We defined four decoding targets

(classes) based on block type (4 contexts: RD, MM, MP, OR). Sound that were

preceded by a rare (10% of the times) omission were discarded, whereas all the

omissions were kept in the omission entropy decoding.

2. *Sound-to-sound decoding*: To test whether we could classify carrier frequency in

general, we defined four targets (classes) for the decoding related to the carrier

frequency of the sound presented on each trial (4 carrier frequencies). In order to

avoid any potential carry over effect from the previous sound, the classifier was

trained only on the random (RD) sounds. Also here the sounds preceded by an

omission were discarded. To avoid further imbalance, the number of trials preceding

a target sound were equalized, e.g. the 928 Hz, sound trials were preceded by the

same number of 200 Hz, 430 Hz, 928 Hz and 2000 Hz trials (N-1, N balancing).

3. *Sound-to-omission decoding*: To test whether omission periods contain carrier frequency specific neural activity, omission trials were labeled according to the carrier frequency of the sound which would have been presented. As with the sound-to-sound decoding, the random sounds trials were used to train the classifier, which was subsequently applied to a test set of trials where sounds were not presented, i.e. omissions, using the same balancing schemes as before.

Using a Multiclass Linear Discriminant Analysis (LDA) classifier, we performed a decoding analysis at each time point around stimulus / omission onset. A five-fold cross-validation scheme, repeated for five times, was applied for entropy-level and the random sound-to-sound decoding, whereas for the training on the RD sound - testing on MM, MP, OR sounds as well as for the sound-to-omission decoding no cross-validation was performed, given the cross decoding nature of the latter testing of the classifier. For the sound-to-omission decoding analysis, the training set was restricted to random sound trials and the testing set contained only omissions. In all cases, training and testing partitions always contained different sets of data.

Classification accuracy for each subject was averaged at the group level and reported to depict the classifier's ability to decode over time (i.e. time-generalization analysis at sensor level). The time generalization method was used to study the ability of each LDA classifier across different time points in the training set to generalize to every time point in the testing set [46]. For the sound-to-sound and sound-to-omissions decoding, time generalization was calculated for each entropy level separately, resulting in four generalization matrices, one for each entropy level. This was necessary to assess whether the contextual sound sequence influences classification accuracy on a systematic level..

Decoding weights projection analysis

For relevant time frames the training decoders weights were extracted and projected in the source space in the following way. For each participant, realistically shaped, single-shell headmodels[68] were computed by co-registering the participants' headshapes either with their structural MRI (15 participants) or – when no individual MRI was available (18 participants) – with a standard brain from the Montreal Neurological Institute (MNI, Montreal, Canada), warped to the individual headshape. A grid with 1.5 cm resolution based on an MNI template brain was morphed to fit the brain volume of each participant. LCMV spatial filters were computed starting from the preprocessed data of the training random sounds (for the sound-to-sound and sound-to-omission decoding), or from all the sound or omission data (for the entropy decoding) [69]. Instead of, as commonly done in the field, multiplying the the sensor level single-trial time-series by the filter obtained above to create the so called virtual sensors, we multiplied the sensor level "corrected by the covariance matrix" [70] decoding weights time-series by the spatials filters to obtain a "informative activity" pattern [47]. Baseline normalization was performed only for visualization purposes (subtraction of 50-ms pre-stimulus activity).

Statistical analysis

For the sound and omission decoding part, we tested the dependence on entropy level using a regression test (*depsamplesregT* in Fieldtrip). Results for sounds and omissions were sorted from random to ordered respectively. In order to account for multiple comparisons, we used a nonparametric cluster permutation test [71] as implemented in Fieldtrip using 10000 permutations and a $p < .025$ to threshold the clusters, on a pseudo time-frequency (testing time vs training time - accuracy 2D structure). Moreover, to

investigate the temporal dynamics of the single entropy levels in the regression we ran the statistics above averaging across two different time windows ("avgoverfreq' in fieldtrip), namely an early time window "W1" (from 75 ms to 125 ms post random sound onset in the training set) for the sound-to-sound decoding, and a later window "W2" (from 125 ms to 333 ms).

Acknowledgments

We thank Dr. Anne Hauswald, Dr. Anne Weise and Mrs. Marta Partyka for helpful comments on earlier versions of the ms, and Miss Hayley Prins for proof reading it. We thank Mr. David Opferkuch and Mr. Manfred Seifter for the help with the measurements.

Author information

Affiliations

1 - Centre for Cognitive Neuroscience and Division of Physiological Psychology, University of Salzburg, Hellbrunnerstraße 34, 5020 Salzburg, Austria

2 - Lyon Neuroscience Research Center, Brain Dynamics and Cognition team, INSERM UMRS 1028, CNRS UMR 5292, Université Claude Bernard Lyon 1, Université de Lyon, F-69000, Lyon, France

Contributions

G.D and N.W. designed the study; G.D. performed the experiments; G.D., G.S. and N.W. designed and performed the analyses; G.D., G.S. and N.W. wrote the manuscript.

Competing interests

637 The authors declare no competing interests.

Corresponding author

639 gianpaolo.demarchi@sbg.ac.at (G.D.)

Correspondence and requests for materials

641 Further information and requests for resources should be directed to and will be fulfilled by

642 Gianpaolo Demarchi (gianpaolo.demarchi@sbg.ac.at).

References

1. Ekman, M., Kok, P. & De Lange, F. P. Time-compressed preplay of anticipated events in human primary visual cortex. *Nat. Commun.* **8**, 15276 (2017).

2. Felleman, D. J. & Van Essen, D. C. Distributed hierarchical processing in the primate cerebral cortex. *Cereb Cortex* **1**, 1–47 (1991).

3. Ungerleider, L. G. & Haxby, J. V. 'What' and 'where' in the human brain. *Curr. Opin. Neurobiol.* **4**, 157–165 (1994).

4. Rauschecker, J. P. & Tian, B. Mechanisms and streams for processing of "what" and "where" in auditory cortex. *Proc Natl Acad Sci U A* **97**, 11800–11806 (2000).

5. Plakke, B. & Romanski, L. M. Auditory connections and functions of prefrontal cortex. *Front Neurosci* **8**, 199 (2014).

6. Fontolan, L., Morillon, B., Liegeois-Chauvel, C. & Giraud, A.-L. The contribution of frequency-specific activity to hierarchical information processing in the human auditory cortex. *Nat. Commun.* **5**, 4694 (2014).

7. Recasens, M., Gross, J. & Uhlhaas, P. J. Low-Frequency Oscillatory Correlates of Auditory Predictive Processing in Cortical-Subcortical Networks: A MEG-Study. *Sci. Rep.* **8**, 14007 (2018).

8. Rao, R. P. N. & Ballard, D. H. Predictive coding in the visual cortex: A functional interpretation of some extra-classical receptive-field effects. *Nat. Neurosci.* **2**, 79–87 (1999).

9. Friston, K. A theory of cortical responses. *Philos. Trans. R. Soc. B Biol. Sci.* **360**, 815–836 (2005).

10. Wolpert, D. M., Ghahramani, Z. & Jordan, M. I. An internal model for sensorimotor

integration. *Science* **269**, 1880–1882 (1995).

11. Wolpert, D. M. & Miall, R. C. Forward Models for Physiological Motor Control. *Neural*

Netw **9**, 1265–1279 (1996).

12. Spratling, M. W. A review of predictive coding algorithms. *Brain Cogn.* **112**, 92–97

(2017).

13. Alink, A., Schwiedrzik, C. M., Kohler, A., Singer, W. & Muckli, L. Stimulus predictability

reduces responses in primary visual cortex. *J Neurosci* **30**, 2960–2966 (2010).

14. Todorovic, A. & de Lange, F. P. Repetition Suppression and Expectation Suppression

Are Dissociable in Time in Early Auditory Evoked Fields. *J. Neurosci.* **32**, 13389–13395

(2012).

15. Summerfield, C. & de Lange, F. P. Expectation in perceptual decision making: neural and

computational mechanisms. *Nat. Rev. Neurosci.* **15**, 745–756 (2014).

16. Smith, F. W. & Muckli, L. Nonstimulated early visual areas carry information about

surrounding context. *Proc Natl Acad Sci U A* **107**, 20099–20103 (2010).

17. Peelen, M. V. & Kastner, S. A neural basis for real-world visual search in human

occipitotemporal cortex. *Proc Natl Acad Sci U A* **108**, 12125–12130 (2011).

18. Kok, P., Failing, M. F. & de Lange, F. P. Prior Expectations Evoke Stimulus Templates in

the Primary Visual Cortex. *J. Cogn. Neurosci.* **26**, 1546–1554 (2014).

19. Kok, P., Jehee, J. F. M. & de Lange, F. P. Less Is More: Expectation Sharpens

Representations in the Primary Visual Cortex. *Neuron* **75**, 265–270 (2012).

20. Kok, P., Mostert, P. & Lange, F. P. de. Prior expectations induce prestimulus sensory

templates. *Proc. Natl. Acad. Sci.* 201705652 (2017). doi:10.1073/pnas.1705652114

21. Jaramillo, S. & Zador, A. M. The auditory cortex mediates the perceptual effects of

acoustic temporal expectation. *Nat Neurosci* **14**, 246–251 (2011).

22. Rubin, J., Ulanovsky, N., Nelken, I. & Tishby, N. The Representation of Prediction Error

in Auditory Cortex. *PLoS Comput. Biol.* **12**, e1005058 (2016).

23. Fritz, J. B., Elhilali, M., David, S. V. & Shamma, S. A. Does attention play a role in dynamic receptive field adaptation to changing acoustic salience in A1? *Hear Res* **229**, 186–203 (2007).

24. Eliades, S. J. & Wang, X. Sensory-motor interaction in the primate auditory cortex during self-initiated vocalizations. *J Neurophysiol* **89**, 2194–2207 (2003).

25. Schneider, D. M. & Mooney, R. Motor-related signals in the auditory system for listening and learning. *Curr. Opin. Neurobiol.* **33**, 78–84 (2015).

26. Eliades, S. J. & Wang, X. Neural substrates of vocalization feedback monitoring in primate auditory cortex. *Nature* **453**, 1102–1106 (2008).

27. Schneider, D. M., Sundararajan, J. & Mooney, R. A cortical filter that learns to suppress the acoustic consequences of movement. *Nature* **561**, 391 (2018).

28. Whitford, T. J. *et al.* Neurophysiological evidence of efference copies to inner speech. *Elife* **6**, (2017).

29. Leonard, M. K., Baud, M. O., Sjerps, M. J. & Chang, E. F. Perceptual restoration of masked speech in human cortex. *Nat Commun* **7**, 13619 (2016).

30. Kraemer, D. J. M., Macrae, C. N., Green, A. E. & Kelley, W. M. Musical imagery: sound of silence activates auditory cortex. *Nature* **434**, 158 (2005).

31. Müller, N. *et al.* You can't stop the music: reduced auditory alpha power and coupling between auditory and memory regions facilitate the illusory perception of music during noise. *Neuroimage* **79**, 383–393 (2013).

32. Voisin, J., Bidet-Caulet, A., Bertrand, O. & Fonlupt, P. Listening in silence activates auditory areas: a functional magnetic resonance imaging study. *J Neurosci* **26**, 273–278 (2006).

33. Berlot, E., Formisano, E. & De Martino, F. Mapping Frequency-Specific Tone Predictions in the Human Auditory Cortex at High Spatial Resolution. *J. Neurosci.* **38**, 4934–4942 (2018).

34. Heilbron, M. & Chait, M. Great expectations: Is there evidence for predictive coding in

auditory cortex? *Neuroscience* (2017). doi:10.1016/j.neuroscience.2017.07.061

35. Näätänen, R., Paavilainen, P., Rinne, T. & Alho, K. The mismatch negativity (MMN) in

basic research of central auditory processing: A review. *Clin. Neurophysiol.* **118**,

2544–2590 (2007).

36. SanMiguel, I., Saupe, K. & Schröger, E. I know what is missing here:

electrophysiological prediction error signals elicited by omissions of predicted "what" but

not "when". *Front. Hum. Neurosci.* **7**, (2013).

37. SanMiguel, I., Widmann, A., Bendixen, A., Trujillo-Barreto, N. & Schröger, E. Hearing

Silences: Human Auditory Processing Relies on Preactivation of Sound-Specific Brain

Activity Patterns. *J. Neurosci.* **33**, 8633–8639 (2013).

38. Recasens, M. & Uhlhaas, P. J. Test-retest reliability of the magnetic mismatch negativity

response to sound duration and omission deviants. *Neuroimage* **157**, 184–195 (2017).

39. Bendixen, A., Scharinger, M., Strauß, A. & Obleser, J. Prediction in the service of

comprehension: Modulated early brain responses to omitted speech segments. *Cortex*

53, 9–26 (2014).

40. Raij, T., McEvoy, L., Mäkelä, J. P. & Hari, R. Human auditory cortex is activated by

omission of auditory stimuli. *Brain Res.* **745**, 134–143 (1997).

41. Wacongne, C. *et al.* Evidence for a hierarchy of predictions and prediction errors in

human cortex. *Proc. Natl. Acad. Sci.* **108**, 20754–20759 (2011).

42. Todorovic, A., van Ede, F., Maris, E. & de Lange, F. P. Prior Expectation Mediates Neural

Adaptation to Repeated Sounds in the Auditory Cortex: An MEG Study. *J. Neurosci.* **31**,

9118–9123 (2011).

43. Chennu, S. *et al.* Silent Expectations: Dynamic Causal Modeling of Cortical Prediction

and Attention to Sounds That Weren't. *J Neurosci* **36**, 8305–8316 (2016).

44. Auksztulewicz, R. *et al.* The Cumulative Effects of Predictability on Synaptic Gain in the

Auditory Processing Stream. *J. Neurosci.* **37**, 6751–6760 (2017).

45. Barascud, N., Pearce, M. T., Griffiths, T. D., Friston, K. J. & Chait, M. Brain responses in

humans reveal ideal observer-like sensitivity to complex acoustic patterns. *Proc Natl*

Acad Sci U A **113**, E616-25 (2016).

46. King, J. R. & Dehaene, S. Characterizing the dynamics of mental representations: The

temporal generalization method. *Trends Cogn. Sci.* **18**, 203–210 (2014).

47. Marti, S. & Dehaene, S. Discrete and continuous mechanisms of temporal selection in

rapid visual streams. *Nat. Commun.* **8**, (2017).

48. Fritz, J. B., Elhilali, M. & Shamma, S. A. Adaptive changes in cortical receptive fields

induced by attention to complex sounds. *J Neurophysiol* **98**, 2337–2346 (2007).

49. Bendixen, A., Schroger, E. & Winkler, I. I Heard That Coming: Event-Related Potential

Evidence for Stimulus-Driven Prediction in the Auditory System. *J. Neurosci.* **29**,

8447–8451 (2009).

50. Grootswagers, T., Wardle, S. G. & Carlson, T. A. Decoding Dynamic Brain Patterns from

Evoked Responses: A Tutorial on Multivariate Pattern Analysis Applied to Time Series

Neuroimaging Data. *J. Cogn. Neurosci.* **29**, 677–697 (2017).

51. Saenz, M. & Langers, D. R. M. Tonotopic mapping of human auditory cortex. *Hear. Res.*

307, 42–52 (2014).

52. Lütkenhöner, B., Krumbholz, K. & Seither-Preisler, A. Studies of tonotopy based on wave

N100 of the auditory evoked field are problematic. *Neuroimage* **19**, 935–949 (2003).

53. Santolin, C. & Saffran, J. R. Constraints on Statistical Learning Across Species. *Trends*

Cogn. Sci. **22**, 52–63 (2018).

54. Frost, R., Armstrong, B. C., Siegelman, N. & Christiansen, M. H. Domain generality

versus modality specificity: the paradox of statistical learning. *Trends Cogn. Sci.* **19**,

117–125 (2015).

55. Cichy, R. M., Ramirez, F. M. & Pantazis, D. Can visual information encoded in cortical

columns be decoded from magnetoencephalography data in humans? *NeuroImage* **121**, 193–204 (2015).

56. Cichy, R. M., Ramirez, F. M. & Pantazis, D. Can visual information encoded in cortical columns be decoded from magnetoencephalography data in humans? *NeuroImage* **121**, 193–204 (2015).

57. Kaiser, D., Oosterhof, N. N. & Peelen, M. V. The Neural Dynamics of Attentional Selection in Natural Scenes. *J. Neurosci.* **36**, 10522–10528 (2016).

58. Chouiter, L. *et al.* Experience-based Auditory Predictions Modulate Brain Activity to Silence as Do Real Sounds. *J. Cogn. Neurosci.* **27**, 1968–1980 (2015).

59. Wacongne, C., Changeux, J.-P. & Dehaene, S. A Neuronal Model of Predictive Coding Accounting for the Mismatch Negativity. *J. Neurosci.* **32**, 3665–3678 (2012).

60. Barascud, N., Pearce, M. T., Griffiths, T. D., Friston, K. J. & Chait, M. Brain responses in humans reveal ideal observer-like sensitivity to complex acoustic patterns. *Proc. Natl. Acad. Sci.* **113**, E616–E625 (2016).

61. Nastase, S., Iacovella, V. & Hasson, U. Uncertainty in visual and auditory series is coded by modality-general and modality-specific neural systems. *Hum. Brain Mapp.* **35**, 1111–1128 (2014).

62. Brainard, D. H. The Psychophysics Toolbox. *Spat Vis* **10**, 433–436 (1997).

63. Cichy, R. M. & Pantazis, D. Multivariate pattern analysis of MEG and EEG: A comparison of representational structure in time and space. *NeuroImage* **158**, 441–454 (2017).

64. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J. M. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput. Intell. Neurosci.* **2011**, (2011).

65. Gross, J. *et al.* Good practice for conducting and reporting MEG research. *NeuroImage* **65**, 349–363 (2013).

66. Blankertz, B., Lemm, S., Treder, M., Haufe, S. & Müller, K.-R. Single-trial analysis and

classification of ERP components — A tutorial. *NeuroImage* **56**, 814–825 (2011).

67. Treder, M. S., Porbadnigk, A. K., Shahbazi Avarvand, F., Müller, K. R. & Blankertz, B.

The LDA beamformer: Optimal estimation of ERP source time series using linear

discriminant analysis. *NeuroImage* **129**, 279–291 (2016).

68. Nolte, G. The magnetic lead field theorem in the quasi-static approximation and its use

for magnetoenchephalography forward calculation in realistic volume conductors. *Phys.*

Med. Biol. **48**, 3637–3652 (2003).

69. Van Veen, B. D., van Drongelen, W., Yuchtman, M. & Suzuki, A. Localization of brain

electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Trans*

Biomed Eng **44**, 867–880 (1997).

70. Haufe, S. *et al.* On the interpretation of weight vectors of linear models in multivariate

neuroimaging. *NeuroImage* **87**, 96–110 (2014).

71. Maris, E. & Oostenveld, R. Nonparametric statistical testing of EEG-and MEG-data. *J.*

Neurosci. Methods **164**, 177–190 (2007).