Prospective identification of cross-reactive human peptide-MHC ligands for T cell receptor based therapies

Ron S. Gejman[1,2,3], Martin G. Klatt[2], Aaron Chang[2,3], Heather F. Jones[2,3], Claire Y. Oh[2,3], Smita S. Chandran[4,5], Tatiana Korontsvit[3], Viktoriya Zakahleva[3], Tao Dao[3], Christopher A. Klebanoff[2,4,5], David A. Scheinberg[2,3,5]

1. Tri-Institutional MD-PhD Program (MSKCC, Rockefeller University, Weill Cornell Medical College), New York, NY 10065, USA.
2. Weill Cornell Medicine, New York, NY, USA.
3. Molecular Pharmacology Program, Memorial Sloan Kettering Cancer Center, New York, NY, USA
4. Center for Cell Engineering and Department of Medicine, MSKCC, New York, NY, USA.
5. Parker Institute for Cancer Immunotherapy, MSKCC, New York, NY, USA.

T cell receptor (TCR)-based therapeutic cells and agents (e.g. adoptive T cell[1,2], TCR-engineered T cells[3], ImmTACs[4], TCR mimic antibodies[5], neoantigen vaccines[6,7]) have emerged as a new class of effective and selective cancer therapeutics against intracellular cancer-associated proteins. These agents rely on presentation of short peptides derived from cellular, viral or phagocytosed proteins on major histocompatibility complex (MHC). However, cross-reactivities of these agents to off-target cells and tissues are poorly understood, difficult to predict, and have resulted in serious, sometimes fatal, adverse events[8]. Here we have developed a mammalian, minigene-based method (termed "PresentER") that encodes MHC-I peptide ligands for functional immunological assays as well as for determining the reactivities and potential cross-reactivities of TCR-like therapeutic drugs against libraries of MHC-I ligands. This system is highly specific to, and entirely dependent on, the genetically encoded MHC peptide sequence, because it does not require proteasome cleavage, transporter associated with antigen processing (TAP) or processing, for immune presentation. For the expression of defined MHC-I ligands, this system is superior to expression of full-length cDNA. PresentER-expressing cells can be bound by TCR and TCR mimic (TCRm) antibodies, activate antigen-specific T cells, lead to antigen-specific cell death *in vitro* and tumor rejection in wild-type mice. Using PresentER in a pooled library screen, we find dozens of MHC-I ligands encoded in the human proteome that are cross-reactive with two TCR mimic antibodies and are not predictable by other methods. PresentER has broad and immediate utility for both basic and translational studies in immunology and oncology.

TCR based therapeutics are structurally similar to the TCR on CD8 T cells and thus share both their potential advantages and challenges. For instance, CD8 T cells can theoretically discern whether any 8-12 amino acid peptide is self, foreign or altered-self. Yet, the number of possible peptide sequences that can be encoded by the twenty proteinogenic amino acids is significantly larger than the number of circulating T cells in the human body. In order to account for this discrepancy, TCR can be highly cross-reactive: a single TCR may be capable of recognizing over 1 million distinct pMHC[9]. Thymic selection *in vivo* is critical to ensure that few circulating T cells are auto-reactive. However, TCR-based therapeutics that do not undergo negative selection for the human pMHC repertoire or are further engineered for high affinity binding can be auto-reactive. A prominent example is an affinity-enhanced engineered anti-MAGE-A3 T cell, which induced lethal cardiotoxicity in two patients during a phase I clinical trial. Extensive preclinical testing failed to uncover off-target reactivity of the anti-MAGE TCR, but afterwards it was discovered that an epitope derived from the Titin protein (a structural protein highly expressed by cardiomyocytes), was cross-reactive with the MAGE-A3 TCR[8]. Hence, a major challenge to the development of safe TCR based therapeutics is the prospective identification of off-tumor off-targets[10].

Identifying off-tumor off-targets is challenging for three reasons: (1) the scope and extent of the repertoire of MHC ligands in humans is unknown, despite multiple reports of isolation by mass spectrometry, (2) cross-reactive pMHC are not readily predictable from crystal structures or alanine scanning and (3) animal models of cross-reactivity are not possible due to the species-specific peptide processing[11] and structure of MHC. Methods to search the MHC-I ligandome for TCR targets have been developed with yeast[12,13], insect-baculovirus[14] and tetramer[15,16] technologies. These systems are

powerful and have been used to discover cross-reactive targets of some TCRs, in addition to
elucidating fundamental biology of TCRs[17]. However, the cellular systems use a synthetic covalent
linker to enforce peptide-MHC proximity, which allows presentation of peptides that would not ordinarily
be presented and may distort the structure of the epitope. The tetramer-based screening systems rely
on peptide synthesis, which is expensive and time consuming. Furthermore, tetramer, yeast and insect
systems cannot be used for *in vitro* and *in vivo* immunology assays, such as T cell recognition and
cytotoxicity. The PresentER system, by relying on the native MHC-I molecules of mammalian cells
avoids these difficulties.

We designed a TAP independent antigen presentation minigene by encoding an endoplasmic reticulum
(ER) signal sequence upstream of an MHC-I ligand, building on the methods of other groups[18]. We
modified a Mouse mammary tumor virus ER signal sequence to include SfiI (type IIS) restriction sites
flanking a removable cassette for efficient, directional cloning of DNA sequences **(Figure 1a** and **Sup.
Fig. 1)**. PresentER was designed for pooled screening applications; therefore, the amino acid
sequence corresponding to a peptide is encoded only once per minigene.

We have previously isolated and characterized two anti-cancer TCR mimic (TCRm) antibodies (ESK1[5]
and Pr20[19]), which bind to the HLA-A*02:01 ligands RMFPNAPYL (WT1:126-134), and ALYVDSLFFL
(PRAME:300-309), respectively. These antibodies selectively bound T2 cells expressing their
respective PresentER epitopes but not irrelevant epitopes **(Fig. 1b-c and Suppl. Fig. 2a-b)**.

While ESK1 and Pr20 have nanomolar affinity to pMHC, TCR typically have micromolar affinity to their
targets, making functional binding assays difficult to perform. T2 cells expressing PresentER-
NLVPMVATV (Cytomegalovirus pp65:495-503; HLA-A*02:01 ligand) were specifically bound by a TCR
multimer directed to this epitope (**Fig 1d**). In addition, an A6 TCR[20] tetramer showed specific binding to
T2 cells expressing its target MHC-I ligand PresentER LLFGYPVYV (HTLV-1 Tax:11-19; HLA-A*02:01
ligand) (Sup. **Fig. 2c**).

PresentER also was useful in presenting antigens corresponding to different MHC-I alleles. For
example, an antibody against mouse MHC-I H2-Kb/SIINFEKL bound the correct epitope in the TAP2
deficient mouse RMA-S cell line expressing PresentER SIINFEKL (Chicken Ovalbumin 257-264) or
MSIIFFLPL (PEDF:271–279). (**Fig. 1d)**.

PresentER can be used in functional immunology assays. Genetically engineered T cells expressing
the F5 TCR, specific to an HLA-A*02:01 peptide from MART-1 (27-35:AAGIGILTV)[3,21] released IFN-γ
when exposed to peptide pulsed T2 cells or T2 cells expressing the PresentER-MART-1 minigene (**Fig.
1e**). F5 T cells specifically killed PresentER MART-1 expressing T2 cells in an *in vitro* co-culture assay
(**Fig. 1f**). Finally, we challenged wild type C56B6/N mice with RMA/S cells expressing either foreign or
wild type H2-Kb ligands to test whether PresentER minigenes could mediate antigen-specific tumor
rejection *in vivo*. We found that tumors expressing foreign MHC-I ligands were rejected, in contrast to
tumors expressing non-immunogenic self-peptide (**Fig. 1g-h**).

We hypothesized that the PresentER system would proteasomal cleavage and peptide transport in the
generation of peptide-MHC. To test whether any peptide processing was occurring, we
immunoprecipitated peptide-MHC complexes from T2 cells expressing PresentER-RMF or PresentER-
ALY and identified bound peptides by mass spectrometry. RMFPNAPYL and ALYVDSLFFL were
identified only in cells encoding those PresentER constructs. Truncated versions of the RMF and ALY
peptides were not found, nor were peptides derived from the ER signal sequence identified (**Sup. Fig.
3a**). We scrambled the ER signal sequence to test whether peptides were associating with MHC via a
signal-sequence independent mechanism and found no binding to cells expressing these constructs
(Sup. Fig. 3b). Importantly, by performing low MOI transductions, we show that the PresentER system
is capable of driving sufficient antigen presentation from a single copy of the retroviral minigene,
thereby enabling its use in a pooled screen **(Sup. Fig. 4)**.
108

Pr20 and ESK1 have known cross-reactivities to off-target MHC-I ligands. Based on alanine/residue
scanning and structural[22] data, we determined that ESK1 binding to RMFPNAPYL depended primarily
on the R1 and P4 residues and for Pr20, the C-terminus of the peptide was important[19]. Therefore we
constructed a biased library of possible ESK1 cross-reactive targets by searching the human proteome
in silico for 9-mer peptide sequences containing arginine in position 1 and proline in position 4. We
performed a similar search for Pr20 cross-reactive targets using a 10-mer peptide motif based on prior
biochemical data **(Fig. 2a)**. We located 1,157 and 24,539 potential cross-reactive peptides of ESK1 and
Pr20, respectively, with NetMHCPan[23] predicted HLA-A*02:01 affinity of less than 500nM. We
synthesized a pool of 12,472 oligonucleotides that together encoded all of the ESK1 cross reactive
peptides and half of the Pr20 cross-reactive targets plus the one amino acid mutants of RMF and ALY
(termed "CR-ESK1" and "CR-Pr20", respectively), as well as positive/negative controls **(Fig. 2b)**. The
oligonucleotides were cloned into the PresentER vector and Illumina sequencing was performed to
demonstrate that library representation was maintained during cloning. The flow-based screen assaying
for cross-reactivity was performed as follows: T2 cells were transduced at low multiplicity of infection
(MOI) and transductants selected with puromycin. Library representation was kept at >1,000x at all
stages before the final sort. Transduced T2 cells were stained with ESK1 or Pr20, sorted and
sequenced (**Fig. 2c**).
126
The abundance of each minigene in the unsorted samples was well correlated (r = 0.93), indicating that
sample handling and the days between sorts did not affect the library representation (**Sup. Fig. 5a)**.
Minigenes encoding previously known ESK1 ligands had higher enrichment scores than those
encoding non-ligands (p=0.032), suggesting that the flow-based screen was able to separate ESK1
binders from non-binders (**Sup. Fig. 6a)**. Surprisingly, several of the most enriched peptides that
emerged in the ESK1 screen were CR-Pr20 peptides (**Fig. 2d)**. Although these peptides are 10-mers,
some bear sequence similarity to the target ligand of ESK1.
134
135 To validate screen hits, we synthesized a subset of the enriched peptides and assayed them for binding
136 to ESK1 by pulsing T2 cells. Of the 27 peptides tested, 22 (81%) showed increased binding to ESK1,
137 including several which had originally been selected for Pr20 cross-reactivity and did not contain a
138 proline in position 4 (**Fig. 2e)**. These unusual targets could not have been predicted from either the
139 crystal structure of ESK1 or the alanine scanning data.
140
Recently, large databases of HLA-A*02:01 peptide ligands isolated from tumors and normal tissue have
become available[24-27]. Within these databases (including personal correspondence with Department of
Immunology members at Tübingen) we found 48 nine-mer peptides with R1 and P4. Forty-five had
been included in the library of CR-ESK1 peptides identified *in silico* from the human proteome. We
synthesized 27 of these and found that 17 (63%) bound to ESK1 when pulsed on T2 cells. The median
ESK1 enrichment in the flow-based screen of these 17 peptides was 1.8 whereas the median
enrichment of the non-binders was 0.86 (**Sup. Fig. 6b**), indicating that these ESK1 ligands were
enriched in the screen, even if they were not among the top hits.
149
Two WT1-negative[28] cell lines contained known MHC-I ligands corresponding to the ESK1 off-targets
discovered in the PresentER screen (TPC-1: RLPPPFPGL, RVMPSSFFL, RLGPVPPGL, JY:
KLYNPENIYL, RLVPFLVEL). RMFPNAPYL was not found among the MHC-I ligands
immunoprecipitated from these lines. We tested ESK1 binding in each of these lines and found that JY
cells bound ESK1 at high levels while TPC-1 was marginally positive for ESK1 binding (**Fig. 2f-g**).
Thus, PresentER may be used to identify both theoretical and, in some cases, actually presented TCR
mimic off-targets.
157
A screen of Pr20 cross-reactive ligands was performed in the same manner as described above. The
abundance of each minigene in the unsorted samples was highly correlated (r=0.90-0.94) (**Sup. Fig.
5b)**. Positive control Pr20 binders were not enriched relative to the negative controls (p=0.71) (**Sup.
Fig. 7a)**. Twenty peptides more than 5-fold enriched for Pr20 binding with predicted ic50s of less than
100nM (**Sup. Fig. 7b)**. An additional 47 peptides that matched the Pr20 ligand consensus motif were

located in the HLA-A*02:01 ligand databases and 13 had been included in the CR-Pr20 library. These
peptides were not overall enriched for Pr20 binding (0.73-1.65 enrichment scores).
165
166 45 peptides were synthesized, including those enriched in the Pr20 off-target screen and several HLA-
167 A*02:01 ligands identified by mass spectrometry that matched the Pr20 ligand consensus sequence. Of
168 these, 28 (62%) were found to be Pr20 ligands when the peptides were pulsed onto T2 cells, including
169 3 known HLA-A*02:01 ligands. The 28 validated peptides were more enriched for Pr20 binding in the
170 screen than the 17 non-validated peptides (**Sup. Fig. 7c**).
171
Examining only the CR-ESK1 subset of peptides, we noticed that the peptides most enriched for ESK1
binding were also predicted to have the highest affinity for HLA-A*02:01 (**Fig. 3a**). The peptides that are
≥5-fold enriched for ESK1 binding have a median affinity to HLA-A*02:01 of 31nM, compared to 95nM
for the library as a whole and 102nM for ≥5-fold depleted peptides (**Fig. 3b**). We found the same result
in the Pr20 screen: the most enriched Pr20 ligands also had the highest affinity to MHC-I. (**Fig. 3c**). We
cloned minigenes for four of the most enriched CR-ESK1 (RLFPLAWTV 31.8x; KLMGAISFFI 41.9x)
and CR-Pr20 (WLLGDSSFFL 6.5x; LLIQEGPFFV 6.6x) peptides and tested them for binding to ESK1
and Pr20. Compared to RMFPNAPYL and ALYVDSLFFL, cells expressing these peptides bound ESK1
and Pr20 at much higher levels (**Fig. 3d**).
181
Previously known ligands of ESK1 and Pr20 were not all identified by flow-based screening of
PresentER minigenes. These ligands were originally identified by pulsing T2 cells with saturating
quantities of each peptide. We speculate that some of these peptides may not be well presented when
expressed genetically, either because of inefficient loading onto MHC-I or negative selection during
peptide editing, e.g. by TAPBPR[29]. The skew we observed in both ESK1 and Pr20-enriched minigenes
towards high-affinity HLA-A*02:01 ligands suggests that genetic expression of peptides selects for
presentation of ligands with the highest affinities for HLA-A*02:01. This is be an unexpected feature of
PresentER, as affinity to MHC-I is the most important factor in determining if a peptide is presented on
MHC-I (although high expression levels may overcome low affinity)[30]. This suggests that methods that
incorporate a flexible linker to covalently retain a peptide in the vicinity of MHC may lead to artificial
presentation of peptides that would never be presented endogenously. Additional study of the
difference between genetic expression of MHC-I ligands and peptide pulsing should be pursued to help
investigators decide which cross-reactive peptides are likely to be endogenous MHC-I ligands and thus
pose a risk to patients in a clinical setting.
196
Preclinical evaluation methods for novel therapeutic agents directed towards peptide-MHC have been
insufficient to prevent harmful off-tumor off-target toxicity. In PresentER, we have developed a
mammalian screening approach to prospectively identify cross-reactive MHC-I ligands. While
PresentER may not detect all naturally expressed cross-reactive epitopes and may detect epitopes that
are never presented in an endogenous setting, the system can help to identify potential cells and
tissues at risk for closer clinical surveillance.
203
In this report we have shown that PresentER can be used for biochemical evaluation of potentially
therapeutic TCR based agents. PresentER can also be used as an immune presentation platform *in*
vitro and *in vivo*; thus, this work can be expanded to recapitulate the MHC restricted antigenic diversity
of human cancer. Libraries of MHC-I ligands could be used to ask how tumor neoantigen heterogeneity
affects progression and treatment of tumors in immunocompetent animals, and address areas such as
neoantigen immunogenicity and clonality, cancer vaccination and immunoediting.
210

Acknowledgements
We thank Eureka Therapeutics for their generous gift of the ESK1 and Pr20 antibodies; Brian Baker
and Lance Hellman for the gift of A6 TCR plasmids and Yael David for assistance in expressing and
refolding; Scott Lowe and Eusebio Manchado Robles for the gift of the MSCV backbone and assistance
with pooled library cloning; Steven A. Rosenberg for the F5 TCR. We thank the Integrated Genomics

Operation at Memorial Sloan Kettering Cancer Center and the Flow Cytometry core facility for their
assistance.
218
219 **Funding**

220 The project was supported by the Parker Institute for Cancer Immunotherapy and the Functional
221 Genomics Initiative at Memorial Sloan Kettering Cancer Center. RSG is supported by NCI F30
222 CA200327 and NIGMS T32GM07739. DAS is supported by NCI PO1 CA 55349, CA23766. MM is
223 supported by Deutsche Forschungsgemeinschaft Grant no. KL 3118/1-1.

224
225
226
227

References

1. O'Reilly, R. J., Dao, T., Koehne, G., Scheinberg, D. & Doubrovina, E. Adoptive transfer of unselected or leukemia-reactive T-cells in the treatment of relapse following allogeneic hematopoietic cell transplantation. *Semin. Immunol.* **22,** 162–172 (2010).
2. Rosenberg, S. A. & Restifo, N. P. Adoptive cell transfer as personalized immunotherapy for human cancer. *Science* **348,** 62–68 (2015).
3. Johnson, L. A., Heemskerk, B., of, D. P. T. J.2006. Gene Transfer of Tumor-Reactive TCR Confers Both High Avidity and Tumor Reactivity to Nonreactive Peripheral Blood Mononuclear Cells and Tumor-Infiltrating Lymphocytes. *Am Assoc Immnol* **177,** 6548–6559 (2006).
4. Oates, J., Hassan, N. J. & Jakobsen, B. K. ImmTACs for targeted cancer therapy: Why, what, how, and which. *Mol. Immunol.* **67,** 67–74 (2015).
5. Dao, T. *et al.* Targeting the intracellular WT1 oncogene product with a therapeutic human antibody. *Sci Transl Med* **5,** 176ra33 (2013).
6. Sahin, U. *et al.* Personalized RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer. *Nature* **547,** 222–226 (2017).
7. Ott, P. A. *et al.* An immunogenic personal neoantigen vaccine for patients with melanoma. *Nature* **547,** 217–221 (2017).
8. Cameron, B. J. *et al.* Identification of a Titin-derived HLA-A1-presented peptide as a cross-reactive target for engineered MAGE A3-directed T cells. *Sci Transl Med* **5,** 197ra103–197ra103 (2013).
9. Singh, N. K. *et al.* Emerging Concepts in TCR Specificity: Rationalizing and (Maybe) Predicting Outcomes. *J Immunol* **199,** 2203–2213 (2017).
10. Chang, A. Y. *et al.* Opportunities and challenges for TCR mimic antibodies in cancer therapy. *Expert Opin Biol Ther* **16,** 979–987 (2016).
11. Kotturi, M. F. *et al.* Of mice and humans: how good are HLA transgenic mice as a model of human immune responses? *Immunome Res* **5,** 3 (2009).
12. Birnbaum, M. E. *et al.* Deconstructing the peptide-MHC specificity of T cell recognition. *Cell* **157,** 1073–1087 (2014).
13. Gee, M. H. *et al.* Antigen Identification for Orphan T Cell Receptors Expressed on Tumor-Infiltrating Lymphocytes. *Cell* (2017). doi:10.1016/j.cell.2017.11.043
14. Crawford, F. *et al.* Use of baculovirus MHC/peptide display libraries to characterize T-cell receptor ligands. *Immunol. Rev.* **210,** 156–170 (2006).
15. Hadrup, S. R. *et al.* Parallel detection of antigen-specific T-cell responses by multidimensional encoding of MHC multimers. *Nat Meth* **6,** 520–526 (2009).
16. Bentzen, A. K. *et al.* Large-scale detection of antigen-specific T cells using peptide-MHC-I multimers labeled with DNA barcodes. *Nat Biotechnol* **34,** 1037–1045 (2016).
17. Adams, J. J. *et al.* Structural interplay between germline interactions and adaptive recognition determines the bandwidth of TCR-peptide-MHC cross-reactivity. *Nat Immunol* **17,** 87–94 (2016).
18. Bacik, I., Cox, J. H., Anderson, R., Yewdell, J. W. & Bennink, J. R. TAP (transporter associated with antigen processing)-independent presentation of endogenously synthesized peptides is enhanced by endoplasmic reticulum insertion sequences located at the amino- but not carboxyl-terminus of the peptide. *The Journal of Immunology* **152,** 381–387 (1994).
19. Chang, A. Y. *et al.* A therapeutic T cell receptor mimic antibody targets tumor-associated PRAME peptide/HLA-I antigens. *J Clin Invest* **127,** 2705–2718 (2017).
20. Garboczi, D. N. *et al.* Assembly, specific binding, and crystallization of a human TCR-alphabeta with an antigenic Tax peptide from human T lymphotropic virus type 1 and the class I MHC molecule HLA-A2. *The Journal of Immunology* **157,** 5403–5410 (1996).
21. Johnson, L. A. *et al.* Gene therapy with human and mouse T-cell receptors mediates cancer regression and targets normal tissues expressing cognate antigen. *Blood* **114,** 535–546 (2009).

22. Ataie, N. *et al.* Structure of a TCR-Mimic Antibody with Target Predicts Pharmacogenetics.
J. Mol. Biol. **428,** 194–205 (2016).
23. Hoof, I. *et al.* NetMHCpan, a method for MHC class I binding prediction beyond humans.
Immunogenetics **61,** 1–13 (2009).
24. Schuster, H. *et al.* The immunopeptidomic landscape of ovarian carcinomas. *Proc Natl*
Acad Sci USA **114,** E9942–E9951 (2017).
25. Bassani-Sternberg, M. *et al.* Direct identification of clinically relevant neoepitopes
presented on native human melanoma tissue by mass spectrometry. *Nat Commun* **7,**
13404 (2016).
26. Bassani-Sternberg, M. *et al.* Deciphering HLA-I motifs across HLA peptidomes improves
neo-antigen predictions and identifies allostery regulating HLA specificity. *PLoS Comput*
Biol **13,** e1005725 (2017).
27. Bassani-Sternberg, M., Pletscher-Frankild, S., Jensen, L. J. & Mann, M. Mass
spectrometry of human leukocyte antigen class I peptidomes reveals strong effects of
protein abundance and turnover on antigen presentation. *Molecular & Cellular Proteomics*
14, 658–673 (2015).
28. Li, Z. *et al.* Identification of a WT1 protein-derived peptide, WT1, as a HLA-A 0206-
restricted, WT1-specific CTL epitope. *Microbiol. Immunol.* **52,** 551–558 (2008).
29. Hermann, C. *et al.* TAPBPR alters MHC class I peptide presentation by functioning as a
peptide exchange catalyst. *Elife* **4,** 26 (2015).
30. Abelin, J. G. *et al.* Mass Spectrometry Profiling of HLA-Associated Peptidomes in Mono-
allelic Cells Enables More Accurate Epitope Prediction. *Immunity* **46,** 315–326 (2017).
304

Figure 1: Design and characterization of PresentER. **(A)** The PresentER minigene encodes a signal sequence upstream of a peptide antigen and followed by a stop signal. **(B)** T2 cells expressing a PresentER minigene encoding the peptide RMFPNAPYL are bound by fluorescently labeled ESK1, a TCR mimic antibody to the complex of RMFPNAPYL/HLA-A*02:01. The color of each sample in the histogram correspond to the color of each sample in the barplot **(C)** of the same data. Antibody binding is quantified by gating all events with fluorescence intensity higher than the irrelevant-antigen controls. **(D)** A multimerized and fluorescently labeled TCR binds to T2 cells expressing its target peptide CMV pp65/NLVPMVATV in complex with HLA. **(E)** ELISPOT of genetically engineered T cells expressing the F5 TCR directed against the HLA-A*02:01 ligand MART-1 27-35 (AAGIGILTV) challenged with T2 cells pulsed with peptide, expressing the PresentER MART-1 minigene or irrelevant NY-ESO-1 157-165 (SLLMWITQC) peptide at Effector:Target ratios from 1:1 to 4:1. **(F)** The results of an *in vitro* co-culture killing assay where F5 expressing T cells were incubated with PresentER MART-1 or PresentER NY-ESO-1 minigene (GFP) expressing T2s mixed with PresentER ALYVDSLFFL (mCherry) expressing T2s. The fraction of GFP positive cells is reported at 21h and 45h, normalized to the percentage in the wells without T cells. **(G)** A fluorescently labeled antibody to SIINFEKL/H-2Kb (clone 25-D1.16) binds to RMA-S cells expressing PresentER-SIINFEKL, but not to PresentER-MSIIFFLPL. **(H)** C57BL/6N mice were injected subcutaneously with 5x10^6 RMA-S cells expressing PresentER-SIINFEKL (immunogenic; black; n=14) or PresentER-MSIIFFLPL (not immunogenic; gray; n=10). Spider plots of tumor growth across several independent experiments are plotted.

Figure 2: Design of PresentER library and ESK1 screen. **(A)** Design of sequence constraints on the target peptide library based on prior biochemical data. The human exome was mined for peptides matching the specified ESK1 and Pr20 consensus motifs. Asterisks indicate any amino acid is allowed. Red characters indicate prohibited amino acids at that position and black characters indicate allowed amino acids at that position. **(B)** The library that was constructed included 13 control peptides, 1,337 CR-ESK1 peptides and 11,083 CR-Pr20 peptides. **(C)** Schematic of the flow-based screen. T2s are transduced at low MOI (<0.3) with retrovirus encoding a pool of PresentER minigenes. Transduced cells are selected by puromycin and then cultured until sufficient cells are obtained. Cells are stained with the TCR mimic antibody and fluorescent activated cell sorting is used to sort binding and non-binding populations of cells. Genomic DNA is extracted from sorted cells and sequenced with Illumina sequencing. **(D)** Scatterplot of the ESK1 library screen. Each point is a unique peptide minigene with the x-axis indicating minigene enrichment for ESK1 binding (with 1 set as no enrichment) and y-axis indicating the peptide's predicted ic50 (in nM) to HLA-A*02:01. Lower ic50 indicates higher affinity. Marked control peptides and known ESK1 targets are plotted as triangles; CR-ESK1 as circles and CR-Pr20 as squares. **(E)** 27 peptides that were highly enriched for ESK1 binding and had high predicted affinity to HLA-A*02:01 (from Figure 2C) were synthesized at microgram scale, pulsed onto T2 cells and stained with a fluorescently labeled ESK1. Previously identified cross-reactive targets were included as positive controls. The median fluorescence intensity (MFI) of ESK1 binding is plotted, normalized to RMFPNAPYL, set at 100 units. **(F)** Representative ESK1 and isotype staining of the JY cell line. **(G)** Quantification of ESK1 and isotype staining of the JY and TPC1 cell lines.

Figure 3: Peptides enriched in TCRm screening are high affinity MHC-I ligands **(A)** Scatterplot of the ESK1 library screen with only CR-ESK1 peptides (and controls) plotted. Each point is a unique peptide minigene with the x-axis indicating minigene enrichment for ESK1 binding (with 1 set as no enrichment) and y-axis indicating the peptide's predicted ic50 (in nM) to HLA-A*02:01. Lower ic50 indicates higher affinity. Marked control peptides and known ESK1 target peptides are plotted as triangles and CR-ESK1 peptides as circles. **(B)** The netMHCPan predicted HLA-A*02:01 affinity in ic50 (nM) of all screened peptides compared to peptides which were ≥5-fold depleted in the ESK screen and peptides that were ≥5-fold enriched for ESK1 binding. **(C)** The netMHCPan predicted HLA-A*02:01 affinity in ic50 (nM) of all screened peptides compared to peptides that were ≥5-fold, ≥3-fold or ≥2-fold enriched in the Pr20 screen and peptides that were ≤3 or ≤2-fold depleted for Pr20 binding. **(D)** ESK1 and Pr20 staining of 6 PresentER minigenes.