Generation of tumorigenic porcine pancreatic ductal epithelial cells:
toward a large animal model of pancreatic cancer

Neeley Remmers[1,2], Jesse L. Cox[3], James A. Grunkemeyer[4], Shruthi Aravind[1,2], Christopher K.
Arkfeld[1,2], Michael A. Hollingsworth[4], Mark A. Carlson[1,2,5*]

[1]Department of Surgery, University of Nebraska Medical Center, Omaha, NE, USA

[2] Surgery Department, VA Nebraska-Western Iowa Health Care System, Omaha, NE, USA

[3]Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha,
NE, USA

[4]Eppley Cancer Institute, University of Nebraska Medical Center, Omaha, NE, USA

[5]Department of Genetics, Cell Biology and Anatomy, University of Nebraska Medical Center,
Omaha, NE, USA

18 *Corresponding author
19 Email: macarlso@unmc.edu (MAC)

20
21

Abstract

Background. A large animal model of pancreatic cancer would permit development of diagnostic and interventional technologies not possible in murine models, and also would provide a more biologically-relevant platform for penultimate testing of novel therapies, prior to human testing. Here, we describe our initial studies in the development of an autochthonous, genetically-defined, large animal model of pancreatic cancer, using immunocompetent pigs.

Methods. Primary pancreatic epithelial cells were isolated from pancreatic duct of domestic pigs; epithelial origin was confirmed with immunohistochemistry. Three transformed cell lines subsequently were generated from these primary cells using expression of oncogenic KRAS and dominant negative p53, with/without knockdown of p16 and SMAD4. We tested these cell lines using *in vitro* and *in vivo* assays of transformation and tumorigenesis.

Results. The transformed cell lines outperformed the primary cells in terms proliferation, population doubling time, soft agar growth, 2D migration, and Matrigel invasion, with the greatest differences observed when all four genes (KRAS, p53, p16, and SMAD4) were targeted. All three transformed cell lines grew tumors when injected subcutaneously in nude mice, demonstrating undifferentiated morphology, mild desmoplasia, and staining for both epithelial and mesenchymal markers. Injection into the pancreas of nude mice resulted in distant metastases, particularly when all four genes were targeted.

Conclusions. Tumorigenic porcine pancreatic cell lines were generated. Inclusion of four genetic

"hits" (KRAS, p53, p16, and SMAD4) appeared to produce the best results in our *in vitro* and *in*

vivo assays. The next step will be to perform autologous or syngeneic implantation of these cell

lines into the pancreas of immunocompetent pigs. We believe that the resultant large animal

model of pancreatic cancer could supplement existing murine models, thus improving preclinical

research on diagnostic, interventional, and therapeutic technologies.

Introduction

In the United States in 2016, approximately 53,000 people (48% female) were diagnosed with pancreatic cancer (~3.1% of all new cancer diagnoses), and there were ~42,000 deaths (49% female) from pancreatic cancer (~7.0% of all cancer deaths) [1, 2]. The lifetime risk for pancreatic cancer is approximately 1 in 65 [1, 2]. The incidence of pancreatic cancer has been gradually increasing since the mid-1990's, and generally is higher in the African-American population [1, 2]. Pancreatic cancer is now is the fourth most common cause of cancer-related death in both men and women (after lung, prostate, and colorectal cancer, or lung, breast, and colorectal cancer, respectively) [1, 2]. Despite apparent advances in treatment modalities and strategies [3, 4], mortality from pancreatic cancer has not decreased [1, 2]. As of 2012, the U.S. overall 5-year survival rate from pancreatic cancer was 7.7%; 5-year survival rates in localized, regional (nodal spread), or metastatic disease were 29.3, 11.1, and 2.6%, respectively [1, 2]. So there remains a need for improved early diagnosis and therapy for pancreatic cancer.

Rodent models of pancreatic cancer may not accurately reflect human biology because of differences in physiology, anatomy, immune response, and genetic sequence between the two species [5-8]. Remarkably, only 5-8% of anti-cancer drugs that emerged from preclinical studies and entered clinical studies have been ultimately approved for clinical use [9, 10]. The cause of this low approval rate is multifactorial, but likely includes the less-than-optimal predictive ability of some murine models (e.g., tumor xenografting into immunosuppressed mice) to determine the efficacy of various therapeutics in humans [5-7, 11-15]. Moreover, there are a number of genes for which the genotype-phenotype relationship is discordant between mice and human, including $CFTR^{-/-}$ and $APC^{+/-}$ [16, 17]. Incidentally, both the porcine $CFTR^{-/-}$ and $APC^{+/-}$ mutants reiterate

the human phenotype (pulmonary/GI disease and rectal polyposis, respectively) [*16-18*], in

contradistinction to the murine mutants.

In fairness, the recent trend to employ genetically-engineered mouse models (GEMM),

patient-derived xenografts (PDX), humanized mice, and *in vivo* site-directed CRISPR/Cas9

gene-edited mice in the testing of anti-cancer therapeutics may yield murine models with better

predictive ability than obtained with previously [*7, 19-23*]. Though promising, these more

advanced murine models come with increased cost and complexity [*21*], and experience with

them still is early. Importantly, all murine models have limited utility in the development of

diagnostic or interventional technology that requires an animal subject whose size approximates

a human. So at present, there remains a need for improved animal models of pancreatic cancer

that (1) are more predictive of human response to anti-cancer therapy [*21, 23*], and (2) are of

adequate size for development of specific technologies. Herein we describe some initial steps

taken in the development of a genetically-defined, autochthonous model of pancreatic cancer in

immunocompetent pigs.

Materials and Methods

Standards, rigor, reproducibility, and transparency

The animal studies of this report were designed, performed, and reported in accordance with both the ARRIVE recommendations (Animal Research: Reporting of *In Vivo* Experiments [*24*]) and the National Institutes of Health Principles and Guidelines for Reporting Preclinical Research [*25, 26*]; for details, refer to Tables S1 and S2, respectively.

Materials and animal subjects

All reagents were purchased through Thermo Fisher Scientific (www.thermofisher.com) unless otherwise noted. Short DNA sequences for vector construction, mutagenesis, and amplification purposes are shown in Table S3. Antibody information is given in Table S4. Wild type domestic swine (male and female; age 3 months at time of purchase; 30-32 kg) were purchased from the Animal Research and Development Center of the University of Nebraska Lincoln (ardc.unl.edu). Athymic homozygous nude mice (Crl:NU(NCr)-*Foxn1^{nu}*; female; 8-9 weeks old) were purchased from Charles River Laboratories, Inc. (www.criver.com). Primers utilized in this report (Table S3) were synthesized by Integrated DNA Technologies, Inc. (www.idtdna.com). DNA sequencing was performed by the UNMC Genomics Core Facility (www.unmc.edu/vcr/cores/vcr-cores/genomics).

Animal welfare

The animals utilized to generate data for this report were maintained and treated in

accordance with the *Guide for the Care and Use of Laboratory Animals* (8th ed.) from the

National Research Council and the National Institutes of Health [*27*], and also in accordance

with the Animal Welfare Act of the United States (U.S. Code 7, Sections 2131 – 2159). The

animal protocols pertaining to this manuscript were approved by the Institutional Animal Care

and Use Committee (IACUC) of the VA Nebraska-Western Iowa Health Care System (ID

numbers 00927, 00937, 00998, and 01017) or by the IACUC of the University of Nebraska

Medical Center (ID number 16-133-11-FC). All procedures were performed in animal facilities

approved by the Association for Assessment and Accreditation of Laboratory Animal Care

International (AAALAC; www.aaalac.org) and by the Office of Laboratory Animal Welfare of

the Public Health Service (grants.nih.gov/grants/olaw/olaw.htm). All surgical procedures were

performed under isoflurane anesthesia, and all efforts were made to minimize suffering.

Euthanasia was performed in accordance with the AVMA Guidelines for the Euthanasia of

Animals [*28*].

125   Isolation of porcine pancreatic ductal epithelial cells

A detailed protocol for isolation of porcine pancreatic ductal epithelial cells is provided

in the Supporting Information. In brief, the intact pancreas from male porcine research subjects

(age 5 mo) was harvested within 5 min after euthanasia, which was accomplished by transection

of the intrathoracic inferior vena cava and exsanguination while under deep isoflurane

anesthesia. These pigs had been on a protocol to study biomaterials within skin wounds of the

dorsum. The subject had not received any recent medication other the anesthetics given for

euthanasia; buprenorphine and cefovecin sodium had been given 4 weeks prior to euthanasia.

Immediately after explantation of the pancreas, the main pancreatic duct was dissected sterilely

with micro instruments from the organ body under 3.5x loupe magnification. The duct then was

mechanically digested by passage through a 70 μm sieve (Corning™ Sterile Cell Strainers,

Thermo Fisher Scientific, cat. no. 07-201-431).

The collected fragments were enzymatically digested with 1 mg/mL of Collagenase D at

37˚C for 1 h with gentle shaking. The cells were pelleted (600 g x 5 min), the supernatant was

discarded, the cell pellet was resuspended in whole media, which was defined as: DMEM (high

glucose with L-glutamine; Thermo Fisher Scientific, cat. no. 12100-046) supplemented with

10% (final concentration) fetal bovine serum (FBS; Thermo Fisher Scientific, cat. no. 26140079)

and 1% Antibiotic-Antimycotic Solution (Corning Inc., cat. no. 30-004-CI; cellgro.com). Cell

concentration in the resuspension was determined with a hemocytometer, and cells then were

diluted and pipetted into a 96-well plate (1-10 cell/well, 100-200 μL/well). After 5-7 days of

culture under standard conditions (whole media, 37˚C, 5% CO_2), wells that contained cells with

epithelial-like morphology were trypsinized and re-plated into a new 96-well plate, in order to

dilute out any fibroblasts. Cells were passaged in this fashion at least four times, until no cells

with fibroblast morphology were present. The resulting cells were passaged up to a T25 flask,

and maintained with standard conditions.

Generation of p53 and KRAS mutants and construction of

expression vector

In order to generate the porcine p53[R167H] mutant, wild-type p53 cDNA first was amplified

from cervical lymph node tissue, which was obtained <5 min after euthanasia of a 4 month old

male domestic swine that had been on an unrelated research protocol. In brief, fresh nodal tissue

was flash-frozen in liquid N_2 and then pulverized with a mortar and pestle, with continual

addition of liquid N_2 during pulverization. The frozen powder then was placed into the first

buffer solution of the QIAGEN RNEasy Mini Kit (cat. no. 74104; www.qiagen.com), and total

RNA was isolated per the manufacturer's instructions.

After isolation, the total RNA underwent reverse transcription to cDNA with a Verso

cDNA Synthesis Kit (Thermo Fisher Scientific, cat. no. AB1453A), per the manufacturer's

instructions. The wild type p53 sequence was amplified out of the cDNA using the PCR primers

shown in Table S3, which flanked the p53 cDNA with *Sal*I and *Bam*HI restriction sites.

Successful amplification of the wild-type p53 cDNA was confirmed by inserting the amplified

candidate sequence into the TOPO® vector (TOPO® TA Cloning® Kit; Invitrogen™/Life

Technologies™, Thermo Fisher Scientific, cat. no. K202020) per the manufacturer's

instructions, followed by sequencing.

Site-directed mutation of wild-type p53 into p53^{R167H} was performed using Agilent

Technologies' QuickChange II Site-Directed Mutagenesis Kit (cat. no. 200523;

www.genomics.agilent.com) with the mutagenic primers shown in Table S3, per the

manufacturer's instructions. Presence of the p53^{R167H} mutation was verified by sequencing as

described above. The multiple cloning site of a pIRES2-AcGFP1 Vector (Takara Bio USA, Inc.,

cat. no. 632435; www.clontech.com; manufacturer's vector information included as **Fig. S1**) was

cut with *Sal*I and *Bam*HI, and the p53^{R167H} sequence then was ligated into this plasmid.

The source of the porcine KRAS^{G12D} mutant was the plasmid used to generate the

p53/KRAS Onco-pig [*29, 30*]. The KRAS^{G12D} cDNA was amplified out of this plasmid with

primers (see Table S3) that flanked the sequence with *Xho*I and *Pst*I restriction sites. The

amplified product was inserted into the TOPO vector and verified by sequencing, as described

above. The above pIRES2-AcGFP1 Vector (already containing the p53^{R167H} sequence) then was

cut with *Xho*I and *Pst*I, and the KRAS^{G12D} sequence was ligated into this plasmid, producing a

pIRES2-AcGFP1 Vector which contained both mutant cDNAs within its multiple cloning site

(KRAS^{G12D} upstream).

The newly-constructed plasmid, hereafter designated as GKP (G = AcGFP1; K =

KRAS^{G12D}; P = p53^{R167H}), was transformed into One Shot™ Stbl3™ Chemically Competent *E.*

coli (Invitrogen™/Thermo Fisher Scientific, cat. no. C737303), per the manufacturer's

instructions, and plasmid DNA subsequently was isolated using a QIAGEN Plasmid Maxi Kit

(cat. no. 12162), per the manufacturer's instructions. This plasmid then was transfected into

Takara's Lenti-X™ 293T cells (Clontech, cat. no. 632180), using Takara's Xfect™ Transfection

Reagent (Clontech, cat. no. 631317), per the manufacturer's instructions, to generate infectious

lentiviral particles that would direct expression of AcGFP1, KRAS^{G12D}, and p53^{R167H} mutants in

transduced cells.

Generation of shRNA-expressing vectors

Short hairpin RNA (shRNA) constructs targeting the porcine SMAD4 and p16 genes

were created using InvivoGen's siRNA Wizard™ software (www.invivogen.com/sirnawizard).

Three targeting sequences for each gene along with scrambled controls initially were generated

and tested. The shRNA construct that demonstrated the best target knock-down in preliminary

experimentation (as verified by PCR, data not shown) was utilized for subsequent experiments

(see Table S3 for sequences ultimately selected for the shRNA constructs). Primers were cloned

into the psiRNA-h7SKhygro G1 and psiRNA-h7SKneo G1 vectors (InvivoGen, cat. no. ksirna3-

h21 and ksirna3-n21, respectively; www.invivogen.com) and plasmids then were isolated, all per

the manufacturer's protocol.

Cell transformations

Primary porcine pancreatic epithelial cells in T75 flasks were grown to 80% confluency

under standard conditions. The media then was exchanged with 2-3 mL of supernatant from non-

lysed Lenti-X™ 293T cells (containing GKP viral particles) with 2 µg/mL polybrene (cat. no.

TR1003, Thermo Fisher Scientific). After 24-48 h at 37˚C, treated epithelial cells were re-seeded

into 6-well plates under standard conditions and grown to 80% confluency. An exchange with

whole media containing 2 µg/mL G418 aminoglycoside antibiotic then was performed; the G418

dose was chosen based on preliminary dose-response studies against non-treated epithelial cells.

After 24 h, a whole media exchange was done, and the presence of transduced cells was

determined with inverted GFP fluorescent microscopy of living cells. Subsequent transfections

for RNAi were done with the above plasmids employing shRNA sequences against SMAD4

and/or p16, and using the LyoVec™ reagent (InvivoGen, cat. no. lyec-12), all per the

manufacturer's protocol. Transfected cells then were selected for expression of the shRNA

vector using the appropriate aminoglycoside antibiotic (G418 or hygromycin B).

PCR

Cell and tissue RNA was isolated using the QIAGEN RNEasy Mini Kit. Purified RNA

then was used to generate cDNA using the Verso cDNA Synthesis Kit. The Platinum® Blue

PCR Supermix (Invitrogen™/Life Technologies, cat. no. 12580) subsequently was used for all

PCR reactions. Amplified products were separated with agarose gel electrophoresis, and then

visualized using a UV-light box. qPCR was performed using the PowerUp™ SYBR® Green

Master Mix (Applied Biosystems™/ Thermo Fisher Scientific, cat. no. A25741) per

manufacturer's protocol, and run on an Applied Biosystems™ 7500 Fast Dx Real-Time PCR

Instrument. Fold changes in gene expression were calculated using the comparative C_T method

[31]. All primers used are listed in Table S3.

Immunoblotting

231 Western blot analysis was performed to confirm overexpression of the mutant p53 protein

232 (see Table S4 for a list of antibodies used), as previously described [32]. An antibody specific for

233 the mutant KRAS protein was not commercially available. Antibody expression was visualized

234 using the Li-Cor Odyssey Electrophoresis Imaging System (www.licor.com).

Soft agar assay

A standard soft agar assay [33] was used to determine anchorage independent growth. A

base layer of 1% agarose was plated into 6-well plates. A total of 2,500 cells/well were mixed

with 0.7% agarose and plated on top of the base layer. The plates were incubated under standard

conditions for 21 days. The cells then were stained with crystal violet, and counted using an

inverted microscope. Cells were plated in triplicate, and total counts from all three wells were

averaged.

Migration assay

245    A standard scratch assay (monolayer wounding) [34] was performed to determine cellular

246 migration rate. Cells were plated in triplicate into 6-well plates. A horizontal scratch using a 10

247 μL pipet tip was made in each well. After washing away scratched-off cells, baseline images

248 along the scratch were obtained, the plates were incubated under standard conditions, and

249 subsequent images were captured at 3, 6, 9, 12, and 15 h after the initial scratch. ImageJ software

250 (imagej.nih.gov/ij) was used to measure the distance between the two migrating cellular fronts

251 (scratch edges) at 3-5 locations along the scratch. Average distance at each time point was

252 plotted to generate the migratory rate (μm/h).

Invasion assay

255    BioCoat™ Matrigel™ Invasion Chambers (Corning™, Thermo Fisher Scientific, cat. no.

256 08-774) were plated with 50,000 cells (upper chamber) in triplicate, and incubated under

257 standard conditions for 24 h. The media from the upper chamber then was removed, and any

258 cells remaining in the upper chamber were removed using a cotton swab. Cells that had migrated

259 to the bottom of the membrane were stained using a Kwik-Diff™ kit (Shandon™, Thermo Fisher

260 Scientific, cat. no. 9990701). Membranes were mounted onto glass slides, and cells per high-

261 power field were counted using ImageJ software.

Population doubling assay

264    Cells were plated in 6-well plates (20,000 cell/well), and cultured under standard

265 conditions. Triplicate plates then were trypsinized on days 1, 2, 3, 4, 6, and 8, and cells were

266 counted with a hemocytometer. Cell number *vs*. day was plotted to determine the day range in

267 which linear growth was achieved. The data from this linear growth phase were used to

268   determine population doubling time (DT) using the formula: $DT = (\Delta t) \times \ln(2) \div \ln(N_f/N_i)$

269   where Δt = time interval between initial and final cell count, N_f = cell count at final time, and N_i

270   = cell count at initial time.

Proliferation assay

Relative cell proliferation rates were assayed using an MTT (3-(4,5-dimethylthiazol-2-

yl)-2,5-diphenyltetrazolium bromide) assay kit (Vybrant™ MTT Cell Proliferation Assay Kit,

Invitrogen™, Thermo Fisher Scientific, cat. no. V13154). Cells were plated in triplicate in a 96-

well plate (5,000 cell/well), and cultured under standard conditions for 48 h. MTT reagent then

was added to the cells per the manufacturer's instructions, followed by addition of the solvent

solution 3.5 h later. Absorbance was measured with a plate reader 3.5 h after solvent addition.

Mean absorbance was normalized to absorbance from wild type pancreatic ductal epithelial cells

to calculate fold-difference in proliferation.

Immunofluorescence and immunohistochemistry

283   Antibodies used in immunofluorescent and immunohistochemical experiments are listed

284   in Table S4. Agilent Dako EnVision kits (www.agilent.com) were used for all IHC analyses per

285   the manufacturer's instructions.

Subcutaneous tumorigenic cell injection

Subcutaneous implantation of tumorigenic cells was performed as previously described

[35], with some modifications. Tumorigenic porcine pancreatic ductal epithelial cells were

trypsinized, counted, and resuspended in DMEM at a concentration of 1×10^7 viable cells/mL.

Nude mice (N = 10 per treatment group) then were injected with 5×10^6 cells (500 µL) into the

right hind flank. Tumors were allowed to grow for 6 weeks or until they reached 1.8 cm in

diameter, as measured with a caliper. At necropsy all gross tumor was measured and collected,

portions underwent formalin fixation and paraffin embedding, and sections subsequently

underwent H&E or immunohistochemical staining as described above. An independent, blinded

pathologist analyzed the stained sections to determine whether tumors were epithelial in origin,

and if they displayed malignant features.

Orthotopic tumorigenic cell injection

Orthotopic implantation of tumorigenic cells was performed as previously described [*35*]

to analyze metastases and desmoplasia. In brief, tumorigenic porcine pancreatic ductal epithelial

cells were trypsinized and counted, and 1×10^4 viable cells in 20 µL DMEM were injected with

a 20 gauge needle into the pancreas of nude mice (N = 10-12 per treatment group) through a 5

mm incision in the left upper quadrant. Mice were euthanized 6 weeks after injection, and tumors

and organs were harvested for gross and histologic analysis, as described in the previous

paragraph.

Statistical analysis

309   Data are reported as mean ± standard deviation. Groups of continuous data were

310   compared with ANOVA and the unpaired t-test.

Results

Isolation of primary porcine pancreatic ductal epithelial cells

Cells cultured from micro-dissected pancreatic ducts displayed epithelial morphology under phase microscopy and stained for CK19 (an established marker of pancreatic ductal epithelium [36]; **Fig. 1A-B**). Based on these results, we were confident that we had a population of pancreatic epithelial cells that we could use to generate tumorigenic cell lines.

Generation of tumorigenic cell lines

In order to transform our primary porcine pancreatic ductal epithelial cells, we first generated a lentiviral construct containing $KRAS^{G12D}$ and $TP53^{R167H}$, genes previously identified [29] as the porcine equivalents to the mutant $KRAS$ and $TP53$ which are present in multiple human cancers [37-41]; in the mouse, expression of these mutants was the basis for the KRAS/p53 genetically engineered murine model of pancreatic cancer [42]. For our model, we chose to use a lentiviral platform for the vector, because its genome would be large enough to accommodate insertion of both mutant genes; in addition, we believed that a lentivirus would be optimal for transforming primary cells.

Since initial sequencing of the porcine genome has been accomplished [43], we were able to utilize the National Library of Medicine's nucleotide BLAST® database (blast.ncbi.nlm.nih.gov/blast.cgi) to determine the porcine genetic equivalents for human $SMAD4$ and $CDKN2A$. We then designed primers (Table S3) to amplify these two genes from genomic DNA isolated from skin of a healthy domestic pig. We sequenced our amplification products,

and accessed the BLAST® database to confirm that our products aligned with the porcine

SMAD4 and *CDKN2A* gene sequences. We then proceeded to generate lentiviral constructs to

transform primary porcine pancreatic ductal epithelial cells into cell lines expressing various

combinations of mutant KRAS and p53, SMAD4 shRNA, and p16[Ink4A] shRNA (see cell line

definitions in Table 1), as described under Materials and Methods.

Primary porcine epithelial cells next were transduced with the GKP lentivirus to generate

transformed cell lines. Overexpression of KRAS[G12D] and p53[R167H] was confirmed in these cell

lines with qPCR and immunoblotting (**Fig. 1C-E**). Of note, we could not obtain a reliable

antibody to detect porcine KRAS with immunoblotting, so we had to rely on qPCR results and

expression of GFP as markers of KRAS[G12D] expression. Preliminary *in vitro* analyses to probe

the tumorigenic properties of these GKP-transformed cell lines demonstrated modest increases in

soft agar colony formation and migration speed over wild type cells (**Fig. 2A & B**). Interestingly,

the cell line (2.22) with the best performance in the soft agar and migration assays had only

modest overexpression of KRAS[G12D] and p53[R167H] (**Fig. 1 & 2**); in contrast, the cell lines with

the highest mutant overexpression performed relatively poorly in these *in vitro* assays of

"tumorigenesis" (i.e., evidence of transformed behavior in cell culture).

While *in vitro* experiments were being performed, *in vivo* pancreatic tumor induction was

attempted utilizing a transgenic mini-pig available from the National Swine Resource and

Research Center (NSRRC; nsrrc.missouri.edu). Known as the "Onco-pig," [*30*], this subject

carries an LSL-cassette containing the dominant negative *TP53*[R167H] and the activated *KRAS*[G12D]

sequences [*30, 44, 45*]; i.e., this subject is the porcine analog of the KRAS/p53 mouse [*42, 46*].

As demonstrated previously, site-specific expression of Cre recombinase in the Onco-pig

resulted in localized p53 inhibition and KRAS activation, while subcutaneous injection of AdCre

produced mesenchymal tumors at the injection sites [*30*]. We injected Cre recombinase into the

pancreas of five Onco-pigs (**Fig. S2**). After four months, we observed no gross tumors. However,

there was immunohistochemical evidence of transgene expression at the pancreatic injection

sites along with numerous microscopic proliferative lesions (**Fig. S2**).

Based on the modest evidence of *in vitro* transformation and the lack of gross *in vivo*

tumorigenesis using expression of p53^{R167H} and KRAS^{G12D} only, we decided that additional

oncogenic stress might be helpful to increase the tumor-like properties of transformed pancreatic

ductal epithelial cells. Utilizing cell line 2.22 (hereafter referred to PGKP; see Table 1), which

had relatively good performance in the soft agar and migration assays (**Fig. 2A & B**), sequential

transduction with lentiviral constructs expressing shRNA against SMAD4 and then p16^{Ink4A} was

performed to generate cell lines PGKPS and PGKPSC (see Table 1), respectively. RT-PCR then

was used to confirm knockdown of the targeted transcripts in these two cell lines (**Fig. 2C**).

In vitro tumorigenic properties of transformed cells

The *in vitro* "tumorigenic" properties of the PGKP, PGKPS, and PGKPSC cell lines first

were compared with the soft agar and migration assays (**Fig. 2D & E**). Addition of SMAD4 ±

p16^{Ink4A} knockdown enhanced the ability of transformed cells to form colonies in soft agar and

increased their migration speed (2D wounding assay), particularly when both transcripts were

targeted (i.e., the PGKPSC line). We then compared population doubling time, proliferation

(metabolic dye conversion) , and Matrigel® invasion ability among the three transformed lines

with respect to wild type cells (**Fig. 3A-C**). Both the PGKPS and PGKPSC cell lines had greater

proliferation and invasive ability compared to either wild type cells or the PGKP cell line (**Fig.**

3B, C). The doubling time for all three transformed cell lines was approximately the same at ~15

h, compared to the ~4 d doubling time of wild type cells (**Fig. 3A**). Based on the *in vitro* assays

of tumorigenesis, we suspected that all three of our transformed cell lines had the potential to

form tumors *in vivo*, albeit to varying degrees. We subsequently decided to compare the *in vivo*

tumorigenicity among all three cell lines in an immunodeficient mouse model.

Subcutaneous and orthotopic tumor transplants in immunodeficient
386

mice

In order to determine if our cell lines retained their tumorigenic properties *in vivo*, we

utilized homozygous athymic mice to generate subcutaneous and orthotopic cell implantation

models. The subcutaneous model was used to assess *in vivo* tumor growth. All three cell lines

grew sizeable tumors (>1 cm diameter) within 6 weeks; growth rates were not statistically

different among the three cell lines (**Fig. 4A & B**). The subcutaneous tumors were well-

vascularized and mucinous in gross appearance (**Fig. 4A**). Confirmation of GKP lentiviral

transduction was demonstrated with immunohistochemistry of the $p53^{R167H}$ mutant in

subcutaneous tumors (**Fig. S3**).

Tumors from the subcutaneous implant model then underwent immunohistochemical

staining with an array of epithelial and mesenchymal markers (**Fig. 5**). The distribution of

staining for the epithelial markers (E-cadherin, epithelial cell adhesion molecule, pan-

cytokeratin, cytokeratin-19) generally was more diffuse than the mesenchymal marker staining.

In some regions the epithelial marker staining was clustered and intense. The overall abundance

of staining for the epithelial markers appeared greater in the PGKPS and PGKPSC lines with

respect to the PGKP line. The distribution of staining for the mesenchymal markers (α-smooth

muscle actin, vimentin, and type I collagen) was variable, sometimes appearing in cords or

strands in some sections, reminiscent of the desmoplastic reaction in human pancreatic cancer

[*47, 48*]. In other sections, mesenchymal marker staining was minimal. Similar to the epithelial

markers, the overall abundance of staining for the mesenchymal markers appeared greater in the

PGKPS and PGKPSC lines compared to the PGKP line.

With the immunohistochemical studies of the subcutaneous tumors demonstrating some

epithelial characteristics, the nude mouse orthotopic implantation model was used next to assess

the metastatic potential of all three cell lines. The percentage of mice implanted with each cell

line that subsequently developed metastasis in the small bowel, diaphragm, liver, lung, lymph

node, peritoneum, or spleen is shown in **Fig. 4C.** The degree of metastatic spread was minimal;

nodal tissue was the only metastatic site common to all three lines. The PGKPCS cell line

exhibited the greatest array of metastatic spread, with the most common site being the spleen

(though it was not clear in two of five mice with splenic disease after PGKPSC implantation

whether the spleen was involved with extension from the primary tumor, or from direct seeding,

or whether these were true metastases). Interestingly, one of the two subjects with liver

metastasis after PGKPSC implantation had primary tumor within the gallbladder, rather than

within the pancreas. Similar to the subcutaneous tumor model, most tumors in the orthotopic

model were well-vascularized and mucinous in gross appearance.

Discussion

Our overall goal with this project was to generate tumorigenic pancreatic cell lines that could be used in an immunocompetent porcine model of pancreatic cancer. Current murine models for pancreatic cancer will continue to be helpful, particularly for the study of molecular mechanisms. However, murine models are limited in their ability to replicate human biology and size, so a large animal model of pancreatic cancer likely would enhance our ability to develop and test new diagnostic and treatment modalities for this disease. The data presented herein demonstrated that wild type porcine pancreatic ductal epithelium can be transformed with modulation of common tumor-associated target genes, and that these transformed cells subsequently can grow tumors in immunodeficient mice. These data provide a pathway for the construction of an autochthonous porcine model of pancreatic cancer, namely, orthotopic implantation of tumorigenic pancreatic cells.

The rationale to build a porcine model of pancreatic cancer is (1) to have a platform for diagnostic/therapeutic device development otherwise not achievable in murine models; and (2) to have a highly predictive preclinical model in which anti-cancer therapies (including immunotherapies) could be vetted/optimized prior to a clinical trial. Porcine biomedical models have been used for decades in the fields of trauma and hemostasis [49], xenotransplantation [50], dermal healing [51], toxicology [52], atherosclerosis [53], and cardiac regeneration [54], to name a few. The rationale to use the pig in this modeling effort is that this species mimics human genomics [55-59], epigenetics [60], physiology [52, 55, 61, 62], metabolism [55, 62, 63], inflammation and immune response [59, 64-68], and size [62, 69] remarkably well (in particular,

better than mice), with reasonable compromises towards cost and husbandry [62]. Recently, a

porcine genome map was generated [43], and further coverage, annotation, and confirmation is

ongoing [57]. Genetic manipulation of pigs (including knockouts, tissue-specific transgenics,

inducible expression [29, 70-77]) with similar tools as used in the mouse is becoming more

routine, with new gene-edited porcine models emerging in 2015-2017 for diseases such as

atherosclerosis, cystic fibrosis, Duchenne muscular dystrophy, and ataxia telangiectasia [45, 78].

So based on the pig's relatively large size and its proven track record in replicating human

biology which, incidentally, is a demonstrably better replication than can be obtained with

rodents, we selected swine as the model organism for this pancreatic cancer project.

Commonly mutated genes in pancreatic cancer include *KRAS* [*79, 80*], *TP53* [*80-82*],

SMAD4 [*80, 83*], and *CDKN2A* [*80, 84*]. Somatic activation of KRAS via the G12D mutation

($KRAS^{G12D}$) in mice produced widely metastatic pancreatic tumors; survival duration in these

subjects decreased further with p53 inactivation [*42*]. Deletion of *SMAD4* or *CDKN2A* enhanced

tumor growth in a $KRAS^{G12D}$ murine pancreatic cancer model [*85, 86*]. Based on this published

data, we elected to induce porcine pancreatic tumors with targeting of *KRAS*, *TP53*, *SMAD4*, and

CDKN2A, through site-directed expression of mutant genes and shRNA targeting. In the future,

we will gene edit cell lines using the CRISPR/Cas9 system (as opposed to shRNA expression) to

effect specific gene deletions.

Research on immunocompetent large animal cancer models [*87*] includes prostate cancer,

for which there is a canine model [*88*]. A group in Munich has engineered (i) an *APC* mutant pig

that develops rectal polyposis [*17*] and (ii) a pig with Cre-inducible p53 deficiency [*75*]. In

addition, a group in Denmark has engineered a *BRCA* mutant pig [*89*]. In 2012 the NSRRC

(National Swine Resource and Research Center, nsrrc.missouri.edu), an NIH-funded entity

whose mission is to generate transgenic swine models for biomedical research, engineered a

swine (the "Onco-pig"; mini-pig background) which carries an LSL-cassette containing

dominant negative *TP53* with a R167H mutation (p53^{R167H}) and activated *KRAS*^{G12D} [*30, 44, 45*];

i.e., the porcine analog of the KRAS/p53 mouse [*42, 46*]. Site-specific expression of Cre

recombinase in the Onco-pig resulted in localized p53 inhibition and KRAS activation;

subcutaneous injection of AdCre produced mesenchymal tumors at the injection sites [*30*]. In

2017, initial work was published on a Onco-pig-based model of hepatocellular carcinoma [*90*].

476   All three of our cell lines demonstrated *in vitro* transformed behavior and the ability to

form tumors *in vivo*. Although we utilized a relatively high number of cells to obtain tumor

formation, we feel that this is due to the relative young age of our cell lines, which did not

undergo as many passages as other cell lines currently used to study pancreatic cancer. Thus, the

transformed cell lines in this report likely did not undergo an inadvertent selection for the fastest

growing cells, as may occur in older, extensively-passaged cell lines. Moreover, the lack of

desmoplasia in our murine xenograft model was not surprising, since most immunodeficient

murine models of pancreatic cancer do not recapitulate the extent of desmoplasia or metastasis

seen with human disease. We anticipate that future experiments involving the implantation of

transformed pancreatic ductal epithelial cells into wild type swine (and thus into a "normal"

microenvironment with "normal" immunoediting) will produce tumors with desmoplasia and

metastasis.

References

1. American Cancer Society. Cancer Facts & Figures 2016. Atlanta: American Cancer Society; 2016. Available from: http://www.cancer.org/research/cancerfactsstatistics/cancerfactsfigures2016/index.

2. SEER (Surveillance, Epidemiology, and End Results Program) Stat Fact Sheets: Pancreas Cancer: National Cancer Institute; [Accessed December 12, 2016]. Available from: URL: http://seer.cancer.gov/statfacts/html/pancreas.html.

3. National Comprehensive Cancer Network. Pancreatic Adenocarcinoma, Version 2.2016. 2016. In: NCCN Clinical Practice Guidelines In Oncology (NCCN Guidelines®) [Internet]. NCCN. Available from: www.nccn.org.

4. Costello E, Greenhalf W, Neoptolemos JP. New biomarkers and targets in pancreatic cancer and their application to treatment. *Nat Rev Gastroenterol Hepatol*. 2012;9(8):435-44. doi: 10.1038/nrgastro.2012.119. PubMed ID: 22733351; PubMed Central ID:

5. Seok J, Warren HS, Cuenca AG, Mindrinos MN, Baker HV, Xu W, et al. Genomic responses in mouse models poorly mimic human inflammatory diseases. *PNAS*. 2013;110(9):3507-12. doi: 10.1073/pnas.1222878110. PubMed ID: 23401516; PubMed Central ID:

6. Begley CG, Ellis LM. Drug development: Raise standards for preclinical cancer research. *Nature*. 2012;483(7391):531-3. doi: 10.1038/483531a. PubMed ID: 22460880; PubMed Central ID:

7.   Cook N, Jodrell DI, Tuveson DA. Predictive in vivo animal models and translation to

clinical trials. *Drug Discov Today*. 2012;17(5-6):253-60. doi:

10.1016/j.drudis.2012.02.003. PubMed ID: 22493784; PubMed Central ID:

8.   Le Magnen C, Dutta A, Abate-Shen C. Optimizing mouse models for precision cancer

prevention. *Nat Rev Cancer*. 2016;16(3):187-96. doi: 10.1038/nrc.2016.1. PubMed ID:

26893066; PubMed Central ID:

9.   Reichert JM, Wenger JB. Development trends for new cancer therapeutics and vaccines.

Drug Discov Today. 2008;13(1-2):30-7. doi: 10.1016/j.drudis.2007.09.003. PubMed ID:

18190861; PubMed Central ID:

10.   Sharpless NE, Depinho RA. The mighty mouse: genetically engineered mouse models in

cancer drug development. *Nat Rev Drug Discov*. 2006;5(9):741-54. doi:

10.1038/nrd2110. PubMed ID: 16915232; PubMed Central ID:

11.   Ebos JM, Kerbel RS. Antiangiogenic therapy: impact on invasion, disease progression,

and metastasis. *Nat Rev Clin Oncol*. 2011;8(4):210-21. doi: 10.1038/nrclinonc.2011.21.

PubMed ID: 21364524; PubMed Central ID:

12.   Francia G, Cruz-Munoz W, Man S, Xu P, Kerbel RS. Mouse models of advanced

spontaneous metastasis for experimental therapeutics. *Nat Rev Canc*. 2011;11(2):135-41.

doi: doi:10.1038/nrc3001. PubMed ID: 21258397; PubMed Central ID:

13.   O'Collins VE, Macleod MR, Donnan GA, Horky LL, van der Worp BH, Howells DW.

1,026 experimental treatments in acute stroke. *Ann Neurol*. 2006;59(3):467-77. doi:

10.1002/ana.20741. PubMed ID: 16453316; PubMed Central ID:

14.   Scott S, Kranz JE, Cole J, Lincecum JM, Thompson K, Kelly N, et al. Design, power,

and interpretation of studies in the standard murine model of ALS. *Amyotroph Lat Scler*.

2008;9(1):4-15. doi: 10.1080/17482960701856300. PubMed ID: 18273714; PubMed

Central ID:

15.   Talmadge JE, Singh RK, Fidler IJ, Raz A. Murine models to evaluate novel and

conventional therapeutic strategies for cancer. *Am J Pathol*. 2007;170(3):793-804. doi:

10.2353/ajpath.2007.060929. PubMed ID: 17322365; PubMed Central ID: 1864878.

16.   Rogers CS, Stoltz DA, Meyerholz DK, Ostedgaard LS, Rokhlina T, Taft PJ, et al.

Disruption of the CFTR gene produces a model of cystic fibrosis in newborn pigs.

Science. 2008;321(5897):1837-41. doi: 10.1126/science.1163600. PubMed ID:

18818360; PubMed Central ID: PMC2570747.

17.   Flisikowska T, Merkl C, Landmann M, Eser S, Rezaei N, Cui X, et al. A porcine model

of familial adenomatous polyposis. *Gastroenterology*. 2012;143:1173-5. doi:

10.1053/j.gastro.2012.07.110. PubMed ID: 22864254; PubMed Central ID:

18.   Pezzulo AA, Tang XX, Hoegger MJ, Abou Alaiwa MH, Ramachandran S, Moninger TO,

et al. Reduced airway surface pH impairs bacterial killing in the porcine cystic fibrosis

lung. *Nature*. 2012;487(7405):109-13. doi: 10.1038/nature11130. PubMed ID: 22763554;

PubMed Central ID: PMC3390761.

19.   Maddalo D, Manchado E, Concepcion CP, Bonetti C, Vidigal JA, Han YC, et al. In vivo

engineering of oncogenic chromosomal rearrangements with the CRISPR/Cas9 system.

Nature. 2014;516(7531):423-7. doi: 10.1038/nature13902. PubMed ID: 25337876;

PubMed Central ID: PMC4270925.

20.   Sanchez-Rivera FJ, Papagiannakopoulos T, Romero R, Tammela T, Bauer MR, Bhutkar

A, et al. Rapid modelling of cooperating genetic events in cancer through somatic

genome editing. *Nature*. 2014;516(7531):428-31. doi: 10.1038/nature13906. PubMed ID:

25337879; PubMed Central ID: PMC4292871.

21. Day CP, Merlino G, Van Dyke T. Preclinical mouse cancer models: a maze of

opportunities and challenges. *Cell*. 2015;163(1):39-53. doi: 10.1016/j.cell.2015.08.068.

PubMed ID: 26406370; PubMed Central ID: PMC4583714.

22. Singh M, Murriel CL, Johnson L. Genetically engineered mouse models: closing the gap

between preclinical data and trial outcomes. *Cancer Res*. 2012;72(11):2695-700. doi:

10.1158/0008-5472.CAN-11-2786. PubMed ID: 22593194; PubMed Central ID:

23. Zitvogel L, Pitt JM, Daillere R, Smyth MJ, Kroemer G. Mouse models in

oncoimmunology. *Nat Rev Cancer*. 2016;16(12):759-73. doi: 10.1038/nrc.2016.91.

PubMed ID: 27687979; PubMed Central ID:

24. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience

research reporting: the ARRIVE guidelines for reporting animal research. *PLoS Biol*.

2010;8(6):e1000412. doi: 10.1371/journal.pbio.1000412. PubMed ID: 20613859;

PubMed Central ID: PMC2893951.

25. National Institutes of Health. Principles and Guidelines for Reporting Preclinical

Research [February 5, 2016]. Available from: https://www.nih.gov/research-

training/rigor-reproducibility/principles-guidelines-reporting-preclinical-research.

26. National Institutes of Health. Enhancing Reproducibility through Rigor and

Transparency. NOT-OD-15-103. June 9, 2015. Available from:

http://grants.nih.gov/grants/guide/notice-files/NOT-OD-15-103.html.

27.   Committee for the Update of the Guide for the Care and Use of Laboratory Animals.

578   *Guide for the Care and Use of Laboratory Animals*. Washington, DC: The National

579   Academies Press; 2011.

580   28.   American Veterinary Medical Association Panel on Euthanasia. *AVMA Guidelines for the*

581   *Euthanasia of Animals: 2013 Edition*. Schaumberg, IL: American Veterinary Medical

582   Association; 2013.

583   29.   Adam SJ, Rund LA, Kuzmuk KN, Zachary JF, Schook LB, Counter CM. Genetic

584   induction of tumorigenesis in swine. *Oncogene*. 2007;26(7):1038-45. doi:

585   10.1038/sj.onc.1209892. PubMed ID: 16964292; PubMed Central ID:

586   30.   Schook LB, Collares TV, Hu W, Liang Y, Rodrigues FM, Rund LA, et al. A Genetic

587   Porcine Model of Cancer. *PLOS ONE*. 2015;10(7):e0128864. doi:

588   10.1371/journal.pone.0128864. PubMed ID: 26132737; PubMed Central ID:

589   PMC4488487.

590   31.   Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T)

591   method. *Nat Protoc*. 2008;3(6):1101-8. Epub 2008/06/13. PubMed ID: 18546601;

592   PubMed Central ID:

593   32.   Grinnell F, Zhu M, Carlson MA, Abrams JM. Release of mechanical tension triggers

594   apoptosis of human fibroblasts in a model of regressing granulation tissue. *Exp Cell Res*.

595   1999;248(2):608-19. Epub 1999/05/01. doi: 10.1006/excr.1999.4440. PubMed ID:

596   10222153; PubMed Central ID:

597   33.   Macpherson I, Montagnier L. Agar Suspension Culture for the Selective Assay of Cells

598   Transformed by Polyoma Virus. *Virology*. 1964;23:291-4. Epub 1964/06/01. PubMed ID:

599   14187925; PubMed Central ID:

600   34.   Liang CC, Park AY, Guan JL. In vitro scratch assay: a convenient and inexpensive

601   method for analysis of cell migration in vitro. *Nat Protoc*. 2007;2(2):329-33. Epub

602   2007/04/05. doi: 10.1038/nprot.2007.30. PubMed ID: 17406593; PubMed Central ID:

603   35.   Tsutsumida H, Swanson BJ, Singh PK, Caffrey TC, Kitajima S, Goto M, et al. RNA

604   interference suppression of MUC1 reduces the growth rate and metastatic phenotype of

605   human pancreatic cancer cells. *Clin Cancer Res*. 2006;12(10):2976-87. Epub 2006/05/19.

606   doi: 10.1158/1078-0432.CCR-05-1197. PubMed ID: 16707592; PubMed Central ID:

607   36.   Gioviale MC, Damiano G, Montalto G, Buscemi G, Romano M, Lo Monte AI. Isolation

608   and culture of beta-like cells from porcine Wirsung duct. *Transplant Proc*.

609   2009;41(4):1363-6. doi: 10.1016/j.transproceed.2009.02.062. PubMed ID: 19460560;

610   PubMed Central ID:

611   37.   Govindan R, Ding L, Griffith M, Subramanian J, Dees ND, Kanchi KL, et al. Genomic

612   landscape of non-small cell lung cancer in smokers and never-smokers. *Cell*.

613   2012;150(6):1121-34. doi: 10.1016/j.cell.2012.08.024. PubMed ID: 22980976; PubMed

614   Central ID: PMC3656590.

615   38.   Kandoth C, McLellan MD, Vandin F, Ye K, Niu B, Lu C, et al. Mutational landscape and

616   significance across 12 major cancer types. *Nature*. 2013;502(7471):333-9. Epub

617   2013/10/18. doi: 10.1038/nature12634. PubMed ID: 24132290; PubMed Central ID:

618   39.   Stephens PJ, Tarpey PS, Davies H, Van Loo P, Greenman C, Wedge DC, et al. The

619   landscape of cancer genes and mutational processes in breast cancer. *Nature*.

620   2012;486(7403):400-4. PubMed ID: 22722201; PubMed Central ID:

621   40.   Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Jr., Kinzler KW.

622   Cancer genome landscapes. *Science*. 2013;339(6127):1546-58. doi:

623   10.1126/science.1235122. PubMed ID: 23539594; PubMed Central ID: 3749880.

624   41.   Waddell N, Pajic M, Patch AM, Chang DK, Kassahn KS, Bailey P, et al. Whole genomes

625   redefine the mutational landscape of pancreatic cancer. *Nature*. 2015;518:495-501. doi:

626   10.1038/nature14169. PubMed ID: PMC4523082; PubMed Central ID:

627   42.   Hingorani SR, Wang L, Multani AS, Combs C, Deramaudt TB, Hruban RH, et al.

628   Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely

629   metastatic pancreatic ductal adenocarcinoma in mice. *Cancer Cell*. 2005;7(5):469-83.

630   Epub 2005/05/17. doi: 10.1016/j.ccr.2005.04.023. PubMed ID: 15894267; PubMed

631   Central ID:

632   43.   Groenen MA, Archibald AL, Uenishi H, Tuggle CK, Takeuchi Y, Rothschild MF, et al.

633   Analyses of pig genomes provide insight into porcine demography and evolution. *Nature*.

634   2012;491(7424):393-8. doi: 10.1038/nature11622. PubMed ID: 23151582; PubMed

635   Central ID:

636   44.   Rodrigues F, Hu W, Rund L, Liang Y, Counter C, Schook L. Development of the "Onco-

637   Pig": An Inducible Transgenic Porcine Model for Human Cancer [poster]. 2013

638   Brazilian Genetics Conference; URL:

639   http://www.dbs.illinois.edu/comparativegenomics/userfiles/file/ocnopigPoster_Brazilian

640   %20Genetics%20Conference2013.pdf.

45.   . Conference Proceedings: Swine in Biomedical Research; July 6-8, 2014; Raleigh, NC.

46. DuPage M, Dooley AL, Jacks T. Conditional mouse lung cancer models using adenoviral

or lentiviral delivery of Cre recombinase. *Nat Protoc*. 2009;4(7):1064-72. doi:

10.1038/nprot.2009.95. PubMed ID: 19561589; PubMed Central ID:

47. Bailey JM, Swanson BJ, Hamada T, Eggers JP, Singh PK, Caffery T, et al. Sonic

hedgehog promotes desmoplasia in pancreatic cancer. *Clin Cancer Res*.

2008;14(19):5995-6004. Epub 2008/10/03. doi: 10.1158/1078-0432.CCR-08-0291.

PubMed ID: 18829478; PubMed Central ID: PMC2782957.

48. Handra-Luca A, Hong SM, Walter K, Wolfgang C, Hruban R, Goggins M. Tumour

epithelial vimentin expression and outcome of pancreatic ductal adenocarcinomas. *Br J

Cancer*. 2011;104(8):1296-302. Epub 2011/03/31. doi: 10.1038/bjc.2011.93. PubMed ID:

21448168; PubMed Central ID: PMC3078601.

49. Kheirabadi BS, Mace JE, Terrazas IB, Fedyk CG, Estep JS, Dubick MA, et al. Safety

evaluation of new hemostatic agents, smectite granules, and kaolin-coated gauze in a

vascular injury wound model in swine. *J Trauma*. 2010;68(2):269-78. doi:

10.1097/TA.0b013e3181c97ef1. PubMed ID: 20154537; PubMed Central ID:

50. Cooper DK, Ekser B, Ramsoondar J, Phelps C, Ayares D. The role of genetically

engineered pigs in xenotransplantation research. *J Pathol*. 2016;238(2):288-99. doi:

10.1002/path.4635. PubMed ID: 26365762; PubMed Central ID: PMC4689670.

51. Sullivan TP, Eaglstein WH, Davis SC, Mertz P. The pig as a model for human wound

healing. *Wound Repair Regen*. 2001;9(2):66-76. doi: 10.1046/j.1524-475x.2001.00066.x.

PubMed ID: 11350644; PubMed Central ID:

52. Swindle MM, Makin A, Herron AJ, Clubb FJ, Jr., Frazier KS. Swine as models in

biomedical research and toxicology testing. *Vet Pathol*. 2012;49(2):344-56. Epub

2011/03/29. doi: 10.1177/0300985811402846. PubMed ID: 21441112; PubMed Central

ID:

53.   Xiangdong L, Yuanwu L, Hua Z, Liming R, Qiuyan L, Ning L. Animal models for the

atherosclerosis research: a review. *Protein Cell*. 2011;2(3):189-201. doi: 10.1007/s13238-

011-1016-3. PubMed ID: 21468891; PubMed Central ID: PMC4875304.

54.   Gouadon E, Moore-Morris T, Smit NW, Chatenoud L, Coronel R, Harding SE, et al.

Concise Review: Pluripotent Stem Cell-Derived Cardiac Cells, A Promising Cell Source

for Therapy of Heart Failure: Where Do We Stand? *Stem Cells*. 2016;34(1):34-43. doi:

10.1002/stem.2205. PubMed ID: 26352327; PubMed Central ID:

55.   Kuzmuk KN, Schook LB. Pigs as a Model for Biomedical Sciences. In: Rothschild MF,

Ruvinsky A, editors. *The Genetics of the Pig*. Oxfordshire, UK: CABI; 2011. p. 426-44.

56.   Emes RD, Goodstadt L, Winter EE, Ponting CP. Comparison of the genomes of human

and mouse lays the foundation of genome zoology. *Hum Mol Genet*. 2003;12(7):701-9.

doi: 10.1093/hmg/ddg078. PubMed ID: 12651866; PubMed Central ID:

57.   Walters E, Wolf E, Whyte J, Mao J, Renner S, Nagashima H, et al. Completion of the

swine genome will simplify the production of swine as a large animal biomedical model.

BMC Med Genom. 2012;5(1):55. doi: 10.1186/1755-8794-5-55. PubMed ID: 23151353;

PubMed Central ID:

58.   Dawson HD, McAnulty P, Dayan A, Ganderup N, Hastings K. A comparative assessment

of the pig, mouse and human genomes. In: McAnulty P, Dayan A, Ganderup N, Hastings

K, editors. *The Minipig in Biomedical Research*. Boca Raton, FL: CRC Press; 2012. p.

323-42.

59.   Dawson HD, Smith AD, Chen C, Urban JF, Jr. An in-depth comparison of the porcine,

murine and human inflammasomes; lessons from the porcine genome and transcriptome.

Vet Microbiol. 2017;202:2-15. Epub 2016/06/21. doi: 10.1016/j.vetmic.2016.05.013.

PubMed ID: 27321134; PubMed Central ID:

60.   Schachtschneider KM, Madsen O, Park C, Rund LA, Groenen MA, Schook LB. Adult

porcine genome-wide DNA methylation patterns support pigs as a biomedical model.

BMC Genomics. 2015;16:743. Epub 2015/10/07. doi: 10.1186/s12864-015-1938-x.

PubMed ID: 26438392; PubMed Central ID: PMC4594891.

61.   Vodicka P, Smetana K, Jr., Dvorankova B, Emerick T, Xu YZ, Ourednik J, et al. The

miniature pig as an animal model in biomedical research. *Ann N Y Acad Sci*.

2005;1049:161-71. Epub 2005/06/21. doi: 10.1196/annals.1334.015. PubMed ID:

15965115; PubMed Central ID:

62.   Swindle MM, Smith AC. *Swine in the Laboratory: Surgery, Anesthesia, Imaging, and*

Experimental Techniques. 3rd ed. Boca Raton, FL: CRC Press; 2016.

63.   Spurlock ME, Gabler NK. The development of porcine models of obesity and the

metabolic syndrome. *J Nutr*. 2008;138(2):397-402. Epub 2008/01/22. PubMed ID:

18203910; PubMed Central ID:

64.   Röthkotter HJ. Anatomical particularities of the porcine immune system--a physician's

view. *Dev Comp Immunol*. 2009;33(3):267-72. doi: 10.1016/j.dci.2008.06.016. PubMed

ID: 18775744; PubMed Central ID:

65.   Bailey M. The mucosal immune system: recent developments and future directions in the

pig. *Dev Comp Immunol*. 2009;33(3):375-83. doi: 10.1016/j.dci.2008.07.003. PubMed

ID: 18760299; PubMed Central ID:

66.   Fairbairn L, Kapetanovic R, Sester DP, Hume DA. The mononuclear phagocyte system

of the pig as a model for understanding human innate immunity and disease. *J Leukoc*

Biol. 2011;89(6):855-71. doi: 10.1189/jlb.1110607. PubMed ID: 21233410; PubMed

Central ID:

67.   Meurens F, Summerfield A, Nauwynck H, Saif L, Gerdts V. The pig: a model for human

infectious diseases. *Trends Microbiol*. 2012;20(1):50-7. doi: 10.1016/j.tim.2011.11.002.

PubMed ID: 22153753; PubMed Central ID:

68.   Petersen B, Carnwath JW, Niemann H. The perspectives for porcine-to-human

xenografts. *Comp Immunol Microbiol Infect Dis*. 2009;32(2):91-105. doi:

10.1016/j.cimid.2007.11.014. PubMed ID: 18280567; PubMed Central ID:

69.   Ferrer J, Scott WE, 3rd, Weegman BP, Suszynski TM, Sutherland DE, Hering BJ, et al.

Pig pancreas anatomy: implications for pancreas procurement, preservation, and islet

isolation. *Transplantation*. 2008;86(11):1503-10. doi: 10.1097/TP.0b013e31818bfda1.

PubMed ID: 19077881; PubMed Central ID: 2704055.

70.   Gun G, Kues WA. Current progress of genetically engineered pig models for biomedical

research. *Biores Open Access*. 2014;3(6):255-64. doi: 10.1089/biores.2014.0039.

PubMed ID: 25469311; PubMed Central ID: 4245835.

71.   Fan N, Lai L. Genetically modified pig models for human diseases. *J Genet Genomics*.

2013;40(2):67-73. doi: 10.1016/j.jgg.2012.07.014. PubMed ID: 23439405; PubMed

Central ID:

72.   Prather RS, Lorson M, Ross JW, Whyte JJ, Walters E. Genetically engineered pig models

for human diseases. *Annu Rev Anim Biosci*. 2013;1:203-19. doi: 10.1146/annurev-animal-

031412-103715. PubMed ID: 25387017; PubMed Central ID:

73.   Luo Y, Li J, Liu Y, Lin L, Du Y, Li S, et al. High efficiency of BRCA1 knockout using

734   rAAV-mediated gene targeting: developing a pig model for breast cancer. *Transgenic*

735   *Res*. 2011;20(5):975-88. doi: 10.1007/s11248-010-9472-8. PubMed ID: 21181439;

736   PubMed Central ID:

737   74.   Luo Y, Bolund L, Sorensen CB. Pig gene knockout by rAAV-mediated homologous

738   recombination: comparison of BRCA1 gene knockout efficiency in Yucatan and

739   Gottingen fibroblasts with slightly different target sequences. *Transgenic Res*.

740   2012;21(3):671-6. doi: 10.1007/s11248-011-9563-1. PubMed ID: 22020980; PubMed

741   Central ID:

742   75.   Leuchs S, Saalfrank A, Merkl C, Flisikowska T, Edlinger M, Durkovic M, et al.

743   Inactivation and inducible oncogenic mutation of p53 in gene targeted pigs. *PLoS ONE*.

744   2012;7(10):e43323. doi: 10.1371/journal.pone.0043323. PubMed ID: 23071491; PubMed

745   Central ID:

746   76.   Sieren JC, Meyerholz DK, Wang XJ, Davis BT, Newell JD, Jr., Hammond E, et al.

747   Development and translational imaging of a TP53 porcine tumorigenesis model. *J Clin*

748   *Invest*. 2014;124(9):4052-66. doi: 10.1172/JCI75447. PubMed ID: 25105366; PubMed

749   Central ID: 4151205.

750   77.   Yang L, Guell M, Niu D, George H, Lesha E, Grishin D, et al. Genome-wide inactivation

751   of porcine endogenous retroviruses (PERVs). *Science*. 2015;350(6264):1101-4. Epub

752   2015/10/13. doi: 10.1126/science.aad1191. PubMed ID: 26456528; PubMed Central ID:

753   78.   Beraldi R, Meyerholz D, Savinov A, Kovács A, Weimer J, Dykstra J, et al. Genetic

754   Ataxia Telangiectasia porcine model phenocopies the multisystemic features of the

human disease. *Biochim Biophys Acta*. 2017:ePub ahead of print, 2017 Jul 23. doi:

10.1016/j.bbadis.2017.07.020. PubMed ID: 28746835; PubMed Central ID:

79.   Smit VT, Boot AJ, Smits AM, Fleuren GJ, Cornelisse CJ, Bos JL. KRAS codon 12

mutations occur very frequently in pancreatic adenocarcinomas. *Nucleic Acids Res*.

1988;16(16):7773-82. PubMed ID: 3047672; PubMed Central ID: PMC338489.

80.   Jones S, Zhang X, Parsons DW, Lin JC, Leary RJ, Angenendt P, et al. Core signaling

pathways in human pancreatic cancers revealed by global genomic analyses. *Science*.

2008;321(5897):1801-6. doi: 10.1126/science.1164368. PubMed ID: 18772397; PubMed

Central ID: PMC2848990.

81.   Muller PA, Vousden KH. p53 mutations in cancer. *Nat Cell Biol*. 2013;15(1):2-8. doi:

10.1038/ncb2641. PubMed ID: 23263379; PubMed Central ID:

82.   Olivier M, Hollstein M, Hainaut P. TP53 mutations in human cancers: origins,

consequences, and clinical use. *Cold Spring Harbor Perspectives in Biology*. 2010;2(1).

doi: 10.1101/cshperspect.a001008. PubMed ID: 20182602; PubMed Central ID:

83.   Blackford A, Serrano OK, Wolfgang CL, Parmigiani G, Jones S, Zhang X, et al. SMAD4

gene mutations are associated with poor prognosis in pancreatic cancer. *Clin Cancer Res*.

2009;15(14):4674-9. doi: 10.1158/1078-0432.CCR-09-0227. PubMed ID: 19584151;

PubMed Central ID: PMC2819274.

84.   Caldas C, Hahn SA, da Costa LT, Redston MS, Schutte M, Seymour AB, et al. Frequent

somatic mutations and homozygous deletions of the p16 (MTS1) gene in pancreatic

adenocarcinoma. *Nat Genet*. 1994;8(1):27-32. doi: 10.1038/ng0994-27. PubMed ID:

7726912; PubMed Central ID:

85. Bardeesy N, Cheng KH, Berger JH, Chu GC, Pahler J, Olson P, et al. Smad4 is

dispensable for normal pancreas development yet critical in progression and tumor

biology of pancreas cancer. *Genes Dev*. 2006;20(22):3130-46. doi: 10.1101/gad.1478706.

PubMed ID: 17114584; PubMed Central ID: PMC1635148.

86. Bardeesy N, Aguirre AJ, Chu GC, Cheng KH, Lopez LV, Hezel AF, et al. Both

p16(Ink4a) and the p19(Arf)-p53 pathway constrain progression of pancreatic

adenocarcinoma in the mouse. *Proc Natl Acad Sci U S A*. 2006;103(15):5947-52. doi:

10.1073/pnas.0601273103. PubMed ID: 16585505; PubMed Central ID: PMC1458678.

87. Flisikowska T, Kind A, Schnieke A. The new pig on the block: modelling cancer in pigs.

Transgenic Res. 2013;22(4):673-80. doi: 10.1007/s11248-013-9720-9. PubMed ID:

23748932; PubMed Central ID:

88. Ittmann M, Huang J, Radaelli E, Martin P, Signoretti S, Sullivan R, et al. Animal Models

of Human Prostate Cancer: The Consensus Report of the New York Meeting of the

Mouse Models of Human Cancers Consortium Prostate Pathology Committee. *Canc Res*.

2013;73(9):2718-36. doi: 10.1158/0008-5472.CAN-12-4213. PubMed ID: 23610450;

PubMed Central ID:

89. Luo Y, Kofod-Olsen E, Christensen R, Sorensen CB, Bolund L. Targeted genome editing

by recombinant adeno-associated virus (rAAV) vectors for generating genetically

modified pigs. *J Gen Genom*. 2012;39(6):269-74. doi: 10.1016/j.jgg.2012.05.004.

PubMed ID: 22749014; PubMed Central ID:

90. Schachtschneider K, Schwind R, Darfour-Oduro K, De A, Rund L, Singh K, et al. A

validated, transitional and translational porcine model of hepatocellular carcinoma.

799     *Oncotarget*. 2017:Epub ahead of print, 2017 Jun 29. Epub 2017 Jun 29. doi:

800     10.18632/oncotarget.18872. PubMed ID: 28733541; PubMed Central ID:

Acknowledgements

This study is the result of work supported in part with resources and the use of facilities at the Omaha VA Medical Center (Nebraska-Western Iowa Health Care System). Portions of this study were presented at the 108th Annual Meeting of the American Association for Cancer Research (AACR) in Washington, D.C., April 1-5, 2017. The authors would like to acknowledge the technical assistance of Gerri Siford, Chris Hansen, Kelly O'Connell, and Tom Caffrey. The authors also would like to thank Dr. Lawrence B. Schook and Dr. Laurie A. Rund at the University of Illinois for the gift of their plasmid that contained the $KRAS^{G12D}$ mutation, along with comments, insights and suggestions for its use in this project.

816 **Author Contributions**

Contributor Role	Description	Author
Conceptualization	Ideas; formulation or evolution of overarching research goals and aims.	MAC
Data Curation	Management activities to annotate (produce metadata), scrub data and maintain research data (including software code, where it is necessary for interpreting the data itself) for initial use and later reuse.	NR, JLC, JAG
Formal Analysis	Application of statistical, mathematical, computational, or other formal techniques to analyze or synthesize study data.	NR, MAC
Funding Acquisition	Acquisition of the financial support for the project leading to this publication.	MAC
Investigation	Conducting a research and investigation process, specifically performing the experiments, or data/evidence collection.	NR, JLC, JAG, MAC
Methodology	Development or design of methodology; creation of models.	NR, MAH, MAC
Project Administration	Management and coordination responsibility for the research activity planning and execution.	MAC
Resources	Provision of study materials, reagents, materials, patients, laboratory samples, animals, instrumentation, computing resources, or other analysis tools.	MAH, MAC
Software	Programming, software development; designing computer programs; implementation of the computer code and supporting algorithms; testing of existing code components.	na
Supervision	Oversight and leadership responsibility for the research activity planning and execution, including mentorship external to the core team.	MAH, MAC
Validation	Verification, whether as a part of the activity or separate, of the overall replication/reproducibility of results/experiments and other research outputs.	NR, JLC, JAG
Visualization	Preparation, creation and/or presentation of the published work, specifically visualization/data presentation.	NR, MAC
Draft Preparation	Creation and/or presentation of the published work, specifically writing the initial draft (including substantive translation).	NR, MAC
Review & Editing	Preparation, creation and/or presentation of the published work by those from the original research group, specifically critical review, commentary or revision – including pre- or post-publication stages.	NR, JLC, JAG, MAH, MAC

817
818 na = not applicable.
819

820 **Table 1. Tumorigenic cell lines and their expression products.**
821

Cell line	AcGFP1 protein	KRAS^{G12D} protein	p53^{R167H} protein	SMAD4 shRNA	p16^{Ink4A} shRNA
PGKP	+	+	+		
PGKPS	+	+	+	+	
PGKPSC*	+	+	+	+	+

823 All cell lines based on primary porcine pancreatic ductal epithelial cells. *Abbreviation key: **P** =
824 pancreatic ductal epithelium; **G** = GFP; **K** = KRAS; **P** = p53; **S** = SMAD4; **C** =
825 p16^{Ink4A}/CDKN2A.
826

Figure Legends

Fig. 1. Isolation and transduction of primary porcine pancreatic ductal epithelial cells. (A)

Phase image of cells isolated from porcine pancreatic duct, showing epithelial-like morphology

(bar = 1,000 μm). (B) Immunofluorescent staining for cytokeratin 19 in the cultured primary

cells (bar = 200 μm). (C) Immunoblot for the p53 mutant in nine different cells lines transduced

with the GKP virus (PGKP cells). Pancreas = wild type pancreatic ductal epithelial cells.

Representative blot of three separate experiments. (D-E) qPCR of KRAS and p53 mutants in the

nine PKGP cell lines. Each bar represents mean of three separate experiments. *p<0.05, **

p<0.03, ***p<0.01 (unpaired t-test, compared to wild type).

Fig. 2. Effect of SMAD4 and p16^{Ink4A} knockdown on transformation in the PGKP cell line.

(A) Soft agar assay and (B) migration assay in the nine GKP-transduced cell lines. Pancreas =

wild type pancreatic ductal epithelial cells. Each bar represents mean of three separate

experiments. (C) RT-PCR of SMAD4 and p16^{Ink4A}/CDKN2A mRNA in cell lines expressing

targeted or scramble shRNA. Representative blot of three separate experiments. (D) Soft agar

assay and (E) migration assay of selected cell lines (PGKP, PGKPS, and PGKPSC), showing the

in vitro effect of additional knockdown of SMAD4 ± p16^{Ink4A} on the transformation of GKP-

transduced pancreatic ductal epithelial cells.

Fig. 3. *In vitro* transformation assays comparing the PGKP, PGKPS, and PGKPSC cell

lines. (A) Cell culture population doubling time (count-based assay). Pancreas = wild type

pancreatic ductal epithelial cells. (B) Proliferation rate (metabolic dye-based assay), represented

as fold change, normalized to wild type cells. (C) Invasion (Matrigel®-based assay). Each bar or

data point represents mean of three separate experiments.

Fig. 4. *In vivo* tumorigenesis assays comparing the PGKP, PGKPS, and PGKPSC cell lines.

(A) Sample of a resected tumor from subcutaneous injection in nude mice; note size and

vascularity. (B) Tumor growth curve from the subcutaneous injection in nude mice. Each data

point represents mean of 10 mice. (C) Metastasis after orthotopic implantation in nude mice

Fig. 5. Tumor immunohistochemistry from subcutaneous nude mouse assay. Representative

images of primary tumors derived from the PGKP, PGKPS, and PGKPSC cell lines stained with

an array of epithelial markers, including E-cadherin (Ecad; B, G, L), Epithelial Cell Adhesion

Molecule (EpCAM; C, H, M), Pan-Cytokeratin (PanCK; D, I, N), and Cytokeratin-19 (CK19; E,

J, O); and also stained with mesenchymal markers, including α-Smooth Muscle Actin (SMA; P,

S, V), Vimentin (Vim; Q, T, W), and Type I Collagen (Col I; R, U, X). Bar = 200 μm.

Supporting Information

[upload pending]

Fig. S1. pIRES2-AcGFP1 Vector Information

Fig. S2. Onco-pig

Fig. S3. KRAS/p53 immunohistochemistry of subcutaneous murine tumors

Protocol: Isolation of Epithelial Cells from Porcine Pancreas

Table S1. Responses to the ARRIVE recommendations

Table S2. Responses to the NIH Preclinical Research Guidelines

Table S3. Primers and other short sequences

Table S4. Antibody information

Figure 1

Figure 2

A

B

$ Compared to pancreas
# Compared to PGKP
** p<0.03
*** p<0.01
**** p<0.005

C

* p < 0.05
** p<0.03
*** p<0.001

Figure 3

Figure 4

Figure 5

| | H&E | KRAS | p53 |

Supplementary Figure 3