Generation of tumorigenic porcine pancreatic ductal epithelial cells:
toward a large animal model of pancreatic cancer

Neeley Remmers[1,2], Jesse L. Cox[3], James A. Grunkemeyer[4], Shruthi Aravind[1,2], Christopher K.
Arkfeld[1,2], Michael A. Hollingsworth[4], Mark A. Carlson[1,2,5*]

[1]Department of Surgery, University of Nebraska Medical Center, Omaha, NE, USA

[2] Surgery Department, VA Nebraska-Western Iowa Health Care System, Omaha, NE, USA

[3]Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha,
NE, USA

[4]Eppley Cancer Institute, University of Nebraska Medical Center, Omaha, NE, USA

[5]Department of Genetics, Cell Biology and Anatomy, University of Nebraska Medical Center,
Omaha, NE, USA

14
15
16
17

18 *Corresponding author
19 Email: macarlso@unmc.edu (MAC)

20
21

Abstract

Background. A large animal model of pancreatic cancer would permit development of diagnostic and interventional technologies not possible in murine models, and also would provide a more biologically-relevant platform for penultimate testing of novel therapies, prior to human testing. Here, we describe our initial studies in the development of an autochthonous, genetically-defined, large animal model of pancreatic cancer, using immunocompetent pigs.

Methods. Primary pancreatic epithelial cells were isolated from pancreatic duct of domestic pigs; epithelial origin was confirmed with immunohistochemistry. Three transformed cell lines subsequently were generated from these primary cells using expression of oncogenic KRAS and dominant negative p53, with/without knockdown of p16 and SMAD4. We tested these cell lines using *in vitro* and *in vivo* assays of transformation and tumorigenesis.

Results. The transformed cell lines outperformed the primary cells in terms proliferation, population doubling time, soft agar growth, 2D migration, and Matrigel invasion, with the greatest differences observed when all four genes (KRAS, p53, p16, and SMAD4) were targeted. All three transformed cell lines grew tumors when injected subcutaneously in nude mice, demonstrating undifferentiated morphology, mild desmoplasia, and staining for both epithelial and mesenchymal markers. Injection into the pancreas of nude mice resulted in distant metastases, particularly when all four genes were targeted.

Conclusions. Tumorigenic porcine pancreatic cell lines were generated. Inclusion of four genetic

"hits" (KRAS, p53, p16, and SMAD4) appeared to produce the best results in our *in vitro* and *in*

vivo assays. The next step will be to perform autologous or syngeneic implantation of these cell

lines into the pancreas of immunocompetent pigs. We believe that the resultant large animal

model of pancreatic cancer could supplement existing murine models, thus improving preclinical

research on diagnostic, interventional, and therapeutic technologies.

Introduction

In the United States in 2016, approximately 53,000 people (48% female) were diagnosed with pancreatic cancer (~3.1% of all new cancer diagnoses), and there were ~42,000 deaths (49% female) from pancreatic cancer (~7.0% of all cancer deaths) [1,2]. The lifetime risk for pancreatic cancer is approximately 1 in 65 [1,2]. The incidence of pancreatic cancer has been gradually increasing since the mid-1990's, and generally is higher in the African-American population [1,2]. Pancreatic cancer is now is the fourth most common cause of cancer-related death in both men and women (after lung, prostate, and colorectal cancer, or lung, breast, and colorectal cancer, respectively) [1,2]. Despite apparent advances in treatment modalities and strategies [3], mortality from pancreatic cancer has not decreased [1,2]. As of 2012, the U.S. overall 5-year survival rate from pancreatic cancer was 7.7%; 5-year survival rates in localized, regional (nodal spread), or metastatic disease were 29.3, 11.1, and 2.6%, respectively [1,2]. So there remains a need for improved early diagnosis and therapy for pancreatic cancer.

Rodent models of pancreatic cancer may not accurately reflect human biology because of differences in physiology, anatomy, immune response, and genetic sequence between the two species [4-7]. Remarkably, only 5-8% of anti-cancer drugs that emerged from preclinical studies and entered clinical studies have been ultimately approved for clinical use [8,9]. The cause of this low approval rate is multifactorial, but likely includes the less-than-optimal predictive ability of some murine models (e.g., tumor xenografting into immunosuppressed mice) to determine the efficacy of various therapeutics in humans [4-6,10-14]. Moreover, there are a number of genes for which the genotype-phenotype relationship is discordant between mice and human, including $CFTR^{-/-}$ and $APC^{+/-}$ [15,16]. Incidentally, both the porcine $CFTR^{-/-}$ and $APC^{+/-}$ mutants reiterate

74   the human phenotype (pulmonary/GI disease and rectal polyposis, respectively) [*15-17*], in

75   contradistinction to the murine mutants.

76   In fairness, the recent trend to employ genetically-engineered mouse models (GEMM),

patient-derived xenografts (PDX), humanized mice, and *in vivo* site-directed CRISPR/Cas9

gene-edited mice in the testing of anti-cancer therapeutics may yield murine models with better

predictive ability than obtained with previously [*6,18-22*]. Though promising, these more

advanced murine models come with increased cost and complexity [*20*], and experience with

them still is early. Importantly, all murine models have limited utility in the development of

diagnostic or interventional technology that requires an animal subject whose size approximates

a human. So at present, there remains a need for improved animal models of pancreatic cancer

that (1) are more predictive of human response to anti-cancer therapy [*20,22*], and (2) are of

adequate size for development of specific technologies. Herein we describe some initial steps

taken in the development of a genetically-defined, autochthonous model of pancreatic cancer in

immunocompetent pigs.

Materials and Methods

Standards, rigor, reproducibility, and transparency

The animal studies of this report were designed, performed, and reported in accordance with both the ARRIVE recommendations (Animal Research: Reporting of *In Vivo* Experiments [*23*]) and the National Institutes of Health Principles and Guidelines for Reporting Preclinical Research [*24,25*]; for details, refer to Tables S1 and S2, respectively.

Materials and animal subjects

All reagents were purchased through Thermo Fisher Scientific (www.thermofisher.com) unless otherwise noted. Short DNA sequences for vector construction, mutagenesis, and amplification purposes are shown in Table S3. Antibody information is given in Table S4. Wild type domestic swine (male and female; age 3 months at time of purchase; 30-32 kg) were purchased from the Animal Research and Development Center of the University of Nebraska Lincoln (ardc.unl.edu). Athymic homozygous nude mice (Crl:NU(NCr)-*Foxn1^{nu}*; female; 8-9 weeks old) were purchased from Charles River Laboratories, Inc. (www.criver.com). Primers utilized in this report (Table S3) were synthesized by Integrated DNA Technologies, Inc. (www.idtdna.com). DNA sequencing was performed by the UNMC Genomics Core Facility (www.unmc.edu/vcr/cores/vcr-cores/genomics). Oncopigs [*26*] were purchased from the National Swine Resource and Research Center (NSRRC; www.nsrrc.missouri.edu).

Animal welfare

The animals utilized to generate data for this report were maintained and treated in

accordance with the *Guide for the Care and Use of Laboratory Animals* (8th ed.) from the

National Research Council and the National Institutes of Health [*27*], and also in accordance

with the Animal Welfare Act of the United States (U.S. Code 7, Sections 2131 – 2159). The

animal protocols pertaining to this manuscript were approved by the Institutional Animal Care

and Use Committee (IACUC) of the VA Nebraska-Western Iowa Health Care System (ID

numbers 00927, 00937, 00998, and 01017) or by the IACUC of the University of Nebraska

Medical Center (ID number 16-133-11-FC). All procedures were performed in animal facilities

approved by the Association for Assessment and Accreditation of Laboratory Animal Care

International (AAALAC; www.aaalac.org) and by the Office of Laboratory Animal Welfare of

the Public Health Service (grants.nih.gov/grants/olaw/olaw.htm). All surgical procedures were

performed under isoflurane anesthesia, and all efforts were made to minimize suffering.

Euthanasia was performed in accordance with the AVMA Guidelines for the Euthanasia of

Animals [*28*].

Porcine operative procedures

127   Further details on transgenic porcine subjects and related welfare, safety, husbandry,

128   operative procedures, and perioperative management are given in the Supporting Information.

Isolation of porcine pancreatic ductal epithelial cells

A detailed protocol for isolation of porcine pancreatic ductal epithelial cells is provided

in the Supporting Information (**Protocol S1**). In brief, the intact pancreas from male porcine

research subjects (age 5 mo) was harvested within 5 min after euthanasia, which was

accomplished by transection of the intrathoracic inferior vena cava and exsanguination while

under deep isoflurane anesthesia. These pigs had been on a protocol to study biomaterials within

skin wounds of the dorsum. The subject had not received any recent medication other the

anesthetics given for euthanasia; buprenorphine and cefovecin sodium had been given 4 weeks

prior to euthanasia. Immediately after explantation of the pancreas, the main pancreatic duct was

dissected sterilely with micro instruments from the organ body under 3.5x loupe magnification.

The duct then was mechanically digested by passage through a 70 μm sieve (Corning™ Sterile

Cell Strainers, Thermo Fisher Scientific, cat. no. 07-201-431).

The collected fragments were enzymatically digested with 1 mg/mL of Collagenase D at

37˚C for 1 h with gentle shaking. The cells were pelleted (600 g x 5 min), the supernatant was

discarded, the cell pellet was resuspended in whole media, which was defined as: DMEM (high

glucose with L-glutamine; Thermo Fisher Scientific, cat. no. 12100-046) supplemented with

10% (final concentration) fetal bovine serum (FBS; Thermo Fisher Scientific, cat. no. 26140079)

and 1% Antibiotic-Antimycotic Solution (Corning Inc., cat. no. 30-004-CI; cellgro.com). Cell

concentration in the resuspension was determined with a hemocytometer, and cells then were

diluted and pipetted into a 96-well plate (1-10 cell/well, 100-200 μL/well). After 5-7 days of

culture under standard conditions (whole media, 37˚C, 5% CO_2), wells that contained cells with

epithelial-like morphology were trypsinized and re-plated into a new 96-well plate, in order to

dilute out any fibroblasts. Cells were passaged in this fashion at least four times, until no cells

with fibroblast morphology were present. The resulting cells were passaged up to a T25 flask,

and maintained with standard conditions.

Generation of p53 and KRAS mutants and construction of expression vector

In order to generate the porcine p53^{R167H} mutant, wild-type p53 cDNA first was amplified from cervical lymph node tissue, which was obtained <5 min after euthanasia of a 4-month-old male domestic swine that had been on an unrelated research protocol. In brief, fresh nodal tissue was flash-frozen in liquid N$_2$ and then pulverized with a mortar and pestle, with continual addition of liquid N$_2$ during pulverization. The frozen powder then was placed into the first buffer solution of the QIAGEN RNEasy Mini Kit (cat. no. 74104; www.qiagen.com), and total RNA was isolated per the manufacturer's instructions.

After isolation, the total RNA underwent reverse transcription to cDNA with a Verso cDNA Synthesis Kit (Thermo Fisher Scientific, cat. no. AB1453A), per the manufacturer's instructions. The wild type p53 sequence was amplified out of the cDNA using the PCR primers shown in Table S3, which flanked the p53 cDNA with *Sal*I and *Bam*HI restriction sites. Successful amplification of the wild-type p53 cDNA was confirmed by inserting the amplified candidate sequence into the TOPO® vector (TOPO® TA Cloning® Kit; Invitrogen™/Life Technologies™, Thermo Fisher Scientific, cat. no. K202020) per the manufacturer's instructions, followed by sequencing.

Site-directed mutation of wild-type p53 into p53^{R167H} was performed using Agilent Technologies' QuickChange II Site-Directed Mutagenesis Kit (cat. no. 200523; www.genomics.agilent.com) with the mutagenic primers shown in Table S3, per the manufacturer's instructions. Presence of the p53^{R167H} mutation was verified by sequencing as described above. The multiple cloning site of a pIRES2-AcGFP1 Vector (Takara Bio USA, Inc.,

cat. no. 632435; www.clontech.com; manufacturer's vector information included as **Fig. S1**) was

cut with *Sal*I and *Bam*HI, and the p53^{R167H} sequence then was ligated into this plasmid.

The source of the porcine KRAS^{G12D} mutant was the plasmid used to generate the

p53/KRAS Oncopig [*26,29*]. The KRAS^{G12D} cDNA was amplified out of this plasmid with

primers (see Table S3) that flanked the sequence with *Xho*I and *Pst*I restriction sites. The

amplified product was inserted into the TOPO vector and verified by sequencing, as described

above. The above pIRES2-AcGFP1 Vector (already containing the p53^{R167H} sequence) then was

cut with *Xho*I and *Pst*I, and the KRAS^{G12D} sequence was ligated into this plasmid, producing a

pIRES2-AcGFP1 Vector which contained both mutant cDNAs within its multiple cloning site

(KRAS^{G12D} upstream).

The newly-constructed plasmid, hereafter designated as GKP (G = AcGFP1; K =

KRAS^{G12D}; P = p53^{R167H}), was transformed into One Shot™ Stbl3™ Chemically Competent *E.*

coli (Invitrogen™/Thermo Fisher Scientific, cat. no. C737303), per the manufacturer's

instructions, and plasmid DNA subsequently was isolated using a QIAGEN Plasmid Maxi Kit

(cat. no. 12162), per the manufacturer's instructions. This plasmid then was transfected into

Takara's Lenti-X™ 293T cells (Clontech, cat. no. 632180), using Takara's Xfect™ Transfection

Reagent (Clontech, cat. no. 631317), per the manufacturer's instructions, to generate infectious

lentiviral particles that would direct expression of AcGFP1, KRAS^{G12D}, and p53^{R167H} mutants in

transduced cells.

Generation of shRNA-expressing vectors

Short hairpin RNA (shRNA) constructs targeting the porcine SMAD4 and p16 genes

were created using InvivoGen's siRNA Wizard™ software (www.invivogen.com/sirnawizard).

Three targeting sequences for each gene along with scrambled controls initially were generated

and tested. The shRNA construct that demonstrated the best target knock-down in preliminary

experimentation (as verified by PCR, data not shown) was utilized for subsequent experiments

(see Table S3 for sequences ultimately selected for the shRNA constructs). Primers were cloned

into the psiRNA-h7SKhygro G1 and psiRNA-h7SKneo G1 vectors (InvivoGen, cat. no. ksirna3-

h21 and ksirna3-n21, respectively; www.invivogen.com) and plasmids then were isolated, all per

the manufacturer's protocol.

Cell transformations

Primary porcine pancreatic epithelial cells in T75 flasks were grown to 80% confluency

under standard conditions. The media then was exchanged with 2-3 mL of supernatant from non-

lysed Lenti-X™ 293T cells (containing GKP viral particles) with 2 μg/mL polybrene (cat. no.

TR1003, Thermo Fisher Scientific). After 24-48 h at 37°C, treated epithelial cells were re-seeded

into 6-well plates under standard conditions and grown to 80% confluency. An exchange with

whole media containing 2 μg/mL G418 aminoglycoside antibiotic then was performed; the G418

dose was chosen based on preliminary dose-response studies against non-treated epithelial cells.

After 24 h, a whole media exchange was done, and the presence of transduced cells was

determined with inverted GFP fluorescent microscopy of living cells. Subsequent transfections

for RNAi were done with the above plasmids employing shRNA sequences against SMAD4

and/or p16, and using the LyoVec™ reagent (InvivoGen, cat. no. lyec-12), all per the

manufacturer's protocol. Transfected cells then were selected for expression of the shRNA

vector using the appropriate aminoglycoside antibiotic (G418 or hygromycin B).

PCR

Cell and tissue RNA was isolated using the QIAGEN RNEasy Mini Kit. Purified RNA then was used to generate cDNA using the Verso cDNA Synthesis Kit. The Platinum® Blue PCR Supermix (Invitrogen™/Life Technologies, cat. no. 12580) subsequently was used for all PCR reactions. Amplified products were separated with agarose gel electrophoresis, and then visualized using a UV-light box. qPCR was performed using the PowerUp™ SYBR® Green Master Mix (Applied Biosystems™/ Thermo Fisher Scientific, cat. no. A25741) per manufacturer's protocol, and run on an Applied Biosystems™ 7500 Fast Dx Real-Time PCR Instrument. Fold changes in gene expression were calculated using the comparative C_T method [30]. All primers used are listed in Table S3.

Immunoblotting

Western blot analysis was performed to confirm overexpression of the mutant p53 protein (see Table S4 for a list of antibodies used), as previously described [31]. An antibody specific for the mutant KRAS protein was not commercially available. Antibody expression was visualized using the Li-Cor Odyssey Electrophoresis Imaging System (www.licor.com).

Soft agar assay

A standard soft agar assay [32] was used to determine anchorage independent growth. A base layer of 1% agarose was plated into 6-well plates. A total of 2,500 cells/well were mixed with 0.7% agarose and plated on top of the base layer. The plates were incubated under standard conditions for 21 days. The cells then were stained with crystal violet, and counted using an

inverted microscope. Cells were plated in triplicate, and total counts from all three wells were averaged.

Migration assay

A standard scratch assay (monolayer wounding) [*33*] was performed to determine cellular migration rate. Cells were plated in triplicate into 6-well plates. A horizontal scratch using a 10 µL pipet tip was made in each well. After washing away scratched-off cells, baseline images along the scratch were obtained, the plates were incubated under standard conditions, and subsequent images were captured at 3, 6, 9, 12, and 15 h after the initial scratch. ImageJ software (imagej.nih.gov/ij) was used to measure the distance between the two migrating cellular fronts (scratch edges) at 3-5 locations along the scratch. Average distance at each time point was plotted to generate the migratory rate (µm/h).

Invasion assay

BioCoat™ Matrigel™ Invasion Chambers (Corning™, Thermo Fisher Scientific, cat. no. 08-774) were plated with 50,000 cells (upper chamber) in triplicate, and incubated under standard conditions for 24 h. The media from the upper chamber then was removed, and any cells remaining in the upper chamber were removed using a cotton swab. Cells that had migrated to the bottom of the membrane were stained using a Kwik-Diff™ kit (Shandon™, Thermo Fisher Scientific, cat. no. 9990701). Membranes were mounted onto glass slides, and cells per high-power field were counted using ImageJ software.

Population doubling assay

Cells were plated in 6-well plates (20,000 cell/well), and cultured under standard

conditions. Triplicate plates then were trypsinized on days 1, 2, 3, 4, 6, and 8, and cells were

counted with a hemocytometer. Cell number *vs*. day was plotted to determine the day range in

which linear growth was achieved. The data from this linear growth phase were used to

determine population doubling time (DT) using the formula: $DT = (\Delta t) \times \ln(2) \div \ln(N_f/N_i)$

where Δt = time interval between initial and final cell count, N_f = cell count at final time, and N_i

= cell count at initial time.

Proliferation assay

Relative cell proliferation rates were assayed using an MTT (3-(4,5-dimethylthiazol-2-

yl)-2,5-diphenyltetrazolium bromide) assay kit (Vybrant™ MTT Cell Proliferation Assay Kit,

Invitrogen™, Thermo Fisher Scientific, cat. no. V13154). Cells were plated in triplicate in a 96-

well plate (5,000 cell/well), and cultured under standard conditions for 48 h. MTT reagent then

was added to the cells per the manufacturer's instructions, followed by addition of the solvent

solution 3.5 h later. Absorbance was measured with a plate reader 3.5 h after solvent addition.

Mean absorbance was normalized to absorbance from wild type pancreatic ductal epithelial cells

to calculate fold-difference in proliferation.

Immunofluorescence and immunohistochemistry

288   Antibodies used in immunofluorescent and immunohistochemical experiments are listed

289   in Table S4. Agilent Dako EnVision kits (www.agilent.com) were used for all IHC analyses per

290   the manufacturer's instructions.

Subcutaneous tumorigenic cell injection

Subcutaneous implantation of tumorigenic cells was performed as previously described [34], with some modifications. Transformed porcine pancreatic ductal epithelial cells (the three lineages described in Table 1) were trypsinized, counted, and resuspended in DMEM at a concentration of 1×10^7 viable cells/mL. Nude mice (N = 30; 100% female; maintained in microisolator cages with soft bedding and fed regular chow) were randomized into three treatment groups (representing each transformed cell line in Table 1; N = 10 mice per group, 100% female) using an online randomization tool. Mice then were injected with 5×10^6 cells (500 μL) into the right hind flank under brief isoflurane inhalational anesthesia, administered with a Matrx VMS® small animal anesthesia machine, within a small animal operating room. Tumors were allowed to grow for 6 weeks or until they reached 2 cm in diameter, as measured with a caliper, and then subjects were euthanized using an AVMA-approved [28] method of CO_2 asphyxiation. At necropsy all gross tumor was measured and collected, portions underwent formalin fixation and paraffin embedding, and sections subsequently underwent H&E or immunohistochemical staining as described above. An independent, a blinded pathologist analyzed the stained sections to determine whether tumors were epithelial in origin, and if they displayed malignant features.

Orthotopic tumorigenic cell injection

Orthotopic implantation of tumorigenic cells was performed as previously described [34] to analyze metastases and desmoplasia. In brief, transformed porcine pancreatic ductal epithelial cells (the three lineages described in Table 1) were trypsinized and counted, and 1×10^4 viable cells were suspended into 20 μL DMEM. Nude mice (N = 36; 100% female) housed as described

above were randomized into three treatment groups (representing each transformed cell line in

Table 1; N = 12 per group, 100% female) using an online randomization tool. The 20 µL cell

suspension then was injected with a 20-gauge needle into the pancreas of each nude mouse

through a 5 mm incision in the left upper quadrant, under isoflurane anesthesia within a small

animal operating room. Mice were euthanized 6 weeks after injection using CO_2 asphyxiation as

described above, and tumors and organs were harvested for gross and histologic analysis, as

described in the previous paragraph.

Statistics and power analysis

Data are reported as mean ± standard deviation. Groups of continuous data were

compared with ANOVA and the unpaired t-test. Categorical data were compared with the Fisher

or Chi square test. For the power analysis of the murine subcutaneous tumor implant assay,

tumor diameter was selected as the endpoint. Setting alpha = 0.05 and power = 0.8, ten mice per

group were needed across three groups to detect a difference in means of 30% with the standard

deviation estimated at 20% of the mean. In the orthotopic implantation assay, N = 10 mice per

treatment group across three treatment groups were needed to detect a 100% difference in effect

(+tumor) at a single metastatic site (with alpha set at 0.05 and power = 0.8); or, combining all

seven metastatic sites together, N = 10 mice per group were needed to detect a 40% difference in

effect.

Results

Isolation of primary porcine pancreatic ductal epithelial cells

Cells cultured from micro-dissected pancreatic ducts displayed epithelial morphology under phase microscopy and stained for CK19 (an established marker of pancreatic ductal epithelium [*35*]; **Fig. 1A-B**). Based on these results, we were confident that we had a population of pancreatic epithelial cells that we could use to generate tumorigenic cell lines.

Generation of tumorigenic cell lines

In order to transform our primary porcine pancreatic ductal epithelial cells, we first generated a lentiviral construct containing *KRAS*^{G12D} and *TP53*^{R167H}, genes previously identified [*29*] as the porcine equivalents to the mutant *KRAS* and *TP53* which are present in multiple human cancers [*36-40*]; in the mouse, expression of these mutants was the basis for the KRAS/p53 genetically engineered murine model of pancreatic cancer [*41*]. For our model, we chose to use a lentiviral platform for the vector, because its genome would be large enough to accommodate insertion of both mutant genes; in addition, we believed that a lentivirus would be optimal for transforming primary cells.

Since initial sequencing of the porcine genome has been accomplished [*42*], we were able to utilize the National Library of Medicine's nucleotide BLAST® database (blast.ncbi.nlm.nih.gov/blast.cgi) to determine the porcine genetic equivalents for human *SMAD4* and *CDKN2A*. We then designed primers (Table S3) to amplify these two genes from genomic DNA isolated from skin of a healthy domestic pig. We sequenced our amplification products,

and accessed the BLAST® database to confirm that our products aligned with the porcine

SMAD4 and *CDKN2A* gene sequences. We then proceeded to generate lentiviral constructs to

transform primary porcine pancreatic ductal epithelial cells into cell lines expressing various

combinations of mutant KRAS and p53, SMAD4 shRNA, and p16^{Ink4A} shRNA (see cell line

definitions in Table 1), as described under Materials and Methods.

Primary porcine epithelial cells next were transduced with the GKP lentivirus to generate

transformed cell lines. Overexpression of KRAS^{G12D} and p53^{R167H} was confirmed in these cell

lines with qPCR and immunoblotting (**Fig. 1C-E**). Of note, we could not obtain a reliable

antibody to detect porcine KRAS with immunoblotting, so we had to rely on qPCR results and

expression of GFP as markers of KRAS^{G12D} expression. Preliminary *in vitro* analyses to probe

the tumorigenic properties of these GKP-transformed cell lines demonstrated modest increases in

soft agar colony formation and migration speed over wild type cells (**Fig. 2A & B**). Interestingly,

the cell line (2.22) with the best performance in the soft agar and migration assays had only

modest overexpression of KRAS^{G12D} and p53^{R167H} (**Fig. 1 & 2**); in contrast, the cell lines with

the highest mutant overexpression performed relatively poorly in these *in vitro* assays of

"tumorigenesis" (i.e., evidence of transformed behavior in cell culture).

While *in vitro* experiments were being performed, *in vivo* pancreatic tumor induction was

attempted utilizing a transgenic mini-pig available from the NSRRC. Known as the "Oncopig,"

[*26*], this subject carries an LSL-cassette containing the dominant negative *TP53*^{R167H} and the

activated *KRAS*^{G12D} sequences [*26,43,44*]; i.e., this subject is the porcine analog of the

KRAS/p53 mouse [*41,45*]. As demonstrated previously, site-specific expression of Cre

recombinase in the Oncopig resulted in localized p53 inhibition and KRAS activation, while

subcutaneous injection of AdCre produced mesenchymal tumors at the injection sites [*26*]. We

injected Cre recombinase into the pancreas of five Oncopigs (**Protocol S2**; **Fig. S2**; **Tables S5**

and **S6**). After four months, we observed no gross tumors. However, there was

immunohistochemical evidence of transgene expression at the pancreatic injection sites along

with numerous microscopic proliferative lesions with desmoplastic features (**Fig. S2**).

Based on the modest evidence of *in vitro* transformation and the lack of gross *in vivo*

tumorigenesis using expression of p53^{R167H} and KRAS^{G12D} only, we decided that additional

oncogenic stress might be helpful to increase the tumor-like properties of transformed pancreatic

ductal epithelial cells. Utilizing cell line 2.22 (hereafter referred to PGKP; see Table 1), which

had relatively good performance in the soft agar and migration assays (**Fig. 2A & B**), sequential

transduction with lentiviral constructs expressing shRNA against SMAD4 and then p16^{Ink4A} was

performed to generate cell lines PGKPS and PGKPSC (see Table 1), respectively. RT-PCR then

was used to confirm knockdown of the targeted transcripts in these two cell lines (**Fig. 2C**).

In vitro tumorigenic properties of transformed cells

The *in vitro* "tumorigenic" properties of the PGKP, PGKPS, and PGKPSC cell lines first

were compared with the soft agar and migration assays (**Fig. 2D & E**). Addition of SMAD4 ±

p16^{Ink4A} knockdown enhanced the ability of transformed cells to form colonies in soft agar and

increased their migration speed (2D wounding assay), particularly when both transcripts were

targeted (i.e., the PGKPSC line). We then compared population doubling time, proliferation

(metabolic dye conversion) , and Matrigel® invasion ability among the three transformed lines

with respect to wild type cells (**Fig. 3A-C**). Both the PGKPS and PGKPSC cell lines had greater

proliferation and invasive ability compared to either wild type cells or the PGKP cell line (**Fig.**

3B, C). The doubling time for all three transformed cell lines was approximately the same at ~15

h, compared to the ~4 d doubling time of wild type cells (**Fig. 3A**). Based on the *in vitro* assays

of tumorigenesis, we suspected that all three of our transformed cell lines had the potential to

form tumors *in vivo*, albeit to varying degrees. We subsequently decided to compare the *in vivo*

tumorigenicity among all three cell lines in an immunodeficient mouse model.

Subcutaneous and orthotopic tumor transplants in immunodeficient

mice

In order to determine if our cell lines retained their tumorigenic properties *in vivo*, we

utilized homozygous athymic mice to generate subcutaneous and orthotopic cell implantation

models. The subcutaneous model was used to assess *in vivo* tumor growth. All three cell lines

grew sizeable tumors (>1 cm diameter) within 6 weeks; growth rates were not statistically

different among the three cell lines (**Fig. 4A & B**). The subcutaneous tumors were well-

vascularized and mucinous in gross appearance (**Fig. 4A**). Confirmation of GKP lentiviral

transduction was demonstrated with immunohistochemistry of the p53^{R167H} mutant in

subcutaneous tumors (**Fig. S3**).

Tumors from the subcutaneous implant model then underwent immunohistochemical

staining with an array of epithelial and mesenchymal markers (**Fig. 5**). The distribution of

staining for the epithelial markers (E-cadherin, epithelial cell adhesion molecule, pan-

cytokeratin, cytokeratin-19) generally was more diffuse than the mesenchymal marker staining.

In some regions the epithelial marker staining was clustered and intense. The overall abundance

of staining for the epithelial markers appeared greater in the PGKPS and PGKPSC lines with

respect to the PGKP line. The distribution of staining for the mesenchymal markers (α-smooth

muscle actin, vimentin, and type I collagen) was variable, sometimes appearing in cords or

strands in some sections, reminiscent of the desmoplastic reaction in human pancreatic cancer

[*46,47*]. In other sections, mesenchymal marker staining was minimal. Similar to the epithelial

markers, the overall abundance of staining for the mesenchymal markers appeared greater in the

PGKPS and PGKPSC lines compared to the PGKP line.

With the immunohistochemical studies of the subcutaneous tumors demonstrating some

epithelial characteristics, the nude mouse orthotopic implantation model was used next to assess

the metastatic potential of all three cell lines. The percentage of mice implanted with each cell

line that subsequently developed metastasis in the small bowel, diaphragm, liver, lung, lymph

node, peritoneum, or spleen is shown in **Fig. 4C.** The degree of metastatic spread was minimal;

nodal tissue was the only metastatic site common to all three lines. The PGKPCS cell line

exhibited the greatest array of metastatic spread (p < 0.002, Chi-square), with the most common

site being the spleen (though it was not clear in two of five mice with splenic disease after

PGKPSC implantation whether the spleen was involved with extension from the primary tumor,

or from direct seeding, or whether these were true metastases). Interestingly, one of the two

subjects with liver metastasis after PGKPSC implantation had primary tumor within the

gallbladder, rather than within the pancreas. Similar to the subcutaneous tumor model, most

tumors in the orthotopic model were well-vascularized and mucinous in gross appearance.

Discussion

Our overall goal with this project was to generate tumorigenic pancreatic cell lines that could be used in an immunocompetent porcine model of pancreatic cancer. Current murine models for pancreatic cancer will continue to be helpful, particularly for the study of molecular mechanisms. However, murine models are limited in their ability to replicate human biology and size, so a large animal model of pancreatic cancer likely would enhance our ability to develop and test new diagnostic and treatment modalities for this disease. The data presented herein demonstrated that wild type porcine pancreatic ductal epithelium can be transformed with modulation of common tumor-associated target genes, and that these transformed cells subsequently can grow tumors in immunodeficient mice. These data provide a pathway for the construction of an autochthonous porcine model of pancreatic cancer, namely, orthotopic implantation of tumorigenic pancreatic cells. Proof-of-principle for tumor growth in pigs using subcutaneous implantation of *ex-vivo* transformed autologous fibroblasts was demonstrated in 2007 [*29*].

Porcine biomedical models have been used for decades in the fields of trauma and hemostasis [*48*], xenotransplantation [*49,50*], dermal healing [*51*], toxicology [*52*], atherosclerosis [*53*], and cardiac regeneration [*54*]; the utility of these models is growing. A porcine genome map was generated in 2012 [*42*], and further coverage, annotation, and confirmation is ongoing [*55,56*]. Porcine-centered online tools and databases are now available [*57*]. Genetic manipulation of pigs (including knockouts, tissue-specific transgenics, inducible expression [*29,58-65*]) with similar tools as used in the mouse is becoming more routine, with

new gene-edited porcine models emerging in 2015-2017 for diseases such as atherosclerosis,

cystic fibrosis, Duchenne muscular dystrophy, and ataxia telangiectasia [44,66,67].

The rationale to build a porcine model of pancreatic cancer is (1) to have a platform for

diagnostic/therapeutic device development otherwise not achievable in murine models; and (2) to

have a highly predictive preclinical model in which anti-cancer therapies (including

immunotherapies) could be vetted/optimized prior to a clinical trial [68]. The rationale to use the

pig in this modeling effort is that this species mimics human genomics [55,69-72], epigenetics

[73], physiology [52,69,74,75], metabolism [69,75,76], inflammation and immune response

[72,77-81], and size [75,82] remarkably well (in particular, better than mice), with reasonable

compromises towards cost and husbandry [75]. So based on the pig's relatively large size and its

proven track record in replicating human biology which, incidentally, is a demonstrably better

replication than can be obtained with rodents, we selected swine as the model organism for this

pancreatic cancer project.

Research on immunocompetent large animal cancer models [83-85] includes prostate

cancer, for which there is a canine model [86]. In addition, in 2012 a group in Munich reported

the engineering of (i) an *APC* mutant pig that developed rectal polyposis [16,87] and (ii) a pig

with Cre-inducible p53 deficiency [63]. This group subsequently determined that their p53-null

subjects (*TP53*$^{R167H/R167H}$) developed osteosarcoma by age 7-8 months [88]. Other p53-deficient

pigs have been engineered since this initial report [64,89]; in the report from Iowa, half (5 out of

10) of p53-deficient (*TP53*$^{R167H/R167H}$) pigs developed lymphoma or osteogenic tumor at age 6-18

months [64]. A group in Denmark reported the creation of a *BRCA* mutant pig in 2012. [90].

A KRAS/p53 "Oncopig" was reported in 2015 [26,43,44,84]. This subject has a somatic

LSL-cassette that can express dominant negative p53 (R167H mutation) and activated KRAS

(G12D mutation); i.e., the porcine analog of the KRAS/p53 mouse [*41,45*]. Site-specific

expression of Cre recombinase in the Oncopig resulted in localized p53 inhibition and KRAS

activation; subcutaneous injection of AdCre produced mesenchymal tumors at the injection sites

[*26*]. In 2017, initial work was published on a Oncopig-based model of hepatocellular carcinoma

[*91*]. Also in 2017, another genetic porcine model of intestinal neoplasia was reported [*92*],

utilizing inducible expression of KRAS^{G12D}, c-Myc, SV40 large T antigen, and retinoblastoma

protein (pRb). One out of three pigs total in this model developed duodenal neuroendocrine

carcinoma with lymph node metastasis at two months after induction. A porcine model of

pancreatic cancer has not yet been reported, other than some preliminary data presented by us in

2017 [*93*].

The process we used for developing transformed porcine pancreatic ductal epithelial cell

lines for future orthotopic implantation was somewhat iterative, in that we modified our strategy

along the way based on our early results. Commonly mutated genes in pancreatic cancer include

KRAS [*94,95*] and *TP53* [*95-97*]. In mice, somatic activation of KRAS via the G12D mutation

(*KRAS*^{G12D}) produced widely metastatic pancreatic tumors; survival duration in these subjects

decreased further with p53 inactivation [*41*]. Based on this murine model, and the published

success with subcutaneous tumor induction in the KRAS/p53 Oncopig [*26*], we elected to

transform pancreatic ductal epithelial cells with expression of activated KRAS^{G12D} and p53^{R167H}

only. However, the initial results from our *in vitro* transformation assays with these two gene

edits (i.e., the PGKP cell line of Table 1) were somewhat underwhelming. Combined with the

finding of no gross tumor four months after pancreatic AdCre injection in five Oncopig subjects,

we decided that additional genetic "hits" might be necessary for transformation of porcine

pancreatic ductal epithelial cells. Of note, Schook et al. [*29*] found that porcine dermal

fibroblasts required six genetic edits (human telomerase reverse transcriptase, dominant negative

p53, cyclin D1, activated cyclin dependent kinase, oncogenic c-Myc, and oncogenic H-Ras) to

optimize the tumorigenic phenotype of this particular cell.

Other commonly mutated genes in pancreatic cancer include *SMAD4* [*95,98*] and

CDKN2A [*95,99*]. Deletion of *SMAD4* or *CDKN2A* in a *KRAS*[G12D] murine pancreatic cancer

model enhanced tumor growth [*100,101*]. Based on these published data and our above

transformation results with just the *KRAS* and *TP53* edits, we elected to add knockdown of

SMAD4 and p16 to our list of hits for transformation of porcine pancreatic ductal epithelial cells.

Ultimately, all three of our cell lines (Table 1) demonstrated transformed behavior *in vitro* and

the ability to form tumors *in vivo* (nude mice), with perhaps some enhancement by the addition

of SMAD4 and p16 knockdown. In the future, we intend to utilize CRISPR/Cas9 editing to

disrupt the genes of these and/or other targets.

Although we utilized a relatively high number of cells to obtain tumor formation in our

nude mice experiments, we feel that this is due to the relative young age of our cell lines, which

did not undergo as many passages as other cell lines which have been used to study pancreatic

cancer. Thus, the transformed cell lines in this report likely did not undergo an inadvertent

selection for the fastest growing cells, as may occur in older, extensively-passaged lines.

Regarding the lack of desmoplasia in our murine xenograft model, we did not consider this

surprising, in that most immunodeficient murine models of pancreatic cancer do not recapitulate

the extent of desmoplasia or metastasis seen with human disease. We anticipate that future

experiments involving the implantation of transformed pancreatic ductal epithelial cells into wild

type swine (and thus into a "normal" microenvironment with "normal" immunoediting) will

produce tumors with desmoplasia and metastasis.

Of note, the pathologic findings in our five Oncopig subjects, while not macroscopically

tumorous, displayed tissue architecture reminiscent of desmoplasia. Regarding the lack of

macroscopic tumor in these transgenic pigs, it is conceivable that the induction process was not

optimal secondary to an inadequate dose of AdCre, inadequate recombinase activity, inadequate

tissue delivery of the enzyme, or some other technical issue. Our future plans in this respect will

involve additional attempts at induction of pancreatic tumor in the Oncopig with and without

introduction of some additional gene edits, such as local disruption of *SMAD4* and *CDKN2A*

(e.g., with *in vivo* CRISPR/Cas9 gene editing [*102,103*]).

References

1.   American Cancer Society. *Cancer Facts & Figures 2016*. Atlanta: American Cancer

Society; 2016. Available from:

http://www.cancer.org/research/cancerfactsstatistics/cancerfactsfigures2016/index.

2.   *SEER (Surveillance, Epidemiology, and End Results Program) Stat Fact Sheets:*

Pancreas Cancer. National Cancer Institute. Accessed December 12, 2016. URL:

http://seer.cancer.gov/statfacts/html/pancreas.html.

3.   National Comprehensive Cancer Network. Pancreatic Adenocarcinoma, Version 1.2018.

In: NCCN Clinical Practice Guidelines In Oncology (NCCN Guidelines®). NCCN; 27

April 2018. Available from: www.nccn.org.

4.   Seok J, Warren HS, Cuenca AG, Mindrinos MN, Baker HV, Xu W, et al. Genomic

responses in mouse models poorly mimic human inflammatory diseases. *PNAS* 2013;

110(9): 3507-12. PMID: 23401516. doi: 10.1073/pnas.1222878110.

5.   Begley CG, Ellis LM. Drug development: Raise standards for preclinical cancer research.

Nature 2012; 483(7391): 531-3. PMID: 22460880. doi: 10.1038/483531a.

6.   Cook N, Jodrell DI, Tuveson DA. Predictive in vivo animal models and translation to

clinical trials. *Drug Discov Today* 2012; 17(5-6): 253-60. PMID: 22493784. doi:

10.1016/j.drudis.2012.02.003.

7.   Le Magnen C, Dutta A, Abate-Shen C. Optimizing mouse models for precision cancer

prevention. *Nat Rev Cancer* 2016; 16(3): 187-96. PMID: 26893066. doi:

10.1038/nrc.2016.1.

8.   Reichert JM, Wenger JB. Development trends for new cancer therapeutics and vaccines.

566   *Drug Discov Today* 2008; 13(1-2): 30-7. PMID: 18190861. doi:

567   10.1016/j.drudis.2007.09.003.

568   9.   Sharpless NE, Depinho RA. The mighty mouse: genetically engineered mouse models in

569   cancer drug development. *Nat Rev Drug Discov* 2006; 5(9): 741-54. PMID: 16915232.

570   doi: 10.1038/nrd2110.

571   10.   Ebos JM, Kerbel RS. Antiangiogenic therapy: impact on invasion, disease progression,

572   and metastasis. *Nat Rev Clin Oncol* 2011; 8(4): 210-21. PMID: 21364524. doi:

573   10.1038/nrclinonc.2011.21.

574   11.   Francia G, Cruz-Munoz W, Man S, Xu P, Kerbel RS. Mouse models of advanced

575   spontaneous metastasis for experimental therapeutics. *Nat Rev Canc* 2011; 11(2): 135-41.

576   PMID: 21258397. doi: doi:10.1038/nrc3001.

577   12.   O'Collins VE, Macleod MR, Donnan GA, Horky LL, van der Worp BH, Howells DW.

578   1,026 experimental treatments in acute stroke. *Ann Neurol* 2006; 59(3): 467-77. PMID:

579   16453316. doi: 10.1002/ana.20741.

580   13.   Scott S, Kranz JE, Cole J, Lincecum JM, Thompson K, Kelly N, et al. Design, power,

581   and interpretation of studies in the standard murine model of ALS. *Amyotroph Lat Scler*

582   2008; 9(1): 4-15. PMID: 18273714. doi: 10.1080/17482960701856300.

583   14.   Talmadge JE, Singh RK, Fidler IJ, Raz A. Murine models to evaluate novel and

584   conventional therapeutic strategies for cancer. *Am J Pathol* 2007; 170(3): 793-804.

585   PMID: 17322365. doi: 10.2353/ajpath.2007.060929.

586   15.   Rogers CS, Stoltz DA, Meyerholz DK, Ostedgaard LS, Rokhlina T, Taft PJ, et al.

587   Disruption of the CFTR gene produces a model of cystic fibrosis in newborn pigs.

588   *Science* 2008; 321(5897): 1837-41. PMID: 18818360. doi: 10.1126/science.1163600.

589   16.   Flisikowska T, Merkl C, Landmann M, Eser S, Rezaei N, Cui X, et al. A porcine model

590   of familial adenomatous polyposis. *Gastroenterology* 2012; 143: 1173-5. PMID:

591   22864254. doi: 10.1053/j.gastro.2012.07.110.

592   17.   Pezzulo AA, Tang XX, Hoegger MJ, Abou Alaiwa MH, Ramachandran S, Moninger TO,

593   et al. Reduced airway surface pH impairs bacterial killing in the porcine cystic fibrosis

594   lung. *Nature* 2012; 487(7405): 109-13. PMID: 22763554. doi: 10.1038/nature11130.

595   18.   Maddalo D, Manchado E, Concepcion CP, Bonetti C, Vidigal JA, Han YC, et al. In vivo

596   engineering of oncogenic chromosomal rearrangements with the CRISPR/Cas9 system.

597   *Nature* 2014; 516(7531): 423-7. PMID: 25337876. doi: 10.1038/nature13902.

598   19.   Sanchez-Rivera FJ, Papagiannakopoulos T, Romero R, Tammela T, Bauer MR, Bhutkar

599   A, et al. Rapid modelling of cooperating genetic events in cancer through somatic

600   genome editing. *Nature* 2014; 516(7531): 428-31. PMID: 25337879. doi:

601   10.1038/nature13906.

602   20.   Day CP, Merlino G, Van Dyke T. Preclinical mouse cancer models: a maze of

603   opportunities and challenges. *Cell* 2015; 163(1): 39-53. PMID: 26406370. doi:

604   10.1016/j.cell.2015.08.068.

605   21.   Singh M, Murriel CL, Johnson L. Genetically engineered mouse models: closing the gap

606   between preclinical data and trial outcomes. *Cancer Res* 2012; 72(11): 2695-700. PMID:

607   22593194. doi: 10.1158/0008-5472.CAN-11-2786.

608   22.   Zitvogel L, Pitt JM, Daillere R, Smyth MJ, Kroemer G. Mouse models in

609   oncoimmunology. *Nat Rev Cancer* 2016; 16(12): 759-73. PMID: 27687979. doi:

610   10.1038/nrc.2016.91.

611   23.   Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience

612   research reporting: the ARRIVE guidelines for reporting animal research. *PLoS Biol*

613   2010; 8(6): e1000412. PMID: 20613859. doi: 10.1371/journal.pbio.1000412.

614   24.   National Institutes of Health. *Principles and Guidelines for Reporting Preclinical*

615   *Research*. Accessed 5 February 2016. https://www.nih.gov/research-training/rigor-

616   reproducibility/principles-guidelines-reporting-preclinical-research.

617   25.   National Institutes of Health. *Enhancing Reproducibility through Rigor and*

618   *Transparency*. NOT-OD-15-103. June 9, 2015. http://grants.nih.gov/grants/guide/notice-

619   files/NOT-OD-15-103.html

620   26.   Schook LB, Collares TV, Hu W, Liang Y, Rodrigues FM, Rund LA, et al. A Genetic

621   Porcine Model of Cancer. *PLOS ONE* 2015; 10(7): e0128864. PMID: 26132737. doi:

622   10.1371/journal.pone.0128864.

623   27.   Committee for the Update of the Guide for the Care and Use of Laboratory Animals.

624   *Guide for the Care and Use of Laboratory Animals*. Washington, DC: The National

625   Academies Press; 2011.

626   28.   American Veterinary Medical Association Panel on Euthanasia. *AVMA Guidelines for the*

627   *Euthanasia of Animals*. Schaumberg, IL: American Veterinary Medical Association;

628   2013.

29.   Adam SJ, Rund LA, Kuzmuk KN, Zachary JF, Schook LB, Counter CM. Genetic

induction of tumorigenesis in swine. *Oncogene* 2007; 26(7): 1038-45. PMID: 16964292.

doi: 10.1038/sj.onc.1209892.

30.   Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T)

method. *Nat Protoc* 2008; 3(6): 1101-8. PMID: 18546601. doi:

31.   Grinnell F, Zhu M, Carlson MA, Abrams JM. Release of mechanical tension triggers

apoptosis of human fibroblasts in a model of regressing granulation tissue. *Exp Cell Res*

1999; 248(2): 608-19. PMID: 10222153. doi: 10.1006/excr.1999.4440.

32.   Macpherson I, Montagnier L. Agar Suspension Culture for the Selective Assay of Cells

Transformed by Polyoma Virus. *Virology* 1964; 23: 291-4. PMID: 14187925. doi:

33.   Liang CC, Park AY, Guan JL. In vitro scratch assay: a convenient and inexpensive

method for analysis of cell migration in vitro. *Nat Protoc* 2007; 2(2): 329-33. PMID:

17406593. doi: 10.1038/nprot.2007.30.

34.   Tsutsumida H, Swanson BJ, Singh PK, Caffrey TC, Kitajima S, Goto M, et al. RNA

interference suppression of MUC1 reduces the growth rate and metastatic phenotype of

human pancreatic cancer cells. *Clin Cancer Res* 2006; 12(10): 2976-87. PMID:

16707592. doi: 10.1158/1078-0432.CCR-05-1197.

35.   Gioviale MC, Damiano G, Montalto G, Buscemi G, Romano M, Lo Monte AI. Isolation

and culture of beta-like cells from porcine Wirsung duct. *Transplant Proc* 2009; 41(4):

1363-6. PMID: 19460560. doi: 10.1016/j.transproceed.2009.02.062.

36.   Govindan R, Ding L, Griffith M, Subramanian J, Dees ND, Kanchi KL, et al. Genomic

landscape of non-small cell lung cancer in smokers and never-smokers. *Cell* 2012;

150(6): 1121-34. PMID: 22980976. doi: 10.1016/j.cell.2012.08.024.

37.   Kandoth C, McLellan MD, Vandin F, Ye K, Niu B, Lu C, et al. Mutational landscape and

653   significance across 12 major cancer types. *Nature* 2013; 502(7471): 333-9. PMID:

654   24132290. doi: 10.1038/nature12634.

655   38.   Stephens PJ, Tarpey PS, Davies H, Van Loo P, Greenman C, Wedge DC, et al. The

656   landscape of cancer genes and mutational processes in breast cancer. *Nature* 2012;

657   486(7403): 400-4. PMID: 22722201. doi:

658   39.   Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Jr., Kinzler KW.

659   Cancer genome landscapes. *Science* 2013; 339(6127): 1546-58. PMID: 23539594. doi:

660   10.1126/science.1235122.

661   40.   Waddell N, Pajic M, Patch AM, Chang DK, Kassahn KS, Bailey P, et al. Whole genomes

662   redefine the mutational landscape of pancreatic cancer. *Nature* 2015; 518: 495-501.

663   PMID: PMC4523082. doi: 10.1038/nature14169.

664   41.   Hingorani SR, Wang L, Multani AS, Combs C, Deramaudt TB, Hruban RH, et al.

665   Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely

666   metastatic pancreatic ductal adenocarcinoma in mice. *Cancer Cell* 2005; 7(5): 469-83.

667   PMID: 15894267. doi: 10.1016/j.ccr.2005.04.023.

668   42.   Groenen MA, Archibald AL, Uenishi H, Tuggle CK, Takeuchi Y, Rothschild MF, et al.

669   Analyses of pig genomes provide insight into porcine demography and evolution. *Nature*

670   2012; 491(7424): 393-8. PMID: 23151582. doi: 10.1038/nature11622.

671   43.   Rodrigues F, Hu W, Rund L, Liang Y, Counter C, Schook L. Development of the "Onco-

672   Pig": An Inducible Transgenic Porcine Model for Human Cancer [poster]. 2013

673   Brazilian Genetics Conference; URL:

http://www.dbs.illinois.edu/comparativegenomics/userfiles/file/ocnopigPoster_Brazilian

%20Genetics%20Conference2013.pdf.

44. Conference Proceedings: Swine in Biomedical Research. July 6-8, 2014. Raleigh, NC.

45. DuPage M, Dooley AL, Jacks T. Conditional mouse lung cancer models using adenoviral

or lentiviral delivery of Cre recombinase. *Nat Protoc* 2009; 4(7): 1064-72. PMID:

19561589. doi: 10.1038/nprot.2009.95.

46. Bailey JM, Swanson BJ, Hamada T, Eggers JP, Singh PK, Caffery T, et al. Sonic

hedgehog promotes desmoplasia in pancreatic cancer. *Clin Cancer Res* 2008; 14(19):

5995-6004. PMID: 18829478. doi: 10.1158/1078-0432.CCR-08-0291.

47. Handra-Luca A, Hong SM, Walter K, Wolfgang C, Hruban R, Goggins M. Tumour

epithelial vimentin expression and outcome of pancreatic ductal adenocarcinomas. *Br J*

Cancer 2011; 104(8): 1296-302. PMID: 21448168. doi: 10.1038/bjc.2011.93.

48. Kheirabadi BS, Mace JE, Terrazas IB, Fedyk CG, Estep JS, Dubick MA, et al. Safety

evaluation of new hemostatic agents, smectite granules, and kaolin-coated gauze in a

vascular injury wound model in swine. *J Trauma* 2010; 68(2): 269-78. PMID: 20154537.

doi: 10.1097/TA.0b013e3181c97ef1.

49. Cooper DK, Ekser B, Ramsoondar J, Phelps C, Ayares D. The role of genetically

engineered pigs in xenotransplantation research. *J Pathol* 2016; 238(2): 288-99. PMID:

26365762. doi: 10.1002/path.4635.

50. Niu D, Wei HJ, Lin L, George H, Wang T, Lee IH, et al. Inactivation of porcine

endogenous retrovirus in pigs using CRISPR-Cas9. *Science* 2017; 357(6357): 1303-7.

PMID: 28798043. doi: 10.1126/science.aan4187.

51.   Sullivan TP, Eaglstein WH, Davis SC, Mertz P. The pig as a model for human wound
697   healing. *Wound Repair Regen* 2001; 9(2): 66-76. PMID: 11350644. doi: 10.1046/j.1524-
698   475x.2001.00066.x.

699   52.   Swindle MM, Makin A, Herron AJ, Clubb FJ, Jr., Frazier KS. Swine as models in
700   biomedical research and toxicology testing. *Vet Pathol* 2012; 49(2): 344-56. PMID:
701   21441112. doi: 10.1177/0300985811402846.

702   53.   Xiangdong L, Yuanwu L, Hua Z, Liming R, Qiuyan L, Ning L. Animal models for the
703   atherosclerosis research: a review. *Protein Cell* 2011; 2(3): 189-201. PMID: 21468891.
704   doi: 10.1007/s13238-011-1016-3.

705   54.   Gouadon E, Moore-Morris T, Smit NW, Chatenoud L, Coronel R, Harding SE, et al.
706   Concise Review: Pluripotent Stem Cell-Derived Cardiac Cells, A Promising Cell Source
707   for Therapy of Heart Failure: Where Do We Stand? *Stem Cells* 2016; 34(1): 34-43.
708   PMID: 26352327. doi: 10.1002/stem.2205.

709   55.   Walters E, Wolf E, Whyte J, Mao J, Renner S, Nagashima H, et al. Completion of the
710   swine genome will simplify the production of swine as a large animal biomedical model.
711   *BMC Med Genom* 2012; 5(1): 55. PMID: 23151353. doi: 10.1186/1755-8794-5-55.

712   56.   Schook LB, Collares TV, Darfour-Oduro KA, De AK, Rund LA, Schachtschneider KM,
713   et al. Unraveling the swine genome: implications for human health. *Annu Rev Anim
714   Biosci* 2015; 3: 219-44. PMID: 25689318. doi: 10.1146/annurev-animal-022114-110815.

715   57.   Dawson HD, Chen C, Gaynor B, Shao J, Urban JF, Jr. The porcine translational research
716   database: a manually curated, genomics and proteomics-based research resource. *BMC
717   Genomics* 2017; 18(1): 643. PMID: 28830355. doi: 10.1186/s12864-017-4009-7.

58.   Gun G, Kues WA. Current progress of genetically engineered pig models for biomedical

research. *Biores Open Access* 2014; 3(6): 255-64. PMID: 25469311. doi:

10.1089/biores.2014.0039.

59.   Fan N, Lai L. Genetically modified pig models for human diseases. *J Genet Genomics*

2013; 40(2): 67-73. PMID: 23439405. doi: 10.1016/j.jgg.2012.07.014.

60.   Prather RS, Lorson M, Ross JW, Whyte JJ, Walters E. Genetically engineered pig models

for human diseases. *Annu Rev Anim Biosci* 2013; 1: 203-19. PMID: 25387017. doi:

10.1146/annurev-animal-031412-103715.

61.   Luo Y, Li J, Liu Y, Lin L, Du Y, Li S, et al. High efficiency of BRCA1 knockout using

rAAV-mediated gene targeting: developing a pig model for breast cancer. *Transgenic Res*

2011; 20(5): 975-88. PMID: 21181439. doi: 10.1007/s11248-010-9472-8.

62.   Luo Y, Bolund L, Sorensen CB. Pig gene knockout by rAAV-mediated homologous

recombination: comparison of BRCA1 gene knockout efficiency in Yucatan and

Gottingen fibroblasts with slightly different target sequences. *Transgenic Res* 2012;

21(3): 671-6. PMID: 22020980. doi: 10.1007/s11248-011-9563-1.

63.   Leuchs S, Saalfrank A, Merkl C, Flisikowska T, Edlinger M, Durkovic M, et al.

Inactivation and inducible oncogenic mutation of p53 in gene targeted pigs. *PLoS ONE*

2012; 7(10): e43323. PMID: 23071491. doi: 10.1371/journal.pone.0043323.

64.   Sieren JC, Meyerholz DK, Wang XJ, Davis BT, Newell JD, Jr., Hammond E, et al.

Development and translational imaging of a TP53 porcine tumorigenesis model. *J Clin*

Invest 2014; 124(9): 4052-66. PMID: 25105366. doi: 10.1172/JCI75447.

65.   Yang L, Guell M, Niu D, George H, Lesha E, Grishin D, et al. Genome-wide inactivation

740   of porcine endogenous retroviruses (PERVs). *Science* 2015; 350(6264): 1101-4. PMID:

741   26456528. doi: 10.1126/science.aad1191.

742   66.   Beraldi R, Meyerholz D, Savinov A, Kovács A, Weimer J, Dykstra J, et al. Genetic

743   Ataxia Telangiectasia porcine model phenocopies the multisystemic features of the

744   human disease. *Biochim Biophys Acta* 2017: ePub ahead of print, 2017 Jul 23. PMID:

745   28746835. doi: 10.1016/j.bbadis.2017.07.020.

746   67.   Perleberg C, Kind A, Schnieke A. Genetically engineered pigs as models for human

747   disease. *Dis Model Mech* 2018; 11(1): dmm030783. PMID: 29419487. doi:

748   10.1242/dmm.030783.

749   68.   Segatto NV, Remiao MH, Schachtschneider KM, Seixas FK, Schook LB, Collares T. The

750   Oncopig Cancer Model as a Complementary Tool for Phenotypic Drug Discovery. *Front*

751   *Pharmacol* 2017; 8: 894. PMID: 29259556. doi: 10.3389/fphar.2017.00894.

752   69.   Kuzmuk KN, Schook LB. Pigs as a Model for Biomedical Sciences. In: Rothschild MF,

753   Ruvinsky A, editors. *The Genetics of the Pig*. Oxfordshire, UK: CABI; 2011. p. 426-44.

754   70.   Emes RD, Goodstadt L, Winter EE, Ponting CP. Comparison of the genomes of human

755   and mouse lays the foundation of genome zoology. *Hum Mol Genet* 2003; 12(7): 701-9.

756   PMID: 12651866. doi: 10.1093/hmg/ddg078.

757   71.   Dawson HD, McAnulty P, Dayan A, Ganderup N, Hastings K. A comparative assessment

758   of the pig, mouse and human genomes. In: McAnulty P, Dayan A, Ganderup N, Hastings

759   K, editors. The Minipig in Biomedical Research. Boca Raton, FL: CRC Press; 2012. p.

760   323-42.

72.   Dawson HD, Smith AD, Chen C, Urban JF, Jr. An in-depth comparison of the porcine,

murine and human inflammasomes; lessons from the porcine genome and transcriptome.

Vet Microbiol 2017; 202: 2-15. PMID: 27321134. doi: 10.1016/j.vetmic.2016.05.013.

73.   Schachtschneider KM, Madsen O, Park C, Rund LA, Groenen MA, Schook LB. Adult

porcine genome-wide DNA methylation patterns support pigs as a biomedical model.

BMC Genomics 2015; 16: 743. PMID: 26438392. doi: 10.1186/s12864-015-1938-x.

74.   Vodicka P, Smetana K, Jr., Dvorankova B, Emerick T, Xu YZ, Ourednik J, et al. The

miniature pig as an animal model in biomedical research. *Ann N Y Acad Sci* 2005; 1049:

161-71. PMID: 15965115. doi: 10.1196/annals.1334.015.

75.   Swindle MM, Smith AC. *Swine in the Laboratory: Surgery, Anesthesia, Imaging, and*

Experimental Techniques. 3rd ed. Boca Raton, FL: CRC Press; 2016.

76.   Spurlock ME, Gabler NK. The development of porcine models of obesity and the

metabolic syndrome. *J Nutr* 2008; 138(2): 397-402. PMID: 18203910. doi:

77.   Röthkotter HJ. Anatomical particularities of the porcine immune system--a physician's

view. *Dev Comp Immunol* 2009; 33(3): 267-72. PMID: 18775744. doi:

10.1016/j.dci.2008.06.016.

78.   Bailey M. The mucosal immune system: recent developments and future directions in the

pig. *Dev Comp Immunol* 2009; 33(3): 375-83. PMID: 18760299. doi:

10.1016/j.dci.2008.07.003.

79.   Fairbairn L, Kapetanovic R, Sester DP, Hume DA. The mononuclear phagocyte system

of the pig as a model for understanding human innate immunity and disease. *J Leukoc*

Biol 2011; 89(6): 855-71. PMID: 21233410. doi: 10.1189/jlb.1110607.

80.   Meurens F, Summerfield A, Nauwynck H, Saif L, Gerdts V. The pig: a model for human

784   infectious diseases. *Trends Microbiol* 2012; 20(1): 50-7. PMID: 22153753. doi:

785   10.1016/j.tim.2011.11.002.

786   81.   Petersen B, Carnwath JW, Niemann H. The perspectives for porcine-to-human

787   xenografts. *Comp Immunol Microbiol Infect Dis* 2009; 32(2): 91-105. PMID: 18280567.

788   doi: 10.1016/j.cimid.2007.11.014.

789   82.   Ferrer J, Scott WE, 3rd, Weegman BP, Suszynski TM, Sutherland DE, Hering BJ, et al.

790   Pig pancreas anatomy: implications for pancreas procurement, preservation, and islet

791   isolation. *Transplantation* 2008; 86(11): 1503-10. PMID: 19077881. doi:

792   10.1097/TP.0b013e31818bfda1.

793   83.   Flisikowska T, Kind A, Schnieke A. Pigs as models of human cancers. *Theriogenology*

794   2016; 86(1): 433-7. PMID: 27156684. doi: 10.1016/j.theriogenology.2016.04.058.

795   84.   Schachtschneider KM, Schwind RM, Newson J, Kinachtchouk N, Rizko M, Mendoza-

796   Elias N, et al. The Oncopig Cancer Model: An Innovative Large Animal Translational

797   Oncology Platform. *Front Oncol* 2017; 7: 190. PMID: 28879168. doi:

798   10.3389/fonc.2017.00190.

799   85.   Watson AL, Carlson DF, Largaespada DA, Hackett PB, Fahrenkrug SC. Engineered

800   Swine Models of Cancer. *Front Genet* 2016; 7: 78. PMID: 27242889. doi:

801   10.3389/fgene.2016.00078.

802   86.   Ittmann M, Huang J, Radaelli E, Martin P, Signoretti S, Sullivan R, et al. Animal Models

803   of Human Prostate Cancer: The Consensus Report of the New York Meeting of the

804   Mouse Models of Human Cancers Consortium Prostate Pathology Committee. *Canc Res*

805   2013; 73(9): 2718-36. PMID: 23610450. doi: 10.1158/0008-5472.CAN-12-4213.

87. Stachowiak M, Flisikowska T, Bauersachs S, Perleberg C, Pausch H, Switonski M, et al.

Altered microRNA profiles during early colon adenoma progression in a porcine model

of familial adenomatous polyposis. *Oncotarget* 2017; 8(56): 96154-60. PMID: 29221194.

doi: 10.18632/oncotarget.21774.

88. Saalfrank A, Janssen KP, Ravon M, Flisikowski K, Eser S, Steiger K, et al. A porcine

model of osteosarcoma. *Oncogenesis* 2016; 5: e210. PMID: 26974205. doi:

10.1038/oncsis.2016.19.

89. Shen Y, Xu K, Yuan Z, Guo J, Zhao H, Zhang X, et al. Efficient generation of P53

biallelic knockout Diannan miniature pigs via TALENs and somatic cell nuclear transfer.

J Transl Med 2017; 15(1): 224. PMID: 29100547. doi: 10.1186/s12967-017-1327-0.

90. Luo Y, Kofod-Olsen E, Christensen R, Sorensen CB, Bolund L. Targeted genome editing

by recombinant adeno-associated virus (rAAV) vectors for generating genetically

modified pigs. *J Gen Genom* 2012; 39(6): 269-74. PMID: 22749014. doi:

10.1016/j.jgg.2012.05.004.

91. Schachtschneider K, Schwind R, Darfour-Oduro K, De A, Rund L, Singh K, et al. A

validated, transitional and translational porcine model of hepatocellular carcinoma.

Oncotarget 2017: Epub ahead of print, 2017 Jun 29. PMID: 28733541. doi:

10.18632/oncotarget.18872.

92. Callesen MM, Arnadottir SS, Lyskjaer I, Orntoft MW, Hoyer S, Dagnaes-Hansen F, et al.

A genetically inducible porcine model of intestinal cancer. *Mol Oncol* 2017; 11(11):

1616-29. PMID: 28881081. doi: 10.1002/1878-0261.12136.

93. Remmers N, Cox J, Grandgenett P, Grunkemeyer J, Rund L, Schook L, et al.

Development of a Porcine Model of Pancreatic Cancer [poster]. Annual Meeting of the

American Association of Cancer Research; Washington, D.C. April 1-5, 2017.

http://cancerres.aacrjournals.org/content/77/13_Supplement/813.short

94. Smit VT, Boot AJ, Smits AM, Fleuren GJ, Cornelisse CJ, Bos JL. KRAS codon 12

mutations occur very frequently in pancreatic adenocarcinomas. *Nucleic Acids Res* 1988;

16(16): 7773-82. PMID: 3047672. doi:

95. Jones S, Zhang X, Parsons DW, Lin JC, Leary RJ, Angenendt P, et al. Core signaling

pathways in human pancreatic cancers revealed by global genomic analyses. *Science*

2008; 321(5897): 1801-6. PMID: 18772397. doi: 10.1126/science.1164368.

96. Muller PA, Vousden KH. p53 mutations in cancer. *Nat Cell Biol* 2013; 15(1): 2-8. PMID:

23263379. doi: 10.1038/ncb2641.

97. Olivier M, Hollstein M, Hainaut P. TP53 mutations in human cancers: origins,

consequences, and clinical use. *Cold Spring Harbor Perspectives in Biology* 2010; 2(1).

PMID: 20182602. doi: 10.1101/cshperspect.a001008.

98. Blackford A, Serrano OK, Wolfgang CL, Parmigiani G, Jones S, Zhang X, et al. SMAD4

gene mutations are associated with poor prognosis in pancreatic cancer. *Clin Cancer Res*

2009; 15(14): 4674-9. PMID: 19584151. doi: 10.1158/1078-0432.CCR-09-0227.

99. Caldas C, Hahn SA, da Costa LT, Redston MS, Schutte M, Seymour AB, et al. Frequent

somatic mutations and homozygous deletions of the p16 (MTS1) gene in pancreatic

adenocarcinoma. *Nat Genet* 1994; 8(1): 27-32. PMID: 7726912. doi: 10.1038/ng0994-27.

100. Bardeesy N, Cheng KH, Berger JH, Chu GC, Pahler J, Olson P, et al. Smad4 is

dispensable for normal pancreas development yet critical in progression and tumor

biology of pancreas cancer. *Genes Dev* 2006; 20(22): 3130-46. PMID: 17114584. doi:

10.1101/gad.1478706.

101.   Bardeesy N, Aguirre AJ, Chu GC, Cheng KH, Lopez LV, Hezel AF, et al. Both

853   p16(Ink4a) and the p19(Arf)-p53 pathway constrain progression of pancreatic

854   adenocarcinoma in the mouse. *Proc Natl Acad Sci U S A* 2006; 103(15): 5947-52. PMID:

855   16585505. doi: 10.1073/pnas.0601273103.

856   102.   Wang K, Jin Q, Ruan D, Yang Y, Liu Q, Wu H, et al. Cre-dependent Cas9-expressing

857   pigs enable efficient in vivo genome editing. *Genome Res* 2017; 27(12): 2061-71. PMID:

858   29146772. doi: 10.1101/gr.222521.117.

859   103.   Ryu J, Prather RS, Lee K. Use of gene-editing technology to introduce targeted

860   modifications in pigs. *J Anim Sci Biotechnol* 2018; 9: 5. PMID: 29423214. doi:

861   10.1186/s40104-017-0228-7.

Acknowledgements

This study is the result of work supported in part with resources and the use of facilities

at the Omaha VA Medical Center (Nebraska-Western Iowa Health Care System). Portions of this

study were presented at the 108th Annual Meeting of the American Association for Cancer

Research (AACR) in Washington, D.C., April 1-5, 2017. The authors would like to acknowledge

the technical assistance of Gerri Siford, Chris Hansen, Kelly O'Connell, and Tom Caffrey. The

authors also would like to thank Dr. Lawrence B. Schook and Dr. Laurie A. Rund at the

University of Illinois for the gift of their plasmid that contained the $KRAS^{G12D}$ mutation, along

with comments, insights and suggestions for its use in this project. Datasets from this manuscript

will be freely shared upon request to the senior author (MAC).

Author Contributions

Contributor Role	Description	Author
Conceptualization	Ideas; formulation or evolution of overarching research goals and aims.	MAC
Data Curation	Management activities to annotate (produce metadata), scrub data and maintain research data (including software code, where it is necessary for interpreting the data itself) for initial use and later reuse.	NR, JLC, JAG, SA
Formal Analysis	Application of statistical, mathematical, computational, or other formal techniques to analyze or synthesize study data.	NR, MAC, SA
Funding Acquisition	Acquisition of the financial support for the project leading to this publication.	MAC
Investigation	Conducting a research and investigation process, specifically performing the experiments, or data/evidence collection.	NR, JLC, JAG, SA, MAC
Methodology	Development or design of methodology; creation of models.	NR, MAH, MAC
Project Administration	Management and coordination responsibility for the research activity planning and execution.	MAC
Resources	Provision of study materials, reagents, materials, patients, laboratory samples, animals, instrumentation, computing resources, or other analysis tools.	MAH, SA, MAC
Software	Programming, software development; designing computer programs; implementation of the computer code and supporting algorithms; testing of existing code components.	na
Supervision	Oversight and leadership responsibility for the research activity planning and execution, including mentorship external to the core team.	MAH, MAC
Validation	Verification, whether as a part of the activity or separate, of the overall replication/reproducibility of results/experiments and other research outputs.	NR, JLC, JAG, SA
Visualization	Preparation, creation and/or presentation of the published work, specifically visualization/data presentation.	NR, MAC
Draft Preparation	Creation and/or presentation of the published work, specifically writing the initial draft (including substantive translation).	NR, MAC
Review & Editing	Preparation, creation and/or presentation of the published work by those from the original research group, specifically critical review, commentary or revision – including pre- or post-publication stages.	NR, JLC, JAG, SA, MAH, MAC

 na = not applicable.

880 **Table 1. Tumorigenic cell lines and their expression products.**
881

Cell line	AcGFP1 protein	KRAS^{G12D} protein	p53^{R167H} protein	SMAD4 shRNA	p16^{Ink4A} shRNA
PGKP	+	+	+		
PGKPS	+	+	+	+	
PGKPSC*	+	+	+	+	+

882
883 All cell lines based on primary porcine pancreatic ductal epithelial cells. *Abbreviation key: **P** =
884 pancreatic ductal epithelium; **G** = GFP; **K** = KRAS; **P** = p53; **S** = SMAD4; **C** =
885 p16^{Ink4A}/CDKN2A.
886

 Figure Legends

Fig. 1. Isolation and transduction of primary porcine pancreatic ductal epithelial cells. (A)

Phase image of cells isolated from porcine pancreatic duct, showing epithelial-like morphology

(bar = 1,000 μm). (B) Immunofluorescent staining for cytokeratin 19 in the cultured primary

cells (bar = 200 μm). (C) Immunoblot for the p53 mutant in nine different cells lines transduced

with the GKP virus (PGKP cells). Pancreas = wild type pancreatic ductal epithelial cells.

Representative blot of three separate experiments. (D-E) qPCR of KRAS and p53 mutants in the

nine PKGP cell lines. Each bar represents mean of three separate experiments. *p<0.05, **

p<0.03, ***p<0.01 (unpaired t-test, compared to wild type).

Fig. 2. Effect of SMAD4 and p16^{Ink4A} knockdown on transformation in the PGKP cell line.

(A) Soft agar assay and (B) migration assay in the nine GKP-transduced cell lines. Pancreas =

wild type pancreatic ductal epithelial cells. Each bar represents mean of three separate

experiments. (C) RT-PCR of SMAD4 and p16^{Ink4A}/CDKN2A mRNA in cell lines expressing

targeted or scramble shRNA. Representative blot of three separate experiments. (D) Soft agar

assay and (E) migration assay of selected cell lines (PGKP, PGKPS, and PGKPSC), showing the

in vitro effect of additional knockdown of SMAD4 ± p16^{Ink4A} on the transformation of GKP-

transduced pancreatic ductal epithelial cells.

Fig. 3. *In vitro* transformation assays comparing the PGKP, PGKPS, and PGKPSC cell

lines. (A) Cell culture population doubling time (count-based assay). Pancreas = wild type

pancreatic ductal epithelial cells. (B) Proliferation rate (metabolic dye-based assay), represented

as fold change, normalized to wild type cells. (C) Invasion (Matrigel®-based assay). Each bar or

data point represents mean of three separate experiments.

Fig. 4. *In vivo* tumorigenesis assays comparing the PGKP, PGKPS, and PGKPSC cell lines.

(A) Sample of a resected tumor from subcutaneous injection in nude mice; note size and

vascularity. (B) Tumor growth curve from the subcutaneous injection in nude mice. Each data

point represents mean of 10 mice. (C) Metastasis after orthotopic implantation in nude mice

Fig. 5. Tumor immunohistochemistry from subcutaneous nude mouse assay. Representative

images of primary tumors derived from the PGKP, PGKPS, and PGKPSC cell lines stained with

an array of epithelial markers, including E-cadherin (Ecad; B, G, L), Epithelial Cell Adhesion

Molecule (EpCAM; C, H, M), Pan-Cytokeratin (PanCK; D, I, N), and Cytokeratin-19 (CK19; E,

J, O); and also stained with mesenchymal markers, including α-Smooth Muscle Actin (SMA; P,

S, V), Vimentin (Vim; Q, T, W), and Type I Collagen (Col I; R, U, X). Bar = 200 μm.

Supporting Information

Fig. S1. pIRES2-AcGFP1 Vector Information

Fig. S2. Studies with the NSRRC KRAS/p53 Oncopig

Fig. S3. KRAS/p53 immunohistochemistry of subcutaneous murine tumors

Protocol S1. Isolation of Epithelial Cells from Porcine Pancreas

Protocol S2. Porcine methodology

Table S1. Responses to the ARRIVE recommendations

Table S2. Responses to the NIH Preclinical Research Guidelines

Table S3. Primers and other short sequences

Table S4. Antibody information

Table S5. Oncopig descriptive data

Table S6. Oncopig serum laboratory testing

Figure 1

Figure 2

A

B

$ Compared to pancreas
# Compared to PGKP
** p<0.03
*** p<0.01
**** p<0.005

C

* p < 0.05
** p<0.03
*** p<0.001

Figure 3

Figure 4

Figure 5