Meta-analysis of hypoxic transcriptomes from public databases

Hidemasa Bono [1],* and Kiichi Hirota [2],*

[1] Database Center for Life Science (DBCLS), Joint Support-Center for Data Science Research, Research Organization of Information and Systems, 1111 Yata, Mishima, Shizuoka 411-8540, Japan; bono@dbcls.rois.ac.jp; bono@dbcls.rois.ac.jp

[2] Department of Human Stress Response Science, Institute of Biomedical Science, Kansai Medical University, Hirakata, Japan; hif1@mac.com

* Correspondence: bono@dbcls.rois.ac.jp; Tel.: +81-4-7135-5508

Abstract: Hypoxia is the insufficiency of oxygen in the cell, and hypoxia-inducible factors (HIFs) are central regulators of oxygen homeostasis. In order to obtain functional insights into the hypoxic response in a data-driven way, we attempted meta-analysis of the RNA-seq data from the hypoxic transcriptome archived in public databases. In view of methodological variability, we first manually curated RNA-seq data from appropriate pairs of transcriptomes before and after hypoxic stress. These included 128 human and 52 murine transcriptome pairs. We classified the results of experiments for each gene into three categories: upregulated, downregulated, and unchanged. Hypoxic transcriptomes were then compared between humans and mice to identify common hypoxia-responsive genes. In addition, meta-analyzed hypoxic transcriptome data were integrated with public ChIP-seq data on the known human HIFs HIF1A and EPAS1 in order to provide insights into hypoxia-responsive pathways involving direct transcription factor binding. This study provides a useful resource for hypoxia research. It also demonstrates the potential of a meta-analysis approach to public gene expression databases for selecting candidate genes from gene expression profiles generated under various experimental conditions.

Keywords: hypoxia; transcriptome; RNA-seq; ChIP-seq; public database; meta-analysis

1. Introduction

The development of high-throughput sequencing technology has enabled cost-effective reading of tens of millions of base pairs in a single run. RNA-seq takes advantage of this technology to elucidate the expression profiles of genes and transcriptomes assayed under particular conditions by producing counts of sequences corresponding to genes of interest.

Published transcriptome data have been archived to two large public repositories, the Gene Expression Omnibus (GEO) in the US National Center for Biotechnology Information (NCBI) [1] and ArrayExpress (AE) in the European Bioinformatics Institute (EBI) [2]. The total size of these databases (DBs) is now over two million samples and near a hundred thousand series. They are open access and thus ready to be utilized for data-driven research. Nevertheless, large-scale comparison among archived transcriptomes has not often been conducted because of the technical challenges presented by the magnitude, complexity, and cumbersome nature of the data. Even if very large amounts of data can be successfully downloaded, it can be difficult to interpret data from different laboratories, as experimental protocols used are not uniform.

Even though oxygen is an essential molecule for life support, higher organisms such as mammals do not have a mechanism to biosynthesize oxygen in the body. Organs and tissues are regularly exposed to risk of "oxygen deficiency," and living organisms have evolved mechanisms to respond to hypoxia. The supply and consumption of oxygen determine its intracellular partial pressure, which is kept in a relatively narrow range. Oxygen deficiency (hypoxia) and oxygen sufficiency (hyper oxygen) cause an adaptive response to maintain oxygen homeostasis at the cellular level. This includes adjustments to the energy metabolism system according to changes in oxygen partial pressure. In addition, higher organisms that are anatomically complex have

specialized mechanisms to acquire necessary and sufficient oxygen for all cells. Proper functioning
and regulation of these systems require the coordinated expression of many genes, and this is
thought to be controlled by transcription factors known as hypoxia-inducible factors (HIFs) [3].
Various groups have published comprehensive data on gene expression in response to hypoxic
stress, but have not produced uniform results, and because of this the pathways involved are not
entirely clear. In addition, there may be other data on hypoxia-related genes that have not been
reported as such, due to the main experimental target not being hypoxic stress [4]. Collective
analysis of gene expression and transcription factor binding information obtained from multiple
studies that also introduces comparative analysis between species has not previously been
attempted in this field. Further, although the analysis of many comprehensive datasets should
enable the elucidation of novel control pathways, comprehensive methods to achieve this are
lacking.
Based on the above background, this study aimed to develop a comparative data analysis
method to identify expression fluctuations and transcription factor binding regions associated with
hypoxic stress, and to use it to perform a meta-analysis of available data in the public DBs, and gene
functional analyses. We first curated hypoxic RNA-seq transcriptome data from GEO and AE to
make a list of hypoxia-normoxia transcriptome pairs. We then did systematic transcriptome
quantification analysis on the datasets we collected. By introducing a new metric that counts
instances of upregulation, downregulation, or unresponsiveness of genes, we determined the
average effect of hypoxia on genes under different experimental conditions. We also compared these
values between genes orthologous in humans and mice. In addition, we examined data from
chromatin immunoprecipitation sequencing (ChIP-seq), which comprehensively elucidates the
regions in which transcription factors bind to genomic DNA.

2. Materials and Methods

2.1. Curation of public gene expression data

In order to screen hypoxia-related gene expression data from public databases, we used a
graphical web tool called All Of gene Expression (AOE; https://aoe.dbcls.jp/) [5]. AOE not only
integrates metadata from the NCBI Gene Expression Omnibus (GEO) [1], EBI ArrayExpress (AE) [2],
and DDBJ Genomic Expression Archive (GEA) [6],   but also those of RNA-seq data archived only in
the Sequence Read Archive (SRA) [7]. Conventional search by the keywords 'hypoxia' or 'hypoxic'
in AOE was adopted to scan the databases initially. The curation of the data, which included paired
'hypoxia' and 'normoxia' experiment entries, and descriptions of cell line and experimental
conditions (oxygen concentration) used, was done manually.
In our previous pilot study, we investigated the human and mouse data produced by
Affymetrix GeneChip [8], but the subsequent accumulation of RNA-seq data enabled this study to
be focused only on RNA-seq. In order to minimize the noise from different sequencing platforms, it
was necessary to exclude data from older sequencing platforms. Consequently, nearly all data were
the product of Illumina sequencing platforms. The complete lists of these paired data are available
from   figshare   (human   https://doi.org/10.6084/m9.figshare.5811987.v2;   mouse
https://doi.org/10.6084/m9.figshare.9948158.v1).

2.2. Gene expression quantification

After hypoxia-related records were sought using AOE, corresponding sequence reads in SRA
were retrieved mainly from the DDBJ FTP site (ftp://ftp.ddbj.nig.ac.jp/). Because the retrieved data
were compressed in the SRA format, these files were converted to FASTQ formatted files for
expression   quantification   using   the   fasterq-dump   program   in   the   SRA   Toolkit
(https://www.ncbi.nlm.nih.gov/sra/docs/toolkitsoft/). Both single-end and paired-end reads were
re-used for the analysis. RNA-seq reads were then quantified using ikra (v1.2.0), an RNA-seq
pipeline centered on Salmon [9]. Ikra automates the RNA-seq data analysis process, which includes
quality control of reads (Trim Galore version 0.4.1 [10] with Cutadapt version 1.9.1 [11]) and

transcript quantification (Salmon version 0.14.0 with reference transcript sets in GENCODE release
30 for human and M21 for mouse). These tools were used with default parameters in ikra. In this
study, the data acquisition and quality control process took around six weeks for the current data
set. Processed transcript quantification from RNA-seq data were also uploaded to figshare and are
publicly   available   (human   https://doi.org/10.6084/m9.figshare.9948170.v1;   mouse
https://doi.org/10.6084/m9.figshare.9948200.v1).
Where hypoxia and normoxia transcriptome data were paired, the ratio of all gene pairs
(termed the HN ratio) was calculated.
105

HN ratio = ([gene expression value in hypoxia] + 1) / ([gene expression value in normoxia] + 1)

106
Values of HN ratios for all experiments were then classified into three groups. When the HN
ratio was over the threshold for upregulation, the gene was regarded as upregulated. Similarly,
when the HN ratio was below the threshold for downregulation, the gene was regarded as
downregulated. If the gene was labeled as neither upregulated nor downregulated, it was classified
as 'unchanged'. Finally, the numbers of counts for up, down, and unchanged were calculated for all
genes. For the up-/downregulated gene classification, several thresholds were tested to optimize the
calibration. For this study, we adopted a two-fold threshold. The number of experiments in which
genes were up/down regulated was counted for all genes in the human genome.
For the evaluation of hypoxia-inducible genes, a hypoxia-and-normoxia score (HN-score) was
calculated for all genes in humans and mice respectively. HN-score was the count of [count of
human RNA-seq UP] - [count of human RNA-seq DOWN]. The same calculation was also done in
mice. Orthologous gene mapping between humans and mice and the corresponding gene
annotations   were   obtained   using   Ensembl   Biomart   [12].   Complete   lists   of   counts
(up/down/unchanged)   after   hypoxic   stress   were   uploaded   to   figshare   (human:
https://doi.org/10.6084/m9.figshare.5812710.v3;   mouse:
https://doi.org/10.6084/m9.figshare.9948233.v2).
All codes and scripts in bash, R, and Perl for the above data processing are freely available
from GitHub at https://github.com/bonohu/chypoxia/ .

2.3. Meta-analysis of ChIP-seq data

Public ChIP-seq data were collected, curated, and pre-calculated for reuse in the ChIP-Atlas
database [13]. Average MACS2 scores for all genes were retrieved using the 'Target Genes' tool in
ChIP-Atlas for Hypoxia-inducible factor 1-alpha (HIF1A) and Endothelial PAS domain-containing
protein 1 (EPAS1, also known as hypoxia-inducible factor-2alpha (HIF-2alpha)) as 'Antigens' with
+-5k for the 'Distance from TSS' parameter. For the integration of ChIP-seq data into RNA-seq data
produced above, the names of genes were used to join two datasets.

2.4. Visualization and the integrated functional analysis of genes

For the visualization of data, TIBCO Spotfire Desktop (version 7.6.0) with TIBCO Spotfire's
"Better   World"   program   license   (TIBCO   Spotfire,   Inc.,   Palo   Alto,   CA,   USA)
(http://spotfire.tibco.com/better-world-donation-program/) was used to produce scatter plots in this
study.
Metascape was used for the gene set enrichment analysis [14]. Conventional 'express
analysis' in Metascape was used to draw histograms. In the gene set enrichment analysis of genes
upregulated and downregulated in hypoxic transcriptomes (Suppl Fig. 1), the HN-scores described
above were used for the extraction of a gene list for Metascape input. Two lists of genes were
generated by extracting genes whose HN-score was over (or below) a threshold, where roughly 1% of
all genes could be listed. These HN-score thresholds were 32 for upregulation (374 genes) and -34 for
downregulation (324 genes), respectively.
144

3. Results

3.1. Curation of hypoxic transriptome data in public databases

We initially mined hypoxia-related gene expression data from public databases using an integrated graphical web tool for gene expression data called AOE, which has been maintained as an index of public gene expression databases. Using the conventional keyword search by 'hypoxia' in AOE, we showed that the number of experiments was very few in most model organisms except humans and mice.

Pairs of samples before and after hypoxic stress were made after careful curation of dataset descriptions. We were able to obtain 128 pairs from 35 data series in humans and 53 pairs from 10 data series in mice in 2018. The complete list of pairs in RNA-seq data by Illumina sequencers is also accessible from figshare (human: https://doi.org/10.6084/m9.figshare.5811987.v2; mouse: https://doi.org/10.6084/m9.figshare.9948158.v1).

The overall procedure of the work described above is depicted in Figure 1.

Figure 1. Schematic view of hypoxic transcriptome meta-analysis. Public databases were searched and hypoxia-related RNA-seq data were manually curated. Following this, meta-analysis was done. In conjunction with meta-analyzed data from ChIP-Atlas, data were visualized and evaluated.

3.2. Meta-analysis of hypoxia-responsive genes

After the quantification of gene expression from RNA-seq data, the number of conditions under which each gene was upregulated, downregulated, and unchanged were counted. The reason for this three-way categorization is that the data are highly series-specific owing to various cell lines and experimental conditions. Complete lists of the meta-analyzed results are accessible from figshare (human: https://doi.org/10.6084/m9.figshare.5812710.v3; mouse: https://doi.org/10.6084/m9.figshare.9948233.v2).

In order to visualize differentially expressed genes, we introduced a value called HN-score.
HN-score is the number of UP counts minus the number of DOWN counts and was calculated for all
genes. Using this score, we were able to quantify the degree to which each gene was affected by
hypoxia. For example, *VEGFA* had 92 UP, 6 DOWN, and 30 unchanged counts. Its HN-score was
thus 86 (= 92-6).

Following this, genes orthologous between humans and mice were related utilizing an
orthologous gene table generated from Ensembl Biomart. This operation yields a list of genes
concurrently upregulated in humans and mice. In order to sort the table, the sum of human and
mouse HN-scores was calculated (called total HN-score). For example, the score for *VEGFA* (human)
was 86 and that for *Vegfa* (mouse) was 23, so the total HN-score for *VEGF* gene was 109. Table 1
shows the top 25 genes with high total HN-score, and a complete merged list is available from
figshare (https://doi.org/10.6084/m9.figshare.9958169.v1).

Figure 2 visualizes this table as a scatter plot of HN-score values for humans and mice. Genes
located in the upper right are those upregulated after hypoxia both in humans and mice. Genes in
that category included previously reported typical hypoxia-responsive genes, for example *PGK1*,
VEGFA, and *EGLN3*, supporting the validity of our method. On the other hand, genes
downregulated both in humans and mice included some not previously identified as
hypoxia-related.

Figure 2. Meta-analysis of hypoxic transcriptomes by RNA-seq.Comparison of human and mouse
hypoxic transcriptomes analyzed by RNA-seq. The X axis shows [count of human RNA-seq
UP]-[count of human RNA-seq DOWN] and the Y axis shows [count of mouse RNA-seq UP]-[count
of mouse RNA-seq DOWN].

We then performed set enrichment analysis using Metascape. The analysis clearly revealed that
some genes upregulated in many samples with high HN-scores are well-known hypoxia-responsive
genes, while some are not (Figure S1A). The latter can be novel candidates for hypoxia-responsive

genes to study signaling pathways in hypoxia research. Metascape also clearly depicted the
functions of downregulated genes with low HN-scores (Figure S1B). Those genes were apparently
related to DNA repair and replication [15].

3.3. Integration of meta-analyzed ChIP-seq data

In order to investigate expression regulation of genes of interest by direct transcription factor
binding, we studied ChIP-seq data. In addition to the public nucleotide sequence databases,
ChIP-seq data processed by the MACS2 program are also available from the ChIP-Atlas database,
which is a database for meta-analysis results from publicly available ChIP-seq data [13]. Thus, we
retrieved ChIP-seq data for Hypoxia-inducible factor 1-alpha (HIF1A) and Endothelial PAS
domain-containing protein 1 (EPAS1, also known as hypoxia-inducible factor-2alpha (HIF-2alpha))
from ChIP-Atlas.

We integrated meta-analysis results of human RNA-seq (hypoxic transcriptome) described
above and human ChIP-seq data for HIF1A and EPAS1. Then, we visualized the results by
conventional scatterplot (Figure 3). The source data are available from figshare
(https://doi.org/10.6084/m9.figshare.9958181.v2). In addition to reported hypoxia-responsive genes
such as *ANKRD37* [16], novel HIF1-target candidate genes were found with high HN-score and high
ChIP-seq score (Figure 3A). Genes with high HN-score but low ChIP-seq score were also found, and
these may belong to the non-HIF1 regulation pathway. *NDRG1*, which encodes a cytoplasmic
protein involved in stress responses, hormone responses, cell growth, and differentiation, is a typical
gene with such a pattern. Furthermore, we found HIF1-specific and HIF2-specific genes in the
scatter plots for HIF1A (Figure 3A) and EPAS1 (Figure 3B). Genes with high ChIP-scores in EPAS1
but not in HIF1A could be HIF2 targets, but there were no distinct genes in this category.

(A)

(B)

Figure 3. Integration of meta-analyzed ChIP-seq peak values to hypoxic transcriptomes. (A) Meta-analyzed hypoxic transcriptomes (RNA-seq) vs. average of HIF1A ChIP-seq peak values. (B) RNA-seq and EPAS1 ChIP-seq.

In order to study the differences between HIF1A and EPAS1 more thoroughly, we then investigated genes with high MACS2 score for HIF1A and EPAS1 (with positive HN-score). Surprisingly, the top 100 genes for HIF1A and EPAS1 were exactly the same, and we therefore increased the number of genes to 300. The top 300 genes for HIF1A (https://doi.org/10.6084/m9.figshare.9958235.v1) and EPAS1 (https://doi.org/10.6084/m9.figshare.9958250.v1) were generated and compared to identify the differences. The two gene lists were analyzed and visualized using the 'calculate and draw custom Venn diagrams' website (http://bioinformatics.psb.ugent.be/webtools/Venn/) (Figure 4A). Gene set enrichment analysis of the intersection of the two gene lists revealed typical features of hypoxia-inducible genes (Figure 4B), while analysis of the HIF1A-specific and EPAS1-specific portions also showed interesting features (Figure 4C, D). Genes with the functional annotations 'M00001: glycolysis (Embden-Meyerhof pathway)' and 'GO:0031167: rRNA methylation' were enriched among HIF1A-specific genes (Figure 4C). On the other hand, genes with 'GO:0040008: regulation of growth' and 'GO:0003151: outflow tract morphogenesis' were enriched among EPAS1-specific genes (Figure 4D). These observations may reflect target gene differences between these two transcription factors.

(A)

(B)

M00001: Glycolysis (Embden-Meyerhof pathway), glucose => pyruvate
GO:0031167: rRNA methylation
R-HSA-8953897: Cellular responses to external stimuli
GO:0010923: negative regulation of phosphatase activity
R-HSA-983169: Class I MHC mediated antigen processing & presentation
GO:0120031: plasma membrane bounded cell projection assembly
GO:0072655: establishment of protein localization to mitochondrion
GO:0070988: demethylation
GO:0032481: positive regulation of type I interferon production
hsa04921: Oxytocin signaling pathway

(C)

GO:0040008: regulation of growth
GO:0003151: outflow tract morphogenesis
GO:0071456: cellular response to hypoxia
GO:0045807: positive regulation of endocytosis
GO:0090287: regulation of cellular response to growth factor stimulus
GO:0008544: epidermis development
GO:0050727: regulation of inflammatory response
GO:0045785: positive regulation of cell adhesion
GO:0048385: regulation of retinoic acid receptor signaling pathway
GO:0002685: regulation of leukocyte migration
M236: PID DELTA NP63 PATHWAY
GO:0010035: response to inorganic substance
R-HSA-9006936: Signaling by TGF-beta family members
hsa05230: Central carbon metabolism in cancer
R-HSA-382551: Transport of small molecules
M167: PID AP1 PATHWAY
GO:0006873: cellular ion homeostasis
GO:0042462: eye photoreceptor cell development
GO:0048511: rhythmic process
GO:1903053: regulation of extracellular matrix organization

(D)

235 **Figure 4**. Comparison and gene set enrichment analysis of HIF1A-specific and EPAS1-specific genes.
236 (A) Venn-diagram of the top 300 genes. (B) Enrichment analysis for 176 intersecting genes. (C)
237 Enrichment analysis for 124 HIF1A-specific genes. (D) Enrichment analysis for 124 EPAS1-specific
238 genes.

4. Discussion

While over two million transcriptome samples have been archived in public databases (NCBI
Gene Expression Omnibus and EBI ArrayExpress), these data were reported by various laboratories
and are thus derived from different populations, analysis platforms, and sampling conditions. The
resultant variability can compromise meaningful comparisons, and it is indispensable to curate the
data manually to do meta-analysis for the study of specific biological stresses.

Using AOE, we made a complete survey of hypoxia-related gene expression data in the public
databases. The large quantity of relevant data available made it possible to do meta-analysis at the
level of transcriptome sequences. In our previous study, only 23 hypoxia-normoxia pairs could be
analyzed due to a lack of available human RNA-seq, and no relevant murine RNA-seq data could be
found. However, although the amount of microarray data available is much greater, most RNA-seq
data are from Illumina platforms (for example HiSeq2500, HiSeq2000, NextSeq500), simplifying
comparison. Thus, we decided to use data only from RNA-seq for our meta-analysis on the hypoxic
transcriptome in the public databases.

We manually curated data by reference to recorded metadata and made pairs of
hypoxia-normoxia data from human and mouse cell lines. It is often troublesome to handle ratio
data that contains a very large number of columns. Thus, we tentatively set the threshold for
upregulation and downregulation in hypoxia, and reduced the ratio information for all samples into
a cumulative upregulation/downregulation count. This conversion made it substantially easier to
interpret genes biologically. After optimization, we determined a threshold of two-fold for this
study.

Our meta-analysis of genes where the count of upregulation was high revealed that genes
involved in 'HIF-1 alpha transcription factor network' (GSEA Gene Set: PID_HIF1_TFPATHWAY)
and 'response to oxygen levels' (Gene Ontology: GO:0070482) were significantly enriched (Figure
S1). This is of course a reasonable result. Detailed analyses of hypoxia-inducible genes in humans
and mice (Table 1) identified a series of hypoxia-inducible genes in a data-driven manner.

In addition, our integration of ChIP-seq into RNA-seq data added substantial information on
hypoxia-inducible genes. High ChIP-seq scores, based on the MACS2 program via the ChIP-Atlas
database, indicated genes whose regulatory regions were directly bound by transcription factors.
For example, *ANKRD37* showed both high HN-score and high ChIP-seq score (Figure 4A), while
NDRG1 had a high HN-score but a low ChIP-seq score. By differentiating genes in this way, we can
generate additional hypotheses about pathways regulating differential regulation. In other words,
the integration of meta-analyzed ChIP-seq and RNA-seq data clearly filtered genes with direct and
indirect regulation, and this information can be a valuable resource for the functional analysis of
hypoxia-inducible genes.

Our future work will involve detailed analyses of co-upregulated genes across many more
species, which will be possible after the compilation of such transcriptome data. In view of the
insights enabled by the integration of ChIP-seq with RNA-seq data, another potential future
approach is the inclusion of additional types of omics data.

All data described in the manuscript are archived in figshare as a collection (Bono, H. figshare
https://doi.org/10.6084/m9.figshare.c.4690397.v1 (2019)). Source codes to replicate our study are
freely available from GitHub (https://github.com/bonohu/chypoxia).

Supplementary Materials:

Figure S1: Gene set enrichment analysis of genes up- and downregulated in hypoxic transcriptomes. (A) Gene
set enrichment analysis of genes with HN-score => 32; 374 genes (around 1% of all genes). (B) Gene set
enrichment analysis of genes with HN-score =< -34; 324 genes (around 1% of all genes)

Author Contributions: conceptualization, H.B. and K.H.; methodology, H.B.; software, H.B.; validation, H.B.;
formal analysis, H.B.; investigation, H.B.; resources, H.B.; data curation, H.B.; writing—original draft
preparation, H.B.; writing—review and editing, K.H.; visualization, H.B.; supervision, H.B.; project
administration, H.B.; funding acquisition, H.B.

Funding: This work was supported by the National Bioscience Database Center of the Japan Science and
Technology Agency (JST).

Acknowledgments: Computing was partly provided by the super computer system at the National Institute of
Genetics (NIG), Research Organization of Information and Systems (ROIS), Japan. We would like to thank
Editage (www.editage.com) for English language editing.

Conflicts of Interest: The authors declare no conflict of interest.
295

References

1.   Barrett, T.; Wilhite, S.E.; Ledoux, P.; Evangelista, C.; Kim, I.F.; Tomashevsky, M.; Marshall, K.A.;
Phillippy, K.H.; Sherman, P.M.; Holko, M.; et al. NCBI GEO: archive for functional genomics data
sets—update. *Nucleic Acids Res* **2013**, *41*, D991–D995.
2.   Athar, A.; Füllgrabe, A.; George, N.; Iqbal, H.; Huerta, L.; Ali, A.; Snow, C.; Fonseca, N.A.; Petryszak, R.;
Papatheodorou, I.; et al. ArrayExpress update - from bulk to single-cell expression data. *Nucleic Acids Res.*
2019, *47*, D711–D715.
3.   Hirota, K. An intimate crosstalk between iron homeostasis and oxygen metabolism regulated by the
hypoxia-inducible factors (HIFs). *Free Radic. Biol. Med.* **2019**, *133*, 118–129.
4.   Hill, D.R.; Huang, S.; Nagy, M.S.; Yadagiri, V.K.; Fields, C.; Mukherjee, D.; Bons, B.; Dedhia, P.H.; Chin,
A.M.; Tsai, Y.-H.; et al. Bacterial colonization stimulates a complex physiological response in the
immature human intestinal epithelium. *Elife* **2017**, *6*.
5.   Bono, H. *All of gene expression (AOE): an integrated index for public gene expression databases*; Bioinformatics,
2019;
6.   Kodama, Y.; Mashima, J.; Kosuge, T.; Ogasawara, O. DDBJ update: the Genomic Expression Archive
(GEA) for functional genomics data. *Nucleic Acids Res.* **2019**, *47*, D69–D73.
7.   Kodama, Y.; Shumway, M.; Leinonen, R.; International Nucleotide Sequence Database Collaboration The
Sequence Read Archive: explosive growth of sequencing data. *Nucleic Acids Res.* **2012**, *40*, D54-56.
8.   Bono, H. Meta-analysis of hypoxic transcriptomes from public databases. *bioRxiv* **2018**, 267310.
9.   Hiraoka, Y.; Yamada, K.; Kawasaki, Y.; Hirose, H.; Matsumoto, Y.; Ishikawa, K.; Yasumizu, Y. *ikra□:*
RNAseq pipeline centered on Salmon; 2019; https://doi.org/10.5281/zenodo.3352573
10.   Krueger, F. *Trim Galore*; 2015; https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/
11.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **2011**, *17*, 10–12.
12.   Kinsella, R.J.; Kähäri, A.; Haider, S.; Zamora, J.; Proctor, G.; Spudich, G.; Almeida-King, J.; Staines, D.;
Derwent, P.; Kerhornou, A.; et al. Ensembl BioMarts: a hub for data retrieval across taxonomic space.
Database (Oxford) **2011**, *2011*, bar030.
13.   Oki, S.; Ohta, T.; Shioi, G.; Hatanaka, H.; Ogasawara, O.; Okuda, Y.; Kawaji, H.; Nakaki, R.; Sese, J.; Meno,
C. ChIP-Atlas: a data-mining suite powered by full integration of public ChIP-seq data. *EMBO Rep.* **2018**,
19.
14.   Zhou, Y.; Zhou, B.; Pache, L.; Chang, M.; Khodabakhshi, A.H.; Tanaseichuk, O.; Benner, C.; Chanda, S.K.
Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun*
2019, *10*, 1523.

15.   Nakamura, H.; Bono, H.; Hiyama, K.; Kawamoto, T.; Kato, Y.; Nakanishi, T.; Nishiyama, M.; Hiyama, E.;
330   Hirohashi, N.; Sueoka, E.; et al. Differentiated embryo chondrocyte plays a crucial role in DNA damage
331   response via transcriptional regulation under hypoxic conditions. *PLoS ONE* **2018**, *13*, e0192136.
332   16.   Benita, Y.; Kikuchi, H.; Smith, A.D.; Zhang, M.Q.; Chung, D.C.; Xavier, R.J. An integrative genomics
333   approach identifies Hypoxia Inducible Factor-1 (HIF-1)-target genes that form the core response to
334   hypoxia. *Nucleic Acids Res.* **2009**, *37*, 4587–4602.