The dyslexia susceptibility KIAA0319 gene shows a highly specific expression pattern during zebrafish development supporting a role beyond neuronal migration.

Monika Gostic[1,2], Angela Martinelli[1,2], Carl Tucker[3], Zhengyi Yang[4], Federico Gasparoli[1], Kishan Dholakia[4,2], Keith Sillar[5], Javier Tello[1,2], and Silvia Paracchini[1,2]

[1]School of Medicine, University of St Andrews, St Andrews, KY16 9TF, UK
[2]Biomedical Sciences Research Complex, University of St Andrews, North Haugh
St Andrews, KY16 9ST, UK
[3]College of Medicine and Veterinary Medicine, The University of Edinburgh, Edinburgh EH16 4TJ, UK
[4]SUPA, School of Physics and Astronomy, University of St Andrews, St Andrews, KY16 9SS, UK
[5]School of Psychology and Neuroscience, University of St Andrews, St Andrews, KY16 9JP, UK

Abstract

Dyslexia is a common neurodevelopmental disorder that affects reading abilities and is predicted to be caused by a significant genetic component. Very few genetic susceptibility factors have been identified so far and amongst those, *KIAA0319* is a key candidate. *KIAA0319* is highly expressed during brain development but its function remains poorly understood. Initial RNA-interference studies in rats suggested a role in neuronal migration whereas subsequent work with double knock-out mouse models for both *Kiaa0319* and its paralogue *Kiaa0319-like* reported effects in the auditory system but not in neuronal migration. To further understand the role of *KIAA0319* during neurodevelopment, we carried out an expression study of the zebrafish orthologue at different embryonic stages. We report particularly high gene expression during the first few hours of development. At later stages, expression becomes localised in well-defined structures such as the otic vesicles, the eyes, the telencephalon and the notochord, supporting a role for *kiaa0319* that is not restricted to brain development. Surprisingly, *kiaa0319-like,* which generally shows higher and ubiquitous expression, was not expressed in the notochord suggesting a role specific to *kiaa0319* in this structure. These data provide a framework to interpret the effects of the dyslexia-associated genetic variants that reside in the *KIAA0319* regulatory sequences and might therefore affect a tightly regulated spatiotemporal expression pattern. This study contributes to our understanding of *KIAA0319* function during embryonic development. In particular, our data support an emerging link with the auditory processing and propose additional roles in the visual system and in the notochord.

Keywords: dyslexia, development, zebrafish, gene expression, sensory organs, notochord.

Introduction

Developmental dyslexia is a specific impairment in learning to read in the absence of any other obvious impairing factors. It affects at least 5% of school-aged children and its heritability is estimated to be above 60% (1). Studying the genetic contribution to dyslexia may help to dissect the underlying neuropsychological mechanisms, which remain hotly debated (2). While a phonological deficit is the most commonly accepted cause for dyslexia, sensory dysfunction in the visual and auditory systems have also been observed in a number of studies (1–3).

The *DYX1C1, DCDC2, ROBO1* and *KIAA0319* genes are known as the classical candidates for dyslexia and they are supported by a number of independent replications studies (4, 5). A role in cortical development, and specifically in neuronal migration, has been proposed for these genes (6), in line with earlier *post-mortem* observations that reported subtle cortical defects in individuals with dyslexia (7, 8). In particular, the *KIAA0319* gene has been found to be associated with dyslexia and reading abilities in multiple clinical and epidemiological cohorts (4, 9). A specific dyslexia-associated variant was shown to affect *KIAA0319* transcription regulation and gene expression levels, providing a mechanism to link genetic variation with the disorder (10, 11). Its paralogous gene, *KIAA0319-LIKE* or *KIAA0319L*, has also been reported to be associated with dyslexia (12). Both *KIAA0319* and *KIAA0319L* are transmembrane proteins (13), but their exact cellular function remains to be fully explained. In parallel, a new role in cilia biology is emerging for dyslexia candidate genes (14, 15). A transcriptome study showed differential regulation for *KIAA0319, DCDC2* and *DYX1C1* in ciliated tissues (16). Functional studies for Dyx1c1 and Dcdc2 showed a role in ciliogenesis in different biological models (17–19) and mutations in both genes have been found in patients with ciliopathies (17, 20). While there is no direct evidence supporting a role for KIAA0319 in cilia, one of its most distinctive features is the presence of five PKD domains (21). Mutations in PKD genes, which play key roles in cilia, lead to ciliopathies and laterality defects (22). *KIAA0319* has been shown to be a target of T-Brain-1 (TBR1), a transcription factor implicated in autism which regulates different brain developmental processes, such as neuronal migration, axon guidance (23) and the determination of left-right asymmetries in bilaterians (24). *KIAA0319* has been shown to be involved in axon growth and regeneration supporting a role also in the adult peripheral nervous system (25).

Both *KIAA0319* and *KIAA0319L* have been implicated in neuronal migration following knockdown experiments that specifically targeted neurons at the early stages of brain development using *in utero* shRNA in rats (11, 26–29). However, knockout (KO) mouse models do not display any cortical abnormalities that could be explained by defective neuronal migration (30, 31). Instead, the KO mice presented auditory system defects (31) in line with observations reported in adult rats that underwent KIAA0319 knock-down *in utero* (27, 32, 33). Therefore, while a role for *KIAA0319* in neurodevelopment is supported by different lines of evidence, its exact function remains largely unknown.

Here, we report a gene expression study for the *kiaa0319* gene during zebrafish development. We observed a specific spatiotemporal expression pattern. *kiaa0319* is highly expressed during the first hours of development. At later stages we observed high expression restricted to specific structures including the eyes, the otic vesicles and the notochord. Surprisingly, *kiaa0319-like,* although presenting a generally higher expression signal consistent with its ubiquitous expression reported in other species, was not expressed in the notochord, suggesting a specific role to *kiaa0319*. These data are suggestive of wider roles for *KIAA0319* during development rather than a specific function restricted to neuronal migration. In particular, our study supports a role outside the brain, specifically in sensory organs and in the notochord.

Materials and Methods

Fish care. Wild type zebrafish (*Danio rerio*) (WIK and AB/TU) and the double transgenic Tg(gfap:GFP);Tg(Oligo2:dsRed) were raised at The Queen's Medical Research Institute at the University of Edinburgh according to standard procedures in a Home Office approved facility. Developmental stages, maintained at 28.5°C, were identified as previously described (34). Animals were handled in accordance with the guidelines from European Directive 2010/63/EU and euthanised in accordance with Schedule 1 procedures of the Home Office Animals (Scientific Procedures) Act 1986. Zebrafish embryos were obtained using the Mass Embryo production system (MEPs) of the wild type line Wik.

PCR and qPCR. Total RNA from developmental stages between 16 – 32 cells, up to 120 hours post-fertilisation (hpf) was extracted using the RNeasy Mini kit according to the manufacturer's instructions (QIAGEN) using at least 50 embryos at each stage. The heart, liver and brain were dissected from 5 adult fish, flash-frozen on dry ice and stored at -80°C until the RNA was extracted. The PrimeScript RT reagent kit (Takara) was used to transcribe the RNA into the cDNA following the manufacturer protocol. The presence of *kiaa0319* transcripts was verified by electrophoresis following PCR amplification. Gene expression level was assessed by quantitative PCR (qPCR) conducted with the Luna Universal RT-qPCR Kit (NEB) and using a Viia7 instrument (Life Technologies, Paisley, UK). For protocol

details see Supplementary materials. Primer sequences and accession numbers are shown in Supplementary Table S1.

Whole-mount in situ hybridization (WISH). WISH was carried out following a previously described protocol (35). Briefly, a DIG-labelled riboprobe targeting *kiaa0319* was transcribed using a T3 Polymerase with a DIG RNA Labelling Mix (Roche) according to the manufacturer's instructions and, as a template, a 1066 bp PCR fragment amplified from cDNA (Supplementary Table S1). Zebrafish embryos were collected and processed at 3 somite, 14 somite, 30 hpf and 48 hpf. Embryos were imaged with a Leica MZ16F or MZFLIII bright field microscopes following treatment with an anti-DIG antibody (Roche, diluted 1:5000 to in blocking buffer) and a staining solution (NBT and BCIP, Roche).

RNAscope. RNAscope ISH (Advanced Cell Diagnostics) was modified from a previously described protocol (36). Samples were fixed in 4% PFA at room temperature for a length of time dependent on the developmental stage (Supplementary Table S2). Samples were hybridized with RNAscope target probes (*kiaa0319l*, nt 545-1425 of ENSDART00000051723.5, Channel 1; *myoD1*, nt 2-1083 of ENSDART00000027661.7, Channel 2; *kiaa0319*, nt 239-1147 of ENSDART00000160645, Channel 3) overnight at 40°C. Images were taken with a Leica TCS SP8 confocal microscope and processed in Leica Application Software X (LAS X) under 20x magnification, unless otherwise specified. Light sheet microscopy (LSM) was conducted with a bespoke microscope built in-house (Supplementary Materials).

Results

As a first step of our analysis, we verified the expression of *kiaa0319* during early zebrafish development (Supplementary Figure S1). The expression of *Kiaa0319* was observed across all the developmental stages that were analysed. In the adult, expression was much higher in the brain compared to heart and liver, where it is barely detectable. The brain-specific expression pattern is consistent with the expression profile observed for KIAA0319 in humans (Supplementary Figure S2A). In contrast, *KIAA0319L* expression is widespread across human tissues (Supplementary Figure S2B). For a more accurate assessment of expression levels, we analysed *kiaa0319* expression by qPCR and compared to *kiaa0319l*, which is expected to be ubiquitously expressed (Figure 1). *kiaa0319* signal could be detected with different intensity across the developmental stages tested and, as predicted, *kiaa0319l* consistently showed higher expression. The highest expression signal was observed for the earliest stages of development (up to 5 hpf) for both genes.

WISH was used to localise the patterns of expression of *kiaa0319* in zebrafish (Figure 2). Consistent with the qPCR data, WISH results show that *kiaa0319* is highly and widely expressed during the early stages of embryonic development (Figure 2A). As development

progresses, this widespread expression becomes restricted to specific structures. At the 14 somite stage (16 hpf), *kiaa0319* expression can be visualized in the developing brain and the body midline (Figure 2A.3 and 2A.4). At 30 hpf, expression is detected in the eye, the otic vesicle and at the midbrain-hindbrain (Figure 2B). The midline expression appears localised to the notochord rather than the spinal cord. At 48 hpf expression becomes weaker in the eyes and otic vesicles and is more pronounced in the telencephalon (Figure 2C).

To achieve higher resolution and specificity we investigated the expression patterns using the highly sensitive RNAscope Fluorescent Multiplex Assay. In particular, we aimed to resolve whether the midline body signal was restricted to the notochord rather than the spinal cord (Figure 3; see Supplementary Figure S3 for the positive and negative controls for these experiments). Consistent with our qPCR and WISH results, at 24 hpf *kiaa0319* expression is widespread but stronger in the brain and body midline. Expression in specific structures, such as the otic vesicles was visible at 48 hpf. RNAscope allowed us to confirm that *kiaa0319* expression was localised to the notochord. For comparison, we analysed the *kiaa0319-like* gene with the same procedure (Figure 3B) and observed a similar expression pattern, with a strong signal in the otic vesicles. However, *kiaa0319-like* gave a much weaker signal in the notochord. Expression of *kiaa0319* in the notochord was further confirmed with fluorescence LSM combined with RNAscope probes targeting *kiaa0319* (Figure 3C). Transverse and lateral images at different developmental stages allowed us to accurately distinguish the spinal cord from the notochord. We detected much higher *kiaa0319* signal intensity in the notochord which became weaker as development progressed. Although weaker, a signal in the spinal cord was also observed. This was strongest at 96 hpf but, rather than increasing or stabilising with development progression, became weaker at 120 hpf. Finally, analysis in the double Tg(gfap:GFP);Tg(Oligo2:dsRed) transgenic embryos further confirmed the *kiaa0319* localization to the notochord. This line is ideal for distinguishing these structures as it presents i) secondary motor neurons, interneurons, and oligodendroglia cells labelled with GFP and ii) motor neurons and oligodendrocytes labelled with DsRed (37).

Figure 1. *kiaa0319* is expressed during different stages of development Quantification of the expression of *kiaa0319* (light grey) compared to *kiaa0319l* (dark grey) by qPCR. Two reference scales were used because of the much higher expression up to 5hpf. The RQ values are normalised against the *eef1a1l2* gene. Results are derived from biological triplicates and error bars indicate standard deviation.

Figure 2. *kiaa0319* presents a specific spatiotemporal expression pattern. A) At the 3 somite stage, *kiaa0319* is expressed throughout the embryo with the strongest signal in the head (A1) and to a lesser extent, in the tailbud (A2). At the 14 the somite stage, high expression continues in the head (A3) and is visible along the developing body midline and in the tail (A3 and A4). All images are dorsal views with the head on the left side. **B)** At 30 hpf *kiaa0319* is still expressed throughout the embryo but strong expression emerges in specific structures observed from dorsal (B1) and lateral (B2) views. Details of expression are shown for the eyes (B3; dorsolateral view), the midbrain-hindbrain boundary (B4; dorsal view), the otic vesicles (B5; dorsolateral view) and the notochord (B6; lateral view). **C)** In a dorsal view at 48 hpf, *kiaa0319* expression in the eyes is diminished, while signal in the telencephalon emerges (C1). The signal in the telencephalon is particularly visible in a lateral view (C2) along with expression in the eyes and the region around the notochord. Detailed dorsolateral views of the eye (C3) and otic vesicle (C4) show a specific signal in these structures but at weaker intensity when compared to the pattern observed at 30 hpf.

Figure 3. *kiaa0319* is expressed in the notochord, *kiaa0319l* is not. The expression of *kiaa0319* **(A)** and *kiaa0319l* **(B)** was examined at three embryonic stages (12, 24 and 48 hpf) using the RNAScope Fluorescent Multiplex Assay with results shown for the head (top panels) and the body (lower panels). *kiaa0319* (red) is expressed throughout the three stages. High expression is detected in the brain at early hours of development. At 48 hpf, *kiaa0319* is still highly expressed in the brain, in the otoliths (white arrows and white triangle in reference image) and in the notochord. *Kiaa0319l* shows a very similar pattern of expression, including a strong signal in the otic vesicles, but the expression in the notochord is much weaker compared to *kiaa0319*. Black area in the brain at 48 hpf correspond to the pigmentation of the embryo. **C)** Transverse (top panels) and lateral (lower panels) 3D reconstructions from light-sheet microscopy images at three developmental stages (72, 96 and 120 hpf) on samples labelled with the *kiaa0319* probe. *Kiaa0319* expression localises to the notochord (white triangle) with weakening of the signal as development progresses and as this transient structure disappears. A well-defined signal in the spinal cord (white arrow) is also detected but it weakens at the later stages of development. **D)** Images collected at 42 hpf by confocal microscopy from the Tg(gfap:GFP);Tg(Oligo2:dsRed) embryos which present GFP (green) in the spinal cord. *myoD1* (myogenic differentiation 1, a universal target for myogenic cells (38)) was used as positive control and demonstrates the specificity of the assay. *kiaa0319* (red) clearly shows expression in the notochord. For reference, the bottom right image shows a wild type embryo treated with the *kiaa0319* RNAscope probe (red) together with DAPI staining (blue). The white arrow indicates the spinal cord where only isolated dots are visible while a strong signal is detected throughout the notochord. The scale bar indicates 50 µm in all panels. See Supplementary Figure S3 for the positive and negative controls.

Discussion

We conducted the first zebrafish characterization of the dyslexia susceptibility *kiaa0319* gene. The expression pattern described in our study supports a specific role for this gene in neurodevelopmental processes and adds novel findings towards our understanding of its role in dyslexia. We found that *kiaa0319* is highly expressed at the very first stages of embryonic development with a pattern that becomes increasingly restricted to specific structures such as the developing eyes, otic vesicles, as well as the telencephalon and the notochord. We analysed for comparison also the *kiaa0319-like* gene, which showed particularly high expression in the otic vesicles but only a very weak signal in the notochord. This observation is surprising given the generally higher and ubiquitous expression of *KIAA0319-LIKE* reported in human tissues (Supplementary Figure S2) and suggests a specific role to kiaa0319. To the best of our knowledge, this is the first study reporting expression of *kiaa0319* during the very first hours of development and clearly showing its expression at specific structures including sensory organs.

Such specific expression pattern is likely to result from a fine-tuned regulation. We showed previously that the *KIAA0319* genetic variants associated with dyslexia reside in the gene regulatory region (11) and could be pinpointed to a single polymorphism (10). The dyslexia-associated allele of this polymorphism was shown to reduce expression of the *KIAA0319* gene in human cells. It is therefore possible that regulatory variants at the same locus might also affect the spatiotemporal regulation of expression.

The function of the KIAA0319 protein has been studied in human cell lines and in rodent models, however it has not been fully elucidated yet. The first functional characterization was conducted in rats and suggested a role in neuronal migration (11) while more recent studies in mouse indicated an involvement in biological processes beyond brain development (25, 31, 39).

While we observed expression in the brain, which supports a role for *kiaa0319* in brain development, we also observed expression in other organs. The specific expression of *kiaa0319* in the otic vesicles and the notochord (Figure 2 and 3) are consistent with the recent studies conducted in mice. In particular, Guidi et al (31) generated a double KO mouse model for the *Kiaa0319* and *Kiaa0319l* genes and the most notable phenotype reported was an impairment of the auditory system. Analysis of individual KO for both genes showed mild effects for Kiaa0319l but no effects for Kiaa0319 alone. In our study, the strong expression of both *Kiaa0319* and *Kiaa0319l* in the otic vesicles (Figure 2 and Figure 3) are in agreement with the data by Guidi and colleagues (31) pointing towards an evolutionarily conserved function. Rodent models for other dyslexia candidate genes (i.e. *DCDC2* and *DYX1C1*) have also suggested an impairment in auditory processing (40, 41). Given the eye-specific expression observed here, it would be expected for vision-related phenotypes to

have also been observed in the rodent models. Accordingly, we recommend conducting a thorough visual assessment in future studies of Kiaa0319 knock-out models.

Whether dyslexia is the result of a deficit in sensory systems, as predicted by the magnocellular theory (42), remains highly debated (15). Defects in both the visual and auditory systems have been reported in individuals with dyslexia across different studies, but heterogeneity and inconsistency across studies remain significant challenges (2). In the context of this debate, the *kiaa0319* expression in the eye and otic vesicles during zebrafish development is in line with a role of sensory organs. While it would be tempting to reach conclusions, it is worth noting that it is not possible to generalise and make strong assumptions based on observations for genes analysed in isolation. Moreover, the *KIAA0319* genetic associations (as most genetic associations with complex traits) explains only a small fraction of dyslexia heritability (15).

In addition to a signal in the brain, the eyes and the otic vesicles, *kiaa0319* shows high expression in the notochord, which instead was very weak for *kiaa0319-like.* The notochord is a transient embryonic structure essential for guiding the development and patterning of the early embryo (43). It is a source of signalling to the surrounding tissues to guide structural development, particularly for the spinal cord. For example, the notochord is the source of sonic hedgehog (SHH) signalling which controls many processes including the development of motor neurons, the establishment of the dorsal-ventral axis and left/right asymmetries (44, 45). The notochord also controls in a highly specific spatiotemporal manner the trajectories of dorsal root ganglion (DRG) axons through repressive signals mediated by aggrecan, one of the chondroitin sulfate proteoglycans (CSPGs) specifically found in the cartilage (46). A similar repressive role has been described for *Kiaa0319* in mice, reporting a role in repressing axon growth in hippocampal and DRG neurons (25). The same study also showed that *Kiaa0319* was expressed in sensory and spinal cord neurons in post-natal and adult mice. Our data are consistent with these findings and support a role for *kiaa0319* both in sensory organs and in body midline structures.

While our study did not investigate the cellular function of *kiaa0319*, the tissues where we observed specific expression (i.e. eye, otic vesicles and notochord) are characterised by a prominent role of cilia (47). This is consistent with previous transcriptome studies reporting higher expression of dyslexia candidate genes, including *KIAA0319,* in ciliated tissues (16) and with the emerging role of these genes in cilia biology (15).

In summary, our characterization of the *KIAA0319* dyslexia susceptibility gene in zebrafish reveals a specific pattern of expression during development. In addition to the expected expression in the brain, we show for the first time high embryonic expression during the first hours of development and, later on, specific expression in the eyes, the otic vesicles, and in the notochord. These data contribute to our understanding of the role of

KIAA0319 during development. Our data support the emerging role for *KIAA0319* in sensory organs adding to the debate around the different theories aimed at explaining the pathophysiology of developmental dyslexia. In particular, this study will contribute to the ongoing discussion around the role of neuronal migration in dyslexia (39), first proposed in the eighties and corroborated by the initial characterization of susceptibility genes, including *KIAA0319*.

Acknowledgements

SP is a Royal Society University Research Fellow. This work was supported by Royal Society [RG160373], Carnegie Trust [50341] and Northwood Trust awards to SP. MG was supported by a 600th Anniversary University of St Andrews PhD scholarship and is the recipient of EuFishBioMed scholarship used to visit the Centre de Biologie du Développement (CBD), in Toulouse, France. KD and ZY were supported by a UK Engineering and Physical Sciences Research Council (EPSRC) grant [EP/R004854/1].The authors are grateful to Prof Caterina Becker for access to zebrafish facilities, to Prof. Bruce Appel for the *Tg(gfap:GFP)*;*Tg(Oligo2:dsRed)* transgenic line, to Drs Patrick Blader and Julie Batut for mentoring on zebrafish experiments during MG stay at the CBD and to Dr Luiz Guidi for comments to the manuscript.

References

1. Shaywitz,S.E. and Shaywitz,B.A. (2005) Dyslexia (Specific Reading Disability). *Biol. Psychiatry*, **57**, 1301–1309.

2. Goswami,U. (2014) Sensory theories of developmental dyslexia: three challenges for research. *Nat. Rev. Neurosci.*, **16**, 43–54.

3. Ramus,F. and Ahissar,M. (2012) Developmental dyslexia: The difficulties of interpreting poor performance, and the importance of normal performance. *Cogn. Neuropsychol.*, **29**, 104–122.

4. Newbury,D.F., Monaco,A.P. and Paracchini,S. (2014) Reading and language disorders: the importance of both quantity and quality. *Genes (Basel)*, **5**, 285–309.

5. Carrion-Castillo,A., Franke,B. and Fisher,S.E. (2013) Molecular Genetics of Dyslexia: An Overview. *Dyslexia*, **19**, 214–240.

6. Paracchini,S., Scerri,T. and Monaco,A.P. (2007) The genetic lexicon of dyslexia. *Annu Rev Genomics Hum Genet*, **8**, 57–79.

7. Humphreys,P., Kaufmann,W.E. and Galaburda,A.M. (1990) Developmental dyslexia in women: neuropathological findings in three patients. *Ann Neurol*, **28**, 727–738.

8. Galaburda,A.M., LoTurco,J., Ramus,F., Fitch,R.H. and Rosen,G.D. (2006) From genes to behavior in developmental dyslexia. *Nat Neurosci*, **9**, 1213–1217.

9. Paracchini,S. (2011) Dissection of genetic associations with language-related traits in population-based cohorts. *J Neurodev Disord*, **3**, 365–373.

10. Dennis,M.Y., Paracchini,S., Scerri,T.S., Prokunina-Olsson,L., Knight,J.C., Wade-Martins,R., Coggill,P., Beck,S., Green,E.D. and Monaco,A.P. (2009) A common variant associated with dyslexia reduces expression of the KIAA0319 gene. *PLoS Genet*, **5**, e1000436.

11. Paracchini,S., Thomas,A., Castro,S., Lai,C., Paramasivam,M., Wang,Y., Keating,B.J., Taylor,J.M., Hacking,D.F., Scerri,T., *et al.* (2006) The chromosome 6p22 haplotype associated with dyslexia reduces the expression of KIAA0319, a novel gene involved in neuronal migration. *Hum Mol Genet*, **15**, 1659–1666.

12. Couto,J.M., Gomez,L., Wigg,K., Cate-Carter,T., Archibald,J., Anderson,B., Tannock,R., Kerr,E.N., Lovett,M.W., Humphries,T., *et al.* (2008) The KIAA0319-Like *(KIAA0319L)* Gene on Chromosome 1p34 as a Candidate for Reading Disabilities. *J. Neurogenet.*, **22**, 295–313.

13. Velayos-Baeza,A., Toma,C., da Roza,S., Paracchini,S. and Monaco,A.P. (2007) Alternative splicing in the dyslexia-associated gene KIAA0319. *Mamm Genome*, **18**, 627–634.

14. Brandler,W.M. and Paracchini,S. (2014) The genetic relationship between handedness and neurodevelopmental disorders. *Trends Mol Med*, **20**, 83–90.

15. Paracchini,S., Diaz,R. and Stein,J. (2016) Chapter Two – Advances in Dyslexia Genetics—New Insights Into the Role of Brain Asymmetries. *Adv. Genet.*, **96**, 53–97.

16. Ivliev,A.E., t Hoen,P.A., van Roon-Mom,W.M., Peters,D.J. and Sergeeva,M.G. (2012) Exploring the transcriptome of ciliated cells using in silico dissection of human tissues. *PLoS One*, **7**, e35618.

17. Tarkar,A., Loges,N.T., Slagle,C.E., Francis,R., Dougherty,G.W., Tamayo,J. V, Shook,B., Cantino,M., Schwartz,D., Jahnke,C., *et al.* (2013) DYX1C1 is required for axonemal dynein assembly and ciliary motility. *Nat Genet*, **45**, 995–1003.

18. Chandrasekar,G., Vesterlund,L., Hultenby,K., Tapia-Paez,I. and Kere,J. (2013) The zebrafish orthologue of the dyslexia candidate gene DYX1C1 is essential for cilia growth and function. *PLoS One*, **8**, e63123.

19. Massinen,S., Hokkanen,M.E., Matsson,H., Tammimies,K., Tapia-Paez,I., Dahlstrom-Heuser,V., Kuja-Panula,J., Burghoorn,J., Jeppsson,K.E., Swoboda,P., *et al.* (2011) Increased expression of the dyslexia candidate gene DCDC2 affects length and signaling of primary cilia in neurons. *PLoS One*, **6**, e20580.

20. Schueler,M., Braun,D.A., Chandrasekar,G., Gee,H.Y., Klasson,T.D., Halbritter,J., Bieder,A., Porath,J.D., Airik,R., Zhou,W., *et al.* (2015) DCDC2 Mutations Cause a Renal-Hepatic Ciliopathy by Disrupting Wnt Signaling. *Am J Hum Genet*, **96**, 81–92.

21. Velayos-Baeza,A., Toma,C., Paracchini,S. and Monaco,A.P. (2008) The dyslexia-associated gene KIAA0319 encodes highly N- and O-glycosylated plasma membrane and secreted isoforms. *Hum Mol Genet*, **17**, 859–871.

22. Hildebrandt,F., Benzing,T. and Katsanis,N. (2011) Ciliopathies. *N Engl J Med*, **364**, 1533–1543.

23. Chuang,H.-C., Huang,T.-N. and Hsueh,Y.-P. (2015) T-Brain-1 - A Potential Master Regulator in Autism Spectrum Disorders. *Autism Res.*, **8**, 412–426.

24. Kitaguchi,T., Mizugishi,K., Hatayama,M., Aruga,J. and Mikoshiba,K. (2002) Xenopus Brachyury regulates mesodermal expression of Zic3, a gene controlling left-right asymmetry. *Dev Growth Differ*, **44**, 55–61.

25. Franquinho,F., Nogueira-Rodrigues,J., Duarte,J.M.J.M., Esteves,S.S.S.S., Carter-Su,C., Monaco,A.P., Molnár,Z., Velayos-Baeza,A., Brites,P. and Sousa,M.M.M. (2017) The dyslexia-susceptibility protein KIAA0319 inhibits axon growth through Smad2 signaling. *Cereb Cortex*, **27**, In press.

26. Peschansky,V.J., Burbridge,T.J., Volz,A.J., Fiondella,C., Wissner-Gross,Z., Galaburda,A.M., Lo Turco,J.J. and Rosen,G.D. (2010) The effect of variation in expression of the candidate dyslexia susceptibility gene homolog Kiaa0319 on neuronal migration and dendritic morphology in the rat. *Cereb. Cortex*, **20**, 884–97.

27. Szalkowski,C.E., Fiondella,C.G., Galaburda,A.M., Rosen,G.D., Loturco,J.J. and Fitch,R.H. (2012) Neocortical disruption and behavioral impairments in rats following in utero RNAi of candidate dyslexia risk gene Kiaa0319. *Int J Dev Neurosci*, **30**, 293–302.

28. Adler,W.T., Platt,M.P., Mehlhorn,A.J., Haight,J.L., Currier,T.A., Etchegaray,M.A., Galaburda,A.M. and Rosen,G.D. (2013) Position of neocortical neurons transfected at different gestational ages with shRNA targeted against candidate dyslexia susceptibility genes. *PLoS One*, **8**, e65179.

29. Platt,M.P., Adler,W.T., Mehlhorn,A.J., Johnson,G.C., Wright,K.A., Choi,R.T., Tsang,W.H., Poon,M.W., Yeung,S.Y., Waye,M.M., *et al.* (2013) Embryonic disruption of the candidate dyslexia susceptibility gene homolog Kiaa0319-like results in neuronal migration disorders. *Neuroscience*, **248**, 585–593.

30. Martinez-Garay,I., Guidi,L.G., Holloway,Z.G., Bailey,M.A.G., Lyngholm,D., Schneider,T., Donnison,T., Butt,S.J.B., Monaco,A.P., Molnár,Z., *et al.* Normal radial migration and lamination are maintained in dyslexia-susceptibility candidate gene homolog Kiaa0319 knockout mice. *Brain Struct. Funct.*, **222**.

31. Guidi,L.G., Mattley,J., Martinez-Garay,I., Monaco,A.P., Linden,J.F., Velayos-Baeza,A. and Molnár,Z. (2017) Knockout Mice for Dyslexia Susceptibility Gene Homologs KIAA0319 and KIAA0319L have Unaffected Neuronal Migration but Display Abnormal Auditory Processing. *Cereb. Cortex*, 10.1093/cercor/bhx269.

32. Centanni,T.M., Chen,F., Booker,A.M., Engineer,C.T., Sloan,A.M., Rennaker,R.L., LoTurco,J.J. and Kilgard,M.P. (2014) Speech sound processing deficits and training-induced neural plasticity in rats with dyslexia gene knockdown. *PLoS One*, **9**, e98439.

33. Centanni,T.M., Booker,A.B., Sloan,A.M., Chen,F., Maher,B.J., Carraway,R.S., Khodaparast,N., Rennaker,R., LoTurco,J.J. and Kilgard,M.P. (2014) Knockdown of the dyslexia-associated gene Kiaa0319 impairs temporal responses to speech stimuli in rat primary auditory cortex. *Cereb Cortex*, **24**, 1753–1766.

34. Kimmel,C.B., Ballard,W.W., Kimmel,S.R., Ullmann,B. and Schilling,T.F. (1995) Stages of embryonic development of the zebrafish. *Dev. Dyn.*, **203**, 253–310.

35. Thisse,C. and Thisse,B. (2008) High-resolution in situ hybridization to whole-mount zebrafish embryos. *Nat. Protoc.*, **3**, 59–69.

36. Gross-Thebing,T., Paksa,A. and Raz,E. (2014) Simultaneous high-resolution detection of multiple transcripts combined with localization of proteins in whole-mount embryos. *BMC Biol.*, **12**, 55.

37. Shin,J., Park,H.-C., Topczewska,J.M., Mawdsley,D.J. and Appel,B. (2003) Neural cell fate analysis in zebrafish using olig2 BAC transgenics Neural cell fate analysis in

zebrafish using olig2 BAC transgenics. *Methods Cell Sci.*, 10.1023/B.

38. Weinberg,E.S., Allende,M.L., Kelly,C.S., Abdelhamid,A., Murakami,T., Andermann,P., Doerre,O.G., Grunwald,D.J. and Riggleman,B. (1996) Developmental regulation of zebrafish MyoD in wild-type, no tail and spadetail embryos. *Development*, **122**, 271–80.

39. Guidi,L.G., Monaco,A.P., Velayos-Baeza,A., Paracchini,S., Bishop,D.V.M. and Molnár,Z. (2018) The neuronal migration hypothesis of dyslexia: a critical evaluation thirty years on. *submitted*.

40. Truong,D.T., Che,A., Rendall,A.R., Szalkowski,C.E., LoTurco,J.J., Galaburda,A.M. and Holly Fitch,R. (2014) Mutation of Dcdc2 in mice leads to impairments in auditory processing and memory ability. *Genes Brain Behav*, **13**, 802–811.

41. Szalkowski,C.E., Booker,A.B., Truong,D.T., Threlkeld,S.W., Rosen,G.D. and Fitch,R.H. (2013) Knockdown of the candidate dyslexia susceptibility gene homolog dyx1c1 in rodents: effects on auditory processing, visual attention, and cortical and thalamic anatomy. *Dev. Neurosci.*, **35**, 50–68.

42. Stein,J. (2001) The magnocellular theory of developmental dyslexia. *Dyslexia*, **7**, 12–36.

43. Stemple,D.L. (2005) Structure and function of the notochord: an essential organ for chordate development. *Development*, **132**, 2503–12.

44. Echelard,Y., Epstein,D.J., St-Jacques,B., Shen,L., Mohler,J., McMahon,J.A. and McMahon,A.P. (1993) Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell*, **75**, 1417–30.

45. Schilling,T.F., Concordet,J.-P. and Ingham,P.W. (1999) Regulation of Left–Right Asymmetries in the Zebrafish by Shh and BMP4. *Dev. Biol.*, **210**, 277–287.

46. Masuda,T., Fukamauchi,F., Takeda,Y., Fujisawa,H., Watanabe,K., Okado,N. and Shiga,T. (2004) Developmental regulation of notochord-derived repulsion for dorsal root ganglion axons. *Mol. Cell. Neurosci.*, **25**, 217–227.

47. Malicki,J., Avanesov,A., Li,J., Yuan,S. and Sun,Z. (2011) Analysis of Cilia Structure and Function in Zebrafish. In *Methods in cell biology.*Vol. 101, pp. 39–74.