Understanding policing as a mechanism of cheater control in cooperating bacteria

Tobias Wechsler[1], Rolf Kümmerli[1] and Akos Dobay[2,3]

[1]Department of Plant and Microbial Biology, University of Zurich, Zurich, Switzerland.

[2]Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland.

[3]Zurich Institute of Forensic Medicine, University of Zurich, Zurich, Switzerland.

Email addresses: tobias.wechsler@uzh.ch, rolf.kuemmerli@uzh.ch, akos.dobay@uzh.ch

Corresponding author: Akos Dobay, Zurich Institute of Forensic Medicine, University of Zurich, Winterthurerstrasse 190/52, CH-8057 Zurich, Switzerland

Tel. +41 (0) 44 635 76 72, Fax +41 (0) 44 635 54 90, Email: Akos.Dobay@uzh.ch

Running head: Policing & cheater control in cooperating bacteria

Keywords: individual-based simulation, public goods dilemma, evolution of cooperation, cheater control mechanism, sociomicrobiology

Abstract

Policing occurs in insect, animal and human societies, where it is used as a conditional strategy to prevent cheating and enforce cooperation. Recently, it has been suggested that policing might even be relevant in enforcing cooperation in much simpler organisms such as bacteria. Here, we used individual-based modelling to develop an evolutionary concept for policing in bacteria, and identify the conditions under which it can be adaptive. We modelled interactions between cooperators, producing a beneficial public good, cheaters exploiting the public good without contributing to it, and public good producing policers that secrete a toxin to selectively target cheaters. We found that toxin-mediated policing is favored when (i) toxins are potent and durable, (ii) cheap to produce, (iii) cell and public good diffusion is intermediate, and (iv) toxins diffuse farther than the public good. Overall, we show that toxin-mediated policing can enforce cooperation, but the parameter space where it is beneficial seems quite narrow. Moreover, we found that policing decays when the genetic linkage between public good and toxin production breaks. This is because policing is itself a public good, offering protection to toxin-resistant mutants that still produce public goods, yet no longer invest in toxins. Our work suggests that very specific environmental conditions are required for genetically fixed policing mechanisms to evolve in bacteria, and offers empirically testable predictions for their evolutionary stability.

Introduction

Cooperation, a process where individuals act to increase the fitness of others at an immediate cost to themselves, is common across the tree of life (Sachs et al. 2004; West et al. 2007b). While cooperation is widespread in humans (Fehr and Fischbacher 2003) and higher organisms such as insects and vertebrates (Clutton-Brock 2002; Ratnieks et al. 2006), we have only recently recognized that microbes also evolved adaptive cooperative behavior (West et al. 2007a; Nadell et al. 2009; Ross-Gillespie and Kümmerli 2014; Cavaliere et al. 2017). Types of microbial cooperation include the formation of biofilms, where individuals secrete polymeric compounds to build a protective extracellular matrix (Nadell et al. 2009), the formation of fruiting bodies where some individuals sacrifice themselves to enable the dispersal of others (Velicer and Vos 2009); and the secretion of shareable metabolites, including proteases to digest nutrients (Diggle et al. 2007; Sandoz et al. 2007), siderophores to scavenge extra-cellular iron (Griffin et al. 2004; Kümmerli et al. 2010), and biosurfactants enabling group motility (de Vargas Roditi et al. 2013).

Although cooperation is thought to provide benefits for the collective as a whole, it is intrinsically vulnerable to exploitation by cheating mutants, which benefit from the cooperative acts performed by others, but refrain from contributing themselves to the welfare of the group (Ghoul et al. 2014). Thus, there has been great interest in identifying mechanisms that maintain cooperation and prevent the invasion of cheaters (West et al. 2007b; Asfahl and Schuster 2017; Özkaya et al. 2017). Work on microbes have proved particularly useful in this context because cheating mutants can easily be engineered and factors important for cooperation can be experimentally manipulated in laboratory settings. Studies following this approach revealed a plethora of mechanisms promoting cooperation, including: (i) limited dispersal ensuring that

64   cooperators stay together (Kümmerli et al. 2009; van Gestel et al. 2014); (ii) molecular

mechanisms allowing the recognition of other cooperators (Mehdiabadi et al. 2006; Smukalla

66   et al. 2008; Rendueles et al. 2015); (iii) regulatory linkage between multiple traits imposing

additional costs on cheaters (Jousset et al. 2009; Dandekar et al. 2012; Ross-Gillespie et al.

68   2015; Mitri and Foster 2016; Popat et al. 2017); and (iv) mechanisms reducing the cost of

cooperation such as public good recycling (Kümmerli and Brown 2010) or the use of

70   superfluous nutrients for public goods production (Xavier et al. 2011; Sexton and Schuster

2017).

In addition, several studies suggested that bacteria have evolved policing mechanisms, which

74   enable cooperators to directly sanction cheaters and thereby enforce cooperation (Manhes and

Velicer 2011; Inglis et al. 2014; Wang et al. 2015; Majerczyk et al. 2016; Evans et al. 2018).

76   For example, Wang et al. (2015) showed that in the opportunistic pathogen *Pseudomonas

aeruginosa*, the synthesis of a publicly shareable protease is regulatorily coupled to the

78   synthesis of the toxin cyanide in such a way that cheaters, deficient for protease production,

simultaneously lose immunity against cyanide. Thus, the costly synthesis of a harmful toxin,

80   which selectively targets non-cooperative cheaters, can be understood as a policing mechanism.

While it seems intriguing that organisms as simple as bacteria can perform policing behavior,

82   multiple questions regarding the evolution of microbial policing have remained unaddressed.

For one thing, we know little about the environmental conditions promoting policing via toxin

84   production. In this context, one would expect that the spatial structure of the environment,

which affects the diffusion of cells, public goods and toxins, should play a crucial role (Driscoll

86   and Pepper 2010; Inglis et al. 2011; Dobay et al. 2014). Moreover, it is unknown how potent a

toxin must be and how much it can cost in order to efficiently fight cheaters. Finally, it remains

88   unclear whether policing is an evolutionary stable strategy or whether non-policing cooperators,

    which are immune against toxins can exploit policers as second order cheaters (Inglis et al.

90   2014), as it is the case in higher organisms (Boyd et al. 2003).

92   Here, we address these issues by using realistic individual-based models that simulate

    interactions between cooperating, policing and cheating bacteria on a toroidal two-dimensional

94   surface (Dobay et al. 2014). In our *in-silico* approach, bacteria are modeled as discs and seeded

    in low numbers onto the surface of their habitat, where they can consume resources, grow,

96   divide, disperse, and secrete compounds according to specified parameters. Important to note

    is that both bacteria and public good molecules are modelled as individual agents and are free

98   to diffuse on a continuous landscape, closely mimicking biological conditions (Figs. 1 + S1).

    For our main analysis, we considered two bacterial strains that match those used in the empirical

100 policing paper by Wang et al. (2015). Specifically, we implemented a policing strain P that

    produces a public good together with a tightly linked toxin and an immunity protein; and a

102 cheating strain C that is deficient for public good production and, due to the regulatory linkage,

    is also unable to produce the toxin and its immunity protein. We further considered a

104 cooperating control wildtype strain W, which produces the public good but features no policing

    mechanism. Finally, we also considered a second-order policing cheater R, which produces the

106 public good and the immunity protein, but not the toxin. For this strain, we assume that the

    genetic linkage between the three traits can be broken, such that R directly evolves from P.

    In a first set of simulations, we examined the performance of the cooperator, cheater and

110 policing strains in monoculture to understand how the relative costs and benefits of public good

    and toxin production affect strain fitness. Next, we competed wildtype cooperators against

112   cheaters across a range of bacterial and public good diffusion coefficients to determine the

parameter space where cooperation is favored in the absence of policing (Dobay et al. 2014).

114   Subsequently, we competed the cheater against the policing strain across the same parameter

space, to test whether policing extends the range of conditions across which cooperation is

116   favored, and to identify properties of the toxin system (diffusion, potency, durability) to

guarantee efficient policing. Finally, we simulated the situation where toxin-resistant public

118   good producers evolve from the policing strain, and ask whether policing via toxin production

itself constitutes an exploitable public good.

Methods

122   *IN-SILICO* **HABITAT AND BACTERIA**

The *in-silico* habitat consists of a two-dimensional continuous toroidal surface, with no

124   boundaries. The size of the surface is 60 x 60 μm^2 = 3,600 μm^2. Bacteria are modeled as discs

with an initial radius *r* of 0.5 μm. Bacteria can consume resources, which are unlimited in our

126   system, grow at a basic growth rate μ, and divide when reaching the threshold radius of 1.0 μm.

Bacteria can disperse on the landscape according to a specific cell diffusion coefficient D_c

128   (μm^2/s), and are not bound to a grid, but can freely move on the continuous landscape (i.e. we

used an off-lattice model with double-precision floating-point format). At the beginning of a

130   simulation, we randomly seed one founder bacteria of each strain onto the surface, free to grow

and divide according to its life cycle (see Figs. 1 + S1 for a visualization).

In addition to the basic growth rate μ, the growth of a bacterium is influenced by its social

134   behavior and its interaction with other community members. Costs, reducing growth, incur to

individuals producing public goods, toxins and immunity proteins. Additional costs incur to

susceptible cells taking up toxins. The uptake of a public good, meanwhile, has a positive effect on growth for the beneficiary. Accordingly, the growth of our four strains W (public good producing wildtype strain), P (policing strain), C (toxin-sensitive cheating strain), and R (second-order toxin-resistant cheater strain) are defined by the following set of functions:

$$G_W(t+1) = (\mu - c_{pg} + b\sum P_j)G_W(t) \tag{1}$$

$$G_P(t+1) = (\mu - c_{pg} - c_{tox} - c_{res} + b\sum P_j)G_P(t) \tag{2}$$

$$G_C(t+1) = (\mu + b\sum P_j)(1 - (\frac{\sum T_i}{\theta_T})^\kappa)G_C(t) \tag{3}$$

$$G_R(t+1) = (\mu - c_{pg} - c_{res} + b\sum P_j)G_R(t) \tag{4}$$

where c_{pg}, c_{tox}, c_{res} are the production costs per public good, toxin and the immunity protein, respectively. The term $\sum P_j$ stands for the number of public goods consumed by an individual and b for the benefit derived from this action. The term $\sum T_i$ represents the number of consumed toxins. Toxins decrease the overall growth rate and lead to death when they accumulate beyond the threshold value θ_T. The parameter θ_T can thus be understood as a measure of toxin potency. The negative effect toxins have on growth is further defined by the latency parameter κ. If $\kappa = 1.0$ then there is no latency and each toxin molecule has a linear additive negative effect on growth rate (Fig. S2). With $\kappa > 1.0$, susceptible cells can tolerate toxins at low uptake rates, while the negative effects of toxins on growth accelerate with increased toxin uptake. This accounts for the common phenomenon that bacteria possess non-specific resistant mechanisms

(e.g. efflux pumps), thereby often tolerating low toxin concentrations (Fernández and Hancock

156   2012; Nikaido and Pagès 2012).

158   The public good producing strains W and P constitutively secrete one molecule per time step,

whereas P additionally secretes toxins at the same rate. The diffusion of cells, public goods and

160   toxins (described by the diffusion coefficients D_c, d_{pg} and d_{tox}, respectively) were modeled

according to a Gaussian random walk, with a Gaussian random number generator based on the

162   Box-Muller transform that converts uniformly distributed random numbers to normally

distributed random numbers. Following diffusion, public goods and toxins were consumed

164   whenever there was co-localization of molecules and cells.

166   Both, public goods and toxins were represented by points on the landscape up to the precision

of the computer (double precision) (Figs. 1 + S1). The molecules remain in the simulation until

168   they were either consumed or decayed. The probability of decaying was determined by a

durability value δ and an exponential decay function

$$P_{decay} = 1 - e^{-\omega \Delta T / \delta} \tag{5}$$

170   where ΔT is the current lifespan of a molecule and $\omega = 0.1$ the steepness of the decay.

172   On our off-lattice landscape, individuals can physically overlap following cell growth and

diffusion. To remove the overlaps, we implemented a procedure where cells are moved slightly

174   apart from each other by a random factor scaled by a maximum pulling distance. In a final step,

our stimulation involves a life-dead-control of each individual followed by the removal of dead

176   cells that have passed the toxin threshold value θ_T. Fig. 2 depicts the order in which all the

actions in our simulation were executed per time step (representing one second). This

178   simulation cycle continues until the total number of cells reaches the carrying capacity of $K =$ 1000 cells. For each of the simulated parameter settings, we performed 50 independent

180   replicates. At each time step, we recorded the relative strain frequency. At the end of a simulation, we measured the final strain frequencies, the mean time between two cell divisions,

182   and the *per capita* public good and toxin uptake for each strain.

184   **EXPLORED PARAMETER SPACE**

Table 1 provides an overview of all the parameters in our simulations and the value ranges

explored. One key focus of our study is to understand how cell dispersal and molecule diffusion affect cooperation and policing. For cell dispersal D_c, we considered active but random bacterial

motility, and varied this parameter from 0 $\mu m^2/s$ (no dispersal) to 3.5 $\mu m^2/s$ (high dispersal) in steps of 0.5. For molecule diffusion d_{pg} and d_{tox}, we implemented passive random diffusion

processes and varied parameters from 1.0 $\mu m^2/s$ (low diffusion) to 7.0 $\mu m^2/s$ (high diffusion) in steps of one. Another important aim of our study is to explore how toxin properties affect

policing. In addition to toxin diffusion, we varied: (a) toxin durability δ_{tox} from 50 (rapid decay) to 500 (intermediate decay) to 5000 (low decay) time steps; (b) toxin threshold θ_T from

600 (few molecules required to kill) to 1800 (many molecules required to kill) in steps of 400; and (c) toxin latency κ from 1.0 (low tolerance and immediate linear adverse effect) to 6.0 (high

tolerance and delayed exponential adverse effect) in steps of 0.5.

Finally, the outcome of our simulations necessarily depends on the cost and benefit parameters. We implemented a basic growth μ to ensure that all cells can grow even if they do not produce

public goods. This ensures that cells can only die because of toxins and not because they experience negative growth. $\mu = 1.0$ results in cell division occurring every 1,200 time steps, in

the absence of molecule secretion. We fixed the cost and benefit of public good production to

c_{pg} = 0.001 and b = 0.01. These values are based on our previous experience with the system

204   (Dobay et al. 2014) and ensure that public production generates a substantial net fitness benefit,

thus significantly reducing cell division time (see also Fig. 3). Of key interest is how costly

206   policing ($c_{tox+res}$) can be relative to c_{pg} in order to maintain a net benefit of cooperation. To

address this question, we gradually varied the cost ratio of these two traits $[c_{pg}/(c_{tox} + c_{res})]$

208   from 0.25 to 4.0. For all simulations, we assumed $c_{tox} = c_{res}$.

210   **STRAIN PERFORMANCE IN MONOCULTURE**

In a first set of simulations, we assessed the performance of the strains W (wildtype cooperator)

212   P (policing cooperator) and C (public-good cheater) in monoculture. These assays allow us to

quantify (i) the net benefit of cooperation, and (ii) the acceptable boundary costs of policing to

214   ensure a net benefit of cooperation. A net benefit of cooperation is given when the policing

strain P grows better than strain C, which neither produces public goods nor toxins. Because

216   we know that cell dispersal and secreted molecule diffusion influence the efficiency of

cooperation (Dobay et al. 2014), we performed simulations under three different diffusivity

218   regimes, including high (D_c = 4.0 μm²/s, $d_{pg} = d_{tox}$= 7.0 μm²/s), medium (D_c = 2.0 μm²/s, d_{pg}

= d_{tox} = 3.5 μm²/s) and low (D_c = 0.1 μm²/s, $d_{pg} = d_{tox}$ = 1.0 μm²/s) diffusivity. We then

220   compared the time needed by the three strains to reach carrying capacity.

222   **PAIRWISE STRAIN COMPETITION**

We first simulated competitions between the strains W and C across the indicated range of

224   dispersal and diffusion parameters to understand the conditions under which cooperation can

be maintained in the absence of policing. Next, we competed P against C. In this scenario, C

226   can still exploit the public good produced by *P*, but at the same time is harmed by the toxins

secreted by this opponent. In a first set of simulations, we implemented intermediate default

228   toxin properties (d_{tox} = 4.0 μm^2/s, δ_{tox} = 500 s, θ_T = 1000, and κ = 3.5) to examine whether

policing extends the range of conditions across which cooperation is favored. In a second set of

230   simulations, we then individually manipulated each of the four different toxin properties to

identify the features required for toxin-mediated policing to be efficient.

THREE-WAY STRAIN INTERACTIONS

234   To examine what happens when the genetic linkage between cooperation and policing breaks,

we performed three-way competitions between strains *P*, *C*, and *R*. The latter strain no longer

236   produces the toxin, but is resistant to it. Because we do not necessarily expect an overall winner

in these competitions, as strain frequency could potentially follow cyclical patterns (Kerr et al.

238   2002), we extended our simulation code by implementing a random cell removal event, which

is activated as soon as more than 30 % of the surface is covered with cells. This allowed us to

240   keep population size at roughly *K*/2, and to follow strain frequency over extended periods of

time (i.e. 80,000 time steps).

Results

244   **STRAIN PERFORMANCE IN MONOCULTURE**

The growth of strain *C* (neither producing public goods nor toxins) was not affected by the

246   diffusivity of the environment, and was solely determined by the basic growth rate μ, reaching

carrying capacity after 12,000 time steps (Fig. 3). Strain *W* (producing public goods) grew

248   significantly better than *C*, demonstrating the benefit of public goods cooperation. The growth

of this strain was reduced in more diffusive environments, where the likelihood of public good

250   sharing and consumption declines. The performance of strain *P* (producing toxins in addition

to public goods) greatly varied both in response to the relative costs of policing and the

252   diffusivity of the environment. In environments characterized by low diffusion, *P* always

outperformed *C* even when the cost of toxin production was four times higher than the cost of

254   public good production (triangles in Fig. 3). This was no longer the case in environments

characterized by intermediate (diamonds in Fig. 3) or high diffusivity (dots in Fig. 3), where *P*

256   only grew better than *C* when toxins were cheaper than public goods. If this condition was not

met, then the high costs of policing combined with reduced public good consumption decreased

258   cell growth to a point that prevented populations from reaching carrying capacity.

260   Interestingly, the relationship between the cost ratio $c_{pg}/(c_{tox} + c_{res})$ and the time needed to

reach carrying capacity (τ_K) was best captured by the Monod equation (Monod 1949), a

262   hyperbolic equation initially used to explain the exponential growth rate as a function of nutrient

concentration. Applied to our system, we found that the function

$$R(\tau_K) = R_K \frac{\tau_K}{R_\tau + \tau_K},$$ (6)

264   provides a fair approximation to relate r to τ_K, where R_K represents the ratio limit for reaching

carrying capacity and R_τ the time at which the carrying capacity is half the maximum (Fig. S3).

LOW CELL DISPERSAL FAVORS COOPERATION WITHOUT POLICING

268   When competing *W* against *C* without a policing mechanism in place, cooperation was only

strongly favored when cells did not disperse (Fig. 4A). In other words, if cooperator cells grew

270   as microcolonies, physically separated from the cheaters, then efficient public good sharing

among cooperators is promoted. We further found that cooperators and cheaters could coexist

272   when cell and public good diffusion was low, but greater than zero (Fig. 4A). Under all other

conditions, cheaters strongly dominated the community and pushed cooperators to very low

274   frequencies. These results are in line with our previous simulations, showing that there is only

a narrow parameter space within which public good cooperation is favored (Dobay et al. 2014).

TOXIN-MEDIATED POLICING HAS POSITIVE AND NEGATIVE EFECTS FOR

278   **COOPERATION**

The introduction of a policing mechanism, which operates via the secretion of a toxin that

280   selectively targets cheaters, had multiple dramatic effects on the competitive outcome between

the cheater C and the cooperating policer P (Fig. 4B). Policing strongly increased selection for

282   cooperation under conditions where cooperators could previously only coexist with cheaters

(compare Fig. 4A and 4B for combinations of low public good diffusion and cell dispersal).

284   This finding demonstrates that policing can indeed extend the parameter space across which

cooperation is favored. Conversely, we found that policing also had negative effects and

286   drastically accelerated selection against cooperation, especially under conditions where

cheaters previously experienced only a moderate selective advantage (compare Fig. 4A and 4B

288   for combinations of intermediate public good diffusion and cell dispersal). These two opposing

effects led to a sharp transition between conditions that either completely favor P or C, leaving

290   very few conditions where coexistence between the two strains is possible.

292   **HOW TOXIN PROPERTIES AFFECT POLICING EFFICIENCY**

We found that higher toxin diffusion d_{tox} significantly increased selection for cooperation (Fig.

294   5A, 5E and 5B), especially under conditions of intermediate public good diffusion and cell

dispersal. These results show that policing is particularly efficient when toxins are sent away to

296   target more distant competitors whilst keeping the public good more local for preferential

sharing among producers. Our simulations further revealed that high toxin durability δ_{tox}

298   increases policing efficiency and thus selection for cooperation (Fig. 5C, 5E, 5D). These

findings can be explained by the fact that more durable toxins are more likely to reach target

300   cells. Next, we examined the role of toxin potency θ_T (i.e. the number of toxins needed to kill

a target cell), and found that high potency is crucial for policing to promote cooperation (Fig.

302   5F, 5E, 5G). While this finding seems trivial at first sight, the dramatic effects we observed

when decreasing toxin potency are remarkable. For instance, a reduction of toxin potency by

304   45 % (from $\theta_T = 1000$ to $= 1800$) already completely negated any benefit of policing, and even

increased the parameter space across which cooperation was selected against. Finally, we

306   explored the role of toxin latency κ for policing. Toxin latency is a measure of target cell

tolerance: with low values of κ, the fitness of target cells is immediately affected in a linear

308   way, whereas large values of κ mean that target cells can tolerate a certain level of toxins and

negative effects only kick in later, but accelerate with higher toxin uptake rates (Figs. S2). We

310   found that low κ values dramatically increased the efiiciency of policing and selection for

cooperation (Fig. 5H, 5E, 5I).

POLICING GOES EXTINCT WHEN THE GENETIC LINKAGE BETWEEN TRAITS

314   **BREAKS**

Next, we simulated the case where the genetic linkage between cooperation and policing breaks.

316   Specifically, we studied the performance of strain *R*, which evolved directly from *P* by losing

toxin production but keeping resistance, in competition with *P* and *C* across a range of

318   intermediate environmental diffusivities. We found that the presence of *R* consistently drove *P*

to extinction under all environmental conditions tested (Fig. 6). Our simulations, which kept

320 populations at half the carrying capacity ($K/2 \sim 500$ cells) for an extended period of time

(roughly 10 times longer than in the previous pairwise competition assays), allowed us to

322 distinguish three distinct competition phases. The first phase comprises the time frame in which

the community grows from three cells to $K/2$. During this phase, we observed cyclical

324 fluctuations of strain frequencies with a general tendency for C to increase, P to decrease, and

R to remain stable. The cyclical patterns observed here are reminiscent of the rock-paper-

326 scissors dynamics described by previous studies on bacterial social interactions (Kerr et al.

2002; Kelsic et al. 2015; Inglis et al. 2016), where strains chasing each other in non-transitive

328 interactions in circles with no overall winner. The second phase was characterized by a

pronounced dip in C frequency accompanied by a strong increase in R frequency, and a

330 moderate increase in P frequency in eight of the nine diffusion conditions (Fig. 6). This pattern

is most likely explained by the accumulation of toxins in the environment, which efficiently

332 suppressed C, but at the same time gave R leverage, as it could benefit from the effect of

policing without paying the cost for it. During the third phase, we observed the extinction of P,

334 the concomitant recovery of C followed by a decrease of R (Fig. 6). These patterns arise because

as P decreases, toxin concentration declines allowing the recovery of C, which then efficiently

336 exploit the public goods produced by R. In all cases, our simulations returned to a simple

cooperator-cheater scenario, with the relative success of the two strategies being determined by

338 the diffusivity of the environment (similar to the results in Fig. 4A).

340 **Discussion**

Understanding how policing can repress competition and foster cooperation in social groups of

342 higher organisms has attracted the attention of evolutionary biologists for decades (Clutton-

Brock and Parker 1995; Frank 1995, 2003; West et al. 2007b; Ratnieks and Wenseleers 2008;

344 Kümmerli 2011; Raihani et al. 2012; Cant et al. 2013). Here, we explored whether policing

could also be an effective way to enforce cooperation in groups of microbes, sharing beneficial

346   public goods. In our models, policing is exerted via the secretion of a toxin that specifically

targets cheater cells, which do not contribute to the pool of public goods. Our simulations reveal

348   that policing is most conducive under conditions of intermediate cell dispersal and public good

diffusion, where it can extend the parameter space, under which cooperation is favored. We

350   further found that an effective policing mechanism entails a toxin that is: (i) cheaper to produce

than the public good; (ii) more diffusible than the public good in order to effectively reach

352   cheaters; (iii) durable; and (iv) potent in killing. However, our simulations also reveal two major

downsides of policing. First, policing can accelerate the loss of cooperation under conditions

354   where cheaters and cooperators can coexist in the absence of policing. This leads to a sharp

state transition between conditions where policing either favors or disfavors cooperation.

356   Second, policing is selected against if the genetic linkage between public good and the toxin-

anti-toxin production can be broken. If this occurs then toxin-resistant mutants that produce

358   public goods, but no longer contribute to toxin production, derail policing, showing that

microbial policing itself constitutes an exploitable public good (Inglis et al. 2014).

Several studies have recently proposed toxin-based policing mechanisms to keep cheaters in

362   check (Inglis et al. 2014; Wang et al. 2015; Majerczyk et al. 2016; Evans et al. 2018). While

the idea of bacteria being able to punish cheating community members is exciting, our

364   simulations reveal that the ecological spectrum under which policing can be favored is actually

quite narrow. For one thing, we found that policing is not required when environmental

366   diffusivity is low, conditions that promote cooperation *per se*. The reason for this is that low

public good diffusion and low bacterial dispersal lead to significant spatial segregation of

368   strains within bacterial community, which promotes the local sharing of public goods among

cooperators (Kümmerli et al. 2009; Julou et al. 2013; Mitri and Foster 2013; van Gestel et al.

370 2014; Weigert and Kümmerli 2017). Moreover, we show that policing is not favorable when

environmental diffusivity is high, conditions that break any spatial association between

372 cooperators and their public goods. Toxin-mediated policing is detrimental here because: (i)

cheaters can freely exploit public goods; (ii) many toxins get lost due to the high diffusion, and

374 thus never reach their targets; (iii) and the high level of cell mixing reduces the efficiency and

selectivity of policing, as cooperators are hit by toxins as often as cheaters (Inglis et al. 2011).

376 Overall, it turns out that intermediate levels of environmental diffusivity proved most beneficial

for policing. The issue with this relatively narrow parameter space is that environmental

378 diffusivity is likely to vary both temporally and spatially under natural conditions, which could

quickly shift the selective balance for or against policing. It thus remains to be seen whether

380 policing can indeed evolve under fluctuating environmental conditions.

382 Another important point that has received little attention so far concerns the question whether

the reported policing mechanisms have indeed evolved for this very purpose or whether they

384 represent by-products of regulatory linkage of traits for other reasons than policing (West et al.

2007c). For instance, in the case of *P. aeruginosa* it is well conceivable that cyanide primarily

386 serves as a broad-spectrum toxin to target inter-specific competitors under high cell density

(Bernier et al. 2016). This might be the reason why the expression of cyanide is controlled by

388 quorum sensing, and why it is regulatorily linked to other public good traits, such as protease

production, whose expression is also controlled in a density-dependent manner (Darch et al.

390 2012). Consequently, the observed cyanide-mediated policing exerted by wildtype strains on

protease-deficient strains could be a mere by-product of this regulatory linkage. Alternatively,

392 it is also possible that an initial co-incidental regulatory linkage between cyanide and protease

later proved useful as a policing mechanism and evolved as such through cooption (Foster

394   2011). Clearly, further research is needed to uncover the evolutionary history of these putative

policing mechanisms, and care must be taken to distinguish between mechanistic (proximate)

396   and evolutionary (ultimate) explanation of observed behavioral patterns (West et al. 2007c).

398   The potential policing mechanisms reported for microbes and the one implemented in our

simulations differ in one important aspect from the policing systems found in higher organisms.

400   Specifically, the difference is that the microbial policing mechanisms are genetically fixed (i.e.

strains are either cooperating policers or cheaters), whereas in higher organisms policing is

402   perceived as a conditional strategy, which can be applied to sanction cheaters only if required

(Clutton-Brock and Parker 1995; Frank 1995, 2003; Kiers et al. 2003; West et al. 2007b;

404   Ratnieks and Wenseleers 2008; Kümmerli 2011; Raihani et al. 2012). In the latter scenario,

individuals can take decisions on whether to cheat or to cooperate, and whether to impose

406   sanctions or not. In certain cases, it was found that the mere threat of policing was sufficient to

coerce individuals to cooperate and prevent cheating in the first place (Wenseleers and Ratnieks

408   2006; Kümmerli 2011; Cant et al. 2013). Microbes clearly lack cognitive abilities for such

conditional behaviors, and it is thus not surprising that cheating, cooperating and policing

410   strategies are genetically fixed in these organisms. Nonetheless, we argue that it seems fair to

use the term 'policing' for the reported behaviors, but also to keep in mind the difference

412   between conditional and fixed strategies.

414   In summary, our work contributes to the development of an evolutionary concept for policing

in microbial systems. It shows how ecological factors, in particular the diffusivity of the

416   environment, interact with the properties of a toxin-mediated policing system, to define the

parameter space in which policing can be favored. It further demonstrates how realistic

418 individual-based modelling, tracking both cells and their public goods over time and across

space, can be used to deepen our understanding of social interactions in microbes.

AUTHOR CONTRIBUTIONS

422 T. W., R. K. and A. D. conceived the study. T. W. and A. D. constructed the model. T. W.

performed the simulations. T. W., R. K. and A. D. analyzed the data and wrote the paper.

ACKNOWLEDGMENTS

426 We thank Barbara König and Marta Manser for comments, the Services and Support for Science

IT (S3IT) at the University of Zurich for technical support with the simulations on the

428 ScienceCloud, and the Swiss National Science Foundation (grant no. PP00P3-139164) and the

European Research Council (ERC-CoG grant no. 681295) for funding. The authors declare no

430 conflict of interest with this manuscript.

LITERATURE CITED

434 Asfahl, K. L. and M. Schuster. 2017. Social interactions in bacterial cell-cell signaling. FEMS
Microbiolol. Ecol. 41:92-107.

436 Bernier, S. P., M. L. Workentine, X. Li, N. A. Magarvey, G. A. O'Toole, and M. G. Surette.
2016. Cyanide toxicity to *Burkholderia cenocepacia* is modulated by polymicrobial

438 communities and environmental factors. Front. Microbiol. 7:1-14.

Boyd, R., H. Gintis, S. Bowles, and P. J. Richerson. 2003. The evolution of altruistic

440 punishment. Proc. Natl. Acad. Sci. U.S.A. 100:3531-3535.

Cant, M. A., H. J. Nichols, R. A. Johnstone, and S. J. Hodge. 2013. Policing of reproduction by

442 hidden threats in a cooperative mammal. Proc. Natl. Acad. Sci. U.S.A. 111:326-330.

Cavaliere, M., S. Feng, O. S. Soyer, and J. I. Jimenez. 2017. Cooperation in microbial communities and their biotechnological applications. Environ. Microbiol. 19:2949-2963.

Clutton-Brock, T. H. 2002. Breeding together: kin selection and mutualism in cooperative vertebrates. Science 296:69-72.

Clutton-Brock, T. H. and G. A. Parker. 1995. Punishment in animal societies. Nature 373:209-216.

Dandekar, A. A., S. Chugani, and E. P. Greenberg. 2012. Bacterial quorum sensing and metabolic incentives to cooperate. Science 338:264-266.

Darch, S. E., S. A. West, K. Winzer, and S. P. Diggle. 2012. Density-dependent fitness benefits in quorum-sensing bacterial populations. Proc. Natl. Acad. Sci. U.S.A. 109:8259-8263.

de Vargas Roditi, L., K. E. Boyle, and J. B. Xavier. 2013. Multilevel selection analysis of a microbial social trait. Mol. Syst. Biol. 9:684.

Diggle, S. P., A. S. Griffin, G. S. Campell, and S. A. West. 2007. Cooperation and conflict in quorum-sensing bacterial populations. Nature 450:411-414.

Dobay, A., H. C. Bagheri, A. Messina, R. Kümmerli, and D. J. Rankin. 2014. Interaction effects of cell diffusion, cell density and public goods properties on the evolution of cooperation in digital microbes. J. Evol. Biol. 27:1869-1877.

Driscoll, W. W. and J. W. Pepper. 2010. Theory for the evolution of diffusible external goods. Evolution 64:2682-2687.

Evans, K. C., S. Benomar, L. A. Camuy-Vélez, E. B. Nasseri, X. Wang, B. Neuenswander, and J. R. Chandler. 2018. Quorum-sensing control of antibiotic resistance stabilizes cooperation in *Chromobacterium violaceum*. ISME J. in press.

Fehr, E. and U. Fischbacher. 2003. The nature of human altruism. Nature 425:785- 791.

Fernández, L. and R. E. W. Hancock. 2012. Adaptive and mutational resistance: role of porins and efflux pumps in drug resistance. Clin. Microbiol. Rev. 25:661-681.

Foster, K. 2011. The sociobiology of molecular systems. Nat. Rev. Genet. 12:193-203.

Frank, S. A. 1995. Mutual policing and repression of competition in the evolution of cooperative groups. Nature 377:520-522.

Frank, S. A. 2003. Perspective: repression of competition and the evolution of cooperation. Evolution 57:693-705.

474   Ghoul, M., A. S. Griffin, and S. A. West. 2014. Toward and evolutionary definition of cheating. Evolution 68:318-331.

476   Griffin, A., S. A. West, and A. Buckling. 2004. Cooperation and competition in pathogenic bacteria. Nature 430:1024-1027.

478   Inglis, R. F., J. M. Biernaskie, A. Gardner, and R. Kümmerli. 2016. Presence of a loner strain maintains cooperation and diversity in well-mixed bacterial communities. Proc. R. Soc.
480   B 283:20152682.

Inglis, R. F., P. G. Roberts, A. Gardner, and A. Buckling. 2011. Spite and the scale of
482   competition in *Pseudomonas aeruginosa*. Am. Nat. 178:276-285.

Inglis, R. F., S. West, and A. Buckling. 2014. An experimental study of strong reciprocity in
484   bacteria. Biol. Lett. 10:20131069.

Jousset, A., L. Rochat, M. Péchy-Tarr, C. Keel, S. Scheu, and M. Bonkowski. 2009. Predators
486   promote defence of rhizosphere bacterial populations by selective feeding on non-toxic cheaters. ISME J. 3:666-674.

488   Julou, T., T. Mora, L. Guillon, V. Croquette, I. J. Schalk, D. Bensimon, and N. Desprat. 2013. Cell-cell contacts confine public goods diffusion inside *Pseudomonas aeruginosa*
490   clonal microcolonies. Proc. Natl. Acad. Sci. U.S.A. 110:12577-12582.

Kelsic, E. D., J. Zhao, K. Vetsigian, and R. Kishony. 2015. Counteraction of antibiotic
492   production and degradation stabilizes microbial communities. Nature 521:516-519.

Kerr, B., M. A. Riley, M. W. Feldman, and B. J. M. Bohannan. 2002. Local dispersal promotes
494   biodiversity in a real-life game of rock-paper-scissors. Nature 418:171-174.

Kiers, E. T., R. A. Rousseau, S. A. West, and R. F. Denison. 2003. Host sanctions and the
496   legume-rhizobium mutualism. Nature 425:78-81.

Kümmerli, R. 2011. A test of evolutionary policing theory with data from human societies.
498   PLoS One 6:e24350.

Kümmerli, R. and S. P. Brown. 2010. Molecular and regulatory properties of a public good
500   shape the evolution of cooperation. Proc. Natl. Acad. Sci. U.S.A. 107:18921-18926.

Kümmerli, R., A. S. Griffin, S. A. West, A. Buckling, and F. Harrison. 2009. Viscous medium
502   promotes cooperation in the pathogenic bacterium *Pseudomonas aeruginosa*. Proc. R. Soc. B 276:3531-3538.

504   Kümmerli, R., P. van den Berg, A. S. Griffin, S. A. West, and A. Gardner. 2010. Repression of competition promotes cooperation: experimental evidence from bacteria. J. Evol. Biol.
506   23:699-706.

Majerczyk, C., E. Schneider, and E. P. Greenberg. 2016. Quorum sensing control of Type VI
508   secretion factors restricts the proliferation of quorum-sensing mutants. eLife 5:e14712.

Manhes, P. and G. J. Velicer. 2011. Experimental evolution of selfish policing in social bacteria.
510   Proc. Natl. Acad. Sci. U.S.A. 108:5357-5362.

Mehdiabadi, N. J., C. N. Jack, T. T. Farnham, T. G. Platt, S. E. Kalla, G. Shaulsky, D. C.
512   Queller, and J. E. Strassmann. 2006. Kin preferences in a social microbe. Nature 442:881-882.

514   Mitri, S. and K. R. Foster. 2013. The genotypic view of social interactions in microbial communities. Annu. Rev. Genet. 47:247-273.

516   Mitri, S. and K. R. Foster. 2016. Pleiotropy and the low cost of individual traits promote cooperation. Evolution 70:488-494.

518   Monod, J. 1949. The growth of bacterial cultures. Annu. Rev. Microbiol. 3:371-394.

Nadell, C. D., J. B. Xavier, and K. R. Foster. 2009. The sociobiology of biofilms. FEMS
520   Microbiol. Rev. 33:206-224.

Nikaido, H. and J.-M. Pagès. 2012. Broad-specificity efflux pumps and their role in multidrug
522   resistance of Gram-negative bacteria. FEMS Microbiol. Rev. 36:340-363.

Özkaya, Ö., K. B. Xavier, F. Dionisio, and R. Balbontin. 2017. Maintenance of microbial
524   cooperation mediated by public goods in single and multiple traits scenarios. J. Bacteriol.

526   Popat, R., F. Harrison, A. C. da Silva, S. A. Easton, L. McNally, P. Williams, and S. P. Diggle. 2017. Environmental modification via a quorum sensing molecule influences the social
528   landscape of siderophore production. Proc. R. Soc. B 284.

Raihani, N. J., A. Thornton, and R. Bshary. 2012. Punishment and cooperation in nature. Trends
530   Ecol. Evol. 27:288-295.

Ratnieks, F. L. W., K. R. Foster, and T. Wenseleers. 2006. Conflict resolution in insect
532   societies. Annu. Rev. Entomol. 51:581-608.

Ratnieks, F. L. W. and T. Wenseleers. 2008. Altruism in insect societies and beyond: voluntary
534   or enforced? Trends Ecol. Evol. 23:45-52.

Rendueles, O., P. C. Zee, I. Dinkelacker, M. Amherd, S. Wielgoss, and G. J. Velicer. 2015. Rapid and widespread de novo evolution of kin discrimination. Proc. Natl. Acad. Sci. U.S.A. 112:9076-9081.

Ross-Gillespie, A., Z. Dumas, and R. Kümmerli. 2015. Evolutionary dynamics of interlinked public goods traits: an experimental study of siderophore production in *Pseudomonas aeruginosa*. J. Evol. Biol. 28:29-39.

Ross-Gillespie, A. and R. Kümmerli. 2014. Collective decision-making in microbes. Front. Microbiol. 5:54.

Sachs, J. L., U. G. Mueller, T. P. Wilcox, and J. J. Bull. 2004. The evolution of cooperation. Q. Rev. Biol. 79:135-160.

Sandoz, K. M., S. M. Mitzimberg, and M. Schuster. 2007. Social cheating in *Pseudomonas aeruginosa* quorum sensing. Proc. Natl. Acad. Sci. U.S.A. 104:15876-15881.

Sexton, D. J. and M. Schuster. 2017. Nutrient limitation determines the fitness of cheaters in bacterial siderophore cooperation. Nat. Commun. 8:230.

Smukalla, S., M. Caldara, N. Pochet, A. Beauvais, S. Guadagnini, C. Yan, M. D. Vinces, A. Jansen, M. C. Prevost, J.-P. Latgé, G. R. Fink, K. R. Foster, and K. J. Verstrepen. 2008. *FLO1* Is a variable green beard gene that drives biofilm-like cooperation in budding yeast. Cell 135:726-737.

van Gestel, J., F. J. Weissing, O. P. Kuipers, and A. T. Kovács. 2014. Density of founder cells affects spatial pattern formation and cooperation in *Bacillus subtilis* biofilms. ISME J. 8:2069-2079.

Velicer, G. J. and M. Vos. 2009. Sociobiology of the *Myxobacteria*. Annu. Rev. Microbiol. 63:599-623.

Wang, M., A. L. Schaefer, A. A. Dandekar, and E. P. Greenberg. 2015. Quorum sensing and policing of *Pseudomonas aeruginosa* social cheaters. Proc. Natl. Acad. Sci. U.S.A. 112:201500704.

Weigert, M. and R. Kümmerli. 2017. The physical boundaries of public goods cooperation between surface-attached bacterial cells. Proceedings of the Royal Society B: Biological Sciences 284:20170631.

Wenseleers, T. and F. L. W. Ratnieks. 2006. Enforced altruism in insect societies. Nature 444:50.

566   West, S. A., S. P. Diggle, A. Buckling, A. Gardner, and A. S. Griffin. 2007a. The social lives of microbes. Annu. Rev. Ecol. Evol. Syst. 38:53-77.

568   West, S. A., A. S. Griffin, and A. Gardner. 2007b. Evolutionary explanations for cooperation. Curr. Biol. 17:R661-R672.

570   West, S. A., A. S. Griffin, and A. Gardner. 2007c. Social semantics: altruism, cooperation, mutualism, strong reciprocity and group selection. J. Evol. Biol. 20:415-432.

572   Xavier, J. B., W. Kim, and K. R. Foster. 2011. A molecular mechanism that stabilizes cooperative secretions in *Pseudomonas aeruginosa*. Mol. Microbiol. 79:166-179.

Table 1

Item	Parameter	Description	Range
Cell	r	radius	0.5 - 1 µm
	D_c	dispersal	0 - 3.5 µm²s⁻¹
	μ	growth rate	1.0
Public good	p_{pg}	production rate	1.0 s⁻¹
	c_{pg}	production cost	0.001
	b	benefit	0.01
	d_{pg}	diffusion	1.0 – 7.0 µm²s⁻¹
	δ_{pg}	durability	500 s
Toxin	p_{tox}	production rate	1.0 s⁻¹
	$c_{tox} + c_{res}$	production & immunity cost	0.00025 - 0.004[*]
	c_{tox}	production cost	$0.000\bar{3}$[†]
	c_{res}	immunity cost	$0.000\bar{3}$[†]
	d_{tox}	diffusion	1.0 – 7.0 µm²s⁻¹
	δ_{tox}	durability	50 - 5000 s
	θ_T	potency	600 - 1800
	κ	latency	1 – 6

[*] varied in the simulations shown in Fig. 3.

578   [†] based on the simulations shown in Fig. 3, c_{tox} and c_{res} were fixed to these values for all subsequent analyses.

FIGURES

Time step: 00095 s   Time step: 03295 s   Time step: 04790 s

Figure 1. Point-in-time snapshots showing the spatial distribution of cooperating policers (in blue) and cheaters (in yellow) during competition. The point-in-time snapshots were taken during the early, intermediate and late simulation phase, depicting the growth of digital bacteria in their simulated environment. The density of public goods (in green) and toxins (in red) produced by the cooperating policers are indicated by the intensity level of their respective color. The specific parameters values for this example are: $D_c = 0.0$ µm^2/s, $d_{pg} = 1.0$ µm^2/s, $d_{tox} = 2.0$ µm^2/s, $\delta_{tox} = 500$ s, $\theta_T = 1000$ and $\kappa = 3.5$.

Figure 2. Flow diagram of the computer simulation showing the order of each procedure called during a single time step (i.e. one second). Our platform allows for two different types of culturing conditions: batch-culture growth, where simulations stop when populations reach the carrying capacity $K = 1000$ (solid lines); and continuous growth in a chemostat, where simulations stop after a given time period (dashed lines). To guarantee continuous growth, the chemostat cycle comprised a random cell removal mechanism, such that population size was kept constant at $K/2$. Each box describes a key element in a cell's life cycle, affecting its fitness: i.e. cells grew according their fitness equations (see main text) and divided when reaching the threshold radius of 1 µm (two times the initial radius). Since cells are free to move on a continuous landscape, they can overlap following diffusion. To remove cell overlap, we used a procedure based on the physics of hard-core interactions used in molecular dynamics and cell elasticity.

Figure 3. Growth in monoculture and the relative cost of policing. We examined how large the cost of policing ($c_{tox} + c_{res}$) can be relative to the cost of public good production (c_{pg}) to guarantee a net benefit of cooperation. To address this question, we compared the growth of the policing strain (producing toxins and public goods, in blue) with the growth of the cooperator strain (producing only public goods, in red) and the cheater strain (producing neither toxins nor public goods, in yellow) in monoculture for a range of cost ratios, under three different diffusion regimes. The yellow bar indicates that the time needed by the cheater strain to reach carrying capacity was not affected by the diffusivity of the environment. The red bars show that the time needed by the cooperator strain to reach carrying capacity increased with higher diffusivity of the environment (plain line: low diffusivity, $D_c = 0.0 \ \mu m^2/s$, $d_{pg} = d_{tox} = 1.0 \ \mu m^2/s$; dashed line: intermediate diffusivity, $D_c = 2.0 \ \mu m^2/s$, $d_{pg} = d_{tox} = 3.5 \ \mu m^2/s$; dotted line: high diffusivity, $D_c = 4.0 \ \mu m^2/s$, $d_{pg} = d_{tox} = 7.0 \ \mu m^2/s$). For policing to evolve, the time needed by the policer strain to reach carrying capacity must be within these boundaries (low diffusivity: blue triangles; intermediate diffusivity: diamonds, high diffusivity: dots). Values represent averages of 50 independent simulations. Standard errors of the mean are smaller than the size of the markers and therefore not shown. We set $c_{pg} = 0.001$, while varying $c_{tox} + c_{res}$ from 0.00025 to 0.004. The black encircled values correspond to $c_{pg}/(c_{tox} + c_{res}) = 1.5$, the relative policing cost used for all subsequent simulations.

Figure 4. Competition between cheaters and cooperators across a range of cell dispersal D_c and public good diffusion d_{pg} coefficients. Heat maps depict the frequency of the cooperator strain after the community reached stationary phase. (A) Outcome of competitions between the public-good-producing cooperator W and the cheater C in the absence of a policing mechanism. (B) Outcome of competitions between the policing cooperator P and the cheater C. D_c varied from 0.0 to 3.5 µm²/s, whereas d_{pg} varied from 1.0 - 7.0 µm²/s. The other parameters were set to intermediate values: $d_{tox} = 4.0$ µm²/s, $\delta_{tox} = 500$ s, $\theta_T = 1000$, $\kappa = 3.5$.

Figure 5. Heat maps exploring the effect of different toxin properties on the success of cooperation. We varied toxin diffusion d_{tox}, toxin durability δ_{tox}, toxin potency θ_T and toxin latency κ across a range of values (see Table 1) and show here heat maps of the frequency of the cooperator strain after the community reached carrying capacity for the lowest, highest and intermediate parameter values. At the center of the figure we placed a reference heat map (E), where all the parameters were set to intermediate values (d_{tox}= 4.0 μm²/s, δ_{tox} = 500 s, θ_T= 1000, κ = 3.5). We then varied toxin diffusion from 1.0 to 7.0 μm²/s (A, E and B), toxin durability from 50 to 5000 (C, E and D), toxin potency from 600 to 1800, (F, E and G) and toxin latency κ from 2.0 to 6.0 (H, E and I). We only varied a single parameter at the time, and kept all others at their intermediate default value used for (E).

Figure 6. Policing is selected against when the genetic linkage between public good and toxin production breaks. Panels show simulated three-way interactions between cheaters *C* (yellow lines), cooperating policers *P* (blue lines), and toxin-resistant non-policing cooperators *R* (red lines) across a range of public good and cell dispersal values. We assume that *R* can directly evolve from the policing strain *P* when the genetic linkage between public good and toxin production breaks. If this happens, our simulations reveal that policers are always selected against. Trajectories show strain frequency fluctuations in continuous cultures over 80,000 time steps. Values depict means of 50 independent replicates with the standard error of the mean (transparent areas). For all simulations, we kept toxin properties constant: d_{tox}= 7.0 µm²/s, δ_{tox} = 500 s, θ_T= 1000, κ = 3.5.

SUPPLEMENTARY FIGURES

Figure S1

Cell dispersal:	2.5 µm² s⁻¹
Public good diffusion:	3.0 µm² s⁻¹
Toxin diffusion:	4.0µm² s⁻¹
Toxin potency:	1000
Toxin durability:	500 s
Toxin latency:	3.5

Figure S1. Point-in-time snapshots showing the spatial distribution of cooperating policers (in blue) and cheaters (in yellow) during competition under three different starting conditions. The point-in-time snapshots were taken during the early, intermediate and late simulation phase depicting the growth of digital bacteria in their simulated environments. The only parameter we varied in these examples was public good diffusion d_{pg} = 3. The density of public goods and toxins are indicated by the intensity level of the background color. In green are the distribution of public goods and in red the toxins.

Figure S2

Figure S2. Effect of toxin latency κ on the growth rate of sensitive cells. Small values of κ lead to an immediate negative impact on growth, whereas larger values of κ extend the latency phase during which the growth of sensitive cells is not affected.

Figure S3

Figure S3. Monod's equation explains the relationship between the relative cost of policing and strain fitness. We simulated the growth of the policing strain (producing a toxin and a public good) in monoculture for a range of $c_{pg}/(c_{tox} + c_{res})$ ratios, under three different diffusion regimes. We kept the cost of public good production constant at $c_{pg} = 0.001$, while varying the cost of toxins ($c_{tox} + c_{res}$) from 0.00025 to 0.004. The fitness of the policing strain (measured as the time required to reach carrying capacity) declined with higher relative toxin costs. The decline was adequately captured by Monod's equation under conditions of (A) low diffusion (coefficient of correlation R = 0.999); (B) medium diffusion (R = 0.991); and (C) high diffusion (R = 0.997). However, the equation does not fully capture the observed pattern when the ratio approaches one, namely when $(c_{tox} + c_{res}) \approx c_{pg}$, under conditions of low environmental diffusivity. This is partially understandable since $R(\tau_K)$ does not directly take the viscosity of the medium into account. Details on diffusion parameter values can be obtained from Figure 3. Open circles show means of 50 independent replicates with the standard error of the mean.