Cell type-dependent control of p53 transcription and enhancer activity by p63

Gizem Karsli Uzunbas, Faraz Ahmed, and Morgan A. Sammons*

Department of Biological Sciences, State University of New York at Albany, Albany, NY 12222

* Corresponding Author: masammons@albany.edu

Abstract

Transcriptional activation by p53 provides powerful, organism-wide tumor suppression. In this work, we demonstrate that the p53-induced transcriptome is dependent on pre-establishment of cell type-dependent *cis*-regulatory networks. The epithelial-specific p53 transcriptome is strongly dependent on the p53 family member p63, which acts as a pioneer factor for epithelial-specific enhancers. These results suggest a broad mechanism for regulating the p53-dependent cellular response to stress through differential regulation of *cis*-regulatory elements and identify p63 as a direct and critical regulator of enhancer activity in epithelial cell types.

Introduction

The p53 family of transcription factors regulate highly diverse cellular functions, including tumor suppression and control of cell specification and identity (Levrero et al. 2000). p53 is a master tumor suppressor that protects organismal fidelity after exposure to cellular stress like DNA damage. This activity is dependent on p53's ability to activate transcription of a canonical network of genes involved in DNA damage repair, cell cycle arrest, apoptosis, and senescence (Kastenhuber and Lowe 2017). Surprisingly, many of these transcriptional pathways are individually dispensable for tumor suppression, suggesting p53 regulates a redundant and not-yet fully characterized transcriptional network (Andrysik et al. 2017; Li et al. 2012).

Genetic inactivation of the p53 tumor suppressor pathway is highly recurrent across cancer types. p53 mutations, while frequent, vary depending on the tumor type with additional genetic and epigenetic mechanisms proposed to inactivate the p53 pathway in the presence of wild-type p53 (Zhu et al. 2016; Ciriello et al. 2013; Kandoth et al. 2013). p63 and p73, p53 family members, function similarly to p53 in the stress response, although their precise roles in tumor suppression are unresolved (Flores et al.

2002; Botchkarev and Flores 2014; DeYoung and Ellisen 2007). p63 and p73 are

primarily lineage restricted to epithelial cell types where each serves critical and non-

overlapping roles in cell identity and self-renewal (Melino et al. 2015; Levrero et al.

2000). Mutations within p63 are causative for a number of epithelial-associated human

developmental disorders independent of p53 activity, and mutations in p63 target genes

underlie similar phenotypes (Fakhouri et al. 2014; Brunner et al. 2002). A significant and

still outstanding question involves dissection of specific roles and functional interplay

between p53 family members during development, in the regulation of cellular

homeostasis, and in the etiology of disease.

The similarity between DNA binding motifs suggested that competition for binding

sites serves a central role in the regulation of p53 family member activity (Moll et al.

2001; Billant et al. 2016). Indeed, previous studies implicate the ΔNp63 isoforms,

lacking transactivation domains, as direct repressors of the p53-induced transcriptome

through a binding site competitive mechanism (Yang et al. 2006, 1998; Mundt et al.

2010). Amplification of p63 is thought to drive a number of epithelial cancers,

particularly squamous cell carcinomas, although both p53-dependent and independent

mechanisms have been proposed for this driver activity (Saladi et al. 2017). As the

majority of cancers are derived from epithelial tissues, the mechanisms of p53 family

dependent tumor suppression in those tissues is of special interest (Kandoth et al.

2013).

Fate choice after p53 activation, be it apoptosis, temporary/permanent cell cycle

arrest, or continued proliferation, is variable across cell types suggesting that p53-

dependent transcription is also cell type-dependent (Kastenhuber and Lowe 2017). A

bevy of sensitive methodological approaches have been used to identify p53 binding

sites and gene targets across transformed and primary cell lines in an effort to explain

these terminal cell fate choices and p53-dependent tumor suppression. A recent set of

meta-analyses have suggested that p53 binding to the genome is invariant (Verfaillie et

al. 2016), proposing that p53 acts independently to drive gene expression of a core

tumor suppressor network across all cell types due to the low enrichment of other

transcription factor motifs at p53-bound enhancers and the reported pioneer factor

activity of p53 (Verfaillie et al. 2016; Andrysik et al. 2017; Younger and Rinn 2017).

Conversely, similar experimental approaches demonstrate cell type-specific p53 binding

and gene targets although the mechanisms driving these differential activities are

unknown (Hafner et al. 2017). Additional experimental evidence and models are

required to unravel these mutually exclusive p53 regulatory mechanisms.

We previously proposed that the local chromatin environment, including variable

chromatin accessibility and enhancer activity, contributes to novel p53 activities across

cell types (Sammons et al. 2015). To address this question, we performed genomewide

transcriptome, epigenome, and p53 cistrome profiling in primary dermal foreskin

fibroblasts (SkFib) and breast epithelial (MCF10A) cell lines, two cell types with varying

enhancer activity at p53 binding sites (Sammons et al. 2015). Our results directly

implicate differential *cis*-regulatory element activity as a mediator of the p53 network,

with both promoter and enhancer activity contributing to p53-dependent gene

expression differences. Further, we have identified the p53 family member p63 as one

factor that drives the epithelial-specific p53 transcriptome through an enhancer

maintenance activity. We further propose that p63 serves as licensing factor for a set of

epithelial-specific enhancers. Thus, these data support a mechanism whereby co-

operating transcription factors control cell type-dependent *cis*-regulatory networks that

regulate p53 activity.

Results and Discussion

We used polyA-enriched RNA-seq to assess gene expression dynamics in model

cell lines (MCF10A and SkFib) after p53 activation in response to Nutlin-3A treatment

(Fig. 1A, Supplemental Fig. S1A). We reasoned Nutlin-3A, a non-genotoxic activator of

p53, would better reflect p53-dependent transcriptional activity relative to genotoxic

activation methods as DNA damage-dependent, but p53-independent, signaling would

not confound our analysis (Vassilev et al. 2004; Allen et al. 2014). Consistent with a cell

type-dependent activity of p53, MCF10A cells displayed a broad and specific set of

Nutlin-3A-induced genes, whereas SkFib showed a markedly smaller, albeit specific,

p53-activated transcriptome (Fig. 1B). Orthogonal validation by qRT-PCR confirmed a

stringent cell type-specificity of each set of genes and demonstrated that canonical p53

targets, like *CDKN1A* and *BTG2* are indeed shared across cell types (Supplemental Fig.

S1B-D). Decreasing the fold-change cutoff yielded an increased number of differentially

regulated genes for each cell type, but maintained the trend that p53-dependent gene

targets are more abundant in MCF10A (Supplemental Fig. S1E). Genes upregulated in

both cell lines are significantly associated with the core p53 response (Supplemental

Fig. S1F), as expected. MCF10A-specific upregulated genes are enriched in gene

ontology groups related to establishment of the skin barrier and p53-dependent

processes like programmed cell death and homeostasis. In contrast, SkFib-specific

targets are enriched in other cellular stress-related pathways including those associated

with the hypoxia response and catabolic processes (Supplemental Fig. S1F,

Supplemental Table S1). The majority of the MCF10A-specific targets represent novel

p53-dependent genes and over 20% of SkFib-specific targets have not been observed

previously (Supplemental Fig. S1G, Supplemental Tables S2-3 (Fischer 2017)). These

initial analyses reveal that the p53-activated transcriptome is highly variable between

non-transformed cell types and may reflect the need for tailored, cell specific responses

to p53 activation after cellular stress.

We next confirmed that a set of cell type-specific targets were p53-dependent

(Fig. 1C-D, Supplemental S1H). Indeed, knockdown of p53 in SkFib led to sharp

reduction in both basal and Nutlin-3A-induced expression of the SkFib-specific genes,

GDNF and *TRIM55* (Fig. 1D). Depletion of p53 in MCF10A (Fig. 1E, Supplemental Fig.

S1I) led to a loss of Nutlin-3A-induced *RIC3* and *IL1B* expression relative to a non-

targeting control shRNA (Fig. 1F), indicating these genes are *bona fide* p53 target

genes in epithelial cells.

Recent analyses reached opposing conclusions with regard to the cell type-

dependence of p53 engagement with the genome (Verfaillie et al. 2016; Hafner et al.

2017). We thus assessed whether differences in p53 engagement with the genome in

our model cell lines might explain our observations of cell type-dependent

transcriptomes. Consistent with cell type-dependence, genomewide ChIP-seq revealed

a highly-enriched set of MCF10A-specific p53 binding sites and a smaller number of

SkFib-specific sites, mirroring our gene expression observations (Fig. 1G). p53 binding

at common sites as measured by ChIP-seq mirrors both the basal and induced

expression of p53 (Supplemental Fig. S1J). p53 protein expression and binding do not

correlate with the ability to activate common targets as SkFib displayed dramatically

increased Nutlin-3A induced gene activation compared to MCF10A (Supplemental Fig.

S1K). Common p53 binding events are primarily within 10kb of genes upregulated in

both cell types, similar to previous results suggesting that many direct p53 targets have

proximal p53 binding (Supplemental Fig. S1L). MCF10A-specific genes have a higher

percentage of proximal p53 binding relative to SkFib. Notably, we observe both

MCF10A and SkFib p53 binding proximal to SkFib-specific genes. Cell type-specific

binding events displayed lower p53 occupancy than at common sites, suggesting that

common sites might represent more stable or higher affinity p53 binding sites

(Supplemental Fig. S1M). Taken together, these data provide evidence that the p53-

dependent transcriptome is indeed cell type-specific and that p53 engagement with the

genome is variable across cell types.

We next wanted to identify potential mechanisms driving the strong, cell type-

specificity within the p53 transcriptome and cistrome. The most straightforward

hypothesis is that differential p53 binding drives cell type-specific transcriptome

differences, but our data suggest that this might only explain MCF10A behavior (Fig.

1G, Supplemental Fig. S1L). Thus, we examined *GDNF*, a SkFib-specific gene, and

observed proximal p53 binding upstream of the *GDNF* transcriptional start site (Fig. 2A).

We also noted p53 binding to this region in MCF10A, even though p53-dependent

GDNF expression is SkFib-specific (Fig. 2A). Similar p53 proximal binding was

observed at SkFib-specific genes *TRIM55* and *MAMDC2* (Supplemental Fig. S2A),

suggesting that differential proximal p53 binding cannot explain the cell type-specificity

observed in SkFib relative to MCF10A. We and others previously proposed that

enhancer activity is critical for regulating p53-dependent gene expression(Sammons et

al. 2015; Sullivan et al. 2018). As enhancers also have a strong cell type-

dependence(Calo and Wysocka 2013; Heinz et al. 2015), we performed ChIP-seq

experiments for canonical promoter (H3K4me3) and enhancer-associated (H3K4me2,

H3K27ac) histone modifications to determine whether differentially active promoters or

enhancers might explain the observed cell type specificity of the p53 transcriptome.

The local chromatin environment at the *GDNF* proximal p53 binding site is

enriched for H3K27ac and H3K4me2, but lacks H3K4me3, suggestive of an active

enhancer in both cell types (dashed box, Fig. 2A). Consistent with high basal and p53-

inducible expression of *GDNF* in SkFib, the *GDNF* promoter is strongly enriched for the

canonical promoter-associated histone modification H3K4me3 (solid box, Fig. 2A) which

is completely absent from MCF10A. We observed differential promoter activity at other

SkFib-specific p53 target promoters (Supplemental Fig. S2A) without observed

differences in p53 binding. It is currently unclear whether the lack of activity in MCF10A

is driven by lack of promoter-bound cofactors, through repressive chromatin-associated

mechanisms, or both. Repressive DNA and histone methylation have been implicated in

differential control of p53 activity in cancer cell lines (Hermeking et al. 1997; Mungamuri

et al. 2012; París et al. 2008). These observations suggest that promoter competence,

independent of p53 binding, potently controls cell specific p53 transcription responses.

We then sought to better understand the mechanisms driving the epithelial-

specific p53 target genes. p53 binding is pervasive in MCF10A relative to SkFib (Fig.

1G), therefore we investigated the connection between *de novo* epithelial p53 binding

and the induction of specific targets. Examination of the MCF10A-specific targets *RIC3*

and *MYBPHL* revealed MCF10A-specific binding of p53 binding to putative intragenic

enhancers specifically in MCF10A (Fig. 2B, Supplemental Fig. S2B). This is consistent

with the lack of basal expression of *RIC3* mRNA in SkFib and p53-dependent activation

in MCF10A. We speculated that MCF10A-specific p53 binding might correlate with

enhancer signatures specific to epithelial cell types. MCF10A-specific p53 binding is

flanked by enrichment of MCF10A-specific H3K27ac and H3K4me2 (Fig. 2C). These

sites also lack H3K4me3 (Supplemental Fig. S2C), suggesting they are putative

enhancers. Common p53 binding sites are enriched for enhancer-associated chromatin

modifications in both cell types whereas cell type-specific p53 binding overlaps cell

type-specific enhancer modifications (Supplemental Fig. S2D-G). Taken together, these

data demonstrate that cell specific p53 binding is correlated with enhancers unique to

that cell type. These observations thus provide a putative mechanism for cell type-

dependent responses. Transcriptional activation requires both gene-proximal p53

binding and pre-establishment of *cis*-regulatory elements. While both are necessary,

neither is sufficient to confer p53-dependent gene activation.

We next investigated what might drive the cell type-dependence and activity of

p53-bound enhancers. Enhancer specification is generally believed to be regulated by

the combined effort of multiple common and lineage-dependent transcription factors

(Heinz et al. 2015). Multiple recent reports suggest that p53 activity at *cis*-regulator

elements is independent of any accessory transcription factors, primarily due to low

enrichment/specificity of other DNA motifs (Younger and Rinn 2017; Verfaillie et al.

2016; Sullivan et al. 2018). Indeed, MCF10A enhancers are highly enriched for

underlying p53/p63 motifs relative to SkFib, consistent with results from the FANTOM

consortium and a lack of any obvious epithelial-specific transcription factor motifs

(Supplemental Fig. S2H, Supplemental Table S4, (The FANTOM Consortium et al.

2014)). As the p53 family shares a common, although not identical, response element

motif, we speculated that p53 family members might regulate MCF10A-specific p53

binding events or enhancer activity and might explain previous observations.

Epithelial cells highly express p63 and are dependent on p63 activity for self-

renewal and epidermal commitment *in vivo* (Yang et al. 1999). p63 also bookmarks

enhancers during epidermal differentiation (Kouwenhoven et al. 2015) and may regulate

chromatin accessibility through interactions with chromatin modifiers (Bao et al. 2015;

Lin-Shiao et al. 2018). p63 is expressed only in MCF10A which do not express the other

p53 family member p73 (Supplemental Fig. S3A). Therefore, we focused on the function

of p63 in epithelial enhancer activity by performing genomewide ChIP-seq of p63 in

MCF10A after DMSO or Nutlin-3A treatment. Examination of *IL1B* and *RNASE7*,

MCF10A-specific p53 targets, displayed proximal p63 binding to distinct enhancer

regions co-occupied by p53 in an MCF10A-specific manner (Fig. 3A, Supplemental Fig.

S3B). Globally, p63 binds to significantly more genomic loci than does p53 (Fig. 3B), but

over 75% of epithelial-specific p53 binding events overlap p63 binding. Shared binding

sites display higher occupancy and are more likely to have enhancer-associated

chromatin modifications than do sites unique to either p53 or p63 (Fig. 3C,

Supplemental Fig. S3C), suggesting that both p53 and p63 might be necessary for full

enhancer activity.

Overall, p63 binding correlates with epithelial-specific enhancer modifications

and co-occupies the majority of p53 sites (Fig. 3B, Supplemental Fig. S3D). The modal

binding of p63 is within 10kB of the TSS of Nutlin-3A-induced genes, further suggesting

that p63 might co-regulate these specific target genes (Supplemental Fig. S3D). In

order to test whether p63 regulates p53 activity, we created MCF10A cells stably-

expressing shRNA against p63 (Fig. 3D, Supplemental Fig. S3E). We then performed

RNA-seq analysis of MCF10A cells expressing non-targeting (scr), p53, or p63 shRNA

to determine their relative contributions to epithelial-specific target genes. Loss of p53

expression strongly inhibited Nutlin-3A-induced gene expression consistent with this set

of genes representing true p53 targets (Fig. 3E). Knockdown of p63 showed more

pleiotropic effects. Globally, Nutlin-3A-induced target gene activation is inhibited in p63

knockdown relative to non-targeting controls, suggesting p63 activity is required for a

set of epithelial p53 targets (Fig. 3E). Whereas 40% of p53 target genes show defective

induction in the absence of p63, the remainder of the genes are either unaffected or

display increased p53-dependent transactivation (Supplemental Fig. S3F). This

observation is consistent with previous reports of direct repression of p53-dependent

transactivation through binding site competition (Mundt et al. 2010; Yang et al. 1998).

Our data suggest that p63 also functions to co-activate a set of p53-dependent

genes specifically in epithelial cell types. We then validated these results by qRT-PCR

on a panel of epithelial target genes after p53 or p63 knockdown. *RIC3,* previously

demonstrated to be a p53-dependent gene (Fig. 1F), is also derepressed and shows

higher p53-dependent transactivation in the absence of p63 (Fig. 3F). Conversely, both

IL1A and *IL1B* are dependent on p53 and p63 for full transactivation, with *IL1B* showing

a strong dependence on p63 for basal expression (Fig. 3F). Knockdown of p63 also led

to basal gene expression changes for p53 targets (Supplemental Fig. S3G) and genes

related to cell cycle and cell adhesion/extracellular matrix (Supplemental Table S5), as

has previously been described (Carroll et al. 2006).

In total, these results suggest that p63 functions as both a co-activator and

repressor of p53-dependent transcription in MCF10A, but the mechanisms driving these

activities were not clear. p53 and p63 engagement to the genome in epithelial cells

correlates with enhancer activity and the p53 family are putative pioneer factors with

enhancer bookmarking activity (Kouwenhoven et al. 2015; Bao et al. 2015; Sammons et

al. 2015; Cui and Zhurkin 2014). Therefore, we examined the effect of p53 and p63 on

enhancer identity in MCF10A. Globally, loss of p63 led to a dramatic reduction of

H3K27ac and H3K4me2 at p63-bound enhancers (Fig. 4A-B). Enhancer-associated

modifications were unaffected by loss of p53, nor were the genomewide complement of

enhancers affected by loss of either p53 or p63 (Fig. 4A-B), suggesting this is a

phenomenon specific to p63-bound enhancers. Expression of *RRM1*, a member of the

ribonucleotide reductase enzyme complex, was significantly downregulated by loss of

p63 and unaffected in the absence of p53. p63-bound enhancers upstream of *RRM1*

and *EDN2*, a p53 target gene, showed dramatic loss of H3K27ac and H3K4me2

enrichment in p63-depleted cells, with similar loss of enhancer-associated modifications

observed broadly at p63 binding sites (Fig. 4A-C, Supplemental Fig. S4A). In total, over

30% and 20% of p63-bound H3K27ac and H3K4me2 sites, respectively, showed at

least a 2-fold reduction in MCF10A-p63sh cells relative to scrambled shRNA (Fig. 4D).

The loss of enhancer-associated histone modifications at p63 binding sites proximal to

genes dependent on p63 expression suggests that p63 directly controls enhancer

activity (Supplemental Fig. S4B).

We then investigated additional mechanisms that might contribute to our

observations of p63-dependent enhancer activity. p63-bound enhancers with the

highest-dependence on p63 for H3K4me2 enrichment were significantly more likely to

overlap epithelial-specific DNase hypersensitive sites than those bound by p63, but not

affected by p63 depletion (Supplemental Fig. S4C, Supplemental Table S6). MCF10A-

specific p63 binding sites are strongly enriched for enhancer-associated modifications

across epithelial cell types (Supplemental Fig. S4D). The p63-insensitive enhancers

were broadly accessible across cell types, suggesting that these enhancers, while

bound by p63, are not p63 dependent (Fig. 4E). Indeed, analysis of transcription factor

motifs within p63-bound enhancers revealed a strong enrichment of ETS family

transcription factors in the p63-insensitive enhancers (Fig. 4F, Supplemental Table S7).

These data suggest that the activity of p63-insensitive enhancers is dependent on other

transcription factors. Therefore, our results demonstrate that p63 activity is critical for

local H3K27ac and H3K4me2 enrichment at a set of epithelial-specific enhancers and

that p63 binds to another set of cell type-independent enhancers putatively dependent

on transcription factor families like ETS. Combinatorial transcription factor binding at

enhancers is critical for appropriate spatial and temporal gene expression (Reiter et al.

2017; Shlyueva et al. 2014), and the collaborators at p53 family-bound enhancers are

relatively understudied compared to promoters(Sullivan et al. 2018). Recent *in vivo*

dissection of p53 enhancer activity implicated the general enhancer-associated AP-1

family member CEBPB as a direct regulator of an enhancer upstream of *CDKN1A*

(Korkmaz et al. 2016).

Our data indicates that p63 might be a pioneer factor for a set of epithelial-

specific enhancers, but not the entire complement of p63-bound enhancers. A number

of studies have demonstrated that p53 has partial pioneer factor activity as it can

directly bind its consensus sequence in the context of a nucleosome. p53 may possess

pioneering activity, but the large majority of p53 binding sites are constitutively closed or

are within accessible chromatin (Younger and Rinn 2017; Sammons et al. 2015;

Andrysik et al. 2017). Therefore, we examined whether p53 might serve pioneer or

maintenance roles at enhancers. Basal H3K4me2 and H3K27ac enrichment at p53

binding sites was essentially unaffected by the loss of p53 (Fig 4G). p63 knockdown led

to a loss of H3K27ac and H3K4me2 enrichment at p53 binding sites (Fig. 4G,

Supplemental Fig. S4E), an expected result as approximately 75% of p53 binding

events in MCF10A cells are also p63 binding sites (Fig. 3B). Examination of p53+/p63-

binding events displayed a similar outcome, with both H3K27ac and H3K4me2

unaffected by p53 loss (Supplemental Fig. S4F). Surprisingly, enhancer-associated

histone modifications at these sites were still partially dependent on p63 (Supplemental

Fig. S4F), potentially due to sub-peak threshold binding of p63. p53 is required for *de*

novo Nutlin-3A-induced H3K27ac levels at p53 binding sites (Fig. 4G-H). p63 depletion,

while affecting basal enrichment, has a milder effect on Nutlin-3A-induced H3K27ac

(Fig. 4G). p53 activation restores H3K27ac to near wild-type levels in p63-depleted

MCF10A (Fig. 4G). Consistent with p63 as both an activator and repressor of p53-

dependent transcription, basal levels of H3K27ac are reduced in the absence of p63 but

are derepressed upon p53 activation (Fig. 4G-H).

Epithelial cell types like MCF10A express multiple p53 family members that might

redundantly regulate enhancer activity. We therefore extended our analysis to isogenic

WT or *TP53-/-* HCT116 human colon carcinoma cell lines and WT or *Trp53-/-* mouse

embryonic fibroblasts (MEF). Both cell models lack endogenous p63 and thus should

allow us to define the role of p53 in enhancer maintenance and activity. Similar to our

observations in MCF10A cells, basal enhancer-associated H3K4me1/2 methylation at

p53-bound enhancers is unaffected by the loss of p53 (Fig. 4I-J). These data suggest

that p53 is not an enhancer maintenance factor and contributes primarily to *de novo*

histone acetylation and transcriptional activation at already established enhancers. p63,

on the other hand, is critical for maintenance of epithelial-specific enhancers and

positively influences p53 transcriptional activity through these *cis*-regulatory regions.

Therefore, p53 and p63 have both their previously reported antagonistic relationship

and a more broadly cooperative partnership than expected (Botchkarev and Flores

2014; Flores et al. 2002). The requirement of p63 for epithelial-specific p53 activity

supports a model whereby the p53-dependent transcriptome is licensed by the

enhancer regulatory abilities of other transcription factors. This model is a particularly

attractive avenue for further inquiry as it provides a straightforward mechanism for

driving cell type-dependent tumor suppressor and homeostatic activities of p53.

Materials and Methods

Cell Culture and Treatment

MCF10A mammalian epithelial cells and skin foreskin fibroblast cells (AG22153, Coriell Institute) were cultured at 37˚C in 5% CO_2 in HuMEC Complete media (Gibco, #12752010) and DMEM media (with 10% FBS and 1% Pen/Strep, Gibco), respectively. Wild-type and p53-knockout HCT116 colon carcinoma and mouse embryonic fibroblasts were kind gifts of Carol Prives and Jing Huang, respectively. To induce p53 activation, cells were treated with Nutlin-3A (5 μM final) or DMSO (vehicle control) for 6 hours.

Western Blotting

Cells were lysed in RIPA buffer (10 mM Tris-HCl (pH 8.0), 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS and 140 mM NaCl, supplemented freshly with protease inhibitors) and probed with antibodies against p53 (BD Biosciences; BD554293), p63 (Cell Signaling; 13102), and GAPDH (Cell Signaling; 5174).

Other detailed experimental procedures are described in the Supplemental Materials.

Data Availability

All raw data associated with this study are pending at NCBI Gene Expression Omnibus and accession numbers will be available upon publication.

Acknowledgments

We are grateful to Allison Catizone and Enrique Lin Shiao for critical feedback on this manuscript. We thank Jason Herschkowitz for the MCF10A cell line and the University at Albany CFG for sequencing support. This work is supported by start-up funds from the State University of New York at Albany.

References

Allen MA, Andrysik Z, Dengler VL, Mellert HS, Guarnieri A, Freeman JA, Sullivan KD, Galbraith MD, Luo X, Kraus WL, et al. 2014. Global analysis of p53-regulated transcription identifies its direct targets and unexpected regulatory mechanisms. *Elife* **3**: e02200.

Andrysik Z, Galbraith MD, Guarnieri AL, Zaccara S, Sullivan KD, Pandey A, MacBeth M, Inga A, Espinosa JM. 2017. Identification of a core TP53 transcriptional program with highly distributed tumor suppressive activity. *Genome Res* **27**: 1645–1657.

Bao X, Rubin AJ, Qu K, Zhang J, Giresi PG, Chang HY, Khavari PA. 2015. A novel ATAC-seq approach reveals lineage-specific reinforcement of the open chromatin landscape via cooperation between BAF and p63. *Genome Biol* **16**: 284.

Billant O, Leon A, Le Guellec S, Friocourt G, Blondel M, Voisset C. 2016. The dominant-negative interplay between p53, p63 and p73: A family affair. *Oncotarget* **7**: 69549–69564.

Botchkarev VA, Flores ER. 2014. p53/p63/p73 in the epidermis in health and disease. *Cold Spring Harb Perspect Med* **4**.

Brunner HG, Hamel BC, Van Bokhoven H. 2002. The p63 gene in EEC and other syndromes. *J Med Genet* **39**: 377–81.

Calo E, Wysocka J. 2013. Modification of Enhancer Chromatin: What, How, and Why? *Mol Cell* **49**: 825–837.

Carroll DK, Carroll JS, Leong C-O, Cheng F, Brown M, Mills AA, Brugge JS, Ellisen LW. 2006. p63 regulates an adhesion programme and cell survival in epithelial cells. *Nat Cell Biol* **8**: 551.

Ciriello G, Miller ML, Aksoy BA, Senbabaoglu Y, Schultz N, Sander C. 2013. Emerging landscape of oncogenic signatures across human cancers. *Nat Genet* **45**: 1127–33.

Cui F, Zhurkin VB. 2014. Rotational positioning of nucleosomes facilitates selective binding of p53 to response elements associated with cell cycle arrest. *Nucleic Acids Res* **42**: 836–47.

DeYoung MP, Ellisen LW. 2007. p63 and p73 in human cancer: defining the network. *Oncogene* **26**: 5169–5183.

Fakhouri WD, Rahimov F, Attanasio C, Kouwenhoven EN, Ferreira De Lima RL, Felix TM, Nitschke L, Huver D, Barrons J, Kousa YA, et al. 2014. An etiologic

regulatory mutation in IRF6 with loss- and gain-of-function effects. *Hum Mol*
Genet **23**: 2711–20.

Fischer M. 2017. Census and evaluation of p53 target genes. *Oncogene* **36**: 3943–
3956.

Flores ER, Tsai KY, Crowley D, Sengupta S, Yang A, McKeon F, Jacks T. 2002. p63
and p73 are required for p53-dependent apoptosis in response to DNA damage.
Nature **416**: 560–564.

Hafner A, Lahav G, Stewart-Ornstein J. 2017. Stereotyped p53 binding tuned by
chromatin accessibility. *bioRxiv* 177667.

Heinz S, Romanoski CE, Benner C, Glass CK. 2015. The selection and function of cell
type-specific enhancers. *Nat Rev Mol Cell Biol* **16**: 144–154.

Hermeking H, Lengauer C, Polyak K, He T-C, Zhang L, Thiagalingam S, Kinzler KW,
Vogelstein B. 1997. 14-3-3σ Is a p53-Regulated Inhibitor of G2/M Progression.
Mol Cell **1**: 3–11.

Kandoth C, McLellan MD, Vandin F, Ye K, Niu B, Lu C, Xie M, Zhang Q, McMichael JF,
Wyczalkowski MA, et al. 2013. Mutational landscape and significance across 12
major cancer types. *Nature* **502**: 333–9.

Kastenhuber ER, Lowe SW. 2017. Putting p53 in Context. *Cell* **170**: 1062–1078.

Kenzelmann Broz D, Spano Mello S, Bieging KT, Jiang D, Dusek RL, Brady CA, Sidow
A, Attardi LD. 2013. Global genomic profiling reveals an extensive p53-regulated
autophagy program contributing to key p53 responses. *Genes Dev* **27**: 1016–31.

Korkmaz G, Lopes R, Ugalde AP, Nevedomskaya E, Han R, Myacheva K, Zwart W,
Elkon R, Agami R. 2016. Functional genetic screens for enhancer elements in
the human genome using CRISPR-Cas9. *Nat Biotechnol* **34**: 192–198.

Kouwenhoven EN, Oti M, Niehues H, van Heeringen SJ, Schalkwijk J, Stunnenberg
HG, van Bokhoven H, Zhou H. 2015. Transcription factor p63 bookmarks and
regulates dynamic enhancers during epidermal differentiation. *EMBO Rep* **16**:
863–78.

Levrero M, Laurenzi VD, Costanzo A, Gong J, Wang JY, Melino G. 2000. The
p53/p63/p73 family of transcription factors: overlapping and distinct functions. *J*
Cell Sci **113**: 1661–1670.

Li T, Kon N, Jiang L, Tan M, Ludwig T, Zhao Y, Baer R, Gu W. 2012. Tumor
suppression in the absence of p53-mediated cell-cycle arrest, apoptosis, and
senescence. *Cell* **149**: 1269–83.

Lin-Shiao E, Lan Y, Coradin M, Anderson A, Donahue G, Simpson CL, Sen P, Saffie R,
Busino L, Garcia BA, et al. 2018. KMT2D regulates p63 target enhancers to
coordinate epithelial homeostasis. *Genes Dev* **32**: 181–193.

Melino G, Memmi EM, Pelicci PG, Bernassola F. 2015. Maintaining epithelial stemness
with p63. *Sci Signal* **8**: re9.

Moll UM, Erster S, Zaika A. 2001. p53, p63 and p73--solos, alliances and feuds among
family members. *Biochim Biophys Acta* **1552**: 47–59.

Mundt HM, Stremmel W, Melino G, Krammer PH, Schilling T, Müller M. 2010. Dominant
negative (DeltaN) p63alpha induces drug resistance in hepatocellular carcinoma
by interference with apoptosis signaling pathways. *Biochem Biophys Res
Commun* **396**: 335–341.

Mungamuri SK, Benson EK, Wang S, Gu W, Lee SW, Aaronson SA. 2012. p53-
mediated heterochromatin reorganization regulates its cell fate decisions. *Nat
Struct Mol Biol* **19**: 478–484.

París R, Henry RE, Stephens SJ, McBryde M, Espinosa JM. 2008. Multiple p53-
independent gene silencing mechanisms define the cellular response to p53
activation. *Cell Cycle* **7**: 2427–2433.

Reiter F, Wienerroither S, Stark A. 2017. Combinatorial function of transcription factors
and cofactors. *Curr Opin Genet Dev* **43**: 73–81.

Saladi SV, Ross K, Karaayvaz M, Tata PR, Mou H, Rajagopal J, Ramaswamy S, Ellisen
LW. 2017. ACTL6A Is Co-Amplified with p63 in Squamous Cell Carcinoma to
Drive YAP Activation, Regenerative Proliferation, and Poor Prognosis. *Cancer
Cell* **31**: 35–49.

Sammons MA, Zhu J, Drake AM, Berger SL. 2015. TP53 engagement with the genome
occurs in distinct local chromatin environments via pioneer factor activity.
Genome Res **25**: 179–88.

Shlyueva D, Stampfel G, Stark A. 2014. Transcriptional enhancers: from properties to
genome-wide predictions. *Nat Rev Genet* **15**: 272–286.

Sullivan KD, Galbraith MD, Andrysik Z, Espinosa JM. 2018. Mechanisms of
transcriptional regulation by p53. *Cell Death Differ* **25**: 133–143.

The FANTOM Consortium, Andersson R, Gebhard C, Miguel-Escalada I, Hoof I,
Bornholdt J, Boyd M, Chen Y, Zhao X, Schmidl C, et al. 2014. An atlas of active
enhancers across human cell types and tissues. *Nature* **507**: 455–461.

Vassilev LT, Vu BT, Graves B, Carvajal D, Podlaski F, Filipovic Z, Kong N, Kammlott U,
Lukacs C, Klein C, et al. 2004. In vivo activation of the p53 pathway by small-
molecule antagonists of MDM2. *Science* **303**: 844–848.

Verfaillie A, Svetlichnyy D, Imrichova H, Davie K, Fiers M, Kalender Atak Z, Hulselmans
G, Christiaens V, Aerts S. 2016. Multiplex enhancer-reporter assays uncover
unsophisticated TP53 enhancer logic. *Genome Res* **26**: 882–895.

Yang A, Kaghad M, Wang Y, Gillett E, Fleming MD, Dötsch V, Andrews NC, Caput D,
McKeon F. 1998. p63, a p53 Homolog at 3q27–29, Encodes Multiple Products
with Transactivating, Death-Inducing, and Dominant-Negative Activities. *Mol Cell*
2: 305–316.

Yang A, Schweitzer R, Sun D, Kaghad M, Walker N, Bronson RT, Tabin C, Sharpe A,
Caput D, Crum C, et al. 1999. p63 is essential for regenerative proliferation in
limb, craniofacial and epithelial development. *Nature* **398**: 714–8.

Yang A, Zhu Z, Kapranov P, McKeon F, Church GM, Gingeras TR, Struhl K. 2006.
Relationships between p63 Binding, DNA Sequence, Transcription Activity, and
Biological Function in Human Cells. *Mol Cell* **24**: 593–602.

Younger ST, Rinn JL. 2017. p53 regulates enhancer accessibility and activity in
response to DNA damage. *Nucleic Acids Res* **45**: 9889–9900.

Zhu J, Dou Z, Sammons MA, Levine AJ, Berger SL. 2016. Lysine methylation represses
p53 activity in teratocarcinoma cancer cells. *Proc Natl Acad Sci U S A* **113**:
9822–9827.

Figure 1

Figure 1. The p53 cistrome and transcriptome are cell type-dependent *(A)* Western blot of cell lysates from MCF10A or SkFib cells at 6 hours after DMSO or Nutlin-3A treatment. *(B)* Venn diagram indicating differentially upregulated genes (Nutlin-3a/DMSO, fold change >2, adjusted p<0.05) in MCF10A or SkFib cells. *(C)* Immunoblotting for p53 expression at 6 hours of DMSO (D) or Nutlin-3A (N) treatment in SkFib cells stably expressing shRNA against p53 (p53 sh) or a non-targeting control shRNA (scr). *(D)* qRT-PCR analysis of SkFib-specific gene targets, GDNF and TRIM55, in SkFib cells stably expressing an shRNA to p53 (p53 sh) or a non-targeting control shRNA (scr). Gene expression is normalized to GAPDH expression. Error bars represent SEM; ****p<0.0001, ***p<0.001 and **p<0.01, calculated by Student's t-test. *(E)* Immunoblotting for p53 expression at 6 hours of DMSO (D) or Nutlin-3A (N) treatment in response to p53 knockdown in MCF10A cells stably expressing shRNA against p53 or a non-targeting control shRNA (scr). *(F)* qRT-PCR analysis of MCF10A-specific Nutlin-3A-induced genes, RIC3 and IL1B, in MCF10A cells stably expressing an shRNA to p53 (p53 sh) or a non-targeting control shRNA (scr). Expression is normalized to GAPDH expression. Error bars represent SEM; ****p<0.0001, calculated by Student's t-test. *(G)* Venn diagram depicting overlap between significantly-enriched (p<0.01, MACS v2) Nutlin-3A-induced p53 peaks in MCF10A or SkFib.

Figure 2

Figure 2. *Cis*-regulatory activity and differential p53 binding drives cell type-specific target gene expression. *(A)* Representative UCSC Genome Browser track view of the GDNF locus, a SkFib-specific p53 target. A p53-bound, putative enhancer (H3K-27ac+, H3K4me2+, H3K4me3-) is bound by a dashed box, while the putative GDNF promoter (H3K27ac/H3K4me2/H3K4me3+) is bound by a solid box. The y-axis is scaled to the maximum intensity between MCF10A or SkFib for each feature. *(B)* Representative UCSC Genome Browser track view at the RIC3 locus, illustrating p53 binding to an MCF10A-specific enhancer signature (H3K27ac+, H3K4me2+, H3K4me3-) in response to Nutlin-3A treatment. The y-axis is scaled to the maximum intensity between MCF10A or SkFib for each feature. *(C)* Heatmap plots at MCF10A-specific p53 binding sites for p53, H3K27ac, and H3K4me2 enrichment in MCF10A or SkFib within a 2000-bp window (-/+ 1000 bp from the p53 motif; '-1' indicates 1000bp downstream from the motif, '0' indicates the motif center, and '+1' indicates 1000bp upstream from the motif).

Figure 3

Figure 3. p63 engages with p53-bound enhancers and regulates p53-dependent trans-activation of epithelial target genes. *(A)* Representative UCSC Genome Browser track view of the IL1B locus, illustrating three MCF10A-specific putative enhancers bound by p53 and p63 in response to Nutlin-3A treatment (H3K27ac+, H3K4me2+, H3K4me3-; dashed box). The y-axis is scaled to the maximum intensity for each data set. *(B)* Venn diagram representation of the overlap between enriched p53 and p63 peaks (input-normalized, p53/p63 motif-positive, MACS v2, p<0.01) in MCF10A when treated with Nutlin-3A. *(C)* Heatmap plots of p53 and p63 enrichment at p53 only, shared, and p63 only binding sites in MCF10A cells within a 2000-bp window (-/+ 1000 bp from the p53 motif) in response to Nutlin-3A treatment. *(D)* Immunoblotting for p63 in response to p63 knockdown in MCF10A cells stably expressing shRNA to p63 or a non-targeting control shRNA (scr) after 6 hours of DMSO (D) or Nutlin-3A (N) treatment. *(E)* Fold-change (Nutlin-3A/DMSO, log2, median in black) of MCF10A-specific Nutlin-3A-induced genes in MCF10A cells expressing shRNA targeting either a non-targeting control (scr), p53, or p63. ****p<0.0001 and ***p<0.001 *(F)* qRT-PCR analysis of MCF10A-specific Nutlin-3A-induced genes, RIC3, IL1A and IL1B. Expression is normalized to GAPDH expression. Error bars represent SEM; ****p<0.0001 and **p<0.01, calculated by Student's t-test.

Figure 4

Figure 4: p63, and not p53, maintains enhancer chromatin identity. Input-subtracted *(A)* H327ac or *(B)* H3K4me2 enrichment at p63 binding sites (left, -/+ 250bp from p63 peak center) or at all remaining enhancers (H3K27ac+, H3K4me2+, H3K4me3-, right) in MCF10A cells expressing non-targeting (Scr), p53, or p63 shRNA. *(C)* Representative UCSC Genome Browser track view of the *RRM1* locus, illustrating three MCF10A-specific putative enhancers bound by p63 that are lost in response to p63 depletion (H3K27ac+, H3K4me2+, H3K4me3-; dashed box). The y-axis is scaled to the maximum intensity for each data set. *(D)* Bar graphs depicting the percent of p63 peaks that show 2-fold gains or losses of H3K27ac (left) or H3K4me2 (right) in response to either p53 or p63 depletion relative to non-targeting control shRNA. *(E)* Heatmap depicting the number of p63 binding sites that lose H3K4me2 (left) or remain constant (right) after p63 depletion that are found within regions of DNAse hypersensitivity (DHS) across multiple cell lines. *(F)* Heatmap illustrating the –log(10) p-value of motif enrichment found at p63 binding sites that lose H3K4me2 (left) or remain constant (right) after p63 depletion. *(G)* H3K27ac enrichment at p53 binding sites in MCF10A cells expressing the represented shRNA molecules. *(H)* Bar graph displaying the number of p53 binding sites with more than 2-fold change in H3K27ac enrichment after Nutlin-3A treatment of MCF10 cells expressing the indicated shRNA. *(I)* H3K4me2 enrichment at p53 binding sites (Verfaillie et al. 2016) in HCT116 *TP53+/+* or -/- cells in response to DMSO or Nutlin-3A treatment. *(J)* H3K4me1 and H3K4me2 enrichment at p53 binding sites (Kenzelmann Broz et al. 2013) in *Trp53+/+* or *Trp53-/-* mouse embryonic fibroblasts.

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure S1. Cell type-dependent p53 transcription in MCF10A and SkFib.
(A) Experimental design for RNA-sequencing analysis in response to p53 activation by DMSO or 5µM Nutlin-3A for 6 hours. (B-D) qRT-PCR validation of two representative *(B)* common, *(C)* MCF10A, and *(D)* SkFib-specific Nutlin-3A-induced targets. Target gene expression is normalized to GAPDH and data represent technical replicates from three independent biological replicates. Error bars represent SEM with p values calculated by Student's t-test, ****p<0.0001. *(E)* Venn diagram indicating differentially upregulated genes (Nutlin-3a/DMSO, fold change >2, adjusted p<0.05) in MCF10A or SkFib cells. *(F)* Gene ontology analysis of differentially expressed genes from MCF10A and SkFib cells showing upregulated genes for top three significant biological processes. *(G)* Intersection between SkFib (top) or MCF10A (bottom) p53 targets identified in this work compared to the meta-analysis of p53 targets identified in at least 3 independent experiments (Fischer, M 2017). *(H)* qRT-PCR analysis of p53 expression at 6 hours of DMSO (D) or Nutlin-3A (N) treatment in SkFib cells stably expressing shRNA against p53 (p53 sh) or a non-targeting control shRNA (scr). p53 expression is normalized to GAPDH expression (Relative expression). Error bars represent SEM; ****p<0.0001 and ***p<0.001, calculated by Student's t-test. *(I)* qRT-PCR analysis of p53 expression at 6 hours of DMSO (D) or Nutlin-3A (N) treatment in response to p53 knockdown in MCF10A cells stably expressing shRNA against p53 or a non-targeting control shRNA (scr). p53 RNA expression is normalized to GAPDH expression. Error bars represent SEM; ****p<0.0001, calculated by Student's t-test. *(J)* Heatmap of input-subtracted p53 enrichment for MCF10A (blue, left) or SkFib (green, right) within a 2000-bp window (-/+ 1000 bp from the p53 motif; '-1' indicates 1000bp downstream from the motif, '0' indicates the motif center, and '+1' indicates 1000bp upstream from the motif). *(K)* Fold change ratio of Nutlin-3A/DMSO of common input-subtracted p53 enrichment for MCF10A or SkFib. *(L)* The percentage of p53 peaks common in MCF10A and SkFib (left), or only in MCF10A (middle), or only in SkFib (right) in response to Nutlin-3A treatment within varying distances to the nearest transcription start site (TSS) of a Nutlin-3A-induced gene. *(M)* Boxplot analysis of the input-subtracted p53 enrichment for MCF10A (blue) or SkFib (green) cells for common-(left), or MCF10A-specific-(middle), or SkFib-specific-p53 binding sites (right) in response to DMSO (D) or Nutlin-3A (N) treatment.

Supplemental Figure S2. Epigenomic analysis of cell type-dependent p53 activation. *(A)* Representative UCSC Genome Browser track view of TRIM55 and MAMDC2 loci, two fibroblast-specific p53 targets. p53-bound, putative enhancers (H3K27ac+, H3K4me2+, H3K4me3-) illustrated by dashed box for TRIM55 and MAMDC2, while the putative MAMDC2 promoter (H3K27ac/H3K4me2/H3K4me3+) is bound by a solid box. The y-axis is scaled to the maximum intensity between MCF10A or SkFib for each feature. *(B)* Representative UCSC Genome Browser track view at the MYBPHL locus, demonstrating an MCF10A-specific enhancer signature (H3K27ac+,H3K4me2+, H3K4me3-) at a p53 peak in response to Nutlin-3A treatment. The y-axis is scaled to the maximum intensity between MCF10A or SkFib for each feature. *(C)* The percentage of MCF10A p53 binding sites overlapping H3K4me2, H3K4me3, or H3K27ac under basal (DMSO-treated) conditions. (D-F) The percentage of intersecting H3K27ac peaks (input-normalized, MACS v2, $p<0.01$) with p53 peaks (input-normalized, MACS v2, $p<0.01$) that are *(D)* common to MCF10A and SkFib, *(E)* MCF10A-specific, or *(F)* SkFib-specific in response to DMSO (D) or Nutlin-3A (N) treatment. Adjacent boxplots depict H3K27ac enrichment over a 500bp window from the p53-binding site. *(G)* The percentage of intersecting H3K4me3 peaks (input-normalized, MACS v2, $p<0.01$) with p53 peaks (input-normalized, MACS v2, $p<0.01$) that are common to MCF10A and SkFib (top), MCF10A-specific (middle), or SkFib-specific (bottom) in response to DMSO (D) or Nutlin-3A (N) treatment. *(H)* Homer-derived transcription factor motif enrichment found in MCF10A (top) or SkFib (bottom) specific enhancers

Supplemental Figure S3. Epithelial-specific p53 binding sites correlate with p63 binding. *(A)* qRT-PCR analysis of p53, p63 or p73 of cell lysates from MCF10A, SkFib, HUVEC and HaCaT cells at 6 hours after DMSO (D) or Nutlin-3A (N) treatment. Expressions detected by qRT-PCR are normalized to GAPDH expression. HaCaT cells used as positive control for p63 and p73 expression. Error bars represent SEM, p values were calculated by Student's t-test, ****$p<0.0001$. *(B)* Representative UCSC Genome Browser track view of the RNASE7 locus, illustrating a MCF10A-specific putative enhancer bound by p53 and p63 in response to Nutlin-3A treatment (H3K27Ac+, H3K4me2+, H3K4me3-; dashed box). The y-axis is scaled to the maximum intensity for each data set. *(C)* The percentage of intersecting H3K27ac peaks (input-normalized, MACS v2, $p<0.01$) with p63 peaks observed in MCF10A cells (input-normalized, MACS v2, $p<0.01$) in response to DMSO (D) or Nutlin-3A (N) treatment. *(D)* The percentage of p63 binding sites observed in MCF10A cells at varying distances to the nearest TSS of all RefSeq genes (white) or Nutlin-3A induced genes (red). *(E)* qRT-PCR (right) analysis of p53 and p63 in response to p63 knockdown in MCF10A cells stably expressing shRNA to p63 or a non-targeting control shRNA (scr) after 6 hours of DMSO (D) or Nutlin-3A (N) treatment. Target gene expression is normalized to GAPDH for qRT-PCR analysis. Error bars represent SEM, p values were calculated by Student's t-test, ****$p<0.0001$, ***$p<0.001$ and **$p<0.01$. *(F)* K-means clustering of Nutlin-3A/DMSO fold-change values of previously identified MCF10A-specific p53 targets in MCF10A cells expressing either control (scr), p53, or p63-targeted shRNA. *(G)* The number and percent of p53 target genes showing downregulated, upregulated, or unchanged basal mRNA expression in MCF10A-p63KD cell lines.

Supplemental Figure S4. p63 is an enhancer maintenance factor.
(A) Representative UCSC Genome Browser track view of the EDN2 locus, illustrating three MCF10A-specific putative enhancers bound by p63 that are lost in response to p63 depletion (H3K27ac+, H3K4me2+, H3K4me3-; three separate dashed box). The y-axis is scaled to the maximum intensity for each data set. *(B)* The number of p63-sensitive, H3K4me2-marked enhancers (out of 1496 total) overlapping DNAse hypersensitive sites (DHS) across epithelial and non-epithelial cell types analyzed by the ENCODE project. Error bars represent the median and 95% confidence interval. *(C)* The percentage of all MCF10A p63 binding sites intersecting ChromHMM-derived "Enhancer" chromatin states in epithelial and non-epithelial cell types. *(D)* The distance (base pairs, log2) between p63-sensitive (left) or p63-insensitive (right) H3K4me2-marked enhancers and genes upregulated or downregulated in response to p63 depletion in MCF10A cells. *(E)* H3K4me2 enrichment (Input-subtracted H3K4me2, -/+ 250bp from p53 motif center) at p53 binding sites in MCF10A cells expressing control (scr), p53, or p63-targeted shRNA in response to DMSO or Nutlin-3A treatment. *(F)* Enrichment of H3K4me2 (left) or H3K27ac (right) at MCF10A p53+/p63- binding sites in in MCF10A cells expressing control (scr), p53, or p63-targeted shRNA in response to DMSO treatment. H3K4me2 and H3K27ac enrichment are reported as input-subtracted histone modification within a -/+ 250bp window from the p53 motif center.

Supplemental Materials for Karsli Uzunbas et al.
Methods

RNA-sequencing sample and library preparation

Total mRNA was extracted with E.Z.N.A. Total RNA kit (Omega; R6834-02) and Poly(A)+ RNA was isolated by double selection with poly-dT beads, using 2.5 μg total RNA, which is then followed by first- and second-strand synthesis. Sequencing libraries were prepared using NEXTflex Rapid Illumina DNA-Seq Library Prep Kit (Bioo Scientific). Samples were single-end sequenced on an NextSeq 500. RNA-seq reads were aligned via STAR (Dobin et al. 2013) to Ensembl v.75/hg19. Count tables were generated using STAR-aligned BAM files and HTseq (Anders et al. 2015). Count tables were then used to call differentially expressed genes using DEseq2 (Love et al. 2014).

Quantitative Real-Time PCR

Total RNA was extracted as per RNA-sequencing protocol and cDNA was synthesized using 1ug of total RNA as a template and the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems; 413760). Relative standard qPCR was performed using iTaq Universal SYBR Green Supermix (Bio-Rad; 172-5124) with primers shown in Supplemental Table S8 on an ABI 7900HT instrument.

Lentivirus production, purification and transduction

Lentiviral shRNAs were produced using HEK293T cells that were seeded in 6 well plates. shRNA sequences are as follows: p53 (GTCCAGATGAAGCTCCCAGAA), p63 (CCGTTTCGTCAGAACACACAT). 1 μg of pLKO plasmid having either scramble shRNA or p53 shRNA or p63 shRNA was combined with 1 μg mix of packaging plasmids (pMD2 and psPAX2) and the mixture was diluted into jetPRIME buffer (Polyplus Transfection; 89129-924) and reagents, following the manufacturer's protocol. Lentivirus containing supernatants were collected at 24 hours and 48 hours post-transfection, and filtered through a 0.45 μm membrane and stored in aliquots at -80°C. MCF10A or SkFib cells were transfected with lentivirus supplemented with 8 μg/ml

polybrene. At 24 hour post-infection with lentivirus, media was replaced with the proper puromycin selection (0.5 μg/ml for MCF10A and 2 μg/ml for SkFib).

ChIP-sequencing sample and library preparation

Cells were crosslinked at 80% confluency with 1% formaldehyde for 10 min at room temperature. Crosslinking was quenched with 125mM glycine and the resulting pellet was washed twice with cold PBS and lysed as previously described (Lee et al. 2006). Samples were subjected to sonication with Diagenode Bioruptor Plus for 40 cycles (30 sec on/off at high setting) for shearing chromatin to 150-500 bp average size.

Immunoprecipitation reactions for SkFib, MCF10A, and HCT116 cell lines were performed with Diagenode IP-Star Compact Automated System, with the exception of p53 ChIP experiments. Mouse embryonic fibroblast and p53 ChIP-seq were performed as described in Sammons et al 2015 (Sammons et al. 2015). Antibodies were preconjugated to Protein G beads (Invitrogen) against p63 (Cell Signaling; 13102), H3K4me1 (Abcam, ab8895), H3K4me3 (Active Motif; 39159), H3K4me2 (Millipore; 07-030), H3K27ac (Active Motif, 39133), p53 monoclonal antibody DO1 (BD Biosciences; BD554293). Immunoprecipitated DNA was reverse-crosslinked at 65°C for 4 hours, eluted, purified using SPRI beads, and used to construct sequencing libraries.

Sequencing libraries were prepared using NEBNext Ultra DNA Library Prep Kit for Illumina (New England Biolabs). Prior to sequencing, library quality control was performed with Qubit (Thermo Fisher Scientific), Bioanalyzer (Agilent) and qPCR quantification. All ChIP samples, including input, were single-end sequenced on an NextSeq 500 at the University at Albany Center for Functional Genomics. Uniquely aligned reads (up to one mismatch) were aligned to NCBI37/hg19 using Bowtie2 (Langmead and Salzberg 2012).

ChIP-seq Peak Calling

Significant regions of transcription factor (p53, p63) and histone modification (H3K4me2, H3K4me3, H3K27ac) enrichment were called using macs v.2.1.0 (Zhang et al. 2008) preserving only peaks with an adjusted P-value < 0.01. All peak calling was performed with sheared chromatin input. p53 and p63 motif analysis was performed using p53scan (Smeenk et al. 2008) and all macs-derived peaks lacking a p53/p63

consensus motif were removed from further analysis. A full list of p53/p63 peaks can be found in Supplemental Table S9. Global transcription factor motif enrichment experiments were performed using the findMotifsGenome.pl script from HOMER (Heinz et al. 2010).

Analysis of chromatin enrichment
Chromatin enrichment analyses were performed using the annotatePeaks.pl script of HOMER. Read-depth-scaled chromatin tag enrichment within a specified window was input-normalized (target – input) and then further scaled to total genomewide enrichment of peaks for that histone modification.

Computational Analysis and Plotting
Peak file intersections were performed using the intersectBed package of BedTools (Quinlan and Hall 2010), scoring positive intersections as those with at least 1bp of overlap (-f 1E-9). The original peak file (-a) was reported only once (-u) if any intersection exists with the query file (-b). The closestBed package of BedTools was used to measure distances between features with the distance reported using the –d option. Figures were generated using Graphpad Prism or Datagraph. Bigwig (bw) files were generated with the bamCoverage package of deepTools v.2.5.4 (Ramírez et al. 2016) with a bin size of 1 (--binSize 1) and read extension (--extendReads 300). Heatmaps were then generated using bigwig files for a -/+ 1000bp region from the p53/p63 peak center using the computeMatrix (reference-point, -a 1000, -b 1000, --binSize 10) and plotHeatmap functions. Genome browser tracks (bedGraph) were generated using the makeUCSCfile.pl package of HOMER (Heinz et al. 2010). K-means clustering was performed using cluster 3.0 for Mac.

Statistical Testing
Statistical testing was performed using Graphpad Prism (v. 7) or using built-in statistical parameters in *R* (v.3.4.0).

ChromHMM chromatin state and DNAse hypersensitive site (DHS) enrichment analysis

ChromHMM analysis (Ernst and Kellis 2012) was performed using the 25 state model (http://egg2.wustl.edu/roadmap/web_portal/), with at least 50% of the p53/p63 motif required to overlap a single chromatin enrichment term (closestBed –a p53.file –b ChromHMM.file –d –f 0.51). The 25 chromatin state models for ChromHMM terms were combined into 4 regulatory terms: enhancer, repressive, quiescent, or transcription. A full list of p53, p63, and p53/p63 binding site regulatory region inferences across the 127 cell types of the ChromHMM analysis can be found in Supplemental Table 10. DHS data were downloaded from http://hgdownload.cse.ucsc.edu/goldenpath/hg19/encodeDCC/wgEncodeRegDnaseClustered/. Intersection of p53/p63 peaks with DHS data was performed using the closetBed package of BedTools, with 100% of the p53/p63 motif required to overlap the DHS (-f 1).

Supplemental References

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinforma Oxf Engl* **31**: 166–169.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinforma Oxf Engl* **29**: 15–21.

Ernst J, Kellis M. 2012. ChromHMM: automating chromatin-state discovery and characterization. *Nat Methods* **9**: 215–216.

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK. 2010. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**: 576–589.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357–359.

Lee TI, Johnstone SE, Young RA. 2006. Chromatin immunoprecipitation and microarray-based analysis of protein location. *Nat Protoc* **1**: 729–748.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**: 841–842.

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**: W160–W165.

Sammons MA, Zhu J, Drake AM, Berger SL. 2015. TP53 engagement with the genome occurs in distinct local chromatin environments via pioneer factor activity. *Genome Res* **25**: 179–188.

Smeenk L, van Heeringen SJ, Koeppel M, van Driel MA, Bartels SJ, Akkers RC, Denissov S, Stunnenberg HG, Lohrum M. 2008. Characterization of genome-wide p53-binding sites upon stress response. *Nucleic Acids Res* **36**: 3639–54.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, et al. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**: R137.