Influenza A virus environmental persistence is driven by the hemagglutinin and the neuraminidase

Corresponding author: India Leclercq & Thomas Labadie

1- Thomas Labadie, thomas.labadie@pasteur.fr,[a,b]
2- Christophe Batéjat, christophe.batejat@pasteur.fr, [a]
3- Jean-Claude Manuguerra, jean-claude.manuguerra@pasteur.fr,[a, *]
4- India Leclercq, india.leclercq@pasteur.fr,[a,b, *]

[a] Institut Pasteur, Unité Environnement et Risques Infectieux, Cellule d'Intervention Biologique d'Urgence (CIBU), 28 rue du Dr Roux, Paris, France

[b] Université Paris Diderot, Sorbonne Paris Cité (Cellule Pasteur), rue du Dr Roux 75015 Paris, France

* co-last authors

Abstract

The transmission routes of Influenza A viruses (IAVs) submit virus particles to a wide range of environmental conditions that affect their transmission. In water, temperature, salinity and pH are important factors modulating viral persistence in a strain-dependant manner, and the viral factors driving IAV persistence remained to be described. We used an innovative method based on a real-time cell system analysis to quantify viral decay in an environmental model. Thus, we identified the viral hemagglutinin (HA) and neuraminidase (NA) as the main proteins driving the environmental persistence by comparing the inactivation slopes of several reassortant viruses. We also introduced synonymous and non-synonymous mutations in the HA or in the NA that modulated IAV persistence. Our results demonstrate that HA stability and expression level, as well as calcium-binding sites of the NA protein are molecular determinants of viral persistence.

27 Finally, IAV particles could not trigger membrane fusion after environmental exposure,

28 stressing the importance of the HA and the NA for environmental persistence.

Introduction

Influenza A viruses (IAVs) has a wide host range, which allows them to spread almost

everywhere on the planet. In nature, the H1N1 subtype infects several hosts such as

domestic and aquatic birds, humans, swine or dogs and thus spreads through a wide

range of environmental conditions. In humans, a pandemic H1N1 virus emerged in 2009

after the reassortment between a swine H1N1 virus from the Eurasian lineage and a swine

H2N1 virus descended from a triple reassortment between a swine H1N1 virus from the

North American lineage, an avian virus and the human H3N2 virus. In aquatic birds, which

are the main reservoir of these viruses, IAVs spread mainly by a fecal-oral route through

water. In poultry such as chicken and turkeys, or in mammalian species, IAVs mainly have

a respiratory tropism, and the virus spreads by contact between infected and susceptible

hosts or by contaminated fomites, as well as through aerosol or respiratory droplets. In

any case, the transmission routes of IAVs submit virus particles to a wide range of

environmental conditions, which more or less rapidly affect them. In water, the time for

IAV inactivation depends on widely studied abiotic factors such as temperature (Brown

et al. 2009, 2007; Dublineau et al. 2011; Keeler et al. 2014; Nazir et al. 2010; Poulson et

al. 2016; D.E Stallknecht and Brown 2009; H. Zhang et al. 2014), water salinity (Brown et

al. 2009; Dublineau et al. 2011; Keeler et al. 2014; Poulson et al. 2016) and pH (Brown et

al. 2009; Keeler et al. 2014; Poulson et al. 2016) and can range from a few days in saline

water (35 g.L^{-1} NaCl) at 35°C to several years at 4°C (Dublineau et al. 2011). The

environment acts both as a reservoir (Roche et al. 2014; David E. Stallknecht et al. 2010)

and as a bottleneck for IAVs evolution. It was experimentally demonstrated that IAV

persistence in water is extremely variable among strains and subtypes (Dublineau et al.

2011; Lebarbenchon et al. 2012; D.E Stallknecht and Brown 2009). However, inactivation

processes of avian IAVs in the environment are far from being well understood

(Lebarbenchon et al. 2010). Only a few viral factors, such as the viral envelope origin and

composition (Shigematsu et al. 2014; Bajimaya et al. 2017) and the pH of activation of the

HA (Reed et al. 2010) are known to affect virus survival. On the other hand, the viral

envelope and the viral genome are affected much more slowly than the actual loss of

infectivity in water, suggesting that other viral factors drive IAV survival (Shigematsu et al.

2014; Dublineau et al. 2011). The envelope of IAV structure is made of two external

glycoproteins, the hemagglutinin (HA) and the neuraminidase (NA), and a proton channel

M2, which are inserted into a lipid bilayer surrounding a matrix M1 protein layer. Inside

viral particles, the segmented and negative strand RNA genome surrounded by a

nucleocapsid (NP) interacts with the viral polymerase made of the three proteins PB1,

PB2 and PA to constitute the ribonucleoprotein (RNP) complex. The 8 RNPs interact

closely with the M1 proteins.

In order to identify viral genetic drivers of the persistence phenotype, we generated

reassortants from H1N1 viruses, which do not have the same persistence in an

environmental model, and compared their inactivation slopes. As a model, we used saline

(35 g.L^{-1} NaCl) water at 35°C, because it is the average salt concentration in the ocean

and this temperature allows observing differences of viral persistence more rapidly. Using

fluorescence imaging microscopy, we also wanted to understand how our environmental

model disrupts viral functions and infectivity of IAVs. We identified molecular determinants

of the persistence phenotype in the environment, by introducing codon-optimized

synonymous mutations or non-synonymous mutations in the HA or the NA gene.

Altogether, our results demonstrate that the survival of Influenza viral particles is

predominantly driven by the two external glycoproteins, and that environmental conditions

affect the HA mediated steps during viral entry.

Results

Influence of viral proteins on H1N1 strains persistence outside the host

Influenza A viruses, such as the seasonal H1N1 virus of 1999 and the pandemic H1N1

virus of 2009, show different persistence in water (Dublineau et al. 2011; Lebarbenchon

et al. 2012; D.E Stallknecht and Brown 2009). This study aimed at characterizing viral

molecular drivers associated with variations of persistence in saline water (35 g.L^{-1} NaCl)

at 35°C used as an environmental model. These temperature and salinity conditions were

selected because they allow observing variations between strains in a short period of time

(Dublineau et al. 2011), and 35 g.L^{-1} NaCl is the average salt concentration in the oceans

(Millero et al. 2008). In order to understand how viral proteins influence the persistence of

H1N1 influenza particles, reassortant viruses between a pandemic H1N1 virus, the

A/Bretagne/7608/2009 strain (whole/2009) and a pre-pandemic H1N1 virus, the

A/WSN/1933 strain (whole/1933) were generated using a plasmid-based reverse genetics

system. We rescued reassortant viruses containing either the HA/NA segments, the

PB1/PB2/PA segments, the M segment or the NS/NP segments from

A/Bretagne/7608/2009 strain with the other segments from A/WSN/1933 as a

complementary genomic backbone. We conversely rescued the same mirroring set of

viruses with the same pool of segments but from the A/WSN/1933 strain and with the

genomic backbone constituted by A/Bretagne/7608/2009 virus segments. The

composition of each virus is detailed in tables 1 and 2. All reassortants grew except the

NS-NP/1933 virus. Thus, we generated seven different reassortant viruses and compared

their persistence with a real-time cell analysis system (RTCA), as described in the

Methods section (Fig. 1A). To quantify viral decay, we monitored CIT_{50} values after 1, 24

and 48 hours of exposure in saline water at 35°C. The increase of CIT_{50} value over time

reflects the loss of infectivity and the progressive inactivation of viral particles (Fig. 1B).

We calculated inactivation slopes from experimental CIT_{50} values (Fig. 1C). The mean

inactivation slope of the whole/2009 virus in saline water at 35°C was 4.4 $CIT_{50}.day^{-1}$ (Fig.

1D), which is twice more stable than the whole/1933 virus that has a mean inactivation

slope of 8.3 $CIT_{50}.day^{-1}$ (Fig. 1E). When compared with the whole/2009 virus (Fig. 1D),

replacing the 2009 HA and NA by the 1933 HA and NA (HA-NA/1933 virus), or the 2009

M segment by a 1933 M segment (M/1933 virus) significantly destabilized the virus with

mean inactivation slopes of 6.9 and 11.3 $CIT_{50}.day^{-1}$ respectively. On the contrary, the

Pol/1933 virus persistence was not significantly different from that of the whole/2009 virus,

with a mean inactivation slope of 3.3 $CIT_{50}.day^{-1}$. Compared with the whole/1933 virus

(Fig. 1E), replacement of the 1933 HA and NA by the 2009 HA and NA (HA-NA/2009

virus) significantly increased the persistence of the virus with a mean inactivation slope of

6.7 $CIT_{50}.day^{-1}$. The replacement of the 1933 M or NS and NP segments by their 2009

counterparts (M/2009 and NS-NP/2009 viruses) did not change significantly their

persistence. More surprisingly, replacing the polymerase PB1, PB2 and PA segments of

1933 by the polymerase segments of 2009 (Pol/2009 virus) significantly destabilized this

reassortant virus with a mean inactivation slope of 12.4 $CIT_{50}.day^{-1}$.

Thus, once the HA and NA were isolated from their genomic context into a new virus, the

persistence phenotype of this new virus tended to be the same as the virus from which

HA and NA were originated. The matrix protein and to a certain extent the viral polymerase

only induced a decrease of viral persistence without reflecting the phenotype of their

original virus. Altogether, these results suggested that HA and NA are the most important

viral proteins driving the persistence phenotype.

Molecular determinants of the HA driving environmental persistence

In order to understand how the HA protein modulates environmental persistence, we

rescued two reassortant viruses bearing a HA and NA either from the H1N1

A/NewCaledonia/20/1999 strain (HA-NA/1999) or from the pandemic A/Paris/2590/2009

strain (HA-NA/2009) (Table 2), both with the same genomic backbone belonging to the

A/WSN/1933 strain. We selected these HA and NA because the two strains have a

different persistence phenotype in saline water at 35°C (Dublineau et al. 2011) in favour

of the pandemic strain. Using a third strain also allowed us to compare the HA and NA of

the 2009 pandemic strain and the pre-pandemic strain independently of their parental

genomic context. We wanted to assess whether introducing changes in the HA sequence

had an impact on viral persistence. We thus synthesized a codon-optimized gene

encoding the 1999 HA (HA_{opti}), producing the same HA but containing 24.4% of

synonymous substitutions. We then generated the corresponding reassortant virus

bearing this HA_{opti} and the 1999 NA in the A/WSN/1933 backbone (HA_{opti}-NA/2009, Table

2). The slope of inactivation was significantly lower in saline water at 35°C for the

HA-NA/2009 virus compared with the HA-NA/1999 virus inactivation slope (Fig. 2A),

respectively 5.9 and 9.9 $CIT_{50}.day^{-1}$. This result is in agreement with that of a previous

study (Dublineau et al. 2011) on the persistence phenotype of the wild-type

non-reassortant A/Paris/2590/2009 and the A/NewCaledonia/20/1999 strains, and

confirms that HA and NA are indeed the main viral factors leading the persistence of viral

particles. Interestingly, HA_{opti}-NA/1999 was as stable as HA-NA/2009 and thus

significantly more stable than HA-NA/1999 (Fig. 2A) despite the identical amino acid

composition of HA.

We also introduced non-synonymous mutations in the HA nucleotide sequence to study

the impact of a single HA amino-acid change on virus persistence. For this purpose, we

compared the HA amino-acid sequences of the HA-NA/1999 and HA-NA/2009 viruses

and, based on a preliminary mutation screening performed using a lentiviral

pseudo-particles system (Sawoo et al. 2014), we selected five residues in the HA protein

that might influence virus particle persistence outside the host (Fig. 2B). Ten reassortant

viruses bearing HA with amino-acid substitutions or insertions/deletions were generated

after site-directed mutagenesis of the pPol-HA plasmid. We then exposed these mutated

HA-NA/2009 and HA_{opti}-NA/1999 viruses to saline water at 35°C (Fig. 2C and 2D). The

HA/Y454F substitutions or HAΔK147 deletion in the HA of HA-NA/2009 virus induced a

significant increase of the mean inactivation slope, respectively to 9.2 and 10.3

$CIT_{50}.day^{-1}$, thus generating very unstable mutants. Substitutions HA/K300Q and

HA/K53S also destabilized the HA-NA/2009 virus but to a lesser extent, whereas the

HA/T327A substitution did not affect its persistence, with mean inactivation slopes of 9.2,

7.3 and 6.6 $CIT_{50}.day^{-1}$ respectively. The insertion of HA::K147 or substitutions A326T,

F453Y and Q299K in the HA of HA_{opti}-NA/1999 virus greatly decreased the persistence of

the virus, with a mean inactivation slope of 9.9, 8.0, 9.8 and 9.6 $CIT_{50}.day^{-1}$, respectively.

The S53K substitution however had less impact on the persistence, with a mean

inactivation slope of 7.6 $CIT_{50}.day^{-1}$. Altogether, these results demonstrate that

synonymous mutations or a single amino acid change in the HA is sufficient to negatively

affect the viral persistence outside the host.

Viral particles cannot trigger membrane fusion after being exposed to saline water

at 35°C

Since the loss of hemagglutination titre of viral particles is slower than their loss of

infectivity (Chu 1948) (Fig. 3A and Fig. S1), we decided to evaluate whether viruses were

still able to bind their cellular receptor. For this purpose, we immuno-labelled the viral

nucleoprotein (NP), which encapsidates the viral genome to form the ribonucleoprotein,

and used confocal microscopy for its detection in infected MDCK cells. In cells infected

with non-exposed HA-NA/2009 or HA-NA/1999 viruses, the NP protein was concentrated

within the nucleus 2 hours after infection (Fig. 3B). On the contrary, in MDCK cells infected

with exposed HA-NA/2009 or HA-NA/1999 viruses for 5 days to saline water at 35°C, we

detected the NP protein close to the cell membrane but not in the nucleus after 2 hours of

infection. We confirmed that NP localization of exposed viruses was similar to that of

non-exposed viruses after 20 min of infection, when virus entry is not completely achieved

(Fig. S2). We thus assessed the HA-triggered fusion of the viral membrane in infected

MDCK cells using R18-labelled viruses at self-quenched concentration, a widely used

technique to detect fusion of the viral membrane with the endosomal membrane (Pohl,

Edinger, and Stertz 2014; Schelker et al. 2016). Fusion events were detected 20 min after

infection by confocal microscopy in cells infected with viruses exposed 5 days to saline

water at 35°C or non-exposed viruses (Fig. 3C). We detected almost no R18 dequenching

signal in cells infected with HA-NA/2009 or HA-NA/1999 exposed viruses, whereas we

observed numerous fusion events in cells infected with the non-exposed viruses. This

result demonstrates that the loss of infectivity of H1N1 viruses in saline water at 35°C is a

consequence of the HA inability to trigger membrane fusion.

HA stability at low pH is not the only determinant of viral persistence in water

A link between HA stability at low pH and stability to increasing temperatures has already

been proposed (Imai et al. 2012; Krenn et al. 2011). Therefore, we measured the

persistence of HA-NA/2009, HA-NA/1999 and mutated viruses in various pH-adjusted

PBS solutions for 1 hour at room temperature (Fig. 4A). In addition, we quantified HA cell

surface expression by flow cytometry in MDCK cells at 24 hours post infection (Fig. 4B).

The most unstable viruses (HA/Y454F and HAΔK147 viruses; Fig. 2C) showed a lower

HA surface expression level (Fig. 4B) and had a higher sensitivity at low pH compared

with HA-NA/2009 virus. For the stable HA$_{opti}$-NA/1999 virus, we observed a higher HA

surface expression level compared with the unstable HA-NA/1999 virus, but they both had

a high sensitivity at low pH. Similarly, the whole/1933 virus, which was more stable than

the Pol/2009 virus (Fig. 1E) even if they both have the same HA and NA, had much higher

HA expression levels in infected cells (Fig. 4C). Altogether, our results support the idea

that HA expression level is a viral driver of environmental persistence when the HA has a

high pH of inactivation.

Molecular determinants of the NA driving environmental persistence

Based on the results obtained with our reassortant viruses, the NA might be as important as the HA in driving the persistence phenotype (Fig. 1B). Previous studies have shown that the NA stability is strain dependent, and is linked to the calcium binding by the protein (Baker and Gandhi 1976; Burmeister, Cusack, and Ruigrok 1994). The 2009 pandemic NA has three calcium binding sites (Air 2012), as well as the 1918 NA (Xu et al. 2008), and thus presumably the 1999 NA. After alignment of the 1999 and 2009 NA protein sequences (Fig. 5A), we observed differences in the amino acids directly involved in the second and the third calcium binding sites (Air 2012), respectively near the active site and the side chains (D381, D387, and D379) (Fig 5A). We then decided to substitute the amino acids around the positions DGAD/341/NGAN (substitutions D341N and D344N), and/or DTDSD/382/GDTNN (substitutions D382G, S385N and D386N) (Fig 5A). We generated reassortant viruses with a mutated NA either in position 341 (HA-NA $_{(341)}$/1999 virus), or in position 382 (HA-NA $_{(382)}$/1999 virus) or in both positions (HA-NA $_{(341/382)}$/1999 virus), all with a 1999 HA and a genomic backbone from the 1933 H1N1 virus. We determined the persistence of these mutant viruses in saline water at 35°C by measuring a mean inactivation slope of 10.2 $CIT_{50}.day^{-1}$ for the HA-NA $_{(341)}$/1999 virus, 5.6 $CIT_{50}.day^{-1}$ for the HA-NA $_{(382)}$/1999 virus and 4.3 $CIT_{50}.day^{-1}$ for the HA-NA $_{(341/382)}$/1999 virus. In comparison, the HA-NA/1999 virus had a mean inactivation slope of 9.9 $CIT_{50}.day^{-1}$ (Fig 5B). We thus observed a positive effect of mutations at the third calcium binding site towards long-lasting persistence for the HA-NA $_{(382)}$/1999 virus. A cumulative effect of mutations at both calcium binding sites was also observed, leading to a significant increase of HA-NA $_{(341/382)}$/1999 virus survival compared with HA-NA $_{(382)}$/1999 virus. The

results demonstrated that amino-acids at the calcium binding sites in the NA are important

molecular determinants of virus environmental persistence. Viruses with a high

environmental persistence also had a stable NA, as shown in Fig. 5C and Fig. 5D,

measuring the NA activity of viruses exposed or not to saline water at 35°C. Viruses with

an important environmental persistence also presented a high NA activity (Fig. 5E).

Unfortunately, the NA activity for the HA-NA $_{(341)}$/1999 virus was below our detection

threshold and could not be measured.

Discussion

Differences in environmental persistence among influenza virus strains of the same

subtype were previously described (D.E Stallknecht and Brown 2009; Dublineau et al.

2011; Lebarbenchon et al. 2012), without providing a clear molecular basis to this

phenomenon. It was also demonstrated that their viral genome was not degraded and that

the viral envelope remained intact after a few days in saline water at 35°C (Dublineau et

al. 2011; Shigematsu et al. 2014). The results produced in our controlled experimental

model with H1N1 viral strains are the first to describe molecular basis of the IAVs

persistence outside the host. Comparing the persistence of multiple engineered

reassortants allowed us to identify the HA and NA proteins as major viral drivers of the

virus survival outside the host (Fig. 1D and 1e and Fig. 2A).

Codon optimization of the HA gene increased considerably the viral persistence of the

HA$_{opti}$-NA/1999 virus compared with the non-optimized HA-NA/1999 virus (Fig. 2A) while

they both harboured an identical HA protein. Codon usage bias that favours synonymous

mutations of HA exists in nature, particularly at the more functionally constrained residues

(Plotkin and Dushoff 2003). It would be interesting to investigate whether the viral

codon-optimization in nature also confers adaption of the persistence phenotype to the

environment. In addition, we found amino-acid changes in HA that affected the

persistence of the whole virus, but without a reciprocal effect between the amino-acid

changes in the 1999 or the 2009 HA. The two mutations responsible for the greatest

decrease of viral persistence were located near the receptor binding site (position 147)

and in the internal stem (position 454 or 453) while the others were located in the external

loops of the HA1 subdomain (Fig. 3A). Some mutations affecting HA stability in low pH

conditions were previously described (Russier et al. 2016; Byrd-Leotis et al. 2015) and

few mutations were associated with a higher persistence outside the host (Reed et al.

2010) and to heat inactivation (Krenn et al. 2011). Similarly, we observed a link between

pH stability of our 2009 HA-mutated viruses (Fig. 5A) and their persistence in water.

Our results also demonstrated that a virus with a high sensitivity to low pH still has the

potential to be stable in the environment. Indeed, HA-NA/1999 and HA_{opti}-NA/1999 viruses

both displayed high sensitivity to low pH although they have a very different persistence

phenotype in water. We also observed a higher HA surface expression levels in cells

infected with HA_{opti}-NA/1999 virus (Fig. 4B), in agreement with a previous study using

codon optimized HA (32). Altogether, our results suggest that the survival likelihood for a

virus increases when its HA surface level is higher, warranting to examine further the

correlation between HA level at the cell surface and HA level at the virion surface. Results

obtained with the Pol/2009 and the whole/1933 reassortant viruses, for which we expected

different transcription and replication efficiencies of the HA segment, are in agreement

with this finding (Fig. 4C and Fig. 1E). Although HA-NA/2009 and HA-NA/1933 are

respectively more stable and less stable, compared with their parental whole/1933 and

whole/2009 viruses, they have the same inactivation slope (Fig. 1D and 1E). Different

replication efficiency of the polymerase complexes resulting in various HA expression

levels might explain this observation.

Similarly, the less stable HAΔK147 and HA/Y454F mutant viruses (Fig. 3B) have a high

sensitivity to low pH and lower HA expression level in infected cells compared to the other

mutants (Fig. 4A and 4B). On the other hand, the whole/2009 and the Pol/1933 viruses

had the same persistence but significant differences of HA expression levels (Fig. 4C).

They both harbour the same HA of 2009, which is stable at low pH. Overall, these results

suggest that the persistence of a virus with a HA resistant to low pH does not vary with

HA expression level.

After losing their infectivity, HA-NA/2009 and HA-NA/1999 viruses were still able to attach

to cells but could not trigger viral fusion after endocytic uptake. Three or four neighbouring

HA trimers are probably mandatory to reach the hemifusion step occurring during the

fusion process (33). This number might not be reached for viruses that do not have a

sufficient density of functional HA at their surface and could explain the observed absence

of fusion after exposure to a hostile environment. This observation might in turn provide

an explanation for how viruses with higher HA surface levels stay stable longer, as the HA

inactivation is probably progressive in the environment as suggested by our results on the

loss hemagglutination activity (Fig. 3A).

We provided evidences that the NA protein is also a driver of IAVs environmental

persistence. Our results highlighted the importance of amino acid positions in the calcium

binding sites for both the NA stability and virus survival. It was previously shown that the

NA from the 2009 and 1918 H1N1 pandemic viruses possess a third calcium-binding site

coordinated by side chains (Xu et al. 2008; Air 2012), suggesting that all the human N1

may contain three calcium-binding sites. Based on a sequence alignment between the NA

of the 2009 and 1999 H1N1 viruses and observations of the structural features, we

substituted key amino acids involved in the second and the third calcium-binding sites.

These mutations induced an important increase of the environmental persistence of both

the HA-NA $_{(382)}$/1999 and HA-NA $_{(341/382)}$/1999 viruses compared with the HA-NA/1999

virus (Fig. 5B). Our results raised new questions on this putative third calcium-binding site

in the human N1, which may not be present or possess a weaker affinity for calcium ions

in the NA of the pre-pandemic H1N1 of 1999. Moreover, we observed that the more stable

virus in the environment, such as HA$_{opti}$-NA/1999 virus and the double mutant HA-NA

$_{(341/382)}$/1999 virus, induced a high HA surface expression level or had a stable and high

NA activity (Fig. 4B and Fig. 5E). The mutations introduced in the HA or the NA probably

modified the viral HA/NA balance, which might be an important molecular determinant of

environmental persistence, possibly by increasing sialic acid binding, which was

suggested to stabilize the influenza virus (36, 37). These results raised new questions

about the role of the NA protein in the viral entry, which is not yet fully understood.

Based on previous published results, we found that a log-linear model drives influenza A

virus inactivation in our experimental model (Dublineau et al. 2011). We observed that

the distributions of the inactivation slopes among the different viruses were significantly

different (Bartlett's statistic, $P<0.05$), so we used a non-parametric

Wilcoxon-Mann-Whitney test to compare virus persistence. Moreover, we observed that

the standard deviation of the mean inactivation slopes tends to increase in the

experiments performed with low persistence viruses compared with experiments with high

persistence viruses (Fig. S3). It is possibly because of minor genetic variants or

morphological variants, present in every viral population, which could have a higher

persistence than the average population. Minor variants proportion may increase during

the viral decay monitoring of low persistence viruses. Indeed, the average population of

high persistence viruses remained infectious for a longer period and is more

homogeneous after a long time at 35°C compared with low persistence viruses that are

submitted to a more important bottleneck.

The putative role of particle persistence in the environment for IAV transmission has been

discussed in reviews (39, 40). Using computational approaches, it was suggested that the

virus persistence in the environmental reservoir is an important parameter impacting

overwintering of IAVs, infection probability of migratory ducks in low density population

areas as well as spatial variations of IAV spreading (9, 41, 42). IAVs persistence could

also explain their evolutionary dynamic. It was shown that hemagglutinin diversity found

in avian IAVs is positively correlated to viral persistence (Roche et al. 2014). This could

allow viral strains to adapt to local environmental conditions (Lebarbenchon et al. 2012).

In addition, evidence of IAV persistence was shown in Antarctica (43) and in Siberian ice

lakes (44) where viruses are most likely stable for years (Dublineau et al. 2011; Brown et

al. 2007), justifying to further investigate the role of long-term persistence on virus

reintroduction. The present study provides for the first time experimental data highlighting

the role of the HA variability in driving influenza persistence in the environment.

In conclusion, the molecular drivers of influenza virus persistence that we identified in the

present study could help refining ecological model of IAVs transmission and their genetic

diversity in the environment. Our results are providing an experimental basis to further

investigate the role of the HA and NA proteins in driving the phenotype of persistence

among all influenza viral subtypes and establish the impact of their persistence on viral

transmission in the environment.

Methods

Cells and viruses

Madin-Darby canine kidney epithelial (MDCK) cells were maintained in Modified Eagle's

Medium (MEM) (GIBCO, ThermoFisher Scientific), supplemented with 10% foetal calf

serum (FCS) and antibiotics (100 units.mL^{-1} penicillin, 100 mg.mL^{-1} streptomycin, GIBCO,

Life Technologies). Human embryonic kidney cells 293T (HEK-293T) were maintained in

Dulbecco's Modified Eagle's medium (DMEM) (GIBCO, ThermoFisher Scientific)

supplemented with 10% FCS. All cells were incubated at 37°C in humidified 5% CO_2

incubator. MDCK cells were infected at a multiplicity of infection (MOI) of 10^{-4}

plaque-forming units per cell (pfu.cell^{-1}) and maintained in MEM without FCS in the

presence of 1 µg.mL^{-1} of TPCK-trypsin (Trypsin/L-1-Tosylamide-2-phenylethyl

chloromethyl Ketone, Whortington Biochemical Corporation) at 35°C for 3 days to

generate stock viruses. The clarified supernatants were harvested, aliquoted and stored

at -80°C.

Generation of reassortant viruses and mutated viruses

Recombinant viruses were rescued after co-transfection of MDCK and 293T co-cultivated

cells using FuGENE HD transfection reagent (Promega) at a ratio of 3:1 (µL:µg).

A/WSN/1933 recombinant virus was rescued by reverse genetics using a twelve plasmids

transfection system, with eight pPol transcription plasmids containing an individual

genomic segment under the control of a truncated human RNA polymerase I promoter

and upstream the hepatitis Delta virus ribozyme as well as four pCDNA expression

plasmids containing either the NP, PA, PB1 or PB2 gene under the control of a

cytomegalovirus promoter as described previously (45). A/Bretagne/7608/2009

recombinant virus was rescued by reverse genetics using an 8 plasmids transfection

system as described previously (Hoffmann et al. 2000). Reassortant viruses between

A/WSN/1933 and A/Bretagne/7608/2009 strains were rescued using plasmids from those

two reverse genetics systems (Reassortant viruses are listed in table 1). Reassortant

viruses harbouring HA and NA from A/Paris/2590/2009 or A/NewCaledonia/20/1999

strains with the internal genes of A/WSN/1933 were rescued using the twelve plasmids

transfection system for A/WSN/1933. Nucleotide changes were introduced into the

pPol-HA plasmid by using the QuikChange II site-directed mutagenesis kit (Agilent) in

accordance with the manufacturer's instructions. Because the pPol-HA plasmid

expressing the A/NewCaledonia/20/1999 HA segment was unstable in bacteria, the HA

DNA sequence was replaced by a codon-optimized sequence and used for transfection.

Virus identity and absence of unintended mutations were confirmed by Sanger

sequencing using BigDye Terminator v1.1 cycle sequencing kit (Applied Biosystems) and

a 3730 DNA Analyzer (Applied Biosystems).

Virus persistence in saline water at 35°C

Virus persistence in saline water at 35°C was studied as follow: viral suspensions were

diluted 10 times in saline distilled water (35 $g.L^{-1}$ NaCl) and placed in a humidified

incubator (5% CO2, 35°C) for 1 hour, 24 hours or 48 hours. The pH of the saline water

did not vary between experiments. In order to quantify their residual infectivity, MDCK

cells were seeded on a 16-well microtiter plate (30000 cells per well) coated with

microelectrode sensors in the xCELLigence® Real-Time Cell Analysis (RTCA) DP

instrument (ACEA Bioscience, Inc.) and grown for 24 hours (5% CO_2, 37°C). Cells were

then infected by exposed viruses and cell impedance, expressed as an arbitrary unit called

the Cell Index (CI), was measured through the electrodes every 15 min. Cell Index

decrease due to virus-induced cytopathogenic effect was quantified with the CIT_{50} value,

which is the necessary time in hours to measure a 50% decrease from the initial CI, always

set as the CI value 5 hours after infection. CIT_{50} values are linearly correlated to the $TCID_{50}$

titres of the viral suspensions (47) (Fig. S4). Thus, the increase of the CIT_{50} values over

time reflects the loss of infectivity. For each exposed viral suspension, CIT_{50} values were

obtained at different exposure times and plotted to calculate a linear regression slope

referred to as the inactivation slope. All data shown on viral persistence have been

obtained at different periods and with different viral stock productions. Each replicate value

of the inactivation slopes refers to a replicate virus suspension tested for persistence.

Virus concentration for functional analyses

For the experiments described below, harvested supernatants of stock viruses were

clarified and concentrated using a vivaspin 20 centrifugal concentrator (1 000 000 MWCO,

Sartorius), to reduce the initial volume by 10 times. Concentrated supernatants were then

diluted in saline distilled water (35g.L^{-1} NaCl) at a ratio of 1:10 and placed either at 4°C (0

day exposure) or at 35°C for 5 days and then kept at 4°C until used in further analysis.

Hemagglutination assay

Exposed viruses were diluted in two-fold dilution steps with PBS in a 96-wells plate and

mixed with an equal volume of a 0.75% suspension of fresh guinea pig erythrocytes. The

mixture was incubated for 1 hour at room temperature before observing erythrocytes

aggregation.

Neuraminidase activity assay

Neuraminidase activity of exposed viruses was measured using a NA-Fluor kit (Applied

Biosystems) according to manufacturer's protocol. Viruses were diluted in two-fold dilution

steps and mixed with an equal volume of MUNANA

(2'- (4-methylumbelliferyl)-alpha-D-N-acetylneuraminic acid) substrate. After 1 or 2 hours,

the level of fluorescence was quantified with a microplate fluorimeter (TriStar^2 LB942,

Berthold).

Fluorescence microscopy

For influenza virus nucleoprotein (NP) labelling, MDCK cells were seeded on glass

coverslips in 6-well plates for 24 hours before inoculation with exposed viruses diluted

1:10 in MEM. Twenty minutes or 2 hours post infection, cells were washed twice with MEM

and fixed with paraformaldehyde 2% (w/v) for 10 min. Cell membranes were labelled using

Alexa Fluor 488 conjugated wheat germ agglutinin (ThermoFisher Scientific) at 4 µg.mL^{-1}

for 10 min and permeabilized during 20 min in PBS buffer containing 0.2% Triton X-100.

Fixed cells were incubated at 4°C overnight in the presence of an anti-NP antibody

(MAB8257, Merck Millipore) diluted in PBS buffer containing 2% BSA at a concentration

of 1µg.mL^{-1}, then incubated at room temperature for 2 hours with an Alexa Fluor 647 F

(ab')2-Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary antibody (ThermoFisher

Scientific) diluted at a concentration of 2 µg.mL^{-1}. For Influenza virus envelope labelling

and endosomal fusion analyses, concentrated viral suspensions were incubated in the

presence of rhodamine B (R18, ThermoFisher Scientific) and Dioc$_{18}$ (3)

(3,3'-Dioctadecyloxacarbocyanine Perchlorate, ThermoFisher Scientific) for 2 hours at

room temperature both at a final concentration of 20 µM. Viral suspensions were then

applied on a PD-10 desalting column (GE Healthcare) using gravity flow according to the

manufacturer's instructions to remove fluorophore in suspension. MDCK cells seeded on

glass coverslips in 6-well plates for 24 hours were placed at 4°C for 5 min prior to infection

in order to synchronize viral fusion (Schelker et al. 2016) and were then inoculated with

labelled viral suspensions. After infection, the cells were incubated at 35°C for 20 min and

then washed twice with MEM and fixed with paraformaldehyde 2% (w/v) for 10 min. In all

experiments, cell nuclei were labelled using Hoescht 33342 diluted in PBS at a

concentration of 5 µg.mL^{-1} and incubated for 10 min. Glass coverslips were then mounted

on a glass slide with Prolong gold antifade reagent (ThermoFisher Scientific). Confocal

laser scanning of fluorescence was performed using LSM700 inverted microscope (Zeiss)

equipped with a plan apochromat 10X objective with a numerical aperture (N.A) of 0.45

and an enhanced contrast plan-neofluar 40X oil objective with a N.A of 1.3.

HA stability to pH

Three independent virus replicates were diluted in a pH-controlled PBS solution (at a

resolution of 0.2) with citric acid and incubated 1 hour at room temperature. Residual

infectivity of each replicate was then quantified by a TCID$_{50}$ method, as described

previously (Dublineau et al. 2011).

HA expression level by flow-cytometry

Cell surface HA expression of infected cells was measured by immunofluorescence using

flow cytometry. MDCK cells were infected for 24 hours in 6-well plates at a MOI of 0.5

pfu.cell^{-1} in the presence of TPCK-trypsin. Cell monolayers were washed twice with PBS,

re-suspended using trypsin-EDTA (ThermoFisher Scientific), fixed with paraformaldehyde

2% (w/v) for 10 min and incubated with an anti-HA monoclonal antibody (Sinobiological)

at a concentration of 1 µg.mL^{-1} at 4°C overnight. Cells were then incubated for 30 min at

room temperature in the presence of a species-specific Alexafluor 488 conjugated

secondary antibody (ThermoFisher Scientific). After centrifugation, cells were

re-suspended in PBS and analysed with an Attune NxT Flow Cytometer (ThermoFisher

Scientific). For each replicate, 50,000 cells were analysed with the same parameters.

Statistical analyses and software

Numerical data were analysed with Prism software version 6.07 (Graph Pad Software). If

needed, a Grubb's test was performed to identify and remove outliers. ANOVA tests were

performed for statistical comparison when variances were equal across samples

(Bartlett's test). Otherwise a two-tailed, Wilcoxon-Mann-Whitney's test, was used to

compare samples distribution. DNA sequence alignments and Sanger sequencing

analysis were performed using CLC Main Workbench version 7.7.3 (Qiagen).

Hemagglutinin structure was visualized using the PyMOL Molecular Graphics system

version 1.8 (Schrodinger, LLC). Microscopy imaging data were processed using LSM

software Zen Blue edition version 2.3 (Zeiss) and flow cytometry data were analysed with

FlowJo (LLC) software.

Data availability

The data that support the findings of this study are available from the corresponding author

upon request.

Acknowledgements

We thank Olivier Sawoo (Environment and Infectious Risk Unit) for the pre-screen of HA mutants based on lentiviral pseudo-particles. We thank Cyril Barbezange and Sandie Munier (laboratory Molecular Genetics of RNA Viruses, Institut Pasteur Paris) for providing reagents and advices. We are also grateful to Gaëlle Lelandais (Institut de Biologie Intégrative de la Cellule, CNRS UMR 9198) for statistical assistance and to Nathalie Pardigon (Environment and Infectious Risk Unit) for assistance with the manuscript.

This work was supported by the European funded PREDEMICS program (Preparedness, Prediction, and Prevention of Emerging Zoonotic Viruses with Pandemic Potential using Multidisciplinary Approaches) FP7/2007–2013 under Grant Agreement No 278433, the IBEID labex as well as BNP Paribas.

References

Air, Gillian M. 2012. "Influenza Neuraminidase." *Influenza and Other Respiratory Viruses* 6 (4): 245–56. https://doi.org/10.1111/j.1750-2659.2011.00304.x.

Bajimaya, Shringkhala, Tünde Frankl, Tsuyoshi Hayashi, and Toru Takimoto. 2017. "Cholesterol Is Required for Stability and Infectivity of Influenza A and Respiratory Syncytial Viruses." *Virology* 510 (July): 234–41. https://doi.org/10.1016/j.virol.2017.07.024.

Baker, N. J., and S. S. Gandhi. 1976. "Effect of Ca++ on the Stability of Influenza Virus Neuraminidase." *Archives of Virology* 52 (1–2): 7–18.

Brown, Justin D., Ginger Goekjian, Rebecca Poulson, Steve Valeika, and David E. Stallknecht. 2009. "Avian Influenza Virus in Water: Infectivity Is Dependent on PH, Salinity and Temperature." *Veterinary Microbiology* 136 (1): 20–26. https://doi.org/10.1016/j.vetmic.2008.10.027.

Brown, Justin D., David E. Swayne, Robert J. Cooper, Rachel E. Burns, and David E. Stallknecht. 2007. "Persistence of H5 and H7 Avian Influenza Viruses in Water." *Avian Diseases* 51 (s1): 285–89. https://doi.org/10.1637/7636-042806R.1.

Burmeister, Wilhelm Pascal, Stephen Cusack, and Rob W. H. Ruigrok. 1994. "Calcium Is Needed for the Thermostability of Influenza B Virus Neuraminidase." *Journal of General Virology* 75 (2): 381–88. https://doi.org/10.1099/0022-1317-75-2-381.

Byrd-Leotis, Lauren, Summer E. Galloway, Evangeline Agbogu, and David A. Steinhauer. 2015. "Influenza Hemagglutinin (HA) Stem Region Mutations That Stabilize or Destabilize the Structure of Multiple HA Subtypes." *Journal of Virology* 89 (8): 4504–16. https://doi.org/10.1128/JVI.00057-15.

Chu, C. M. 1948. "Inactivation of Haemagglutinin and Infectivity of Influenza and Newcastle Disease Viruses by Heat and by Formalin." *The Journal of Hygiene* 46 (3): 247–51.

Dublineau, Amélie, Christophe Batéjat, Anthony Pinon, Ana Maria Burguière, India Leclercq, and Jean-Claude Manuguerra. 2011. "Persistence of the 2009 Pandemic Influenza A (H1N1) Virus in Water and on Non-Porous Surface." *PLoS ONE* 6 (11): e28043. https://doi.org/10.1371/journal.pone.0028043.

Fang, Ying, Peifang Ye, Xiaobo Wang, Xiao Xu, and William Reisen. 2011. "Real-Time Monitoring of Flavivirus Induced Cytopathogenesis Using Cell Electric Impedance Technology." *Journal of Virological Methods* 173 (2): 251–58. https://doi.org/10.1016/j.jviromet.2011.02.013.

Fodor, Ervin, Louise Devenish, Othmar G. Engelhardt, Peter Palese, George G. Brownlee, and Adolfo García-Sastre. 1999. "Rescue of Influenza A Virus from Recombinant DNA." *Journal of Virology* 73 (11): 9679–82.

Hirose, Ryohei, Takaaki Nakaya, Yuji Naito, Tomo Daidoji, Yohei Watanabe, Hiroaki Yasuda, Hideyuki Konishi, and Yoshito Itoh. 2017. "Mechanism of Human Influenza Virus RNA Persistence and Virion Survival in Feces: Mucus Protects Virions From Acid and Digestive Juices." *The Journal of Infectious Diseases* 216 (1): 105–9. https://doi.org/10.1093/infdis/jix224.

Hoffmann, Erich, Gabriele Neumann, Yoshihiro Kawaoka, Gerd Hobom, and Robert G. Webster. 2000. "A DNA Transfection System for Generation of Influenza A Virus

from Eight Plasmids." *Proceedings of the National Academy of Sciences* 97 (11): 6108–13. https://doi.org/10.1073/pnas.100133697.

Hurt, Aeron C., Yvonne C. F. Su, Malet Aban, Heidi Peck, Hilda Lau, Chantal Baas, Yi-Mo Deng, et al. 2016. "Evidence for the Introduction, Reassortment, and Persistence of Diverse Influenza A Viruses in Antarctica." *Journal of Virology* 90 (21): 9674–82. https://doi.org/10.1128/JVI.01404-16.

Imai, Masaki, Tokiko Watanabe, Masato Hatta, Subash C. Das, Makoto Ozawa, Kyoko Shinya, Gongxun Zhong, et al. 2012. "Experimental Adaptation of an Influenza H5 HA Confers Respiratory Droplet Transmission to a Reassortant H5 HA/H1N1 Virus in Ferrets." *Nature* 486 (7403): 420–28. https://doi.org/10.1038/nature10831.

Ivanovic, Tijana, Jason L Choi, Sean P Whelan, Antoine M van Oijen, and Stephen C Harrison. 2013. "Influenza-Virus Membrane Fusion by Cooperative Fold-Back of Stochastically Induced Hemagglutinin Intermediates." *ELife* 2 (February). https://doi.org/10.7554/eLife.00333.

Jiang, Yongping, Kangzhen Yu, Hongbo Zhang, Pingjing Zhang, Chenjun Li, Guobin Tian, Yanbing Li, et al. 2007. "Enhanced Protective Efficacy of H5 Subtype Avian Influenza DNA Vaccine with Codon Optimized HA Gene in a PCAGGS Plasmid Vector." *Antiviral Research* 75 (3): 234–41. https://doi.org/10.1016/j.antiviral.2007.03.009.

Keeler, Shamus P., Melinda S. Dalton, Alan M. Cressler, Roy D. Berghaus, and David E. Stallknecht. 2014. "Abiotic Factors Affecting the Persistence of Avian Influenza Virus in Surface Waters of Waterfowl Habitats." *Applied and Environmental Microbiology* 80 (9): 2910–17. https://doi.org/10.1128/AEM.03790-13.

Krenn, Brigitte M., Andrej Egorov, Ekaterina Romanovskaya-Romanko, Markus Wolschek, Sabine Nakowitsch, Tanja Ruthsatz, Bettina Kiefmann, et al. 2011. "Single HA2 Mutation Increases the Infectivity and Immunogenicity of a Live Attenuated H5N1 Intranasal Influenza Vaccine Candidate Lacking NS1." *PLOS ONE* 6 (4): e18577. https://doi.org/10.1371/journal.pone.0018577.

Lang, Andrew S., Anke Kelly, and Jonathan A. Runstadler. 2008. "Prevalence and Diversity of Avian Influenza Viruses in Environmental Reservoirs." *Journal of General Virology* 89 (2): 509–19. https://doi.org/10.1099/vir.0.83369-0.

Lebarbenchon, Camille, Chris J. Feare, François Renaud, Frédéric Thomas, and Michel Gauthier-Clerc. 2010. "Persistence of Highly Pathogenic Avian Influenza Viruses in Natural Ecosystems." *Emerging Infectious Diseases* 16 (7): 1057–62. https://doi.org/10.3201/eid1607.090389.

Lebarbenchon, Camille, Srinand Sreevatsan, Thierry Lefèvre, My Yang, Muthannan A. Ramakrishnan, Justin D. Brown, and David E. Stallknecht. 2012. "Reassortant Influenza A Viruses in Wild Duck Populations: Effects on Viral Shedding and Persistence in Water." *Proceedings of the Royal Society B: Biological Sciences* 279 (1744): 3967–75. https://doi.org/10.1098/rspb.2012.1271.

Millero, Frank J., Rainer Feistel, Daniel G. Wright, and Trevor J. McDougall. 2008. "The Composition of Standard Seawater and the Definition of the Reference-Composition Salinity Scale." *Deep Sea Research Part I: Oceanographic Research Papers* 55 (1): 50–72. https://doi.org/10.1016/j.dsr.2007.10.001.

Nazir, Jawad, Renate Haumacher, Anthony Ike, Petra Stumpf, Reinhard Böhm, and Rachel E. Marschang. 2010. "Long-Term Study on Tenacity of Avian Influenza

Viruses in Water (Distilled Water, Normal Saline, and Surface Water) at Different Temperatures." *Avian Diseases* 54 (s1): 720–24. https://doi.org/10.1637/8754-033109-ResNote.1.

Plotkin, Joshua B., and Jonathan Dushoff. 2003. "Codon Bias and Frequency-Dependent Selection on the Hemagglutinin Epitopes of Influenza A Virus." *Proceedings of the National Academy of Sciences* 100 (12): 7152–57. https://doi.org/10.1073/pnas.1132114100.

Pohl, Marie O., Thomas O. Edinger, and Silke Stertz. 2014. "Prolidase Is Required for Early Trafficking Events during Influenza A Virus Entry." *Journal of Virology* 88 (19): 11271–83. https://doi.org/10.1128/JVI.00800-14.

Poulson, R. L., S. M. Tompkins, R. D. Berghaus, J. D. Brown, and D. E. Stallknecht. 2016. "Environmental Stability of Swine and Human Pandemic Influenza Viruses in Water under Variable Conditions of Temperature, Salinity, and PH." *Applied and Environmental Microbiology* 82 (13): 3721–26. https://doi.org/10.1128/AEM.00133-16.

Reed, Mark L., Olga A. Bridges, Patrick Seiler, Jeong-Ki Kim, Hui-Ling Yen, Rachelle Salomon, Elena A. Govorkova, Robert G. Webster, and Charles J. Russell. 2010. "The PH of Activation of the Hemagglutinin Protein Regulates H5N1 Influenza Virus Pathogenicity and Transmissibility in Ducks." *Journal of Virology* 84 (3): 1527–35. https://doi.org/10.1128/JVI.02069-09.

Roche, Benjamin, John M. Drake, Justin Brown, David E. Stallknecht, Trevor Bedford, and Pejman Rohani. 2014. "Adaptive Evolution and Environmental Durability Jointly Structure Phylodynamic Patterns in Avian Influenza Viruses." *PLOS Biology* 12 (8): e1001931. https://doi.org/10.1371/journal.pbio.1001931.

Russier, Marion, Guohua Yang, Jerold E. Rehg, Sook-San Wong, Heba H. Mostafa, Thomas P. Fabrizio, Subrata Barman, et al. 2016. "Molecular Requirements for a Pandemic Influenza Virus: An Acid-Stable Hemagglutinin Protein." *Proceedings of the National Academy of Sciences of the United States of America* 113 (6): 1636–41. https://doi.org/10.1073/pnas.1524384113.

Sawoo, Olivier, Amélie Dublineau, Christophe Batéjat, Paul Zhou, Jean-Claude Manuguerra, and India Leclercq. 2014. "Cleavage of Hemagglutinin-Bearing Lentiviral Pseudotypes and Their Use in the Study of Influenza Virus Persistence." *PLoS ONE* 9 (8): e106192. https://doi.org/10.1371/journal.pone.0106192.

Schelker, Max, Caroline Maria Mair, Fabian Jolmes, Robert-William Welke, Edda Klipp, Andreas Herrmann, Max Flöttmann, and Christian Sieben. 2016. "Viral RNA Degradation and Diffusion Act as a Bottleneck for the Influenza A Virus Infection Efficiency." *PLOS Comput Biol* 12 (10): e1005075. https://doi.org/10.1371/journal.pcbi.1005075.

Shigematsu, Sayuri, Amélie Dublineau, Olivier Sawoo, Christophe Batéjat, Toshifumi Matsuyama, India Leclercq, and Jean-Claude Manuguerra. 2014. "Influenza A Virus Survival in Water Is Influenced by the Origin Species of the Host Cell." *Influenza and Other Respiratory Viruses* 8 (1): 123–30. https://doi.org/10.1111/irv.12179.

Sooryanarain, Harini, and Subbiah Elankumaran. 2015. "Environmental Role in Influenza Virus Outbreaks." *Annual Review of Animal Biosciences* 3 (1): null. https://doi.org/10.1146/annurev-animal-022114-111017.

Souris, Marc, Daniel Gonzalez, Witthawat Wiriyarat, Kamlang Chumpolbanchorn,
Supaluk Khaklang, Suwannapa Ninphanomchai, Weena Paungpin, et al. 2015.
"Potential Role of Fresh Water Apple Snails on H5N1 Influenza Virus Persistence
and Concentration in Nature." *Air & Water Borne Diseases* 4 (1).
https://doi.org/10.4172/2167-7719.1000119.
Stallknecht, David E., Virginia H. Goekjian, Benjamin R. Wilcox, Rebecca L. Poulson, and
Justin D. Brown. 2010. "Avian Influenza Virus in Aquatic Habitats: What Do We
Need to Learn?" *Avian Diseases* 54 (1 Suppl): 461–65.
https://doi.org/10.1637/8760-033109-Reg.1.
Stallknecht, D.E, and J. D. Brown. 2009. "Tenacity of Avian Influenza Viruses." *Revue
Scientifique et Technique de l'OIE* 28 (1): 59–67.
Vittecoq, Marion, Hermann Gauduin, Thibault Oudart, Olivier Bertrand, Benjamin Roche,
Matthieu Guillemain, and Olivier Boutron. 2017. "Modeling the Spread of Avian
Influenza Viruses in Aquatic Reservoirs: A Novel Hydrodynamic Approach Applied
to the Rhône Delta (Southern France)." *Science of The Total Environment* 595
(October): 787–800. https://doi.org/10.1016/j.scitotenv.2017.03.165.
Weber, Thomas P, and Nikolaos I Stilianakis. 2008. "Inactivation of Influenza A Viruses in
the Environment and Modes of Transmission: A Critical Review." *The Journal of
Infection* 57 (5): 361–73. https://doi.org/10.1016/j.jinf.2008.08.013.
Xu, Xiaojin, Xueyong Zhu, Raymond A. Dwek, James Stevens, and Ian A. Wilson. 2008.
"Structural Characterization of the 1918 Influenza Virus H1N1 Neuraminidase."
Journal of Virology 82 (21): 10493–501. https://doi.org/10.1128/JVI.00959-08.
Zhang, Gang, Dany Shoham, David Gilichinsky, Sergei Davydov, John D. Castello, and
Scott O. Rogers. 2006. "Evidence of Influenza A Virus RNA in Siberian Lake Ice."
Journal of Virology 80 (24): 12229–35. https://doi.org/10.1128/JVI.00986-06.
Zhang, Hongbo, Yan Li, Jianjun Chen, Quanjiao Chen, and Ze Chen. 2014. "Perpetuation
of H5N1 and H9N2 Avian Influenza Viruses in Natural Water Bodies." *Journal of
General Virology* 95 (7): 1430–35. https://doi.org/10.1099/vir.0.063438-0.
665

 Tables and figures legends

Reassortant name	A/Bretagne/7608/2009	A/WSN/1933
Whole/2009	HA, NA, M, PB1, PB2, PA, NS, NP	-
HA-NA/2009	HA, NA	M, PB1, PB2, PA, NS, NP
M/2009	M	HA, NA, PB1, PB2, PA, NS, NP
Pol/2009	PB1, PB2, PA	HA, NA, M, NS, NP
NS-NP/2009	NS, NP	HA, NA, M, PB1, PB2, PA
Whole/1933	-	HA, NA, M, PB1, PB2, PA, NS, NP
HA-NA/1933	M, PB1, PB2, PA, NS, NP	HA, NA
M/1933	HA, NA, PB1, PB2, PA, NS, NP	M
Pol/1933	HA, NA, M, NS, NP	PB1, PB2, PA

674 **Table 1** Engineered reassortants between A/Bretagne/7608/2009 H1N1 virus and

675 A/WSN/1933 H1N1 virus

Reassortant name	A/Paris/2590/2009	A/NewCaledonia/20/1999	A/WSN/1933
HA-NA/2009	HA, NA	-	
HA-NA/1999	-	HA, NA	M, PB1, PB2, PA, NS, NP
HA$_{opti}$-NA/1999	-	HA (codon optimized), NA	

682 **Table 2** Engineered reassortants between A/Paris/2590/2009 H1N1 virus and

683 A/NewCaledonia/20/1999 H1N1 virus with the A/WSN/1933 H1N1 virus used as a

684 genomic backbone

FIG 1 Influence of viral proteins on H1N1 viruses persistence in saline water at 35°C.

(A) Viral particles were diluted in saline water (35 g.L^{-1} NaCl) and exposed to a

temperature of 35°C for 0, 1 or 2 days. MDCK cells were infected by exposed viruses and

impedance was monitored continuously and plotted as "Cell Index" (CI). (B) CI decrease

due to virus-induced cytopathogenic effect was quantified with the CIT$_{50}$ value, which is

the necessary time in hours to measure a 50% decrease from the initial CI, always set as

the cell index value 5 hours after infection (symbolized as a vertical dashed line). (C) The

CIT$_{50}$ values of an exposed virus increase with the duration of exposure in saline water at

35°C. It reflects the loss of infectivity and a linear regression analysis allowed calculating

the viral inactivation slope. (D) and (E) Inactivation slopes comparison between (D) the

2009 wild-type H1N1 virus (reference; whole/2009) and reassortant viruses harbouring

different genomic segments of the 1933 H1N1 strain or (E) between the 1933 wild-type

H1N1 virus (reference; whole/1933) and the reassortant viruses harbouring genomic

segments of the 2009 H1N1 strain. Means of inactivation slopes (horizontal lines) were

compared using a Wilcoxon-Mann-Whitney test (NS, $P>0.1$; *$P<0.05$; ***$P<0.0005$). Each

dot represents a single inactivation slope (d and e) and vertical lines represent the

standard deviation. Dashed lines represent the mean inactivation slope of the virus used

as a reference for comparison.

FIG 2 Synonymous and non-synonymous substitutions in the HA change environmental persistence. (A) Inactivation slopes of reassortant viruses bearing a HA and a NA either from 2009 (HA-NA/2009), 1999 (HA-NA/1999) or 1999 with 24% of the HA mutated by synonymous substitutions (HA$_{opti}$-NA/1999). Each dot represents a single inactivation slope and their means (horizontal lines) were compared using an ANOVA test (NS, $P>0.05$; ****$P<0.0001$). **(B)** Selected mutations either on the HA of 2009 or the optimized HA of 1999 are indicated by red spheres and arrows on the crystal structure of a 2009 H1N1 virus trimeric HA (3LZG) (side or top-down views). (C) and (D) Mean

inactivation slopes of mutated viruses were compared to the HA-NA/2009 virus (C) or

HA$_{opti}$-NA/1999 virus (D). Each dot represents a single inactivation slope value and their

means (horizontal lines) were compared using a Wilcoxon-Mann-Whitney test (NS,

P>0.05; *P<0.02; **P<0.005; ***P=0.0001; ****P<0.0001). Vertical lines represent the

standard deviation and dashed lines represent the mean inactivation slope of the virus

used as a reference for comparison.

FIG 3 The HA of virus particles is still able to bind its cellular receptor but cannot trigger membrane fusion after being exposed. (A) Evolution of the residual

hemagglutination titre (%) during exposure of reassortant viruses in saline water at 35°C.

Vertical lines represent the standard deviation (N=3). (B) Confocal immunofluorescence

microscopy (x40) of (A) MDCK cells 2 hours after infection by either the HA-NA/2009 virus

(left), or the HA-NA/1999 virus (right), which have been exposed (bottom) or not (top) for

5 days to saline water (35 g.L^{-1} NaCl) at 35°C. Signal of the influenza NP immunolabelling

is shown in orange. WGA labelling signal at the cell membrane is shown in green and the

nucleus labelling with Hoescht dye is shown in blue. For each condition a merge of all

channels (top left), the NP immunolabelling acquisition channel only (top right) or a Z-stack

of sectional views at a distance of 0.42 µM (bottom) are shown. (C) MDCK cells (x10

magnification) 20 min after infection with both HA-NA/2009 and HA-NA/1999 viruses

previously labelled with the R18 fluorophore at a self-quenched concentration. R18

dequenching signal due to the fusion of the viral envelope with the endosomal membrane

is shown in red. Scale bars represent 20 µm (B, merged and NP views), 10 µm (B,

sectional views) and 100 µm (C).

FIG 4 Virus stability to lower pH exposure and HA expression in infected cells. (A)

H1N1 reassortant viruses either with a 1999 HA (wild-type or optimized) or a 2009 HA

(wild-type or with a single amino-acid substitution) were exposed to pH-adjusted PBS

buffer for one hour and the residual infectivity values (in percentage from the initial TCID$_{50}$

titre) were calculated. Spheres represent the mean residual infectivity and vertical lines

represent the standard deviation (N=3). (B) and (C), HA surface expression in infected

743   MDCK cells analysed by flow cytometry detecting Alexafluor488 signal 24 hours after

744   infection. Horizontal lines represent the median of fluorescence intensities and HA surface

745   expression levels were compared using a Wilcoxon-Mann-Whitney test (NS, $P>0.05$;

746   *$P<0.02$; ****$P<0.0001$).

FIG 5 Non-synonymous substitutions in the NA change environmental persistence.

(A) Alignment of amino acid sequences of NA from 1999 and 2009 H1N1 viruses. Putative

amino acids involved in the second (red) and third (green) calcium-binding site are

highlighted. Selected mutations on the 1999 NA are indicated by red (NGAN/341/DGAD)

or green (GTDNN/382/DTDSD) spheres on the crystal structure of a 2009 H1N1 virus

tetrameric NA (4B7Q) (top-down view). (B) Mean inactivation slopes of viruses after

exposure to saline water at 35°C. Each dot represents a single inactivation slope and their

means (horizontal lines) were compared using an ANOVA test (*$P<0.05$; ****$P<0.0001$).

(C) and (D) Evolution of residual NA activity (%) during exposure of reassortant viruses in

saline water at 35°C. (C) and (D) represent two individual experiments. Vertical lines

represent the standard deviation (N=3). (E) NA titer of reassortant viruses measured with

a two-fold serial dilution steps and then calculated for viral titer of 8 $\log_{10}$ ($TCID_{50}.mL^{-1}$).