Title: Optimizing the conservation of migratory species over their full annual cycle

Authors: Schuster R.[a,b,*], Wilson S.[a,c], Rodewald, A. D.[d,e], Arcese P.[f], Fink, D.[d], Auer, T.[d],

Bennett, J. R[a]

[a] Department of Biology, 1125 Colonel By Drive, Carleton University, Ottawa ON, K1S 5B6

Canada.

[b] Ecosystem Science and Management Program, 3333 University Way, University of Northern

British Columbia, Prince George BC, V2N 4Z9 Canada.

[c] Environment and Climate Change Canada, National Wildlife Research Centre, 1125 Colonel

By Drive, Ottawa, Ontario, Canada, K1S 5B6.

[d]Cornell Lab of Ornithology, 159 Sapsucker Woods Rd., Ithaca, NY 14850, USA

[e] Department of Natural Resources, Cornell University, Fernow Hall, #111, Ithaca, NY 14853,

USA.

[f] Department of Forest and Conservation Sciences, 2424 Main Mall, University of British

Columbia, Vancouver BC, V6T 1Z4 Canada.

[*] Corresponding author: Richard Schuster, Department of Biology, 1125 Colonel By Drive,

Carleton University, Ottawa ON, K1S 5B6 Canada. 250-635-2321. mail@richard-

schuster.com

Keywords: migratory species, conservation, spatial optimization, citizen-science, land-sharing,

land-sparing

Abstract

Limited knowledge of the distribution, abundance, and habitat associations of migratory species

introduces uncertainty about the most effective conservation actions. We used Neotropical

migratory birds as a model group to evaluate contrasting approaches to land prioritization to

support ≥30% of the global abundances of 117 species throughout the annual cycle in the

Western hemisphere. Conservation targets were achieved in 43% less land area in plans based on

annual vs. weekly optimizations. Plans agnostic to population structure required comparatively

less land area to meet targets, but at the expense of representation. Less land area was also

needed to meet conservation targets when human-dominated lands were included rather than

excluded from solutions. Our results point to key trade-offs between efforts minimizing the

opportunity costs of conservation vs. those ensuring spatiotemporal representation of

populations, and demonstrate a novel approach to the conservation of migratory species based on

leading-edge abundance models and linear programming to identify portfolios of priority

landscapes and inform conservation planners.

Land-use change is a key threat to the conservation of biodiversity, ecosystems[1], and the services

they provide globally[2,3], and migratory species are particularly vulnerable to such change given

the vast geographic areas they occupy over the annual cycle[4,5]. Indeed, a recent global

assessment indicated that protected areas adequately protect the ranges of just 9% of migratory

bird species[5]. Strategic approaches to identify and conserve habitats critical to the persistence of

migratory species are therefore sorely needed.

Unfortunately, substantial gaps in knowledge of the abundance, distribution, and

demography of most migratory species[6] have hampered strategic planning and led to uncertainty

about the optimal allocation of conservation effort[5,7]. Given that populations of many migratory

species continue to decline[4,8], there is an urgent need to identify portfolios of lands critical to the

persistence of target species, and amenable to management in support of species conservation

without compromising human well-being.

Multi-species decision support tools can facilitate the identification of areas crucial to the

conservation of migratory species but have remained intractable due to limits on knowledge and

computing power. We capitalized on advances in crowd-sourced models of bird species

abundance and distribution[9,10] and linear programing techniques[11] to develop a robust multi-

species planning tool to estimate the land area needed to conserve 117 Nearctic-Neotropical

migratory songbirds throughout the annual cycle (SI Table 1). Specifically, we combined fine-

scale, predictive models of distribution and abundance estimated weekly throughout the year

with spatial optimization techniques[12] to identify the amount and type of land needed to reach

our conservation targets given alternative planning scenarios at hemispheric scales.

We first estimated the abundance and distribution of 117 migratory bird species weekly,

using spatiotemporal exploratory models[9,13] to calculate the relative abundance of each species

throughout the annual cycle (SI Fig. 1). We next recorded and compared the geographic area

requirements and land cover types selected when optimizing during each week of the annual

cycle (hereafter, "weekly"), versus simultaneously over the entire annual cycle (hereafter, "full

annual cycle"). Because all existing conservation plans consider stationary phases of the

breeding and non-breeding periods separately[14,15], our analysis is the first example of spatial

optimization scenarios which track populations over their full annual cycle.

We next created area-optimized solutions designed to conserve lands used by 30% of the

global populations of all 117 species in each of 52 weeks by sampling species a) over their entire

range, without accounting for population structure, or b) by sampling within 5 regional

population clusters, identified weekly to accommodate spatial variation in population structure

and migratory connectivity. Our 30% target is arbitrary, but intermediate to the 17% of terrestrial

ecosystems targeted by the Convention on Biodiversity[16] and 50% targets suggested by

comparative analysis[17], and it can be easily modified to reflect strategic goals[18].

Last, we compared area-based conservation plans designed to represent different

perspectives about the potential contribution of human-modified lands to the conservation of

migratory birds. Our 'land-sparing' approach emphasized the protection of relatively intact

habitat as indicated by a low human footprint index[19] (SI Fig. 2), whereas our 'land sharing'

approach permitted the inclusion of landscapes converted to more intensive use by humans[20].

Exploring such constraints represents a critical step in conservation planning, given that human

cultural history, values, and well-being can all affect conservation success and represent critical

inputs into structured decisions about the most efficacious actions[21–23].

Results and Discussion

The land area required to achieve conservation targets declined by 56% on average when

prioritizations were conducted over the full annual cycle rather than weekly (range = 49% to

65%; Table 1). Full annual cycle solutions also resulted in less land area being prioritized in

land-sharing and land-sparing scenarios as compared to solution based on weekly approaches

(62% and 49%, respectively; Table 1, Fig. 1, 2). These area reductions under full annual cycle

planning generally result from cases such as the inclusion of sites used by a single species in two

or more weeks of the year, or by two or more species in during two or more weeks.

Because population structure – let alone its consequences for movement or connectivity –

is poorly understood in most migratory species[24], we developed an innovative approach to

account for structure statistically. Specifically, we delineated the populations of each species into

5 spatial clusters and stratified our weekly sampling among clusters to capture the full

geographic distribution of each species. As expected, the area required to reach our conservation

targets increased when we accommodated population structure, though relatively less so under a

land-sparing (13% increase) compared to a land-sharing (26% increase) scenario (Table 1, Figs.

1, 2). Although we currently lack empirical data with which our spatial clusters can be validated,

our predictions can be tested directly as tracking and genetic mapping techniques improve to

allow comparisons of observed and predicted migration routes. That said, our current method

provides a useful approach to ensure geographic representation of population structure of a broad

suite of species using publicly-available citizen science data in spatial planning tools.

Many conservation interventions, including land protection, are constrained by limits on

fiscal or human resources and the opportunity costs of development. Our results indicate that

land area represents one of the key trade-offs in conservation designed to account for population

structure and migratory connectivity. In particular, we showed that sampling populations across

the species range each week required almost twice the amount of land compared to plans based

on the relative abundance of species mapped over the full annual cycle. Our work thus offers the

first empirical evidence to support recent calls to assess conservation needs of migratory species

across the annual cycle in ways that conserve regional representation, species diversity, and

adaptive potential[5,7,10,25]. These findings suggest a need to re-evaluate conservation planning

processes based on less precise methods. For example, government and non-governmental

organizations allocate up to $1 billion annually to bird conservation based on aspatial targets and

expert elicitation, with most actions directed to breeding habitat[14,15]. Our results suggest an

alternative approach that stands to meet conservation targets at lower land management cost and,

potentially, more compatible with human-dominated lands potentially serving a dual purpose of

supporting migratory species and human livelihoods.

Another key result of our work is that incorporating conservation objectives in human-

dominated habitats may dramatically improve the efficiency of conservation area designs if the

demographic performance of migrants is similar in 'working' and 'intact' landscapes. We found

that land-sharing approaches required 27% and 18% less land area, respectively, than land-

sparing approaches including or ignoring population structure (Table 1). Our findings thus add to

a growing body of literature indicating the need to broaden the lens through which we view

conservation to both accommodate human livelihoods and conserve valued species[21–23].

Our comparisons of land-sharing and land-sparing approaches identified other

geographical or ecosystem-related factors that might influence conservation decisions. Most

notably, land-sparing approaches selected larger areas of needle-leaved forest in boreal and

mountainous zones of western Canada, and more broad-leaved evergreen forest in the eastern

Andes and western Amazon basin (Fig. 1,2; Table 2). Weekly and full annual cycle approaches

to land-sparing resulted in geographically similar outcomes (Fig. 1,2), but also differed in land

cover types selected (Table 2). Whereas annual cycle planning with land sparing consistently

increased the amount of land prioritized over most types of land cover, weekly approaches with

and without land sparing resulted in large increases in area requirement for some cover-types and

decreases in others (e.g., Table 2 and Table S1). In particular, a weekly, land-sparing approach

favored broadleaf evergreen over mixed and broadleaf deciduous forest. Overall, these results

illustrate potential trade-offs that conservation practitioners considering optimized portfolios

must consider as additional targets and constraints are identified and incorporated in higher-level

management models[23]. Even without consensus among conservation practitioners on which

scenario to focus on, there is still a considerable amount of land selected in at least six of the

eight scenarios investigated, illustrating priority areas that most approaches agree on (126, 000

km^2, Figure 3).

Several additional caveats arise from our results, particularly with respect to land-sharing

and sparing. Implementing conservation action in working landscapes may be more challenging

than in areas with less human activity if the opportunity costs of management are higher in

developed than undeveloped landscapes. For example, even if identified as a high-priority site

for conservation in our land-sharing scenarios, land already converted to human use may be

more vulnerable to degradation in the future than more intact areas[26]. Such habitat degradation,

especially if combined with other anthropogenic stressors that may directly or indirectly reduce

survival or performance of wildlife[27], could make it difficult to reach population goals for

species even if area needs are lower compared to less developed landscapes. In practice, both

approaches are likely to be utilized given that target species will differ in their reliance on more

or less developed habitats[28]. Therefore, our approach to prioritization provides planners with

guidance on the approximate locations and requirements for land needed to meet our stated

targets under a range of scenarios. With such portfolios in hand, planners can then more readily

assess the cost-effectiveness of alternate approaches to land management and socio-economic

policies most favorable to conservation and human well-being[21–23]. We also emphasize that the

30% target used here is illustrative only. In some cases, higher targets may be needed to avoid

range contraction or the local extinction of sub-populations, to conserve ecological function such

as seed dispersal or pest control[29], or to maintain the evolutionary potential of locally-adapted

populations[25,30]. Nevertheless, our 30% target returned solutions in all cases which vastly exceed

the areal extent of existing conservation plans in support of Neotropical migrant birds.

Conclusion

Ongoing declines in the abundance and distribution of many migratory species amid severe

constraints on financial and human resources[31] points to an urgent need for area-based plans that

optimize the efficiency of conservation investments in ways that achieve conservation targets

while minimizing the opportunity costs of land conservation and impacts on human livelihood[21–

23,32]. Our solutions minimized the total land area prioritized for conservation to provide an area-

efficient portfolio of lands for further consideration by conservation planners. Three key lessons

can be derived from our results. First, scenarios based on the distributions of abundance of all

117 species over the entire annual cycle required less land area to meet conservation targets than

scenarios based on optimizations that used the weekly distributions of those species throughout

the year. Second, accounting for population structure through stratified sampling across the

entire distribution of species increased the total land area required to achieve conservation

targets. Despite requiring more land area, ensuring geographic representation may be necessary

to the long-term persistence of species, particularly in widely-distributed species with population

genetic structure potentially reflecting local adaptation to climatic conditions[25,30]. Third, area-

based plans that accommodated human activity (land-sharing) were more efficient than land-

sparing approaches that avoided areas with a high human footprint. However, because migrants

vary spatially and temporally in their tolerance of human-impacted landscapes[33], achieving

conservation goals will likely require a portfolio of sites located in both intact and disturbed

landscapes. Third, although our planning scenarios focused on Neotropical migratory birds, our

approach could be easily adjusted and replicated in other migratory species and systems with

sufficient data. In the case of birds, citizen science data and advanced prioritization tools allowed

us to reveal marked efficiencies in area-based plans spanning the full annual cycle and multiple

jurisdictions to conserve 117 individual species simultaneously.

Table 1. Area requirements to meet a 30% population target for 117 Neotropical migrant bird

species for different prioritization approaches under land sharing and land sparing scenarios.

Area Constraint	Single Population		Population Structure	
	Weekly	Annual Cycle	Weekly	Annual Cycle
Land Sharing	14.38	5.51	20.03	6.93
Land Sparing	14.54	7.45	16.44	8.45

Table entries show the area needed to meet targets (million km2). Weekly prioritizes the most

efficient target for each week of the year independently and sums the total area across all weeks.

Annual cycle prioritizes the most efficient target for all weeks combined. Single population

identifies the 30% area target for each species from anywhere within the species range.

Population structure identifies population sub-structure using a clustering approach to ensure

representation from across the range of each species in each week of the year.

Table 2. Area selected (1000 km^2) for major land cover types using full annual cycle planning for land sharing vs. sparing scenarios

and for single population vs population structure approaches.

Land cover	Area available	Single Population			Population Structure		
		Land Sparing	Land Sharing	% reduction	Land Sparing	Land Sharing	% reduction
Cropland/Mosaic Cropland	2269	339	313	8	445	439	1
Grassland	5555	1198	1088	9	1313	1238	6
Urban areas	205	9	95	-956	25	74	-196
Broadleaf Deciduous Forest	1994	627	637	-2	619	548	11
Broadleaf Evergreen Forest	6921	1595	735	54	2024	1433	29
Needleleaf Forest	4599	1359	1006	26	1395	1160	17
Mixed Forest	966	310	246	21	311	285	8
Mosaic Forest	934	207	160	23	229	194	15
Flooded Forest	540	148	97	34	162	136	16
Shrubland	4226	912	643	29	1135	864	24
Wetland	468	144	74	49	159	107	33
Barren	1053	207	79	62	208	109	48
Total	**31615**	**7055**	**5174**	**27**	**8025**	**6586**	**18**

200 Area available is the total amount of each land cover available based on all cells throughout the year where ≥ 1 species was present. %

201 reduction is the percentage decrease in the area required for each land cover type with land sharing in comparison to land sparing. Not

202 all land cover classes are included in the table and therefore individual land cover values do not sum to the total in each column. Land

203 cover data was extracted from the global land cover map for 2015 (300m resolution)[50]. See Supplemental Information Table 3 for

204 equivalent land area estimates under weekly planning scenarios.

Figure 1. Comparison of areas prioritized for weekly and full annual cycle planning under a land sharing approach allowing for the inclusion of human dominated landscapes versus a land sparing approach that excludes areas of high human footprint. The prioritization is based on a target of 30% of global populations of 117 species of Neotropical migratory birds when each species range is considered as a single population. a) = land sharing, weekly, b) = sharing annual cycle, c) = land sparing weekly, d) = land sparing annual cycle. A more detailed version of this figure focusing on northern South America is SI Figure 4.

Figure 2. Comparison of areas prioritized for weekly and full annual cycle planning under a land

sharing approach allowing for the inclusion of human dominated landscapes versus a land

sparing approach that excludes areas of high human footprint. The prioritization is based on a

target of 30% of global populations of 117 species of Neotropical migratory birds when each

species range is considered with population structure (five regional clusters). a) = land sharing,

weekly, b) = sharing annual cycle, c) = land sparing weekly, d) = land sparing annual cycle. A

more detailed version of this figure focusing on northern South America is SI Figure 5.

Figure 3. Range of agreement between the eight scenarios investigated. Darker blue indicates

that most or all scenarios selected specific areas across the Western Hemisphere, and lighter

yellow indicates areas of high scenario specificity. Scenario types considered: i) summing

scenarios for each species in each week of the year vs. optimizing over all weeks and species in a

full annual cycle, ii) including vs. ignoring spatial variation in population structure and migratory

connectivity, and iii) incorporating vs. avoiding human-dominated landscapes in solutions.

Methods

Species selection

We included 117 species of Neotropical migratory passerines for our analysis

(Supplemental Information Table 1). These species fell into two broad groups based on their

breeding and stationary non-breeding ranges: 1) species where individuals breed in North

America north of the US-Mexico border and migrate south of the Tropic of Cancer during the

non-breeding period (n=101 species, SI Table 1), and 2) species with both migratory and resident

populations or subspecies, for which individuals from migratory populations north of the US-

Mexico border move south of the Tropic of Cancer during the non-breeding period (n=16

species).

Approaches to conservation prioritization

We created 8 planning scenarios using weekly STEM models for each of 117 focal

species and incorporating different assumptions about temporal scale and cost metrics employed

in prioritization. First, we contrasted scenarios optimizing during each week of the year

separately versus simultaneously over the entire annual cycle. We next created area-optimized

solutions to conserve 30% of the global populations of all species in each week by sampling each

species a) over their entire range, without accounting for population structure, or b) as 5 regional

population clusters identified weekly to accommodate spatial variation in population structure

and migratory connectivity. Third, we compared area-based conservation plans designed to

represent different perspectives about the potential contribution of human-modified landscapes to

the conservation of migratory birds, while including either the unrestricted cost metric or the

human footprint cost metric, to create a total of 8 scenarios (SI Fig. 3). We used the prioritzr[34] R

package for the analysis, which interfaces with the Gurobi[35] optimization software.

Spatial prioritization approach

Here we use the concept of systematic conservation planning[36], to inform choices about

areas to protect, in order to optimize outcomes for biodiversity while minimizing societal costs[37].

To achieve the goal to optimize the trade-off between conservation benefit and socioeconomic

cost, i.e. to get the most benefit for limited conservation funds, we strive to minimize an

objective function over a set of decision variables, subject to a series of constraints. Integer linear

programming (ILP) is the subset of optimization algorithms used here to solve reserve design

problems. The general form of an ILP problem can be expressed in matrix notation as:

$$Minimize\ c^T x\ subject\ to\ Ax\ \geq\ b$$

Where x is a vector of decision variables, c and b are vectors of known coefficients, and A is the

constraint matrix. The final term specifies a series of structural constraints where relational

operators for the constraint can be either $\geq$ the coefficients. In the minimum set cover problem, c

is a vector of costs for each planning unit, b a vector of targets for each conservation feature, the

relational operator would be $\geq$ for all features, and A is the representation matrix with $A_{ij}=r_{ij}$, the

representation level of feature i in planning unit j. We set an objective to find the solution that

fulfills all the targets and constraints for the smallest area, which we use as our measure of cost

[11]. This objective is similar to that used in Marxan, the most widely used spatial conservation

planning tool[38].

Spatiotemporal exploratory models

We used spatiotemporal exploratory models (STEM)[9,13,39] to generate estimates of

relative abundance for each species. STEM is a type of species distribution model created as an

ensemble of local regression models generated from a spatiotemporal block subsampling design.

Repeatedly subsampling and partitioning the study extent into grids of spatiotemporal blocks,

and then fitting independent regression models (base models) in each block produces an

ensemble of partially overlapping local models. Estimates at a given location and date are made

by averaging across all the local models that contain the location and date. Combining estimates

across the ensemble controls for inter-model variability[40] and adapts to non-stationary predictor–

response relationships[13]. To account for spatial variation in the density of the bird observation

data[41], smaller spatiotemporal blocks ($10° \times 10° \times 30$ continuous days) were used north of $12°$

latitude and larger blocks ($20° \times 20° \times 30$ continuous days) were used in the southern portion of

the study extent.

The bird observation data used to implement STEM came from the eBird citizen-science

database[42]. The data included species counts from complete checklists collected under the

"traveling", "stationary", and "areal" protocols from January 1, 2004 to December 31, 2016

within the spatial extent bounded by $180°$ to $30°$ W Longitude (as well as Alaska between $150°$ E

and $180°$ E). This resulted in a dataset consisting of 14 million checklists collected at 1.7 million

unique locations, of which 10% were withheld for model validation.

Within each base model, species' occupancy and abundance was assumed to be

stationary. We fit zero-inflated boosted regression trees[9] to predict the observed counts

(abundance) of species based on three general classes of predictors: i) spatial predictors to

account for spatial (and spatiotemporal) patterns; ii) temporal predictors to account for trends;

and iii) predictors that describe the observation/detection process, which account for variation in

detection rates, a nuisance when making inference about species occupancy and abundance.

Spatial information was captured using elevation[43] and NASA MODIS land[44] and water cover

data. The MODIS data were summarized as the proportion and spatial configuration of each of

the 19 cover classes within 2.8×2.8 km (784 hectare) pixels centered at each eBird location

using FRAGSTATS[45] and SDMTools[46]. Summarizing the land-cover information at this

resolution reduced the impact of erroneous cover classifications, and reduced the impact of

inaccurate eBird checklist locations. The time of day was used to model variation in availability

for detection; e.g., diurnal variation in behavior, such as participation in the "dawn chorus"[47].

Day of the year (1-366) was used to capture day-to-day changes in occupancy, and year was

included to account for year-to-year differences. Finally, to account for variation in detection

rates variables for the number of hours spent searching for species, the length of the transect

traveled during the search, and the number of people in the search party were included in each

base model.

Estimates of relative abundance and occupancy were rendered at weekly temporal

resolution and 8.4×8.4 km spatial resolution. Each estimate was calculated as an ensemble

average across 50-100 base models. The quantity estimated was either the expected number of

birds of a given species (abundance) or the probability of the species being reported (occupancy)

by a typical eBird participant on a search starting from the center of the pixel from 7:00 to 8:00

AM while traveling 1 km.

Sampling for Population Structure and Migratory Connectivity

Many of the species used here are represented by multiple sub-species or populations known or

suspected to follow different migratory pathways and use different breeding or wintering

habitats[5,18,48]. However, in the absence of detailed knowledge on migration pathways for the vast

majority of species, we developed a system of stratified sampling to represent the weekly

distribution and spatial structure of each of 117 focal species to insure representation across their

range throughout the annual cycle. To do so, we first conducted cluster analyses of weekly

distribution maps for all 117 species to identify 5 clusters of equal abundance that encompassed

the entire species range to insure representation across it. Our cluster analysis was based on a

dissimilarity matrix of geographic locations and abundances (which were weighted by 1/3 to

primarily focus on geographic effects and not bias cluster delineation toward spatially separated

abundance clusters), and used the CLARA algorithm, which is an extension of the k-medoids

technique for large datasets[49]. Our use of 5 clusters was arbitrary but flexible, and could be

adjusted by the number of sub-species, races or sub-populations of interest.

Land use constraints

We used two metrics to constrain our systematic conservation prioritization. First, we

used a constant cost metric, where each planning unit was assigned a cost value of 1. Second, we

used human footprint (2009; 1 km resolution)[19] to identify areas more and less subject to human

use, access or development pressures; specifically, we calculated the mean human footprint value

for each 8.4 x 8.4 km pixel in our study area and used it as the 'cost' of each pixel during

prioritization.

Land cover representation

After the prioritization analyses, we summarized the major land cover types for each

scenario that we generated. We used the 2015 data set of the global land cover map[50] at a 300m

resolution and clipped the original data to the study area. For each scenario, we used the

geospatial data abstraction library[51] to warp the selected cells from the prioritization onto the

raster grid of the land cover dataset. There were 37 land cover classes identified across scenarios

and the frequency and area amount of each was summarized for all scenarios. As a final step we

combined similar land cover classes into broader classes (SI Table 2) and we used these to

examine differences in area and land cover types selected under single season vs. full annual

cycle planning and for land sparing vs land sharing scenarios (Table 2).

Code availability

All computer code used in analysis, files generated from the analysis and outputs such as

figures and tables have been deposited and will be made publicly available on publication here:

https://osf.io/58hgs/?view_only=4bddcf37b95e470da3d3d90ba0f260de. The STEM model

outputs used as inputs to the analysis will be made publicly available shortly by the Cornell Lab

of Ornithology.

Acknowledgments: RS is supported by a Liber Ero Fellowship, ADR by a Garvin endowment,

and JRB and PA by the Natural Sciences and Engineering Research Council of Canada. We also

thank the eBird participants for their contributions and eBird team for their support. This work

was funded by The Leon Levy Foundation, The Wolf Creek Charitable Foundation, NASA

(NNH12ZDA001N-ECOF), Microsoft Azure Research Award (CRM: 0518680), and the

National Science Foundation (ABI sustaining: DBI-1356308; computing support from CNS-

1059284 and CCF-1522054). Data, analysis scripts and full results are archived here:

https://osf.io/58hgs/?view_only=4bddcf37b95e470da3d3d90ba0f260de.

Author Contributions RS, SW, ADR, JRB and PA conceived the study. RS, DF and TA

collected data and conducted analyses. All authors contributed to writing and editing the paper.

References

1. Maxwell, S. L., Fuller, R. A., Brooks, T. M. & Watson, J. E. M. Biodiversity: The ravages of

guns, nets and bulldozers. *Nature* **536**, 143–145 (2016).

2. Bauer, S. & Hoye, B. J. Migratory Animals Couple Biodiversity and Ecosystem Functioning

Worldwide. *Science* **344**, (2014).

3. Semmens, D. J., Diffendorfer, J. E., López-Hoffman, L. & Shapiro, C. D. Accounting for the

ecosystem services of migratory species: Quantifying migration support and spatial

subsidies. *Ecological Economics* **70**, 2236–2242 (2011).

4. Wilcove, D. S. & Wikelski, M. Going, Going, Gone: Is Animal Migration Disappearing.

PLoS Biology **6**, e188 (2008).

5. Runge, C. A., Martin, T. G., Possingham, H. P., Willis, S. G. & Fuller, R. A. Conserving

mobile species. *Frontiers in Ecology and the Environment* **12**, 395–402 (2014).

6. Zuckerberg, B., Fink, D., La Sorte, F. A., Hochachka, W. M. & Kelling, S. Novel seasonal

land cover associations for eastern North American forest birds identified through dynamic

species distribution modelling. *Diversity and Distributions* **22**, 717–730 (2016).

7. Runge, C. A., Tulloch, A. I. T., Possingham, H. P., Tulloch, V. J. D. & Fuller, R. A.

Incorporating dynamic distributions into spatial prioritization. *Diversity and Distributions*

22, 332–343 (2016).

8. Harris, G., Thirgood, S., Hopcraft, J., Cromsight, J. & Berger, J. Global decline in

aggregated migrations of large terrestrial mammals. *Endangered Species Research* **7**, 55–76

(2009).

9. Johnston, A. *et al.* Abundance models improve spatial and temporal prioritization of

conservation resources. *Ecological Applications* **25**, 1749–1756 (2015).

10. La Sorte, F. A. *et al.* Global change and the distributional dynamics of migratory bird

populations wintering in Central America. *Global Change Biology* (2017).

doi:10.1111/gcb.13794

11. Beyer, H. L., Dujardin, Y., Watts, M. E. & Possingham, H. P. Solving conservation planning

problems with integer linear programming. *Ecological Modelling* **328**, 14–22 (2016).

12. Moilanen, A., Wilson, K. A. & Possingham, H. P. *Spatial conservation prioritization:*

quantitative methods and computational tools. **6**, (Oxford University Press Oxford, UK,

2009).

13. Fink, D. *et al.* Spatiotemporal exploratory models for broad-scale survey data. *Ecological*

Applications **20**, 2131–2147 (2010).

14. Species At Risk Act. *Bill C-5, An act respecting the protection of wildlife species at risk in*

Canada. (2002).

15. U.S. Congress. *Endangered Species Act of 1973 (16 U.S.C. 1531–1544, 87 Stat. 884)*.

(1973).

16. MacKinnon, D. *et al.* Canada and Aichi Biodiversity Target 11: understanding `other

effective area-based conservation measures' in the context of the broader target. *Biodiversity*

and Conservation **24**, 3559–3581 (2015).

17. Noss, R. F. *et al.* Bolder thinking for conservation. *Conservation Biology* **26**, 1–4 (2012).

18. Wilson, S. *et al.* Prioritize diversity or declining species? Trade-offs and synergies in spatial

planning for the conservation of migratory birds. *bioRxiv* 429019 (2018).

doi:10.1101/429019

19. Venter, O. *et al.* Sixteen years of change in the global terrestrial human footprint and

implications for biodiversity conservation. *Nature Communications* **7**, (2016).

20. Phalan, B., Onial, M., Balmford, A. & Green, R. E. Reconciling food production and

biodiversity conservation: land sharing and land sparing compared. *Science* **333**, 1289–1291

(2011).

21. Barrett, C. B. & Arcese, P. Are integrated conservation-development projects (ICDPs)

sustainable? On the conservation of large mammals in sub-Saharan Africa. *World*

development **23**, 1073–1084 (1995).

22. Ban, N. C. *et al.* A social–ecological approach to conservation planning: embedding social

considerations. *Frontiers in Ecology and the Environment* **11**, 194–202 (2013).

23. Schwartz, M. W. *et al.* Decision Support Frameworks and Tools for Conservation.

Conservation Letters **11**, e12385 (2018).

24. Webster, M. S., Marra, P. P., Haig, S. M., Bensch, S. & Holmes, R. T. Links between

worlds: unraveling migratory connectivity. *Trends in Ecology & Evolution* **17**, 76–83 (2002).

25. Bay, R. A. *et al.* Genomic signals of selection predict climate-driven population declines in a

migratory bird. *Science* **359**, 83–86 (2018).

26. Haddad, N. M. *et al.* Habitat fragmentation and its lasting impact on Earth{\textquoteright}s

ecosystems. *Science Advances* **1**, (2015).

27. Loss, S. R., Will, T. & Marra, P. P. Direct Mortality of Birds from Anthropogenic Causes.

Annual Review of Ecology, Evolution, and Systematics **46**, 99–120 (2015).

28. Jenkins, C. N., Pimm, S. L. & Joppa, L. N. Global patterns of terrestrial vertebrate diversity

and conservation. *Proceedings of the National Academy of Sciences* **110**, E2602–E2610

(2013).

29. Kenis, M., Hurley, B. P., Hajek, A. E. & Cock, M. J. W. Classical biological control of

insect pests of trees: facts and figures. *Biological Invasions* 1–17 (2017).

30. Arcese, P. & Keller, L. Population Structure. in *Ornithology: Foundation, Analysis, and*

Application (eds. Morrison, M. L., Rodewald, A. D., Voelker, G., Colón, M. R. & Prather, J.

F.) (JHU Press, 2018).

31. McCarthy, D. P. *et al.* Financial costs of meeting global biodiversity conservation targets:

current spending and unmet needs. *Science* **338**, 946–949 (2012).

32. Bottrill, M. C. *et al.* Is conservation triage just smart decision making? *Trends in Ecology &*

Evolution **23**, 649–654 (2008).

33. Faaborg, J. *et al.* Recent advances in understanding migration systems of New World land

birds. *Ecological Monographs* **80**, 3–48 (2010).

34. Hanson, J. *et al.* prioritizr: Systematic Conservation Prioritization in R, Version 3.0.3.

(2017).

35. Gurobi Optimization Inc. Gurobi Optimizer Reference Manual, Version 7.5.1. (2017).

36. Margules, C. R. & Pressey, R. L. Systematic conservation planning. *Nature* **405**, 243–53

(2000).

37. McIntosh, E. J., Pressey, R. L., Lloyd, S., Smith, R. & Grenyer, R. The Impact of Systematic

Conservation Planning. *Annual Review of Environment and Resources* **42**, annurev-environ-

102016-060902 (2017).

38. Ball, I. R. R., Possingham, H. P. P. & Watts, M. E. E. Marxan and relatives: Software for

spatial conservation prioritisation. in *Spatial conservation prioritisation: Quantitative*

methods and computational tools. (eds. Moilanen, A., Wilson, K. & Possingham, H. P.) 185–

195 (Oxford University Press, 2009).

39. Fink, D. *et al.* Crowdsourcing meets ecology: hemisphere-wide spatiotemporal species

distribution models. *AI magazine* **35**, 19–30 (2014).

40. Efron, B. Estimation and accuracy after model selection. *Journal of the American Statistical*
Association **109**, 991–1007 (2014).

41. Fink, D., Damoulas, T. & Dave, J. Adaptive Spatio-Temporal Exploratory Models:
Hemisphere-wide species distributions from massively crowdsourced eBird data. in *AAAI*
(2013).

42. Sullivan, B. L. *et al.* The eBird enterprise: an integrated approach to development and
application of citizen science. *Biological Conservation* **169**, 31–40 (2014).

43. Amatulli, G. *et al.* A suite of global, cross-scale topographic variables for environmental and
biodiversity modeling. *Scientific Data* **In press**, (2017).

44. Friedl, M. A. *et al.* MODIS Collection 5 global land cover: Algorithm refinements and
characterization of new datasets. *Remote sensing of Environment* **114**, 168–182 (2010).

45. McGarigal, K., Cushman, S. A. & Ene, E. FRAGSTATS v4: spatial pattern analysis program
for categorical and continuous maps. University of Massachusetts, Amherst, Massachusetts,
USA. *goo. gl/aAEbMk* (2012).

46. VanDerWal, J., Falconi, L., Januchowski, S., Shoo, L. & Storlie, C. SDMTools: Species
Distribution Modelling Tools: Tools for processing data associated with species distribution
modelling exercises. *R package version* 1 (2014).

47. Diefenbach, D. R., Marshall, M. R., Mattice, J. A. & Brauning, D. W. Incorporating
availability for detection in estimates of bird abundance. *The Auk* **124**, 96–106 (2007).

48. Martin, T. G. *et al.* Optimal conservation of migratory species. *PLoS One* **2**, 751 (2007).

49. Kaufman, L. & Rousseeuw, P. J. Partitioning around medoids (program pam). *Finding*
groups in data: an introduction to cluster analysis 68–125 (1990).

50. ESA Climate Change Initiative. Global land cover map 300m resolution for 2015. (2017).

Available at: http://maps.elie.ucl.ac.be/CCI/viewer/download.php.

51. Warmerdam, F. The geospatial data abstraction library. in *Open source approaches in*

spatial data handling 87–104 (Springer, 2008).

489