1

A sub-movement time scale of human motor adaptation

Abbreviated Title: Adaptive Control of Reaching

F. Crevecoeur[1,2], J.-L. Thonnard[1,2,3], P. Lefèvre[1,2]

[1]*Institute of Communication and Information Technologies, Electronics and Applied Mathematics*

(ICTM), UCLouvain, Belgium

[2]*Institute of Neuroscience, UCLouvain, Belgium*

[3]*Cilniques Universitaires Saint-Luc, UCLouvain, Belgium*

Corresponding Author:

F. Crevecoeur

4 avenue Georges Lemaitre

1348 Louvain-la-Neuve, Belgium

email: frederic.crevecoeur@uclouvain.be

Abstract

Humans and animals adapt motor commands to predictable disturbances within tens of trials in laboratory conditions. However in natural conditions, adaptation likely occurs continuously and it remains unknown how the brain handles novel disturbances across time. Because motor commands and sensory feedback together carry continuous information about limb dynamics, we hypothesized that the nervous system could adapt to unexpected disturbances online. We tested this hypothesis in two reaching experiments during which opposing velocity-dependent force fields were randomly applied. We found that within-movement feedback corrections gradually improved, despite the fact that the perturbations were unexpected. Moreover, when participants were instructed to stop at a via-point, the application of a force field prior to the via-point induced mirror-image after-effects after the via-point, indicative of within-trial adaptation to the unexpected dynamics. These findings highlight a fast time-scale of motor learning, which supplements feedback control and supports adaptation of an ongoing movement.

Introduction

Neural plasticity in the sensorimotor system enables adaptive internal representations of

movement dynamics, which supports motor adaptation and acquisition of motor skills with practice

[1]. In the context of reaching movements, studies have documented that healthy humans and animals

can learn to anticipate the impact of a force field applied to the limb, and recover straight reach

paths within tens to hundreds of trials [2-7]. Importantly, studies on motor learning have consistently

highlighted the presence of an after-effect, which mirrors the initial movement deviation and

indicates that adaptation was supported by a novel internal model of movement dynamics [8-10].

To date, motor learning and adaptation have been exclusively studied on a trial-by-trial basis,

highlighting context-dependent learning rates and memory dynamics across movements [11-15].

Although this approach has revealed several fundamental properties of the motor system, it has left

largely unexplored the problem of online control during early exposure to the force field. More

precisely, it remains unknown how the nervous system controls reaching movements in the presence

of unexpected dynamics, or errors in the internal models, as occurs when a force field is introduced

experimentally. To address this question, we sought to investigate whether within-movement error

corrections could be mediated by rapid adaptation, allowing partial compensation for model errors

in real-time.

In fact, it is often assumed that unexpected disturbances during movements are

automatically countered by the limb's mechanical impedance and by reflexes [3,16,17]. However, recent

work suggests that muscles viscoelastic properties, as well as the gain of the spinal reflex (latency

~20ms for upper limb muscles) are low at spontaneous levels of muscle activation [18]. Furthermore,

long-latency feedback (latency ≥50ms) is also based on internal models of the limb and

environmental dynamics [19-21], even when correcting for very small disturbances [22]. Thus, the

presence of model errors is equally challenging for rapid feedback control as it is for movement

planning, and yet healthy humans seem to handle the presence of an unexpected force field

relatively well.

In theory, an approximate internal model can be deduced during movement because motor

commands and sensory feedback together carry information about the underlying dynamics. This

problem in engineering were studied in the framework of *adaptive control*[23-25]. The idea of adaptive

control is to adapt the model continuously based on sensory feedback about the unexpected

movement error. During reaching experiments, the unexpected introduction of a force field produces

a model error, which generates a hand path deviation that can be used to estimate the underlying

dynamics in addition to the state of the limb. It is important to distinguish adaptive control from

standard (extended) stochastic optimal control. Indeed, stochastic optimal control is based on the

assumption that there is no fixed model error, and that the disturbances in the process follow a

known distribution [26,27]. The goal of adaptive control is to correct for the possible presence of model

errors continuously.

Considering the very rich repertoire of rapid sensorimotor loops [28], we explored the

possibility that mechanical disturbances to the limb can update the internal models online following

the principles of adaptive control. Online estimation of a task-related parameter was reported in the

context of visuomotor rotations [29], where participants had to estimate the mapping between hand

and cursor motion. Here we address this problem in the context of reaching in a force field to probe

specifically the internal representations of limb dynamics. We found that movements displayed

online adjustments consistent with real-time motor adaptation. We further illustrate in a simple

model that these effects, as well as standard learning curves, can be simulated in the framework of

adaptive control. We suggest that adaptive control likely constitutes an important component of

motor learning, as it bridges online motor corrections with learning across few trials. We also

highlight the associated computational challenges for the nervous system that have yet to be fully

characterized.

Results

Participants grabbed the handle of a robotic device and performed forward reaching

movements towards a visual target (Fig. 1a). In the first experiment, clockwise and counterclockwise

orthogonal force fields (FF) were randomly applied as catch trials, such that neither the occurrence

nor the direction of the force field could be anticipated. Reach paths were of course strongly

impacted by the presence of the force field, however the online correction became smoother with

practice (compare trials #1 and 20 in Fig. 1b). To quantify this, we extracted the maximum hand

deviation along the direction of the force field, and the target overshoot near the end of the

movement (Fig. 1b: Max. and Osh., respectively). We found that the maximum deviation did not

evolve significantly across force field trials (Fig. 1c), whereas the overshoot displayed a significant

exponential decay (Fig. 1d, $R^2 = 0.17$ and 0.56 for clockwise and counterclockwise force fields,

respectively). We also observed that exposure to a single force field trials induced significant after-

effect on the next trial, as previously reported in a random adaptation paradigm [30] (Supplementary

figure S1). The absence of a clear change in the maximum deviation confirms that there was no

anticipation, or default strategy that could counter the unexpected disturbance, yet the online

corrections improved across trials.

100　　　　　　　　　　　　　　　　(Figure 1 about here)

The reduction in target overshoot resulted from a smoother correction for the lateral

displacement. Indeed, there was no significant change in the peak forward hand velocity across trials

for both clockwise (rmANOVA, $F_{(23,299)} = 0.56$, $P = 0.9$) and counterclockwise force fields ($F_{(22,286)} =$

0.87, $P = 0.62$). Thus the perturbation applied to the limb remained statistically similar. However, the

absolute peak hand force near the end of the movement decreased significantly. This is illustrated in

Fig. 2a for average continuous traces, and in Fig. 2b showing the significant exponential decay of the

terminal peak hand force ($R^2 = 0.17$ and 0.11 for clockwise and counterclockwise perturbations). As

a consequence, the correlation between the measured lateral force and the force field perturbation,

corresponding to the forward velocity multiplied by the scaling coefficient of the force field,

significantly increased across trials (Fig. 2c). In the case of perfect compensation, the measured force

should equal the action of the force field, and a correlation of 1 is expected. Thus, the increase in

correlation between the measured force and the force field within trials indicated that the online

correction became progressively tuned to the properties of the force field.

116   (Figure 2 about here)

These results are expected in the framework of adaptive control. However it remains

possible that participants learned to produce smooth corrective movements without learning about

the force field specifically. To address this, we sought a more direct link between the online

correction and the force field by using a via-point located at two thirds of the reach path. In this

experiment, participants were given scores for good trials, and bonuses when they stopped at the

via-point. The bonus was awarded if the hand velocity inside the via-point dropped below 3cm/s.

Force field trials were randomly interleaved, with and without via-points. In a random subset of

trials, the force field was turned off at the via-point. We reasoned that if the feedback correction

during force field trials reflected an online update of the internal models, then these corrections

should evoke almost instantaneous after-effects for the remainder of the movement from the via

point to the goal target.

Our results confirmed this prediction. First we reproduced the observations made in the first

experiment for the trials without via-point (see supplemental figure S2). Second, we found a

systematic after-effect between the via-point and the goal target that mirrored the impact of the

force field experienced between the start target and the via-point (Fig. 3a for average hand paths

and 3b). The stopping at the via-point was necessary for two reasons: first it allowed us to turn off

the force field unnoticeably because the velocity was close to zero. Second, it also ensured that the

after-effects were not due to momentum following the online correction, because the velocity was

again very close to zero at the via-point. Thus the after-effects between the via-point and the goal

target were due to a re-acceleration in the direction opposite to the perturbation induced by the

force field. We extracted the minimum lateral hand velocity in the via-point and the lateral hand

velocity at the second peak (Fig. 3c: squares and open discs on exemplar traces). As expected, there

was no difference across clockwise, baseline, and counterclockwise trials for the minimum velocity

(Fig. 3d). In contrast, the lateral hand velocity measured at the second maximum displayed a very

strong modulation consistent with an after-effect (Fig. 3d, rmANOVA on average lateral hand speed

across force field: $F_{(2,14)}$ = 17.23, P < 0.001).

145   (Figure 3 about here)

Finally we were interested to see how early the modulation of the lateral hand velocity

following the via-point occurred. In addition to the main effect of the force field, we found a

significant interaction between the force field and the block number ($F_{(10,70)}$ = 2.34, P <0.05).

Strikingly, the effect reported in Fig. 3 was already present in the first block (Fig. 4b, paired t-test on

data from block 1: t_8 = 3.6, P = 0.007, effect size: 1.2). This difference increased across blocks,

indicating an increase in the rate of online adjustments of control (data from block 6: t_8 = 5.8, P <

0.001, effect size: 1.95).

155   (Figure 4 about here)

The presence of a significant modulation in the first block indicated that online adaptation

occurred during the first few trials (there were 5 trials with via-point and switching-off of the force

field per direction and per block). This striking result warranted further investigation: can these

adjustments occur within the very first trial? Our dataset cannot provide a definitive answer because,

due to randomization, some via-point trials were preceded by force field trials without via-point for

some participants. Nevertheless, we used a statistical model of the modulation of lateral velocity as a

function of trial number to obtain the beginnings of an answer.

We first changed the sign of velocity during CW trials such that an after effect would

correspond to a positive velocity for all trials. Then, we accounted for idiosyncrasy by including

participants as a random factor in a mixed linear model. We found a significant positive correlation

between the lateral hand velocity and the trial index. Most importantly, the regression included an

offset that was significantly distinct from 0. The intercept value was $0.014ms^{-1}$, and the *P*-values for

the slope and intercept were 0.014 and 0.006, respectively. The same regression performed on the

baseline trials, where no force field was applied prior to the vial point, revealed an offset of -0.007,

which was not significant (*P* = 0.06). Thus, we can reject the hypothesis that the lateral hand velocity

was zero in the very first trial. In other words, our dataset supports the hypothesis that online

adjustments occurred even during the very first trial. Lastly, we compared the lateral hand velocity

from the very first successful trial in each direction and from each participant, and also found a

significant modulation of hand velocity following the via-point based on a single trial in each

direction. Indeed the lateral velocity following CCW perturbations was significantly higher than

following CW perturbations (one-tail paired t-test, $t_{(8)}$ = 1.91, *P* = 0.044).

These results indicate that the internal representation dynamics was updated during

movements, and could modify the previously planned movement from the via-point to the goal

target in a way that was consistent with a near-instantaneous after-effect. Furthermore, statistical

modeling suggested that such adjustments occurred during the very first trial. It is possible that

participant used three internal models, one for null-field trials and one per force-field direction, and

switched between them dependent on the ongoing movement. Although plausible, this view does

not easily explain how these models are learned in the first place nor how they are refined across

trials to account for the larger amounts of online adaptation observed across the blocks (Fig. 4). A

more compelling view in our opinion is the hypothesis that the nervous system uses adaptive control,

describing how sensory feedback and motor commands are used to track model parameters online.

To illustrate this, we now present the theoretical model and simulations based on adaptive

control. First it is important to highlight that adaptive changes in the internal models differ from the

sensorimotor loops that correct for perturbations altering the state of the limb (Fig. 5a: inner loop,

solid). In the inner loop, sensory feedback is used to estimate the state of the system through an

observer (e.g. Kalman filter), and the controller generates motor commands that correct for motor

errors. This loop is responsible for adjusting motor commands following mechanical disturbances to

the limb or following target jumps, two situations requiring an update of the ongoing command to

comply with the goal of the task.

197   (Figure 5 about here)

In the framework of adaptive control, there is a second loop (Fig. 5a: outer loop, dashed),

which estimate the model based on current sensory data and motor commands [23-25]. This estimation

is indirect because there is no direct measurement of the model parameters, thus the motor

command and sensory feedback must be used to deduce the structure of the underlying dynamics.

The simulations performed in this study are based on iterative least square identification (RLS) [25], but

other techniques such as expectation-maximization may be considered [31]. However it is important to

note that the increase in after-effect observed across the blocks of Experiment 2 required the

presence of a parameter to change the online learning rate, which is why we privileged the RLS

formalism as it includes such parameter by design (see Methods, γ). The outer loop updates the

model used in the estimator and in the controller of the inner loop (Fig. 5a, oblique arrows). The

second experiment was designed to highlight the presence of this loop: at the via-point with near-

zero velocity, there is no deviation or error in the state of the limb that requires correction by the

inner loop. Yet, participants did not simply reach straight ahead because of change in the internal

models performed by the outer loop during the first portion of the reaching movement.

Using this framework, we reproduced standard learning curves in computer simulations, with

exponential decay of the lateral hand displacement, mirror-image catch trials, and unlearning

following the catch trials as in typical human learning experiments (Fig. 5b) [6]. This model also

reproduced the reduction in peak end force occurring near the end of movement as observed in

Experiment 1 by assuming that the online learning rate increased across force field trials (Fig. 5c, see

also Methods), that is the online adjustments were smaller in the first trials than in the last trials.

This idea is consistent with the presence of savings, characterizing a faster re-learning upon exposure

to a novel environment experienced before [11,13]. Here we simply hypothesized a similar faster

relearning when exposed to a previously experienced perturbation based on the continuous inflow

of sensory information and an increased online learning rate (γ parameter). Finally, we also

reproduced the presence of after-effects after switching off the force field at the via-point and

extracting the lateral velocity exactly as in Experiment 2 (Fig. 3c and 5d). Again, the modulation of

online adaptation documented across trials and blocks could be reproduced by increasing the online

learning rate similar to the faster trial-by-trial re-adaptation observed in previous learning

experiments [13].

Discussion

Controlling movements in the presence of model errors is a challenging task for robotics as well as for the nervous system. However healthy humans can handle model errors like those arising when a force field is introduced experimentally during reaching. To understand this, we explored the possibility that the nervous system reduces the impact of model errors online following the principles of adaptive control. We found that participants' feedback corrections to unexpected force fields became gradually better (Experiment 1), and evoked within-trial after-effects for an ongoing sequence of movements (Experiment 2). We showed that these behaviors, as well as standard learning curves, were captured in a simple adaptive control model in which motor commands, state estimates, and sensory feedback are used to update the internal models of dynamics during movement. The strengths of the theoretical model were that it reproduced some aspects of our experimental findings, along with the typical exponential decay observed in human motor adaptation, and based on online continuous processing of sensory feedback. Thus it constitutes a powerful candidate model for linking online control with motor learning.

We should emphasize that there is no disagreement between adaptive control and stochastic optimal control (LQG, [26]), but instead there is complementarity between these models. Stochastic optimal control is based on the assumption that the system dynamics are known exactly, and that the process disturbance is known in distribution [32]. The unexpected introduction of a force field introduces a fixed error in the model parameters, which is not a stochastic (random) disturbance. This violates the assumption of stochastic optimal control and warrants the use of control design that explicitly considers the possibility of an error in the model parameters. Adaptive control aims at solving this problem.

These results provide evidence that separate motor memories of opposite perturbations can be learned and retrieved within movement. Previous work showed that distinct planning conditions were critical to learn opposite perturbations, such as explicit cues [33], distinct representations of movements [34], or even distinct prior or follow-through movements associated with the same

movement [35,36]. In our study, there was no cue given about the upcoming perturbation, and

movements' internal representations were identical across baseline and force field trials. Thus, it was

the physical state of the limb during movement execution that evoked changes in neural

representations of reach dynamics.

The retrieval of distinct motor memories was highlighted by the increase in online adaptation

rate responsible for the gradual increase in lateral modulation of hand velocity (Fig. 4). This result

highlights that participants were able to recognize a previously experienced perturbation, or

assumed that the ongoing perturbation was the same as the previously experienced ones, and

increased the within-trial adaptation rate. However, it may not be necessary to assume that

previously learned internal models are retrieved. Indeed, our data indicated that adjustments could

happen within the very first trial, and this is captured in the model in which adaptation can occur in

real-time through parameter tracking without assuming a pre-stored alternative model. We suggest

that online tracking in real-time combined with a change in online learning rate upon re-exposure to

a previously experienced perturbation is a parsimonious account of our data.

It remains unclear how quickly these adjustments occurred within movements, and this

represents an important but challenging question for future research. Anatomically, long-latency

corrections for perturbation to the limb (~60ms) are supported by a distributed network through

cerebellum [37] primary somatosensory area, motor and pre-motor cortices, and parietal cortex [28,38,39].

This network clearly overlaps with the main regions associated with short-term plasticity [1].

Behaviorally, long-latency feedback exhibits exquisite sensitivity to even very small disturbances [22].

Thus, the network of brain regions engaged in trial-to-trial learning is also likely recruited very early

during unexpected force field trials, and it is conceivable that rapid adjustments occurred within this

rapid pathway.

Computationally, however, the correction of internal models involves a less direct mapping

between sensory feedback and the internal representations of movement dynamics, in addition to a

change in the neural controller. This computational load may require longer processing times

associated with model updates in comparison with feedback responses following perturbations to

the limb or to the movement goal. As an absolute upper bound, we may consider that adjustments in

Experiment 2 were performed between reach onset and the moment when participants exited the

via-point, which was typically less than 0.5s. We expect that future work aim at measuring the

latency of this outer loop more accurately.

285   Another clear challenge for future work is to unravel the underlying neural mechanism. The

key point is to understand how the brain generates a near instantaneous after effect like those

observed in Experiment 2. Indeed, at the via-point with near-zero velocity across all trials, there was

no motor error to correct; yet the subsequent motor output reflected previously encountered

disturbances. One possibility to account for this result is through synaptic plasticity [1]: that is the gain

of the same sensorimotor loop changes due to re-afference of sensory prediction errors inducing

changes in synaptic strength. Another possibility is that the unexpected force field generated

changes in the state of the sensorimotor system along a dimension which does not influence the

motor output [40,41], allowing similar biomechanical state (i.e. stopping at the via-point) followed by a

distinct motor output across force field trials when leaving the via-point. Such mechanism does not

necessarily engage changes in synaptic weights, which is computationally advantageous [23].

Unraveling the latency of the model updates will likely set physiological constraints on the candidate

neural underpinnings.

298   Finally, our study also opens questions from a theoretical standpoint. Indeed, we must recall

299   that our model was excessively simple: we considered the translation of a point-mass in the plane

300   with full state information (see Methods and Supplementary Material). We thus expect that future

301   modeling work unveil the computational challenges that arise when considering more realistic

302   models of the neuro-musculo-skeletal system, including nonlinear dynamics, sensorimotor noise, and

303   temporal delays.

Methods

Experimental Procedures

Healthy volunteers (n=24) provided informed written consent and the experimental procedures were approved by the ethics committee at the host institution (UCLouvain). Participants grabbed the handle of a robotic device (KINARM, BKIN Technologies, Kingston, Canada) and were instructed to perform reaching movements towards a visual target. The movement onset was cued by filling in the goal target (Fig. 1a), and feedback was provided about movement timing (good trials were between 600ms and 800ms following the go signal). In Experiment 1 (n = 14), baseline trials were randomly interleaved with orthogonal force field trials mapping the forward hand velocity onto a lateral force ($F_x = \pm L\dot{y}, L = 13\ Nsm^{-1}$). Movements consisted of 15cm reaches in the forward direction. To make the task less predictable, other trial types were also randomly interleaved including trials with a via-point situated on the side of the straight reach path, and trials with a constant background load applied to the hand ($F_x = \pm 4N$). Participants performed six blocks of 80 trials including 4 force field trials per direction per block, summing to a total of 24 force field trials per direction and participant.

In Experiment 2, movements consisted of 18cm reaches in the forward direction, and the via-point (radius: 1cm) was located at 12cm on the straight line joining the start and goal target. Participants performed six blocks of 80 trials composed of 45 baseline trials, 5 force field trials, 5 baseline trials with a via-point, 5 force field trials with a via-point, and 5 force field trials with a via-point and with a switching off of the force field. All trials and force field directions were randomly interleaved. The experimental setup monitored the hand velocity in real time allowing to turn off the force field if (1) the hand cursor was in the via-point and (2) the hand velocity dropped below 3cm/s while in the via-point (determined based on preliminary testing). Feedback about a successful stopping at the via-point was given online. This task was difficult and we recorded an average of 21, 25 and 17 trials for counterclockwise, baseline and clockwise via-point trials meeting these two conditions, respectively (range across subjects: [5, 30], [18, 29], [2, 30]).

Analysis and Statistical Design

Position and force signals were sampled at 1kHz. Velocity signals were derived from numerical differentiation (2nd order finite difference algorithm). The presence of significant effects was assessed with repeated-measures ANOVAs (rmANOVA). In experiment 1, one participant missed a force-field trial such that the analyses were performed on 23 trials for the corresponding force field direction. Post-hoc tests were performed based on paired t-test with Bonferroni correction for multiple comparisons. The exponentials were fitted in the least-square sense and the significance of the fit was determined based on whether the 95% confidence interval of the exponent responsible for the curvature of the fit included or not the value of zero. An R^2 statistics was derived as follows:

$$R^2 = 1 - \frac{V_{RES}}{V_{TOT}}, \tag{1}$$

where V_{RES} denotes the variance of the residuals and V_{TOT} is the variance of the data. We adopted Cohen's d to quantify the effect size for paired comparison as the mean difference between two populations divided by the standard deviation of the paired difference between the populations [42]. The mixed linear model of lateral hand velocity as a function of trial number was based on standard techniques [43]. We changed the sign of the lateral velocity for clockwise force field trials such that for all trials, a positive modulation reflected the presence of an after effect. The dependent variable was the lateral hand velocity measured at the moment of the second velocity peak, the fixed factor was the trial index, and the random effect was the participant.

Model

The model is based on the translation of a point-mass with unity mass in the horizontal plane. The state-space representation is:

$$\ddot{x} = -k\dot{x} + F_x + F_{ENV}, \tag{2}$$

$$\ddot{y} = -k\dot{y} + F_y, \tag{3}$$

$$\tau\dot{F}_x = u_x - F_x, \tag{4}$$

$\tau \dot{F}_y = u_y - F_y,$ (5)

where k is a dissipative constant (0.1 Nsm), Equations 4 and 5 capture the first order muscle

dynamics with time constant set to 0.1s, and F_{ENV} represents the unknown environmental

perturbation: $F_{ENV} = 0$ for the null field, or $F_{ENV} = \pm L\dot{y}$ for force field trials.

This system of differential equations can be transformed in a discrete-time state space

representation allowing us to add stochastic disturbances. Let x^* and y^* denote the coordinates of

the goal target relative to the start, $\boldsymbol{z}_t := [x_t, \ y_t, \ \dot{x}_t, \ \dot{y}_t, \ F_{x,t}, \ F_{y,t}, \ x^*, \ y^*]^T$ represent the state

vector at the time step t, and $\boldsymbol{u}_t := [u_{x,t}, u_{y,t}]^T$ represent the two-dimensional control vector, then

the system can be written as follows:

$\boldsymbol{z}_{t+1} = (\hat{A}_t + \Delta A)\boldsymbol{z}_t + B\boldsymbol{u}_t + \boldsymbol{\varepsilon}_t,$ (6)

where $\hat{A}_t$ is the current (time varying) expected dynamics, ΔA is the unknown model error containing

the unmodeled environmental disturbance (F_{ENV}), and $\boldsymbol{\varepsilon}_t$ is a Gaussian disturbance with zero-mean

and covariance $\Sigma_\varepsilon := BB^T$. We denote the true unforced dynamics by the matrix $A := \hat{A}_t + \Delta A$.

Details about these matrices can be found in the Supplementary Material.

We assumed for simplicity that the controller has perfect state measurement, meaning that

$\boldsymbol{z}_t$ is known (fully observable case). The problem is that the controller does not know ΔA. This

problem was approached in the context of sensorimotor control assuming that ΔA followed a known

(Gaussian) distribution [27], which induces state-dependent noise. Observe that this assumption

implies that the force field intensity varies randomly over time, which is not consistent with our

experiment for which the force field varied randomly across trials, but remained fixed within trials. In

other words, stochastic optimal control does not apply in the context of this paper because ΔA

depends on F_{ENV}, which is not a stochastic variable but instead fixed bias or error in the model. Thus

the initial controller is derived assuming that the system dynamics corresponds to A. To further

simplify the problem, we assume that the controller knows which parameter is unknown (L). Let $\hat{A}_t$

denote the expected dynamics at time t dependent on the current estimate of the force field

intensity noted $\hat{L}_t$, a Linear-Quadratic-Gaussian controller can be derived giving a time series of

control signals based on linear feedback gains that depend on $\hat{A}$:

$$\boldsymbol{u}_t = -C_t(\hat{A}_t)\boldsymbol{z}_t. \qquad\qquad (7)$$

Now, under this assumption, we may predict the next state vector, which we designate by $\hat{\boldsymbol{z}}_t$.

When the true dynamics is distinct from the expected dynamics ($\Delta A = A - \hat{A}_t \neq 0$), there is a

prediction error, $\boldsymbol{e}_t := \boldsymbol{z}_t - \hat{\boldsymbol{z}}_t$, which can be used to correct the estimate of the model. We follow

standard Recursive Least Square identification techniques (RLS), and use the following adaptive rule

to update the estimate of L [25]:

$$\hat{L}_{t+1} = \hat{L}_t + \gamma \frac{\partial \hat{\boldsymbol{z}}_t}{\partial L} \boldsymbol{e}_t,$$

390   (8)

where γ is the online learning rate. This new value of estimated force field intensity is then used to

update in $\hat{A}_{t+1}$, and a new series of feedback gains is computed and applied to the next state as

control law: $\boldsymbol{u}_{t+1} = -C_{t+1}(\hat{A}_{t+1})\boldsymbol{z}_{t+1}$. In all, the closed loop controller consists in iteratively deriving

optimal control gains based on an adaptive representation of the unknown parameters recursively

estimated using Eqn 8. It can be shown in theory that under reasonable assumptions, this adaptive

optimal control scheme robustly stabilizes the closed loop system [25].

The free parameters of the model are the cost-function used to derive the LQG controller,

and the online learning rate γ. We fixed the cost-function to generate smooth movements with bell-

shape velocity profiles comparable with human movements qualitatively and allowed γ to vary in

order to capture the greater online adjustments observed across trials in Experiment 1. This

parameter was fixed for the simulations of the learning curve, and for the simulation of the via-point

experiment. The simulation of a learning paradigm (Fig. 5b) was performed by letting $\hat{A}_t$ evolve

during movements and storing the value obtained at the end of the trial for the beginning of the next

trial ($\gamma = 0.05$). The reduction in peak end-force observed in Experiment 1 was simulated by using

low or high values of γ during movement ($\gamma = 0.01$ and 0.1, respectively, Fig. 5c). Finally Experiment

3 was simulated as a series of null field trials interleaved with force-field trials with switching-off at

the via-point ($\gamma = 0.25$, Fig. 5c). Fig. 5c reports the force field trials, and the baseline trials preceded

by at least 3 baseline trials to washout the impact of each force field trial.

Figure Legends

Figure 1: Experiment 1, Behavior. a: Illustration of a successful trial in Experiment 1. Participants waited for the go signal, and were instructed to reach for the goal target within 0.6 s to 0.8 s of the go signal. Feedback about movement timing was provided to encourage participants to adjust their movement speed, but all trials were included in the dataset. b: Hand paths from all participants for the first force field trial (top) and the force field trial number 20 (bottom), were selected to illustrate the change in online control. Counter-clockwise (CCW) and clock-wise (CW) FF trials are depicted in blue and red, respectively. The black traces (baseline) are randomly selected trials from each participant. The baseline trials were performed in the null field, following at least 3 null-field trials. The maximum hand path deviation in the direction of the force field (Max.) and the maximum target overshoot in the opposite direction (Osh.) were extracted. c: Maximum hand path deviation for CCW and CW force fields across trials. The dashed traces illustrate the non-significant exponential fits. d: Maximum target overshoot across FF trials. Solid traces indicate significant exponential decay. Shaded blue and red areas around the means indicate the SEM across participants (n=14).

Figure 2. Improvement of Online Corrections. a: lateral force during the first (dashed/light) and last (solid/dark) trials. Shaded areas represent one SEM across participants (n=14) with the same color code as in Fig.1. The reduction in peak end force is highlighted with the black arrows. b: Peak end force across force field trials (mean±SEM). The solid traces show the significant exponential decay. c: Correlation between the lateral hand velocity and the force generated by the force field during the time interval corresponding to the gray rectangle in panel a (from 100ms to 700ms). The correlations were averaged across participants; the shaded area represents one SEM. The significant linear regression of the correlations across blocks is illustrated.

Figure 3: Experiment 2 Behavior. a: Average hand traces from Experiment 2 for trials with a via-point

with the same color code as in Experiment 1 (black: null-field, red: CW, blue: CCW). For all trials, the

second portion of the reach, from the via-point to the goal target, was performed without force field

(FF Off). b: Average x-coordinate as a function of time across participants (n=10). Shaded areas

represent one SEM across participants. The gray rectangle illustrates the portion of the movement

spent in the via-point approximately. Observe the displacement following the via-point in the

opposite direction as the perturbation-related movement. c: Illustration of the trial-by-trial

extraction of parameters on five randomly selected trials from one participant. The force field was

turned off when the hand speed dropped below 3 cm/s (dashed trace). The parameters extracted

were the lateral hand velocity at the moment of the hand speed minimum (squares), and the lateral

hand velocity at the moment of hand speed maximum (open discs). d: Lateral hand velocity at the

moment of hand speed minimum or at the moment of the second hand speed maximum averaged

across trials for each participant (gray dots), and averaged across participants (blue, black and red,

mean±SEM). Individual means across the three trial types were subtracted from the data for

illustration purpose. The two stars represent significant pair-wise differences at the level $P<0.01$ with

Bonferroni correction for multiple comparisons.

Figure 4: Learning across Blocks. a: Lateral hand velocity measured at the second peak hand speed

following CCW and CW force-fields (red and blue, respectively), and baseline trials (black) across

blocks. There were blocks in which participants did not succeed any force-filed trial, thus displays are

the mean±SEM across participants in each block with $8 \leq n \leq 10$. The insets highlight the first and last

block. b: Individual average across trials for each participant (gray dots), and group mean±SEM in the

first and 6th block (n = 9 in both cases). The two stars highlight significant difference based on paired

t-test ($P < 0.01$).

Figure 5. Model Simulations. a: Schematic illustration of an adaptive controller: sensory feedback is

used to estimate both the state of the system and the model, which in turn updates the controller

and estimator online. The solid arrows represent the inner loop correcting for state deviations and

the dashed arrows represent the outer loop correcting for model errors. b: Reproduction of a

standard learning experiment (hand paths and maximum lateral deviation across trials), with mirror-

image catch trials (ct), and unlearning following the catch trials (difference between ct-1 and ct+1).

The trial-by-trial changes were reproduced with online computations exclusively (inspired by

Thoroughman and Shadmehr [6]). c: Simulation of the results from Experiment 1: the adaptive

controller reduces the peak end force by increasing the online learning rate, which improves the

tuning of online corrections to the properties of the force field. d: Simulations of behavior from

Experiment 2, the force field was turned off at the via-point, and the second portion of the reach

displays an after effect (see Methods).

Acknowledgements

476   FC is supported by a grant from the Belgian Fund for Scientific Research (F.R.S.-FNRS,

477   Chercheur Qualifié, grant number 1.C.033.18F). This work was supported by grants from PRODEX and

478   by the Interuniversity Attraction Pole of the Belgian Science Policy (BELSPO).

References

Dayan, E. & Cohen, L. G. Neuroplasticity subserving motor skill learning. *Neuron* **72**, 443-454, doi:10.1016/j.neuron.2011.10.008 (2011).

Lackner, J. R. & DiZio, P. Rapid adaptation to Coriolis-force perturbations of arm trajectory. *Journal of Neurophysiology* **72**, 299-313 (1994).

Shadmehr, R. & Mussa-Ivaldi, F. A. Adaptive representation of dynamics during learning of a motor task. *Journal of Neuroscience* **14**, 3208-3224 (1994).

Krakauer, J. W., Ghilardi, M. F. & Ghez, C. Independent learning of internal models for kinematic and dynamic control of reaching. *Nature Neuroscience* **2**, 1026-1031 (1999).

Wagner, M. J. & Smith, M. A. Shared Internal Models for Feedforward and Feedback Control. *Journal of Neuroscience* **28**, 10663-10673, doi:10.1523/jneurosci.5479-07.2008 (2008).

Thoroughman, K. A. & Shadmehr, R. Learning of action through adaptive combination of motor primitives. *Nature* **407**, 742-747 (2000).

Singh, K. & Scott, S. H. A motor learning strategy reflects neural circuitry for limb control. *Nature Neuroscience* **6**, 399-403, doi:10.1038/nn1026 (2003).

Lackner, J. R. & DiZio, P. Motor control and learning in altered dynamic environments. *Current Opinion in Neurobiology* **15**, 653-659, doi:10.1016/j.conb.2005.10.012|ISSN 0959-4388 (2005).

Shadmehr, R., Smith, M. A. & Krakauer, J. W. Error Correction, Sensory Prediction, and Adaptation in Motor Control. *Annual Review of Neuroscience, Vol 33* **33**, 89-108 doi:10.1146/annurev-neuro-060909-153135 (2010).

Wolpert, D. M., Diedrichsen, J. & Flanagan, J. R. Principles of sensorimotor learning. *Nature Reviews Neuroscience* **12**, 739-751, doi:10.1038/nrn3112 (2011).

Smith, M. A., Ghazizadeh, A. & Shadmehr, R. Interacting adaptive processes with different timescales underlie short-term motor learning. *Plos Biology* **4**, 1035-1043, doi:10.1371/journal.pbio.0040179 (2006).

Kording, K. P., Tenenbaum, J. B. & Shadmehr, R. The dynamics of memory as a consequence of optimal adaptation to a changing body. *Nat Neurosci* **10**, 779-786, doi:10.1038/nn1901 (2007).

Gonzalez Castro, L. N., Hadjiosif, A. M., Hemphill, M. A. & Smith, M. A. Environmental consistency determines the rate of motor adaptation. *Curr Biol* **24**, 1050-1061, doi:10.1016/j.cub.2014.03.049 (2014).

Wei, K. & Koerding, K. Uncertainty of feedback and state estimation determines the speed of motor adaptation. *Frontiers in Computational Neuroscience* **4**, doi:10.3389/fncom.2010.00011 (2010).

Diedrichsen, J., White, O., Newman, D. & Lally, N. Use-dependent and error-based learning of motor behaviors. *J Neurosci* **30**, 5159-5166, doi:10.1523/JNEUROSCI.5406-09.2010 (2010).

Franklin, D. W. & Wolpert, D. M. Computational Mechanisms of Sensorimotor Control. *Neuron* **72**, 425-442, doi:10.1016/j.neuron.2011.10.006 (2011).

Burdet, E. *et al.* A method for measuring endpoint stiffness during multi-joint arm movements. *Journal of Biomechanics* **33**, 1705-1709, doi:10.1016/s0021-9290(00)00142-1 (2000).

Crevecoeur, F. & Scott, S. H. Beyond Muscles Stiffness: Importance of State Estimation to Account for Very Fast Motor Corrections. *PLoS Computational Biology* **10**, e1003869 (2014).

Kurtzer, I. L., Pruszynski, J. A. & Scott, S. H. Long-latency reflexes of the human arm reflect an internal model of limb dynamics. *Current Biology* **18**, 449-453, doi:10.1016/j.cub.2008.02.053 (2008).

Cluff, Y. & Scott, S. H. Rapid feedback responses correlate with reach adaptation and properties of novel upper limb loads. *Journal of Neuroscience* **33**, 15903-15914 (2013).

21 Crevecoeur, F. & Scott, S. H. Priors Engaged in Long-Latency Responses to Mechanical
Perturbations Suggest a Rapid Update in State Estimation. *Plos Computational Biology* **9**,
doi:10.1371/journal.pcbi.1003177 (2013).
22 Crevecoeur, F., Kurtzer, I. & Scott, S. H. Fast corrective responses are evoked by
perturbations approaching the natural variability of posture and movement tasks. *Journal of*
Neurophysiology **107**, 2821-2832, doi:10.1152/jn.00849.2011 (2012).
23 Fortney, K. & Tweed, D. B. Computational Advantages of Reverberating Loops for
Sensorimotor Learning. *Neural Computation* **24**, 611-634 (2012).
24 Ioannou, P. & J, S. *Robust Adaptive Control.* (Prentice Hall Inc., 1996).
25 Bitmead, R. R., Gevers, M. & Wertz, V. *Adaptive optimal control : the thinking man's GPC*.
(Prentice Hall, 1990).
26 Todorov, E. Stochastic optimal control and estimation methods adapted to the noise
characteristics of the sensorimotor system. *Neural Computation* **17**, 1084-1108 (2005).
27 Izawa, J., Rane, T., Donchin, O. & Shadmehr, R. Motor adaptation as a process of
reoptimization. *Journal of Neuroscience* **28**, 2883-2891, doi:10.1523/jneurosci.5359-07.2008
(2008).
28 Scott, S. H. A Functional Taxonomy of Bottom-Up Sensory Feedback Processing for Motor
Actions. *Trends Neurosci* **39**, 512-526, doi:10.1016/j.tins.2016.06.001 (2016).
29 Braun, D. A., Aertsen, A., Wolpert, D. M. & Mehring, C. Learning Optimal Adaptation
Strategies in Unpredictable Motor Tasks. *Journal of Neuroscience* **29**, 6472-6478,
doi:10.1523/jneurosci.3075-08.2009 (2009).
30 Scheidt, R. A., Dingwell, J. B. & Mussa-Ivaldi, F. A. Learning to move amid uncertainty. *Journal*
of Neurophysiology **86**, 971-985 (2001).
31 Ghahramani, Z. & Hinton, G. E. Parameter estimation for linear dynamical systems. *Technical*
Report CRG-TR-96-2, University of Toronto, Dept. of Computer Sceince (1996).
32 Astrom, K. J. *Introduction to stochastic control theory*. (Academic Press, 1970).
33 Schaal, S., Sternad, D., Osu, R. & Kawato, M. Rhythmic arm movement is not discrete. *Nat*
Neurosci **7**, 1136-1143 (2004).
34 Hirashima, M. & Nozaki, D. Distinct Motor Plans Form and Retrieve Distinct Motor Memories
for Physically Identical Movements. *Current Biology* **22**, 432-436,
doi:10.1016/j.cub.2012.01.042 (2012).
35 Howard, I. S., Ingram, J. N., Franklin, D. W. & Wolpert, D. M. Gone in 0.6 seconds: the
encoding of motor memories depends on recent sensorimotor states. *J Neurosci* **32**, 12756-
12768, doi:10.1523/JNEUROSCI.5909-11.2012 (2012).
36 Sheahan, H. R., Franklin, D. W. & Wolpert, D. M. Motor Planning, Not Execution, Separates
Motor Memories. *Neuron* **92**, 773-779, doi:10.1016/j.neuron.2016.10.017 (2016).
37 Hore, J. & Flament, D. Changes in motor cortex neural discharge associated with the
development of cerebellar limb ataxia. *J Neurophysiol* **60**, 1285-1302 (1988).
38 Pruszynski, J. A. *et al.* Primary motor cortex underlies multi-joint integration for fast feedback
control. *Nature* **478**, 387-390, doi:10.1038/nature10436 (2011).
39 Omrani, M., Murnaghan, C. D., Pruszynski, J. A. & Scott, S. H. Distributed task-specific
processing of somatosensory feedback for voluntary motor control. *Elife* **5**,
doi:10.7554/eLife.13141 (2016).
40 Kaufman, M. T., Churchland, M. M., Ryu, S. I. & Shenoy, K. V. Cortical activity in the null
space: permitting preparation without movement. *Nat Neurosci* **17**, 440-448,
doi:10.1038/nn.3643 (2014).
41 Stavisky, S. D., Kao, J. C., Ryu, S. I. & Shenoy, K. V. Motor Cortical Visuomotor Feedback
Activity Is Initially Isolated from Downstream Targets in Output-Null Neural State Space
Dimensions. *Neuron* **95**, 195-208 e199, doi:10.1016/j.neuron.2017.05.023 (2017).
42 Lakens, D. Calculating and reporting effect sizes to facilitate cumulative science: a practical
primer for t-tests and ANOVAs. *Front Psychol* **4**, 863, doi:10.3389/fpsyg.2013.00863 (2013).

43   Laird, N. M. & Ware, J. H. Random-effects models for longitudinal data. *Biometrics* **38**, 963-
580   974 (1982).

a

Time course of events:

1	**2**	**3**
Wait [2, 4] s	Go t = 0	Good if 0.6s ≤ t ≤ 0.8s

y
x

b

CCW
$F_x = -L\dot{y}$

CW
$F_x = +L\dot{y}$

FF trial #1

Max.
Osh.
Baseline

FF trial #20

5cm

c **Maximum**

x-coordinate [cm]
6
0
-3
-6
1   12   24
Trial

d **Overshoot**

NS
$P < 0.05$

x-coordinate [cm]
3
0
-3
1   12   24
Trial

Figure 1

Figure 2

Figure 3

Figure 4

a

Goal → Controller → Command

Controller ↔ $\hat{z}_t$ State E.

$\hat{A}_t$ Model E. ← Feedback

b

ct+1
ct-1
ct
ct: catch trial

5 cm

Max. deviation [cm]
ct-1
ct+1
ct
Trial Number

c

Experiment 1: CW

Peak End Force

x-force [N]

Online Rate:
Low
High

Time [s]

d

Experiment 2

x [cm]
FF Off

Time [s]

Lateral Vel. [cm/s]

@Min   @2Max

Figure 5