TITLE: Phosphatidylethanolamine made in the inner mitochondrial membrane is essential for

yeast cytochrome bc_1 complex function

AUTHORS: Elizabeth Calzada[1], J. Michael McCaffery[2], and Steven M. Claypool[1*]

AFFILIATIONS:

[1] Department of Physiology, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[2]Integrated Imaging Center, Department of Biology, Johns Hopkins University, Baltimore, MD

7
CORRESPONDING AUTHOR: Steven M. Claypool, Dept. of Physiology, Johns Hopkins School

of Medicine, 725 N. Wolfe St., Baltimore, MD 21225-2185. Tel. (410) 614-1786; Fax. (410) 955-

0461; E-mail: sclaypo1@jhmi.edu

ABSTRACT:

Of the four separate PE biosynthetic pathways in eukaryotes, one occurs in the

mitochondrial inner membrane (IM) and is executed by phosphatidylserine decarboxylase

(Psd1p). Deletion of Psd1, which is lethal in mice, compromises mitochondrial function. We

hypothesize that this reflects inefficient import of non-mitochondrial PE into the IM. To test this,

we re-wired PE metabolism in yeast by re-directing Psd1p to the outer mitochondrial membrane

or the endomembrane system. Our biochemical and functional analyses identified the IMS as

the greatest barrier for PE import and demonstrated that PE synthesis in the IM is critical for

cytochrome bc_1 complex (III) function. Importantly, mutations predicted to disrupt a conserved

PE-binding site in the complex III subunit, Qcr7p, impaired complex III activity similar to *PSD1*

deletion. Collectively, these data demonstrate that PE made in the IM by Psd1p is critical to

support the intrinsic functionality of complex III and establish one likely mechanism.

INTRODUCTION:

The evolutionary design of living cells has revealed redundancies in various metabolic

pathways to promote survival. Importantly however, the sequestration of enzymes and their

substrates into different membrane compartments allows for the enrichment and regulation of

metabolite synthesis in regions of the cell where they are essential. Such is the case for the

biosynthesis of the abundant membrane phospholipid, phosphatidylethanolamine (PE). In

mammalian cells PE is synthesized by four separate pathways, three of which localize to the

endoplasmic reticulum (ER) [1]. A final pathway is dependent on phosphatidylserine

decarboxylase (Psd1p) which is spatially isolated within the mitochondrial inner membrane (IM)

[2-5]. The predominant pathways for PE production include the Kennedy pathway, which

synthesizes PE through the stepwise conjugation of CDP-ethanolamine to diacylglycerol (DAG),

and the Psd pathway which utilizes phosphatidylserine (PS) as a substrate to generate PE [1]. In

multicellular organisms, preference for either the Kennedy pathway or the Psd pathway varies

between tissues and cell types [6]. Notably, deletion of either pathway is lethal during murine

embryogenesis, highlighting the importance of PE generation in both the ER and mitochondrial

compartments for development [7,8].

Conservation of the Psd pathway from bacteria to humans likely reflects the

endosymbiotic origin of mitochondria which in turn suggests that mitochondrial PS and PE

metabolism has been preserved to optimize mitochondrial performance [6]. Indeed, deletion of

phosphatidylserine decarboxylase (*Pisd* in mouse and humans and *PSD1* in yeast) in eukaryotic

cells decreases cellular growth, impairs oxidative phosphorylation, produces aberrant

mitochondrial morphology, and diminishes PE levels in cells and mitochondria [7,9-12]. The Psd

pathway is the predominant pathway for PE production in *Saccharomyces cerevisiae* and

produces up to 70% of PE in the cell [13]. In contrast to mammals, yeast additionally contain

Psd2p which localizes to either Golgi or endosomal compartments [5,14]. Deletion of *PSD2* alone

does not recapitulate any of the mitochondrial defects associated with loss of *PSD1*, further

emphasizing the importance of the mitochondrial PE biosynthetic pathway [5]. The combined

absence of *PSD1* and *PSD2* produces a strain that is auxotrophic for exogenous ethanolamine

supplementation which is used to generate PE through the Kennedy pathway. As the severity of

PE depletion can be genetically and metabolically adjusted in yeast, this organism has served

as an invaluable model for studying the importance of the mitochondrial Psd pathway for cellular

respiration.

Characterization of the Psd pathway over more than 50 years of research has revealed

detailed mechanistic insight into the biogenesis of the mature Psd1p enzyme [15]. In yeast, *PSD1*

is a nuclear encoded gene whose transcript is translated on cytosolic ribosomes which produce

a full-length zymogen that is catalytically inactive. Upon its mitochondrial import, two matrix

metalloproteases sequentially process the N-terminus of Psd1p which then undergoes a third

processing event that is executed internally by the enzyme itself at the conserved C-terminal

LGST motif [2,3]. This unique autocatalytic processing event severs Psd1p into a large

membrane-anchored β subunit and a small α subunit and is performed by a self-contained

catalytic triad typical of serine proteases [16,17]. Self-processing endows the small α subunit with

an N-terminal pyruvoyl prosthetic group that is crucial for its decarboxylase activity. Post-

cleavage, the two Psd1p subunits remain non-covalently associated within the inner membrane

where they decarboxylate PS to PE [17].

The substrate of Psd1p, PS, is synthesized on the mitochondrial-associated membrane

(MAM) of the ER by phosphatidylserine synthase (Cho1p) [18]. Thus, the amphipathic PS must be

imported from the ER and traverse through two aqueous compartments, the cytosol and the

mitochondrial intermembrane space (IMS), to reach Psd1p at the IM [19]. While the redundant

roles played by ER-mitochondria membrane tethers in PS mitochondrial import continue to be

elucidated [20-24], recent evidence in yeast indicates that Mic60p, a subunit of the mitochondrial

contact site and cristae organizing system (MICOS), works in conjunction with the soluble lipid

carrier, Ups2p, to expose PS for Psd1p decarboxylation in the IMS [25,26]. Whether or not a

parallel pathway exists for PE import into the IM remains unclear. The lethal consequence of

pisd deletion in mice and the failure of supplemental ethanolamine to rescue the respiratory

defects of *psd1Δ* yeast were taken as evidence that PE made outside of the mitochondrion is

unable to compensate for the absence of Psd1p [1,7,11,27]. However, it was recently reported that

PE made by the Kennedy pathway can in fact restore the impaired oxidative phosphorylation of

psd1Δ yeast suggesting that extra-mitochondrial PE can compensate for the absence of the Psd

pathway [28].

Here, we sought to determine if either the cytosol and/or the IMS is a barrier that

prevents PE made outside of the IM from rescuing the impaired oxidative phosphorylation that

occurs in *S. cerevisiae* lacking Psd1p. Previously, we generated a chimeric Psd1p protein, ER-

Psd1p, which is targeted to the endosomal compartment and is catalytically active [29]. In the

current study, we further characterized the cell biology of ER-Psd1 yeast, together with an OM-

targeted chimeric Psd1p (OM-Psd1p), to test if the cytosol, IMS, or both, are barriers that

prevent non-mitochondrially produced PE from functionally rescuing the absence of PE made in

the IM. Alongside strains expressing these re-directed Psd1p constructs, we compared the

mitochondrial function of *psd1Δpsd2Δ* yeast grown in the non-fermentable carbon source,

lactate, with or without exogenous ethanolamine supplementation, to evaluate the ability of the

ER-localized Kennedy pathway to support mitochondrial function. In contrast to cytochrome *c*

oxidase (complex IV)*,* cytochrome bc_1 respiratory complex (complex III) activity was impaired in

the absence of IM-synthesized PE even after supplementation with ethanolamine. Intriguingly,

the positive effect of ethanolamine on complex IV activity reflected an increase in mitochondrial

levels of cardiolipin (CL), and not PE. Notably, the absence of Cho1p disrupted complex III and

IV activities without altering Psd1p steady state levels. These results highlight a crucial role of

PE for complex III function and suggest that the IMS is the greatest obstacle preventing extra-

mitochondrially produced PE from functionally substituting for IM-fabricated PE. Structures of

yeast and human complex III have each revealed a bound PE in close proximity to the complex

III subunit, Qcr7p [30,31]. Strikingly, mutations predicted to disrupt PE-binding by Qcr7p impaired

complex III activity to a similar level as in the absence of Psd1p. Altogether, we demonstrate

that PE made in the IM by Psd1p is critical to support full complex III function and provide the

first molecular evidence of the functional importance of a PE-binding site in complex III.

RESULTS

The impaired respiratory growth of psd1Δ yeast is not rescued by PE synthesis through the

Kennedy pathway

The ability of supplemental

ethanolamine to rescue the respiratory

growth defect of *psd1Δ* yeast has

been reported by some groups [28] but

not others [1,11,27]. This is an important

controversy to resolve. If extra-

mitochondrially produced PE can

replace PE normally made in the IM,

then this would imply that the

mitochondrial Psd1 pathway *per se*, is

not necessary for mitochondrial

function. By extension, it would

suggest that mechanisms to move

extra-mitochondrially produced PE into

Fig 1. Ethanolamine supplementation fails to rescue the growth defect of *psd1Δ* on respiratory medium in 4 separate yeast strains. (A, B) The indicated strains were pre-cultured at 30°C in YPD and spotted onto (A) synthetic complete dextrose (SCD) or ethanol-glycerol (SCEG) medium +/- 2mM ethanolamine (+E) or (B) SCEG +/- 10mM ethanolamine and incubated at 30°C for 3 days.

mitochondria are robust enough to compensate for the absence of Psd1p. Therefore, we tested

the growth of wildtype (WT), *psd1Δ*, *psd2Δ,* and *psd1Δpsd2Δ* in synthetic complete ethanol-

glycerol (SCEG) medium with or without 2mM ethanolamine supplementation (Fig 1A).

Consistent with previous findings [1,11,27], we found that exogenous ethanolamine

supplementation restored respiratory growth of *psd1Δpsd2Δ* yeast to *psd1Δ* levels but failed to

fully restore the respiratory defect associated with *psd1Δ* yeast. Importantly, this basic result

was confirmed in *psd1Δ* yeast made in three additional yeast strain backgrounds (Fig 1B, note

that the supplemental ethanolamine concentration was increased to 10mM), although subtle

differences between the strains were noted. Overall, these findings indicate that the Kennedy

pathway cannot fully compensate for the absence of Psd1p function, potentially due to

inefficient trafficking of PE from the ER to mitochondrial OM and/or from the OM to the IM.

Validation of constructs that re-direct Psd1p to OM or ER membranes

To interrogate whether the cytosol and/or the IMS is a barrier that prevents extra-

mitochondrially produced PE from replacing PE normally made in the IM, we generated chimeric

Psd1p constructs that are localized to either the ER or OM to redirect PS and PE metabolism.

As depicted in Fig 2A, these two constructs, and the WT IM-localized Psd1p control (referred to

as IM-Psd1p to distinguish it from strains expressing endogenous Psd1p), contain a C-terminal

3XFLAG tag to track autocatalytic function of these chimeras by immunodetecting the released

Psd1p α subunit. To re-direct Psd1p to the mitochondrial OM, the mitochondrial targeting

sequence and transmembrane domain of Psd1p were replaced by the equivalent domains of

the single-pass OM resident protein, Tom20p. ER-Psd1p, which is directed to the secretory

pathway, was generated by replacing the mitochondrial targeting sequence of Psd1p with the

NH_2-terminal signal sequence of carboxypeptidase Y (CPY) [29,32]. The ER-Psd1p construct

additionally contains an *N*-glycosylation signal immediately downstream of the CPY leader

sequence to track its topology. OM-Psd1p and ER-Psd1p each yielded mature β and α subunits

as detected by immunoblot of yeast cell extracts demonstrating that self-processing of Psd1p

was not impaired by these modifications to its NH_2-terminus (Fig 2B). All three integrated

constructs were similarly expressed relative to each other and over-expressed compared to

endogenous Psd1p. Importantly, both OM-Psd1p and ER-Psd1p rescued the ethanolamine

auxotrophy of *psd1Δpsd2Δ* yeast indicating that these constructs are fully functional and

capable of generating levels of PE necessary for cellular growth (Fig 2C).

Previously, localization of ER-Psd1p was established by virtue of its enrichment in the

40,000 x *g* pellet (P40) after subcellular fractionation by gravity centrifugation and its sensitivity

to endoglycosidase H, which revealed a mobility shift following SDS-PAGE post-treatment [29]. To

confirm the OM localization of OM-Psd1p, its protease accessibility was determined in intact

mitochondria, OM-ruptured mitoplasts, and detergent-solubilized mitochondria and compared to

IM-Psd1p. Protease treatment of intact mitochondria expressing IM-Psd1p showed that IM-

Psd1p, like the IM control Tim54p, was protected against degradation (Fig 2D). In contrast,

Fig 2. OM-Psd1p and ER-Psd1p constructs are functional and localize to the OM and ER, respectfully. (A) Schematic of IM-Psd1p, OM-Psd1p, and ER-Psd1p. All three constructs contain a 3XFLAG tag at the C-terminus (yellow). The Tom20 residues (1-100) that replace the mitochondrial targeting sequence (MT) and transmembrane (TM) domain of IM-Psd1p (green) are shown for OM-Psd1p (blue), and the carboxypeptidase Y signal sequence (residues 1-37) as well as an NXS motif are indicated for ER-Psd1p (purple). (B) The β and α subunits of Psd1p were detected in yeast whole cell extracts of the indicated strains by immunoblot. Tom70p served as a loading control. (C) The indicated strains were spotted onto synthetic complete dextrose (SCD) medium +/- 2mM ethanolamine (+E) and incubated at 30°C for 4 days. (D) Protease protection assay in intact mitochondria (Mito), osmotically ruptured mitochondria (MP), or deoxycholate-solubilized mitochondria (Det.). Following incubation -/+ 100μg proteinase K (Prot. K) for 30 minutes, samples were collected, resolved by SDS-PAGE and immunoblotted for Psd1p (β and α subunits), and the mitochondrial compartment-specific markers Tom70p (OM), Tim54p (IM), and Abf2p (matrix). The exposures of the indicated antibodies were linearly adjusted relative to the set of samples for each strain, and are boxed individually to reflect this. Three biological replicates were performed using two separate batches of isolated mitochondria from each strain. (E) Illustration indicating the topology of (1) IM-Psd1p, (2) OM-Psd1p, and (3) ER-Psd1p.

similar to the OM control Tom70p, OM-Psd1p was completely degraded in intact mitochondria

verifying that it was successfully re-localized to the OM with the bulk of the enzyme facing the

cytosol. As expected given the presence of inter-organelle contact sites, a proportion of ER-

Psd1p co-fractionated with crude mitochondria (Supplementary Fig 1) and demonstrated

protease-sensitivity in intact mitochondria (Fig 2D), a topology that is consistent with its N-

glycosylation status (Fig 2E). Thus, a portion of ER-Psd1p is retained in the ER and/or resides

in an endosomal compartment that is co-purified with mitochondria.

OM-Psd1p and ER-Psd1p generate levels of PE in mitochondrial membranes that exceed WT

Next, the lipid content of cellular and mitochondrial membranes was assessed in WT,

psd1Δ, *psd2Δ*, *psd1Δpsd2Δ*, IM-Psd1, OM-Psd1 and ER-Psd1 yeast (Fig 3 and supplementary

Fig 2). The absence of Psd1p resulted in reduced levels of cellular and mitochondrial PE; the

combined absence of Psd1p and Psd2p resulted in an additive effect on the steady state

abundance of PE (Fig 3A and 3G). In *psd1Δ* and *psd1Δpsd2Δ* membranes, the levels of

phosphatidylinositol (PI) was increased (Figure 3B and 3H) and CL decreased (Fig 3C and 3I),

consistent with previous reports [9,28,33]. A compensatory increase in PS was notably absent in

these yeast strains (Fig 3D and 3J). While mitochondrial PE levels were modestly decreased in

the absence of Psd2p (Fig 3G), this decrease failed to result in a respiratory growth defect (Fig

1). Combined, these results indicate that Psd2p contributes to the pool of PE associated with

mitochondria which is nonetheless unable to functionally replace PE made by Psd1p.

Interestingly, OM-Psd1 and ER-Psd1 yeast contained significantly higher relative

amounts of PE than IM-Psd1 in both cellular and mitochondrial membranes (Fig 3A and 3G). As

IM-Psd1p, OM-Psd1p, and ER-Psd1p are similarly over-expressed, this suggests that OM-

Psd1p and ER-Psd1p, which are either present in compartments where PS is made (ER-Psd1p)

or through which PS must traffic to reach IM-Psd1p (OM-Psd1p), have short-circuited normal

mitochondrial PE metabolism. In support of this interpretation, the steady state levels of Cho1p

(Supplementary Fig 3) and mitochondrial PS (Fig 3J) were increased by OM-Psd1p but not ER-

Fig 3. OM-Psd1 and ER-Psd1 contain robust levels of PE in both cellular and mitochondrial membranes. (A-F) Cellular and (G-L) mitochondrial phospholipids from the indicated strains were labeled overnight with $^{32}P_i$ and separated by TLC. All graphs show the mean ± S.E.M. for n=6 biological replicates. Significant differences compared to WT were calculated by one-way ANOVA. (M) Key for symbols used for statistical analysis interpretation by one-way ANOVA when comparing samples *versus* WT (*), *psd1Δpsd2Δ* (#), or IM-Psd1 ($).

Psd1p, consistent with a mitochondrial trafficking requirement for the former, but not the latter,

chimera. Of note, OM-Psd1p and ER-Psd1p both normalized the absolute amount of Cho1p and

its phosphorylated pool which were significantly increased in *psd1Δpsd2Δ* yeast grown in

respiratory conditions (Supplementary Fig 3A, and 3D-F). As Cho1p phosphorylation has been

shown to both inhibit enzyme activity and stabilize the polypeptide [34], these changes suggest

that feedback mechanisms are activated by the severe PE depletion present in *psd1Δpsd2Δ*

yeast. The relative abundance of phosphatidylcholine (PC) was reduced in OM-Psd1 and ER-

Psd1 (Fig 3E and 3K), and although CL levels were significantly increased compared to

psd1Δpsd2Δ mitochondria, they were still lower than in WT (Fig 3C and 3I). The reduced levels

of PC in OM-Psd1 and ER-Psd1 is notable as it might have been predicted that an increased

production of PE would have resulted in augmented PC synthesis by Pem1p and Pem2p,

methyltransferases which reside in the ER and convert PE to PC [35]. Importantly, the steady

state level of Kar2p, the yeast equivalent of the Hsp70 chaperone BiP [36], was not increased in

OM-Psd1 or ER-Psd1 (Supplementary Fig 3G), demonstrating that their altered membrane

compositions did not induce ER stress. In contrast, Kar2p was significantly elevated in both

psd1Δpsd2Δ and *cho1Δ* strains.

Overall, the increased levels of PE detected in OM-Psd1 and ER-Psd1 suggests that

both chimeras have increased access to substrate compared to IM-Psd1. This conclusion is

bolstered both by the fact that overexpressed IM-Psd1p only restored PE to WT levels (Fig 3A

and 3G), as well as results from a prior study that used a plasmid-based overexpression

system [1]. Thus, access to substrate is a major regulatory component that determines Psd1p

activity.

A limitation of the radiolabeling-based phospholipid analysis is that it utilized crude

mitochondria isolated after physical disruption of intact yeast with glass beads. As such,

phospholipid analyses were additionally performed using sucrose gradient purified mitochondria

derived from non-radiolabeled yeast cultures. Sucrose purification resulted in mitochondria that

were enriched in mitochondrial proteins compared to crude mitochondria (Supplementary Fig

4A). Subsequent phospholipid analyses of the sucrose purified mitochondria provided a profile

that was consistent with results derived using crude mitochondria (Fig 3), with one notable

exception: the PE levels were roughly equal in OM-Psd1 and ER-Psd1 sucrose purified

mitochondria (Supplementary Fig 4E). The reduced amounts of mitochondrial PE in ER-Psd1

following sucrose purification likely reflects their increased purity and provides evidence that PE

made in the ER can access mitochondrial membranes. Importantly, there were no dramatic

differences in the mitochondrial architectures of strains with elevated or decreased PE levels

(Supplementary Fig 5), consistent with previous observations in *psd1Δ* yeast [28]. Overall, re-

routing Psd1p to either the OM or ER results in a robust increase in cellular and mitochondrial

PE levels which may or may not reach the mitochondrial IM.

OM-Psd1 and ER-Psd1 phenocopy the respiratory defect of psd1Δ yeast

225   We initially evaluated oxidative phosphorylation in these strains by determining their

growth on synthetic mediums containing dextrose +/- 2mM ethanolamine, lactate, or ethanol-

glycerol +/- 2mM ethanolamine (Fig 4A). Compared to IM-Psd1p, OM-Psd1p and ER-Psd1p

only partially improved growth of *psd1Δpsd2Δ* yeast on respiratory carbon sources (lactate and

ethanol-glycerol). In fact, they supported respiratory growth that was similar to (ER-Psd1p) or

slightly better than (OM-Psd1p) *psd1Δ* yeast, but still significantly reduced compared to the WT

strain. Interestingly, respiratory growth was slightly better for *psd1Δpsd2Δ* yeast expressing

OM-Psd1p *versus* ER-Psd1p. The respiratory phenotype of OM-Psd1 and ER-Psd1 yeast

suggests that PE made in either the OM or ER cannot functionally replace PE produced in the

IM. Indeed, ethanolamine supplementation did not further improve the respiratory growth of

either OM-Psd1 or ER-Psd1 yeast (Fig 4A).

236   To directly assess OXPHOS capacity in these strains, oxygen consumption was

monitored in isolated mitochondria using an O_2 electrode after the addition of ADP and

ascorbate tetramethyl-*p*-phenyldiamine (TMPD) which promotes proton pumping by complex IV

(Fig 4B-4E). To determine maximal respiratory capacity, carbonyl cyanide *m*-chlorophenyl

hydrazine (CCCP) was added after ADP was depleted and mitochondrial respiration had

returned from the ADP-stimulated (State 3) to the resting (State 4) respiratory rate. Mitochondria

lacking Psd1p, but not Psd2p, had reduced maximal uncoupled respiratory rate compared to

WT mitochondria. Even though *psd2Δ* mitochondria consumed O_2 like WT, the combined

absence of Psd1p and Psd2p caused a more severe respiratory defect than was observed

when only Psd1p was missing. This indicates that in the absence of Psd1p, PE made by Psd2p

has some capacity to compensate by supporting limited complex IV activity. Interestingly, the

uncoupled respiratory rate for OM-Psd1 and ER-Psd1 was significantly improved compared to

psd1Δpsd2Δ mitochondria but still impaired relative to IM-Psd1 (Fig 4B). The respiratory control

Fig 4. OM-Psd1 and ER-Psd1 OXPHOS function phenocopies *psd1Δ*. (A) The indicated strains were spotted and incubated at 30°C for 2 days on SCD +/- 2mM ethanolamine (+E) and for 5 days on rich lactate (RL), SC lactate (SC-LAC), and SCEG +/- 2mM ethanolamine (+E). (B-E) O_2 consumption measurements from mitochondria isolated from yeast grown in rich lactate using ascorbate-TMPD as a substrate. (B) The maximal respiratory rate was recorded after the addition of CCCP, (C) state 3 respiration was assessed after addition of ADP, and (D) state 4 respiration was recorded following ADP depletion. (E) The respiratory control ratio (RCR) is calculated by dividing state 3 by state 4 respiratory rates. Analysis *versus* WT (*), *psd1Δpsd2Δ* (#), or IM-Psd1 ($) was performed by one-way ANOVA $\pm$ S.E.M. for n=4.

ratio (RCR) is an indication of how well respiration is coupled to ATP synthesis and is calculated

by dividing the ADP-stimulated respiration rate by the resting respiration rate (State 3/State 4).

RCR ratios were significantly decreased in *psd1Δ* and *psd1Δpsd2Δ* mitochondria but

perplexingly, OM-Psd1 and ER-Psd1 displayed a ratio similar to WT and IM-Psd1 levels (Fig

4C). For *psd1Δpsd2Δ* yeast expressing OM-Psd1p and ER-Psd1p, the normal RCR stems from

the fact that they improved State 3 O_2 consumption more than State 4. Similar to Psd2p in the

context of *psd1Δ* yeast, OM-Psd1p and ER-Psd1p significantly improved *psd1Δpsd2Δ*

respiratory rates to roughly *psd1Δ* levels, indicating that extra-mitochondrial PE can indeed play

an important role in enhancing respiration. Combined, these results suggest that defective

complex IV function may contribute to the reduced respiratory growth observed when Psd1p is

absent from the IM.

Complex IV function in psd1Δpsd2Δ mitochondria is rescued by ER-Psd1p

When grown in dextrose, *psd1Δ* and *psd1Δpsd2Δ* yeast lose their mitochondrial

genome, visualized by formation of petite colonies, at a high frequency [1,28,37]. Due to this, we

harvested mitochondria from cultures grown in rich lactate to select for growth of respiratory

competent cells. As expected, mitochondrial DNA (mtDNA) levels were equivalent between

strains in these growth conditions (Fig 5A; ρ^0 yeast, which are devoid of mtDNA, served as a

negative control). Next, isolated complex IV activity measurements were recorded by monitoring

the rate of oxidation of reduced cytochrome *c* in mitochondria solubilized in 0.5% *n*-Dodecyl-β-

D-maltoside (DDM). Complex IV activity was significantly decreased in *psd1Δ*, *psd1Δpsd2Δ* and

OM-Psd1 mitochondria but surprisingly, ER-Psd1 retained WT function (Fig 5B). To determine if

the different complex IV activities associated with OM-Psd1 and ER-Psd1 mitochondria

reflected changes in its expression, we analyzed the steady state amounts of both nuclear and

mtDNA encoded subunits of complex IV (Fig 5C). While the levels of the mtDNA-encoded

subunit, Cox2p, was significantly decreased in *psd1Δpsd2Δ*and OM-Psd1 strains, the steady

Fig 5. Complex III and IV activities are impaired when Psd1p is absent in the IM. (A) mtDNA was isolated from the indicated strains, normalized, and quantified by qPCR. Analysis was performed by one-way ANOVA ± S.E.M. for n=3. (B) Spectrophotometric assay following the oxidation of cytochrome *c* at 550nM in isolated mitochondria solubilized in 0.5% (w/v) DDM. Analysis *versus* WT (*), *psd1Δpsd2Δ* (#), or IM-Psd1 ($) was performed by one-way ANOVA ± S.E.M. for n=6 or n=3 for ρ⁰. P values for decreases that didn't achieve significance are reported in red and were analyzed by student *t*-test *versus* WT. (C) Mitochondria from the indicated strains were immunoblotted for subunits of complex III (CIII), complex IV (CIV), complex V (CV), and the Coq synthome, cytochrome *c*, and markers of each mitochondrial compartment. (D, E) Blue native-PAGE analysis of respiratory supercomplexes (RSCs) using mitochondrial extracts solubilized in 1.5% (w/v) digitonin. (D) Complex IV assembly was monitored by immunoblot against the nuclear-encoded subunit Cox4p and (E) Complex III assembly was monitored by immunoblot against the nuclear-encoded subunit Rip1p. Mitochondria lacking CL (*crd1Δ*) were used as a positive control for RSC destabilization[53]. (F) Spectrophotometric assay following thereduction of cytochrome *c* at 550nM in isolated mitochondria solubilized in 0.5% (w/v) DDM. Analysis *versus* WT (*), *psd1Δpsd2Δ* (#), or IM-Psd1 ($) was performed by one-way ANOVA ± S.E.M for n=6 or n=3 for ρ⁰. P values for decreases that didn't achieve significance are reported in red and were analyzed by student *t*-test versus WT.

state abundance of the two additional mtDNA-encoded subunits, Cox1p and Cox3p, were not

significantly changed and the amount of a constituent encoded in the nucleus, Cox4p, was

decreased in OM-Psd1 but not *psd1Δpsd2Δ* (Supplementary Fig 6). Further, blue native-PAGE

analyses indicated that complex IV assembly into respiratory supercomplexes that consist of a

complex III dimer affiliated with either one or two complex IV monomers [38], was normal

regardless of the absence of *PSD1* or *PSD2*, singly or in combination, as reported by others [28],

or whether Psd1p was expressed in the IM, OM, or ER (Fig 5D).

Cytochrome bc_1 complex function is impaired when Psd1p is absent from the IM

The ability of ER-Psd1p, but not OM-Psd1p, to rescue complex IV activity to WT levels

was surprising given that neither chimeric construct restored respiratory growth of the

psd1Δpsd2Δ strain to this degree (Fig 4A). Therefore, we postulated that the incomplete

respiratory growth rescue of OM-Psd1 and ER-Psd1 could reflect a defect at the level of

complex III. Indeed, complex III activity was reduced in *psd1Δ* and significantly decreased in

psd1Δpsd2Δ mitochondrial extracts (Fig 5F). While ER-Psd1p significantly improved complex IV

function over that detected in *psd1Δpsd2Δ* mitochondria, neither ER-Psd1p or OM-Psd1p

supported complex III activity that resembled WT. The reduced complex III activity in *psd1Δ*,

psd1Δpsd2Δ, and *psd1Δpsd2Δ* expressing OM-Psd1p or ER-Psd1p did not reflect alterations in

the steady state abundance of its subunits (Fig 5C, Cor2p, Rip1p, Qcr6p, and Qcr7p, quantified

in Supplementary Fig 6) or its assembly into supercomplexes (Fig 5E) although there was

proportionately more of the smaller supercomplex ($III_2IV > III_2IV_2$) detected in *psd1Δpsd2Δ*

mitochondrial extracts. Furthermore, the steady state levels of the complex III electron acceptor

cytochrome *c* were normal (Fig 5C and Supplementary Fig 6M). Similarly, subunits of the

coenzyme Q (CoQ) synthome, a macromolecular complex that catalyzes the synthesis of the

complex III electron donor CoQ [39], were equal with one exception (Fig 5C and Supplementary

Fig 7A). In *psd1Δpsd2Δ* mitochondria, Coq1p was increased which could represent an attempt

to diminish membrane stress as observed in bacteria [40]. Moreover, CoQ_6 supplementation,

which is capable of rescuing strains with reduced CoQ biosynthesis [39], failed to improve

respiratory growth of *psd1Δ* or *psd1Δpsd2Δ* yeast (Supplementary Fig 8). Lastly, *psd1Δ*,

psd1Δpsd2Δ, OM-Psd1, and ER-Psd1 respiratory growth was not further impaired in medium

lacking *para*-amino benzoic acid (Supplementary Fig 9), a molecule that can be used to produce

CoQ by a secondary pathway [41]. In total, our results demonstrate that CoQ is not limiting for

respiratory function in any of these strains. As such, they favor the hypothesis that the impaired

complex III activity of *psd1Δ*, *psd1Δpsd2Δ*, OM-Psd1, and ER-Psd1 is intrinsic to the multi-

subunit holoenzyme itself.

PE made by the Kennedy pathway rescues complex IV but not complex III function

As ER-Psd1p restores complex IV, but not complex III, function, we isolated

psd1Δpsd2Δ mitochondria from cultures grown in rich lactate, rich lactate + 2mM choline, and

rich lactate +2mM ethanolamine, to evaluate the potential impact of PC and PE generation by

the Kennedy pathways on complex IV activity (Fig 6A-B). Similar to ER-Psd1, supplementation

of *psd1Δpsd2Δ* with ethanolamine, but not choline, restored complex IV (Fig 6A), but not

complex III (Fig 6B), function to WT levels. Combined, our results indicate that PE, but not PC,

made in the ER by the Kennedy pathway (Fig 6A) or in the endosomal system by either Psd2p

or ER-Psd1p (Fig 5B), can significantly rescue the severe complex IV dysfunction that occurs in

psd1Δpsd2Δ yeast. Surprisingly, when we evaluated the phospholipid composition of

psd1Δpsd2Δ yeast supplemented with choline or ethanolamine, we found that the inclusion of

ethanolamine did not alter mitochondrial PE levels (Fig 6C-D) and only modestly and yet

significantly increased cellular PE abundance (Fig 6E-F), consistent with the slight increases

observed in [1,33] but not [28]. Intriguingly, ethanolamine, but not choline, supplementation

significantly increased CL in *psd1Δpsd2Δ* yeast to *psd1Δ* levels (Fig 6C and 6G). Significant

Fig 6. The Kennedy Pathway for PE synthesis can rescue the activity of complex IV but not complex III. Spectrophotometric assay following the (A) oxidation and (B) reduction of cytochrome *c* at 550nM in isolated mitochondria solubilized in 0.5% (w/v) DDM. Analysis *versus* WT (*) or *psd1Δpsd2Δ* (#) was performed by one-way ANOVA $\pm$ S.E.M. for n=6 or n=4 for ρ^0. P values for decreases that didn't achieve significance are reported in red and were analyzed by student *t*-test. (C-G) Mitochondrial phospholipids from the indicated strains were labeled overnight with $^{32}P_i$ and separated by TLC. (C) Representative TLC plate for mitochondrial $^{32}P_i$ lipids. (D-G) Graphs representing quantitation of (D) mitochondrial PE and (G) CL levels show the mean $\pm$ S.E.M. for n=6 biological replicates. Significant differences compared to (*) WT or (#) *psd1Δpsd2Δ* were calculated by one-way ANOVA. (E) Representative TLC plate for cellular $^{32}P_i$ lipids. (F) Quantitation of cellular PE levels representing the mean $\pm$ S.E.M. for n=6 biological replicates. Significant differences compared to (*) WT or (#) *psd1Δpsd2Δ* were calculated by student *t*-test.

changes in the abundance of other phospholipid species in *psd1Δpsd2Δ* yeast supplemented

with either choline or ethanolamine were not observed (Supplementary Fig 10). As such, these

results indicate that the Kennedy Pathway for PE but not PC production, is metabolically-linked

to CL biosynthesis and/or stability. Moreover, these results suggest that the ability of

ethanolamine to improve complex IV activity in *psd1Δpsd2Δ* yeast is due to its unanticipated

capacity to increase CL levels, a phospholipid known to be important for complex IV function [42].

These findings further underscore that PE made within the IM is necessary for the full activity of

Fig 7

Fig 7. Depletion of PE through deletion of Cho1p impairs complex III and complex IV activities. (A) Metabolic pathways tied to PS synthesis by Cho1p in yeast. (B) Detection of the β subunit of Psd1p and Cho1p expression were verified in yeast whole cell extracts of the indicated strains by immunoblot. Kgd1p served as a loading control. *Cho1p, phosphorylated Cho1p. (C) The indicated strains were spotted onto synthetic complete dextrose (SCD) medium +/- 2mM ethanolamine (+E) and incubated at 30°C for 2 days. (D-J) Mitochondrial phospholipids from the indicated strains were labeled overnight with $^{32}P_i$, separated by TLC, and quantitated by phosphoimaging. All graphs show the mean $\pm$ S.E.M. for n=6 biological replicates. Significant differences compared to (*) WT or (#) *psd1Δpsd2Δ* were calculated by one-way ANOVA. (K) The indicated strains were spotted and incubated at 30°C for 2 days on YPD and for 3 days on rich lactate (RL), and SCEG +/- 2mM ethanolamine (+E). (L,M) Spectrophotometric assay following the (L) reduction and (M) oxidation of cytochrome *c* at 550nM in isolated mitochondria solubilized in 0.5% (w/v) DDM. Analysis *versus* WT by one-way ANOVA $\pm$ S.E.M. for n=6 or n=3 for ρ⁰. (N) Steady state expression of mitochondrial proteins in mitochondria isolated from the indicated strains.

complex III which is otherwise normally expressed, fully assembled, and not limited by the

amount of either of its mobile electron carriers.

Non-enzymatic functions of Psd1p independent of PE biosynthesis are not required for complex

III activity

Cho1p mediates the production of PS in the MAM of the ER through conjugation of free

serine with CDP-DAG [18]. In yeast, this feeds into both the Psd1p and Psd2p PS decarboxylation

pathways (Fig 7A). We generated deletion strains of Cho1p in the WT and *psd1Δpsd2Δ* (labeled

cho1Δ1Δ2Δ in Fig 7) genetic backgrounds as a way to deplete mitochondrial PS/PE levels while

preserving Psd1p expression. Consistent with previous data, deletion of Cho1p did not affect

Psd1p accumulation or maturation [29] (Fig 7B) and resulted in an ethanolamine auxotrophy [43]

(Fig 7C). As anticipated, PS and PE levels were drastically reduced in strains lacking Cho1p

(Fig 7D-F). In comparison to *psd1Δpsd2Δ*, *cho1Δpsd1Δpsd2Δ* yeast had a significant increase

in CL and PI levels (Fig 7G and 7H), the former of which may be associated with its enhanced

respiratory growth (Fig 7K). Deletion of *cho1Δ* in the *psd1Δpsd2Δ* background restored PC to

WT levels (Fig 7J) but failed to increase the levels of the CL precursor PA (Fig 7I). Growth of

cho1Δ was decreased compared to WT in YPD and rich lactate medium and similar to

psd1Δpsd2Δ in SCEG supplemented with ethanolamine (Fig 7K). Notably, the activities of

complex III (Fig 7L) and IV (Fig 7M) were reduced in the absence of Cho1p. Since Psd1p and

essential subunits of complexes III and IV were expressed normally in the absence of Cho1p,

singly or in combination with Psd1p and Psd2p (Fig 7N), our combined results directly implicate

mitochondrial PE depletion as the cause for the reduced respiratory function of *psd1Δ*,

psd1Δpsd2Δ, and *cho1Δ* yeast.

Glu82 of Qcr7p may coordinate the headgroup of a functionally important PE molecule

associated with complex III

PE was identified in crystal structures of the yeast and mammalian cytochrome bc_1

complex in association with the essential mtDNA-encoded catalytic subunit cytochrome *b*

(Cob1p) as well as the nuclear-encoded subunit Qcr7p [30,31]. Qcr7p is associated with the matrix-

facing surface of Cob1p and it is postulated that hydrogen bonding interactions between the

headgroup of PE and Glu82 of Qcr7p may help position the complex vertically within the bilayer

(Fig 8A). To test the importance of this residue in forming hydrogen bonds with the amine group

of PE, we introduced a charge reversal at this position by mutating Glu82 to Arg and also

created a strain expressing Asp82 to test the potential effect of shortening the distance of this

interaction. Importantly, the amount of Qcr7p^{E82R} or Qcr7p^{E82D} detected in cell and mitochondrial

lysates (Fig 8B and 8C) was similar to WT indicating that neither mutation compromised Qcr7p

stability (the Qcr7p^{E82R} variant was consistently upshifted compared to WT following SDS-

PAGE). Further, Qcr7p^{E82R} and Qcr7p^{E82D} supported respiratory growth in rich or minimal

medium (Fig 8D). Despite being sufficiently functional to promote respiratory growth, complex III

activity was significantly decreased for Qcr7p^{E82R} and Qcr7p^{E82D} to a similar degree as when

Psd1p is missing (Fig 8E). Surprisingly, complex IV activity was also decreased in Qcr7p^{E82R} but

not Qcr7p^{E82D} (Fig 8F). The impaired respiratory complex activity for Qcr7p^{E82R} and Qcr7p^{E82D}

was independent of any changes in the steady state amount of their subunits, (Fig 8C). These

results provide the first molecular evidence of the functional importance of a conserved PE-

binding site identified in the structures of yeast and human complex III. Collectively, these data

demonstrate that PE made in the IM by Psd1p is critical to support the intrinsic functionality of

complex III and establish one likely mechanism.

Fig 8. A PE-coordinating residue in Qcr7p is important for complex III activity. (A) The crystal structure of yeast cytochrome bc_1 that modeled associated lipids was downloaded using PDB ID: 1KB9. Using PyMOL, the region containing of the catalytic subunit Cob1p (magenta) near the matrix facing surface was enlarged to demonstrate hydrophobic interactions between this subunit and the acyl chains of PE. Arg51 of Qcr8p (orange) also shows hydrophobic interactions with a carbon atom from the ethanolamine headgroup. Glu82 of Qcr7p (blue) was predicted to form a hydrogen bonding interaction (3.4 Å distance) with the amine group of PE, whose atoms are depicted as spheres *gray*: carbon, *red*: oxygen, and *blue*: nitrogen (hydrogen atoms are not represented). (B) WT and mutant Qcr7p was detected in yeast whole cell extracts on the indicated strains by immunoblot; Aac2p serves as a loading control. (C) Mitochondria from the indicated strains were immunoblotted for subunits of complex III and complex IV as well as markers for the indicated mitochondrial compartments. (D)The indicated strains were spotted and incubated at 30°C for 2 days on YPD and for 3 days on rich lactate (RL) and SCEG. (E, F) Spectrophotometric assay following the (E) reduction or (F) oxidation of cytochrome *c* at 550nm in isolated mitochondria solubilizedin 0.5% (w/v) DDM. Analysis *versus* WT (*) was performed by one-way ANOVA ± S.E.M. for n=6 or n=3 for ρ⁰. P values for decreases that didn't achieve significance are reported and were evaluated by student t-test *versus* WT.

DISCUSSION

PE levels have been shown to be altered in models of the neurodegenerative disorders

Parkinson's disease [44] and Alzheimer's disease [45], and recent evidence has implicated the

turnover of mammalian Pisd as an important regulatory mechanism that diminishes the

proliferation of breast cancer cells [46]. Thus, regulation of the Psd pathway is not only important

for mitochondrial function but also organismal health and physiology.

Using an assortment of biochemical assays that measure both the combined and

individual activities of the respiratory chain complexes, we sought to clarify the respiratory

defect associated with *psd1Δ* yeast and whether or not PE made by the Kennedy pathway could

rescue it. Two different groups using the same yeast strain background have reported that

supplemental ethanolamine rescues [28] or fails to rescue [11] respiratory function of *psd1Δ* yeast.

Given these conflicting results, we first tested the ability of ethanolamine supplementation to

improve the respiratory growth phenotype of yeast lacking Psd1p. While we observed that

growth of *psd1Δpsd2Δ* could be rescued on respiratory medium supplemented with

ethanolamine, growth after 3 days was delayed and impaired similarly to *psd1Δ* yeast.

Perplexingly, supplementation of *psd1Δpsd2Δ* with ethanolamine to generate PE through the

Kennedy pathway did not significantly increase mitochondrial PE levels and instead resulted in

a significant, albeit modest, increase in CL (Fig 6C-F), which nonetheless failed to completely

restore respiratory growth. Importantly, the inability of ethanolamine to fully restore *psd1Δ*

respiratory growth was verified in four distinct yeast strain backgrounds suggesting that genetic

differences are not the source for the disparate results reported previously [1,11,27,28]. This notion is

further supported by the fact that two groups that each utilized *psd1Δ* derived from the same

parental strain came to opposite conclusions regarding the rescuing capacity of PE made by the

Kennedy pathway [11,28].

Interestingly, lyso-PE supplementation (PE containing only one acyl chain) can rescue

psd1Δ respiratory growth but acts via a separate pathway than the Kennedy pathway [27].

Further, lyso-PE has also been demonstrated to reverse the mitochondrial defects associated

with RNAi silencing of mammalian *Pisd* in cell culture [10], and can increase the proliferation rate

of the MCF7-RAS breast cancer cell line when Pisd levels are depleted [46]. Thus, the different

ER-resident PE producing pathways appear to differ with respect to their ability to compensate

for the absence of the mitochondrial Psd pathway suggesting that pathway-specific

mitochondrial import mechanisms for PE may exist.

Our oxidative phosphorylation analyses demonstrated that both OM-Psd1 and ER-Psd1

phenocopy the respiratory defect of *psd1Δ* mitochondria. This implies that the IMS is the

greatest barrier preventing PE import into the inner membrane. Moreover, we have identified a

key oxidative phosphorylation component that is uniquely dependent on PE made by Psd1p in

the IM: respiratory complex III. Specifically, we demonstrated that while PE synthesized in the

endosomal system–by ER-Psd1p, Psd2p, or the Kennedy pathway– can rescue the impaired

complex IV activity that occurs in the absence of Psd1p in the IM, it fails to do so for complex III

(Fig 5F and Fig 6B). Since complex III was impaired in yeast deficient in mitochondrial PS and

PE that contained a normal amount of fully processed Psd1p (*cho1Δ*, Fig 7K), our results

demonstrate that the complex III functional defect results from an insufficient supply of PE and

not from a non-PE related activity of Psd1p.

What is the basis for the capacity of extra-mitochondrially produced PE to rescue

complex IV but not complex III function? As both complex III and complex IV are IM residents,

one possibility is that these respiratory complexes require different levels of PE within the inner

membrane to function appropriately. If true, this would imply that the relative levels of PE in the

IM in the strains analyzed are equivalent between WT, *psd2Δ*, and IM-Psd1, similarly depleted

in ER-Psd1 and *psd1Δ*, less than *psd1Δ* in OM-Psd1, and almost completely absent in

psd1Δpsd2Δ. Unfortunately, while it is clear that OM-Psd1 and ER-Psd1 mitochondria contain

robust levels of PE, we have yet to determine its steady state distribution between the outer and

inner membranes. Another possible explanation for the different dependence of complex III and

IV function on IM-produced PE could reflect the nature of the PE made in the mitochondrion

versus elsewhere. Evidence that polyunsaturated 38:4 and 38:5 PE species are enriched in the

mitochondrial inner membrane of HeLa cells has been previously reported [47]. While both the

Kennedy and Psd pathways have the capacity to produce all diacyl-PE species, the Kennedy

pathway preferentially synthesizes mono 34:2 or di-unsaturated 36:3 PE and Pisd mainly

generates polyunsaturated 38:4 or 38:5 PE [48]. However, *S. cerevisiae* is unable to synthesize

complex unsaturated fatty acids and the PE acyl chain pattern is the same between whole cells

and isolated mitochondria [49,50]. It is interesting to note that, even though the Kennedy and Psd

pathways both share a preference for generating both mono- and di-unsaturated PE in yeast,

there is a 10% enrichment of 34:2 PE that contains oleic and palmitoleic acid in crude

mitochondria versus ER-derived microsomes [49]. Another intriguing possibility is that the

synthesis of PE by Psd1p in the IM is somehow directly coupled to the incorporation of this lipid

into partially or fully assembled complex III. Recently, a subunit of complex IV was found to

associate with the MICOS subunit, Mic19, in mammalian cells by EM tomography and

immunoprecipitation [51]. As the MICOS complex was suggested to work in concert with Ups2p to

regulate mitochondrial PS/PE metabolism in yeast [26], contact sites between the OM and IM

could potentially facilitate the transport of ER-derived PE to complex IV more directly than it can

to complex III.

Intriguingly, the improved complex IV functionality provided by ethanolamine correlated

best with an increased abundance of CL and not PE (Fig 6C-F), a phenomenon that has been

observed previously [1,28], suggesting that this rescue is actually CL-dependent. The importance

of CL in the assembly and function of the respiratory complexes and supercomplexes is well

documented [42,52,53]. Our results strongly indicate that when Psd1p is missing or mis-localized, a

minimal threshold of PE is not achieved within the IM that is necessary to accumulate normal

levels of CL and promote the full activity of the OXPHOS machinery. Although it is presently

unclear how the Kennedy Pathway for PE production promotes CL accumulation, it is known

that the Ups1p and Ups2p lipid trafficking proteins have an inverse relationship with respect to

CL/PE metabolism suggesting that this may be linked to PS/PA trafficking into the IM [3,54,55].

Depletion of PS in the *psd1Δpsd2Δ* background restored CL levels to that of *psd1Δ* and *cho1Δ*

which were still comparatively reduced *vs* WT (Fig 7G). This increase in CL also coincided with

improved growth for *cho1Δpsd1Δpsd2Δ* yeast (Fig 7K). It is possible that in the absence of PS,

metabolic pathways that either promote PA formation in the ER [56] or promote PA import to the

IM [57,58] are stimulated. Moving forward, it will be important to distinguish between these non-

mutually exclusive models.

Another outstanding question raised by our study is exactly how PE made by Psd1p in

the IM promotes complex III function. Since *psd1Δpsd2Δ* yeast contain normal amounts of

cytochrome *c* and CoQ_6 is not limiting, the underlying respiratory defect is intrinsic to complex

III. We demonstrated that mutations in a residue of Qcr7p predicted to bind PE impaired

complex III activity (Fig 8E) establishing a likely mechanistic basis for the requirement of PE for

its function. To our knowledge, this is the first molecular evidence demonstrating the functional

importance of a specific interaction of PE with complex III, which until now had only been

postulated from crystal structures of yeast complex III and human respiratory supercomplexes

[30,31]. A second PE is found adjacent to the dimer interface of this complex; as such, the acyl

chains of PE are thought to interact with both monomers and assist in dimer formation.

Additionally, the acyl chains of PE at the dimer interface could potentially promote quinol-

quinone exchange at the Q_i and Q_0 sites. More broadly, during quinol-quinone exchange,

sidechain movement of Cob1p is thought to be necessary to transfer protons from His202 to

ubiquinone, and perhaps depletion of PE hinders this movement and delays proton flux [59].

Further, electron-transfer reactions occur within microsecond-to-millisecond time scales

and electron-tunneling between complex III monomers is thought to maximize the efficiency of

coupled redox reactions between monomers [60]. The generation of the free radical semiquinone

during the Q cycle performed by complex III is proposed to be diminished by a mechanism that

delays quinol oxidation in the Q_o site when electron transfer is pharmacologically impaired [61]. If

depletion of PE or mutagenesis of residues that interact with this lipid diminish the efficiency of

electron transfer between complex III monomers or between complex III and its substrates, this

could result in reduced complex III function as a means to prevent superoxide production.

These structural observations will guide future efforts to determine the role(s) of PE as it relates

to complex III activity.

485
486   **MATERIALS AND METHODS**

487
Yeast strains and growth conditions

All yeast strains used in this study are listed in Table 1 and were derived from GA74-1A unless

otherwise noted. Deletion strains were generated by PCR-mediated gene replacement of the

entire reading frame as previously described [52]. Psd1p containing a COOH-terminal 3XFLAG

tag subcloned into pRS305 has been described [17,29]. To re-direct Psd1p to the mitochondrial

OM, the first 100 amino acids of Psd1p, encompassing its mitochondrial targeting sequence and

transmembrane domain, were replaced by the equivalent domains (amino acids 1-34) of the

single-pass OM resident protein, Tom20p. ER-Psd1p, which is directed to the secretory

pathway, was generated by replacing the first 57 amino acids of Psd1p, encompassing its

mitochondrial targeting signal (MTS), with the N-terminal signal sequence (amino acids 1-23) of

carboxypeptidase Y, as previously described [29]. Additionally, the ER-Psd1p construct contains

an NXS *N*-glycosylation signal immediately downstream of the CPY leader sequence to track its

topology. The IM-Psd1, OM-Psd1, and ER-Psd1 constructs, which all contained the C-terminal

3XFLAG tag, were subcloned into the pRS305 plasmid, linearized, and integrated into the *LEU2*

locus in the *psd1Δpsd2Δ* background. Clones were selected on synthetic dropout medium

(0.17% (w/v) yeast nitrogen base, 0.5% (w/v) ammonium sulfate, 0.2% (w/v) dropout mixture

synthetic-leu, 2% (w/v) dextrose) and verified by immunoblot.

For strains that were genetically modified by homology-integrated clustered regulatory

interspaced short palindromic repeats (CRISPR)-Cas (HI-CRISPR) [62], CRISPR-Cas9 gene

blocks were designed to target *CHO1* or *QCR7* and cloned into the pCRCT plasmid [62], as

previously described [17]. The spacer sequences (including the CRISPR-Cas9 target (20-bp) and

the homology repair template (50-bp) homology arms on both sides (total 100-bp) flanking the

Cas9 recognition sequence) were ordered as gBlocks (Integrated DNA Technologies).

Specifically, *CHO1* knockout was achieved by incorporating an 8-bp deletion within the

homology repair template to induce a frameshift mutation by removal of nucleotides 38-45

downstream of the start ATG sequence of the *CHO1* open reading frame (ORF). The CRISPR

construct was designed to target the protospacer adjacent motif (PAM) sequence encoded by

nucleotides 40-42 on the reverse strand of the *CHO1* ORF. Single point mutations were

introduced into the *QCR7* ORF at positions 244-246 downstream of the ATG start site to mutate

Glu82 (GAG) to Asp82 (GAC) or Arg82 (AGA) through the design of homology repair templates

encoding these mutations. Specificity of gene block integration into this locus was accomplished

by targeting the PAM sequence encoded by nucleotides 209-211 on the reverse strand of the

QCR7 ORF. Both *QCR7* gene blocks encoded for silent alanine mutations at position 208-210

to mutate the PAM sequence and prevent re-cleavage by Cas9 after homology-directed repair,

and position 214-216 to prevent hairpin formation of gBlocks. Additionally, the homology arms

adjacent to the PAM sequence were extended to 125-bp on each side for these constructs.

Integration of gene blocks encoding for these mutations were verified by sequencing of yeast

genomic DNA using primers specific for *QCR7*.

Yeast were grown on YPD (1% (w/v) yeast extract, 2% (w/v) tryptone, 2% (w/v) dextose)

plates. To assess the function of the assorted re-directed Psd1p constructs, overnight cultures

grown in synthetic complete dextrose (SCD; 0.17% (w/v) yeast nitrogen base, 0.5% (w/v)

ammonium sulfate, 0.2% (w/v) complete amino acid mixture, 2% (w/v) dextrose) supplemented

with 2mM ethanolamine hydrochloride were spotted on SCD plates in the absence or presence

of 2mM ethanolamine hydrochloride, or spotted onto rich lactate (1% (w/v) yeast extract, 2%

(w/v) tryptone, 0.05% (w/v) dextrose, 2% (v/v) lactic acid, 3.4mM $CaCl_2$-$2H_2O$, 8.5mM NaCl,

2.95mM $MgCl_2$-$6H_2O$, 7.35mM KH_2PO_4, 18.7mM NH_4Cl, pH 5.5), or synthetic complete ethanol

glycerol (SCEG; 0.17% (w/v) yeast nitrogen base, 0.5% (w/v) ammonium sulfate, 0.2% (w/v)

complete amino acid mixture, 1% (v/v) ethanol, 3% (v/v) glycerol) or synthetic complete lactate

(SC-LAC; 0.17% (w/v) yeast nitrogen base, 0.5% (w/v) ammonium sulfate, 0.2% (w/v) complete

amino acid mixture, 0.05% (w/v) dextrose, 2% (v/v) lactic acid, 3.4mM $CaCl_2$-$2H_2O$, 8.5mM

NaCl, 2.95mM $MgCl_2$-$6H_2O$, 7.35mM KH_2PO_4, 18.7mM NH_4Cl, pH 5.5) in the absence or

presence of 2mM ethanolamine hydrochloride to test respiratory growth.

For CoQ_6 supplementation experiments, starter cultures were grown in SCD + 2mM

ethanolamine overnight and diluted to 0.025 OD_{600} in 500μL of either SCEG, SCEG + 2mM

ethanolamine, SCEG + 2μM CoQ_6 (Avanti Polar Lipids, Inc), SCEG + 2μM CoQ_6 + 2mM

ethanolamine, SCEG + 10μM CoQ_6, or SCEG + 10μM CoQ_6 + 2mM ethanolamine in duplicate

in a 48 well plate. OD_{600} measurements were then recorded every 30 minutes for a period of 48

hours at 30°C using a Tecan Infinite 200 Pro instrument. For experiments designed to test the

importance of the pABA pathway for cell growth, synthetic media lacking pABA was utilized for

liquid and solid growth of yeast cells. -pABA media consisted of 790 mg/mL CSM Mixture

Complete (Formedium LTD) and 6.9g/L yeast nitrogen base lacking amino acids and *para*-

amino benzoic acid (Formedium LTD). For +pABA media, the same mixture was used but

included the addition of 100μM *para*-amino benzoic acid (Research Products International, Inc)

dissolved in water and sterile filtered using a 0.20μM filter (Corning, Inc). Individual colonies of

yeast were used to inoculate starter cultures in -pABA medium containing 2% (w/v) glucose and

2mM ethanolamine. Solid growth of yeast on agar plates was evaluated in -/+pABA medium

containing 1% (v/v) ethanol and 3% (v/v) glycerol -/+ 2mM ethanolamine.

Mitochondrial Isolation

Isolation of crude mitochondria was performed as previously described [63]. Yeast were selected

on rich lactate plates and grown on rich lactate media to prevent the loss of mitochondrial DNA.

To improve growth on rich lactate, *psd1Δpsd2Δ* yeast were grown in the presence of 2mM

choline prior to harvesting mitochondria (except where indicated in Fig 6A,B and in Fig 7).

Purification of mitochondria by sucrose step gradient was performed as previously described [29].

Electron microscopy

Cells were grown in rich lactate medium and harvested at mid-log phase by centrifugation. Cells

were fixed in 3% glutaraldehyde contained in 0.1 M sodium cacodylate, pH 7.4, 5 mM $CaCl_2$, 5

mM $MgCl_2$, and 2.5% (w/v) sucrose for 1 hr at room temperature with gentle agitation,

spheroplasted, embedded in 2% ultra-low temperature agarose (prepared in water), cooled, and

subsequently cut into small pieces ($\sim$1 mm^3) as previously described [52]. The cells were then

post-fixed in 1% OsO_4, 1% potassium ferrocyanide contained in 0.1 M sodium cacodylate, 5

mM $CaCl_2$, pH 7.4, for 30 min at room temperature. The blocks were washed thoroughly four

times with double distilled H_2O, 10 min total, transferred to 1% thiocarbohydrazide at room

temperature for 3 min, washed in double distilled H_2O (four times, 1 min each), and transferred

to 1% OsO_4, 1% potassium ferrocyanide in 0.1 M sodium cacodylate, pH 7.4, for an additional 3

min at room temperature. The cells were washed four times with double distilled H_2O (15 min

total), stained *en bloc* in Kellenberger's uranyl acetate for 2 hr to overnight, dehydrated through

a graded series of ethanol, and subsequently embedded in Spurr resin. Sections were cut on a

Reichert Ultracut T ultramicrotome, post-stained with uranyl acetate and lead citrate, and

observed on an FEI Tecnai 12 transmission electron microscope at 100 kV. Images were

recorded with a Soft Imaging System Megaview III digital camera, and figures were assembled

in Adobe Photoshop with only linear adjustments in contrast and brightness.

mtDNA quantitation

DNA was extracted as described [64]. In brief, yeast cells were grown for 2 days in rich lactate

and the collected cell pellets were vortexed at level 10 for 3 minutes with 200μL breaking buffer

(2% (v/v) Triton X-100, 1% (v/v) SDS, 100mM NaCl, 10mM Tris pH8.0, 1mM EDTA pH 8.0),

0.3g glass beads, and 200μL phenol/chloroform/isoamyl alcohol at room temperature. The

solution was neutralized with the addition of 200μL of Tris-EDTA (TE) buffer pH 8.0 and phases

were separated by centrifugation at 21,000 x g for 5 minutes. The aqueous phase was collected

and DNA was precipitated by the addition of 100% ethanol and collected in the pellet after

centrifugation at 21,000 x g for 3 minutes. The pellets were resuspended in 400μL TE buffer, pH

8.0 and RNA was digested with the addition of 3 μL of 10mg/mL RNAse A and incubation at

37°C for 5 minutes before addition of 10 μL of 4M Ammonium acetate and 1 mL 100% ethanol.

DNA pellets were recovered by centrifugation at 21,000 x g for 3 minutes, dried, and

resuspended in 30 μL TE buffer pH 8.0. DNA was stored at -80°C until ready for use,

quantitated, and a portion diluted to 10ng/μL to be used as template in the qPCR reaction. The

FastStart Universal SYBR Green Master Rox (Roche) was used for qPCR performed according

to the manufacturer's instructions. 50ng of genomic DNA was used as a template and the

following primers were used at 100nM concentration in a 20μL reaction: *COX1* forward (5′-

CTACAGATACAGCATTTCCAAGA-3′), *COX1* reverse (5′- GTGCCTGAATAGATGATAATGGT-

3′), *ACT1* forward (5′- GTATGTGTAAAGCCGGTTTTG-3′), and *ACT1* reverse (5′-

CATGATACCTTGGTGTCTTGG -3′). The reactions were performed in technical duplicate with

three biological replicates. After completion of thermocycling in a QuantStudio 6 Flex Real-Time

PCR System (Thermo Fisher), melting-curve data was collected to verify PCR specificity. The

absence of primer dimers and the Ct value difference between the nuclear (*ACT1*) and

mitochondrial (*COX1*) target were computed as a measure of the mitochondrial DNA copy

number relative to the nuclear genome.

Mitochondrial respiration measurements

Mitochondrial oxygen consumption was measured using a Clark-type oxygen electrode in a

magnetically stirred, thermostatically controlled 1.5 mL chamber at 25°C (Oxytherm; Hansatech)

as previously described [65] with some variations. 100μg of mitochondria were resuspended in

0.25M sucrose, 0.25mg/mL BSA, 20mM KCl, 20mM Tris-Cl, 0.5mM EDTA, 4mM KH_2PO_4, and

3mM $MgCl_2$, pH 7.2. After addition of 1mM ascorbate + 0.3mM TMPD, state 2 rate was

monitored for approximately 30 sec. State 3 respiration was initiated by addition of 50μM ADP.

After state 4 rate was measured, 10μM CCCP was added to induce uncoupled respiration, and

the rate was followed for either 2 minutes or until oxygen level reached zero.

Complex III and IV Activity Measurements

Complex III and IV activities were measured as described [66,67]. To measure complex III activity,

25μg of mitochondria solubilized in 0.5% (w/v) *n*-dodecyl-β-D-maltoside were added to reaction

buffer (50mM KP_i, 2mM EDTA, pH 7.4) with 0.008% (w/v) horse heart cytochrome *c* and 1mM

KCN. The reaction was started by adding 100μM decylubiquinol, and the reduction of

cytochrome *c* followed at 550nM. Complex IV activity was initiated by adding 5μg of solubilized

mitochondria to reaction buffer with 0.008% ferrocytochrome *c* and measured by recording

cytochrome *c* oxidation at 550nm.

Antibodies

Most antibodies used in this study were generated by our laboratory or in the laboratories of J.

Schatz (University of Basel, Basel, Switzerland) or C. Koehler (UCLA) and have been described

previously [3,63,65,67-69]. Other antibodies used were mouse anti-Sec62p (kind gift of Dr. David

Meyers (UCLA)), mouse anti-FLAG (clone M2, Sigma), mouse anti-Dpm1p (113686, Abcam),

rabbit anti-Qcr7p [70], rabbit antisera reactive to Coq1p [71], Coq4p [72], Coq7p [73], or Coq9p [74], rabbit

antisera raised against the C-terminus of Cho1p [34], rabbit anti-Kar2p [36] and horseradish

peroxidase-conjugated (Thermo Fisher Scientific) or IRDye 800CW (LI-COR) secondary

antibodies.

Miscellaneous

Preparation of yeast cell extracts, submitochondrial fractionation, phospholipid analysis, 1D BN-

PAGE, and immunoblotting were performed as described previously [17,29]. Immunoblots using IR

800 CW secondary antibodies were imaged using an Odyssey CLx Imaging System.

Immunoblots and TLC plates were quantitated by Quantity One Software by Bio-Rad

Laboratories. Statistical comparisons (ns, $P > 0.05$; 1 symbol $P \leq 0.05$; 2 symbols $P \leq 0.01$; 3

symbols $P \leq 0.001$; 4 symbols $P \leq 0.0001$) were performed by student t test or one-way analysis

of variance (ANOVA) with Holm-Sidak pairwise comparison using SigmaPlot 11 software

(Systat Software, San Jose, CA); $P \leq 0.05$ were deemed significant. In some cases, replicates

of samples were loaded on the same SDS-PAGE gel, and thus borders of neighboring samples

may be detected on the borders of some immunoblots. All graphs show the mean $\pm$ S.E.M. At

least three biological replicates represent each of the experiments performed in this study,

unless otherwise indicated.

643
REFERENCES
1   Birner, R., Burgermeister, M., Schneiter, R. & Daum, G. Roles of
phosphatidylethanolamine and of its several biosynthetic pathways in Saccharomyces
cerevisiae. *Mol Biol Cell* **12**, 997-1007 (2001).
2   Horvath, S. E. *et al.* Processing and topology of the yeast mitochondrial
phosphatidylserine decarboxylase 1. *J Biol Chem* **287**, 36744-36755,
doi:10.1074/jbc.M112.398107 (2012).
3   Tamura, Y. *et al.* Phosphatidylethanolamine biosynthesis in mitochondria:
phosphatidylserine (PS) trafficking is independent of a PS decarboxylase and
intermembrane space proteins UPS1P and UPS2P. *J Biol Chem* **287**, 43961-43971,
doi:10.1074/jbc.M112.390997 (2012).
4   Zborowski, J., Dygas, A. & Wojtczak, L. Phosphatidylserine decarboxylase is located on
the external side of the inner mitochondrial membrane. *FEBS Lett* **157**, 179-182 (1983).
5   Trotter, P. J. & Voelker, D. R. Identification of a non-mitochondrial phosphatidylserine
decarboxylase activity (PSD2) in the yeast Saccharomyces cerevisiae. *J Biol Chem* **270**,
6062-6070 (1995).
6   Calzada, E., Onguka, O. & Claypool, S. M. Phosphatidylethanolamine Metabolism in
Health and Disease. *Int Rev Cell Mol Biol* **321**, 29-88, doi:10.1016/bs.ircmb.2015.10.001
(2016).

7   Steenbergen, R. *et al.* Disruption of the phosphatidylserine decarboxylase gene in mice
causes embryonic lethality and mitochondrial defects. *J Biol Chem* **280**, 40032-40040,
doi:10.1074/jbc.M506510200 (2005).
8   Fullerton, M. D., Hakimuddin, F. & Bakovic, M. Developmental and metabolic effects of
disruption of the mouse CTP:phosphoethanolamine cytidylyltransferase gene (Pcyt2). *Mol*
Cell Biol **27**, 3327-3336, doi:10.1128/MCB.01527-06 (2007).
9   Bottinger, L. *et al.* Phosphatidylethanolamine and cardiolipin differentially affect the
stability of mitochondrial respiratory chain supercomplexes. *J Mol Biol* **423**, 677-686,
doi:10.1016/j.jmb.2012.09.001 (2012).
10   Tasseva, G. *et al.* Phosphatidylethanolamine deficiency in Mammalian mitochondria
impairs oxidative phosphorylation and alters mitochondrial morphology. *J Biol Chem* **288**,
4158-4173, doi:10.1074/jbc.M112.434183 (2013).
11   Chan, E. Y. & McQuibban, G. A. Phosphatidylserine decarboxylase 1 (Psd1) promotes
mitochondrial fusion by regulating the biophysical properties of the mitochondrial
membrane and alternative topogenesis of mitochondrial genome maintenance protein 1
(Mgm1). *J Biol Chem* **287**, 40131-40139, doi:10.1074/jbc.M112.399428 (2012).
12   Joshi, A. S., Thompson, M. N., Fei, N., Huttemann, M. & Greenberg, M. L. Cardiolipin and
mitochondrial phosphatidylethanolamine have overlapping functions in mitochondrial
fusion in Saccharomyces cerevisiae. *J Biol Chem* **287**, 17589-17597,
doi:10.1074/jbc.M111.330167 (2012).
13   Burgermeister, M., Birner-Grunberger, R., Nebauer, R. & Daum, G. Contribution of
different pathways to the supply of phosphatidylethanolamine and phosphatidylcholine to
mitochondrial membranes of the yeast Saccharomyces cerevisiae. *Biochim Biophys Acta*
1686, 161-168, doi:10.1016/j.bbalip.2004.09.007 (2004).
14   Gulshan, K., Shahi, P. & Moye-Rowley, W. S. Compartment-specific synthesis of
phosphatidylethanolamine is required for normal heavy metal resistance. *Mol Biol Cell* **21**,
443-455, doi:10.1091/mbc.E09-06-0519 (2010).
15   Di Bartolomeo, F., Wagner, A. & Daum, G. Cell biology, physiology and enzymology of
phosphatidylserine decarboxylase. *Biochim Biophys Acta* **1862**, 25-38,
doi:10.1016/j.bbalip.2016.09.007 (2017).
16   Choi, J. Y., Duraisingh, M. T., Marti, M., Ben Mamoun, C. & Voelker, D. R. From Protease
to Decarboxylase: THE MOLECULAR METAMORPHOSIS OF PHOSPHATIDYLSERINE
DECARBOXYLASE. *J Biol Chem* **290**, 10972-10980, doi:10.1074/jbc.M115.642413
(2015).
17   Ogunbona, O. B., Onguka, O., Calzada, E. & Claypool, S. M. Multitiered and Cooperative
Surveillance of Mitochondrial Phosphatidylserine Decarboxylase 1. *Mol Cell Biol* **37**,
doi:10.1128/MCB.00049-17 (2017).
18   Vance, J. E. Phospholipid synthesis in a membrane fraction associated with mitochondria.
J Biol Chem **265**, 7248-7256 (1990).
19   Voelker, D. R. Phosphatidylserine functions as the major precursor of
phosphatidylethanolamine in cultured BHK-21 cells. *Proc Natl Acad Sci U S A* **81**, 2669-
2673 (1984).
20   Elbaz-Alon, Y. *et al.* A dynamic interface between vacuoles and mitochondria in yeast.
Dev Cell **30**, 95-102, doi:10.1016/j.devcel.2014.06.007 (2014).
21   Kojima, R., Endo, T. & Tamura, Y. A phospholipid transfer function of ER-mitochondria
encounter structure revealed in vitro. *Scientific reports* **6**, 30777, doi:10.1038/srep30777
(2016).
22   Kornmann, B. *et al.* An ER-mitochondria tethering complex revealed by a synthetic biology
screen. *Science* **325**, 477-481, doi:10.1126/science.1175088 (2009).

712 23 Lahiri, S. *et al.* A conserved endoplasmic reticulum membrane protein complex (EMC)
713   facilitates phospholipid transfer from the ER to mitochondria. *PLoS Biol* **12**, e1001969,
714   doi:10.1371/journal.pbio.1001969 (2014).
715 24 Nguyen, T. T. *et al.* Gem1 and ERMES do not directly affect phosphatidylserine transport
716   from ER to mitochondria or mitochondrial inheritance. *Traffic* **13**, 880-890,
717   doi:10.1111/j.1600-0854.2012.01352.x (2012).
718 25 Miyata, N., Watanabe, Y., Tamura, Y., Endo, T. & Kuge, O. Phosphatidylserine transport
719   by Ups2–Mdm35 in respiration-active mitochondria. *The Journal of Cell Biology* **214**, 77-
720   88, doi:10.1083/jcb.201601082 (2016).
721 26 Aaltonen, M. J. *et al.* MICOS and phospholipid transfer by Ups2-Mdm35 organize
722   membrane lipid synthesis in mitochondria. *J Cell Biol* **213**, 525-534,
723   doi:10.1083/jcb.201602007 (2016).
724 27 Riekhof, W. R. & Voelker, D. R. Uptake and utilization of lyso-phosphatidylethanolamine
725   by Saccharomyces cerevisiae. *J Biol Chem* **281**, 36588-36596,
726   doi:10.1074/jbc.M608851200 (2006).
727 28 Baker, C. D., Basu Ball, W., Pryce, E. N. & Gohil, V. M. Specific requirements of nonbilayer
728   phospholipids in mitochondrial respiratory chain function and formation. *Mol Biol Cell* **27**,
729   2161-2171, doi:10.1091/mbc.E15-12-0865 (2016).
730 29 Onguka, O., Calzada, E., Ogunbona, O. B. & Claypool, S. M. Phosphatidylserine
731   decarboxylase 1 autocatalysis and function does not require a mitochondrial-specific
732   factor. *J Biol Chem* **290**, 12744-12752, doi:10.1074/jbc.M115.641118 (2015).
733 30 Guo, R., Zong, S., Wu, M., Gu, J. & Yang, M. Architecture of Human Mitochondrial
734   Respiratory Megacomplex I2III2IV2. *Cell* **170**, 1247-1257 e1212,
735   doi:10.1016/j.cell.2017.07.050 (2017).
736 31 Lange, C., Nett, J. H., Trumpower, B. L. & Hunte, C. Specific roles of protein-phospholipid
737   interactions in the yeast cytochrome bc1 complex structure. *EMBO J* **20**, 6591-6600,
738   doi:10.1093/emboj/20.23.6591 (2001).
739 32 Johnson, L. M., Bankaitis, V. A. & Emr, S. D. Distinct sequence determinants direct
740   intracellular sorting and modification of a yeast vacuolar protease. *Cell* **48**, 875-885
741   (1987).
742 33 Storey, M. K. *et al.* Phosphatidylethanolamine has an essential role in Saccharomyces
743   cerevisiae that is independent of its ability to form hexagonal phase structures. *J Biol
744   Chem* **276**, 48539-48548, doi:10.1074/jbc.M109043200 (2001).
745 34 Choi, H. S., Han, G. S. & Carman, G. M. Phosphorylation of yeast phosphatidylserine
746   synthase by protein kinase A: identification of Ser46 and Ser47 as major sites of
747   phosphorylation. *J Biol Chem* **285**, 11526-11536, doi:10.1074/jbc.M110.100727 (2010).
748 35 Kodaki, T. & Yamashita, S. Yeast phosphatidylethanolamine methylation pathway.
749   Cloning and characterization of two distinct methyltransferase genes. *J Biol Chem* **262**,
750   15428-15435 (1987).
751 36 Huyer, G. *et al.* Distinct machinery is required in Saccharomyces cerevisiae for the
752   endoplasmic reticulum-associated degradation of a multispanning membrane protein and
753   a soluble luminal protein. *J Biol Chem* **279**, 38369-38378, doi:10.1074/jbc.M402468200
754   (2004).
755 37 Trotter, P. J., Pedretti, J. & Voelker, D. R. Phosphatidylserine decarboxylase from
756   Saccharomyces cerevisiae. Isolation of mutants, cloning of the gene, and creation of a
757   null allele. *J Biol Chem* **268**, 21416-21424 (1993).
758 38 Schagger, H. & Pfeiffer, K. Supercomplexes in the respiratory chains of yeast and
759   mammalian mitochondria. *EMBO J* **19**, 1777-1783, doi:10.1093/emboj/19.8.1777 (2000).
760 39 He, C. H., Xie, L. X., Allan, C. M., Tran, U. C. & Clarke, C. F. Coenzyme Q supplementation
761   or over-expression of the yeast Coq8 putative kinase stabilizes multi-subunit Coq

polypeptide complexes in yeast coq null mutants. *Biochim Biophys Acta* **1841**, 630-644,
doi:10.1016/j.bbalip.2013.12.017 (2014).

40   Sevin, D. C. & Sauer, U. Ubiquinone accumulation improves osmotic-stress tolerance in
Escherichia coli. *Nature chemical biology* **10**, 266-272, doi:10.1038/nchembio.1437
(2014).

41   Stefely, J. A. *et al.* Mitochondrial protein functions elucidated by multi-omic mass
spectrometry profiling. *Nat Biotechnol* **34**, 1191-1197, doi:10.1038/nbt.3683 (2016).

42   Jiang, F. *et al.* Absence of cardiolipin in the crd1 null mutant results in decreased
mitochondrial membrane potential and reduced mitochondrial function. *J Biol Chem* **275**,
22387-22394, doi:10.1074/jbc.M909868199 (2000).

43   Atkinson, K. D. *et al.* Yeast mutants auxotrophic for choline or ethanolamine. *J Bacteriol*
141, 558-564 (1980).

44   Wang, S. *et al.* Phosphatidylethanolamine deficiency disrupts alpha-synuclein
homeostasis in yeast and worm models of Parkinson disease. *Proc Natl Acad Sci U S A*
111, E3976-3985, doi:10.1073/pnas.1411694111 (2014).

45   Nesic, I. *et al.* Alterations in phosphatidylethanolamine levels affect the generation of
Abeta. *Aging Cell* **11**, 63-72, doi:10.1111/j.1474-9726.2011.00760.x (2012).

46   Keckesova, Z. *et al.* LACTB is a tumour suppressor that modulates lipid metabolism and
cell state. *Nature* **543**, 681-686, doi:10.1038/nature21408 (2017).

47   Kainu, V., Hermansson, M., Hanninen, S., Hokynar, K. & Somerharju, P. Import of
phosphatidylserine to and export of phosphatidylethanolamine molecular species from
mitochondria. *Biochim Biophys Acta* **1831**, 429-437, doi:10.1016/j.bbalip.2012.11.003
(2013).

48   Bleijerveld, O. B., Brouwers, J. F., Vaandrager, A. B., Helms, J. B. & Houweling, M. The
CDP-ethanolamine pathway and phosphatidylserine decarboxylation generate different
phosphatidylethanolamine molecular species. *J Biol Chem* **282**, 28362-28372,
doi:10.1074/jbc.M703786200 (2007).

49   Burgermeister, M., Birner-Grunberger, R., Heyn, M. & Daum, G. Contribution of different
biosynthetic pathways to species selectivity of aminoglycerophospholipids assembled into
mitochondrial membranes of the yeast Saccharomyces cerevisiae. *Biochim Biophys Acta*
1686, 148-160, doi:10.1016/j.bbalip.2004.09.005 (2004).

50   Schneiter, R. *et al.* Electrospray ionization tandem mass spectrometry (ESI-MS/MS)
analysis of the lipid molecular species composition of yeast subcellular membranes
reveals acyl chain-based sorting/remodeling of distinct molecular species en route to the
plasma membrane. *J Cell Biol* **146**, 741-754 (1999).

51   Sastri, M. *et al.* Sub-mitochondrial localization of the genetic-tagged mitochondrial
intermembrane space-bridging components Mic19, Mic60 and Sam50. *J Cell Sci* **130**,
3248-3260, doi:10.1242/jcs.201400 (2017).

52   Baile, M. G. *et al.* Unremodeled and remodeled cardiolipin are functionally
indistinguishable in yeast. *J Biol Chem* **289**, 1768-1778, doi:10.1074/jbc.M113.525733
(2014).

53   Pfeiffer, K. *et al.* Cardiolipin stabilizes respiratory chain supercomplexes. *J Biol Chem* **278**,
52873-52880, doi:10.1074/jbc.M308366200 (2003).

54   Miyata, N., Goda, N., Matsuo, K., Hoketsu, T. & Kuge, O. Cooperative function of Fmp30,
Mdm31, and Mdm32 in Ups1-independent cardiolipin accumulation in the yeast
Saccharomyces cerevisiae. *Sci Rep* **7**, 16447, doi:10.1038/s41598-017-16661-2 (2017).

55   Tamura, Y., Endo, T., Iijima, M. & Sesaki, H. Ups1p and Ups2p antagonistically regulate
cardiolipin metabolism in mitochondria. *J Cell Biol* **185**, 1029-1045,
doi:10.1083/jcb.200812018 (2009).

56 Lu, Y. W. & Claypool, S. M. Disorders of phospholipid metabolism: an emerging class of
mitochondrial disease due to defects in nuclear genes. *Front Genet* **6**, 3,
doi:10.3389/fgene.2015.00003 (2015).
57 Connerth, M. *et al.* Intramitochondrial transport of phosphatidic acid in yeast by a lipid
transfer protein. *Science* **338**, 815-818, doi:10.1126/science.1225625 (2012).
58 Watanabe, Y., Tamura, Y., Kawano, S. & Endo, T. Structural and mechanistic insights into
phospholipid transfer by Ups1-Mdm35 in mitochondria. *Nat Commun* **6**, 7922,
doi:10.1038/ncomms8922 (2015).
59 Hunte, C., Koepke, J., Lange, C., Rossmanith, T. & Michel, H. Structure at 2.3 A resolution
of the cytochrome bc(1) complex from the yeast Saccharomyces cerevisiae co-crystallized
with an antibody Fv fragment. *Structure* **8**, 669-684 (2000).
60 Swierczek, M. *et al.* An electronic bus bar lies in the core of cytochrome bc1. *Science* **329**,
451-454, doi:10.1126/science.1190899 (2010).
61 Quinlan, C. L., Gerencser, A. A., Treberg, J. R. & Brand, M. D. The mechanism of
superoxide production by the antimycin-inhibited mitochondrial Q-cycle. *J Biol Chem* **286**,
31361-31372, doi:10.1074/jbc.M111.267898 (2011).
62 Bao, Z. *et al.* Homology-Integrated CRISPR–Cas (HI-CRISPR) System for One-Step
Multigene Disruption in Saccharomyces cerevisiae. *ACS Synthetic Biology* **4**, 585-594,
doi:10.1021/sb500255k (2015).
63 Claypool, S. M., McCaffery, J. M. & Koehler, C. M. Mitochondrial mislocalization and
altered assembly of a cluster of Barth syndrome mutant tafazzins. *The Journal of cell*
biology **174**, 379-390. PMCID: PMC2064234, doi:jcb.200605043 [pii]
10.1083/jcb.200605043 (2006).
64 Hoffman, C. S. Preparation of yeast DNA. *Curr Protoc Mol Biol* **Chapter 13**, Unit13.11,
doi:10.1002/0471142727.mb1311s39 (2001).
65 Claypool, S. M., Oktay, Y., Boontheung, P., Loo, J. A. & Koehler, C. M. Cardiolipin defines
the interactome of the major ADP/ATP carrier protein of the mitochondrial inner
membrane. *J Cell Biol* **182**, 937-950, doi:10.1083/jcb.200801152 (2008).
66 Tzagoloff, A., Akai, A. & Needleman, R. B. Assembly of the mitochondrial membrane
system: isolation of nuclear and cytoplasmic mutants of Saccharomyces cerevisiae with
specific defects in mitochondrial functions. *J Bacteriol* **122**, 826-831 (1975).
67 Baile, M. G., Whited, K. & Claypool, S. M. Deacylation on the matrix side of the
mitochondrial inner membrane regulates cardiolipin remodeling. *Mol Biol Cell* **24**, 2008-
2020, doi:10.1091/mbc.E13-03-0121 (2013).
68 Whited, K., Baile, M. G., Currier, P. & Claypool, S. M. Seven functional classes of Barth
syndrome mutation. *Hum Mol Genet* **22**, 483-492, doi:10.1093/hmg/dds447 (2013).
69 Riezman, H. *et al.* Import of proteins into mitochondria: a 70 kilodalton outer membrane
protein with a large carboxy-terminal deletion is still transported to the outer membrane.
EMBO J **2**, 2161-2168 (1983).
70 Hildenbeutel, M. *et al.* Assembly factors monitor sequential hemylation of cytochrome
b to regulate mitochondrial translation. *The Journal of Cell Biology* **205**, 511-
524, doi:10.1083/jcb.201401009 (2014).
71 Gin, P. & Clarke, C. F. Genetic Evidence for a Multi-subunit Complex in Coenzyme Q
Biosynthesis in Yeast and the Role of the Coq1 Hexaprenyl Diphosphate Synthase.
Journal of Biological Chemistry **280**, 2676-2681, doi:10.1074/jbc.M411527200 (2005).
72 Belogrudov, G. I. *et al.* Yeast COQ4 Encodes a Mitochondrial Protein Required for
Coenzyme Q Synthesis. *Archives of Biochemistry and Biophysics* **392**, 48-58,
doi:https://doi.org/10.1006/abbi.2001.2448 (2001).
73 Tran, U. C. *et al.* Complementation of Saccharomyces cerevisiae coq7 Mutants by
Mitochondrial Targeting of the Escherichia coli UbiF Polypeptide: TWO FUNCTIONS OF

YEAST COQ7 POLYPEPTIDE IN COENZYME Q BIOSYNTHESIS. *Journal of Biological*
Chemistry **281**, 16401-16409, doi:10.1074/jbc.M513267200 (2006).
74 Hsieh, E. J., Dinoso, J. B. & Clarke, C. F. A tRNATRP gene mediates the suppression of
cbs2-223 previously attributed to ABC1/COQ8. *Biochemical and Biophysical Research*
Communications **317**, 648-653, doi:https://doi.org/10.1016/j.bbrc.2004.03.096 (2004).
866

ACKNOWLEDGEMENTS

We would like to thank Drs. Carla Koehler (UCLA), Cathy Clarke (UCLA), Martin Ott (Stockholm

University), Susan Michaelis (Johns Hopkins University School of Medicine), and George

Carman (Rutgers University) for antibodies, Drs. Ya-Wen Lu, Ouma Onguka, Seun Ogunbona,

and Matthew G. Baile for technical assistance, and Erica Avery for revision of this manuscript.

This work was supported by the National Institutes of Health (R01GM111548 to S.M.C. and

T32GM007445 to E.C.). The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

E. C. and S. M. C. co-wrote and performed experiments for this manuscript. J. M. M. performed

all electron microscopy sample preparation, sectioning, and imaging.

COMPETING INTERESTS

The authors declare no competing interests in the design and/or interpretation of this study.

MATERIALS & CORRESPONDENCE

Correspondence and material requests should be directed to S.M.C.

FIGURE LEGENDS

Fig 1. Ethanolamine supplementation fails to rescue the growth defect of *psd1Δ* on respiratory medium in 4 separate yeast strains. (A, B) The indicated strains were pre-cultured at 30°C in YPD and spotted onto (A) synthetic complete dextrose (SCD) or ethanol-glycerol (SCEG) medium +/- 2mM ethanolamine (+E) or (B) SCEG +/- 10mM ethanolamine and incubated at 30°C for 3 days.

Fig 2. OM-Psd1p and ER-Psd1p constructs are functional and localize to the OM and ER,

respectfully. (A) Schematic of IM-Psd1p, OM-Psd1p, and ER-Psd1p. All three constructs contain

a 3XFLAG tag at the C-terminus (yellow). The Tom20 residues (1-100) that replace the

mitochondrial targeting sequence (MT) and transmembrane (TM) domain of IM-Psd1p (green)

are shown for OM-Psd1p (blue), and the carboxypeptidase Y signal sequence (residues 1-37) as

well as an NXS motif are indicated for ER-Psd1p (purple). (B) The β and α subunits of Psd1p

were detected in yeast whole cell extracts of the indicated strains by immunoblot. Tom70p served

as a loading control. (C) The indicated strains were spotted onto synthetic complete dextrose

(SCD) medium +/- 2mM ethanolamine (+E) and incubated at 30°C for 4 days. (D) Protease

protection assay in intact mitochondria (Mito), osmotically ruptured mitochondria (MP), or

deoxycholate-solubilized mitochondria (Det.). Following incubation -/+ 100μg proteinase K (Prot.

K) for 30 minutes, samples were collected, resolved by SDS-PAGE and immunoblotted for Psd1p

(β and α subunits), and the mitochondrial compartment-specific markers Tom70p (OM), Tim54p

(IM), and Abf2p (matrix). (E) Illustration indicating the topology of (1) IM-Psd1p, (2) OM-Psd1p,

and (3) ER-Psd1p.

Fig 3. OM-Psd1 and ER-Psd1 contain robust levels of PE in both cellular and mitochondrial

membranes. (A-F) Cellular and (G-L) mitochondrial phospholipids from the indicated strains were

labeled overnight with $^{32}P_i$ and separated by TLC. All graphs show the mean $\pm$ S.E.M. for n=6

biological replicates. Significant differences compared to WT were calculated by one-way

ANOVA. (M) Key for symbols used for statistical analysis interpretation by one-way ANOVA when

comparing samples *versus* WT (*), *psd1Δpsd2Δ* (#), or IM-Psd1 ($).

Fig 4. OM-Psd1 and ER-Psd1 OXPHOS function phenocopies *psd1Δ*. (A) The indicated

strains were spotted and incubated at 30°C for 2 days on SCD +/- 2mM ethanolamine (+E) and

for 5 days on rich lactate (RL), SC lactate (SC-LAC), and SCEG +/- 2mM ethanolamine (+E). (B-

E) O_2 consumption measurements from mitochondria isolated from yeast grown in rich lactate

using ascorbate-TMPD as a substrate. (B) The maximal respiratory rate was recorded after the

addition of CCCP, (C) state 3 respiration was assessed after addition of ADP, and (D) state 4

respiration was recorded following ADP depletion. (E) The respiratory control ratio (RCR) is

calculated by dividing state 3 by state 4 respiratory rates. Analysis *versus* WT (*), *psd1Δpsd2Δ*

(#), or IM-Psd1 ($) was performed by one-way ANOVA $\pm$ S.E.M. for n=4.

Fig 5. Complex III and IV activities are impaired when Psd1p is absent in the IM. (A) mtDNA

was isolated from the indicated strains, normalized, and quantified by qPCR. Analysis was

performed by one-way ANOVA $\pm$ S.E.M. for n=3. (B) Spectrophotometric assay following the

oxidation of cytochrome *c* at 550nM in isolated mitochondria solubilized in 0.5% (w/v) DDM.

Analysis *versus* WT (*), *psd1Δpsd2Δ* (#), or IM-Psd1 ($) was performed by one-way ANOVA $\pm$

S.E.M. for n=6 or n=3 for ρ^0. P values for decreases that didn't achieve significance are reported

in red and were analyzed by student *t*-test *versus* WT. (C) Mitochondria from the indicated strains

were immunoblotted for subunits of complex III (CIII), complex IV (CIV), complex V (CV), the Coq

synthome, cytochrome *c*, and markers of each mitochondrial compartment. (D, E) Blue native-

PAGE analysis of respiratory supercomplexes (RSCs) using mitochondrial extracts solubilized in

1.5% (w/v) digitonin. (D) Complex IV assembly was monitored by immunoblot against the nuclear-

encoded subunit Cox4p and (E) Complex III assembly was monitored by immunoblot against the

nuclear-encoded subunit Rip1p. Mitochondria lacking CL (*crd1Δ*) were used as a positive control

for RSC destabilization[53]. (F) Spectrophotometric assay following the reduction of cytochrome *c*

at 550nM in isolated mitochondria solubilized in 0.5% (w/v) DDM. Analysis *versus* WT (*),

psd1Δpsd2Δ (#), or IM-Psd1 ($) was performed by one-way ANOVA $\pm$ S.E.M for n=6 or n=3 for

ρ^0. P values for decreases that didn't achieve significance are reported in red and were analyzed

by student *t*-test *versus* WT.

Fig 6. The Kennedy Pathway for PE synthesis can rescue the activity of complex IV but not

complex III. Spectrophotometric assay following the (A) oxidation and (B) reduction of

cytochrome *c* at 550nM in isolated mitochondria solubilized in 0.5% (w/v) DDM. Analysis *versus*

WT (*) or *psd1Δpsd2Δ* (#) was performed by one-way ANOVA $\pm$ S.E.M. for n=6 or n=4 for ρ^0. P

values for decreases that didn't achieve significance are reported in red and were analyzed by

student *t*-test. (C-G) Mitochondrial phospholipids from the indicated strains were labeled overnight

with $^{32}P_i$ and separated by TLC. (C) Representative TLC plate for mitochondrial $^{32}P_i$ lipids. (D-G)

Graphs representing quantitation of (D) mitochondrial PE and (G) cardiolipin levels show the

mean $\pm$ S.E.M. for n=6 biological replicates. Significant differences compared to (*) WT or (#)

psd1Δpsd2Δ were calculated by one-way ANOVA. (E) Representative TLC plate for cellular $^{32}P_i$

lipids. (F) Quantitation of cellular PE levels representing the mean $\pm$ S.E.M. for n=6 biological

replicates. Significant differences compared to (*) WT or (#) *psd1Δpsd2Δ* were calculated by

student *t*-test.

Fig 7. Depletion of PE through deletion of Cho1p impairs complex III and complex IV

activities. (A) Metabolic pathways tied to PS synthesis by Cho1p in yeast. (B) Detection of the

β subunit of Psd1p and Cho1p expression were verified in yeast whole cell extracts of the

indicated strains by immunoblot. Kgd1p served as a loading control. *Cho1p, phosphorylated

Cho1p. (C) The indicated strains were spotted onto synthetic complete dextrose (SCD) medium

+/- 2mM ethanolamine (+E) and incubated at 30°C for 2 days. (D-J) Mitochondrial phospholipids

from the indicated strains were labeled overnight with $^{32}P_i$, separated by TLC, and quantitated

by phosphoimaging. All graphs show the mean $\pm$ S.E.M. for n=6 biological replicates.

Significant differences compared to (*) WT or (#) *psd1Δpsd2Δ* were calculated by one-way

ANOVA. (K) The indicated strains were spotted and incubated at 30°C for 2 days on YPD and

for 3 days on rich lactate (RL), and SCEG +/- 2mM ethanolamine (+E). (L,M)

Spectrophotometric assay following the (L) reduction and (M) oxidation of cytochrome *c* at

550nM in isolated mitochondria solubilized in 0.5% (w/v) DDM. Analysis *versus* WT by one-way

ANOVA $\pm$ S.E.M. for n=6 or n=3 for ρ^0. (N) Steady state expression of mitochondrial proteins in

mitochondria isolated from the indicated strains.

Fig 8. A PE-coordinating residue in Qcr7p is important for complex III activity. (A) The

crystal structure of yeast cytochrome bc_1 that modeled associated lipids was downloaded using

PDB ID: 1KB9. Using PyMOL, the region containing the catalytic subunit Cob1p (magenta) near

the matrix facing surface was enlarged to demonstrate hydrophobic interactions between this

subunit and the acyl chains of PE. Arg51 of Qcr8p (orange) also shows hydrophobic interactions

with a carbon atom from the ethanolamine headgroup. Glu82 of Qcr7p (blue) was predicted to

form a hydrogen bonding interaction (3.4 Å distance) with the amine group of PE, whose atoms

are depicted as spheres; *gray*: carbon, *red*: oxygen, and *blue*: nitrogen (hydrogen atoms are not

represented). (B) WT and mutant Qcr7p was detected in yeast whole cell extracts of the indicated

strains by immunoblot; Aac2p served as a loading control. (C) Mitochondria from the indicated

strains were immunoblotted for subunits of complex III and complex IV as well as markers for the

indicated mitochondrial compartments. (D)The indicated strains were spotted and incubated at

30°C for 2 days on YPD and for 3 days on rich lactate (RL) and SCEG. (E, F) Spectrophotometric

assay following the (E) reduction or (F) oxidation of cytochrome *c* at 550nm in isolated

mitochondria solubilized in 0.5% (w/v) DDM. Analysis *versus* WT (*) was performed by one-way

ANOVA $\pm$ S.E.M. for n=6 or n=3 for ρ^0. P values for decreases that didn't achieve significance are

reported and were evaluated by student t-test *versus* WT.

TABLES

TABLE 1 - Strains used in this work		
Strain	**Genotype**	**Source**
Parental		
GA74-1A	*MAT*a, *his3-11,15, leu2, ura3, trp1, ade8 [rho+, mit+]*	Carla Koehler
Deletions		
psd1Δ	*MAT*a, psd1Δ::HIS3MX6, leu2, ura3, trp1, ade8 [rho+, mit+]*	Ref. 29
psd2Δ	*MAT*a, psd2Δ::HIS3MX6, leu2, ura3, trp1, ade8 [rho+, mit+]*	Ref. 29
psd1Δpsd2Δ	*MAT*a, psd2Δ::HIS3MX6, leu2, ura3, psd1Δ::TRP1, ade8 [rho+, mit+]*	Ref. 29
crd1Δ	*MAT*a, his3-11,15, leu2, ura3, crd1Δ::TRP1, ade8 [rho+, mit+]*	Ref. 67
ρ⁰	*MAT*a, his3-11,15, leu2, ura3, trp1, ade8 [rho-, mit+]*	This study
cho1Δ	*MAT*a, cho1Δ::HIS3MX6, leu2, ura3, trp1, ade8 [rho+, mit+]*	Ref. 29
Integrations		
psd1Δpsd2Δ::IM-Psd1	*MAT*a, psd2Δ::HIS3MX6, Psd3XFLAG::LEU2, ura3, psd1Δ::TRP1, ade8 [rho+, mit+]*	Ref. 29
psd1Δpsd2Δ::OM-Psd1	*MAT*a, psd2Δ::HIS3MX6, Tom20-Psd3XFLAG::LEU2, ura3, psd1Δ::TRP1, ade8 [rho+, mit+]*	This study
psd1Δpsd2Δ::ER-Psd1	*MAT*a, psd2Δ::HIS3MX6, CPY*mPsd3XFLAG::LEU2, ura3, psd1Δ::TRP1, ade8 [rho+, mit+]*	Ref. 29
HI-CRISPR Edited		
cho1Δpsd1Δpsd2Δ	*MAT*a, psd2Δ::HIS3MX6, leu2, ura3, psd1Δ::TRP1, *cho1 HI-CRISPR ade8 [rho+, mit+]*	This study
Qcr7p E82R	*MAT*a, his3-11, 15, leu2, ura3, trp1, ade8 [rho⁺, mit⁺] *Qcr7pᴱ⁸²ᴿ HI-CRISPR*	This study
Qcr7p E82D	*MAT*a, his3-11, 15, leu2, ura3, trp1, ade8 [rho⁺, mit⁺] *Qcr7pᴱ⁸²ᴰ HI-CRISPR*	This study
Parental/Deletion W303	*MAT*a, his3-11, ura3-1,15, trp-1-1, ade2-1, can1-100 [rho+, mit+]*	Cathy Clarke
psd1Δ W303	*MAT*a, psd1Δ::HIS3MX6, ura3-1,15, trp1-1, ade2-1, can1-100*	This study
D2730-10B	*MAT*a [rho+, mit+]*	Carla Koehler
psd1Δ D273-10B	*MAT*a, psd1Δ::KANMX6, [rho+, mit+]*	This study
BY4741	*MAT*a, his3-1, le2-0, met15-0, ura3-0 [rho+, mit+]*	Euroscarf
psd1Δ BY4741	*MAT*a, psd1Δ::KANMX4,his3-1, le2-0, met15-0, ura3-0 [rho+, mit+]*	Euroscarf

Supplementary Figures

A. Cell Fractionation

SM - starting material
P13 - 13,000 x g pellet (mitochondria)
P40 - 40,000 x g pellet (ER/endosomes)
S40 - 40,000 x g supernatant (cytosol)

Supplementary Fig 1. ER-Psd1p co-fractionates with ER/endosomal compartment. (A) Following growth in rich lactate medium to late log phase, fractions were collected from the indicated yeast strains by differential gravity centrifugation. Equal protein amounts from each collected fraction were resolved by SDS-PAGE and immunoblotted for Psd1p (β and α subunits) and markers for each compartment (Kgd1p for mitochondria/P13, Sec62p for the ER/P40, and Hsp70 for the cytosol/S40).

Supplementary Fig 2. TLC images of $^{32}P_i$ labeled lipids. Representative TLC plates from (A) Cellular and (B) mitochondrial lipid extractions. The migration of different phospholipid classes is indicated.

991
992

Supplementary Fig 3. Cho1p expression and phosphorylation state is increased in *psd1Δpsd2Δ* yeast, which also indicate hallmarks of ER stress through elevated levels of Kar2p. Yeast strains were cultured in rich lactate medium for 2 days at 30°C. Cells were harvested by centrifugation, lysed, protein extracted, resuspended in Laemelli buffer, and analyzed by western blot. (B-I) The expression of the indicated proteins was normalized relative to WT and statistical differences compared to WT (*) determined by one-way ANOVA ± S.E.M. for n=6.

Supplementary Fig 4. Phospholipid analysis of sucrose gradient purified mitochondria indicates equivalent levels of PE in OM-Psd1 and ER-Psd1 mitochondria. (A) After crude isolation of mitochondria by gravity centrifugation, samples were overlaid on a two-step sucrose gradient containing 32% and 60% sucrose in EM buffer and subjected to ultracentrifugation at 134,000 x *g* for 1 hour at 4°C. Purification of mitochondria was monitored by using the mitochondrial marker Tom70p and the endosomal marker Dpm1p. (B) Total phospholipid content/mitochondrial protein (mean ± S.E.M., n =4) in sucrose purified mitochondria. (C-F) Quantitation of phospholipid levels after separation by TLC and visualization by molybdenum blue staining. Analysis *versus* WT (*), *psd1Δpsd2Δ* (#), or IM-Psd1 ($) was performed by one-way ANOVA ± S.E.M. for n=4. (G) Representative TLC plate of phospholipid extraction from sucrose purified mitochondria followed by visualization using molybdenum blue reagent.

Supplementary Fig 5. Mitochondrial morphology is not overtly affected by disruption or alteration of Psd1 pathway. Cells from the indicated strains were analyzed by transmission electron microscopy. A) GA74-1A parental wildtype strain, B) *psd1Δ*, C) *psd2Δ*, D) *psd1Δpsd2Δ*, E) *psd1Δpsd2Δ*::IM-Psd1, F) *psd1Δpsd2Δ*::OM-Psd1, G) *psd1Δpsd2Δ*::ER-Psd1. *n,* nucleus; *m,* mitochondria; and *v*, vacuole. *Bars,* 0.5 μm.

Supplementary Fig 6 (continued on next page)

Supplementary Fig 7 (continued on next page)

Supplementary Fig 7. Quantitation of CoQ synthome subunits and additional mitochondrial proteins. Densitometry analysis of steady state protein amounts in isolated mitochondria (30 μg) from the indicated strains (representative immunoblots shown in Fig 5F). Analysis *versus* WT (*) or *psd1Δpsd2Δ* (#) was performed by one-way ANOVA $\pm$ S.E.M. for n=4.

Supplementary Fig 8 (continued on next page)

Supplementary Fig 8. CoQ$_6$ supplementation does not rescue the *psd1*Δ and *psd1*Δ *psd2*Δ growth defects on respiratory medium. OD$_{600}$ measurements were recorded every 30 minutes for a period of 48 hours at 30°C for yeast grown in (A) SCEG, (B) SCEG + 2mM ethanolamine (eth), (C) SCEG + 2μM CoQ$_6$, (D) SCEG + 2μM CoQ$_6$ + 2mM ethanolamine, (E) SCEG + 10μM CoQ$_6$, and (F) SCEG + 10μM CoQ$_6$ + 2mM ethanolamine. ± S.E.M. for n=2.

1001

Supplementary Fig 9. The *para*-amino benzoic acid pathway for CoQ$_6$ biosynthesis is not necessary for *psd1Δ* growth. (A) Pathways for CoQ Biosynthesis; 4-hydroxy phenylpyruvate (4-HBz), 4-hydroxybenzoate (4-HB), 3-polyprenyl-4-hydroxybenzoate (PPHB), CoQ$_6$H$_2$, 3-hexaprenyl-4-aminobenzoate (PPAB), *para*-amino-benzoate (PABA). The indicated Spearman correlation coefficient (rs) for 4-HB molecules that share a negative correlation with Psd1p molecules, was derived from the Yeast 3 Thousand (Y3K) dataset [41]. (B) The indicated strains were spotted and incubated at 30°C for 4 days on SCEG +/-PABA and +/- 2mM ethanolamine (+E).

1002

Cellular Steady State Phospholipids - Choline/Ethanolamine Supplemented

Mitochondrial Steady State Phospholipids - Choline/Ethanolamine Supplemented

Supplementary Fig 10. Cellular and mitochondrial phospholipid profiles from choline and ethanolamine supplemented *psd1Δpsd2Δ* yeast. (A-D) Cellular and (E-H) mitochondrial phospholipids from the indicated strains were labeled overnight with $^{32}P_i$ and separated by TLC. All graphs show the mean ± S.E.M. for n=6 biological replicates. Significant differences *versus* WT (*) or *psd1Δpsd2Δ* (#) were calculated by one-way ANOVA. Key for number of symbols represented for statistical analysis interpretation 1 symbol = $p < 0.05$, 2 symbols = $p < 0.01$, 3 symbols = $p < 0.001$, and 4 symbols = $p < 0.0001$.

1003
1004