Cytokinin perception in potato: New features of canonic players

Sergey N. Lomin[1], Yulia A. Myakushina[1], Oksana O. Kolachevskaya[1], Irina A. Getman[1], Dmitry V. Arkhipov[1], Ekaterina M. Savelieva[1], Dmitry I. Osolodkin[1-3] and Georgy A. Romanov[1,4] *

[1] Timiryazev Institute of Plant Physiology, Russian Academy of Sciences, Botanicheskaya 35, 127276 Moscow, Russia

[2] Institute of Pharmacy and Translational Medicine, Sechenov First Moscow State Medical University, Trubetskaya 8, 119991 Moscow, Russia

[3] Institute of Poliomyelitis and Viral Encephalitides, Chumakov FSC R&D IBP RAS, Poselok Instituta Poliomelita 8 bd. 1, Poselenie Moskovsky, Moscow 108819, Russia

[4] Belozersky Institute of Physico-Chemical Biology, Lomonosov Moscow State University, Leninskie Gory 1, Bld. 40, 119992 Moscow, Russia

* To whom correspondence should be addressed. E-mail: gromanov@yahoo.com or gar@ippras.ru.
Tel.: 7(499)6785432

Running title: **Cytokinin perception in potato**

Abstract

Potato is the most economically important non-cereal food crop. Tuber formation in potato is regulated by phytohormones, cytokinins in particular. The present work was aimed to study cytokinin signal perception in potato. The sequenced potato genome of doubled monoploid Phureja was used for bioinformatic analysis and as a tool for identification of cytokinin receptors from autotetraploid potato cv. Désirée. All basic elements of multistep phosphorelay (MSP) required for cytokinin signal transduction were identified in Phureja genome, including three genes orthologous to three cytokinin receptor genes (*AHK 2-4*) of Arabidopsis. As distinct from Phureja, autotetraploid potato contains at least two allelic isoforms of each receptor type. Putative receptor genes from Désirée plants were cloned, sequenced and expressed, and main characteristics of encoded proteins, firstly their consensus motifs, structure models, ligand-binding properties, and the ability to transmit cytokinin signal, were determined. In all studied aspects the predicted sensor histidine kinases met the requirements for genuine cytokinin receptors. Expression of potato cytokinin receptors was found to be organ-specific and sensitive to growth conditions, particularly to sucrose content. Our results provide a solid basis for further in-depth study of cytokinin signaling system and biotechnological improvement of potato.

Key words: CHASE domain-containing histidine kinase, cytokinin, cytokinin receptor, cytokinin

signaling, gene expression, hormone perception, potato, *Solanum tuberosum.*

Abbreviations: BA, 6-benzyladenine; CHASE, Cyclases/Histidine kinases Associated SEnsory; CHK,

CHASE domain-containing histidine kinases; CRF, cytokinin response factor; cZ, *cis*-zeatin; DI,

dimerization interphase; DZ, dihydrozeatin; GFP, green fluorescent protein; HK, histidine kinase; iP,

isopentenyladenine; LacZ, galactosidase; LD, long day; MSP, multistep phosphorelay; RR, response

regulator; SNP, single nucleotide polymorphism; TD, thidiazuron; TM, transmembrane; tZ, *trans*-zeatin.

Introduction

Potato is a widespread practically important crop, its tuber formation is controlled by phytohormones (reviewed in Aksenova *et al*., 2012, 2014). Previous studies have shown that cytokinins and auxins can accelerate and enhance potato tuber formation (Aksenova *et al*., 2000; Romanov *et al*., 2000; Roumeliotis *et al*., 2012; Kolachevskaya *et al*., 2015, 2017). In non-tuberizing plants (tobacco, tomato), increased doses of active cytokinins stimulate morphogenesis, in many aspects resembling tuber formation (Guivarc'h *et al*., 2002; Eviatar-Ribak *et al*., 2013). Cytokinin signaling is also involved in the formation of nodules on the roots of legumes (reviewed in Frugier *et al*., 2008; Miri *et al*., 2016). Cytokinins largely determine the nature of source-sink relationships in the whole plant, enhancing the attracting ability of the tubers (Abelenda and Prat, 2013). Elevated doses of cytokinins affect the overall architectonics of potato plants, suppressing the root development (Aksenova *et al*., 2000). In addition, cytokinins participate in plant defense against biotic and abiotic adverse factors (Zwack and Rashotte, 2015; Brütting *et al*., 2017; Thu *et al*., 2017). All the above indicates the important role of cytokinins in both the formation of tubers and the general development and resistance of potato plants.

The molecular mechanism of cytokinin action on a plant cell has been established using mainly the Arabidopsis model (reviewed in Hutchison and Kieber, 2002, Hwang *et al*., 2002; Kakimoto, 2003; Heyl and Schmülling, 2003; Sakakibara, 2006; Müller and Sheen, 2007). This mechanism is based on multistep phosphorepay (MSP) and uses three protein species to bring the cytokinin signal up to the primary response genes: (i) transmembrane catalytic receptors with histidine kinase activity, (ii) mobile phosphotransmitters circulating between the cytoplasm and nucleus, and (iii) nuclear transcription factors, B-type response regulators. Other proteins (CRFs, pseudophosphotransmitters, A-type response regulators) affect the intensity of the cytokinin signaling through the main transmission pathway (Kieber and Schaller, 2014).

Receptors are key factors in the perception and transduction of hormonal signals. In the case of cytokinins, receptors are sensory hybrid histidine kinases largely homologous to bacterial sensory histidine kinases, members of two-component signal transduction system. Known cytokinin receptors are multidomain proteins located mainly in ER membranes (Caesar *et al*., 2011; Lomin *et al*., 2011, 2018; Wulfetange *et al*., 2011; Daudu *et al*., 2017; Ding *et al*., 2017) with N-terminal hormone-binding sensory module localized in the ER lumen and the central and C-terminal catalytic domains protruding in the cytosol (Steklov *et al*., 2013; Lomin *et al*., 2018). Until now, cytokinin receptors have been studied in a few vascular plant species, primarily and most detailed in Arabidopsis and maize (Kakimoto, 2003; Yonekura-Sakakibara *et al*., 2004; Romanov *et al*., 2006; Lomin *et al*., 2011, 2012, 2015; 2018; Stolz *et al*., 2011; Heyl *et al*., 2012; Steklov *et al*., 2013; Wang et al., 2014). In recent years, cytokinin receptor studies have been extended to new species including rice (Choi *et al*., 2012;

Ding *et al*., 2017), barrelclover (Held *et al*., 2014), *Medicago truncatula* (Laffont *et al*., 2015; Boivin *et al*., 2016), oilseed rape (Kuderová *et al*., 2015), *Nicotiana attenuata* (Schäfer *et al*., 2015), and apple (Daudu *et al*., 2017). These studies have demonstrated that the cytokinin perception apparatus in some aspects is species-specific. Potato differs from most plant species by its ability to form tubers. This process, sensitive to various cues including cytokinins, makes the study of cytokinin receptors of potato especially intriguing. So far, to our knowledge, there have been no scientific reports on such studies.

In this paper, we have examined potato cytokinin receptors of a homozygous doubled monoploid Phureja (DM1-3 516 R44) whose genome was sequenced several years ago (Potato Genome Sequencing Consortium, 2011). Cloning and expression of receptor encoding genes were conducted using the commercial autotetraploid potato cv. Désirée. The presence of all necessary MSP elements in potato was demonstrated and main characteristics of CHASE domain-containing cytokinin receptors, primarily their consensus motifs, 3D structures, ligand-binding properties, and the ability to transmit the signal by MSP were ascertain. In contrast to the Phureja monoploid, distinct alleles for each of the three main forms of receptors were found in the Désirée potato. Expression of cytokinin receptor genes was shown to be organ-specific and affected by sucrose. The obtained results might serve as a framework for new biotechnological approaches in improving potato productivity and stress resistance.

Materials and methods

Sequence analysis

Nucleotide/polypeptide sequences of cytokinin receptors and other proteins related to the cytokinin signaling were retrieved from databases NCBI (National Center for Biotechnology Information, http://www.ncbi.nlm.nih.gov), Phytozome 11 (https://phytozome. jgi.doe.gov/pz/portal.html), MSU Rice Genome Annotation Project Release 7 (http://rice.plantbiology.msu.edu/) and congenie.org (http://congenie.org/) using the BLASTP tool and *AHK2* (AT5G35750), *AHK3* (AT1G27320), *AHK4* (AT2G01830) and other cytokinin-related genes of *Arabidopsis thaliana* as templates. Domain structure of proteins was determined with PROSITE (http://prosite.expasy.org/). Transmembrane domains were determined using MESSA service (http://prodata.swmed.edu/MESSA/MESSA.cgi) (Cong and Grishin, 2012). Domain visualization was performed using the MyDomains – Image Creator service (http://prosite.expasy.org/mydomains/).

Phylogenetic analysis was performed using the MEGA6.0 (Tamura *et al*., 2013). Alignment of nucleotide sequences (CDS, codon mode) was performed by ClustalW algorithm. Method of maximum likelihood was employed for phylogenetic reconstruction. The search for key amino acids in receptor domains by alignment and visualization of protein sequences was carried out in Clustal X2.1 (Larkin *et al*., 2007) and Jalview (Clamp *et al*., 2004), respectively.

Homology modeling

Search of templates for homology modeling was performed at SWISS-MODEL web-service (https://swissmodel.expasy.org/) (Biasini *et al*., 2014). Modeling of potato (*Solanum tuberosum* L.) protein structures was accomplished in Modeller 9.15 (https://salilab.org/modeller/) (Sali and Blundell, 1993) using *automodel* class for comparative modeling. For each protein, 200 models were built, and the best model was selected according to DOPE score value (Shen and Sali, 2006) as determined by Modeller. For StHK receiver domains, chain A AHK5rd-AHP1 complex structure (Bauer *et al*., 2013) (PDB ID: 4EUK) and CKI1rd structure (Pekárová *et al*., 2011) (PDB ID: 3MMN) were used as templates. ERS1 histidine kinase domain (PDB ID: 4MT8) and ETR1 H-ATPase domain (PDB ID: 4PL9) structures (Mayerhofer *et al*., 2015) were used as templates for StHK catalytic module. StHK sensor modules were modeled using the structure of AHK4 receptor (PDB ID: 3T4L) as the template (Hothorn *et al*., 2011). After adding hydrogen atoms, models were energy minimized in USCF Chimera 1.10.2 (http://www.cgl.ucsf.edu/chimera/) (Pettersen *et al*., 2004) using AMBER ff14SB force field (Maier *et al*., 2015) with 300 steps of steepest descent and 300 steps of conjugate gradient; step size was 0.02 Å in both cases. Stereochemical quality of the models was assessed with ProCheck (Laskowski *et al*., 1993) implemented in PDBsum Web service (www.ebi.ac.uk/pdbsum), ProSA-web (https://prosa.services.came.sbg.ac.at/prosa.php) (Wiederstein and Sippl, 2007) and QMEAN server (https://swissmodel.expasy.org/qmean/help) (Benkert *et al*., 2009). Visualization and superposition of the models were accomplished with UCSF Chimera.

Promoter analysis

Promoter regions of *Arabidopsis thaliana* cytokinin receptor genes (*AHK2*, *AHK3* and *AHK4*) were obtained from TAIR database (https://www.arabidopsis.org). Identification of promoter regions of cytokinin receptor genes (*StHK2*, *StHK3* and *StHK4*) of potato was performed using Phytozome 11 and NCBI databases. DNA sequence of 1000 nucleotides long upstream the gene transcription start was taken as a promoter region. The search for *cis*-regulatory elements in promoters of studied genes was carried out using PLACE (http://www.dna.affrc.go.jp/htdocs/PLACE/) and PlantCARE (http://bioinformatics.psb.ugent.be/webtools/plantcare/html/) programs.

Receptor cloning

Experiments were performed with autotetraploid potato (*Solanum tuberosum*) plantlets of Désirée variety. Plants were propagated by *in vitro* cloning on Murasige-Skoog agarous medium supplemented with 1.5% sucrose, at 20 °C and 16 h photoperiod in a controlled climate chamber with luminescent white light illumination (Kolachevskaya *et al*., 2015, 2017). Total RNA was isolated from fresh or immediately frozen plant material. To remove genomic DNA, RNA samples were treated with RNase-

free DNase I (Thermo Scientific). For reverse transcription, 2 μg of RNA and 200 units of RevertAid ™ were used according to the manufacturer's instructions (Thermo Scientific). Total DNA was isolated from the leaves of the potato using the CTAB method. The resulting cDNA and total DNA were used to amplify genes encoding predicted potato cytokinin receptors with high-precision Phusion High-Fidelity DNA polymerase (Thermo Scientific). The primer design was performed to amplify the full-length and truncated (sensory modules with flanking transmembrane helices) CDS of cytokinin receptors according to sequences in the NCBI Genbank XM_015303261.1, XM_006352114.2 and XM_006354988.2. PCR products were gel purified and cloned using the PCR Cloning Kit (Thermo Scientific) into the plasmid pJET1.2/blunt according to the manufacturer's instructions followed by transformation of *E. coli* strain DH10B (Invitrogen). *StHK4* was amplified using StHK4_truncated primers. The product was inserted into the construction of pB7FWG2-AHK3 instead of *AHK3*. The latter was removed at the BcuI and EcoRI restriction sites (Lomin *et al*., 2015). The nucleotide sequences of the cloned genes were confirmed by DNA sequencing.

StHK2 and StHK3 sequences were subcloned into the plasmid pDONR^{TM}221 in BP reaction with Gateway® BP Clonase® II Enzyme mix (Thermo Scientific). Then, using the LR reaction with the LR Clonase® II Plus enzyme (Thermo Scientific), the cloned sequence was transferred into the expression vector pB7FWG2 (Karimi *et al*., 2007) where it was fused at the 3'-terminus to the *eGFP* gene. For expression in *E. coli*, *StHK2* and *StHK4* were amplified using primers StHK2_COLD and StHK4_COLD, respectively. The product was then inserted into the plasmid pCOLD IV (Takara BioInc.) at the XhoI and XbaI restriction sites for *StHK2* and SacI and EcoRI restriction sites for *StHK4*, followed by transformation of the *E. coli* DH10B strain.

Transient expression of receptor genes in tobacco plants

The transient transformation of tobacco (*Nicotiana benthamiana*) leaves was accomplished according to Sparkles *et al*. (2006). Eight week-old tobacco plants were infiltrated with a mixture of *Agrobacterium tumefaciens* carrying cytokinin receptor genes fused to GFP and the *A. tumefaciens* helper strain p19 (Voinnet *et al*., 2003), and the expression of receptor genes was checked after 5–6 days on fluorescence microscope Axio Imager Z2 (Carl Zeiss Microscopy GmbH) before leaves were proceeded further for microsome isolation.

Plant membrane isolation

All manipulations were done at 4 °C. Tobacco leaves 6 days after infiltration were homogenized in buffer (3 ml per 1 g of fresh weight) containing 100 mM Tris-HCl (pH 8.0), 2 mM Na$_2$-EDTA, 50 mM KCl, 1 mM DTT and 1 mM PMSF. The homogenate was filtrated through Miracloth (Calbiochem, San Diego, USA), and the filtrate was centrifuged for 5 min at 5000 *g*. Then supernatant was centrifuged for

40 min at 15000 *g*. The microsome pellet was resuspended in 50 mM KCl, 10% glycerol and then

microsome suspension was stored at -70 °C.

Hormone binding assays

Ligand binding studies were performed in PBS as described previously (Romanov *et al*., 2005; Lomin *et*

al., 2015). Studies of pH influence on hormone binding were performed in 50 mM MES-KOH (pH 5–7)

or Tris-HCl (pH 7–9) buffers with 50 mM KCl. K_D for [³H]*t*Z binding to different receptors were

determined in saturation assays followed by data analysis in Scatchard plots.

Assessment of receptor functionality

Plasmids pCOLD IV with *StHK2* or *StHK4* coding sequences were transferred for the expression into *E.*

coli strain KMI001 (Suzuki *et al*., 2001). In this strain, HK receptor→YojN→RcsB→*cps::lacZ* pathway

can be activated by external cytokinin (Takeda *et al*., 2001). The activation of the signaling pathway

was monitored by measuring β-galactosidase activity of *E. coli* cells. Cultivation of clones on Petri

dishes containing 40 mM glucose, 40 µg ml⁻¹ X-gal, 100 µM IPTG, 50 µg ml⁻¹ ampicillin at 15 °C was

performed for 4 days. The individual clones were then streaked onto new Petri dishes containing 40 mM

glucose, 40 µg ml⁻¹ X-gal, 100 µM IPTG, 50 µg ml⁻¹ ampicillin ± *trans*-zeatin at a concentration of 0.5

µM. The clones were grown for 3 days at 15 °C. Expression of the *cps::lacZ* construct was evaluated by

blue staining of bacterial clones.

Gene expression analysis

Potato (*Solanum tuberosum* cv. Désirée) plants were cultivated under standard *in vitro* conditions at a

long (16 h) day for 5-6 weeks on liquid MS medium containing 1.5% or 5% sucrose. For hormone

treatment, the medium was replaced with the same one supplemented with 6-benzyladenine (BA, 1 µM).

Tubes were inverted several times to assure uniform plant wetting and then incubated for 1 h under

standard conditions. Finally plant organs (leaves, stems, roots, tubers) were isolated and immediately

frozen in liquid nitrogen. Control plants were treated in a same way only without hormone. Total RNA

was isolated by Trizol method (Brenner *et al*., 2005), this RNA served template for cDNA synthesis by

reverse transcription (Invitrogen). All RNA samples were treated with RNase-free DNase I. The

resulting cDNA was checked for the genomic DNA contamination by PCR with primers differentiating

cDNA and genomic DNA. The band derived from genomic DNA was absent in the separating gel.

Expression of genes encoding predicted proteins of cytokinin signaling system was determined by qRT

PCR. Potato housekeeping genes *StEF1*α (elongation factor 1-α, AB061263) and *StCYC* (cyclophilin,

AF126551) were used as reference genes (Nicot *et al*., 2005).

Statistical analysis

Statistical analysis was accomplished using the Student's t-test. *P*-value <0.05 was considered as
statistically significant. In tables and graphics, mean values with standard errors are presented.

Results

Monoploid Phureja genome analysis

Potato has everything necessary for cytokinin signaling via the MSP pathway

The search for protein sequences and encoding genes involved in cytokinin signaling was performed on
the basis of the duplicated potato monoploid Phureja genome (Potato Genome Sequencing Consortium,
2011). In general, all potential components of the canonical cytokinin signaling system described in
Arabidopsis and other plant species with a sequenced genome (Kieber and Schaller, 2014) were
identified in potato too. Potential cytokinin-related genes found in potato encode homologs of CHASE
domain-containing histidine kinases (CHK), phosphotransmitters (HPt), and response regulators of A
(RR-A) and B (RR-B) types (Table 1). This indicates the MSP functioning in potato cells for cytokinin
signal transduction, involving proteins of a two-component system. In the potato monoploid proteome
three predicted protein-coding sequences XP_015158747.1, XP_006352176.1 and XP_006355050.1,
orthologous to Arabidopsis receptors AHK2, AHK3 and CRE1/AHK4, respectively, were detected. By
analogy with the Arabidopsis orthologs, these proteins were annotated in NCBI as StHK2, StHK3, and
StHK4. They correspond to mRNA sequences XM_015303261.1, XM_006352114.2 and
XM_006354988.2. Deduced proteins StHK2, StHK3, and StHK4 share 59.35%, 67.75%, and 67.52%
sequence similarity with the Arabidopsis orthologs. The lengths of *StHK2*, *StHK3* and *StHK4* genes are
5345, 4216, and 3810 bp, respectively, and predicted proteins are 1263, 1032, and 992 amino acids long
(Table 1).

Phylogenetic analysis classified StHKs into three clades

The phylogenetic analysis was performed to compare the conserved and unique features of potato
cytokinin receptors with the features of Arabidopsis, rice, tomato and other species receptors (Fig. 1).
Cytokinin receptors of flowering plants can be grouped into three main clades, corresponding to the
Arabidopsis AHK2, AHK3, and CRE1/AHK4 receptors (Pils and Heyl, 2009; Lomin *et al*., 2012;
Steklov *et al*., 2013). Potato and tomato receptors are unequivocally distributed among these three

clades. Evolutionally, they are closer to Arabidopsis than to rice receptors, what was expected since

potato, tomato and Arabidopsis are dicots whereas rice is a monocot.

Multiple alignments revealed common and unique features of StHKs

We investigated the modular architecture of predicted potato cytokinin receptors. The exon-intron

structure of the cognate genes as well as occurrence and position of functional domains in the receptor

proteins were analyzed. Known cytokinin receptors share a common organization, including (from N to

C termini) sensory module with CHASE domain, catalytic module with histidine kinase and ATPase

domains, and receiver module with pseudoreceiver and receiver domains (Kakimoto, 2003; Steklov *et*

al., 2013). The sensory module is flanked by predicted transmembrane (TM) α-helices. There is always

a single TM-helix downstream of module while the number of TM-helices upstream of module is

variable. Number of upstream TM-helices is usually maximal (up to 3-4) in AHK2 clade members,

minimal (1) in the AHK4 clade and intermediate in the AHK3 clade (Steklov *et al*., 2013). The domain

structure of potato receptors fully corresponds to the canonical one (Fig. 2).

At the N-termini of potato cytokinin receptors, the number of upstream TM helices is 3, 2, and 1 in

StHK2, StHK3, and StHK4, respectively. Cytokinin receptor genes share similar exon-intron

organization. The exon boundaries in the receptor genes of different species coincide in most cases. A

multiple alignment of receptor sequences from potato, rice and Arabidopsis was carried out (Fig. 3). All

canonical motifs present in known cytokinin receptors were also found in the potato orthologs. In the

catalytic domain of potato receptors, H̲, N, G1, F and G2 motifs were identified, in the receiver domain

– DD-D̲-K motifs. Conserved sequences contain phosphorylatable histidine (H̲) and aspartate (D̲)

residues. StHK2 has a conserved aspartate in its receiver-like domain (Rec-like), similarly to orthologs

from Arabidopsis (AHK2), tomato (SlHK2) and rice (OsHK3 and OsHK5). However, DD-D̲-K motif in

the Rec-like domain has little in common with the respective sequence in the Rec domain (Fig. 3C).

Highly conserved motifs were earlier found in sensory modules and downstream TM segments of

cytokinin receptors (Steklov *et al*., 2013). These motifs are obviously important for ligand binding and

transmembrane signal transfer. In potato receptors, these motifs are also present, although with some

peculiarities. In particular, StHK2 has a deviation from the canonic motif in CHASE domain, where

either Glu or Asp is located at position 90, while StHK2 has Gln at this position. StHK3 has a deviation

at the position 177, strongly conserved in the HK3-clade. This position is occupied by Phe in the

canonic motif, while in StHK3 by Leu. In the general HK-motif, there is either Phe or Tyr at position

177. StHK4 is distinguished by positions 83 (Ser instead of Ala) and 172 (Phe instead of Tyr) in

conserved motifs. Note that counterparts of Gln90, Leu177 and Ser83 are present also in tomato

genome, so these substitutions may be characteristic of Solanaceae family. Phe172 seems to be unique

for potato.

StHK functional domains adopt canonical 3D structures

We have built homology models of StHK receptors' sensory, catalytic, and receiver domain structure

(Fig. 4). High structural similarity of potato receptors with their Arabidopsis orthologs was observed as

expected. Key functional regions, such as ligand-binding sites, phosphorylation sites, ATP-binding sites

and dimerization interfaces, are particularly conserved. Sensory modules consisting of dimerization,

PAS and pseudo-PAS domains (the latter two comprise the CHASE domain) are very similar in

Arabidopsis and potato. StHK2 and StHK3 differ from StHK4 by an insertion of 14 and 17 residues,

respectively, in the region adjacent to the C-terminus of α3-helix (the first α-helix of the PAS domain).

This insertion apparently does not participate in the ligand recognition site and is unlikely to directly

affect the ligand-binding properties of the protein. Similar insertions are also present in AHK2 and

AHK3 receptors from Arabidopsis.

The catalytic modules include histidine kinase domains and H-ATPase domains. Histidine kinase

domains are formed by two α-helices and contain dimerization interface and phosphorylation site

(conserved histidine). H-ATPase domains including ATP-binding sites have a sophisticated structure

based on parallel/antiparallel β-sheets and α-helices. A large insert at the β2-β3 linker (more than 50

residues long) differs cytokinin receptors from bacterial histidine kinases and H-ATPase domain of the

ethylene receptor. This insert is located, however, on the opposite side from the ATP binding site. This

structural feature is inherent not only for potato receptors but also for cytokinin receptors of other

species.

The CKI1 histidine kinase receiver domain (RD), used as a template for cytokinin receptor RD,

adopts a fold typical for the REC (or CheY-like) superfamily proteins. It is formed by five α-helices and

β-sheet of five parallel strands. Two α-helices are located on one side of the β-sheet, and remaining

three on the other side. The same fold is characteristic of the RD model of the Arabidopsis CRE1/AHK4

receptor. As distinct from this, an additional small helix is present in the region between α3 helix and β4

sheet in RD models of other potato and Arabidopsis receptors except CRE1/AHK4. Conserved aspartate

residue, serving as a phosphate acceptor in RD, is located at the N terminus of the β3 sheet (Fig. 4).

Deviations from canonic CHASE motifs in sensory modules of potato cytokinin receptors evidently

do not alter 3D structures of the modules. Unusual Gln90 is far from the ligand-binding pocket of

StHK2, with sidechain directed to the dimerizing interface. Although the unusual Leu177 of StHK3 is

localized in the ligand-binding site, its sidechain is oriented to the opposite direction. The substitutions

in StHK4 seem to be more functional than in other potato receptors. Ser73 and Phe172 are localized in

the ligand-binding pocket periphery and their sidechains are oriented inwards. Hence, these latter

substitutions might somehow influence the ligand specificity of the receptor.

Experimental studies on autotetraploid potato cv. Désirée

Potato cv. Désirée possesses multiple alleles of **StHK** *genes*

A homozygous doubled monoploid Phureja (DM1-3 516 R44) is an artificial form of potato

phenotypically differing from commonly known diploid/tetraploid potato varieties (Potato Genome

Sequencing Consortium, 2011). Such differences in phenotype are underlain by considerable sequence

and structural genome variations between potato haplotypes. Therefore, the results of genome study of

monoploid Phureja do not mirror exactly more complex genomes of common potato cultivars.

Our experimental study of cytokinin receptors was performed using the autotetraploid potato cv.

Désirée, widely used for commercial purposes, as well as a model laboratory plant (Aksenova *et al*.,

2000; Kolachevskaya *et al*., 2015). We cloned the putative receptor genes using primers designed

according to Phureja gene sequence data. Distinct from Phureja genome, at least six genes of putative

cytokinin receptors were cloned from cDNA of Désirée plants. All these genes share a typical

module/domain structure characteristic of hybrid sensor histidine kinases (Figs. 2–4). According to their

sequence, encoded proteins fall pairwise into three known clades of cytokinin receptors (Table 2, Fig.

1). Thus, each form of cytokinin receptors from potato cv. Désirée consists of at least two close

isoforms encoded by natural receptor alleles. Sequencing of cloned genes revealed traits of both

similarity and divergence between Phureja and Désirée plants. The nucleotide sequences of HK2-clade

members *StHK2a* and *StHK2b* differ from the orthologous Phureja sequence by five and four

nucleotides (5 and 4 SNPs), respectively. At the protein level, StHK2a and StHK2b have three and two

amino acid substitutions, respectively, relative to Phureja receptor (Table 2).

Of two cloned genes of HK3-clade, *StHK3a* is identical to its counterpart of Phureja, whereas

StHK3b differs by 20 SNPs together with a 3-nucleotide deletion. These differences result in the

absence of one amino acid and nine substitutions in StHK3b compared to its Phureja ortholog. Similar

data were obtained for HK4-clade: *StHK4a* was fully identical to that of Phureja whereas *StHK4b*

differed by 28 SNPs and a 3-nucleotide deletion. Correspondingly, StHK4b differs from its Phureja

ortholog, as well as from StHK4a, by deletion of one amino acid and substitution of 13 ones (Table 2).

Analysis of amino acid sequences of the proteins showed that all putative histidine kinases of Désirée

potato retain the domains and consensus sequences typical for cytokinin receptors, despite aa

substitutions (Fig. 2). This indicates that all proteins encoded by the cloned StHK genes of tetraploid

potato plants can successfully function as cytokinin receptors.

StHKs have typical cytokinin-binding properties except StHK3 with distinct ligand specificity

To analyze ligand-binding properties of the receptors, a recently developed plant membrane assay system (Lomin *et al*., 2015) was used. Potato cytokinin receptor genes were cloned into pB7FWG2 vectors for transient expression in tobacco leaves. In the case of *StHK2* and *StHK4* genes, the full-length cDNA sequences were expressed, but in the case of *StHK3*, expression of the full-length receptor failed for unknown reasons. Instead of full-length receptor, we cloned a genomic sequence of the StHK3a sensory module flanked with transmembrane domains. From the transiently transformed tobacco leaves, a microsomal fraction enriched with individual potato receptors was obtained. The binding assays were conducted using this fraction and tritium-labeled cytokinin. In aggregate, we tested four receptors belonging to all three clades: StHK2a, StHK3a (sensory module, further designated as StHK3a$_{SM}$), StHK4a, and StHK4b.

First, we determined the pH-dependence of hormone binding to these receptors within the pH range of 5–9 (Fig. 5). All receptors exhibited maximal *trans*-zeatin binding at the neutral-mildly basic pH: StHK2a at pH 7.5, StHK3a$_{SM}$ at pH 7, StHK4a at pH 7.5–8, and StHK4b at pH 8–9. All receptors showed a decrease in ligand binding at acid pH: StHK2a and StHK3a$_{SM}$ reduced their binding at pH 5 compared to pH 7 by a factor of 2 and 5, respectively. Ligand binding by StHK4a and StHK4b decreased at pH 5 about three times compared to maximal values. Although the StHK3a receptor was represented in this study only by its sensory module, a control experiment with the full-length receptor StHK2a and its sensory module showed a similar pH-dependence of hormone binding (data not shown). This means that an isolated sensory module is sufficient to determine the pH-dependence of hormone binding by the receptor.

The interaction of a hormone with a receptor is characterized by the equilibrium dissociation constant (K_d) of the ligand-receptor complex. K_d values were determined by the dose-dependent binding of labeled *trans*-zeatin to receptors, the results were processed by the Scatchard method (Lomin and Romanov, 2008). All receptors demonstrated high affinity for *trans*-zeatin, with similar K_d at the nanomolar level (Table 3). The determined K_d values were close to the values of analogous constants for cytokinin receptors of other species (Lomin *et al*., 2012, 2015, Kuderova *et al*., 2015) and were well correlated with concentrations of active cytokinins *in planta* (Hirose *et al*., 2008).

Different cytokinins are usually present in the plant: *trans*- and *cis*-zeatins, isopentenyladenine, and dihydrozeatin. In addition to natural cytokinins, there are many synthetic ones. Receptors exhibit different affinities for these substances. We studied the ligand specificity of receptors in competitive experiments where binding of labeled cytokinin was carried out in the presence of various concentrations of certain unlabeled ligands. Based on the obtained competition curves, the apparent K_d values were determined for each ligand as was described (Lomin and Romanov, 2008). We analyzed the

interaction with the receptors of six cytokinins, including five natural ones as well as synthetic urea-type cytokinin thidiazuron (Table 3). The ligand specificity of the receptors showed much in common. All analyzed receptors had a high and nearly equal affinity for *trans*-zeatin and isopentenyladenine, apparent K_d ranging from 2.1 to 5.2 nM. All receptors bound *cis*-zeatin significantly weaker, with K_d over 100 nM. 6-Benzyladenine exhibited an intermediate affinity with K_d ranging from 40 to 60 nM. Regarding the two remaining cytokinins, the receptors showed significant differences. StHK3a$_{SM}$ bound dihydrozeatin with K_d ~21 nM, much stronger than other potato receptors (K_d ~170-230 nM). StHK2 and StHK3a$_{SM}$ showed a high affinity for thidiazuron (K_d=1.4 nM and 2.3 nM, respectively), whereas its affinity for StHK4a and StHK4b was much lower (K_d=12.6 nM and 17.2 nM, respectively). The cytokinin affinity ranking for the potato receptors are as follows: StHK2, TD>iP=tZ>BA>cZ>DZ; StHK3, TD>iP=tZ>DZ>BA>cZ; StHK4, iP=tZ>TD>BA>cZ>DZ. The preference profiles of StHK2 and StHK3a$_{SM}$ differ by DZ position, and from (almost identical) StHK4 isoforms by TD position. The greatest differences (in TD and DZ positions) were revealed between the StHK3 and StHK4 receptors. With respect to ligand specificity, StHK2 and StHK4 have much in common with their Arabidopsis orthologs, whereas StHK3 binds isopentenyladenine and 6-benzyladenine with much higher affinity than AHK3. Although the StHK3a receptor was represented in this study only by its sensory module, previous data showed that sensory module is sufficient to characterize the ligand preference of the full-length receptor (Stolz *et al*., 2011; Lomin *et al*., 2015).

StHKs are able to trigger signaling via MSP

The ability of the putative potato receptors to trigger cytokinin signaling was tested on *E. coli* ΔrcsC mutant devoid of its own RcsC hybrid histidine kinase and equipped with the *cps:LacZ* construct with the *LacZ* reporter gene driven by *cps* promoter (Suzuki *et al*., 2001; Takeda *et al*., 2001). This design allows assessment of the ability of hybrid histidine kinases to initiate signaling over the MSP pathway. Activation of MSP signaling in the bacteria leads to the expression of the reporter galactosidase (LacZ), whose activity is manifested by blueing of clones growing on X-Gal-supplemented medium. We expressed the cloned genes of the putative potato cytokinin receptors in *E. coli* ΔrcsC. In the clones expressing the receptors but not in the control clone, blue staining was observed (Fig. 6). The degree of blueing was greatly increased in the presence of cytokinin. It confirms the ability of the cloned potato receptors to transmit the cytokinin signal to the primary response genes via the canonic MSP pathway.

StHKs exhibit **in planta** *organ-specific expression pattern which has unique properties*

To assess the functionality of a gene *in vivo*, it is important to know the level and pattern of its expression in the living organism. We studied the expression of putative cytokinin receptor genes in organs of potato plants grown *in vitro* under conditions favorable for either vegetative growth (1.5% sucrose) or for tuber formation (5% sucrose). The mRNA contents of the *StHK2*, *StHK3* and *StHK4* genes was determined by the qRT PCR method. For the quantitative comparison of the expression profiles, intra-exon primers were selected for each tested gene. These primers were complementary to both alleles of the same clade owing to a great similarity of these gene sequences. The relative amounts of putative receptors of distinct clades in potato organs were judged by comparing the levels of transcripts of the cognate genes.

Expression levels differed significantly depending on receptor, organ and growth conditions (Fig. 7). Expression patterns were different in plants grown on media with low (1.5%) or high (5%) sucrose content. In the case of 1.5% sucrose medium, the highest expression of *StHK3* genes was observed in roots, while in the case of 5% sucrose medium, the maximal expression of *StHK3* tended to occur in leaves. In the low sucrose grown plants, *StHK4* gene was much weaker expressed in leaves than in stems or roots, whereas at the higher sucrose content levels of *StHK4* expression in different organs were equalized. In the *StHK2* group, noticeable organ-specific differences were detected when plants were grown on 5%, but not on 1.5% sucrose. The lowest expression level of all groups of receptors was usually observed in tubers compared to other organs (Fig. 7A).

Within each organ, expression of *StHK3* undoubtedly dominated in leaves, regardless of the sucrose content (Fig. 7B). Expression of *StHK2* and especially *StHK4* genes in leaves was much weaker. In stems grown on 1.5% sucrose, expression of *StHK4* prevailed, while the lowest expression was characteristic of *StHK2* genes. In the roots, expression of *StHK2* genes was relatively weak, whereas the genes of *StHK3* and *StHK4* clades were expressed actively and in almost equal proportions. A dissimilar pattern of expression was observed in plants grown on 5% sucrose. Here in addition to leaves, in all other organs tested (stems, roots, tubers) the expression of *StHK3* alleles prevailed too, though to a lesser extent than in leaves. Compared to the low-sucrose medium, 5% sucrose increased the relative expression of *StHK2* genes (in stems and roots), while decreasing the level of *StHK4* expression. Thus, unlike Arabidopsis, in potato plants there is evidently no dominance of StHK4 receptors in the roots, on the contrary, StHK3 receptors seem to dominate there when cultivating plants on tuber-inductive 5% sucrose. A common feature of potato and Arabidopsis is a very low expression of HK4 orthologs in leaves.

Although the primers used for qRT PCR did not distinguish closely related isoforms of the cytokinin receptor genes, it is still possible to approximately estimate the relative expression of these alleles. To achieve this goal, data on cDNA clone numbers can be used (Table 2). Within the same clade, relative quantity of cDNA clones harboring a distinct isoform should reflect the relative occurrence of cognate

mRNAs. According to the last column of Table 2 corresponding to aerial part of potato seedlings, two
mRNA isoforms of the HK2 clade were in the 1:1 ratio; among mRNA isoforms of HK3 clade, *StHK3a*
was approx. two-fold more frequent than *StHK3b*; in the case of HK4 clade, *StHK4b* was expressed
about one order of magnitude more intensively than *StHK4a*.

StHK promoter activity is hardly affected by cytokinins, in accordance with low* cis*-element content

Treatment of potato plants with 6-benzyladenine had a small effect on the expression of the cytokinin
receptor genes, and the hormonal impact, when occurred, was only local and not always reproduced. At
1.5% sucrose, the upregulation (on average, 2.5-fold) of *StHK4* expression was regularly recorded, but
only in leaves (Fig. 8). It can be stated that the effect of cytokinin on the expression of potato receptor
genes, if any, is mostly limited to *StHK4* and depends on both organ/tissue type and conditions of plant
cultivation.

To validate the results of cytokinin treatment experiments, the effect of cytokinin administration on
the transcript level of the genes of type A response regulator (*RR*-A) genes was analyzed. These genes in
other species (Arabidopsis, maize) represent genes of primary response to cytokinin, so it might be
expected that in potato too they would be responsive to cytokinin. Indeed, our experiments showed a
rapid and reliable increase in the expression of *StRR*-A genes, in contrast to the receptor genes, after
plant treatment with BA (Fig. 8). These results prove the reliability of design and implementation of
experiments and, on the other hand, corroborate the common mode of functioning of the cytokinin
signaling system in different plant species.

Analysis of promoter structures of the studied genes (Fig. 9) was generally consistent with the gene
expression data. Long cytokinin-sensitive *cis*-regulatory elements or blocks of 4 or more short elements
near the transcription start (~300 bp area) were found in promoters of almost all *StRR*-A, but not *StHK*
genes. Among the receptor genes, only *StHK4* has a block of 3 short cytokinin-sensitive *cis*-elements
near the start of transcription. It is possible that this block determines the responsiveness of *StHK4* to
cytokinin under certain conditions, as shown in Fig. 8. Though this promoter analysis was accomplished
using the genome sequence of var. Phureja, our experimental analysis of sequenced promoters of
Désirée plants showed a complete similarity in this respect of the promoters from these two potato lines.

Discussion

453 Plant morphogenesis, in particular tuberization, is based on spatiotemporal cell proliferation and
454 differentiation. The main biological effect of plant hormones cytokinins is the induction of cell divisions
455 (Sakakibara, 2006; Romanov, 2009), therefore cytokinins are important participants of morphogenetic

processes. Indeed, with regard to potato development, cytokinins were reported to accelerate and scale up tuber formation (Aksenova *et al*., 2000; Romanov *et al*., 2000). In non-potato plants, cytokinins alone were able to induce the emergence of tuber-like structures (Guivarc'h *et al*., 2002; Eviatar-Ribak *et al*., 2013; Frugier *et al*., 2008; Miri *et al*., 2016). Apart the impact on the formation of tubers, cytokinins are known to regulate overall plant architecture, biomass partitioning as well as resistance to biotic and abiotic stress-factors (Aksenova *et al*., 2000; Abelenda and Prat, 2013; Zwack and Rashotte, 2015; Brütting *et al*., 2017; Thu *et al*., 2017). All the above points to the importance to investigate cytokinin signaling system in plants, in particular in bulbous crops like potato.

Herein, we present first results of detailed study of cytokinin receptors from potato plants. Two different potato forms were examined: doubled monoploid Phureja and tetraploid potato of Désirée variety. Phureja plants possess, like Arabidopsis, three cytokinin receptor orthologs. By contrast, in Désirée plants two allelic forms of each receptor type (StHK2a/b, StHK3a/b and StHK4a/b) have been found belonging to the three known phylogenetic clades. It is not excluded that the real number of receptor alleles in Désirée plants is somewhat higher. Within each group, receptor isomers differ by a few amino acid substitutions which do not affect most conserved motifs. However, some recently revealed consensus motifs in sensory module (Steklov *et al*., 2013) are distinctive in receptors of potato plants. The reason for such peculiar properties is not yet clear. Molecular modeling was employed to build all main domains of potato cytokinin receptors. In general, potato cytokinin receptors share similar domain structure with crystallized hybrid histidine kinases from other species. Note that such a complete characterization of all main domains of cytokinin receptors is presented for the first time.

The ligand-binding properties of individual potato receptors have been determined: affinity constants for active cytokinins, pH-dependence of ligand binding, ligand specificity. Two of the studied receptors (StHK3a and StHK4a) are identical in potato cv. Désirée and var. Phureja. All receptors have high affinity for tZ, significantly lower for BA, and relatively low for cZ. StHK3 differs from other potato receptors by relatively high affinity for DZ. All receptors bind cytokinin stronger in basic (pH 7–9) than acidic (pH 5–7) pH range. This indicates the intracellular functioning of potato cytokinin receptors (Romanov *et al*., 2018). The ligand-binding properties of StHK3 differ from those of orthologs in Arabidopsis, maize and oilseed rape: contrary to the other orthologs of this clade, StHK3 strongly binds iP with the affinity similar to the affinity for tZ. Using the special *E. coli* strain, the functionality of cloned potato receptors was confirmed by testing their ability to transduce cytokinin signal via MSP up to the target gene.

Expression of cytokinin receptor genes in various organs of potato plants and under different growing conditions was studied. The predominant expression of the *StHK3* genes was revealed in leaves, as well as in other organs of plants grown on 5% sucrose, although the degree of dominance of StHK3 was less pronounced in stems, roots and tubers. When plants were grown on 1.5% sucrose, *StHK4* expression

predominated in stems while in roots the expression levels of *StHK3* and *StHK4* were relatively high
and nearly equal. In contrast to other species (Romanov, 2009; Lomin *et al*., 2012), no prevalent
expression of HK4 orthologs in roots was found. Exogenous cytokinin had little effect on the expression
of cytokinin receptors in potato plants except *StHK4* which can be rapidly upregulated in leaves.
Analysis of promoter structures showed a correlation between the occurrence of *cis*-regulatory elements
and the cytokinin sensitivity of gene expression.

Thus, the totality of our results left no doubt that studied StHK proteins are genuine cytokinin
receptors in potato. The observed unique structural features refine and broaden our notion on the
properties of cytokinin receptors. The revealed peculiarities of cytokinin perception apparatus in potato
might be associated with the ability of this crop to produce tubers. The obtained results create a solid
basis for further in-depth study of the role of the cytokinin signaling system in potato ontogenesis and
provide new biotechnological tools to optimize hormonal regulation of tuber formation.

Acknowledgements

504 This work was supported by the Russian Science Foundation, grants no. 14-14-01095 (before 31.12
505 2016, bioinformatic and some experimental data) and 17-74-20181 (in 2017, conclusive experimental
506 results). for providing opportunity to perform some experiments in his laboratory.

References

Abelenda JA, Prat S. 2013. Cytokinins: determinants of sink storage ability. Current Biology **23,** R561–R563.

Aksenova NP, Konstantinova TN, Golyanovskaya SA, Kossmann J, Willmitzer L, Romanov GA. 2000. Transformed potato plants as a model for studying the hormonal and carbohydrate regulation of tuberization. Russian Journal of Plant Physiology **47,** 370–379.

Aksenova NP, Konstantinova TN, Golyanovskaya SA, Sergeeva LI, Romanov GA. 2012. Hormonal regulation of tuber formation in potato plants. Russian Journal of Plant Physiology **59,** 451–466.

Aksenova NP, Sergeeva LI, Kolachevskaya OO, Romanov GA. 2014. Hormonal regulation of tuber formation in potato. In: Ramawat KG, Merillon JM, eds. *Bulbous Plants. Biotechnology*. New York, Oxon UK: CRC Press, 3–36.

Bauer J, Reiss K, Veerabagu M, Heunemann M, Harter K, Stehle T. 2013. Structure-function analysis of Arabidopsis thaliana histidine kinase AHK5 bound to its cognate phosphotransfer protein AHP1. Molecular Plant **6,** 959–970.

Benkert P, Kunzli M, Schwede T. 2009. QMEAN server for protein model quality estimation. Nucleic Acids Research **37,** W510–W514.

Biasini M, Bienert S, Waterhouse A, *et al.* 2014. SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. Nucleic Acids Research **42,** W252–W258.

Boivin S, Kazmierczak T, Brault M, Wen J, Gamas P, Mysore KS, Frugier F. 2016. Different cytokinin histidine kinase receptors regulate nodule initiation as well as later nodule developmental stages in Medicago truncatula. Plant Cell and Environment **39,** 2198–2209.

Brenner WG, Romanov GA, Köllmer I, Bürkle L, Schmülling T. 2005. Immediate-early and delayed cytokinin response genes of Arabidopsis thaliana identified by genome-wide expression profiling reveal novel cytokinin-sensitive processes and suggest cytokinin action through transcriptional cascades. The Plant Journal **44,** 314–333.

Brütting C, Schäfer M, Vanková R, Gase K, Baldwin IT, Meldau S. 2017. Changes in cytokinins are sufficient to alter developmental patterns of defense metabolites in *Nicotiana attenuata*. The Plant Journal **89,** 15–30.

Caesar K, Thamm AM, Witthöft J, Elgass K, Huppenberger P, Grefen C, Horak J, Harter K. 2011. Evidence for the localization of the *Arabidopsis* cytokinin receptors AHK3 and AHK4 in the endoplasmic reticulum. Journal of Experimental Botany **62**, 5571–5580.

Choi J, Lee J, Kim K, Cho M, Ryu H, An G, Hwang I. 2012. Functional identification of OsHk6 as a homotypic cytokinin receptor in rice with preferential affinity for iP. Plant and Cell Physiology **53,** 1334–1343.

Clamp M, Cuff J, Searle S.M, Barton GJ. 2004. The Jalview Java alignment editor. Bioinformatics **20,** 426–427.

Cong Q, Grishin NV. 2012. MESSA: MEta server for sequence analysis. BMC Biology **10,** 82.

Daudu D, Allion E, Liesecke F, *et al*. 2017. CHASE-containing histidine kinase receptors in apple tree: from a common receptor structure to divergent cytokinin binding properties and specific functions. Frontiers in Plant Science **8,** 1614.

Ding W, Tong H, Zheng W, Ye J, Pan Z, Zhang B, Zhu S. 2017. Isolation, characterization and transcriptome analysis of a cytokinin receptor mutant *Osckt1* in rice. Frontiers in Plant Science **8,** 88.

Eviatar-Ribak T, Shalit-Kaneh A, Chappell-Maor L, Amsellem Z, Eshed Y, Lifschitz E. 2013. A cytokinin-activating enzyme promotes tuber formation in tomato. Current Biology **23,** 1057–1064.

Frugier F, Kosuta S, Murray JD, Crespi M, Szczyglowski K. 2008. Cytokinin: secret agent of symbiosis. Trends in Plant Sciences **13,** 115–120.

Guivarc'h A, Rembur J, Goetz M, Roitsch T, Noin M, Schmülling T, Chriqui D. (2002). Local expression of the ipt gene in transgenic tobacco (*Nicotiana tabacum* L. cv. SR1) axillary buds establishes a role for cytokinins in tuberization and sink formation. Journal of Experimental Botany **53,** 621–629.

Held M, Hou H, Miri M, *et al*. 2014. Lotus japonicus cytokinin receptors work partially redundantly to mediate nodule formation. The Plant Cell **26,** 678–694.

Heyl A, Schmülling T. 2003. Cytokinin signal perception and transduction. Current Opinion in Plant Biology **6**, 480–488.

Heyl A, Riefler M, Romanov GA, Schmülling T. 2012. Properties, functions and evolution of cytokinin receptors. European Journal of Cell Biology **91**, 246–256.

Hirose N, Takei K, Kuroha T, Kamada-Nobusada T, Hayashi H, Sakakibara H. 2008. Regulation of cytokinin biosynthesis, compartmentalization and translocation. Journal of Experimental Botany **59**, 75–83.

Hothorn M, Dabi T, Chory J. 2011. Structural basis for cytokinin recognition by Arabidopsis thaliana histidine kinase 4. Nature Chemical Biology **7,** 766–768.

Hutchison CE, Kieber JJ. 2002. Cytokinin signaling in Arabidopsis. The Plant Cell **14 Suppl,** S47–S59.

Hwang I, Chen HC, Sheen J. 2002. Two-component signal transduction pathways in Arabidopsis. Plant Physiology **129,** 500–515.

Kakimoto T. 2003. Perception and signal transduction of cytokinins. Annual Review of Plant Biology **54**, 605–627.

Karimi M, Depicker A, Hilson P. 2007. Recombinational cloning with plant Gateway vectors. Plant Physiology **145,** 1144–1154.

Kieber JJ, Schaller GE. 2014. Cytokinins. *The Arabidopsis Book* **12**. American Society of Plant Biologists, e0168.

Kolachevskaya OO, Alekseeva VV, Sergeeva LI, Rukavtsova EB, Getman IA, Vreugdenhil D, Buryanov YI, Romanov GA. 2015. Expression of auxin synthesis gene *tms1* under control of the tuber-specific promoter enhances potato tuberization *in vitro*. Journal of Integrative Plant Biology **57,** 734–744.

Kolachevskaya OO, Sergeeva LI, Floková K, Getman IA, Lomin SN, Alekseeva VV, Rukavtsova EB, Buryanov YI, Romanov GA. 2017. Auxin synthesis gene *tms1* driven by tuber-specific promoter alters hormonal status of transgenic potato plants and their responses to exogenous phytohormones. Plant Cell Reports **36,** 419–435.

Kuderová A, Gallová L, Kuricová K, Nejedlá E, Čurdová A, Micenková L, Plíhal O, Šmajs D, Spíchal L, Hejátko J. 2015. Identification of AHK2- and AHK3-like cytokinin receptors in *Brassica napus* reveals two subfamilies of AHK2 orthologues. Journal of Experimental Botany **66**, 339–353.

Laffont C, Rey T, André O, Novero M, Kazmierczak T, Debellé F, Bonfante P, Jacquet C, Frugier F. 2015. The CRE1 cytokinin pathway is differentially recruited depending on Medicago truncatula root environments and negatively regulates resistance to a pathogen. PLoS One **10,** e0116819.

Larkin MA, Blackshields G, Brown NP, *et al*. 2007. Clustal W and Clustal X version 2.0. Bioinformatics **23,** 2947–2948.

Laskowski RA, MacArthur MW, Moss DS, Thornton JM. 1993. PROCHECK - a program to check the stereochemical quality of protein structures. Journal of Applied Crystallography, **26,** 283–291.

Lomin SN, Krivosheev DM, Steklov MYu, Arkhipov DV, Schmülling T, Romanov GA. 2015. Plant membrane assays with cytokinin receptors underpin the unique role of free cytokinin bases as biologically active ligands. Journal of Experimental Botany **66**, 1851–1863.

Lomin SN, Krivosheev DM, Steklov MY, Osolodkin DI, Romanov GA. 2012. Receptor properties and features of cytokinin signaling. Acta Naturae **4,** 31–45.

Lomin SN, Myakushina YuA, Arkhipov DV, Leonova OG, Popenko VI, Schmülling T, Romanov GA. 2018. Studies of cytokinin receptor-phosphotransmitter interaction provide evidences for the initiation of cytokinin signalling in the endoplasmic reticulum. Functional Plant Biology **45,** 192–202.

Lomin SN, Romanov GA. 2008. The analysis of hormone-receptor interaction. Theoretical and practical aspects. Russian Journal of Plant Physiology **55,** 259–273.

Lomin SN, Yonekura-Sakakibara K, Romanov GA, Sakakibara H. 2011. Ligand-binding properties and subcellular localization of maize cytokinin receptors. Journal of Experimental Botany **62**, 5149–5159.

Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser K.E, Simmerling C. 2015. ff14SB: improving the accuracy of protein side chain and backbone parameters from ff99SB. Journal of Chemical Theory and Computation **11,** 3696–3713.

Mayerhofer H, Panneerselvam S, Kaljunen H, Tuukkanen A, Mertens HD, Mueller-Dieckmann J. 2015. Structural model of the cytosolic domain of the plant ethylene receptor 1 (ETR1). Journal of Biological Chemistry **290,** 2644–2658.

Miri M, Janakirama P, Held M, Ross L, Szczyglowski K. 2016. Into the root: how cytokinin controls rhizobial infection. Trends in Plant Sciences **21,** 178–86.

Müller B, Sheen J. 2007. Advances in cytokinin signaling. Science **318**, 68–69.

Nicot N, Hausman J-F, Hoffman L, Evers D. 2005. Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress. Journal of Experimental Botany **56**, 2907–2914.

Pekárová B, Klumpler T, Třísková O, *et al.* 2011. Structure and binding specificity of the receiver domain of sensor histidine kinase CKI1 from *Arabidopsis thaliana*. The Plant Journal **67,** 827–839.

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera – a visualization system for exploratory research and analysis. Journal of Computational Chemistry **25,** 1605–1612.

Pils B, Heyl A. 2009. Unraveling the evolution of cytokinin signaling. Plant Physiology **151,** 782–791.

Potato Genome Sequencing Consortium. 2011. Genome sequence and analysis of the tuber crop potato. Nature **475,** 189–195.

Romanov GA. 2009. How do cytokinins affect the cell? Russian Journal of Plant Physiology **56**, 268–290.

Romanov GA, Aksenova NP, Konstantinova TN, Golyanovskaya SA, Kossmann J, Willmitzer L. 2000. Effect of indole-3-acetic acid and kinetin on tuberization parameters of different cultivars and transgenic lines of potato *in vitro*. Plant Growth Regulation **32,** 245–251.

Romanov GA, Lomin SN, Schmülling T. 2006. Biochemical characteristics and ligand-binding properties of *Arabidopsis* cytokinin receptor AHK3 compared to CRE1/AHK4 as revealed by a direct binding assay. Journal of Experimental Botany **57**, 4051–4058.

Romanov GA, Lomin SN, Schmülling T. 2018. Cytokinin signaling: from the ER or from the PM? That is the question! New Phytologist, in press.

Romanov GA, Spíchal L, Lomin SN, Strnad M, Schmülling T. 2005. A live cell hormone-binding assay on transgenic bacteria expressing a eukaryotic receptor protein. Analytical Biochemistry **347**, 129–134.

Roumeliotis E, Kloosterman B, Oortwijn M, Kohlen W, Bouwmeester HJ, Visser RGF, Bachem CWB. 2012. The effects of auxin and strigolactones on tuber initiation and stolon architecture in potato. Journal of Experimental Botany **63,** 4539–4548.

Sakakibara H. 2006. Cytokinins: activity, biosynthesis, and translocation. Annual Review of Plant Biology **57**, 431–449.

Sali A, Blundell TL. 1993. Comparative protein modelling by satisfaction of spatial restraints. Journal of Molecular Biology **234,** 779–815.

Schäfer M, Meza-Canales ID, Brütting C, Baldwin IT, Meldau S. 2015. Cytokinin concentrations and CHASE-DOMAIN CONTAINING HIS KINASE 2 (NaCHK2)- and NaCHK3-mediated perception modulate herbivory-induced defense signaling and defenses in *Nicotiana attenuata.* New Phytologist **207,** 645–658.

Shen M, Sali A. 2006. Statistical potential for assessment and prediction of protein structures. Protein Science **15,** 2507–2524.

Sparkes IA, Runions J, Kearns A, Hawes C. 2006. Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. Nature Protocols **1,** 2019–2025.

Steklov MY, Lomin SN, Osolodkin DI, Romanov GA. 2013. Structural basis for cytokinin receptor signaling: an evolutionary approach. Plant Cell Reports **32**, 781–793.

Stolz A, Riefler M, Lomin SN, Achazi K, Romanov GA, Schmülling T. 2011. The specificity of cytokinin signalling in *Arabidopsis thaliana* is mediated by differing ligand affinities and expression profiles of the receptors. The Plant Journal **67**, 157–168.

Suzuki T, Miwa K, Ishikawa K, Yamada H, Aiba H, Mizuno T. 2001. The *Arabidopsis* sensor His-kinase, AHK4, can respond to cytokinins. Plant and Cell Physiology **42,** 107–113.

Takeda S, Fujisawa Y, Matsubara M, Mizuno T. 2001. A novel feature of the multistep phosphorelay in *Escherichia coli*: a revised model of the RcsC→YojN→RcsB signaling pathway implicated in capsular synthesis and swarming behaviour. Molecular Microbiology **40,** 440–450.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. Molecular Biology and Evolution **30**, 2725–2729.

Thu NBA, Hoang XLT, Truc MT, Sulieman S, Thao NP, Tran LP. 2017. Cytokinin signaling in plant response to abiotic stresses. In: Pandey G, Ed. *Mechanism of Plant Hormone Signaling under Stress*. First Edition, vol. 1, chpt. 4. John Wiley & Son, Inc., 71–100.

Voinnet O, Rivas S, Mestre P, Baulcombe D. 2003. An enhanced transient expression system in plants based on suppression of gene silencing by the p19 protein of tomato bushy stunt virus. The Plant Journal **33**, 949–956.

Wang B, Chen Y, Guo B, Kabir MR, Yao Y, Peng H, Xie C, Zhang Y, Sun Q, Ni Z. 2014. Expression and functional analysis of genes encoding cytokinin receptor-like histidine kinase in maize (*Zea mays* L.). Molecular Genetics and Genomics **289,** 501–512.

Wiederstein M, Sippl MJ. 2007. ProSA-web: interactive web service for the recognition of errors in three-dimensional structures of proteins. Nucleic Acids Research **35** (suppl_2), W407–W410.

Wulfetange K, Lomin SN, Romanov GA, Stolz A, Heyl A, Schmülling T. 2011. The cytokinin receptors of Arabidopsis are located mainly to the endoplasmic reticulum. Plant Physiology **156,** 1808–1818.

Yonekura-Sakakibara K, Kojima M, Yamaya T, Sakakibara H. 2004. Molecular characterization of cytokinin-responsive histidine kinases in maize. Differential ligand preferences and response to cis-zeatin. Plant Physiology **134,** 1654–1661.

Zwack PJ, Rashotte AM. 2015. Interactions between cytokinin signalling and abiotic stress responses. Journal of Experimental Botany **66,** 4863–4871.

Table 1. *Proteins and genes predictably related to cytokinin signaling system of potato*

Protein type	Protein name	Gene ID	mRNA	Protein	Protein length, aa
CHK	StHK2	LOC102591086	XM_015303261.1	XP_015158747.1	1263
CHK	StHK3	LOC102587294	XM_006352114.2	XP_006352176.1	1032
CHK	StHK4	LOC102603756	XM_006354988.2	XP_006355050.1	992
HPt	StHP1a	LOC102590747	XM_006365209.2	XP_006365271.1	151
			XM_006365208.2	XP_006365270.1	151
			XM_006365207.2	XP_006365269.1	151
HPt	StHP1b	LOC102603297	XM_006352731.2	XP_006352793.1	152
HPt	StHP1c	PGSC0003DMG40 0028593	PGSC0003DMT4000 73603	PGSC0003DMT4000 73603	148
HPt	StHP6	LOC102601463	XM_006364157.2	XP_006364219.1	156
HPt	StHP4a	LOC102589200	XM_015304066.1	XP_015159552.1	112
			XM_006364659.2	XP_006364721.1	136
HPt	StHP4b	LOC102584884	XM_015315420.1	XP_015170906.1	137
RR-B	StRR1a	LOC102578736	XM_006363517.2	XP_006363579.1	675
			XM_006363518.2	XP_006363580.1	675
RR-B	StRR1b	LOC102586468	XM_006345914.1	XP_006345976.1	663
RR-B	StRR1c	LOC102596771	XM_006349891.2	XP_006349953.1	556
RR-B	StRR14	LOC102606335	XM_006354997.1	XP_006355059.1	653
			XM_006354996.1	XP_006355058.1	656
RR-B	StRR11	LOC102593308	XM_006341706.2	XP_006341768.1	581
			XM_006341705.2	XP_006341767.1	581
			XM_015306278.1	XP_015161764.1	481
RR-B	StRR18a	LOC102598455	XM_006343619.2	XP_006343681.1	681
RR-B	StRR18b	LOC102587717	XM_006350015.2	XP_006350077.1	707
ARR19	StRR19	LOC107060895	XM_015309426.1	XP_015164912.1	371
RR-A	StRR4	LOC102602758	XM_015313344.1	XP_015168830.1	248
RR-A	StRR9a	LOC102590336	XM_006355533.2	XP_006355595.1	163
RR-A	StRR9b	LOC102588738	XM_015314746.1	XP_015170232.1	214
			XM_015314747.1	XP_015170233.1	211
RR-A	StRR9c	LOC102599826	XM_006351210.2	XP_006351272.1	226
RR-A	StRR9d	LOC102601166	XM_006351214.2	XP_006351276.1	226
RR-A	StRR8	LOC102588738	XM_015314747.1	XP_015170233.1	211
			XM_015314746.1	XP_015170232.1	214
RR-A	StRR15	LOC102605280	XM_006344933.2	XP_006344995.1	202
RR-A	StRR17	LOC102583233	XM_006357236.2	XP_006357298.1	156
RR-C	StRR22a	LOC107058083	XM_015303399.1	XP_015158885.1	186
RR-C	StRR22b	LOC107058085	XM_015303400.1	XP_015158886.1	184
RR-C	StRR22c	LOC107059982	XM_015307157.1	XP_015162643.1	137
RR-C	StRR22d	LOC102580685	XM_006361561.2	XP_006361623.2	115

Nomenclature of the NCBI database is represented except StHP1c found only in the Phytozome database. Number of RNA entries exceeds that of proteins due to alternative splicing. Data corresponding to cytokinin receptor proteins/genes and response regulator type A proteins/genes studied in this work are highlighted.

Table 2. *Putative cytokinin receptor genes in potato genomes and encoded proteins*

Receptor clade	Putative cytokinin receptors of potato plants:		SNP number in putative cytokinin receptor genes/proteins of cv. Désirée vs var. Phureja:		Number of Désirée cDNA clones
	Phureja* (length, aa)	Désirée ** (length, aa)	DNA bases	Amino acids	
HK2 orthologs	StHK2 (1263 aa)	StHK2a (1263 aa)	5 SNPs	3 SNPs	17
		StHK2b (1263 aa)	4 SNPs	2 SNPs	17
HK3 orthologs	StHK3 (1032 aa)	StHK3a (1032 aa)	No SNP	No SNP	6
		StHK3b (1031 aa)	20 SNPs, 3 del.	9 SNPs, 1 del.	3
CRE1/AHK4 orthologs	StHK4 (992 aa)	StHK4a (992 aa)	No SNP	No SNP	1
		StHK4b (991 aa)	28 SNPs, 3 del.	13 SNPs, 1 del.	9

*Doubled monoploid, method: total genome sequencing.

** Autotetraploid, method: PCR with cDNA as a template.

Table 3. *The affinity (K_d) of various cytokinins for putative potato receptors*

Cytokinin	Abbreviation	Apparent K_d (nM) for putative receptors:			
		StHK2a	StHK3a$_{SM}$	StHK4a	StHK4b
trans-Zeatin	**tZ**	2.6±0.3	4.7±0.6	2.5±0.7	3.0±0.3
cis-Zeatin	**cZ**	102±7	110±39	106±22	129±19
N^6-Isopentenyladenine	**iP**	2.4±0.2	5.2±0.8	2.1±0.2	2.5±0.3
Dihydrozeatin	**DZ**	169±18	21±3	178±37	227±33
N^6-Benzyladenine	**BA**	45±3.5	49±7	55±7	63±12
Thidiazuron	**TZ**	1.40±0.04	2.3±0.5	12.6±1.9	17.2±2.5

Figure legends

Fig. 1. Phylogenetic tree of cytokinin receptors. Species are: StHK2-4, *Solanum tuberosum*; SlHK2-4, *Solanum lycopersicum*; AHK2-4, *Arabidopsis thaliana*; OsHK3,4,6, *Oryza sativa*; AmbtriHK2,3, *Amborella trichopoda*; PITA 000007449 and PITA 000016046, *Pinus taeda*; MA 101803g0010 and MA 47453g0010, *Picea abies*; CRE1,2-1,2-2, *Selaginella moellendorffii*; CHK1-3, *Physcomitrella patens*. SP – seed plants, Lyco – Lycophyta, Bryo – Bryophyta. Parameters of ClustalW algorithm were: phylogeny test – bootstrap method, no. of bootstrap replications – 100, substitutions type – amino acid, model – equal input model, rates among sites – gamma distributed, no of discrete gamma categories – 3, gaps/missing data treatment – complete deletion, ML heuristic method – subtree-pruning – regrafting.

Fig. 2. Module/domain structures of the predicted potato cytokinin receptors. Protein domains: TM, transmembrane segment; DI, dimerization interphase; CHASE, Cyclase/Histidine kinases Associated SEnsory domain; HisKA, histidine kinase domain; HATPase, adenosine triphosphatase domain; Rec-like, receiver-like domain; Rec, receiver domain. Conserved amino acids and consensus motifs (N, G1, F, G2) are indicated. Scales at the bottom of the structures indicate the length in aa number.

Fig. 3. Cytokinin receptor sequence alignment. Consensus motifs and conserved amino acids are marked. AHK and OsHK correspond to Arabidopsis and rice proteins, respectively. Numbers of not shown amino acids are indicated in brackets.

Fig. 4. Homology models for predicted potato cytokinin receptor domains. Sensor modules and HisKA domain are presented in form of dimers where one of subunits is colored grey.

Fig. 5. pH dependencies of *trans*-zeatin binding to putative potato cytokinin receptors.

Fig. 6. Cytokinin receptors of potato feed MSP signaling pathway in ΔRcsC *E. coli* cells.

Fig. 7. Organ-dependent (A) and clade-dependent (B) patterns of expression of cytokinin receptors in potato plants cultivated on media with different % sucrose. Relative transcript copy number is given as % of the total transcript amount in each plot, taken as 100%.

Fig. 8. Degrees of transcription induction (BA/control) of cytokinin receptor (top) and response regulator type A (bottom) genes after 1 h treatment of potato plants with 1 μM BA. Plants were grown on MS medium with 1.5% sucrose for 5-6 weeks under standard LD conditions. L, S, R signify leaves, stems and roots, respectively. More than two-fold prevalence of transcripts in BA-treated over control plants is considered as significant induction, bars corresponding to induced genes are outlined red.

Fig. 9. Cytokinin-responsive *cis*-regulatory elements in promoters of cytokinin receptor genes (upper part) and response regulators type A genes (lower part) of potato. Elements are shown on both DNA strands. Promoter area proximal to transcription start is boxed.

Fig. 1. Phylogenetic tree of cytokinin receptors. Species are: StHK2-4, *Solanum tuberosum*; SlHK2-4, *Solanum lycopersicum*; AHK2-4, *Arabidopsis thaliana*; OsHK3,4,6, *Oryza sativa*; AmbtriHK2,3, *Amborella trichopoda*; PITA 000007449 and PITA 000016046, *Pinus taeda*; MA 101803g0010 and MA 47453g0010, *Picea abies*; CRE1,2-1,2-2, *Selaginella moellendorffii*; CHK1-3, *Physcomitrella patens*. SP – seed plants, Lyco – Lycophyta, Bryo – Bryophyta. Parameters of ClustalW algorithm were: phylogeny test – bootstrap method, no. of bootstrap replications – 100, substitutions type – amino acid, model – equal input model, rates among sites – gamma distributed, no of discrete gamma categories – 3, gaps/missing data treatment – complete deletion, ML heuristic method – subtree-pruning – regrafting.

Fig. 2. Module/domain structures of the predicted potato cytokinin receptors. Protein domains: TM, transmembrane segment; DI, dimerization interphase; CHASE, Cyclase/Histidine kinases Associated SEnsory domain; HisKA, histidine kinase domain; HATPase, adenosine triphosphatase domain; Rec-like, receiver-like domain; Rec, receiver domain. Conserved amino acids and consensus motifs (N, G1, F, G2) are indicated. Scales at the bottom of the structures indicate the length in aa number.

Fig. 3. Cytokinin receptor sequence alignment. Consensus motifs and conserved amino acids are marked. AHK and OsHK correspond to Arabidopsis and rice proteins, respectively. Numbers of not shown amino acids are indicated in brackets.

Fig. 4. Homology models for predicted potato cytokinin receptor domains. Sensor modules and HisKA domain are presented in form of dimers where one of subunits is colored grey.

Fig. 5. pH dependencies of *trans*-zeatin binding to putative potato cytokinin receptors.

Fig. 6. Cytokinin receptors of potato feed MSP signaling pathway in ΔRcsC *E. coli* cells.

Fig. 7. Organ-dependent (A) and clade-dependent (B) patterns of expression of cytokinin receptors in potato plants cultivated on media with different % sucrose. Relative transcript copy number is given as % of the total transcript amount in each plot, taken as 100%.

Fig. 8. Degrees of transcription induction (BA/control) of cytokinin receptor (top) and response regulator type A (bottom) genes after 1 h treatment of potato plants with 1 μM BA. Plants were grown on MS medium with 1.5% sucrose for 5-6 weeks under standard LD conditions. L, S, R signify leaves, stems and roots, respectively. More than two-fold prevalence of transcripts in BA-treated over control plants is considered as significant induction, bars corresponding to induced genes are outlined red.

Fig. 9. Cytokinin-responsive *cis*-regulatory elements in promoters of cytokinin receptor genes (upper part) and response regulators type A genes (lower part) of potato. Elements are shown on both DNA strands. Promoter area proximal to transcription start is boxed.