Structural diversity across arbuscular mycorrhizal, ectomycorrhizal, and endophytic plant–fungus networks

Hirokazu Toju[1,2*], Hirotoshi Sato[3], Satoshi Yamamoto[4], Akifumi S. Tanabe[5]

[*]Correspondence: toju.hirokazu.4c@kyoto-u.ac.jp

[1]Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2113, Japan

[2]Precursory Research for Embryonic Science and Technology (PRESTO), Japan Science and Technology Agency, Kawaguchi, Saitama 332-0012, Japan

[3]Graduate School of Human and Environmental Studies, Kyoto University, Sakyo, Kyoto 606-8501, Japan

[4]Graduate School of Science, Kyoto University, Kitashirakawa-oiwake-cho, Kyoto 606-8502, Japan

[5]Faculty of Science and Technology, Ryukoku University, 1-5 Yokotani, Seta Oe-cho, Otsu, Shiga 520-2194, Japan

This article includes 7 Additional files (Data S1-7).

Abstract

Background: Below-ground linkage between plant and fungal communities is one of the major drivers of terrestrial ecosystem dynamics. However, we still have limited knowledge of how such plant–fungus associations vary in their community-scale properties depending on fungal functional groups and geographic locations.

Methods: Based on high-throughput sequencing of root-associated fungi in eight forests along the Japanese Archipelago, we performed a comparative analysis of arbuscular mycorrhizal, ectomycorrhizal, and saprotrophic/endophytic associations across a latitudinal gradient from cool-temperate to subtropical regions.

Results: In most of the plant–fungus networks analyzed, host–symbiont associations were significantly specialized but lacked "nested" architecture, which has been commonly reported in plant–pollinator and plant–seed disperser networks. Meanwhile, the structure of arbuscular mycorrhizal networks was differentiated from that of ectomycorrhizal and saprotrophic/endophytic networks, characterized by high connectance. Our data also suggested that geographic factors affected the organization of plant–fungus network structure. For example, the southernmost subtropical site analyzed in this study displayed lower network-level specificity of host–symbiont associations and higher (but still low) nestedness than northern localities.

Conclusions: Our comparative analyses suggest that arbuscular mycorrhizal, ectomycorrhizal, and saprotrophic/endophytic plant–fungus associations often lack nested network architecture, while those associations can vary, to some extent, in their community-scale properties along a latitudinal gradient. Overall, this study provides a basis for future studies that will examine how different types of plant–fungus associations collectively structure terrestrial ecosystems.

Keywords: biodiversity; community ecology; competitive exclusion; host specificity or preference; latitudinal gradients; microbiomes; plant–fungus interactions; plant–soil feedback; species coexistence; mycorrhizal and endophytic symbiosis

Background

Fungi in the below-ground biosphere are key drivers of terrestrial ecosystem processes [1-4].

Mycorrhizal fungi are considered to support land plants not only by provisioning soil nitrogen

and phosphorous [5, 6] but also by increasing plants' resistance to biotic/abiotic stress [7, 8].

Pathogenic fungi in the soil affect the survival/mortality of young plants in a major way,

possibly determining spatial distributions of plant species within forest/grassland ecosystems

[9, 10]. Moreover, recent mycological studies have begun to examine the poorly explored

diversity of endophytic fungi, which can enhance the nutritional conditions and pathogen

resistance of mycorrhizal and non-mycorrhizal plant species [11-16]. Thus, terrestrial biomes

consist of multiple layers of below-ground plant–fungus interactions [17]. Nonetheless, we still

have limited knowledge of the structure of such complex webs of interactions, leaving major

processes in below-ground ecosystems poorly explored.

In enhancing our understanding of community- or ecosystem-level processes of

below-ground plant–fungus interactions, analyses on community-scale properties of such

host–symbiont associations provide essential insights. For example, if a pathogenic fungal

community consists mainly of species with narrow host ranges, it as a whole is expected to

restrict the emergence of dominant plant species through "negative plant–soil feedback",

contributing to the maintenance of plant species diversity within an ecosystem [18-20]. In

contrast, with a high proportion of mycorrhizal fungi with narrow host ranges, their specific

host species, such as Pinaceae plants hosting Suillaceae ectomycorrhizal fungi [21], will

dominate the plant community through positive plant–soil feedback [20, 22]. Meanwhile,

endophytic and arbuscular mycorrhizal fungi with broad host ranges [23-25] may diminish

such negative and positive feedback by interlinking otherwise compartmentalized ecological

dynamics (but see [26]). Therefore, concomitant analyses of community-scale properties of

those multiple plant–fungus associations are of particular importance in understanding how

plant–soil feedbacks organize terrestrial ecological processes.

Since the application of network science to ecology and mycology, researchers have

evaluated the architecture of networks that represent linkage between plant and fungal

communities [27]. Those studies have shown that arbuscular mycorrhizal [28-30],

ectomycorrhizal [31], and ericaceous [32] plant–fungus networks exhibit moderate or low

levels of host–symbiont specificity, while they are structured to avoid overlap of host plant

ranges within fungal communities. In addition, many of those plant–fungus networks [17, 31,

33] are known to lack "nested" architecture (i.e., structure of networks wherein specialist

species interact with subsets of partners of generalist species [34]), which has been commonly

reported in above-ground networks of plant–pollinator and plant–seed-disperser interactions

[34-36] (but see [37]). However, in those previous studies, data of different types of plant-

–fungus networks have been collected from different geographic localities with different

sampling strategy, precluding the chance of simultaneously evaluating the effects of interaction

type and geographic factors. Although comparative studies of published data provide

invaluable insights [27], compiled data often vary in the molecular markers used and they may

differ in appropriate null model assumptions in statistically examining network topological

properties.

In this study, we compared community-scale properties of arbuscular-mycorrhizal,

ectomycorrhizal, and endophytic associations across eight forest sites spanning from

cool-temperate to subtropical regions in Japan. Based on high-throughput sequencing of

root-associated fungi, we obtained network data depicting how multiple plant species are

associated with respective functional groups of fungi in each of the eight forests. We then

examined how structure varied depending on categories of plant–fungus associations and

geographic locations. Overall, this study provides a first step for integrating insights into

community-scale properties of multiple types of below-ground plant–fungus associations and

their ecosystem-level consequences.

Methods

Terminology

In analyzing metadata of community-scale properties of plant–fungus associations, we need to

use consistent terminology that can be applied to a wide range of host–symbiont associations.

While plant–fungus network properties have been compared within a single functional group of

fungi (e.g., arbuscular mycorrhizal or ectomycorrhizal fungi) in most studies, we herein target

not only arbuscular mycorrhizal and ectomycorrhizal fungi but also pathogenic and

saprotrophic/endophytic fungi. Given that those functional groups of fungi vary considerably in

their microscopic structure within plant tissue [8], developing a general criterion for

mutualistic/antagonistic interactions with host plants is impossible. Thus, we targeted all the

fungi detected by high-throughput sequencing and the data described below could contain not

only mutualistic/antagonistic fungi but also commensalistic fungi merely adhering to plant

roots [38]. In this sense, our data represented symbiotic relationships in the broad sense, i.e.,

intimate physical connections between organisms [17, 39].

116 Sampling

We collected root samples at eight forest sites (four cool-temperate, one warm-temperate, and

three subtropical forests) across the entire range of the Japanese Archipelago (45.042–24.407

°N; Fig. 1A; Additional file 1: Data S1). In each forest, 2-cm segment of terminal roots were

collected from 3-cm below the soil surface at 1-m horizontal intervals. In each forest, 383

terminal root samples were collected. Those roots were collected indiscriminately regarding

root morphology or apparent mycorrhizal type so that the samples as a whole represented the

relative frequency of plant–fungal associations in the horizon in each forest [40]. Therefore,

while the sample sets consisted mainly of woody plants, they also included herbaceous plants

(Additional file 2: Data S2). Each root sample was preserved in 70% ethanol and stored at -25

°C until DNA extraction.

128 Molecular analysis

Each root sample was placed in 70% ethanol with 1-mm zirconium balls and it was then shaken

at 15 Hz for 2 min with a TissueLyser II (Qiagen) [23]. The washed roots were subsequently

pulverized by shaking with 4-mm zirconium balls at 25 Hz for 3 min. DNA extraction was then

performed with a cetyltrimethylammonium bromide method [41].

The internal transcribed spacer 1 (ITS1) region of root-associated fungi was amplified with

the primers ITS1F_KYO1 and ITS2_KYO2, which target not only Ascomycota and

Basidiomycota fungi but also diverse non-Dikarya (e.g., Glomeromycota) taxa [42]. We used

the forward primer ITS1F-KYO1 fused with 3–6-mer Ns for improved Illumina sequencing

quality [43] and the forward Illumina sequencing primer (5'- TCG TCG GCA GCG TCA GAT

GTG TAT AAG AGA CAG- [3–6-mer Ns] – [ITS1-KYO2] -3') and the reverse primer

ITS2-KYO2 fused with 3–6-mer Ns and the reverse sequencing primer (5'- GTC TCG TGG

GCT CGG AGA TGT GTA TAA GAG ACA G [3–6-mer Ns] - [ITS2_KYO2] -3'). The DNA

polymerase system of KOD FX Neo (Toyobo) was used with a temperature profile of 94 ℃ for

2 min, followed by 35 cycles at 98 ℃ for 10 s, 50 ℃ for 30 s, 68 ℃ for 50 s, and a final

extension at 68 ℃ for 5 min. The ramp rate was set to 1 ℃/sec to prevent the generation of

chimeric sequences [44]. Illumina sequencing adaptors were then added to each sample in the

subsequent PCR using the forward primers consisting of the P5 Illumina adaptor, 8-mer tags for

sample identification [45], and a partial sequence of the sequencing primer (5'- AAT GAT ACG

GCG ACC ACC GAG ATC TAC AC - [8-mer index] - TCG TCG GCA GCG TC -3') and the

reverse primers consisting of the P7 adaptor, 8-mer tags, and a partial sequence of the

sequencing primer (5'- CAA GCA GAA GAC GGC ATA CGA GAT - [8-mer index] - GTC

TCG TGG GCT CGG -3'). In the reaction, KOD FX Neo was used with a temperature profile

of 94 ℃ for 2 min, followed by 8 cycles at 98 ℃ for 10 s, 55 ℃ for 30 s, 68 ℃ for 50 s, and a

final extension at 68 ℃ for 5 min. The PCR amplicons of 384 samples in each forest (including

one PCR negative control) were pooled with equal volume after a purification/equalization

process with AMPureXP Kit (Beckman Coulter).

For the identification of plants, another set of PCR was performed targeting chloroplast

rbcL region with rbcL_F3 and rbcL_R4 primers [40]. The fusion primer design, DNA

polymerase system, temperature profiles, and purification processes used in the *rbcL* analysis

were the same as those of the fungal ITS analysis. The ITS and *rbcL* libraries were processed in

two Illumina MiSeq runs, in each of which samples of four forest sites were combined (run

center: KYOTO-HE) (2×250 cycles; 15% PhiX spike-in).

Bioinformatics

In total, 17,724,456 and 17,228,848 reads were obtained for the first and second MiSeq runs.

The raw sequencing data were converted into FASTQ files using the program bcl2fastq 1.8.4

provided by Illumina. The FASTQ files were then demultiplexed using the program Claident

v0.2.2016.07.05 [46, 47]. To avoid possible errors resulting from low-quality index sequences,

the sequencing reads whose 8-mer index positions included nucleotides with low (< 30) quality

scores were discarded in this process. As reverse sequences output by Illumina sequencers have

lower quality values than forward sequences, we used only forward sequences after removing

low-quality 3'-ends using Claident (sequencing data deposit: DDBJ DRA accession:

DRA006339). Noisy reads were subsequently discarded and the reads that passed the filtering

process were clustered using VSEARCH [48] as implemented in Claident. The threshold

sequencing similarities in the clustering were set to 97% for fungal ITS and 98% for *rbcL*,

respectively. While sequence similarity values have been set to 97% in most ITS analyses of

Ascomycota and Basidiomycota fungi [49] (see also [50]), a recent study showed that

Glomeromycota fungi generally had much higher intraspecific ITS-sequence variation than

Dikarya fungi [51]. Therefore, we performed an additional clustering analysis with a 94%

cutoff similarity for defining Glomeromycota OTUs. Note that changing cut-off similarities

(81–97%) did not qualitatively change statistical properties of plant–fungus network structure

in a previous study [17]. The taxonomic assignment of the OTUs (Additional files 3-4: Data

S3-4) was conducted based on the combination of the query-centric auto-*k*-nearest neighbor

(QCauto) method [46] and the lowest common ancestor (LCA) algorithm [52] as implemented

in Claident. Note that taxonomic identification results based on the QCauto–LCA approach

were comparable to, or sometimes more accurate than, those with the alternative approach

combining the UCLUST algorithm [53] with the UNITE database [54] [see [32] and [55] for

detailed comparison of the QCautoLCA and UCLUST–UNITE approaches]. The functional

group of each fungal OTU was inferred using the program FUNGuild 1.0 [56]. For 44.1 %

(3560/8080) of fungal OTUs, functional group information was inferred (Additional file 1:

Data S1).

The obtained information of *rbcL* OTUs was used to identify each root sample, although

species-level taxonomic information was unavailable for some plant taxa in each forest due to

the relatively low variability of the chloroplast region [57]. Thus, we also used the information

of the ITS sequencing libraries, which included not only fungal but also host plant sequencing

reads: there were plant taxa that could not be identified to species even with ITS information.

Based on the *rbcL* and ITS information of plant sequences, possibly contaminated samples

were removed from the dataset.

For each of the eight forests, we then obtained a sample (row) × fungal OTU (column) data

matrix, in which a cell entry depicted the number of sequencing reads of an OTU in a sample.

The cell entries whose read counts represented less than 0.1% of the total read count of each

sample were subsequently excluded because those rare entries could derive from

contaminations from soil or PCR/sequencing errors [58]. The filtered matrices were then

rarefied to 1,000 reads per sample using the "rrarefy" function of the vegan 2.4-3 package [59]

of R 3.4.1 [60]. As the number of samples with 1,000 or more reads varied among the eight

forests examined (240–288 samples), it was equalized by randomly sampling 240 samples

without duplication in each forest ("sample-level matrices"; Additional file 2: Data S2).

Based on the sample-level matrices, we obtained another type of matrices, in which a cell

indicated the number of samples representing associations between a plant species/taxa

(row) and a fungal OTU (column) ("species-level matrices"; Additional file 5: Data S5).

In addition to the matrix indicating associations between all fungal OTUs and their host plants

(ALL), a series of partial network matrices representing respective fungal functional groups

were obtained by selecting arbuscular mycorrhizal (AM), ectomycorrhizal (ECM), potentially

pathogenic (PATHO), and saprotrophic/endophytic (SAPENDO) fungal OTUs in each forest

(Additional file 6: Data S6). Due to the limited availability of information of fungal ecology,

functional groups of many fungal OTUs could not be estimated and there were only 9–25

fungal OTUs inferred to be plant pathogens in respective forests (Additional files 1 and 5: Data

S1 and S5).

Data analysis

Based on the sample-level matrices, relationship between the number of samples and that of

observed fungal OTUs was analyzed for each forest using the "specaccum" function of the

vegan package. The community-scale plant–fungus associations represented by the

species-level matrices ("ALL" network matrices; Additional file 5: Data S5) were visualized

using the program GePhi 0.9.1 [61] with "ForceAtlas2" layout algorithm [62]. We then

analyzed the statistical properties of the ALL networks and partial networks (Additional file 6:

Data S6) in terms of the H_2' metric of network-level interaction specificity [63], which has been

frequently used to measure the degree of interaction specificity in host–symbiont networks [64,

65]. The plant–fungus associations were evaluated also by the weighted NODF metric [66] of

network nestedness [34], which measures the degree to which specialists (species with narrow

partner ranges) interact with partners of generalists (species with broad partner ranges) in the

same guild or trophic level. We further examined how host plant ranges were differentiated

within the fungal community of each forest based on checkerboard scores [67]: a high/low

score of the checkerboard index indicates host differentiation/overlap within a guild or trophic

level [65]. Although modularity is another important index frequently used in ecological

network studies [35], its computation was too time-consuming to be applied to randomization

analyses (see below) of our present datasets consisting of more than 1,000 fungal OTUs and

their host plants. Note that we previously found that below-ground plant–fungal associations

generally showed statistically significant but low network modularity [17, 32, 65].

As estimates of network indices could vary depending on species compositions of

examined communities, we standardized the indices as

241   relative index value = $[I_{observed} - \text{mean}(I_{randomized})] / \text{SD}(I_{randomized})$

where $I_{observed}$ was the index estimate of the observed data matrix, and mean($I_{randomized}$) and

SD($I_{randomized}$) were the mean and standardized deviation of the index values of randomized

matrices [65]. Randomized matrices were obtained by shuffling host-plant labels in the

sample-level matrices and subsequently converting the randomized sample-level matrices into

randomized species-level matrices. Although we used two additional methods ["r2dtable" [68]

and "vaznull" [69] methods] of matrix randomization in our previous studies of plant–fungus

networks [17, 65], they were too time-consuming to be used in the present large datasets: note

that the three randomization methods compared in those previous studies yielded qualitatively

similar results [17, 65]. The number of randomizations was set to 1,000 for H_2'/nestedness

analyses and 100 for checkerboard-score analyses, which required substantial computing time.

Based on the network indices, we examined how the community-scale properties of the

plant–fungus associations varied among local forests and fungal functional groups. For each of

interaction specificity (relative H_2'), nestedness (relative weighted NODF nestedness), and

checkerboard index (relative checkerboard values), an ANOVA model was constructed by

incorporating locality (forest sites), fungal functional group, number of plant species/taxa,

number of fungal OTUs, and network connectance (the proportion of non-zero entries in

community matrices) as explanatory variables. The variation in the plant–fungus network

properties was visualized based on a principal component analysis based on a correlation

matrix: the variables included were H_2' interaction specificity, NODF nestedness, checkerboard

index, number of plant species/taxa, number of fungal OTUs, proportion of fungal OTUs to

plant species/taxa, and connectance.

264 **Results**

Total fungal OTU richness was higher in warm-temperate and subtropical forests than in

cool-temperate forests (Figs. 1b and 2). The OTU richness of AM fungi was higher in the three

subtropical forests, while that of ECM fungi decreased in the southern forests (Fig. 3a). The

ratio of the total number of fungal OTUs to the number of plant species/taxa varied among

forests, although there was seemingly no systematic variation between cool-temperate and the

other (warm temperate and subtropical) localities (Fig. 3b). Connectance varied among forests

as well, while it was consistently higher in AM than in ALL, ECM, and SAPENDO

networks/partial networks in seven of the eight study forests (Fig. 3c). The connectance of

PATHO partial networks varied considerably among forests presumably due to low OTU

richness and the resultant uncertainty in index estimation.

The relative H_2' metric of interaction specificity significantly varied among forests but not

among fungal functional groups when the effects of plant diversity, fungal OTU richness, and

connectance were controlled in an ANOVA model (Table 1; Fig. 3d). The relative nestedness of

the ALL matrices of plant–fungus associations was lower than zero in most forests but not in

the southern most subtropical forest (Fig. 3e; Additional file 7: Data S7). Overall, plant–fungus

associations in ALL networks were more specialized (Fig. 3d) and less nested (Fig. 3e) than

those of partial networks. In addition, fungal OTUs in ALL networks displayed stronger

differentiation of host ranges than those in partial networks (Fig. 3f).

After taking into account plant and fungal diversity in an ANOVA model, neither locality

nor fungal functional group explained the variation in relative nestedness (Table 1). The

relative checkerboard scores varied among localities (Fig. 3f), although the effects of locality

were non-significant in an ANOVA model (Table 1). The ANOVA model showed that the

variation in relative checkerboard scores was explained, to some extent, by fungal functional

groups.

In the principal component analysis of network indices, ALL, PATHO and other

networks/partial networks were separated by the first principal component, which represented

high plant diversity, fungal OTU richness, relative H_2', and relative checkerboard scores as well

as low relative nestedness (Fig. 4a). By incorporating the third principal component, which

represented high fungal OTU richness and connectance, the cluster of AM partial networks and

that of ECM and SAPENDO partial networks were grouped with some overlap (Fig. 4b).

Discussion

Our data, which included 17–55 plant species/taxa and more than 1000 fungal OTUs in each of

the eight forests, provided a novel opportunity to evaluate how different types of below-ground

plant–fungus associations varied in their community-scale characteristics along a latitudinal

gradient. We then found that network structural properties differed among different types of

plant–fungus associations (Fig. 3), while geographic factors contributed to the variation found

in network structure (Table 1). Specifically, arbuscular mycorrhizal networks differed in their

architecture from ectomycorrhizal and saprotrophic/endophytic networks, characterized by

high connectance (Fig. 4). We also found that networks consisting of all functional groups of

fungi and their host plants had higher network-level interaction specificity, more differentiated

host ranges between fungi, and lower network nestedness than the partial networks of

arbuscular mycorrhizal, ectomycorrhizal, and saprotrophic/endophytic associations (Figs. 3

and 4). As in previous studies, our data included many fungal OTUs unassigned to functional

groups due to the paucity of the information of fungal functions and guilds in databases [56].

However, by extending findings in previous plant–fungus network studies [30, 31, 65], in

which sampling strategies, interaction type, or geographic factors were not controlled

systematically, this study offers a basis for discussing how different types of below-ground

plant–fungus associations collectively build plant–soil feedback in terrestrial ecosystems.

Among the network indices examined in this study, nestedness showed an idiosyncratic

tendency in light of other types of interaction networks examined in community ecology

[34-36]. We found that below-ground plant–fungus networks often displayed "anti-nested"

architecture, in which scores representing nested network structure were lower than those

expected by chance (i.e., negative values of relative nestedness; Fig. 3e), as suggested also in

previous studies [17, 31, 65]. Although factors organizing anti-nested network architecture

remain to be investigated, competition for host plants among fungal species has been inferred to

decrease nestedness of plant–fungus associations [65]. In addition, a previous comparative

study suggested that plant–fungus network nestedness decreased with increasing annual mean

temperature on a global scale [27].

The prevalence of anti-nested or non-nested network structures is in sharp contrast to

observations on other types of plant–partner networks, which commonly show statistically

significant nested architecture [34]. Specifically, plant–pollinator and plant–seed disperser

interactions are generally characterized by nested network architecture in which overlap of

partner ranges within the same guild are expected to mitigate competition between plant species

[34-36]. Therefore, anti-nested structure of plant–fungus networks provides important insights

in terms of community ecological theory linking network structure and species coexistence [36,

37, 70]. Given that below-ground fungi constitute one of the most species-rich components of

the terrestrial biosphere [3], understanding community-scale properties of below-ground

plant–fungus associations is a major step for disentangling relationship among network

structure, species coexistence, and community stability.

To overcome the inconsistency between theory and observations, we may need to take into

account basic biology of below-ground plant–fungus associations. We here highlight two

backgrounds that need more attention for deepening discussion on ecological networks and

species coexistence. First, in contrast to plant–pollinator or plant–seed disperser networks,

which are often assumed to consist only of mutualistic interactions, below-ground plant–fungus

networks can involve not only mutualistic but also antagonistic and commensalistic

interactions. This diversity of interaction type can lead to high stability of below-ground fungal

and their host plant communities. Specifically, while communities consisting exclusively of

mutualistic interactions are inherently unstable [71], involvement of a small fraction of

antagonistic interactions in those communities can dramatically enhance species coexistence

[72]. Second, because fungi can disperse long distances as spores [73, 74] (but see [75]), their

local species richness (alpha diversity) may be greatly impacted by metacommunity processes

[76]. Interestingly, a recent theoretical study on food webs predicted that strong coupling of

local communities within a metacommunity could result in positive relationship between

species richness and community stability [77]. Such theoretical evaluation of metacommunity

dynamics has been extended to systems involving mutualistic interactions [78], providing

platforms for considering how dispersal abilities of constituent species determine local species

richness/coexistence of different types of plant–partner networks.

To apply a standardized criterion for plant–fungus associations, we did not perform any

data screening based on the "mycorrhizal types" of plant species. As a result, our data included

plant–fungus combinations that could not be classified into well-recognized categories of

mycorrhizal symbioses [8]. For example, ectomycorrhizal fungi were detected not only from

plant species in "ectomycorrhizal" families (e.g., Fagaceae, Pinaceae, and Betulaceae) but also

from other plant species (Fig. 2; Additional file 6: Data S6). In addition, the data included

network links between arbuscular mycorrhizal fungi and ectomycorrhizal plant species

(Additional file 6: Data S6) as reported previously [79]. Although such plant–fungus

associations that do not fall into classic categories of mycorrhizal symbioses seldom attract

attention and they are often removed from high-throughput sequencing datasets, some of them

may represent important ecological interactions. An ectomycorrhizal fungus in the truffle genus

(*Tuber melanosporum*), for instance, is known to cause severe necrosis in root cortices of

non-ectomycorrhizal herbaceous plants [80]. Thus, for the standardization of plant–fungus

network analyses inferred with high-throughput sequencing, we need to take into account the

possibility that network links can represent not only mutualistic but also neutral and

antagonistic interactions [17]. Given also that even well-known combinations of plant–fungus

mycorrhizal interactions can result in antagonistic interaction depending on soil environmental

conditions and host plant nutrition [81, 82], potential diversity of ecological interactions within

a network and its community-scale consequences [72] deserve intensive research.

Our community-scale comparative analysis targeting a broad latitudinal range from

cool-temperate to subtropical regions has some implications for geographic diversity patterns

of plant-associated fungi, although careful interpretation is required given the small number of

study sites. The number of detected ectomycorrhizal fungal OTUs was lower in subtropical

than in temperate forests (Fig. 3a), presumably reflecting geographic variation in the relative

abundance of Fagaceae, Pinaceae, and Betulaceae plants in plant communities as discussed in

previous studies [83-86] (see also [87]). In contrast, the number of arbuscular mycorrhizal

fungal OTUs increased towards south in our data, while a previous meta-analysis detected no

latitudinal diversity gradient regarding the fungal functional group [88] (see also [89]). The

total number of fungal OTUs was also higher in subtropical forests, peaked in the southernmost

site. Interestingly, unlike other study sites, the southernmost sampling site was characterized by

low levels of network-scale interaction specificity and host plant differentiation as well as by

the absence of anti-nested network architecture. Although some pioneering studies have

investigated host preferences of tropical fungi [90-92], it remains a major challenge to examine

whether the observed latitudinal gradient in plant–fungus network structure is extended to

tropical regions.

Conclusions

Based on the large datasets of root-associated fungi, we herein showed how plant–fungus

network architecture vary across a latitudinal gradient across the Japanese Archipelago. For

further understanding the diversity of below-ground pant–fungus associations, more

comparative studies of community-scale characteristics are required especially in the tropics.

Moreover, further data of networks consisting of pathogenic fungi and their host plants are

awaited to discuss community-scale properties of negative plant–soil feedbacks [93]. Given

that the number of pathogenic fungi included in our present analysis was too few to evaluate

statistical features of their networks, selective sampling of pathogen infected plant individuals

may be necessary. Improving reference databases of fungal functions is also an important

challenge towards better understanding of the roles of fungal communities. More

macroecological studies of plant–fungus interactions [75, 94, 95], along with experimental

studies testing functions of poorly characterized fungi [11, 13, 14], will reorganize our

knowledge of terrestrial ecosystem processes.

Abbreviations

ANOVA: analysis of variance; DDBJ: DNA Data Bank of Japan; ITS: internal transcribed

spacer; LCA: lowest common ancestor; OTU: Operational taxonomic unit; QCauto method:

query-centric auto-k-nearest neighbor method.

Acknowledgements

We thank Teshio Experimental Forest (Hokkaido University), Tomakomai Experimental Forest

(Hokkaido University), Sugadaira Research Station (Tsukuba University), Yona Field (Ryukyu

University), Tropical Biosphere Research Center (Ryukyu University), and Forestry Agency of

Japan for the permission of fieldwork and Takayuki Ohgue, Takahiko Koizumi, Miyako

Natsume, Ayu Narita, Yoriko Sugiyama, and Yamato Unnno, Yuko Sawanobori, and Minato

Kodama for their support in molecular experiments.

Funding

This work was financially supported by JSPS KAKENHI Grant (26711026), JST PRESTO

(JPMJPR16Q6), and the Funding Program for Next Generation World-Leading Researchers of

Cabinet Office, the Government of Japan (GS014) to HT.

Availability of data and materials

The Illumina sequencing data were deposited to DNA Data Bank of Japan (DDBJ Sequence

Read Archive: DRA006339). The raw data of fungal community structure and the fungal

community matrices analyzed are available as Additional files 1-6.

Authors' contributions

HT designed the work. HT, HS, SY, and AST conducted fieldwork. HT, HS, and SY performed

the molecular experiments. HT wrote the manuscript with HS, SY, and AST.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable

Ethics approval and consent to participate

Not applicable

Additional files

Additional file 1: Data S1. Information of study sites, taxonomic and functional-group

information of the fungal OTUs detected.

Additional file 2: Data S2. Sample-level matrices of plant–fungus associations.

Additional file 3: Data S3. Sequences of the non-glomeromycete fungal OTUs detected.

Additional file 4: Data S4. Sequences of the glomeromycete fungal OTUs detected.

Additional file 5: Data S5. Species-level matrices of plant–fungus associations.

Additional file 6: Data S6. Network data matrices.

Additional file 7: Data S7. Results of the randomization analysis.

References

1.   van der Heijden MG, Bardgett RD, van Straalen NM: The unseen majority: soil
      microbes as drivers of plant diversity and productivity in terrestrial ecosystems. Ecol
      Lett 2008;11:296-310.

2.   Wardle DA, Bardgett RD, Klironomos JN, Setälä H, van der Putten WH, Wall DH:
      Ecological linkages between aboveground and belowground biota. Science
      2004;304:1629-1633.

3.   Peay KG, Kennedy PG, Talbot JM: Dimensions of biodiversity in the Earth mycobiome.
      Nature Rev Mircobiol 2016;14:434-447.

4.   Blackwell M: The Fungi: 1, 2, 3… 5.1 million species? Am J Bot 2011;98:426-438.

5.   Finlay R, Read D: The structure and function of the vegetative mycelium of
      ectomycorrhizal plants. II. The uptake and distribution of phosphorus by mycelial
      strands interconnecting host plants. New Phytol 1986;103:157-165.

6.   Nara K: Ectomycorrhizal networks and seedling establishment during early primary

succession. New Phytol 2006;169:169-178.

7.   Azcón-Aguilar C, Barea J: Arbuscular mycorrhizas and biological control of soil-borne
plant pathogens–an overview of the mechanisms involved. Mycorrhiza
1997;6:457-464.

8.   Smith SE, Read DJ: Mycorrhizal symbiosis. New York: Academic press; 2008.

9.   Augspurger CK: Seed dispersal of the tropical tree, *Platypodium elegans*, and the
escape of its seedlings from fungal pathogens. J Ecol 1983;71:759-771.

10. Mangan SA, Schnitzer SA, Herre EA, Mack KM, Valencia MC, Sanchez EI et al:
Negative plant-soil feedback predicts tree-species relative abundance in a tropical forest.
Nature 2010;466:752-755.

11. Narisawa K, Kawamata H, Currah RS, Hashiba T: Suppression of *Verticillium wilt* in
eggplant by some fungal root endophytes. Eur J Plant Pathol 2002;108:103-109.

12. Almario J, Jeena G, Wunder J, Langen G, Zuccaro A, Coupland G et al: Root-associated
fungal microbiota of nonmycorrhizal *Arabis alpina* and its contribution to plant
phosphorus nutrition. Proc Natl Acad Sci USA 2017;114:E9403–E9412.

13. Hiruma K, Gerlach N, Sacristán S, Nakano RT, Hacquard S, Kracher B et al: Root
endophyte *Colletotrichum tofieldiae* confers plant fitness benefits that are phosphate
status dependent. Cell 2016;165:464-474.

14. Newsham KK: A meta-analysis of plant responses to dark septate root endophytes. New
Phytol 2011;190:783-793.

15. Addy H, Piercey M, Currah R: Microfungal endophytes in roots. Can J Bot
2005;83:1-13.

16. Jumpponen A: Dark septate endophytes–are they mycorrhizal? Mycorrhiza
2001;11:207-211.

17. Toju H, Guimarães PR, Jr, Olesen JM, Thompson JN: Assembly of complex
plant–fungus networks. Nature Commun 2014;5:5273.

18. Mills KE, Bever JD: Maintenance of diversity within plant communities: soil pathogens
as agents of negative feedback. Ecology 1998;79:1595-1601.

19.   Packer A, Clay K: Soil pathogens and spatial patterns of seedling mortality in a
493   temperate tree. Nature 2000;404:278-281.

494   20.   Bever JD, Dickie IA, Facelli E, Facelli JM, Klironomos J, Moora M et al: Rooting
495   theories of plant community ecology in microbial interactions. Trends Ecol Evol
496   2010;25:468-478.

497   21.   Bruns TD, Bidartondo MI, Taylor DL: Host specificity in ectomycorrhizal
498   communities: what do the exceptions tell us? Integr Comp Biol 2002;42:352-359.

499   22.   McGuire KL: Common ectomycorrhizal networks may maintain monodominance in a
500   tropical rain forest. Ecology 2007;88:567-574.

501   23.   Toju H, Yamamoto S, Sato H, Tanabe AS, Gilbert GS, Kadowaki K: Community
502   composition of root-associated fungi in a *Quercus*-dominated temperate forest:
503   "codominance" of mycorrhizal and root-endophytic fungi. Ecol Evol
504   2013;3:1281-1293.

505   24.   Öpik M, Metsis M, Daniell T, Zobel M, Moora M: Large-scale parallel 454 sequencing
506   reveals host ecological group specificity of arbuscular mycorrhizal fungi in a
507   boreonemoral forest. New Phytol 2009;184:424-437.

508   25.   Allen EB, Allen MF, Helm DJ, Trappe JM, Molina R, Rincon E: Patterns and regulation
509   of mycorrhizal plant and fungal diversity. Plant Soil 1995;170:47-62.

510   26.   Bever JD: Negative feedback within a mutualism: host–specific growth of mycorrhizal
511   fungi reduces plant benefit. Proc R Soc Lond Ser B 2002;269:2595-2601.

512   27.   Põlme S, Bahram M, Jacquemyn H, Kennedy P, Kohout P, Moora M et al: Host
513   preference and network properties in biotrophic plant–fungal associations. New Phytol
514   2018;217:1230-1239.

515   28.   Bennett AE, Daniell TJ, Öpik M, Davison J, Moora M, Zobel M et al: Arbuscular
516   mycorrhizal fungal networks vary throughout the growing season and between
517   successional stages. PLOS ONE 2013;8:e83241.

518   29.   Chagnon PL, Bradley RL, Klironomos JN: Using ecological network theory to evaluate
519   the causes and consequences of arbuscular mycorrhizal community structure. New

Phytol 2012;194:307-312.

30.   Montesinos-Navarro A, Segarra-Moragues JG, Valiente-Banuet A, Verdú M: The

network structure of plant–arbuscular mycorrhizal fungi. New Phytol

2012;194:536-547.

31.   Bahram M, Harend H, Tedersoo L: Network perspectives of ectomycorrhizal

associations. Fungal Ecol 2014;7:70-77.

32.   Toju H, Tanabe A, Ishii H: Ericaceous plant–fungus network in a harsh

alpine–subalpine environment. Mol Ecol 2016;25:3242-3257.

33.   Jacquemyn H, Brys R, Merckx VS, Waud M, Lievens B, Wiegand T: Coexisting orchid

species have distinct mycorrhizal communities and display strong spatial segregation.

New Phytol 2014;202:616-627.

34.   Bascompte J, Jordano P, Melián CJ, Olesen JM: The nested assembly of plant-animal

mutualistic networks. Proc Natl Acad Sci USA 2003;100:9383-9387.

35.   Olesen JM, Bascompte J, Dupont YL, Jordano P: The modularity of pollination

networks. Proc Natl Acad Sci USA 2007;104:19891-19896.

36.   Bastolla U, Fortuna MA, Pascual-García A, Ferrera A, Luque B, Bascompte J: The

architecture of mutualistic networks minimizes competition and increases biodiversity.

Nature 2009;458:1018-1020.

37.   Staniczenko PP, Kopp JC, Allesina S: The ghost of nestedness in ecological networks.

Nature Commun 2013;4:1391.

38.   Caruso T, Rillig MC, Garlaschelli D: On the application of network theory to arbuscular

mycorrhizal fungi–plant interactions: the importance of basic assumptions. New Phytol

2012;194:891-894.

39.   Thompson JN: The geographic mosaic of coevolution. Chicago: University of Chicago

Press; 2005.

40.   Toju H, Yamamoto S, Sato H, Tanabe AS: Sharing of diverse mycorrhizal and

root-endophytic fungi among plant species in an oak-dominated cool-temperate forest.

PLOS ONE 2013;8:e78248.

41.   Sato H, Murakami N: Reproductive isolation among cryptic species in the
ectomycorrhizal genus *Strobilomyces*: population-level CAPS marker-based genetic
analysis. Mol Phyl Evol 2008;48:326-334.

42.   Toju H, Tanabe AS, Yamamoto S, Sato H: High-coverage ITS primers for the
DNA-based identification of ascomycetes and basidiomycetes in environmental
samples. PLOS ONE 2012;7:e40863.

43.   Lundberg DS, Yourstone S, Mieczkowski P, Jones CD, Dangl JL: Practical innovations
for high-throughput amplicon sequencing. Nature Methods 2013;10:999-1002.

44.   Stevens JL, Jackson RL, Olson JB: Slowing PCR ramp speed reduces chimera
formation from environmental samples. J Microbiol Method 2013;93:203-205.

45.   Hamady M, Walker JJ, Harris JK, Gold NJ, Knight R: Error-correcting barcoded
primers for pyrosequencing hundreds of samples in multiplex. Nature Methods
2008;5:235-237.

46.   Tanabe AS, Toju H: Two new computational methods for universal DNA barcoding: a
benchmark using barcode sequences of bacteria, archaea, animals, fungi, and land
plants. PLOS ONE 2013;8:e76910.

47.   Tanabe AS: Claident v0.2.2016.07.05, a software distributed by author at
http://www.fifthdimension.jp/. 2016.

48.   Rognes T, Mahé F, Flouri T, Quince C, Nichols B: Vsearch: program available at
https://github.com/torognes/vsearch. In.; 2014.

49.   Smith DP, Peay KG: Sequence depth, not PCR replication, improves ecological
inference from next generation DNA sequencing. PLOS ONE 2014;9:e90234.

50.   Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M et al: Towards
a unified paradigm for sequence-based identification of fungi. Mol Ecol
2013;22:5271-5277.

51.   Thiéry O, Vasar M, Jairus T, Davison J, Roux C, Kivistik PA et al: Sequence variation in
nuclear ribosomal small subunit, internal transcribed spacer and large subunit regions of

Rhizophagus irregularis and *Gigaspora margarita* is high and isolate-dependent. Mol

Ecol 2016;25:2816-2832.

52.   Huson DH, Auch AF, Qi J, Schuster SC: MEGAN analysis of metagenomic data.
Genome Res 2007;17:377-386.

53.   Edgar RC: Search and clustering orders of magnitude faster than BLAST.
Bioinformatics 2010;26:2460-2461.

54.   Abarenkov K, Henrik Nilsson R, Larsson KH, Alexander IJ, Eberhardt U, Erland S et
al: The UNITE database for molecular identification of fungi–recent updates and future
perspectives. New Phytol 2010;186:281-285.

55.   Toju H, Yamamoto S, Tanabe AS, Hayakawa T, Ishii HS: Network modules and hubs in
plant-root fungal biome. J R Soc Int 2016;13:20151097.

56.   Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J et al: FUNGuild: an
open annotation tool for parsing fungal community datasets by ecological guild. Fungal
Ecol 2016;20:241-248.

57.   CBOL-Plant-Working-Group, Hollingsworth PM, Forrest LL, Spouge JL, Hajibabaei
M, Ratnasingham S et al: A DNA barcode for land plants. Proc Natl Acad Sci USA
2009;106:12794-12797.

58.   Peay KG, Russo SE, McGuire KL, Lim Z, Chan JP, Tan S et al: Lack of host specificity
leads to independent assortment of dipterocarps and ectomycorrhizal fungi across a soil
fertility gradient. Ecol Lett 2015;18:807-816.

59.   Oksanen J, Blanachet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB et al: Vegan:
community ecology package. R package version 2.0-3 available at
http://CRAN.R-project.org/package=vegan. 2012;10:2008.

60.   R-Core-Team: R 3.4.1: A language and environment for statistical computing available
at http://www.R-project.org/. In. Vienna, Austri: R Foundation for Statistical
Computing; 2015.

61.   Bastian M, Heymann S, Jacomy M: Gephi: an open source software for exploring and
manipulating networks. Proc 3rd Int AAAI Conf Weblogs Soc Media 2009;8:361-362.

62.   Jacomy M, Heymann S, Venturini T, Bastian M: ForceAtlas2, a continuous graph layout
algorithm for handy network visualization designed for the Gephi Software. PLOS

ONE 2011;96:e98679.

63.   Blüthgen N, Menzel F, Blüthgen N: Measuring specialization in species interaction
networks. BMC Ecol 2006;6:9.

64.   Blüthgen N, Menzel F, Hovestadt T, Fiala B: Specialization, constraints, and conflicting
interests in mutualistic networks. Curr Biol 2007;17:341-346.

65.   Toju H, Guimarães PR, Jr, Olesen JM, Thompson JN: Below-ground plant–fungus
network topology is not congruent with above-ground plant–animal network topology.
Sci Adv 2015;1:e1500291.

66.   Almeida-Neto M, Ulrich W: A straightforward computational approach for measuring
nestedness using quantitative matrices. Env Model Software 2011;26:173-178.

67.   Stone L, Roberts A: The checkerboard score and species distributions. Oecologia
1990;85:74-79.

68.   Dormann CF, Fründ J, Blüthgen N, Gruber B: Indices, graphs and null models:
analyzing bipartite ecological networks. Open Ecol J 2009;2:7-24.

69.   Vázquez DP, Melián CJ, Williams NM, Blüthgen N, Krasnov BR, Poulin R: Species
abundance and asymmetric interaction strength in ecological networks. Oikos
2007;116:1120-1127.

70.   Rohr RP, Saavedra S, Bascompte J: On the structural stability of mutualistic systems.
Science 2014;345:1253497.

71.   Allesina S, Tang S: Stability criteria for complex ecosystems. Nature
2012;483:205-208.

72.   Mougi A, Kondoh M: Diversity of interaction types and ecological community stability.
Science 2012;337:349-351.

73.   Sato H, Tsujino R, Kurita K, Yokoyama K, Agata K: Modelling the global distribution
of fungal species: new insights into microbial cosmopolitanism. Mol Ecol
2012;21:5599-5612.

74.   Nagarajan S, Singh D: Long-distance dispersion of rust pathogens. Ann Rev
Phytopathol 1990;28:139-153.

75. Talbot JM, Bruns TD, Taylor JW, Smith DP, Branco S, Glassman SI et al: Endemism
and functional convergence across the North American soil mycobiome. Proc Natl
Acad Sci USA 2014;111:6341-6346.

76. Hubbell SP: The unified neutral theory of biodiversity and biogeography, vol. 32.
Princeton: Princeton University Press; 2001.

77. Mougi A, Kondoh M: Food-web complexity, meta-community complexity and
community stability. Sci Rep 2016;6:srep24478.

78. Gravel D, Massol F, Leibold MA: Stability and complexity in model meta-ecosystems.
Nature Commun 2016;7:12457.

79. Dickie IA, Koide RT, Fayish AC: Vesicular-arbuscular mycorrhizal infection of
Quercus rubra seedlings. New Phytol 2001;151:257-264.

80. Plattner I, Hall I: Parasitism of non-host plants by the mycorrhizal fungus *Tuber*
melanosporum. Mycol Res 1995;99:1367-1370.

81. Hoeksema JD, Chaudhary VB, Gehring CA, Johnson NC, Karst J, Koide RT et al: A

meta-analysis of context-dependency in plant response to inoculation with mycorrhizal

fungi. Ecol Lett 2010;13:394-407.

82. Johnson NC, Graham JH: The continuum concept remains a useful framework for
studying mycorrhizal functioning. Plant Soil 2013;363:411-419.

83. Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R et al: Global
diversity and geography of soil fungi. Science 2014;346:1256688.

84. Tedersoo L, Bahram M, Toots M, Diedhiou AG, Henkel TW, Kjøller R et al: Towards
global patterns in the diversity and community structure of ectomycorrhizal fungi. Mol
Ecol 2012;21:4160-4170.

85. Tedersoo L, Nara K: General latitudinal gradient of biodiversity is reversed in
ectomycorrhizal fungi. New Phytol 2010;185:351-354.

86. Põlme S, Bahram M, Yamanaka T, Nara K, Dai YC, Grebenc T et al: Biogeography of
ectomycorrhizal fungi associated with alders (*Alnus* spp.) in relation to biotic and
abiotic variables at the global scale. New Phytol 2013;198:1239-1249.

87. Gao C, Zhang Y, Shi NN, Zheng Y, Chen L, Wubet T et al: Community assembly of
ectomycorrhizal fungi along a subtropical secondary forest succession. New Phytol
2015;205:771-785.

88. Öpik M, Vanatoa A, Vanatoa E, Moora M, Davison J, Kalwij J et al: The online database
MaarjAM reveals global and ecosystemic distribution patterns in arbuscular
mycorrhizal fungi (Glomeromycota). New Phytol 2010;188:223-241.

89. Pärtel M, Öpik M, Moora M, Tedersoo L, Szava-Kovats R, Rosendahl S et al: Historical

biome distribution and recent human disturbance shape the diversity of arbuscular
mycorrhizal fungi. New Phytol 2017;216:227-238.

90. Tedersoo L, Sadam A, Zambrano M, Valencia R, Bahram M: Low diversity and high
host preference of ectomycorrhizal fungi in Western Amazonia, a neotropical
biodiversity hotspot. ISME J 2010;4:465-471.

91. Sato H, Tanabe AS, Toju H: Contrasting diversity and host association of
ectomycorrhizal basidiomycetes versus root-associated ascomycetes in a dipterocarp
rainforest. PLOS ONE 2015;10:e0125550.

92. Husband R, Herre E, Turner S, Gallery R, Young J: Molecular diversity of arbuscular
mycorrhizal fungi and patterns of host association over time and space in a tropical
forest. Mol Ecol 2002;11:2669-2678.

93. LaManna JA, Mangan SA, Alonso A, Bourg NA, Brockelman WY, Bunyavejchewin S
et al: Plant diversity increases with the strength of negative density dependence at the
global scale. Science 2017;356:1389-1392.

94. Arnold AE, Lutzoni F: Diversity and host range of foliar fungal endophytes: are tropical
leaves biodiversity hotspots? Ecology 2007;88:541-549.

95. Schemske DW, Mittelbach GG, Cornell HV, Sobel JM, Roy K: Is there a latitudinal
gradient in the importance of biotic interactions? Ann Rev Ecol Evol Syst
2009;40:245-269.

Table 1 Potential factors contributing to variation in plant–fungus network structure. For each
response variable representing network structure, an ANOVA model including the number of
plant species/taxa, that of fungal OTUs, network connectance, sampling locality, and fungal
functional groups (i.e., categories of plant–fungus networks) was constructed. *P* values
significant after a Bonferroni correction are shown in bold for ANOVA model.

Response variable	Explanatory variable	df	*F*	*P*
Relative interaction specificity	No. plant species/taxa	1	4.6	0.0426
	No. fungal OTUs	1	96.9	**< 0.0001**
	Connectance	1	10.1	**0.0040**
	Locality	7	4.0	**0.0048**
	Functional group	4	2.6	0.0572
Relative nestedness	No. plant species/taxa	1	5.1	0.0322
	No. fungal OTUs	1	39.3	**< 0.0001**
	Connectance	1	2.4	0.1308
	Locality	7	1.5	0.2216
	Functional group	4	2.1	0.1179
Relative checkerboard score	No. plant species/taxa	1	1.0	0.3182
	No. fungal OTUs	1	62.9	**< 0.0001**
	Connectance	1	5.6	0.0262
	Locality	7	1.3	0.2772
	Functional group	4	4.0	**0.0121**

695
696

699 **Fig. 1** Study sites. **a** Map of study sites. In each forest site, a number in a parenthesis indicates

700 the number of plant species/taxa observed in the 240 terminal root samples from which

701 sequencing data were successfully obtained. **b** Relationship between the number of samples

702 and that of plant species/taxa observed. A rarefaction curve obtained from 240 terminal-root

703 samples is shown for each study site.

Site 1
(cool-temperate)

Site 5
(warm-temperate)

Site 2
(cool-temperate)

Site 6
(subtropical)

Site 3
(cool-temperate)

Site 7
(subtropical)

Site 4
(cool-temperate)

Site 8
(subtropical)

- Plant
- Arbuscular mycorrhizal fungal OTU
- Ectomycorrhizal fungal OTU
- Saprotrophic or endophytic fungal OTU
- Plant pathogenic fungal OTU
- Other/unidentified fungal OTU

Fig. 2 Below-ground plant–fungus networks. The "ALL" network involving all the

root-associated fungal OTUs detected and their host plant species/taxa is shown for each forest.

The OTUs/species in the networks are arranged with the "ForceAtlas2" layout algorithm [62].

Size of circles represents betweenness centrality scores compared within plant/fungal

community.

Fig. 3 Network properties. The index scores representing the architecture of plant fungus

networks/partial networks are shown across the eight forests examined. **a** The number of fungal

OTUs. The code numbers of forest sites correspond to those shown in Figure 1. **b** The ratio of

the number of fungal OTUs to that of the plant species/taxa involved in each network/partial

network. **c** Connectance (the proportion of non-zero entries in a species-level matrix). **d**

Network-level interaction specificity (relative H_2'). **e** Nestedness (relative weighted NODF

nestedness). **f** Host range differentiation (relative checkerboard score). For relative interaction

specificity, relative nestedness, and relative checkerboard score (**d-f**), scores higher/lower than

2 roughly indicate that observed network index values are higher/lower than expected by

chance (see Additional file 7: Data S7 for detailed results of the randomization test).

Fig. 4 Principal component analysis of network properties. **a** Principal component 1 vs. principal component 2. Factor loadings of the examined variables are shown on the right. **b** Principal component 1 vs. principal component 3.

731
732