Subsurface microbial habitats in an extreme desert Mars-analogue environment

Kimberley A. Warren-Rhodes[1,2], Kevin C. Lee[3], Stephen D.J. Archer[3], Donnabella C.

Lacap-Bugler[3], Linda Ng Boyle[4], Nathalie Cabrol[1,2], David Wettergreen[5], Kris Zacny[6],

Cecilia Demargasso[7], Jeff Moersch[8], Guillermo Chong[7], Srinivasan Vijayarangan[5], Cinthya

Tebes[7], Michael Wagner[5], Ken Tanaka[9], Trent Hare[9], Christopher Gayle Tate[8], Alian

Wang[10], Jie Wei[10], Greydon Foil[5] and Stephen B. Pointing[11]

[1] NASA Ames Research Center, Moffett Field, CA, USA.

[2] The SETI Institute, Mountain View, CA, USA

[3] School of Sciences, Auckland University of Technology, Auckland, New Zealand.

[4] College of Engineering, University of Washington, Seattle, WA, USA

[5] Institute of Robotics, Carnegie-Mellon University, Pittsburgh, PA, USA

[6] Honeybee Robotics and Spacecraft Mechanisms Corp, Pasadena, CA, USA

[7] Universidad Católica del Norte, Antofagasta, Chile

[8] Department of Earth and Planetary Science, University of Tennessee, Knoxville, TN, USA

[9] United States Geological Survey, Flagstaff, AZ, USA

[10] Department of Chemistry, Washington University, St. Louis, OH, USA

[11] Yale-NUS College and Department of Biological Sciences, National University of

Singapore, Singapore

Summary

Sediments in the hyper-arid core of the Atacama Desert represent one of the closest

terrestrial analogues to Mars' regolith. Understanding the distribution and drivers of life

in the sediment may thus give critical clues on how to search for biosignatures on Mars.

Here, we identify the spatial distribution of highly specialised bacterial communities in

previously unexplored depth horizons of subsurface sediments and their correlation

with moisture and geochemical variables. We deployed an autonomous rover in a

mission-relevant Martian drilling scenario with manual sample validation to recover and

analyse sediments in two Mars-like terrains, a gravel desert pavement and an evaporite-

rich playa. Subsurface communities were distinct from surface-associated communities

and were delineated by depth related to sediment moisture. Geochemical analysis

indicated soluble salts and minerology that influence water bio-availability, particularly

in deeper sediments. Colonisation was also patchy and uncolonised sediment was

associated with indicators of extreme osmotic challenge. Bacterial diversity reflected

strong selection for halotolerant and desiccation tolerant taxa throughout depth

horizons. The most diverse communities occurred within the zone of highest moisture,

whereas deeper sediments were colonized by methylotrophic taxa with limited

diversity. The study provides important baseline microbial ecological data on the linkage

between biocomplexity, moisture and geochemistry in Mars-like sediments at the limit

of habitability and demonstrates feasibility of the rover-mounted drill for future Mars

sample recovery.

Introduction

The surface of Mars is dry, cold, and exposed to high levels of ionising radiation.

However, data accumulated over the past few decades by orbital and landed missions

have demonstrated that early in its history the planet may have been habitable for

microbial life with abundant sources of energy, carbon, nutrients, and shelter [1]. Mars

supported surface and subsurface water and may still do in some circumstances, as well

as organic molecules required for life [2]. As a result, Mars 2020 and ExoMars missions will

be searching for biosignatures [3,4], and the investigation of terrestrial analogues can

provide critical insights for the development and testing of exploration strategies.

Among those, the hyper-arid core of the Atacama Desert in Chile is widely regarded as a

tractable Mars analogue in the field of astrobiology. The Atacama is the driest desert

region on Earth [5] with extremely low moisture inputs and precipitation events that are

stochastic in nature [6]. The region has a long history of climatic stability as an extreme

desert [7,8] resulting in the build-up of evaporates [6,9] and creation of a Mars-like surface.

The desert's hyper-arid core lies at or near the arid limit for soil formation [10] and thus

surface terrain in this region is regarded as sediment.

Animal and plant life are scarce in extreme deserts and instead cyanobacteria-

dominated microbial communities in mineral and rocky refugia including deliquescent

substrates comprise the dominant surface biota and are well-characterised [11,12].

Conversely, evidence for microbial colonisation in hyper-arid Atacama sediment is

scarce, contradictory and almost exclusively limited to surface-associated sediment [9,13–

17] (Supplementary Material, Table S1). Cultivation-based approaches are unreliable as

indicators of environmental microbial diversity and have yielded estimates that varied

by several orders of magnitude [9,14,15,17,18]. Biochemical tests have similarly yielded

inconclusive support for microbial metabolic activity [9,14–17]. High throughput sequencing

of environmental DNA/RNA remains the most reliable indicator for microbial diversity in

extreme desert sediments. This approach has provided critical insight on surface-

associated communities in semi-arid, arid and hyper-arid locations [16]. A recent study

also estimated putative microbial metabolic activity in three subsurface samples after

an unusual rain event and this suggested subsurface sediment may also be a habitat for

microbial communities [17]. Major knowledge gaps persist, and importantly from an

astrobiology perspective the question of how microbial occurrence and diversity may

vary with abiotic variables in sediment depth horizons of a Mars analogue and spatially

within terrain, and whether this can be addressed in a simulated robotic sampling

mission scenario.

To document these questions and support the development and testing of

biosignature exploration strategies, the NASA-funded *Subsurface Life in the Atacama*

Project deployed an autonomous rover-mounted robotic drill in Mars-like desert

pavement and playa terrain within the hyper-arid core of the Atacama under

challenging environmental and logistical constraints (Supplementary Material, Fig. S1-

S3). The drill accessed 32 discreet sediment samples to depths of 800mm along a 50km

transect in a realistic simulation of Martian drilling operations and constraints. Over 60

manually excavated sediment samples were recovered in parallel and validated the

automated sampling for abiotic and biotic components of the sediment habitat and

provide important baseline ecological data on the most Mars-like sediments on Earth.

Extreme habitats in Atacama sediment horizons

The on-board Raman spectrometry revealed surface mineral distributions comprising

feldspar and quartz. Subsurface sediments, and particularly the playa sites, were

elevated in gypsum and anhydrite (Supplementary Material, Table S3, Fig. S5). This was

independently corroborated by laboratory Raman measurements (Supplementary

Material, Table S3) and geochemical analysis that revealed a positive linear correlation

(r=0.78, p<0.001) across all samples between sulfate-sulfur and calcium, the

components of gypsum and anhydrite ($CaSO_4$ +/-$2H_2O$). Anydrite is indicative of drier

conditions at depth, and *in situ* soil moisture sensors validated this for horizons in both

pavement and playa.

Our chemical analysis focused on geochemical reservoirs readily available and

relevant to biological communities and using methodology that was comparable to

other microbial ecology studies. A clear depth-dependent pattern in geochemistry was

revealed for pavement and playa horizons (Fig. 1, Supplementary Material, Table S2, Fig.

S4). Surface sediments were strongly associated with elevated phosphorous

concentrations in both playa and pavement units (Fig. 1). Elevated surface P was

attributed to presence of surface Fe-oxides and calcite that can adsorb P. These P

adsorbing phases were likely present in the subsurface but were diluted by high sulfate

concentrations causing lower recorded P concentrations below the surface. All

subsurface samples displayed very low to undetectable levels of N and low C throughout

horizons. The on-board Raman spectrometer also recorded C as a minor/undetectable

phase in all samples. Electrical conductivity (EC) and Na^+ levels increased with depth in

both substrates indicating increased salinity with depth (p = 0.006). Desert pavement

subsurface sediments separated mainly due to pH and K^+, whilst playa subsurface

sediments displayed elevated levels of sulfate-sulfur, extractable cations (Ca^{2+}, Mg^{2+},

Na^+) and EC indicating increasing osmotic challenge, and most notably in the paleo-playa

Site 10 (Fig. 1, Supplementary Material Table S2).

Average annual subsurface temperatures ranged from 19.9 – 20.9^{o}C in depth

horizons (Supplementary Material, Table S4). At shallow depths (<100mm) temperature

varied between 4 – 34^{o}C, whereas in deeper horizons temperature variation was

somewhat buffered (Supplementary Material, Fig. S6). Moisture values from depth

horizons in representative playa terrain were consistently higher (approximately 4- to 7-

fold) at all depths than those in the desert pavement horizons (Supplementary Material,

Table S4) and this likely reflects local hydrology given the playa was a relatively low-lying

terrain (Supplementary Material, Fig. S1). However, sediment moisture trends in both

substrates broadly indicated the existence of depth groupings into distinct moisture

zones: (a) a surface zone consisting of the top 200mm, where water availability is

typically lowest, except in the short-term following a rain event; (b) a mid-depth zone

(300-500mm), where water availability peaks and persists after a rain event; and (c) a

deep subsurface zone (≥500-800mm), where water availability is typically lower and,

notably for the desert pavement sediments, appeared to be un-impacted by rare large

rainfall events (Supplementary Material, Table S4, Fig. S6).

Depth-related trends in DNA recovery and microbial alpha diversity

Surface colonisation was widespread in desert pavement and playa sites but subsurface

horizons displayed patchy recovery with low yields of quantifiable DNA in the range

0.067-6.5 ng/g sediment, indicating extremely low standing biomass (Supplementary

Material, Table S2, S5). Notably, the paleo-playa site yielded no recoverable DNA at any

depth after repeated efforts at extraction. Our approach using a DNA recovery method

adapted for low-biomass extreme environments highlights for the first time the

patchiness of low biomass microbial colonisation in the most extreme subsurface desert

sediments where micro-habitat conditions are at or near the limit for life [19]. This may

explain, at least in part, why some previous research has concluded this region of the

Atacama was lifeless [9]. Linear Discriminant Analysis was employed to show that

variables most strongly associated with potentially lifeless subsurface sediments where

environmental DNA was irrecoverable were sulfate-sulfur (substrate) (p = 0.001), depth

(p = 0.003), EC (p = 0.006), soluble salts (p = 0.006). This strongly suggests that osmotic

challenge and limited moisture availability are the major extinction drivers in this Mars

analogue sediment. We also demonstrated that substrate-inhibition of DNA recovery

was unlikely to have been a significant factor, since bacterial cell suspensions in the

range reported for desert soils (Supplementary Material, Table S1) added to these

samples were recoverable with 60-80% efficiency compared to a pure laboratory culture

and were not significantly inhibited compared to extractions with spiked sediments that

did originally yield environmental DNA.

Across both terrains any given depth horizon tended to either be colonised

almost throughout or display only patchy near-surface colonisation as measured by

recoverable DNA. Thus, the ubiquitous surface colonisation was not a predictor for

subsurface habitability in these sediments. We postulate that in addition to sediment

moisture, extreme geochemistry also influenced habitability in these sediments as

evidenced by high soluble salts/salinity indicators and abundance of anhydrite and

gypsum mineral signatures (Supplementary Material, Table S2, S3). Temperature

variations were not regarded as significant challenges to microbial colonisation

compared to other variables (Supplementary Material, Fig. S6). The paleo-playa site 10

failed to yield recoverable DNA from any depth, thus highlighting that ancient landforms

created under moisture sufficiency, and with elevated anhydrite levels as indicated by

the rover's Raman spectrometer (Supplementary Material, Fig. S5) may not support

extant genetic biosignatures and do not support recoverable relic DNA. Our overall DNA

recovery success was consistent with our expectations for the driest desert location on

Earth. It is notable that we have observed Antarctic mineral sediments, another Mars

analogue, generally yield higher recovery rates using similar methodologies, e.g. [20,21].

However, this is likely due to enhanced standing biomass due to the less extreme nature

of the growing season in Antarctic desert where long periods of frozen hibernation are

punctuated by periods of moisture sufficiency during the austral summer [22].

Clear patterns were discernible from manually excavated sediment pits. Bacterial

diversity decreased significantly with depth across both terrains (Chao1 richness: r =

0.508, p = 0.044; Shannon's H: r = 0.731, p = 0.001), thus indicating depth as a major

driver of bacterial diversity (Supplementary Material, Fig. S8, Table S6). Bacterial

communities formed six diversity clusters that associated with clearly defined depth

ranges and zones of variability in sediment moisture and geochemistry (Fig. 2a).

Subsurface communities were more distinct between habitat types but also more

heterogenous overall than surface-associated communities, due largely to the lower

bacterial diversity within each depth-defined sediment micro-habitat. Community

evenness displayed little clear pattern with depth for desert pavement but for playa

deep samples displayed very low evenness reflecting their highly specialised diversity

(Supplementary Material, Table S6). In all cases the rover-acquired samples showed a

generally similar pattern in depth profile and diversity clusters although these were less

clearly defined and we attributed this to mixing of sediment during recovery using the

bite-drilling approach (Fig. 2b; Supplementary Material, Fig. S8).

To further unravel the influence of sediment environment on bacterial diversity,

we performed canonical correspondence analysis (CCA) to establish the association of

distinct geochemistry for desert pavement and playa samples on the assembly of

bacterial communities (Fig. 1). Surface communities were strongly influenced by P levels

that were elevated compared to the subsurface (Fig. 1). Bacterial diversity in subsurface

sediment habitats correlated with two groups of geochemical variables: The playa

subsurface community was strongly associated with EC and extractable cations (Ca^{2+},

Mg^{2+}, Na^+) and the rover-mounted Raman data indicated elevated anhydrite and

gypsum (Supplementary Material, Table S3, Fig. S5). This strongly suggests that bio-

availability of water may be limited in this subsurface habitat despite relatively high

moisture levels due to the effects of salt saturation. Conversely the pavement

subsurface community was associated largely with pH, and also C and K^+ although the

magnitude of variation for these variables was low. This pH-dependent structuring of

microbial diversity is consistent with observations for global trends in soil microbial

diversity [23]. The BEST multiple rank correlation routine was employed to further

examine our data and rank the relative association of abiotic variables with the

observed bacterial diversity as follows: extractable cations (Ca^{2+}, Mg^{2+}, Na^+) and

phosphorous>sulfate-sulfur> >EC ($_pw$ = 0.595-0.609, p <0.05), thus further validating the

ordinations and CCA analysis.

Highly specialised sediment bacterial communities

The taxa identified in sediment indicated highly specialised and relatively low diversity

bacterial communites in both desert pavement and playa, and this low diversity reflects

global trends in soil microbial diversity where deserts are considered to be relatively

depauperate [24,25]. Overall the drill samples yielded weaker depth resolution than

manual sampling but still corroborated observations from the manually collected

samples (Fig. 3, Supplementary Material Fig. S9). Bacterial taxonomic diversity varied

significantly with subsurface habitat as compared to a generally consistent surface

community (Fig. 3) [16]. Overall, communities were dominated largely by only three phyla:

Chloroflexi, Actinobacteria and Alphaproteobacteria.

All surface communities were dominated by the AKIW781 lineage of the

photoheterotrophic Chloroflexi, a known desert soil crust taxon (NCBI GenBank

accession number: JQ402146), and overall community structure contributed additional

taxonomic resolution to previous observations of surface sediment communities [16,17].

AKIW781 has also been recorded in desert varnish on rock surfaces [26] as well as a

keystone taxon of hypolithic communities [27] in the Atacama. This indicates a

cosmopolitan distribution and broad habitat preference among surface niches in this

extreme desert. Conversely, the Chloroflexi were minor components of subsurface

communities, decreasing in relative abundance with depth, and mainly comprised an

uncharacterised candidate class Ellin6529, likely adapted to non-phototrophic

metabolism in the subsurface microhabitat.

Low and mid-depth subsurface horizons for both desert pavement and playa

were dominated by the low G-C gram positive taxonomic class Actinobacteria (Fig. 3,

Supplementary Material, Fig. S9). They are typically thick-walled bacteria commonly

encountered in soil. Communities encountered were specific to each depth horizon and

shifts in diversity clearly reflected the moisture zones identified for both pavement and

playa horizons. At shallow depths (<200mm) halotolerant, alkalotolerant, spore-forming

desiccation-tolerant actinobacterial groups were abundant and included the orders

acidimicrobiales, gaiellales, nitriliruptorales and solirubrobactales in desert pavement

and acidimicrobiales and nitriliruptorales in playa. Mid-depth sediments (300-<500 mm)

corresponding to the zone of greatest moisture availability in both substrates supported

highly variable bacterial diversity between samples and also the most diverse

communities (Supplementary Material, Table S6). Of note was the elevated relative

abundance of bacillales at a single pavement site and this may reflect the relatively high

altitude of this site and limited moisture input to the subsurface. The bacillales-

dominated community was more similar to those encountered at greater depths at

other locations. The bacillales are characterised by their ability to form highly resistant

endospores and so may indicate an increasingly challenging micro-habitat in deeper

subsurface sediments.

There were fewer deep sediment samples from which to make comparisons and

this reflected the more challenging subsurface environment, and deeper communities

(500+ mm) generally displayed lowest taxonomic diversity. The deepest desert

pavement community was similar to those at shallower depths but displayed elevated

relative abundance of acidimicrobiales. Conversely the deep playa community shifted to

a fundamentally different composition from shallower depths that was dominated

almost exclusively by a single facultative methylotrophic and desiccation-tolerant

Methylobacterium radiotolerans taxon (NCBI GenBank accession number: LT985974)

from the phylum Alphaproteobacteria. We speculate the C1 metabolism of this taxon

allows it to exploit simple C1 compounds as well as subsurface methane sources[28], a

molecule known to be released from subsurface sources on Mars [29]. An overall picture

thus emerges of highly-specialised bacterial diversity adapted to and reflecting the

challenging subsurface habitat as well as reflecting moisture availability zones

determined at least in part by sediment depth and geochemistry.

Other bacteria typically encountered in desert surface communities and

generally regarded as tolerant to extreme conditions were not major components of

subsurface communities. A complete absence of cyanobacterial taxa was consistent

with their adaptation to surface mineral refugia rather than subsurface sediment

habitats that preclude photoautotrophy [12], and the highly desiccation-tolerant

Deinococcus-Thermus group were represented by only a single lineage of Trueperaceae

candidate genus B-42 recovered in just a few subsurface samples with low abundance

(0.6 – 4.8%), compared to a relatively diverse assemblage of cultivable Deinococci

recovered from surface sediments in other less arid desert locations [30]. This further

supports our evidence for highly specialised subsurface communities selected for by the

distinct geochemistry and microclimate in the subsurface Mars analogue sediments of

the Atacama. They may thus be broadly indicative of the type of microbial consortia that

could exploit Martian subsurface habitats.

In the deepest and least-diverse playa communities biocomplexity was reduced

almost to a single taxon, reflecting extreme selective pressure and also highlighting a

possible lack of resilience to environmental change given that recruitment to deep

subsurface sediments may be limited. The minimum biocomplexity may comprise

multiple ecotypes of a single taxon, each adapted to exploit a given suite of

microclimate and geochemical conditions [31]. They likely exhibit a strong preference for

C1 and/or autotrophic taxa that are somewhat de-linked from their immediate

surroundings in terms of carbon sequestration and reflect the extreme oligotrophic

nature of these microhabitats. Given the highly specialised diversity recovered, the

association of putative physiology with environmental variables, and lack of DNA

recovery in paleo-playa samples; we conclude that our estimates are likely to represent

resident microbial communities rather than relic DNA[17,32].

Implications for detection of biosignatures on Mars

The autonomous rover drilling platform yielded subsurface sediment samples that

allowed for the first time a combined investigation of sediment abiotic properties and

microbial diversity at an unprecedented level of detail. The parallel manual sampling

and analysis demonstrated feasibility and fidelity of the autonomous rover approach.

Highly specialised but low-diversity subsurface bacterial communities were encountered

patchily and this was strongly associated with abiotic variables which suggests that

"follow the water" is only part of the biosignature exploration solution in the search for

potential habitable refuges on Mars. Consideration of subsurface micro-habitat

variability in geochemistry, originating with and adapted to possible water availability

zones may also be key. Whilst the geochemistry of our analogue sites was similar to that

of a habitable Martian regolith, moisture in the Atacama is surface-sourced by fog

and/or rain events [6], whereas on Mars subsurface sources may provide an upward

migration of moisture similar to that observed in Antarctic mineral sediment overlaying

permafrost [33,34]. Thus, extrapolating habitable subsurface locations on Mars would need

to consider this along with the incident radiation regime and other Martian

environmental variables. Based on the exploratory design consideration for this NASA-

funded research, drill depth was constrained at 800mm as a proof of concept study.

However, current plans are for both NASA and ESA to target depths of up to 2m on Mars

in order to target potential subsurface habitats that account for Martian environmental

variables.

The relevance of ecology and microbial habitats to past and possible extant life

on Mars are finally coming to the fore in the robotic search for biosignatures on Mars

[1,35]. As our study suggests, detecting such life or its residual biosignatures may prove

highly challenging, given that in the most extreme deserts on Earth these communities

are extremely patchy in distribution and occur with exceedingly low biomass. The drill

apparatus employed in this study has demonstrated that subsurface sediment

biosignatures can be autonomously recovered, although precise depth delineation

requires refinement with the bite-drilling approach used in this study. Whilst genetic

biosignatures such as DNA used in our study may not ultimately be the primary method

employed to search for traces of life on Mars, they are the most reliable and widely

used method currently available for microbial diversity estimation [24,25]. This approach

provided essential first proof of concept that an incontrovertible biological signature

within the likely range for geochemical variables in a habitable subsurface environment

can be recovered from a Mars-like sediment using an autonomous rover.

Methods

A detailed description of all field and experimental approaches is presented in the

Supplementary Material. Briefly, the automated rover completed a 50km traverse and

used a robotic drill to sample subsurface sediments in depth horizons to 800mm. On-

board mineralogical characterisation was achieved using Raman spectroscopy and

samples were recovered for further laboratory geochemical analysis and estimation of

bacterial diversity from sequencing of environmental DNA. Parallel manually excavated

1 samples were recovered for validation of rover-acquired sampling, and the first detailed

2 characterisation of subsurface bacterial habitats in the Mars analogue sediment.

3 Automated microclimate recording was achieved using dataloggers deployed *in situ* in

4 the depth horizons.

6 **References**

7 1. Cabrol, N. A. The Coevolution of Life and Environment on Mars: An Ecosystem

8   Perspective on the Robotic Exploration of Biosignatures. *Astrobiology* **18,** 1–27

9   (2018).

10 2. Martin-Torres, F. J. *et al.* Transient liquid water and water activity at Gale crater

11   on Mars. *Nat. Geosci* **8,** 357–361 (2015).

12 3. Vago, J. L. *et al.* Habitability on Early Mars and the Search for Biosignatures with

13   the ExoMars Rover. *Astrobiology* **17,** 471–510 (2017).

14 4. Farley, K. & Williford, K. Seeking Signs of Life and More: NASA's Mars 2020

15   Mission. *EOS Trans. - Am. Geophys. Union* **98,**

16   https://doi.org/10.1029/2017EO066153 (2017).

17 5. Peel, M. C. & Finlayson, B. L. Updated world map of the Köppen-Geiger climate

18   classification. *Hydrol. Earth Syst. Sci.* **11,** 1633–1644 (2007).

19 6. McKay, C. P. *et al.* Temperature and Moisture Conditions for Life in the Extreme

20   Arid Region of the Atacama Desert: Four Years of Observations Including the El

21   Nino of 1997-1998. *Astrobiology* **3,** 393–406 (2003).

22 7. Clarke, J. D. A. Antiquity of aridity in the Chilean Atacama Desert. *Geomorphology*

73, 101–114 (2006).

8. Hartley, A. J., Chong, G., Houston, J. & Mather, A. E. 150 million years of climatic stability: evidence from the Atacama Desert, northern Chile. *J. Geol. Soc. London.* **162,** 421–424 (2005).

9. Navarro-Gonzalez, R. *et al.* Mars-like soils in the Atacama Desert, Chile, and the dry limit of microbial life. *Science* **302,** 1018–1021 (2003).

10. Ewing, S. A. *et al.* A threshold in soil formation at Earth's arid–hyperarid transition. *Geochim. Cosmochim. Acta* **70,** 5293–5322 (2006).

11. Pointing, S. B. & Belnap, J. Microbial colonization and controls in dryland systems. *Nat. Rev. Microbiol.* **10,** 551–562 (2012).

12. Azua-Bustos, A., Urrejola, C. & Vicuña, R. Life at the dry edge: Microorganisms of the Atacama Desert. *FEBS Lett.* **586,** 2939–2945 (2012).

13. Drees, K. P. *et al.* Bacterial community structure in the hyperarid core of the Atacama Desert, Chile. *Appl Env. Microbiol* **72,** 7902–7908 (2006).

14. Connon, S. S. A., Lester, E. D. E., Shafaat, H. H. S., Obenhuber, D. C. & Ponce, A. Bacterial diversity in hyperarid Atacama Desert soils. *J Geophys Res - Biogeosci* **112,** G04S17 (2007).

15. Lester, E. D., Satomi, M. & Ponce, A. Microflora of extreme arid Atacama Desert soils. *Soil Biol. Biochem.* **39,** 704–708 (2007).

16. Crits-Christoph, A. *et al.* Colonization patterns of soil microbial communities in the Atacama Desert. *Microbiome* **1,** 28 (2013).

17. Schulze-Makuch, D. *et al.* Transitory microbial habitat in the hyperarid Atacama

Desert. *Proc. Natl. Acad. Sci. U. S. A.* **115,** 2670–2675 (2018).

18. Bagaley, D. R. Uncovering bacterial diversity on and below the surface of a hyper-arid environment, The Atacama Desert, Chile. (Louisiana State University, 2006).

19. McKay, C. P. Requirements and limits for life in the context of exoplanets. *Proc. Natl. Acad. Sci. U. S. A.* **111,** 12628–33 (2014).

20. Pointing, S. B. *et al.* Highly specialized microbial diversity in hyper-arid polar desert. *Proc. Natl. Acad. Sci. U. S. A.* **106,** 19964–19969 (2009).

21. Lee, C. K. *et al.* The Inter-Valley Soil Comparative Survey: the ecology of Dry Valley edaphic microbial communities. *ISME J* **6,** 1046–1057 (2012).

22. Pointing, S. S. B. *et al.* Biogeography of photoautotrophs in the high polar biome. *Front. Plant Sci. Funct. Plant Ecol.* **6,** 692 (2015).

23. Fierer, N. & Jackson, R. B. The diversity and biogeography of soil bacterial communities. *Proc. Natl. Acad. Sci. U. S. A.* **103,** 626–631 (2006).

24. Thompson, L. R. *et al.* A communal catalogue reveals Earth's multiscale microbial diversity. *Nature* **551,** 457–463 (2017).

25. Delgado-Baquerizo, M. *et al.* A global atlas of the dominant bacteria found in soil. *Science.* **359,** 320–325 (2018).

26. Kuhlman, K. R., Venkat, P., La Duc, M. T., Kuhlman, G. M. & McKay, C. P. Evidence of a microbial community associated with rock varnish at Yungay, Atacama Desert, Chile. *J. Geophys. Res.* **113,** G04022 (2008).

27. Lacap, D. C., Warren-Rhodes, K. A., McKay, C. P. & Pointing, S. B. Cyanobacteria and chloroflexi-dominated hypolithic colonization of quartz at the hyper-arid core

of the Atacama Desert, Chile. *Extremophiles* **15,** 31–8 (2011).

28. Kao, Y. *et al.* Scale-dependent key drivers controlling methane oxidation potential

in Chinese grassland soils. *Soil Biol. Biochem.* **111,** 104–114 (2017).

29. Stevens, A. H., Patel, M. R. & Lewis, S. R. Modelled isotopic fractionation and

transient diffusive release of methane from potential subsurface sources on

Mars. *Icarus* **281,** 240–247 (2017).

30. Rainey, F. A. *et al.* Extensive diversity of ionizing-radiation-resistant bacteria

recovered from Sonoran Desert soil and description of nine new species of the

genus *Deinococcus* obtained from a single soil sample. *Appl. Environ. Microbiol.*

71, 5225–5235 (2005).

31. Koeppel, A. *et al.* Identifying the fundamental units of bacterial diversity: a

paradigm shift to incorporate ecology into bacterial systematics. *Proc. Natl. Acad.*

Sci. U. S. A. **105,** 2504–9 (2008).

32. Carini, P. *et al.* Relic DNA is abundant in soil and obscures estimates of soil

microbial diversity. *Nat. Microbiol.* **2,** 16242 (2016).

33. Goordial, J. *et al.* Nearing the cold-arid limits of microbial life in permafrost of an

upper dry valley, Antarctica. *ISME J.* **10,** 1613–1624 (2016).

34. Stomeo, F. *et al.* Abiotic factors influence microbial diversity in permanently cold

soil horizons of a maritime-associated Antarctic Dry Valley. *FEMS Microbiol Ecol*

82, 326–340 (2012).

35. Warren-Rhodes, K. *et al.* Robotic ecological mapping: Habitats and the search for

life in the Atacama Desert. *J. Geophys. Res. Biogeosciences* **112,** (2007).

Acknowledgements

The work was completed with funding from NASA Astrobiology Science and Technology

for Exploring Planets (ASTEP) Program Grant NNX11AJ87G and a Yale-NUS College Start-

Up Grant. The authors are grateful to Brad Sutter (NASA) for advice on interpretation of

geochemical analysis, and Craig Cary (University of Waikato) for assistance with

biosecurity compliance and sample custody.

Author contributions

K.A.W-R., N.C. and S.B.P. conceived the study; D.W., K.Z., S.V., M.W. and G.F. built and

operated the rover; K.A.W-R.; C.D., J.M., G.C., C.T., K.T., T.H., C.G.T., A.W. and J.W.

conducted field operations and sampling; K.C.L., S.D.J.A., D.C.L-B. and S.B.P. processed

sediment samples and conducted geochemical analysis; S.D.J.A. and K.C.L. conducted

molecular biology experiments; K.A.W-R, K.C.L., S.D.J.A., D.C.L-B, L.N-B. and S.B.P.

performed data analysis; K.A.W-R. and S.B.P. wrote the manuscript with inputs from

K.C.L., S.D.J.A., D.W. and N.C. (K.A.W-R. and K.C.L. contributed equally).

Materials & Correspondence

Correspondence and requests for materials should be addressed to K.A.W-R Email:

krhodes@seti.org or S.B.P. Email: stephen.pointing@yale-nus.edu.sg

Figure 1. Correlation of Mars analogue sediment geochemistry with bacterial diversity.

Canonical Correspondence Analysis (CCA) triplot with symmetrical scaling indicating

differences in sediment geochemistry within sediment pits, and influence of these

abiotic variables on bacterial communities and individual taxa. The three most abundant

taxa are labelled (A, B, C). The circle size of each sample indicates species richness index

(Chao 1) of the respective community.

Figure 2. Depth-defined bacterial diversity in Mars analogue sediments of the

Atacama Desert. Non-metric multidimensional scaling (NMDS) ordination of Bray Curtis

similarities for bacterial diversity versus sediment depth from a) manual recovery, and

b) rover recovery. Shaded areas indicate similarity clusters for sediment communities at

the same depth. The size of each symbol (circle or triangle) indicates species richness

index (Chao 1) of the respective community. Mid-range values were used for drill

samples where depth ranges instead of individual depths were generated.

Figure 3. Highly specialised bacterial communities in Mars analogue sediment depth

2 **horizons.** Distribution of bacterial diversity by taxonomic class with sediment depth for

3 manual (M) and autonomous rover drill (D) recovered samples. Coloured shading

4 indicates relative abundance within each community for a given bacterial class. Grey

5 shading indicates no recoverable bacteria.