FERONIA's sensing of cell wall pectin activates ROP GTPase signaling in *Arabidopsis*

Wenwei Lin [1,2], Wenxin Tang [1,2], Charles T. Anderson [3], and Zhenbiao Yang [1,2*]

[1]FAFU-UCR Joint Center for Horticultural Biology and Metabolomics Center, Haixia Institute of Science and Technology, Fujian Agriculture and Forestry University, Fuzhou, Fujian, China
[2]Center for Plant Cell Biology, Institute of Integrative Genome Biology, and Department of Botany and Plant Sciences, University of California, Riverside, CA92521, USA
[3]Department of Biology, The Pennsylvania State University, University Park, PA, 16802, USA

*Correspondence to: Zhenbiao Yang (yang@ucr.edu).

ABSTRACT

Plant cells need to monitor the cell wall dynamic to control the wall homeostasis required for a myriad of processes in plants, but the mechanisms underpinning cell wall sensing and signaling in regulating these processes remain largely elusive. Here, we demonstrate that receptor-like kinase FERONIA senses the cell wall pectin polymer to directly activate the ROP6 GTPase signaling pathway that regulates the formation of the cell shape in the *Arabidopsis* leaf epidermis. The extracellular malectin domain of FER directly interacts with de-methylesterified pectin *in vivo* and *in vitro*. Both loss-of-FER mutations and defects in the pectin biosynthesis and de-methylesterification caused changes in pavement cell shape and ROP6 signaling. FER is required for the activation of ROP6 by de-methylesterified pectin, and physically and genetically interacts with the ROP6 activator, RopGEF14. Thus, our findings elucidate a cell wall sensing and signaling mechanism that connects the cell wall to cellular morphogenesis via the cell surface receptor FER.

INTRODUCTION

Mounting evidence suggests that cell wall polymers provide signals to regulate a large number of plant processes and that plant cells can monitor cell wall dynamics or damages [1-5]. For instance, cell wall biosynthesis and cell wall remodeling-related genes are reprogramed in various cell wall-related mutants[6,7] or in plants treated with cellulose synthesis inhibitor [8] or pectin fragments [9]. Upon the interferences with cell wall status, various compensatory wall responses were induced, including ectopic deposition of lignin and callose [10-12], altered pectin modification status [8,13,14], reactive oxygen species (ROS) production [11], and an elevation in jasmonic acid and ethylene production [11,15,16]. Furthermore, cell wall dynamic plays a regulatory role in the regulation of complex cell shapes [7]. However, the mechanisms for sensing and transducing the wall signals remain largely mysterious. The elucidation of these mechanisms is solely needed to understand how cell expansion and shape changes are coordinated between neighboring cells in a growing tissue or organ.

The spatiotemporal pattern of cell walls (and as such their compositions and structure) is critical for cell expansion and shape formation, because of high cellular turgor pressure in plants [17,18]. Precise coordination and communication via the dynamic cell wall between adjacent cells to regulate these cellular processes are required for organ differentiation, growth, and morphogenesis, and yet the underlying mechanisms remain elusive. The jigsaw puzzle-shaped pavement cells (PCs) of the leaf epidermis serve as an exciting model to investigate the mechanisms for cell-cell coordination of cell shape in a multicellular system [19]. PCs form interdigitated lobes and indentations that are tightly coordinated between the adjacent cells. This process is regulated by two antagonistic RAC/ROP GTPase signaling pathways in an auxin-dependent manner [20]. Locally activated ROP2/ROP4 signaling promotes the outgrowth to form the lobe region, whereas ROP6 at the indenting region promotes the ordering of cortical microtubules (MTs), which restrict radial or lateral cell expansion [21].

Mechanical stress from lobe outgrowth has been implicated in the indentation
reinforcement by promoting cortical MT ordering [22], hinting at a potential regulatory
role of the cell wall in this process. This is further supported by a recent study
suggesting that the cell wall compositions and mechanical heterogeneities across and
along anticlinal cell walls contribute to the regulation of PC interdigitation[7]. The
identification of cell wall sensors is needed to understand the mechanisms by which
the cell wall modulates various plant processes such as PC interdigitation. A large
number of cell surface receptors, such as Wall-associated kinases (WAKs),
Catharanthus roseus RLK1 (CrRLK1)-like family, PR5-like receptor
kinase/thaumatin family, LysM family, L-type lectin RLKs, proline-rich extension-
like receptor kinase (PERK) family, leucine-rich repeat extensins (LRXs), FEI1/2,
LRR-RK MALE DISCOVERER 1-INTERACTING RECEPTOR LIKE KINASE
2/LEUCINE-RICH REPEAT KINASE FAMILY PROTEIN INDUCED BY SALT
STRESS (MIK2/LRR-KISS) and LRR receptor-like protein 44 (RLP44) have been
proposed to be potential wall sensors[1,5,23,24 25-34], but none of them have been clearly
demonstrated to sense and transduce cell wall polymer signals. Here we report that the
FERONIA cell surface receptor senses and transmits cell wall pectin to activate the
ROP signaling pathway to modulate the formation of the puzzle-piece cell shape in
Arabidopsis PCs.
83
84   **RESULTS**
85
86   **The FERONIA (FER) cell surface receptor is associated with the cell wall**
87
Plants have evolved a large number of cell surface receptor-like kinases (RLKs) to
sense various extracellular signals [35]. To identify RLKs that might be involved in the
lobe-indentation coordination during PC interdigitation, we carried out a genetic
screen of T-DNA insertion *rlk* mutants for altered PC morphogenesis and isolated *fer-*
4, a mutant of *FERONIA* [36], which exhibited a severe defect in PC shape (Fig. 1a, b).
Besides root hair growth and development, FER was suggested to be involved in cell
polarization, which is important for the formation of the interdigitated shape in
Arabidopsis leaf epidermal PCs [37]. To further evaluate the role of FER in PC
morphogenesis, we characterized another *fer* allele with an independent T-DNA
insertion (*fer-2* and *fer-5*, Supplementary Fig. 1a)[38,39] and found that both alleles
showed similar defects in PC interdigitation. Both *fer* mutants displayed significantly
wider indentations than wild-type (WT) (Fig. 1a, b, and Supplementary Fig. 1b-d),
reminiscent of a defect in the ROP6 signaling pathway [20]. To further confirm that the
phenotypes observed in the *fer* mutants were due to the mutation in *FER*, we
complemented the *fer-4* mutant with *FER* cDNA fused with yellow fluorescent
protein (*YFP)* under the control of its native promoter. The *FER-YFP* transgene
completely rescued the *fer-4* PC shape defect and restored vegetative growth to wild-
type levels (Fig. 1a, b, and Supplementary Fig. 1e). Together, the data indicate that
FER is required for PC morphogenesis.

**Figure 1. FER is associated with the cell wall and required for *Arabidopsis* epidermal pavement

cells (PCs) morphogenesis. a and b,** PC phenotypes of 2 day-after-germination (DAG) wild-type

(Col-0), *fer-4* and complementary line *pFER::FER-YFP/ fer-4* (a). The degree of PC interdigitation

was determined by the width of indentation (b). Data was generated from the measurement of at least

20 cells collected from 5 different cotyledons from 5 individual seedlings. Asterisk indicates a

significant difference with p≤0.05 (Student's *t*-test) between wild-type and the *fer-4* mutant. Data

are represented as mean ± SE. Scale bar, 50 μm. **c and d,** FER-GFP proteins are enriched in the

insoluble fraction. Soluble (S) and insoluble (P) proteins were fractionated and analyzed by Western

blot (WB) with α-GFP antibody (c) and the ratio of two fractions was quantified (d). The plasma

membrane-localized RLK *BRI1-GFP* transgenic plant was used as a control. The data are shown as

mean ± SE of three repeats. **e,** The cell wall localization of FER-GFP. (*Top* panels) GFP

fluorescence was along with the cell surface of PCs before plasmolysis (-). During plasmolysis (+) GFP

fluorescence localized with the cytoplasm (*Bottom* left panel) and plasma membrane indicated by

overlapping signal with FM4-64 (Merge). A portion of the FER-GFP signal retreated with the

cytoplasm localized on the plasma membrane indicated by overlapping signal with FM4-64 (Merge)

and another portion of the signal remained on the cell surface (*Bottom* right panel). Arrows indicate the

cell wall residue FER-GFP signal. Scale bar, 50 μm.

FER belongs to the *Arabidopsis* subfamily of *Catharanthus roseus* RLK1-like kinases

(CrRLK1Ls) with 17 members all containing a malectin-like domain predicted to bind

carbohydrates (Supplementary Fig. 1a) [3]. In *Arabidopsis*, *FER* was initially found to

control pollen tube perception in the ovule [36] and later shown to regulate a wide range

of growth and developmental processes and responses to the environment [38-46].

Interestingly several CrRLK1L members play important roles in controlling processes

linked to the properties of the cell wall, including cell expansion, polarized growth,

sperm release from pollen tubes, pollen tube integrity maintenance, and defense

responses [24,36,40,41,47-49]. In *Arabidopsis,* FER and its relatives, ANUXR1, ANUXR2,
BUPS1, and BUPS2 are involved in maintaining cell wall integrity and to regulate
cell growth through the recognition of a group of small peptide ligands that derived
from cells named Rapid Alkanilization Factors (RALFs) [24,48,50]. Another CrRLKL1
family member, THESUS1(THE1), identified as a putative cell-wall integrity sensor,
mediates the responses to the perturbation of cellulose synthesis, likely via binding to
the cell wall [3,30]. Therefore, we speculated that FER might have a role as a cell wall
sensor to monitor the dynamics of the cell wall during PCs morphogenesis, and
assessed whether FER is associated with the cell wall. We generated transgenic plants
expressing FER-green fluorescent protein (GFP) fusion driven by its native promoter.
BRI1, a known plasma membrane-localized protein, was used as a control [51]. Soluble
(supernatant) and insoluble (pellet) proteins were fractionated from FER-GFP and
BRI1-GFP transgenic plants by using low-speed centrifugation. As expected, non-
wall-associated BRI1-GFP was enriched in the soluble fraction (Fig. 1c, d). In
contrast, FER-GFP was enriched in the insoluble pellet fraction (Fig. 1c, d). As shown
for wall-associated kinase proteins (WAKs) [28,52], the extraction of FER-GFP from the
insoluble pellet fraction required boiling in the presence of 1% (w/v) SDS and 50 mM
dithiothreitol (DTT) (Supplementary Fig. 1f), indicating that FER is tightly associated
with the cell wall.
153
To further verify the cell wall association of FER, cotyledon PCs from transgenic
seedlings expressing FER-GFP fusion or GFP alone were plasmolyzed (Fig. 1e).
Before plasmolysis, the GFP control exhibited a typical free-GFP localization pattern
in plant cells with a large central vacuole; the signals were largely diffused along the
cell border (cell walls stained by Propidium Iodide, PI) (Supplementary Fig. 1g).
After plasmolysis, the GFP signals retreated with the plasma membrane (plasma
membrane stained by FM4-64) that was detached from the cell walls (Fig. 1e,
Supplementary Fig. 1g, the lower row of panels). By contrast, FER-GFP was mainly
found along the plasma membrane, largely overlapping with the cell wall PI staining
(Supplementary Fig. 1g). After plasmolysis, a portion of the FER-GFP fluorescence
retreated with the plasma membrane (white arrows), while predominant signal
remained at the cell border (Fig. 1e, Supplementary Fig. 1g, the lower row of panels).
These results are consistent with the high enrichment of the FER-GFP protein in the
cell wall fraction, suggesting that FER is a plasma membrane RLK that is tightly
associated with the cell wall.
169

Figure 2. FER binds pectin through the MALA domain to regulate PC morphogenesis. a, FER
was associated with pectin *in vitro*. FER-HA was transiently expressed in *Arabidopsis* protoplasts.
Pull-down (PD) was carried out with pectin and the proteins were determined by Western blot (WB)
with α-HA antibody. *Top* shows that FER-HA pull-down by pectin (PD: pectin; WB: α-HA). *Bottom*
shows the expression of FER-HA proteins (WB: α-HA). **b**, Association of truncated FER with pectin *in
vitro*. Different specific deletion domains of FER-HA were expressed in *Arabidopsis* protoplasts, the
PD assays were carried out with pectin. *Top* shows that truncated FER-HA was pulled-down by pectin
(PD: pectin; WB: α-HA). *Bottom* shows the expression of FER-HA truncated proteins (WB: α-HA). **c,**
FER-ECD directly bound to pectin *in vitro*. His-MBP-FER-ECD was purified from *E. coli*. Pull down
was carried out as above and the proteins were determined by WB with α-MBP antibody. His-MBP-
FER-ECD pull-down by pectin (*Top*). The input His-MBP and His-MBP-FER-ECD proteins (*Bottom*).
d, PC phenotypes in wild-type, *fer-4* and the complementary line *35S::YFP-FER-TCD/ fer-4*. Scale
bar, 20 μm. **e**, The degree of 3 DAG PC interdigitation. Asterisk indicates the average indentation
widths that were significantly different (p≤0.05, Student's *t*-test) between wild-type and the *fer-4*
mutant, wild-type, and *35S:: YFP-FER-TCD* complementation line, respectively. **f**, PC phenotypes in
wild-type, *fer-4* and the complementary line *35S::YFP-FER-ΔMALA/ fer-4. Scale bar, 50* μm. **g**, The
degree of 6 DAG PC interdigitation. Asterisk indicates the average indentation widths that were
significantly different (p≤0.05, Student's *t*-test) between wild-type and *fer-4* mutants, wild and the
35S:: YFP-FER- ΔMALA complementation line, respectively.

The extracellular malectin A domain is responsible for FER's association with pectin and is required for FER's function in PCs

We next determined whether FER is associated with a specific cell wall component
using a semi-*in vitro* pull-down assay. The FER protein fused to an HA tag was

expressed in *Arabidopsis* leaf protoplasts, and proteins isolated from the protoplasts
were incubated with different cell wall components. Insoluble cellulose and
hemicellulose (xylan) were pelleted by centrifugation, but FER-HA was not detected
in either the cellulose or xylan pellets (Supplementary Fig. 2a). In contrast, FER-HA
was detected in the pellet of pectin extracted from apple (50-75% degree
esterification) in a concentration-dependent manner (Fig. 2a), indicating an interaction
between FER and pectin. We then dissected the domains of FER responsible for
binding pectin (Supplementary Fig. 2b). FER-ECD-HA (where the intracellular
domain was deleted) and FER△MALB-HA (deletion of the extracellular MALB
domain) were pulled down by pectin, but FER-TCD-HA (deletion of the entire
extracellular domains) and FER△MALA-HA (deletion of the extracellular MALA
domain) were not (Fig. 2b). Importantly, we found that His-MBP-FER-ECD
recombinant proteins, expressed and purified from *E. coli*, directly bound to pectin i*n*
vitro (Fig. 2c). This suggests that glycosylation is not required for FER's function in
binding to pectin. Furthermore, in agreement with the finding that MALA domain is
required for the interaction (Fig. 2b), we found that His-MBP-FER-MALA domain
recombinant proteins, expressed and purified from *E. coli*, directly bound pectin *in*
vitro (Supplementary Fig. 2c). Thus, these results demonstrate that FER physically
binds to pectin through the extracellular MALA domain.
215
We then determined whether FER's extracellular MALA domain is also required for
the in vivo association with the cell wall and function. We generated *35S::YFP-FER-*
TCD transgenic plants and detected the protein levels in both soluble and insoluble
protein fraction using a GFP antibody. In contrast to the full-length FER-GFP protein,
which was enriched in the insoluble pellet fraction (Fig. 1c, d), YFP-FER-TCD was
enriched in the soluble fraction (Supplementary Fig.2d). Furthermore, the fluorescent
signal of YFP-FER-TCD in plasmolyzed PCs was absent from the cell walls
(Supplementary Fig. 2e). Thus the cytoplasmic domain is not involved in FER's
association with the cell wall. Since the MALA domain bound pectin *in vitro* (Fig. 2b
and Supplementary Fig. 2c), we further tested its requirement for cell wall association
in the *35S::YFP-FER△MALA* transgenic *Arabidopsis* plants by using plasmolysis
assays. In contrast to FER-GFP (Fig. 1e), the majority of YFP-FER△MALA regressed
with the shrunk cytoplasm (Supplementary Fig. 2f), indicating that the MALA
domain is critical for FER's association with the cell wall. Finally, neither YFP-FER-
TCD nor YFP-FER-△MALA was able to fully rescue the PC interdigitation defect in
the *fer-4* knockout mutant (Fig. 2d-g), which was fully rescued by the full-length
FER-YFP (Fig. 1a). Taken together, our results indicate that the extracellular MALA
domain binds to cell wall pectin, allowing FER to associate with the cell wall, and
regulates PC morphogenesis.
235
236 **Pectin methylation level is spatially regulated and important for PC shape**
237
The generation of lobes and indentations in PCs results from the anisotropic growth of
the cell wall and involves the preferential deposition of cellulose in the indentation
region [53]. As expected mutants defective in cellulose synthesis typically exhibit
simpler PC shapes [54] [7]. The pectin composition and structure in the wall in much more

complicated and has been proposed to have a regulatory role[3,55-60]. With the finding
that FER's association with pectin is critical for PC morphogenesis, we sought to
determine the contribution of the cell wall pectin to the formation of PC shapes. In
Arabidopsis, three types of pectic polysaccharides are found: homogalacturonan
(HG), rhamnogalacturonan I (RG-I), and rhamnogalacturonan II (RG-II) [61]. Mutants
with reduced HG levels (*qua1-1* and *qua2-1*) exhibit decreased PC interdigitation[7].
We found that the *arad1arad2* mutant, which has reduced arabinans associated with
rhamnogalacturonan I (RG-I), showed PC shape changes similar to those observed in
the *fer* mutants with wider indentation necks (Fig. 3a, b). These results suggest that
the specific composition and structure of pectic polysaccharides are critical for PC
morphogenesis. This agrees with the importance of fine pectin structures in
modulating other biological roles[7,60].

254
Figure 3. Pectin methylation levels modulate PC morphogenesis and pectin-FER interaction. a,
Cotyledon PC phenotypes in wild-type and the *arad1arad2* mutant. Scale bar, 100 μm. **b,** The degree
of 3 DAG PC interdigitation. Asterisk indicates a significant difference (p≤0.05, Student's *t*-test)
between wild-type and the *arad1arad2* mutant. **c**, Expanding young PCs from 2 DAG cotyledons were
immuno-stained for highly methylesterified or de-methylesterified pectin with LM20 (left) and JIM5
(right) antibodies, respectively. The lobe and indentation regions analyzed were indicated by label or
arrows. Scale bar, 10 μm. **d,** Signal intensity ratios between lobe and indentation regions. The data are
shown as mean ± SE of 20 cells. **e,** Mature PCs from 3 weeks-old seedlings (the third leaf pairs) were
immunostained for highly methylesterified or de-methylesterified with LM20 (left) and JIM5 (right)
antibodies, respectively. The lobe and indentation region analyzed were indicated by arrows. Scale bar,
10 μm. **f,** Signal intensity ratios between lobe and indentation regions. The data are shown as mean ±
SE of 20 cells. **g,** Cotyledon PC phenotypes in wild-type, *pme3* and *PMEI1-OE* mutants. Scale bar, 100
μm. **h,** The degree of 2 DAG PC interdigitation. Asterisks indicate a significant difference (p≤0.05,
Student's *t*-test) between wild type and the mutants. **i,** FER was preferentially associated with de-
methylesterified pectin (PGA) in protoplasts compared to highly methylesterified pectin (pectin). Pull-
down was carried out with PGA and pectin as described above (Fig. 2a). (*Top*) FER-HA proteins from

protoplasts were pull-down by PGA or pectin. (*Bottom*) The His-MBP-FER-ECD proteins were
determined by WB as input control. **j,** FER-ECD preferentially bound de-methylesterified pectin
(PGA) *in vitro* compared to highly methylesterified pectin (pectin). (*Top*) His-MBP tagged FER-ECD
recombinant proteins purified from *E. coli* were pulled down by PGA or pectin and detected by α -His
antibody. (*Bottom)* The His-MBP-FER-ECD proteins were determined by western blotting (WB) as
input control.

Highly methylesterified pectic polysaccharides are synthesized in the Golgi and
secreted into the apoplast [61]. The methyl groups are then selectively removed by
pectin methylesterases (PMEs), which might be spatially regulated by pectic
methylesterase inhibitors (PMEIs) [60]. De-methylesterified HG chains are crosslinked
into a tightly packed conformation by calcium bridges, which contribute to the wall
strength [32,62]. However, in some tissues decline in HG-calcium complexes has been
correlated with a decrease in wall expansibility and an increase in wall stiffening [56,63].
Thus, the diversity and complexity in pectin structure and distribution in different
tissues/cells and in different wall domains within a single cell might reflect functional
diversity in the fine control of pectin gel rheology [60]. Given the importance of pectin
for PC morphogenesis (Fig. 3a, b)[7], we further determined the pattern of pectin
methylesterification in developing PCs (2-day cotyledons) by immunostaining with
the LM20 and JIM5 monoclonal antibodies that recognize highly methylesterified and
highly de-methylesterified pectin, respectively. The intensity of LM20 staining was
greater in the lobe region when compared to the indentation area of the same cell,
indicating a preferential distribution of highly methylesterified pectin in the lobe
region (Fig. 3c, d). In contrast, higher levels of de-methylesterified pectin were found
in the indentation region, comparing to the adjacent cell lobe region as indicated by
the JIM5 immunofluorescence intensity (Fig. 3e, f). Note that this pectin staining
pattern was observed in stage II PCs[20,64] when PCs have just attained their jigsaw
puzzle pattern and are still developing. PCs at early stages (undifferentiated and stage
I cells) or late stage (fully expanded stage III) likely have very different cell wall
structures and composition [20]. Indeed PCs on the late stage (the third leaves of 3-week
old plants) did not exhibit a clear preference in the distribution of highly de-
methylesterified and methylesterified pectin between the lobe and indentation regions
(Fig. 3e, f). These results suggest that dynamic changes in pectin structure occur
during the development of interdigitated PCs.

To investigate the functional significance of pectin de-methylesterification levels in
PC morphogenesis, we characterized PC shape in mutants with reduced pectin de-
methylesterification. We examined PCs phenotype in a knockout mutant for *PME3*
(*AT3G14310*), highly expressed in expanding cotyledons and true leaves [65] [66] and in a
line overexpressing a pectin methylesterase inhibitor (PMEI1-OE) [67]. Both lines
showed interdigitation defects similar to those of the *fer-4* mutant with simpler cell
shapes and wider indentation necks (Fig. 3g, h). These results suggest that pectin
methylesterifcation is spatially regulated at the subcellular level during the early
stages of and is important for PC morphogenesis.

FER preferentially binds de-methylesterified pectin

The significance of pectin methylesterifcation levels described above promoted us to
determine its impact on pectin binding to FER. Interestingly FER proteins were
readily pulled down by de-methylesterified pectin. Both full-length FER protein
expressed from protoplasts (Fig. 3i) and FER-ECD recombinant protein produced
from *E.coli* (Fig. 3j) interacted much more strongly with polygalacturonic acid (PGA,
0% methylesterification) compared with pectin extracted from apple (50-75%
methylesterified) (Fig. 3i, j). Furthermore, *in vitro* pull-down assays showed that
FER's MALA domain preferentially bound de-methylesterifed pectin (PGA)
compared to the highly methyesterified pectin (Supplementary Fig. 3a).
327

Since Ca^{2+} crosslinks de-methlyesterified HG via the carboxy groups, we investigated
the importance of this conformation in binding with FER. We determined the binding
of FER to PGA in the ionic buffer (0.5 mM $CaCl_2$/150 mM NaCl) containing 5 mM,
EDTA, a Ca^{2+} chelator, or in an ionic buffer, in which $CaCl_2$ was replaced with 0.5
mM $MgCl_2$. EDTA prevents the formation of PGA crosslinks, and Mg^{2+} ions are
unable to stabilize PGA crosslinks due to their large size [62]. Both EDTA and $MgCl_2$
treatments greatly suppressed the binding of FER to PGA (Supplementary Fig. 3b).
Thus pectin crosslinks are critical for binding to FER. Taken together, our results
suggest that FER favors binding de-methylesterified crosslinked pectin through the
extracellular MALA domain to regulate PC morphogenesis.
338

339   **Pectin activates ROP6 signaling in a FER-dependent manner**
340

Because both mutations in *FER* and reductions in pectin de-methylesterification result
in a PC phenotype similar to that induced by ROP6 signaling defects [20], we
hypothesized that binding of de-methylesterified pectin to FER activates the ROP6
signaling pathway. We first determined the requirement of FER for ROP6 activation
using an effector binding-based assay [68]. When the assay was conducted using an anti-
ROP6 antibody, we observed dramatic ROP6 activity reduction in the *fer-4* mutant
compared to wild-type (Fig. 4a, b). This reduction was further confirmed by another
assay, in which GFP-ROP6 was introduced into Col-0 and the *fer-4* mutant and
detected with an anti-GFP antibody (Supplementary Fig. 4a, b). Consistent with the
changes in PC phenotypes of *pme3* and *PMEI1-OE* (Fig. 3g, h), ROP6 activation was
also compromised in both *pme3* and *PMEI1-OE* lines (Fig. 4a, b). These results
support the hypothesis that FER's sensing of de-methylesterified pectin leads to the
activation of the intracellular ROP6 signaling pathway. To further test this hypothesis,
we determined whether pectin-mediated activation of ROP6 requires FER. Col-0
protoplasts were treated with PGA, and the activation of ROP6 was determined.
Intriguingly, PGA treatment greatly induced ROP6 activation (Fig. 4c, d), and
importantly this induction was compromised in *fer-4* mutant protoplasts (Fig. 4c, d).
Hence PGA promotes ROP6 activation in a FER-dependent manner.

Figure 4. FER regulates PC morphogenesis through ROP signaling. a and b, Active ROP6 in

wild-type, *fer-4, pme3* and *PMEI1-OE* mutants was analyzed by pull-down with the effector RIC1, as

described previously (44), and the pulled down ROP6 was determined by a ROP6 antibody (a). The

relative active of ROP6 level was quantified (b). 7 days seedlings were used in this assay. **c and d**,

Activation of ROP6 by PGA in *Arabidopsis* protoplasts. ROP6 activities were determined in wild-type

and *fer-4* protoplasts, isolated from 4 weeks adult plants, that were treated with or without 20 μg PGA

(c). The relative active of ROP6 level was quantified (d). Data are mean activity levels from three

independent experiments ±SE (b and d). **e and f**, PC cortical microtubules of wild-type (*GFP-MAP4*)

and the *fer-4* mutant (*fer-4xGFP-MAP4*) (e). *Bottom* panel showed the magnified PC indentation

regions. The degree of cortical microtubules anisotropy was quantified (f). Arrows indicate the

indentation regions. Scale bar, 10 μ m. Data are mean degrees from 20 independent cells ±SE. **g and**

h, PC cortical microtubules of wild-type (*GFP-TUB*) and the *PMEI1-OE* line (*PMEI1-OExGFP-TUB*)

(e). *Bottom* panel showed the magnified PC indentation regions. The degree of cortical microtubules

anisotropy was quantified (h). Arrows indicate the indentation regions. Scale bar, 50 μ m. Data are

mean degrees from 20 independent cells ±SE. Asterisks indicate the significant difference (p≤0.05,

Student's *t*-test) between the wild type and the mutants in above assays.

The activation of ROP6 signaling promotes cortical microtubule organization to

restrict lateral or radial cell expansion, thereby promoting the indentation of PCs [21]. In

agreement with the FER-dependent activation of ROP6 by de-methylesterified pectin

and the phenotypes of *fer-4* PCs with wider indentation necks (Fig. 1a, b and

Supplementary Fig. 1c, d), *fer-4* PCs displayed less bundled and more randomly

arranged cortical MTs, compared with wild-type cells (Fig. 4e, f). Furthermore, a

similar reduction in the ordered arrangement of cortical MTs in the indentation region

of PCs was found in the *PMEI1-OE* line (Fig. 4g, h, and Supplementary Fig. 4c, d)

and *pme3* mutants (Supplementary Fig. 4c, d).

RopGEF14 provides a direct link between FER and ROP6

ROPs are directly activated by ROP guanine nucleotide exchange factors (RopGEFs).
RopGEF1 interacts with the FER kinase domain to activate ROP2 signaling in root
hair development [39]. To assess which RopGEFs might be involved in PC
morphogenesis, we analyzed the expression pattern of all 14 GEFs in different tissues,
and found that only *RopGEF1, 6, 7,* and *14* transcripts were detected in cotyledons
(Supplementary Fig. 5a). Furthermore, the characterization of PC morphogenesis of
T-DNA insertion loss-of-function mutants for these RopGEFs revealed that two
alleles of *gef14* exhibited a defect in PC interdigitation (Fig. 5 a, b, Supplementary
Fig. 5b-i). Cotyledon PCs for these mutant alleles exhibited wider indentation
compared to wild-type, reminiscent of a defect in the ROP6 signaling pathway [20]. The
gef14-2 PC indentation defect was completely rescued by complementing *gef14-2*
with the *GEF14* cDNA fused with a Myc tag and its native promoter (Fig. 5a, b).
Consistently, ROP6 activation (Fig. 5c, d, Supplementary Fig. 5j) and the ordering of
cortical MTs were greatly compromised in the *gef14-2* mutant (Fig. 5e-g). Thus, we
hypothesize that FER activates ROP6 signaling directly through RopGEF14 to
regulate PC morphogenesis.

405
Figure 5. RopGEF14 regulates PC morphogenesis through ROP signaling. a and b, PC
phenotypes of wild-type (Col-0), *gef14-2* and complementary line *pGEF::GEF-4xMyc/ gef14-2* (a).
The degree of PC interdigitation was determined by the width of indentation (b). Asterisk indicates a
significant difference with p≤0.05 (Student's *t*-test) between wild-type and the *gef14-2* mutant. Data
are represented as mean ± SE. Scale bar, 50 μm. **c and d**, Activation of ROP6 in the wild-type and
gef14 mutants were analyzed by pull-down and determined by a ROP6 antibody (c). The relative active
of ROP6 level was quantified (d). **e and f**, PC cortical microtubules of wild-type (*GFP-TUB*) and the
gef14-2 mutant (*GFP-TUB x gef14-2*) (e). *Bottom* panel showed the magnified PC indentation regions.
The degree of cortical microtubules anisotropy was quantified (f). Arrows indicate the indentation
regions. Scale bar, 50 μ m. Data are mean degrees from 20 independent cells ±SE. Asterisks indicate
the significant difference (p≤0.05, Student's *t*-test) between the wild type and the mutants in above
assays.
418

To test this hypothesis, we first determined whether RopGEF14 forms a complex with
FER and ROP6. We performed coimmunoprecipitation (CO-IP) assays in *pGEF14-*
GEF14-Myc transgenic plants. As shown in Fig. 6 a and b, RopGFF14-Myc
coimmunoprecipitated FER and ROP6, which were detected with α-FER[42]and α-
ROP6 antibodies, respectively (Fig. 6a, b). The association of FER with RopGEF14
was further confirmed using *35s-GFP-GEF14* transgenic plants. FER
coimmunoprecipitated GFP-RopGFE14 but not GFP alone from transgenic plants
(Supplementary Fig. 6a). These co-immunoprecipitation results are also consistent
with those of yeast-two-hybrid assay suggesting a direct interaction between GEF14
and FER [39]. Moreover, recombinant MBP-GEF14 proteins specifically pulled down
the full length of FER-HA and FER-TCD-HA expressed in mesophyll protoplasts
(Supplementary Fig. 2b), but not FER-ECD-HA (Supplementary Fig. 2b), further
confirming the interaction between RopGEF14 and FER's intracellular kinase domain
(Supplementary Fig. 6c).

We next conducted a series of experiments to confirm that ROP6 is also a part of the
FER-RopGEF14 complex. Co-IP assays showed that ROP6 associated with
RopGEF14 (Fig. 6b). MBP-GEF14 recombinant proteins efficiently pulled down
ROP6-Flag expressed in protoplasts (Supplementary Fig 6d). Our co-IP assays further
detected the association of FER with ROP6 in mesophyll protoplasts co-expressing
FER-Flag and ROP-GFP (Fig. 6c). FER-Flag proteins were immunoprecipitated by
ROP6-GFP using anti-GFP trap. Complementary results were obtained using FER-
GFP to immunoprecipitate ROP6-Flag (Supplementary Fig. 6b). Taken together, these
data indicate that FER regulates ROP6 signaling by directly interacting with the
RopGEF14 /ROP6 complex.

Our genetic analysis further corroborates the physical interaction among FER,
RopGEF14, and ROP6. Previously, we showed that a constitutively active ROP6
mutant (*CA-rop6*) dramatically suppressed lobe formation and increased the width of
indentation necks in PCs [20]. PC phenotypes of the *CA-rop6 fer-4* double mutant was
identical to those observed in *CA-rop6* (Supplementary Fig. 7a), suggesting that FER
and ROP6 act in the same genetic pathway. Consistently with the reduction ROP6
signaling in the *fer-4* and *gef14-2* mutants (Fig. 4a-f, Fig. 5c-f), overexpression of
ROP6 partially restored the *fer-4* and largely restored the *gef14-2* PC phenotypes,
respectively (Fig. 6d-g). Furthermore, the PC indentation widths of *fer-5gef14-2* and
gef14-2rop6 double mutants were identical to those observed in *fer-5* and *gef14-2*
mutants (Supplementary Fig. 7b-e). These results demonstrate that FER binds de-
methylesterified pectin leading to the activation of the ROP6 signaling pathway
through RopGEF14 and promotes the indentation during PC morphogenesis.

Fig. 6 FER associates with RopGEF14 and ROP6 complex in regulating PC morphogenesis. **a**,

FER associates with RopGEF14 in transgenic plants. Proteins from 10-day-old *pGEF14::GFE14-*

4xMyc/gef14-2 or *gef14-2* seedlings were immunoprecipitated with α-Myc-Trap antibody and analyzed

with Western blot using α-FER antibody (Top). The expression of FER and GEF14-Myc in transgenic

plants are shown (Middle and Bottom). **b**, ROP6 associates with RopGEF14 in transgenic plants.

Proteins from 10-day-old *pGEF14::GFE14-4xMyc/gef14-2* or *gef14-2* seedlings were

immunoprecipitated with α-Myc-Trap antibody and analyzed with Western blot using α-ROP6

antibody (Top). The expression of ROP6 and GEF14-Myc in transgenic plants are shown (Middle and

Bottom). **c**, FER associates with ROP6 in protoplasts. Co-IP was carried out with an α-GFP-Trap

antibody (IP: α-GFP-Trap), and the proteins were analyzed by using Western blot with α-Flag

antibody. Top shows that FER-Flag coimmunoprecipitated with ROP6-GFP (IP: α-GFP-Trap; WB: α-

Flag). Middle and Bottom show the expression of FER-Flag and ROP6-GFP proteins (WB: α-Flag or

α-GFP for input control). **d and e**, ROP6 genetically acts downstream of FER in regulating PC

morphogenesis. PC morphogenesis was characterized by 2 DAG wild-type, *fer-4, ROP6-GFP/fer-4*

and ROP6-GFP seedlings. Overexpression of ROP6-GFP partially restored the PCs defects of the *fer-4*

mutant. Scale bar, 100 μ m. Asterisks indicate the significant difference (p≤0.05, Student's *t*-test)

between the *fer-4* and the *fer-4/ROP6-GFP* in above assays. **f and g**, ROP6 genetically acts

downstream of RopGEF14 in regulating PC morphogenesis. PC morphogenesis was characterized by

wild-type, *gef14-2, ROP6-GFP/gef14-2,* and *ROP6-GFP*. Overexpression of ROP6-GFP largely

restored the PCs defects of the *gef14-2* mutant. Scale bar, 50 μ m. Asterisks indicate the significant

difference (p≤0.05, Student's *t*-test) between the *gef14-2* and the *ROP6-GFP/gef14-2* in above assays.

DISCUSSION

Our comprehensive biochemical, genetic and cell biological data unequivocally

demonstrate that upon sensing cell wall pectin via its extracellular domain the cell

surface receptor kinase FER from the CrRLK1L subfamily directly activates ROP6 GTPase signaling and that this cell wall sensing/signaling pathway regulates PC shape formation in the *Arabidopsis* leaf epidermis. Therefore, FER is at least one of the long-sought cell wall sensors that couple a specific cell wall polysaccharide with intracellular signaling to regulate a particular cellular process. FER connects the cell wall properties to the control of cell expansion, likely by sensing dynamic changes in the cell wall composition and structure. Sensing the changes in pectin modification in the cell wall might be a common mechanism for the regulation of polar cell expansion. ANXUR1/2 and BUPS1/2 from the same family of FER control the cell wall integrity of elongating pollen tubes, which is also dependent on pectin de-methylesterification [48,69]. FER-based monitoring of cell wall integrity and dynamic might also provide a mechanism to coordinate the cell growth between neighboring cells in plant tissues. In line with this, *fer* mutations disrupt coordinated cell growth causing random cell expansion in the root tip [70] and severe defects in the interdigitated PC shape (Fig. 1a, b and Supplementary Fig. 1a, b). During interdigitated cell growth in PCs, the initial ROP2 activation might induce differential modifications of pectin between lobing and indenting sides (e.g, via secretion of PMEs, PMEIs, or pectinase), allowing the coordination between lobing and indenting sides via interactions between de-methylesterified pectin and FER.

By sensing specific wall components, FER and other CrRLK1L family members may directly monitor the cell wall integrity and activate compensatory pathways to maintain wall integrity. FER functions in plant defense response and is proposed to do so by monitoring the disruption of the cell wall caused by pathogen invasion [47], though direct evidence for this role is lacking. Above mentioned THE1 is involved in sensing or signaling disturbances in cell wall cellulose biosynthesis to activate lignin biosynthesis [30]. Interestingly, yeast cells regulate cell wall integrity using a cell surface receptor-Rho GTPase signaling system [71-73]. The cell surface receptor-Rho GTPase signaling systems appear to be a common mechanism for surveying and regulating cell wall integrity.

By binding structural cell wall components such as de-methylesterified pectin, FER, and other CrRLK1L family members might sense the mechanical signals of either internal or external origin. This is consistent with a role for FER in mechanical signal transduction in *Arabidopsis* [70]. Finally, our findings expand the list of FER ligands including RALF [40,41], suggesting that FER and likely its relatives may act as integrators of external signals. Clearly, CrRLK1L/RALF and LRX/RALF complexes are required for cell wall integrity maintenance in pollen tubes [7,48] and roots [39]. Notably, the apical wall of pollen tubes is predominantly composed of pectin, with highly methylesterified pectin deposited at the growing tip and de-methylesterified pectin confined to the shank[69]. Reducing the activity of PME results in pollen tube rupture, reminiscent of the defect in the CrRLK1L/RALF and LRX/RALF pathways [69,74]. PME activity requires an alkaline pH optimum, while de-methylestrification of pectin leads to the cell wall acidification [75,76]. It is intriguing to propose that FER and other CrRLK1L members coordinate the sensing of RALFs (as alkalizing peptides) and pectin methylesterification levels to control the homeostasis of apoplastic pH and

531 pectin status needed to maintain the cell wall integrity. How CrRLK1L members
532 dance with RALFs and pectin to modulate cell growth, morphogenesis, and integrity
533 maintenance is a fascinating question yet to be explored.   Hence our findings here
534 will propel a very fertile field of study on the sensing and signaling of cell wall
535 dynamic, integrity and mechanics in plants.
536
537 **METHODS**
538
Plant Materials and Growth Conditions. The *fer-4* (GABI_GK106A06), *fer-5*
(*Salk_ 029056c*) were ordered from ABRC and *fer-2* was obtained from Ueli
Grossniklaus (University of Zürich, Switzerland). The *pme3* and *PMEI1-OE* were
obtained from Vincenzo Lionetti (Sapienza University of Rome, Italy) and the
arad1arad2 mutant was obtained from J. Paul Knok (University of Leeds, UK). The
*GFP-MAP4*x*fer-4*, *GFP-TUB*x*PMEI1-OE*, *GFP-ROP6*x*fer-4*,*CA-rop6*x*fer-4, fer-*
*5*x*gef14-2,* and *GFP-ROP6*x*gef14-2* mutants were generated by genetic crosses and
confirmed by genotyping or Western blotting. *Arabidopsis* plants were grown in soil
(Sungro S16-281) in a growth room at 23 °C, 40% relative humidity, and 75 μE m^{-2}·s^{-}
^{1} light with a 12-h photoperiod for approximate 4 weeks before protoplast isolations.
To grow *Arabidopsis* seedlings, the seeds were surface sterilized with 50% (vol/vol)
bleach for 10 min, and then placed on the plates with 1/2 MS medium containing
0.5% sucrose, and 0.8% agar at pH 5.7. 2 to 6 days after germination (DAG)
cotyledons were used for pavement cells characterization.
553
Plasmid Construction and Generation of Transgenic Plants. Full-length and
truncated variants *FER*, *BIR1,* and *ROP6* were amplified by PCR from Col-0 cDNA
and cloned into a protoplast transient expression vector (obtained from Libo Shan &
Ping He, Texas A&M) or plant expression vector pGWB641. The *FER* promoter of
1.3 kb was amplified by PCR from Col-0 genomic DNA and introduced into
pGWB641 and pGWB605 binary vectors carrying FER-YFP and FER-GFP
respectively. The truncated FER variants were constructed by overlapping PCR
amplified from Col-0 cDNA and cloned into a protoplast compatible transient
expression vector or plant expression vector pGWB642. FER-ECD and FER-MALA
were amplified by PCR and cloned into proteins expression vector pDEST-HisMBP
(obtained from *Addgene*). Full length of RopGEF14 was amplified from Col-0
genomic DNA and cloned into pGWB516. The GEF14 promoter of 1.5 kb together
with GEF14 was amplified by PCR from Col-0 genomic DNA and introduced into
pGWB516 binary vectors. Full length of RopGEF14 was amplified from Col-0 cDNA
and cloned into pGWB506. All of the constructs were fully sequenced to verify
mutations in the gene coding and promoter region. Stable transgenic lines were
generated by using the standard *Agrobacterium tumefaciens*-mediated transformation
in the *fer-4 , gef14-2* mutants or Col-0.
572
Confocal analysis of *Arabidopsis* cotyledons PC shape. *Arabidopsis* cotyledons
were firstly incubated in PBS buffer with propidium iodide (PI) (2 mg/ml, 20
minutes). Then after washing with PBS buffer for three times, the samples were
observed under confocal microscopy (Leica SP5 Laser Scanning Confocal.). The PCs

images were taken at the mid-region of cotyledons. The widths of indentations were
measured using *LAS AF Lite* software. Each dataset was generated from the
measurement of at least 20 cells collected from 5 different cotyledons from 5
individual seedlings.
581
GFP detection by proteins gel blotting and confocal laser scanning microscopy.
Total proteins extraction and protein gel-blot analysis were performed as described
before with modifications[28]. Soluble proteins were prepared using an extraction buffer
(25 mM Tris-HCl, pH 7.5, 200 mM KCl, 5 mM $MgCl_2$, and 5 mM DTT, 0.1% NP40)
supplemented with a mixture of protease inhibitor (cOmplete Protease inhibitor
cocktail; Roche). After washing and extraction for 15 times, the pellet was used for
insoluble protein preparation. Insoluble proteins were extracted by boiling the pellet
for 10 minutes in the extraction buffer (50 mM Tris-HCl, pH 6.8, 10% glycerol,
0.05% bromphenol blue and 50 mM DTT, 4% SDS). Total GFP was detected using a
GFP polyclonal antibody (Santa Cruze, B-2, 1:1000 dilution) and subjected to
horseradish peroxidase-conjugated mouse secondary antibody, and developed with
enhanced chemiluminescence detection reagents (ThermoFish SuperSignal West Pico
Chemiluminescent Substrate, Cat #. 34080). Ten-day-old seedlings were used for
GFP detection. Whole cotyledons were directly mounted in PI (10 μ g/ml) or FM4-64
(5 μ g/ml) solution and observed with water objectives as described previously. GFP
only transgenic seedlings were used as a control. GFP fluorescence and PI were
excited simultaneously by a blue argon laser (10 mW, 488-nm blue excitation) and
535 nm for red excitation and detected at 515-530 nm for GFP and 600-617nm
wavelengths for PI in a Leica SP5 Laser Scanning Confocal. For plasmolysis,
cotyledons were incubated with PI for 20 minutes and washed with PBS buffer and
subjected to a treatment of a 0.8 M Mannitol solution for 5 minutes before
observation. Images were processed and arranged by ImageJ.
604
Protoplast preparation and transient expression. Protoplasts were prepared
according to the protocol described by Yoo et al[77].Maxiprep DNA for transient
expression was prepared using the Invitrogen PureLink Plasmid Maxiprep Kit. $2x10^5$
protoplasts were transfected with indicated FER-HA or truncated variants and
incubated at room temperature for 10 hours. The protoplasts were collected and stored
at -80 °C for further usage.
611
Coimmunoprecipitate (Co-IP) assay. 2×10^5 protoplasts transfected with
indicated plasmids were lysed with 0.5 mL of extraction buffer (10 mM Hepes at pH
7.5, 100 mM NaCl, 1 Mm EDTA, 10% (vol/vol) glycerol, 0.5% Triton X-100, and
protease inhibitor mixture from Roche). After vortexing vigorously for 30 s, the
samples were centrifuged at 12,470 $\times$ g for 10 min at 4 ° C. The supernatant was
incubated with α -GFP-Trap antibody for 2 h with gentle shaking. The beads were
collected and washed three times with washing buffer (10 mM Hepes at pH 7.5, 100
mM NaCl, 1 mM EDTA, 10% glycerol, and 0.1%Triton X-100) and once with 50
mM Tris • HCl at pH 7.5. The immunoprecipitated proteins were analyzed by
Western blot with α -GFP or α -FLAG antibody. For seedling Co-IP, approximate 1
g of 10-day seedlings were ground in liquid N2 and further ground in 0.5 mL of ice-

cold Co-IP buffer. Samples were centrifuged at 12,470 × g for 10 min at 4 °C. The

resulting supernatant was used to perform the Co-IP assay with the same procedures

as protoplast Co-IP assay with α-Myc-Trap or α-FER (Generated and purified from

Rabbit)[42] antibodies.

In vitro **pull-down assay**. 2×10^5 protoplasts were lysed with 1 mL extraction buffer

(20 mM Tris pH 8.2, NaCl 150 mM, 0.5% Triton X-100, 0.5 mM $CaCl_2$ and protease

inhibitor mixture from Roche)[78]. After vortexing vigorously for 30 seconds, the

samples were centrifuged at 12000 × g for 10 minutes at 4 °C. 20 µl of supernatant

was used as input control and the remainder of the supernatant was incubated with the

indicated amounts of pectin or PGA for 2 -4 hours at 4 °C with gentle shaking. Pectin

or PGA were pelleted and collected by centrifuging at 6000 x g for 10 minutes at 4 °C

and washed three times with washing buffer (20 mM Tris pH 8.2, NaCl 150 mM, 0.5

mM $CaCl_2$ and protease inhibitor mixture from Roche). 20 µl 1x SDS loading buffer

was added to the pull-down pellet and the released proteins were analyzed by Western

blot with a α-HA antibody.

Expression of MBP fusion proteins and affinity purification were performed as

standard protocol. The protein concentration was determined with NanoDrop ND-

1000 spectrophotometer and confirmed by the Bio-Rad Quick Start Bradford Dye

Reagent. 200 ng *E.coli* (Rosetta DE3) produced recombinant proteins were incubated

with 200 µl extraction buffer for 30 minutes and then centrifuged at 6000 x g for 10

minutes at 4 °C, 20 µl of supernatant was used as input control and the remainder of

the supernatant was incubated with the indicated amounts of pectin or PGA for 2

hours at room temperature with gentle shaking. Pull-down was carried out as

described above, and the pull-down proteins were determined by western blot with α-

His or α-MBP antibodies.

Immunolocalization of pectin and MT in PCs. Two DAG cotyledons or the

epidermal layer from 3 weeks old plants the third pair leaves were fixed, frozen,

shattered and permeabilized as described by Burn, J.E [34]. For pectin

immunolocalization, samples were incubated with the primary antibodies JIM5 (1:

100, PlantProbes, University of Leeds, UK) and LM20 (1: 100, PlantProbes,

University of Leeds, UK). Samples were then transferred to a buffer with the

secondary antibody (FITC-conjugated anti-Rat IgG at 1: 100 dilution, Sigma). For

MT immunolocalization, following incubation with a monoclonal anti-α-tubulin-FITC

antibody produced in mouse (Sigma) and washed three times after incubation, the

samples were observed under Leica SP5 Laser Scanning Confocal. For quantification

of pectin in lobes and indentations, the fluorescence intensity was analyzed by

ImageJ. The mean ratio of indention/lobe was calculated to indicate the differential

distribution of de-esterified pectin and esterified pectin in lobe and indentation

regions of PC.

Quantitative analysis of cortical microtubule orientation. The *fer-4* mutant was

crossed to the *GFP-MAP4* line and the *PMEI1-OE* was crossed to *GFP-TUB* line to

enable the visualization of MTs. Images of PC MTs were generated using a Leica SP5

Laser Scanning Confocal. FibrilTool, which is an ImageJ plug-in to quantify fibrillar

structures, was used to analyze the average anisotropy of MT[79]. The indentation
regions were selected with the Polygon tool. With regard to the anisotropy score, "0"
indicates no order (purely isotropic arrays) and "1" indicates perfectly ordered (purely
anisotropic arrays). Each data set was from the measurement of at least 20 cells
collected from 3 different cotyledons from 3 individual seedlings.

Measurement of ROP6 activity. ROP activity measurement was performed as
described by Xu[68]. 10-day-old 1/2 MS grown seedlings or protoplasts isolated from 4-
week-old plants were used in this assay. Total ROP6 and activated ROP6 proteins that
pull down by MBP-RIC1 (GTP-bound ROP6) were detected by Western blots using
ROP6 or GFP and horseradish peroxidase-conjugated rabbit antibodies. The proteins
levels were determined by ImageJ.

RT-PCR Analysis. Total RNA was isolated from wild-type or mutant leaves or
seedlings with TRIzol Reagent (Invitrogen). First-strand cDNA was synthesized from
1 μg of total RNA with reverse transcriptase. The RT-PCR analysis was carried out
by using the synthesized cDNA as templates, and Actin 8 was used as a control gene.
The RT-PCR primer sequences are listed below.

Primers for construct cloning and genotyping

Constructs	Primers	Sequences
FER-TCD (overlapping PCR)	P1S	GGGGACAAGTTTGTACAAAAAAGCAGGCTGCATGAAGA
	P1A	TAGGATTTTCTCTGTTGGAGAGTAATCAGCAG
	P2S	CAGAGAAAATCCTACCGGAATCAGCTCCGTCTAA
	P2A	ATTCTCAAGAGCAGTACTGTTATCAATAGTAACAGAG
	P3S	ACTGCTCTTGAGAATTATGATTCTCTTCTTAATGGAGTG
	P3A	GGGGACCACTTTGTACAAGAAAGCTGGGTCACGTCCCT
FER full length (BamHI/StuI)	F	CG<u>GGATCC</u>ATGAAGATCACAGAGGG
	R	GA<u>AGGCCT</u>ACGTCCCTTTGGATTC
FERΔMALA	F	CATTGAAGTAACCTCCTCTGTTGGAGAGTA
	R	TACTCTCCAACAGAGGAGGTTACTTCAATG
FERΔMALB	F	AACTGGGTTAGGATGCTCAAGAGCAGTACT
	R	AGTACTGCTCTTGAGCATCCTAACCCAGTT
FER-ECD (BamHI/StuI)	F	CG<u>GGATCC</u>ATGAAGATCACAGAGGG
	R	GA<u>AGGCCT</u>GGCCAGAACAACTGC
fer mutants genotyping	*fer*4L	CGGATCCATGAAGATCACAGAGGGACGATTC
	*fer*4R	CGCAGATCTAGCACCAAACACACAAAACCC
	*fer*4LB	GTGGATTGATGTGATATCTCC
	*fer*5L	CGGATCCATGGCTTACCGCAGACGTAAGCGTGG
	*fer*5R	CGAATTCACGTCCCTTTGGATTCATCATCTG
	*fer*5LB	GCGTGGACCGCTTGCTGCAACT
FER full length (pGW641)	FER-F	GGGGACAAGTTTGTACAAAAAAGCAGGCTGCATGAAGA
	FER-R	GGGGACCACTTTGTACAAGAAAGCTGGGTTACGTCCCT

FER ECD and MALA (pDEST-His-MBP)	F-ECD	
	R-ECD	
	MALA-F	gtGGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGGCT
	MALA-F	gtGGGGACCACTTTGTACAAGAAAGCTGGGTCCTAGGCC
	MALA-R	gtGGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGGCT
	MALA-R	gtGGGGACCACTTTGTACAAGAAAGCTGGGTCCTAAATC
FER promoter (HindIII/XbaI)	pFER-F	GGTAAGCTTCGATTTAAGCGAGTTGG
	pFER-R	GCCTCTAGACGATCAAGAGCACTTCTCCGGG
GEF14-Full length (pGWB506)	F-FL	5'GGGGACAAGTTTGTACAAAAAAGCAGGCTGCATGAT
	R-FL	5'GGGGACCACTTTGTACAAGAAAGCTGGGTCAGGAGA
promoter+Full length (pGWB516)	F-pGEF14-GEF14	
	R-pGEF14-GEF14	5' GGGGACAAGTTTGTACAAAAAAGCAGGCTGCTAAT
		5'GGGGACCACTTTGTACAAGAAAGCTGGGTCAGGAGA
RT-PCR	GEF14-F	
	GEF14-R	5' ATGATGCTGATGAGAAGAAGGT 3'
		5' GCACGCATCGAACTAGGA 3'
	GEF1-F	5' ATGGGGAGCTTATCTTCTGAGG 3'
	GEF1-R	5' ATCTCTTTCCGGCGTCACTCCCG 3'
	GEF6-F	5' ATGGAGGATAATAGCTGTATCGG 3'
	GEF6-R	5' ACCCCGGAGATAATTGGCCAATGCT 3'
	GEF7-F	5' ATGGATGGTTCGTCGGAAAATTTGC 3'
	GEF7-R	5' AATCCCAGGATCAAGGTTCGATAC 3'

REFERENCES:

Wolf, S., Hematy, K. & Hofte, H. Growth control and cell wall signaling in plants. *Annual review of plant biology* **63**, 381-407, doi:10.1146/annurev-arplant-042811-105449 (2012).

Cosgrove, D. J. Plant cell wall extensibility: connecting plant cell growth with cell wall structure, mechanics, and the action of wall-modifying enzymes. *Journal of experimental botany* **67**, 463-476, doi:10.1093/jxb/erv511 (2016).

Hofte, H. The yin and yang of cell wall integrity control: brassinosteroid and FERONIA signaling. *Plant & cell physiology* **56**, 224-231, doi:10.1093/pcp/pcu182 (2015).

Peaucelle, A. *et al.* The transcription factor BELLRINGER modulates phyllotaxis by regulating the expression of a pectin methylesterase in Arabidopsis. *Development* **138**, 4733-4741, doi:10.1242/dev.072496 (2011).

Wolf, S. Plant cell wall signalling and receptor-like kinases. *The Biochemical journal* **474**, 471-492, doi:10.1042/BCJ20160238 (2017).

Seifert, G. J. & Blaukopf, C. Irritable walls: the plant extracellular matrix and signaling. *Plant physiology* **153**, 467-478, doi:10.1104/pp.110.153940 (2010).

Majda, M. *et al.* Mechanochemical Polarization of Contiguous Cell Walls Shapes Plant Pavement Cells. *Developmental cell* **43**, 290-304 e294, doi:10.1016/j.devcel.2017.10.017 (2017).

8   Manfield, I. W. *et al.* Novel cell wall architecture of isoxaben-habituated Arabidopsis
suspension-cultured cells: global transcript profiling and cellular analysis. *The Plant journal :*
for cell and molecular biology **40**, 260-275, doi:10.1111/j.1365-313X.2004.02208.x (2004).
9   Moscatiello, R., Mariani, P., Sanders, D. & Maathuis, F. J. Transcriptional analysis of calcium-
dependent and calcium-independent signalling pathways induced by oligogalacturonides.
Journal of experimental botany **57**, 2847-2865, doi:10.1093/jxb/erl043 (2006).
10   Cano-Delgado, A., Penfield, S., Smith, C., Catley, M. & Bevan, M. Reduced cellulose synthesis
invokes lignification and defense responses in Arabidopsis thaliana. *The Plant journal : for cell*
and molecular biology **34**, 351-362 (2003).
11   Denness, L. *et al.* Cell wall damage-induced lignin biosynthesis is regulated by a reactive
oxygen species- and jasmonic acid-dependent process in Arabidopsis. *Plant physiology* **156**,
1364-1374, doi:10.1104/pp.111.175737 (2011).
12   Desprez, T. *et al.* Resistance against herbicide isoxaben and cellulose deficiency caused by
distinct mutations in same cellulose synthase isoform CESA6. *Plant physiology* **128**, 482-490,
doi:10.1104/pp.010822 (2002).
13   Burton, R. A. *et al.* Virus-induced silencing of a plant cellulose synthase gene. *The Plant cell*
12, 691-706 (2000).
14   His, I., Driouich, A., Nicol, F., Jauneau, A. & Hofte, H. Altered pectin composition in primary
cell walls of korrigan, a dwarf mutant of Arabidopsis deficient in a membrane-bound endo-
1,4-beta-glucanase. *Planta* **212**, 348-358 (2001).
15   Ellis, C., Karafyllidis, I. & Turner, J. G. Constitutive activation of jasmonate signaling in an
Arabidopsis mutant correlates with enhanced resistance to Erysiphe cichoracearum,
Pseudomonas syringae, and Myzus persicae. *Molecular plant-microbe interactions : MPMI*
15, 1025-1030, doi:10.1094/MPMI.2002.15.10.1025 (2002).
16   Ellis, C., Karafyllidis, I., Wasternack, C. & Turner, J. G. The Arabidopsis mutant cev1 links cell
wall signaling to jasmonate and ethylene responses. *The Plant cell* **14**, 1557-1566 (2002).
17   Somerville, C. *et al.* Toward a systems approach to understanding plant cell walls. *Science*
306, 2206-2211, doi:10.1126/science.1102765 (2004).
18   Fukuda, H. & Yang, Z. Editorial overview: Cell biology: From signals to cell shape and function.
Current opinion in plant biology **28**, iv-vi, doi:10.1016/j.pbi.2015.11.001 (2015).
19   Yang, Z. Cell polarity signaling in Arabidopsis. *Annual review of cell and developmental*
biology **24**, 551-575, doi:10.1146/annurev.cellbio.23.090506.123233 (2008).
20   Fu, Y., Gu, Y., Zheng, Z., Wasteneys, G. & Yang, Z. Arabidopsis interdigitating cell growth
requires two antagonistic pathways with opposing action on cell morphogenesis. *Cell* **120**,
687-700, doi:10.1016/j.cell.2004.12.026 (2005).
21   Fu, Y., Xu, T., Zhu, L., Wen, M. & Yang, Z. A ROP GTPase signaling pathway controls cortical
microtubule ordering and cell expansion in Arabidopsis. *Current biology : CB* **19**, 1827-1832,
doi:10.1016/j.cub.2009.08.052 (2009).
22   Sampathkumar, A. *et al.* Subcellular and supracellular mechanical stress prescribes
cytoskeleton behavior in Arabidopsis cotyledon pavement cells. *eLife* **3**, e01967,
doi:10.7554/eLife.01967 (2014).
23   Wolf, S. *et al.* A receptor-like protein mediates the response to pectin modification by
activating brassinosteroid signaling. *Proceedings of the National Academy of Sciences of the*
United States of America **111**, 15261-15266, doi:10.1073/pnas.1322979111 (2014).
24   Mecchia, M. A. *et al.* RALF4/19 peptides interact with LRX proteins to control pollen tube
growth in Arabidopsis. *Science* **358**, 1600-1603, doi:10.1126/science.aao5467 (2017).

25   Bai, L. *et al.* Plasma membrane-associated proline-rich extensin-like receptor kinase 4, a
novel regulator of Ca signalling, is required for abscisic acid responses in Arabidopsis
thaliana. *The Plant journal : for cell and molecular biology* **60**, 314-327, doi:10.1111/j.1365-
313X.2009.03956.x (2009).

26   Beneteau, J. *et al.* Binding properties of the N-acetylglucosamine and high-mannose N-glycan
PP2-A1 phloem lectin in Arabidopsis. *Plant physiology* **153**, 1345-1361,
doi:10.1104/pp.110.153882 (2010).

27   Bouwmeester, K. & Govers, F. Arabidopsis L-type lectin receptor kinases: phylogeny,
classification, and expression profiles. *Journal of experimental botany* **60**, 4383-4396,
doi:10.1093/jxb/erp277 (2009).

28   He, Z. H., Fujiki, M. & Kohorn, B. D. A cell wall-associated, receptor-like protein kinase. *The*
Journal of biological chemistry **271**, 19789-19793 (1996).

29   Hematy, K. & Hofte, H. Novel receptor kinases involved in growth regulation. *Current opinion*
in plant biology **11**, 321-328, doi:10.1016/j.pbi.2008.02.008 (2008).

30   Hematy, K. *et al.* A receptor-like kinase mediates the response of Arabidopsis cells to the
inhibition of cellulose synthesis. *Current biology : CB* **17**, 922-931,
doi:10.1016/j.cub.2007.05.018 (2007).

31   Trudel, J., Grenier, J., Potvin, C. & Asselin, A. Several thaumatin-like proteins bind to beta-1,3-
glucans. *Plant physiology* **118**, 1431-1438 (1998).

32   Wang, X., Zafian, P., Choudhary, M. & Lawton, M. The PR5K receptor protein kinase from
Arabidopsis thaliana is structurally related to a family of plant defense proteins. *Proceedings*
of the National Academy of Sciences of the United States of America **93**, 2598-2602 (1996).

33   Xu, S. L., Rahman, A., Baskin, T. I. & Kieber, J. J. Two leucine-rich repeat receptor kinases
mediate signaling, linking cell wall biosynthesis and ACC synthase in Arabidopsis. *The Plant*
cell **20**, 3065-3079, doi:10.1105/tpc.108.063354 (2008).

34   Van der Does, D. *et al.* The Arabidopsis leucine-rich repeat receptor kinase MIK2/LRR-KISS
connects cell wall integrity sensing, root growth and response to abiotic and biotic stresses.
PLoS genetics **13**, e1006832, doi:10.1371/journal.pgen.1006832 (2017).

35   Shiu, S. H. & Bleecker, A. B. Plant receptor-like kinase gene family: diversity, function, and
signaling. *Science's STKE : signal transduction knowledge environment* **2001**, re22,
doi:10.1126/stke.2001.113.re22 (2001).

36   Escobar-Restrepo, J. M. *et al.* The FERONIA receptor-like kinase mediates male-female
interactions during pollen tube reception. *Science* **317**, 656-660,
doi:10.1126/science.1143562 (2007).

37   Li, C. *et al.* Glycosylphosphatidylinositol-anchored proteins as chaperones and co-receptors
for FERONIA receptor kinase signaling in Arabidopsis. *eLife* **4**, doi:10.7554/eLife.06587
(2015).

38   Deslauriers, S. D. & Larsen, P. B. FERONIA is a key modulator of brassinosteroid and ethylene
responsiveness in Arabidopsis hypocotyls. *Molecular plant* **3**, 626-640,
doi:10.1093/mp/ssq015 (2010).

39   Duan, Q., Kita, D., Li, C., Cheung, A. Y. & Wu, H. M. FERONIA receptor-like kinase regulates
RHO GTPase signaling of root hair development. *Proceedings of the National Academy of*
Sciences of the United States of America **107**, 17821-17826, doi:10.1073/pnas.1005366107
(2010).

799 40 Haruta, M., Sabat, G., Stecker, K., Minkoff, B. B. & Sussman, M. R. A peptide hormone and its
800   receptor protein kinase regulate plant cell expansion. *Science* **343**, 408-411,
801   doi:10.1126/science.1244454 (2014).

802 41 Stegmann, M. *et al.* The receptor kinase FER is a RALF-regulated scaffold controlling plant
803   immune signaling. *Science* **355**, 287-289, doi:10.1126/science.aal2541 (2017).

804 42 Chen, J. *et al.* FERONIA interacts with ABI2-type phosphatases to facilitate signaling cross-talk
805   between abscisic acid and RALF peptide in Arabidopsis. *Proceedings of the National Academy*
806   *of Sciences of the United States of America* **113**, E5519-5527, doi:10.1073/pnas.1608449113
807   (2016).

808 43 Duan, Q. *et al.* Reactive oxygen species mediate pollen tube rupture to release sperm for
809   fertilization in Arabidopsis. *Nature communications* **5**, 3129, doi:10.1038/ncomms4129
810   (2014).

811 44 Guo, H. *et al.* Three related receptor-like kinases are required for optimal cell elongation in
812   Arabidopsis thaliana. *Proceedings of the National Academy of Sciences of the United States of*
813   *America* **106**, 7648-7653, doi:10.1073/pnas.0812346106 (2009).

814 45 Huck, N., Moore, J. M., Federer, M. & Grossniklaus, U. The Arabidopsis mutant feronia
815   disrupts the female gametophytic control of pollen tube reception. *Development* **130**, 2149-
816   2159 (2003).

817 46 Yu, F. *et al.* FERONIA receptor kinase pathway suppresses abscisic acid signaling in
818   Arabidopsis by activating ABI2 phosphatase. *Proceedings of the National Academy of*
819   *Sciences of the United States of America* **109**, 14693-14698, doi:10.1073/pnas.1212547109
820   (2012).

821 47 Kessler, S. A. *et al.* Conserved molecular components for pollen tube reception and fungal
822   invasion. *Science* **330**, 968-971, doi:10.1126/science.1195211 (2010).

823 48 Ge, Z. *et al.* Arabidopsis pollen tube integrity and sperm release are regulated by RALF-
824   mediated signaling. *Science*, doi:10.1126/science.aao3642 (2017).

825 49 Lindner, H. *et al.* TURAN and EVAN mediate pollen tube reception in Arabidopsis Synergids
826   through protein glycosylation. *PLoS biology* **13**, e1002139, doi:10.1371/journal.pbio.1002139
827   (2015).

828 50 Boisson-Dernier, A. *et al.* Disruption of the pollen-expressed FERONIA homologs ANXUR1 and
829   ANXUR2 triggers pollen tube discharge. *Development* **136**, 3279-3288,
830   doi:10.1242/dev.040071 (2009).

831 51 Nam, K. H. & Li, J. BRI1/BAK1, a receptor kinase pair mediating brassinosteroid signaling. *Cell*
832   **110**, 203-212 (2002).

833 52 Hou, X. *et al.* Involvement of a cell wall-associated kinase, WAKL4, in Arabidopsis mineral
834   responses. *Plant physiology* **139**, 1704-1716, doi:10.1104/pp.105.066910 (2005).

835 53 Giddings, T. H., Jr. & Staehelin, L. A. Spatial relationship between microtubules and plasma-
836   membrane rosettes during the deposition of primary wall microfibrils in Closterium sp.
837   *Planta* **173**, 22-30, doi:10.1007/BF00394482 (1988).

838 54 Burn, J. E. *et al.* The cellulose-deficient Arabidopsis mutant rsw3 is defective in a gene
839   encoding a putative glucosidase II, an enzyme processing N-glycans during ER quality control.
840   *The Plant journal : for cell and molecular biology* **32**, 949-960 (2002).

841 55 Braybrook, S. A., Hofte, H. & Peaucelle, A. Probing the mechanical contributions of the pectin
842   matrix: insights for cell growth. *Plant signaling & behavior* **7**, 1037-1041,
843   doi:10.4161/psb.20768 (2012).

56 Peaucelle, A. *et al.* Pectin-induced changes in cell wall mechanics underlie organ initiation in
Arabidopsis. *Current biology : CB* **21**, 1720-1726, doi:10.1016/j.cub.2011.08.057 (2011).
57 Peaucelle, A., Wightman, R. & Hofte, H. The Control of Growth Symmetry Breaking in the
Arabidopsis Hypocotyl. *Current biology : CB* **25**, 1746-1752, doi:10.1016/j.cub.2015.05.022
(2015).
58 Pelletier, S. *et al.* A role for pectin de-methylesterification in a developmentally regulated
growth acceleration in dark-grown Arabidopsis hypocotyls. *The New phytologist* **188**, 726-
739, doi:10.1111/j.1469-8137.2010.03409.x (2010).
59 Wolf, S. & Greiner, S. Growth control by cell wall pectins. *Protoplasma* **249 Suppl 2**, S169-
175, doi:10.1007/s00709-011-0371-5 (2012).
60 Wolf, S., Mouille, G. & Pelloux, J. Homogalacturonan methyl-esterification and plant
development. *Molecular plant* **2**, 851-860, doi:10.1093/mp/ssp066 (2009).
61 Atmodjo, M. A., Hao, Z. & Mohnen, D. Evolving views of pectin biosynthesis. *Annu Rev Plant*
Biol **64**, 747-779, doi:10.1146/annurev-arplant-042811-105534 (2013).
62 Liners, F., Thibault, J. F. & Van Cutsem, P. Influence of the degree of polymerization of
oligogalacturonates and of esterification pattern of pectin on their recognition by
monoclonal antibodies. *Plant physiology* **99**, 1099-1104 (1992).
63 Xiao, C., Somerville, C. & Anderson, C. T. POLYGALACTURONASE INVOLVED IN EXPANSION1
functions in cell elongation and flower development in Arabidopsis. *Plant Cell* **26**, 1018-1035,
doi:10.1105/tpc.114.123968 (2014).
64 Fu, Y., Li, H. & Yang, Z. The ROP2 GTPase controls the formation of cortical fine F-actin and
the early phase of directional cell expansion during Arabidopsis organogenesis. *The Plant cell*
14, 777-794 (2002).
65 Wang, M. *et al.* A comparative genome analysis of PME and PMEI families reveals the
evolution of pectin metabolism in plant cell walls. *PloS one* **8**, e72082,
doi:10.1371/journal.pone.0072082 (2013).
66 Guenin, S. *et al.* Identification of pectin methylesterase 3 as a basic pectin methylesterase
isoform involved in adventitious rooting in Arabidopsis thaliana. *The New phytologist* **192**,
114-126, doi:10.1111/j.1469-8137.2011.03797.x (2011).
67 Lionetti, V. *et al.* Overexpression of pectin methylesterase inhibitors in Arabidopsis restricts
fungal infection by Botrytis cinerea. *Plant physiology* **143**, 1871-1880,
doi:10.1104/pp.106.090803 (2007).
68 Xu, T. *et al.* Cell surface- and rho GTPase-based auxin signaling controls cellular
interdigitation in Arabidopsis. *Cell* **143**, 99-110, doi:10.1016/j.cell.2010.09.003 (2010).
69 Bosch, M. & Hepler, P. K. Pectin methylesterases and pectin dynamics in pollen tubes. *The*
Plant cell **17**, 3219-3226, doi:10.1105/tpc.105.037473 (2005).
70 Shih, H. W., Miller, N. D., Dai, C., Spalding, E. P. & Monshausen, G. B. The receptor-like kinase
FERONIA is required for mechanical signal transduction in Arabidopsis seedlings. *Current*
biology : CB **24**, 1887-1892, doi:10.1016/j.cub.2014.06.064 (2014).
71 Cheung, A. Y. & Wu, H. M. THESEUS 1, FERONIA and relatives: a family of cell wall-sensing
receptor kinases? *Current opinion in plant biology* **14**, 632-641,
doi:10.1016/j.pbi.2011.09.001 (2011).
72 Hamann, T. The plant cell wall integrity maintenance mechanism--a case study of a cell wall
plasma membrane signaling network. *Phytochemistry* **112**, 100-109,
doi:10.1016/j.phytochem.2014.09.019 (2015).

73 Hamann, T. The plant cell wall integrity maintenance mechanism-concepts for organization
and mode of action. *Plant & cell physiology* **56**, 215-223, doi:10.1093/pcp/pcu164 (2015).
74 Jiang, L. *et al.* VANGUARD1 encodes a pectin methylesterase that enhances pollen tube
growth in the Arabidopsis style and transmitting tract. *The Plant cell* **17**, 584-596,
doi:10.1105/tpc.104.027631 (2005).
75 Catoire, L., Pierron, M., Morvan, C., du Penhoat, C. H. & Goldberg, R. Investigation of the
action patterns of pectinmethylesterase isoforms through kinetic analyses and NMR
spectroscopy. Implications In cell wall expansion. *The Journal of biological chemistry* **273**,
33150-33156 (1998).
76 Goldberg, R. *et al.* Control of Mung bean pectinmethylesterase isoform activities. Influence
of pH and carboxyl group distribution along the pectic chains. *The Journal of biological*
chemistry **276**, 8841-8847, doi:10.1074/jbc.M001791200 (2001).
77 Yoo, S. D., Cho, Y. H. & Sheen, J. Arabidopsis mesophyll protoplasts: a versatile cell system for
transient gene expression analysis. *Nature protocols* **2**, 1565-1572,
doi:10.1038/nprot.2007.199 (2007).
78 Wagner, T. A. & Kohorn, B. D. Wall-associated kinases are expressed throughout plant
development and are required for cell expansion. *The Plant cell* **13**, 303-318 (2001).
79 Boudaoud, A. *et al.* FibrilTool, an ImageJ plug-in to quantify fibrillar structures in raw
microscopy images. *Nature protocols* **9**, 457-463, doi:10.1038/nprot.2014.024 (2014).
908

909 **ACKNOWLEDGEMENTS**
910

We are grateful to Ueli Grossniklaus (University of Zürich, Switzerland) for *fer-2*
seeds, to Vincenzo Lionetti (Sapienza University of Rome, Italy) for *pme3* and
PMEI1-OE seeds and to J. Paul Knok (University of Leeds, UK) for *arad1 arad2*
seeds. We gratefully acknowledge Eugene Nothnagel (University of California,
Riverside, USA) for helpful discussions. We thank members of the Yang
laboratory for their technical assistance and helpful discussions. This work is
supported by a grant from the U.S. National Institute of General Medical Sciences
to Z.Y. (GM081451) and by Fujian Agriculture and Forestry University.   C. T. A.
is supported by the Center for Lignocellulose Structure and Formation, an Energy
Frontier Research Center funded by the U.S. Department of Energy, Office of
Science, Basic Energy Sciences (award no. DE–SC0001090).

Supplemental Data

Supplemental Figure1. FERONIA is a cell wall-associated protein required for PC
morphogenesis. a, Diagram of FER's predicted domains and T-DNA insertion sites of 3 alleles of *fer*
mutants (*fer-2*, *fer-4*, *fer-5*). **b**, A diagram depicting how the neck widths were measured. A dashed line
segment was drawn between two indentations region, and the distance was decided as the indention
width. **c**, Pavement cell phenotypes of the wild-type and *fer-2*, *fer-4* and *fer-5* mutants. Scale bars, 100
μm. 2 DAG seedlings cotyledon pavement cells were characterized. **d**, The degree of pavement cell
interdigitation was quantified by determining the average indentation widths (Indentation width/ μm).
The stars indicate the average indentation widths were significantly different (p≤0.05) between wild-
type and the *fer* mutants. All data are represented as mean ± SE. **e**, The morphologies of 4 weeks old
soil-grown wild-type, fer-4, and fer-4 complementation line plants. **f**, FER can be extracted by 1%
SDS, 50 mM DTT from the cell fraction. The FER-GFP proteins were determined by Western blot
with a GFP antibody. **g**, The cell wall localization of FER-GFP. (Top panels) GFP fluorescence was
along with the cell surface of PCs before plasmolysis (-) indicated by overlapping signal with PI
(Merge). During plasmolysis (+) GFP fluorescence localized with the cytoplasm (Bottom left panel). A
small portion of the FER-GFP signal retreated with the cytoplasm and majority of the signal remained
on the cell surface (Bottom right panel). Arrows indicate the regressed plasma membrane. Scale bar, 5
μm

Supplemental Figure 2. FER associates with pectin and cell wall with the MALA domain. a, FER

failed to be pulled down by cellulose and xylan *in vitro*. FER-HA was expressed in *Arabidopsis*

protoplasts. Pull-down (PD) was carried out with cellulose and xylan with indicated concentration,

respectively. The FER-HA proteins were determined by Western blot with α-HA antibody. *Top* shows

that FER-HA was not pulled-down by cellulose or xylan (PD: cellulose/xylan; WB: α-HA). *Bottom*

shows the expression of FER-HA proteins (WB: α-HA as input control). **b**, Diagram of the full length

and specific truncated FER-HA constructs. The FER protein contains a signal peptide domain (SP), a

malectin-like extracellular domain A (MALA), a malectin-like extracellular domain B (MALB), an

extracellular juxta-membrane domain (exJM), a transmembrane domain (TM), a juxta-membrane

domain (JM) and a cytosolic kinase domain (Kinase). Numbers are the specific amino acid delineations

between domains. FER-ECD-HA, construct that with the extracellular domain (ECD, includes SP,

MALA, MALB, and exJM) of FER fuses with HA tag, the transmembrane domain and cytosolic

domain (JM and Kinase domain) were deleted. The FER-TCD-HA construct contains FER signal

peptide, a transmembrane domain and cytosolic domain (TCD, includes SP, TM, JM, and Kinase

domain), with the extracellular domain (ECD) deleted and fused with HA tag at the C-terminus. The

FER△MALA-HA construct has the FER MALA domain deleted and an HA tag at the C terminal. The

FER△MALB-HA construct has the FER MALB domain deleted and an HA tag at the C-terminus. **c**,

MALA domain directly binds to pectin *in vitro*. His-MBP-FER-MALA was purified from *E.coli*. PD

was carried out as above and the proteins were determined by Western blot with α-His antibody. His-

MBP-FER-MALA pull-down by pectin (*Top*). The input His-MBP-FER-ECD proteins (*Bottom*). **d**,

Identification of YFP-FER-TCD proteins by Western blot within different fractions. Total soluble (S)

and insoluble (P) proteins were prepared from seedlings of *35S::YFP-FER-TCD* transgenic plants and

analyzed with α-GFP antibody by Western blot. **e**, Deletion of extracellular domain results in loss of

cell wall binding of FER. Determination of the subcellular localization of YFP-FER-TCD by

plasmolysis assay with (+) or without (-) a 0.8 M mannitol solution. Arrows indicated the plasma

membrane region. **f**, MALA domain is required for FER cell wall binding. Determination of the

subcellular localization of YFP- FERΔMALA by plasmolysis assay with (+) or without (-) a 0.8 M

mannitol solution as was described before. Arrows indicated the plasma membrane region.

Supplemental Figure 3. FER preferentially binds to PGA than pectin. a, FER MALA domain was
associated with PGA and pectin *in vitro*. Pull-down was carried out with PGA and pectin as described
previously. (*Top*) His-MBP-FER-MALA proteins produced from *E.coli* were pull-down by PGA or
pectin. (*Bottom)* The His-MBP-FER-MALA proteins were determined by WB as input control. **b**, HG
crosslink formation is required for FER-PGA association. An interaction between FER-HA and PGA
was determined in different ionic solutions of concentrations as indicated. The pull-down was
performed as described previously. *Top* panel shows that FER-HA pull-down by PGA (PD: PGA; WB:
α-HA). *Bottom* panel shows the expression of FER-HA proteins (WB: α-HA for input control).
979

Supplemental Figure 4. FER mediates PCs morphogenesis via ROP6 signal pathway. a and b,

Activation of ROP6 in the wild-type (*GFP-ROP6*) and the *fer-4* mutant (*fer-4xGFP-ROP6*) were

analyzed by pull-down assay and determined by GFP antibody (a). The relative active of ROP6 level

was quantified (b). Protoplasts isolated from 4 weeks old adult plants were used in this assay. **c,**

Cortical microtubule orientation defects of *pme3* and *PMEI1-OE* mutants. Wild-type PCs show highly

ordered transverse cortical microtubules (with high anisotropy) in the indentation region, while

microtubule arrays in the *pme3* and *PMEI1-OE* mutant were mostly present random orientations (Top

panel). Magnified the PC indentation regions (Bottom panel). The microtubules were visualized by

immunostaining with an anti-tubulin antibody. Arrows indicate the indentation region. Scale bar, 10

μm. **d,** The histogram shows that the degree of cortical microtubule anisotropy in wild-type and the

pme3 and *PMEI1-OE* mutants. Anisotropy was calculated using ImageJ. Data are mean degrees from

10 independent cells ±SE. Asterisks indicate the significant difference ($p \leq 0.05$, Student's t-test)

between the wild type and the mutants in above assays.

Supplementary Figure 5. RopGEF14 is required for ROP GTPase activation in regulating PC
morphogenesis. a, The expressions of *RopGEF1, 6, 7* and *14* in cotyledon were determined by RT-
PCR**.** Actin 8 was included as an expression control. **b**, PC phenotypes of wild types (Col-0), the *gef1*
and *gef7* mutants. The degree of 2 DAG PC interdigitation was determined by the width of indentation.
c and d, PC phenotypes of wild types (Col-0), and the *gef6* mutants (c). The degree of 2 DAG PC
interdigitation was determined by the width of indentation (d). **e**, RT-PCR analysis of *GEF1, GEF6*,
GEF7 and *Actin8* (control) in wild-type (Col-0) and the *gef1, gef6* and *gef7* T-DNA insertion mutants.
f, T-DNA insertion sites in the *gef14* mutants with exons (red boxes). **g**, RT-PCR analysis of *GEF14,*
and Actin8 (control) in wild-type (Col-0) the *gef14-2* and *gef14-3* T-DNA insertion mutants. **h and i**,
PC phenotypes of wild types (Col-0), the *gef14-2* and *gef14-3* mutants (g). The degree of 3 DAG PC
interdigitation was determined by the width of indentation (h). **j**, Activation of ROP6 in the wild-type
(*GFP-ROP6*) and *gef14-2* (*gef14-2xGFP-ROP6*) mutant were analyzed by pull-down and determined
by a GFP antibody. Protoplasts isolated from 4 weeks old adult plants were used in this assay.
1010

1012

Supplementary Figure 6. FER, GEF14, and ROP6 genetically act in the same signaling pathway in regulating PC morphogenesis. a, FER associates with RopGEF14 in transgenic plants. Proteins from 10-day-old *35S::GFP-GFE14* or *35S::GFP* transgenic seedlings were immunoprecipitated with α-FER antibody and analyzed with Western blot using α-GFP antibody (Top). The expression of GFP-GEF14, GFP, and FER in transgenic plants are shown (Middle and Bottom). **b,** FER associates with ROP6 in protoplasts. Co-IP was carried out with a α-GFP-Trap antibody (IP: α-GFP-Trap), and the proteins were analyzed by using Western blot with the α-Flag antibody. Top shows that ROP6-Flag coimmunoprecipitated with FER-GFP (IP: α-GFP-Trap; WB: α-Flag). Middle and Bottom show the expression of ROP6-Flag and FER-GFP proteins (WB: α-Flag or α-GFP for input control). **c,** FER associates with RopGEF14 through intracellular kinase domain. The full length of FER-HA, truncated FER-ECD-HA and FER-TCD-HA were expressed in protoplasts and pull down was carried out with recombinant MBP or MBP-GEF14 proteins (PD: α-MBP/MBP-GEF14), and the proteins were analyzed by using Western blot with α-HA antibody. Top shows that FER-HA and FER-TCD-HA coimmunoprecipitated with MBP-GEF14 (PD: α-MBP/MBP-GEF14; WB: α-HA). Middle shows the expression of FER-HA, FER-ECD-HA and FER-TCD-HA proteins (WB: α-HA for input control), and Bottom showed the input recombinant proteins (WB: α-MBP, for input control). **d,** ROP6 associates with RopGEF14. ROP6-Flag was expressed in protoplasts and pull down was carried out with recombinant MBP or MBP-GEF14 proteins (PD: α-MBP/MBP-GEF14), and the proteins were analyzed by using Western blot with α-Flag antibody. Top shows that ROP6-Flag coimmunoprecipitated with MBP-GEF14 (PD: α-MBP/MBP-GEF14; WB: α-Flag). Middle shows the expression of ROP6-Flag proteins (WB: α-Flag for input control), and Bottom showed the input recombinant proteins (WB: α-MBP, for input control).

1035

Supplementary Figure 7. FER, GEF14, and ROP6 genetically act in the same signaling pathway in regulating PC morphogenesis. a ROP6 genetically acts in the same pathway as FER in regulating PC morphogenesis. PC morphogenesis was characterized by wild-type, *fer-4, CA-ROP6,* and *CA-ROP6fer-4.* Scale bar, 50 μ m. **b and c**, GEF14 genetically acts in the same signaling pathway as FER in regulating PC morphogenesis. PC phenotypes were characterized by 2 DAG wild-type, *gef14-2*, *fer-4* and *fer-4gef14-2* mutants (b). The degree of PC interdigitation was determined by the width of indentation (c). **d and e**, ROP6 genetically acts in the same signaling pathway as GFE14 in regulating PC morphogenesis. PC phenotypes were characterized by 2 DAG wild-type, *gef14-2*, *rop6* and *gef14-2rop6* mutants (d). The degree of 2 DAG PC interdigitation was determined by the width of indentation (e). Asterisks indicate the significant difference (p≤0.05, Student's *t*-test) between the wild type and the mutants in above assays.