Streptococcal Lancefield polysaccharides are critical cell wall determinants for human group IIA secreted phospholipase A2 to exert its bactericidal effects

Vincent P. van Hensbergen[1], Elin Movert[2], Vincent de Maat[1], Yoann Le Breton[4], Gérard Lambeau[5], Christine Payré[5], Anna Henningham[6,#], Victor Nizet[6, 7], Jos A.G. van Strijp[1], Fredric Carlsson[2, 3], Kevin S. McIver[4], Nina M. van Sorge[1] *

[1]*Medical Microbiology, University Medical Center Utrecht, Utrecht, University, Utrecht, The Netherlands;* [2]*Department of Experimental Medical Science, Section for Immunology and* [3]*Department of Biology, Section for Molecular Cell Biology, Lund University, Lund, Sweden;* [4]*Department of Cell Biology & Molecular Genetics and Maryland Pathogen Research Institute, University of Maryland, College Park, Maryland, United States of America;* [5]*Université Côte d'Azur, CNRS, Institut de Pharmacologie Moléculaire et Cellulaire, Valbonne, France;* [6]*Department of Pediatrics and* [7]*Skaggs School of Pharmacy & Pharmaceutical Sciences, University of California, San Diego, La Jolla, CA, United States of America.*

Child Healthy Research Centre, The University of Queensland, Brisbane, Australia; a.henningham@uq.edu.au

* Corresponding author: nsorge3@umcutrecht.nl

Short title

Bactericidal mechanism of hGIIA against streptococci

Abstract

Human Group IIA secreted phospholipase A$_2$ (hGIIA) is an acute phase protein with bactericidal activity against Gram-positive bacteria. Infection models in hGIIA transgenic mice have highlighted the importance of hGIIA as an innate defense mechanism against the human pathogens Group A *Streptococcus* (GAS) and Group B *Streptococcus* (GBS). Compared to other Gram-positive bacteria, GAS is remarkably resistant to hGIIA activity. To identify GAS resistance mechanisms, we exposed a highly saturated GAS M1 transposon library to recombinant human hGIIA and compared relative mutant abundance with library input through transposon-sequencing (Tn-seq). Based on transposon prevalence in the output library, we identified nine genes, including *dltA* and *lytR,* conferring increased hGIIA susceptibility. In addition, seven genes conferred increased hGIIA resistance, which included two genes, *gacH* and *gacI* that are located within the Group A Carbohydrate (GAC) gene cluster. Using GAS 5448 wild-type and the isogenic *gacI* mutant and gacI-complemented strains, we demonstrate that loss of the GAC *N*-acetylglucosamine (GlcNAc) side chain in the Δ*gacI* mutant increases hGIIA resistance approximately 10-fold, a phenotype that is conserved across different GAS serotypes. Increased resistance is associated with delayed penetration of hGIIA through the cell wall. Correspondingly, loss of the Lancefield Group B Carbohydrate (GBC) rendered GBS significantly more resistant to hGIIA-mediated killing. This suggests that the streptococcal Lancefield antigens, which are critical determinants for streptococcal physiology and virulence, are required for the human bactericidal enzyme hGIIA to exert its bactericidal function.

Author summary

The human immune system is capable of killing invading bacteria by secreting antimicrobial proteins. Cationic human Group IIA secreted phospholipase A_2 (hGIIA) is especially effective against Gram-positive bacteria by degrading the bacterial membrane. HGIIA requires binding to negatively charged surface structures before it can penetrate through the thick peptidoglycan layer and reach the target phospholipid membrane. Both binding and penetration are dependent on the cationic properties of hGIIA and anionic surface charge of bacteria. HGIIA is constitutively expressed at high concentrations at sites of possible bacterial entry, e.g. in tears, skin and small intestine. In serum, normal concentrations are low but can increase up to 1,000-fold upon inflammation or infection. HGIIA is important for defense against two human pathogens, Group A and Group B *Streptococcus* (GAS, GBS), as shown in mouse infection models. We demonstrate that the Lancefield cell wall polysaccharides that are expressed by these bacteria, the Group A Carbohydrate (GAC) for GAS and the Group B Carbohydrate (GBC) for GBS, are required for optimal hGIIA bactericidal efficacy by facilitating penetration through the peptidoglycan layer. Given the increased hGIIA resistance of antigen-modified or antigen-deficient streptococci, it will be of interest to determine potential regulatory mechanisms regarding expression of streptococcal Lancefield polysaccharides.

Introduction

Many important human bacterial pathogens are also common colonizers of mucosal barriers. Occasionally, such pathogens penetrate these physical barriers to invade the underlying tissues and cause infections. Antimicrobial molecules, sometimes also referred to as 'endogenous antibiotics of the host', are a critical part of the innate immune response to eradicate these intruders and clear the infection. In humans, one of the most potent bactericidal molecules against Gram-positive bacteria is the secreted enzyme human Group IIA phospholipase A2 (hGIIA) [1,2].

HGIIA belongs to a family of 11-12 secreted phospholipase A_2 (sPLA$_2$) enzymes, which are structurally related and hydrolyze various phospholipids [2–5]. In non-inflamed conditions, hGIIA serum levels are low and not sufficient to kill most Gram-positive bacteria [6]. However, sterile inflammation or infection increases hGIIA expression with concentrations reaching up to 1 µg/ml in serum [7], which is sufficient to kill most Gram-positive pathogens *in vitro*. A unique feature of hGIIA compared to other secreted Phospholipase A_2 family members is its high cationic charge, which is required for binding to negatively-charged surface structures and for penetration of the thick peptidoglycan layer surrounding Gram-positive bacteria [2,8,9]. The potent bactericidal activity of hGIIA against Gram-positive bacteria has been demonstrated both *in vitro*, using recombinant hGIIA, as well as *in vivo* using hGIIA transgenic mice [10–16]. To counter the bactericidal effects of hGIIA, pathogens have evolved different resistance mechanisms, for example by suppressing hGIIA expression [17,18] or by increasing the net positive charge of surface structures and membrane. The surface modifications include the addition of positively-charged D-alanine moieties to teichoic acid polymers by the highly conserved *dlt* operon to repulse hGIIA [8] and other cationic antimicrobials [19–22]. In addition, *Staphylococcus aureus* (*S. aureus*) modifies the charge of its bacterial membrane through the molecule MprF [23,24] by adding the cationic amino acid lysine to phosphatidylglycerol (PG), resulting in lysyl-PG [25]. In Group A *Streptococcus* (GAS), the enzyme sortase A (SrtA), a conserved

enzyme in Gram-positive bacteria that recognizes proteins with an LPXTG motif and covalently attaches them to peptidoglycan [26,27], was shown to increase hGIIA resistance [12].

Studies with recombinant hGIIA have highlighted differences in intrinsic hGIIA susceptibility between different Gram-positive species, where *Bacillus subtilis* is killed in the low ng/ml concentration range [28,29], and Group A *Streptococcus* (GAS) is one of the most resistant species known to date [12]. Interestingly, this high resistance is not a common trait of streptococcal pathogens since Group B *Streptococcus* (GBS) is killed by concentrations that are approximately 500 times lower compared to those required to kill GAS [11,12]. Streptococci are historically classified by the expression of structurally different Lancefield antigens [30]. Lancefield antigens are cell wall polysaccharides making up approximately 50% of the dry cell wall mass [31]. All GAS serotypes express the Lancefield Group A carbohydrate (GAC), which consists of a polyrhamnose backbone with alternating *N*-acetylglucosamine (GlcNAc) side chains [31], which are important for virulence [32]. In contrast, all GBS serotypes express the Lancefield Group B carbohydrate (GBC), a multi-antennary structure, containing rhamnose, galactose, GlcNAc, glucitol, and significant amounts of phosphate [33]. Both streptococcal species are important human pathogens as they can cause systemic infections associated with high mortality and morbidity [34–36]. Mouse infection models and *ex vivo* studies on human serum from infected patients demonstrated the importance of hGIIA in defense against lethal infections with GAS and GBS [11,12]. Given the importance of hGIIA in host defense against streptococci, we set out to identify the molecular mechanisms that confer resistance to hGIIA using a comprehensive and unbiased approach.

Results

Globally-disseminated M1T1 GAS is highly resistant to hGIIA

A previous study found that GAS strains are among the most resistant Gram-positive bacteria regarding hGIIA-mediated killing [12]. Mutation of *srtA* in the GAS strain JRS4, an *emm6* serotype, increased hGIIA susceptibility by about 50-fold [12]. GAS M1T1 is a globally-disseminated *emm1* clone that is most often responsible for invasive GAS infections in industrialized countries [37,38] and was not included previously in hGIIA studies [12]. GAS strain 5448, a representative M1T1 isolate, showed concentration-dependent killing by recombinant human hGIIA, with an LD_{50} of 0.05 μg/ml (Fig 1A). Also, GAS M1T1 resistance mechanisms against hGIIA at least partially overlap with GAS JRS4 *emm6*, since mutation of *srtA* rendered GAS M1T1 approximately 35-fold more susceptible to hGIIA (Fig 1A) [12].

To study the activity of hGIIA in a more physiological setting, we spiked pooled normal human serum with different concentrations of recombinant hGIIA. The addition of serum potentiated hGIIA-mediated killing of GAS compared to the purified assay (Fig 1A) as reflected by a 5-fold lower LD_{50} (0.01 ug/ml; Fig 1B). Interestingly, heat inactivation of serum reduced the ability of hGIIA to kill GAS by 10-fold compared to active serum, indicating that there are heat-labile factors in serum that potentiate hGIIA efficacy (Fig 1B). Endogenous hGIIA present in normal human serum does not play a significant role since addition of the hGIIA-specific inhibitor LY311727 [39] did not alter GAS growth in serum (Fig S1).

Identification of GAS genes that affect hGIIA susceptibility using Tn-seq

We set out to identify genes that affect hGIIA susceptibility of GAS M1T1 using the GAS *Krmit* transposon mutant library [40]. To ensure complete coverage of the library in our experiment, we optimized our hGIIA killing assay to support an inoculum of 10^7 CFU, using a final concentration of 0.125 μg/ml hGIIA. The Tn-seq experiment with the GAS *Krmit* transposon mutant library consisted of four non-exposed control samples and four hGIIA-treated samples. Each sample contained on average approximately 30 million reads, of which over 90% of the reads aligned once to the GAS M1T1 5448 reference genome (Table

S1) [41]. To quantify the number of transposon insertions per gene, we divided the reference

genome into 25 nucleotide windows, resulting in 73,182 windows, and mapped each read to

a specific window. More than 48% of the windows had at least one read aligned. We

identified one gene with an exceptionally high number of transposon insertions at a specific

part of the gene (M5005_Spy_1390), suggesting biased insertion of the transposon (Tables

S2, S3 and Fig S2). This gene was therefore excluded from further analysis. No other biased

transposon insertion sites were observed.

We identified 16 genes that contained a significantly different number of transposon

insertions after exposure to hGIIA as indicated by *P*-value of <0.05 (Benjamini-Hochberg

(BH) corrected; Fig 2A, Fig S3, and Tables S2-4). Nine of the 16 genes (56%) showed a

decrease in transposon insertions compared to untreated controls, indicating that the

products of the disrupted genes provide resistance against hGIIA-mediated GAS killing (Fig

2A, Fig S3, Table S3). Three susceptibility genes are located within the *dlt* operon

(M5005_Spy_1070, M5005_Spy_1072, M5005_Spy_1073), which is responsible for D-

alanylation of teichoic acids [42]. Consistently, this operon was previously linked to GAS

resistance against other cationic antimicrobials, such as LL-37 and hGIIA [8,43]. The other

six genes with significant fold decrease in transposon insertions are annotated as

hypothetical proteins (M5005_Spy_0918 and M5005_Spy_1794), a lactoylglutathione lyase

(M5005_Spy_0876), LytR (M5005_Spy_1474) of the LytR/CspA/Psr protein family, the

transcriptional regulator FabT (M5005_Spy_1495), and the NAD glycohydrolase inhibitor

(M5005_Spy_0140) (Fig S3 and Tables S2, S3).

Seven genes showed a relative increase in the number of transposon insertions after

hGIIA exposure, indicating that the products of these genes are important for hGIIA to exert

its bactericidal effect (Fig 2A, Fig. S3, Table S4). Five of the six genes (83%) mapped to two

gene clusters; one gene cluster is annotated as an ABC transporter (M5005_Spy_0939,

M5005_Spy_0940, M5005_Spy_0941) and the other gene cluster is the previously identified

12-gene cluster responsible for biosynthesis of the Group A carbohydrate (GAC) (Fig 2B)

[32]. Within the GAC gene cluster, *gacI* and *gacH* (M5005_Spy_0609 and

M5005_Spy_0610) showed significantly increased number of transposon insertions. The

small downstream gene *gacJ* (M5005_Spy_0611) also demonstrated a 3-fold increase,

however, the BH corrected *P*-value is slightly above 0.05. Other genes within the GAC gene

cluster are essential or crucial as described previously [40,44]. Finally, *guaB*

(M5005_Spy_1857) and the IIC component of a galactose-specific PTS system

(M5005_Spy_1399) were identified as their mutation may confer increased resistance to

hGIIA (Fig S3 and Tables S2, S4). Overall, the transposon library screen identified genes

that confer resistance or are important for the mechanisms of action of hGIIA.

189   HGIIA requires the GAC GlcNAc side chain to exert its bactericidal effect

190   against GAS

To validate the Tn-seq findings, we confirmed the involvement of three genes (*dltA*, *lytR*, and

gacI) by assessing hGIIA-mediated killing of previously generated GAS mutants [32,43,45].

Deletion of *dltA* and *lytR* indeed increased GAS susceptibility to hGIIA-mediated killing by

45-fold and 35-fold, respectively (Fig 3A and 3B). The *dltA* defect could be restored by re-

introduction of the gene on a plasmid (Fig 3A).

In contrast to *dltA* and *lytR*, mutation of *gacI*, which results in loss of the GAC

GlcNAc side chain [46], increased GAS resistance to hGIIA by approximately 10-fold

compared to the parental or *gacI*-complemented (*gacI**) strain (Fig 3C; LD_{50} = 0.5 μg/ml).

The GAC is conserved in all GAS serotypes. We therefore questioned whether deletion of

gacI would have a similar effect on the bactericidal efficacy of hGIIA in four other GAS

serotypes (M2, M3, M4, M28). In all serotypes, deletion of *gacI* increased resistance of GAS

to hGIIA by 5- to 50-fold (Fig 3D), indicating that hGIIA requires the GAC GlcNAc side chain

for optimal bactericidal efficacy in all genetic backgrounds tested.

204   Loss of the GAC GlcNAc side chain delays cell wall translocation of hGIIA

Our observation that GAS Δ*gacI* is more resistant to hGIIA implies that the GAC GlcNAc

moiety is important for the function of hGIIA. To assess whether loss of the GAC GlcNAc

side chain affected hGIIA binding to bacteria, we quantified binding of recombinant hGIIA to

GAS using a sPLA$_2$-specific antibody. At concentrations that are occasionally reached in

plasma during inflamed conditions (up to 1 μg/ml), binding of hGIIA was similar in the three

strains (Fig 4A). Only at concentrations of 5 μg/ml, hGIIA showed reduced interaction with

the *gacI* mutant compared to GAS WT and *gacI**-complemented strains (Fig 4A). The

contribution of differential hGIIA binding to GAS is therefore only relevant to specific

locations such as in tears which contain up to 30 μg/ml hGIIA [28]. Since hGIIA binding is

charge-dependent, we analyzed whether reduced binding at high hGIIA concentrations

could be caused by difference in surface charge. Using the highly cationic protein

cytochrome C, we indeed observed that the *gacI* mutant has a reduced negative surface

charge compared to GAS WT and the *gacI**-complemented strain (Fig 4B). In addition, we

analyzed whether hGIIA bound to the bacterial surface at a specific side using fluorescence

microscopy. hGIIA predominantly localized to the GAS cell poles (Fig 4C).

Cell wall architecture can significantly affect hGIIA cell wall penetration [2]. To assess

how absence of the GAC GlcNAc side chain affected hGIIA cell wall penetration, we

measured changes in membrane depolarization over time using the fluorescent voltage-

sensitive dye DiOC$_2$(3) [47]. In this assay, membrane depolarization results in reduced red

fluorescence. HGIIA required at least 5 minutes to penetrate the GAS cell wall since no

changes in red fluorescence signal were observed at this time point for any of the strains

(Fig S4A). After 5 minutes, membrane depolarization occurred as visualized by diminished

red fluorescence (Fig 4D and Fig S4B, S4C) and was most pronounced at high hGIIA

concentrations for GAS WT and the *gacI**-complemented strain. Compared to these two

strains, the *gacI* mutant exhibits only slight membrane depolarization at all time points (Fig

4D and Fig S4). To determine how membrane depolarization correlates to membrane

permeabilization, we measured influx of the fluorescent DNA dye SYTOX green, which can

only enter damaged membranes [48]. As expected, hGIIA increased the SYTOX signal in

GAS WT and GAS *gacI** after approximately 15 minutes (Fig 4E and Fig S5). SYTOX

intensity continued to increase for 60 to 80 minutes and then plateaued. In sharp contrast,

SYTOX intensity in GAS Δ*gacI* increased at a much slower rate and never reached the

levels of GAS WT and GAS *gacI** after two hours. The fact that SYTOX influx in GAS Δ*gacI*

is significantly slower and reduced compared to GAS WT suggests that the GAC GlcNAc

side chain is essential for efficient trafficking of hGIIA through the GAS cell wall.

A recent study demonstrates that GacI is a membrane protein that is required for the

intracellular formation of undecaprenyl-P-GlcNAc [46]. Therefore, loss of GacI could alter

membrane composition and fluidity to impact the activity of hGIIA on the membrane. To

analyze whether phospholipid hydrolysis is affected in GASΔ*gacI*, we performed the SYTOX

influx assay on protoplasts [49]. Unlike the previous SYTOX results with intact bacteria,

protoplasts from WT, Δ*gacI* and *gacI** strains all became SYTOX positive (Fig 4F and Fig

S6), underlining our conclusion that the presence of the cell wall in the Δ*gacI* limits access of

hGIIA to the streptococcal membrane. Nonetheless, the significantly lower SYTOX in the

Δ*gacI* protoplasts compared to the WT and *gacI**-complemented protoplasts (Fig 4F and Fig

S6), suggest that the absence of GacI reduces degradation by hGIIA.

In summary, these results suggest that hGIIA requires an intact GAC to efficiently

migrate through the cell wall. In addition, the presence of GacI in the cell membrane

increases membrane degradation by hGIIA likely due to a difference in membrane fluidity.

GBC is important for hGIIA bactericidal activity against GBS

We wondered whether the importance of the GAC for hGIIA activity could be extended to

other streptococci such as GBS. As previously described, GBS are generally more sensitive

to hGIIA compared to GAS [12]. Indeed, killing of GBS strain NEM316 occurred at

substantially lower concentrations of hGIIA compared to GAS M1T1 (compare Fig 5A and

1A). We confirmed that killing depends on the catalytic activity of the enzyme since

introduction of an inactivating point mutation (H48Q; Fig 5B) or addition of the hGIIA-specific

inhibitor LY311727 abrogated all killing (Fig 5C). Just as the GAC is the molecular signature

for GAS, GBS uniquely express another Lancefield antigen, known as the Group B

Carbohydrate (GBC). The GBC is a more complicated structure compared to the GAC and

contains significant amounts of phosphate that introduce a negative charge. GBS lacking the GBC, through deletion of *gbcO* [33], are at least 100-fold more resistant to hGIIA compared to GBS WT (Fig 5A-C), and the phenotype is restored upon complementation with *gbcO* on a plasmid (Fig 5A). We could reproduce the Δ*gbcO* phenotype by treating WT GBS with tunicamycin, an inhibitor of *gbcO*-type transferases (Fig 5D) [33,50]. Finally, as observed in GAS, fluorescence microscopy demonstrated that hGIIA bound to the poles of GBS WT (Fig 5E). In conclusion, these results highlight a key role for streptococcal Lancefield antigens in the bactericidal effect of hGIIA.

Discussion

The resistance to acute phase protein hGIIA varies among Gram-positive bacteria, including among closely-related streptococcal species. GAS, an important cause of lethal infection worldwide, is among the most resistant bacteria, whereas GBS, an important cause of neonatal sepsis and meningitis, is killed by hGIIA at concentrations that are approximately 500-fold lower [12]. For GAS, we confirmed the role of Sortase A and DltA and identified LytR as hGIIA resistance factors. Importantly, the streptococcal Lancefield antigens, which are characteristic cell wall polysaccharides, are structural requirements for the activity of hGIIA in both GAS and GBS. Together, these results indicate that the precise cell wall architecture is an important determinant for streptococcal hGIIA susceptibility.

HGIIA is approximately 10-fold more effective against GAS when spiked into normal serum compared to heat-inactivated serum, and 5-fold more effective compared to our 'purified' system. This corresponds to a previous observation where hGIIA activity was approximately 10-fold greater in serum or plasma than in the protein-depleted serum in studies using *S. aureus* as the target pathogen [51]. This suggests the presence of a heat labile protein in plasma that facilitates hGIIA-mediated killing of Gram-positive bacteria. Heat-inactivation of serum is a well-established method to study the influence of the complement system and also abolishes hGIIA activity in acute phase serum [52]. Since the low basal levels of hGIIA in normal human serum are not sufficient to affect GAS viability,

the enhancement could indicate a synergistic effect between hGIIA and the complement

system. A recent study shows formation of the Membrane Attack Complex (MAC) on the

GAS surface without affecting bacterial viability [53]. It is therefore tempting to speculate that

MAC is deposited on Gram-positive bacteria so that bactericidal enzymes, such as hGIIA,

can reach the bacterial membrane more easily.

Sortase A, an enzyme that links LPXTG-containing proteins to peptidoglycan, is

known to be involved in GAS hGIIA resistance [12]. Whether a single or multiple LPXTG

proteins confer resistance is an unresolved question. Our study suggests that Sortase-A

mediated resistance is not caused by a single LPXTG protein since we did not identify a

single LPXTG-encoding gene in the Tn-seq screen (Table S5). Possibly, the underlying

mechanism is similar to the SrtA-dependent resistance of GAS to the antimicrobial peptide

cathelicidin [54], which depends on the accumulation of sorting intermediates at the bacterial

membrane. *SrtA* itself was not identified in the transposon library screen since the mutants

are lost in the competitive environment likely due to inherent defects in growth [40].

We identified and confirmed a role for the protein LytR in GAS hGIIA resistance. LytR

is a member of the LytR-CpsA-Psr (LCP) protein family, a conserved family of cell wall

assembly proteins in Gram-positive bacteria [55]. The GAS genome encodes two members

of this family, *lytR* (M5005_Spy_1474) and *psr* (M5005_Spy_1099). The fact that we only

identified LytR suggests that these proteins have non-redundant, but as yet unidentified,

functions. In several Gram-positive pathogens, including *Streptococcus pneumoniae*,

Staphylococcus aureus and *Bacillus anthracis*, LCP proteins anchor cell wall glycopolymers

such as wall teichoic acid (WTA), lipoteichoic acid (LTA) and capsular polysaccharides to

the cell envelope and are therefore critical for cell envelope assembly and virulence [55–60].

Additionally, *lytR* homologues in *Bacillus subtilis* and *Streptococcus mutans* contribute to cell

wall remodeling by increasing autolysin activity [61,62]. Previously, hGIIA activity has been

linked to autolysins; autolysin-deficient mutants are more resistant to hGIIA than their parent

strain [63]. A suggested mechanism is that hGIIA displaces positively-charged autolysins

from negatively-charged WTA and LTA, resulting in localized peptidoglycan digestion and

facilitated movement of hGIIA through the cell wall. Currently, the role of LytR either in GAS cell wall assembly or in the regulation of autolysin activity is not known, but LytR-deficient GAS display altered membrane integrity and potential [64], which could impact hGIIA susceptibility. Moreover, *lytR* has been linked to GAS virulence in two different studies. In the first study, *lytR* mutants in two different GAS M1 backgrounds showed a more virulent phenotype in a subcutaneous murine model of infection, which was suggested to be a result of increased SpeB activity [64]. LytR-mediated regulation of SpeB is unlikely to play a role in hGIIA-mediated resistance in our experiments, since we used washed bacteria. In a more recent study, *lytR* mutants in GAS 5448 M1T1 showed a competitive disadvantage for fitness *in vivo* upon mixed subcutaneous infection [45]. Unfortunately, there is no information regarding pathology or survival of the mice upon infection with the *lytR* mutant added alone [45].

We also identified genes that render GAS more susceptible to hGIIA. *GacH*, *gacI*, and *gacJ* are located in the biosynthesis gene cluster of the GAC, which may suggest that the GAC is a target for hGIIA on the GAS surface. Mutation of *gacI* and *gacJ* results in loss of the GAC GlcNAc side chain [32,46], whereas mutation of *gacH* does not affect side chain formation [32]. We therefore hypothesize that the GAC provides hGIIA resistance through two distinct mechanisms. First, a *gacI*/*J*-dependent mechanism that works through the GAS GlcNAc side chain as important for binding and penetration of hGIIA to the cell membrane. The second mechanism involves GacH but the underlying molecular aspects remain to be determined. The first mechanism seems to conflict with our previous observations that GlcNAc-deficient GAS have decreased virulence capacity due to increased neutrophil killing and increased susceptibility to antimicrobials in serum including LL-37 [32]. However, hGIIA would not have contributed to *in vitro* assays since we used non-inflamed serum or plasma where basal hGIIA concentrations are too low to affect GAS viability [32]. The fact that *gacI* mutants demonstrate reduced survival *in vivo* suggests that the benefits of expressing the GlcNAc side chain outweigh the increased susceptibility to hGIIA. Since GAS already shows high intrinsic resistance towards hGIIA there is no pressure to lose the GlcNAc side chain. It

might even be detrimental since it makes GAS more vulnerable to effects of other

antimicrobials or yet unidentified host defenses. In contrast to the GAC [31,65], the GBC is a

multi-antennary structure and contains anionic charge due to the presence of phosphate

[33]. For GBS, the increased hGIIA resistance in GBC-negative *gbcO* mutants is therefore

likely explained by the loss of negatively charged groups on the surface. This corresponds to

previous observations in *S. aureus*, where loss of the secondary cell wall glycopolymer

WTA, increased resistance to several antimicrobial proteins, including hGIIA [10].

Contrary to our expectations, fluorescence microscopy analysis showed that hGIIA

bound to the cell poles of both GAS and GBS. This binding pattern does not correspond to

the localization of the GAC and GBC, which are distributed over the entire cell wall as shown

by early electron microscopy studies [66,67]. Binding at the septa of dividing bacteria seems

a preferred binding site for bactericidal agents due to a high turnover of peptidoglycan which

would make penetration easier [68,69]. In addition, the septum is rich in anionic

phospholipids [70], a likely target for cationic hGIIA. Finally, the GAS ExPortal, a unique

microdomain in the GAS membrane that is enriched in anionic lipids, would be another

favored location of binding for the cationic hGIIA [71]. However, the ExPortal is distributed

asymmetrically across the GAS surface and not at the cell poles [71]. The fact that we

observe a similar binding pattern to GBS and GAS, may indicate that GAS and GBS express

a similar protein that localizes at the cell poles and is used by hGIIA as an initial docking

site. The exact reason for hGIIA binding at the cell poles remains to be determined.

Lack of the GAC GlcNAc side chain most profoundly affected penetration of hGIIA

through the cell wall, a mechanism that depends on charge [2,9]. Indeed, membrane

depolarization and permeabilization occurs at a much slower rate in the *gacI* mutant

compared to WT and complemented strains. This implies that the GAC GlcNAc side chain

facilitates penetration of hGIIA through the cell wall in what is referred to as an 'anionic

ladder process' [2]. Interestingly, the GAC does not contain any charged structures.

Therefore, the underlying mechanism may be linked to the previously mentioned autolysin

displacement from interaction with the GAC.

In conclusion, we show that the bactericidal agent hGIIA is able to kill GAS in a

complex serum environment. However, modification or removal of the Lancefield antigen

renders GAS more resistant to the bactericidal activity of hGIIA. Similarly, removing the

Lancefield antigen from GBS renders this species also more resistant to the bactericidal

activity of hGIIA. The Lancefield antigens, previously thought to be solely involved in

physiology, are thus critical cell wall structures for hGIIA to exert its bactericidal effect. The

Tn-seq data discussed in this paper provide exciting new insights into the resistance

mechanisms of GAS and encourage similar experiments in other streptococci species.

Disrupting the resistance mechanisms with therapeutic agents could possibly be sufficient to

provide our own immune system the upper hand in clearing invading streptococcal

pathogens.

Materials and Methods

Bacterial strains

The GAS M1T1 5448 strain was used in this study unless stated otherwise. The 5448Δ*gacI*

knockout and *gacI** complemented strain [32], the 5448Δ*lytR* [45] and the GAS serotypes

M2, M3, M4, and M28 and corresponding Δ*gacI* knockouts [72] were described previously.

Preparation and characterization of the GAS M1T1 5448 transposon library was described

previously by Le Breton et al., 2015 [40]. All GAS strains were grown in Todd-Hewitt broth

(Becton Dickinson) supplemented with 1% yeast extract (Oxoid; THY) as static cultures at

37 °C. Kanamycin (Sigma-Aldrich) was used at a concentration of 300 µg/ml when

appropriate. GBS NEM316 WT, Δ*gbcO* and the complemented strains Δ*gbcO*/pTCV were

kindly provided by Dr. Mistou [33].

Unless stated otherwise, overnight cultures of GAS were diluted and re-grown to

mid-log phase (OD_{600nm} = 0.4), washed and resuspended in HEPES solution (20 mM

HEPES, 2 mM Ca^{2+}, 1% BSA [pH 7.4]) solution at OD_{600nm} = 0.4 (~$1x10^8$ CFU/ml). For GBS

strains, overnight cultures of NEM316 WT, Δ*gbcO* and the complemented strains

Δ*gbcO*/pTCV were diluted in TH broth and grown to mid-log phase (OD_{620nm} = 0.4 for WT

and complemented strains, 0.25 for Δ*gbcO* mutant). Bacteria were then diluted in HEPES

solution and pushed rapidly through a 27-gauge needle, a process repeated three times, to

disrupt bacterial aggregates.

Identification of GAS resistance determinants against hGIIA

Recombinant hGIIA was produced as described previously [73]. The GAS M1T1 *Krmit*

transposon mutant library was grown to mid-log phase in 100 ml THY containing Km and

resuspended in HEPES solution to OD_{600nm}= 0.4. Four experimental replicates of 100 µl (~

1×10^7 CFU) were subsequently incubated in HEPES solution with or without 125 ng/ml

hGIIA for 1 hour at 37 °C. After incubation, 3 ml THY was added to all samples and

incubated at 37 °C until the mid-log phase was reached (recovery step). Cultures were

collected by centrifugation and used for isolation of genomic DNA (gDNA). gDNA was

isolated by phenol-chloroform extraction. Samples were barcoded and prepared for Tn-seq

sequencing as described previously [74]. Tn-seq sequencing was performed on Illumina

NextSeq500 (Sequencing facility University Medical Center, Utrecht, The Netherlands).

Tn-seq data analysis was performed as previously described [74]. In short, barcodes

were split using the Galaxy platform [75] and sequences were mapped to the GAS M1T1

5448 genome [41] using Bowtie 2 [76]. The genome was subsequently divided in 25-bp

windows and each alignment was sorted and indexed by IGV [77]. Insertions were counted

per window and then summed over the genes. Read counts per gene were adjusted to cover

only the first 90% of the gene since transposon insertions in the final 10% potentially do not

cause a knock-out phenotype. Then, read counts were normalized to the total number of

reads that mapped to the genome in each replicate, by calculating the normalized read-

count RKPM (Reads Per Kilobase per Million input reads; RKPM = (number of reads

mapped to a gene x 10^6) / (total mapped input reads in the sample x gene length in kbp)).

Cyber-T [78] was used to perform statistical analysis on the RKPM values. Genes that

contributed to either hGIIA susceptibility or hGIIA resistance were determined when the

Benjamini-Hochberg (BH) corrected *p*-value was <0.05.

hGIIA susceptibility

Mid-log streptococcal suspensions were diluted 1,000 times in HEPES solution and 10 µl

was added to sterile round-bottom 96 well plates (triplicates). Recombinant hGIIA or

catalytically-deficient hGIIA mutant enzyme H48Q was serially diluted in HEPES solution or

human serum and 10 µl aliquots were added to bacteria-containing wells. For hGIIA

inhibition experiments, 50 µM LY311727 was added to the HEPES solution or serum. For

GAS, samples were incubated for 2 hours at 37°C, without shaking, PBS was added and

samples were 10-fold serially diluted and plated on THY agar plates for quantification. For

GBS, bacteria were incubated with hGIIA at 37°C for 30 minutes, the samples were diluted

in PBS and plated onto blood agar plates. After overnight incubation 37°C, colony forming

units (CFU) were counted to calculate the survival rate (Survival (% of inoculum) = (counted

CFU * 100) / CFU count of original inoculum or Survival (%) = (counted CFU * 100) / CFU

count at 0 µg/ml hGIIA). For pharmacological inhibition of GBC expression, NEM316 WT

bacteria were grown to mid-log phase (OD_{620nm} = 0.4) in the presence of 0.5 mg/ml

tunicamycin (Sigma) and used in killing assays as described above.

Membrane potential and permeability assays

Changes in hGIIA-dependent membrane potential were determined using the membrane

potential probe $DiOC_2(3)$ (PromoKine) [47,79]. Bacterial suspensions (OD_{600nm}= 0.4) were

diluted 100 times (~$1x10^6$ CFU/ml), 100 µl aliquots were divided into eppendorf tubes and

incubated with serial dilutions of hGIIA. After incubation at 37°C, 3 mM $DiOC_2(3)$ was added

and incubated at room temperature for 5 minutes in the dark. Changes in green and red

fluorescence emissions were analyzed by flow cytometry.

Bacterial staining with the DNA stain SYTOX Green (Invitrogen) is a measurement

for membrane permeabilization and an indication of bacterial cell death [48]. Serial dilutions

of hGIIA in HEPES solutions were added to wells of a sterile flat-bottom 96 well plate.

Bacteria were resuspended in HEPES solution containing 1 µM SYTOX green (OD_{600nm} =

0.4) and added to hGIIA dilutions in a final volume of 100 μl. Fluorescence over time was

recorded using FLUOstar OPTIMA (green fluorescence 530 nm emission and excitation 488

nm) at 37°C.

Surface charge determination

Bacterial surface charge was determined as previously described [79]. Briefly, exponential

phase bacteria (OD_{600nm} = 0.4) were washed twice in 20 mM MOPS buffer [pH 7.0] and

adjusted to OD_{600nm} = 0.7. After a 10-fold concentration step, 200 μl bacterial aliquots were

added to 200 μg cytochrome c (from *Saccharomyces cerevisiae*, Sigma-Aldrich) in a sterile

96-well round-bottom plate. After 10 minutes at room temperature in the dark, the plate was

centrifuged, the supernatant was transferred to a sterile 96 well flat-bottom plate and

absorbance was recorded at 530 nm. The percentage of bound cytochrome c was

calculated using samples containing MOPS buffer only (100% binding) and samples

containing MOPS buffer and cytochrome c (0% binding).

hGIIA surface binding

To determine hGIIA surface binding, 12.5 μl of bacterial cultures in mid-log phase (OD_{600nm} =

0.4) were added to wells of a sterile 96-well round-bottom plate (triplicates). hGIIA was

serially diluted in HEPES solution without Ca^{2+} and added to the bacteria at indicated

concentrations. After 30 minutes incubation at 4°C, bacteria were collected by centrifugation

and resuspended in HEPES solution without Ca^{2+} containing 1:300 dilution of anti-$sPLA_2$

antibody (Merck Millipore) [28]. After incubation at 4°C for 30 minutes, the samples were

washed and incubated with a 1:1,000 dilution of FITC-labeled goat-anti-mouse IgG

(SouthernBiotech). After washing with HEPES solution without Ca^{2+}, samples were fixed

with 1% paraformaldehyde and fluorescence was recorded by flow cytometry (FACSVerse,

BD Biosciences).

Fluorescence microscopy

To analyze hGIIA surface localization by microscopy, bacteria were grown in 10 ml broth to

mid-log phase and washed with 0.1 M NaHCO$_3$ [pH 9]. For GBS, the bacterial septa were

stained by addition of a 1:1 mixture of Vancomycin bodipy® FL conjugate (Invitrogen,

V34550) and vancomycin (Sigma) at a final concentration of 1.25 μg/ml, during the last

generation time of growth. The surface of GBS was stained with Alexa Fluor® 350

Carboxylic acid Succinimidyl ester (Molecular Probes by Life Technologies, A10168) for 1

hour in room temperature. Bacteria were then resuspended in 500 μl HEPES solution and

the suspension was divided over two tubes. A final concentration of 10 μg/ml hGIIA H48Q

was added to one tube and HEPES solution to the other before a 30 minute incubation at

room temperature. The samples were washed and resuspended in 200 μl HEPES solution,

then again divided to two tubes. A mouse anti-human hGIIA monoclonal antibody (Clone

SCACC353 Cayman Chemical) or an IgG1 isotype control (mouse anti human IgA clone

6E2C1, DAKO) was added to a final concentration of 10 μg/ml to the bacterial suspensions

and incubated at RT. After washing, the samples were incubated with 8 μg/ml of Alexa

Flour® 594 goat anti-mouse IgG1 (Molecular Probes by Life Technologies, A21125). After

30 min incubation, the samples were washed in HEPES solution and fixed in 4%

paraformaldehyde. Ten μl of bacterial suspension were mounted onto microscopic slides

(VWR) using MOWIOL (Sigma) mounting medium before viewing the samples using Zeiss

Axiovert 200M microscope. Pictures were captured using a 63× objective and the

AXIOVISION 4.8 software.

Hydrolysis of membrane phospholipids

To determine hGIIA efficacy in hydrolyzing membrane phospholipids, the membrane

permeabilization assay was modified for protoplasts. Mid-log bacterial suspension were

prepared in in protoplast buffer (20% sucrose, 20 mM Tris-HCl, 10 mM MgCl$_2$, 2 mM CaCl$_2$

[pH 7.4]) containing 1.4 units/μl mutanolysin (Sigma-Aldrich) [49,80,81]. After incubation for

1 hour at 37°C, protoplasts were collected by centrifugation (1,200 rpm 15 minutes) and

resuspended in protoplast buffer to an OD_{600nm} = 0.4. Pore formation by hGIIA was

monitored using SYTOX Green as described above.

Statistical analysis

GraphPad Prism 6 was used to perform statistical analysis. An unpaired two-tailed Student's

t-test was used to compare the means of two groups. A 2-way ANOVA with Bonferroni

multiple comparison test was used to compare multiple groups. Data shown are mean ± SD.

References

1. Nevalainen TJ, Graham GG, Scott KF. Antibacterial actions of secreted
phospholipases A2. Review. Biochim Biophys Acta - Mol Cell Biol Lipids. 2008;1781:
1–9. doi:10.1016/j.bbalip.2007.12.001

2. Weiss JP. Molecular determinants of bacterial sensitivity and resistance to
mammalian Group IIA phospholipase A2. Biochim Biophys Acta - Biomembr. Elsevier
B.V.; 2015;1848: 3072–3077. doi:10.1016/j.bbamem.2015.05.018

3. Gelb MH, Jain MK, Hanel AM, Berg OG. Interfacial enzymology of glycerolipid
hydrolases: lessons from secreted phospholipases A2. Annu Rev Biochem. 1995;64:
653–88. doi:10.1146/annurev.bi.64.070195.003253

4. Lambeau G, Gelb MH. Biochemistry and Physiology of Mammalian Secreted
Phospholipases A $_2$. Annu Rev Biochem. 2008;77: 495–520.
doi:10.1146/annurev.biochem.76.062405.154007

5. Murakami M, Lambeau G. Emerging roles of secreted phospholipase A2 enzymes: An
update. Biochimie. Elsevier Masson SAS; 2013;95: 43–50.
doi:10.1016/j.biochi.2012.09.007

6. Nevalainen T, Kortesuo P. Immunochemical detection of group I and group II
phospholipases A2 in human serum. Clin Chem. 1992;38: 1824–1829. Available:
http://www.clinchem.org/content/38/9/1824.short

7. Green J, Smith G, Buchta R, Lee R, Ho K, Rajkovic I, et al. Circulating phospholipase
A2 activity associated with sepsis and septic shock is indistinguishable from that
associated with rheumatoid arthritis. Inflammation. 1991;15: 355–367.

8. Koprivnjak T, Peschel A, Gelb MH, Liang NS, Weiss JP. Role of charge properties of
bacterial envelope in bactericidal action of human Group IIA phospholipase A2
against *Staphylococcus aureus*. J Biol Chem. 2002;277: 47636–47644.
doi:10.1074/jbc.M205104200

9. Beers SA, Buckland AG, Koduri RS, Cho W, Gelb MH, Wilton DC. The antibacterial
properties of secreted phospholipases A2: A major physiological role for the group IIA
enzyme that depends on the very high pI of the enzyme to allow penetration of the
bacterial cell wall. J Biol Chem. 2002;277: 1788–1793. doi:10.1074/jbc.M109777200

10. Koprivnjak T, Weidenmaier C, Peschel A, Weiss JP. Wall teichoic acid deficiency in

545		*Staphylococcus aureus* confers selective resistance to mammalian group IIA
546		phospholipase A2 and human β-defensin 3. Infect Immun. 2008;76: 2169–2176.
547		doi:10.1128/IAI.01705-07

11. Movert E, Wu Y, Lambeau G, Kahn F, Touqui L, Areschoug T. Secreted group IIA phospholipase A2 protects humans against the Group B *Streptococcus*: Experimental and clinical evidence. J Infect Dis. 2013;208: 2025–2035. doi:10.1093/infdis/jit359

12. Movert E, Wu Y, Lambeau G, Touqui L, Areschoug T. A novel bacterial resistance mechanism against human group IIA-secreted phospholipase A2: role of *Streptococcus pyogenes* sortase A. J Immunol. 2011;187: 6437–46. doi:10.4049/jimmunol.1100499

13. Gelb MH, Valentin E, Ghomashchi F, Lazdunski M, Lambeau G. Cloning and recombinant expression of a structurally novel human secreted phospholipase A2. J Biol Chem. 2000;275: 39823–39826. doi:10.1074/jbc.C000671200

14. Koduri RS, Gronroos JO, Laine VJ, Le Calvez C, Lambeau G, Nevalainen TJ, et al. Bactericidal properties of human and murine groups I, II, V, X, and XII secreted phospholipases A(2). J Biol Chem. 2002;277: 5849–5857. doi:10.1074/jbc.M109699200

15. Piris-Gimenez A, Paya M, Lambeau G, Chignard M, Mock M, Touqui L, et al. In vivo protective role of human group IIa phospholipase A2 against experimental anthrax. J Immunol. 2005;175: 6786–6791. doi:10.4049/jimmunol.175.10.6786

16. Pernet E, Guillemot L, Burgel PR, Martin C, Lambeau G, Sermet-Gaudelus I, et al. *Pseudomonas aeruginosa* eradicates *Staphylococcus aureus* by manipulating the host immunity. Nat Commun. 2014;5. doi:10.1038/ncomms6105

17. Raymond B, Leduc D, Ravaux L, Le Goffic R, Candela T, Raymondjean M, et al. Edema toxin impairs anthracidal phospholipase A2 expression by alveolar macrophages. PLoS Pathog. 2007;3: 1907–1917. doi:10.1371/journal.ppat.0030187

18. Raymond B, Ravaux L, Mémet S, Wu YZ, Sturny-Leclère A, Leduc D, et al. Anthrax lethal toxin down-regulates type-IIA secreted phospholipase A2expression through MAPK/NF-κB inactivation. Biochem Pharmacol. 2010;79: 1149–1155. doi:10.1016/j.bcp.2009.11.023

19. Weidenmaier C, Peschel A, Kempf VAJ, Lucindo N, Yeaman MR, Bayer AS. DltABCD- and MprF-mediated cell envelope modifications of *Staphylococcus aureus* confer resistance to platelet microbicidal proteins and contribute to virulence in a rabbit endocarditis model. Microbiology. 2005;73: 8033–8038. doi:10.1128/IAI.73.12.8033

20. McBride SM, Sonenshein AL. The dlt operon confers resistance to cationic antimicrobial peptides in *Clostridium difficile*. Microbiology. 2011;157: 1457–1465. doi:10.1099/mic.0.045997-0

21. Abachin E, Poyart C, Pellegrini E, Milohanic E, Fiedler F, Berche P, et al. Formation of D-alanyl-lipoteichoic acid is required for adhesion and virulence of *Listeria monocytogenes*. Mol Microbiol. 2002;43: 1–14. doi:10.1046/j.1365-2958.2002.02723.x

22. Fisher N, Shetron-Rama L, Herring-Palmer A, Heffernan B, Bergman N, Hanna P. The dltABCD operon of *Bacillus anthracis* Sterne is required for virulence and resistance to peptide, enzymatic, and cellular mediators of innate immunity. J Bacteriol. 2006;188: 1301–1309. doi:10.1128/JB.188.4.1301-1309.2006

23. Ernst CM, Peschel A. Broad-spectrum antimicrobial peptide resistance by MprF-mediated aminoacylation and flipping of phospholipids. Mol Microbiol. 2011;80: 290–299. doi:10.1111/j.1365-2958.2011.07576.x

24.   Nevalainen TJ, Haapamäki MM, Grönroos JM. Roles of secretory phospholipases A2
in inflammatory diseases and trauma. Biochim Biophys Acta. 2000;1488: 83–90.
doi:10.1016/S1388-1981(00)00112-8

25.   Ernst CM, Kuhn S, Slavetinsky CJ, Krismer B, Heilbronner S, Gekeler C, et al. The
lipid-modifying multiple peptide resistance factor is an oligomer consisting of distinct
interacting   synthase   and   flippase   subunits.   MBio.   2015;6:   1–9.
doi:10.1128/mBio.02340-14

26.   Mazmanian SK. *Staphylococcus aureus* Sortase, an enzyme that anchors surface
proteins   to   the   cell   wall.   Science   1999;285:   760–763.
doi:10.1126/science.285.5428.760

27.   Hendrickx APA, Budzik JM, Oh S-Y, Schneewind O. Architects at the bacterial
surface - sortases and the assembly of pili with isopeptide bonds. Nat Rev Microbiol.
Nature Publishing Group; 2011;9: 166–176. doi:10.1038/nrmicro2520

28.   Qu X, Lehrer RI. Secretory Phospholipase A2 is the principal bactericide for
staphylococci and other Gram-positive bacteria in human tears. Infect Immun.
1998;66: 2791–2797.

29.   Weinrauch Y, Elsbach P, Madsen LM, Foreman A, Weiss J. The potent anti-
Staphylococcus aureus activity of a sterile rabbit inflammatory fluid is due to a 14-kD
phospholipase A2. J Clin Invest. 1996;97: 250–257. doi:10.1172/JCI118399

30.   Lancefield RC. A Serological differentiation of human and other groups of hemolytic
streptococci. J Exp Med. 1933;57: 571–595.

31.   McCarthy M. The lysis of Group A hemolytic streptococci by extracellular enzymes of
Streptomyces albus. II. Nature of the cellular substrate attacked by the lytic Enzymes.
J Exp Med. 1952;96: 569–580.

32.   Van Sorge NM, Cole JN, Kuipers K, Henningham A, Aziz RK, Kasirer-Friede A, et al.
The classical lancefield antigen of group A *Streptococcus* is a virulence determinant
with   implications   for   vaccine   design.   Cell   Host   Microbe.   2014;15:   729–740.
doi:10.1016/j.chom.2014.05.009

33.   Caliot É, Dramsi S, Chapot-Chartier M-P, Courtin P, Kulakauskas S, Péchoux C, et al.
Role of the Group B antigen of *Streptococcus agalactiae*: a peptidoglycan-anchored
polysaccharide involved in cell wall biogenesis. PLoS Pathog. 2012; 8: e1002756.
doi:10.1371/journal.ppat.1002756

34.   Carapetis JR, Steer AC, Mulholland EK, Weber M. The global burden of group A
streptococcal diseases. Lancet Infect Dis. 2005;5: 685–694. doi:10.1016/S1473-
3099(05)70267-X

35.   High KP, Edwards MS, Baker CJ. Group B Streptococcal infections in elderly adults.
Clin Infect Dis. 2005;41: 839–847. doi:10.1086/432804

36.   Edmond KM, Kortsalioudaki C, Scott S, Schrag SJ, Zaidi AK, Cousens S, et al. Group
B streptococcal disease in infants aged younger than 3 months: Systematic review
and meta-analysis. Lancet. 2012;379: 547–556. doi:10.1016/S0140-6736(11)61651-6

37.   Kaul R, McGeer A, Low DE, Green K, Schwartz B, Simor AE, et al. Population-based
surveillance   for   group   A   streptococcal   necrotiziug   fasciitis: Clinical   features,
prognostic indicators, and microbiologic analysis of seventy-seven cases. Am J Med.
1997;103: 18–24. doi:10.1016/S0002-9343(97)00160-5

38.   Davies HD, McGeer A, Schwartz B, Green K, Cann D, Simor AE, et al. Invasive
Group A Streptococcal infections in Ontario , Canada. N Engl J Med. 1996;335: 547–
554.

39.   Attalah HL, Wu Y, Alaoui-El-Azher M, Thouron F, Koumanov K, Wolf C, et al.
Induction of type-IIA secretory phospholipase A2 in animal models of acute lung

injury. Eur Respir J. 2003;21: 1040–1045. doi:10.1183/09031936.03.00093002

40. Le Breton Y, Belew AT, Valdes KM, Islam E, Curry P, Tettelin H, et al. Essential
genes in the core genome of the human pathogen *Streptococcus pyogenes*. Sci Rep.
2015;5: 9838. doi:10.1038/srep09838

41. Fiebig A, Loof TG, Babbar A, Itzek A, Koehorst JJ, Schaap PJ, et al. Comparative
Genomics of *Streptococcus pyogenes* M1 isolates differing in virulence and
propensity to cause systemic infection in mice. Int J Med Microbiol. 2015;305: 532–
543. doi:10.1016/j.ijmm.2015.06.002

42. Peschel A, Otto M, Jack RW, Kalbacher H, Jung G, Götz F. Inactivation of the dlt
operon in *Staphylococcus aureus* confers sensitivity to defensins, protegrins, and
other   antimicrobial   peptides.   J   Biol   Chem.   1999;274:   8405–8410.
doi:10.1074/JBC.274.13.8405

43. Kristian SA, Datta V, Weidenmaier C, Kansal R, Fedtke I, Peschel A, et al. D-
Alanylation of teichoic acids promotes Group A *Streptococcus* antimicrobial peptide
resistance, neutrophil survival, and epithelial cell invasion. J Bacteriol. 2005;187:
6719–6725. doi:10.1128/JB.187.19.6719

44. Le Breton Y, Mistry P, Valdes KM, Quigley J, Kumar N, Tettelin H, et al. Genome-
wide identification of genes required for fitness of Group A *Streptococcus* in human
blood. Infect Immun. 2013;81: 862–875. doi:10.1128/IAI.00837-12

45. Le Breton Y, Belew AT, Freiberg JA, Sundar GS, Islam E, Lieberman J, et al.
Genome-wide discovery of novel M1T1 Group A streptococcal determinants important
for fitness and virulence during soft-tissue infection. PLoS Pathog. 2017;13: 1–28.
doi:10.1371/journal.ppat.1006584

46. Rush JS, Edgar RJ, Deng P, Chen J, Zhu H, van Sorge NM, et al. The molecular
mechanism of N-acetylglucosamine side-chain attachment to the Lancefield Group A
carbohydrate in *Streptococcus pyogenes*. J Biol Chem. 2017;292: jbc.M117.815910.
doi:10.1074/jbc.M117.815910

47. Shapiro HM. Membrane potential estimation by flow cytometry. Methods. 2000;21:
271–279. doi:10.1006/meth.2000.1007

48. Saar-Dover R, Bitler A, Nezer R, Shmuel-Galia L, Firon A, Shimoni E, et al. D-
Alanylation of lipoteichoic acids confers resistance to cationic peptides in Group B
Streptococcus by increasing the cell wall density. PLoS Pathog. 2012;8.
doi:10.1371/journal.ppat.1002891

49. Albertí S, Ashbaugh CD, Wessels MR. Structure of the *has* operon promoter and
regulation of hyaluronic acid capsule expression in Group A *Streptococcus*. Mol
Microbiol. 1998;28: 343–353. doi:10.1046/j.1365-2958.1998.00800.x

50. Campbell J, Singh AK, Santa Maria JP, Kim Y, Brown S, Swoboda JG, et al. Synthetic
lethal compound combinations reveal a fundamental connection between wall teichoic
acid and peptidoglycan biosyntheses in *Staphylococcus aureus*. ACS Chem Biol.
2011;6: 106–116. doi:10.1021/cb100269f

51. Weinrauch Y, Abad C, Liang NS, Lowry SF, Weiss J. Mobilization of potent plasma
bactericidal activity during systemic bacterial challenge: role of Group IIA
Phospholipase A2. J Clin Invest. 1998;102: 633–638. doi:10.1172/JCI3121

52. Grönroos JO, Salonen JH, Viander M, Nevalainen TJ, Laine VJO. Roles of group IIA
phospholipase A2 and complement in killing of bacteria by acute phase serum. Scand
J Immunol. 2005;62: 413–419. doi:10.1111/j.1365-3083.2005.01678.x

53. Berends ETM, Dekkers JF, Nijland R, Kuipers A, Soppe JA, van Strijp JAG, et al.
Distinct localization of the complement C5b-9 complex on Gram-positive bacteria. Cell
Microbiol. 2013;15: 1955–1968. doi:10.1111/cmi.12170

54. Raz A, Tanasescu AM, Zhao AM, Serrano A, Alston T, Sol A, et al. *Streptococcus*
pyogenes sortase mutants are highly susceptible to killing by host factors due to
aberrant envelope physiology. PLoS One. 2015;10: 1–30.
doi:10.1371/journal.pone.0140784

55. Kawai Y, Marles-Wright J, Cleverley RM, Emmins R, Ishikawa S, Kuwano M, et al. A
widespread family of bacterial cell wall assembly proteins. EMBO J. 2011;30: 4931–
4941. doi:10.1038/emboj.2011.358

56. Dengler V, Meier PS, Heusser R, Kupferschmied P, Fazekas J, Friebe S, et al.
Deletion of hypothetical wall teichoic acid ligases in *Staphylococcus aureus* activates
the cell wall stress response. FEMS Microbiol Lett. 2012;333: 109–120.
doi:10.1111/j.1574-6968.2012.02603.x

57. Chan YGY, Frankel MB, Dengler V, Schneewind O, Missiakasa D. *Staphylococcus*
aureus mutants lacking the lytr-cpsa-Psr family of enzymes release cell wall teichoic
acids into the extracellular medium. J Bacteriol. 2013;195: 4650–4659.
doi:10.1128/JB.00544-13

58. Chan YGY, Kim HK, Schneewind O, Missiakas D. The capsular polysaccharide of
Staphylococcus aureus is attached to peptidoglycan by the LytR-CpsA-Psr (LCP)
family of enzymes. J Biol Chem. 2014;289: 15680–15690.
doi:10.1074/jbc.M114.567669

59. Zilla ML, Chan YGY, Lunderberg JM, Schneewind O, Missiakas D. LytR-CpsA-Psr
enzymes as determinants of *Bacillus anthracis* secondary cell wall polysaccharide
assembly. J Bacteriol. 2015;197: 343–353. doi:10.1128/JB.02364-14

60. Harrison J, Lloyd G, Joe M, Lowary TL, Reynolds E, Walters-Morgan H, et al. Lcp1 is
a phosphotransferase responsible for ligating arabinogalactan to peptidoglycan in
Mycobacterium tuberculosis. MBio. 2016;7: 1–12. doi:10.1128/mBio.00972-16

61. Lazarevic V, Margot P, Soldo B, Karamata D. Sequencing and analysis of the *Bacillus*
subtilis lytRABC divergon: a regulatory unit encompassing the structural genes of the
N-acetylmuramoyl-L-alanine amidase and its modifier. J Gen Microbiol. 1992;138:
1949–1961. doi:10.1099/00221287-138-9-1949

62. Chatfield CH, Koo H, Quivey RG. The putative autolysin regulator LytR in
Streptococcus mutans plays a role in cell division and is growth-phase regulated.
Microbiology. 2005;151: 625–631. doi:10.1099/mic.0.27604-0

63. Foreman-Wykert AK, Weinrauch Y, Elsbach P, Weiss J. Cell-wall determinants of the
bactericidal action of group IIA phospholipase A2 against Gram-positive bacteria. J
Clin Invest. 1999;103: 715–21. doi:10.1172/JCI5468

64. Minami M, Ichikawa M, Ohta M, Hasegawa T. The cell envelope-associated protein,
LytR, regulates the cysteine protease SpeB in *Streptococcus pyogenes*. Apmis.
2012;120: 417–426. doi:10.1111/j.1600-0463.2011.02847.x

65. McCarty M. Variation in the group-specific carbohydrate of Group A Streptococci. II.
studies on the chemical basis for serological specificity of the carbohydrates. J Exp
Med. 1956;104: 629–643.

66. Swanson J, Gotschlich EC. Electron microscopy studies on Streptococci II. Group A
Carbohydrate. J Exp Med. 1973;138: 245–258.

67. Wagner M, Wagner B, Kubín VR. Immunoelectron microscopic study of the location of
group-specific and type-specific polysaccharide antigens on isolated walls of group B
streptococci. J Gen Microbiol. 1980;120: 369–76. Available:
http://www.ncbi.nlm.nih.gov/pubmed/7014772

68. Sochacki KA, Barns KJ, Bucki R, Weisshaar JC. Real-time attack on single
Escherichia coli cells by the human antimicrobial peptide LL-37. Proc Natl Acad Sci.

2011;108: E77–E81. doi:10.1073/pnas.1101130108

69.   Rangarajan N, Bakshi S, Weisshaar JC. Localized permeabilization of *E. coli*
membranes by the antimicrobial peptide Cecropin A. Biochemistry. 2013;52: 6584–
6594. doi:10.1021/bi400785j

70.   Mileykovskaya E, Dowhan W. Cardiolipin membrane domains in prokaryotes and
eukaryotes. Biochim Biophys Acta - Biomembr. 2009;1788: 2084–2091.
doi:10.1016/j.bbamem.2009.04.003

71.   Rosch JW, Hsu FF, Caparon MG. Anionic lipids enriched at the ExPortal of
Streptococcus pyogenes. J Bacteriol. 2007;189: 801–6. doi:10.1128/JB.01549-06

72.   Henningham A, Davies MR, Uchiyama S, Van Sorge NM, Lund S, Chen KT, et al.
Virulence role of the GlcNAc side chain of the Lancefield cell wall carbohydrate
antigen in non-M1-serotype Group A *Streptococcus*. MBio. 2018;9: 1–12.

73.   Ghomashchi F, Brglez V, Payré C, Jeammet L, Bezzine S, Gelb MH, et al.
Preparation of the full set of recombinant mouse- and human-secreted
phospholipases A2. 1st ed. Methods in Enzymology. Elsevier Inc.; 2017.
doi:10.1016/bs.mie.2016.10.034

74.   Zhang X, de Maat V, Guzman Prieto AM, Prajsnar TK, Bayjanov JR, de Been M, et al.
Fitness determinants of vancomycin-resistant *Enterococcus faecium* during growth in
human serum. MBC genomics. BMC Genomics; 2017;18: 1–12. doi:10.1101/101329

75.   Goecks J, Nekrutenko A, Taylor J, Galaxy Team T. Galaxy: a comprehensive
approach for supporting accessible, reproducible, and transparent computational
research in the life sciences. Genome Biol. 2010;11: R86. doi:10.1186/gb-2010-11-8-
r86

76.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9: 357–359. doi:10.1038/nmeth.1923

77.   Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Gets G, et al.
Integrative Genomics Viewer. Nat Biotechnol. 2013;29: 24–26. doi:10.1038/nbt0111-
24

78.   Baldi P, Long AD. A Bayesian framework for the analysis of microarray expression
data: regularized t-test and statistical inferences of gene changes. Bioinformatics.
2001;17: 509–519. doi:10.1093/bioinformatics/17.6.509

79.   Carvalho F, Atilano ML, Pombinho R, Covas G. L-Rhamnosylation of *Listeria*
monocytogenes wall teichoic acids promotes resistance to antimicrobial peptides by
delaying   Interaction   with   the   membrane.   2015;   1–29.
doi:10.1371/journal.ppat.1004919

80.   Yokogawa K, Kawata S, Takemura T, Yoshimura Y. Purification and properties of lytic
Enzymes from *Streptomyces globisporus* 1829. Agr Biol Chem. 1975;39: 1533–1543.

81.   Calandra GB, Cole RM. Lysis and protoplast formation of group B streptococci by
mutanolysin. Infect Immun. 1980;28: 1033–1037.

Figure legends

Fig 1. GAS M1T1 is killed by recombinant hGIIA, which is enhanced in human serum.

(A) GAS M1T1 5448 is killed by hGIIA in a dose-dependent manner. Mutation of *srtA*

renders 5448 more susceptible to hGIIA killing. (B) A heat-labile factor in serum enhances

the ability of hGIIA to kill GAS. Data represent mean +/- SD of three independent

experiments. *, $p > 0.05$; ***, $p > 0.001$.

Fig 2. Identifying GAS mutants with different hGIIA susceptibility by Tn-seq analysis.

(A) Average log-fold change of transposon insertions in genes of the hGIIA-treated group

versus the control group. Grey dots represent genes without significant fold change after

hGIIA treatment. Green and orange dots represent genes with significantly increased and

reduced transposon insertions after hGIIA exposure, respectively. Significant hits have a

calculated BH corrected p <0.05. (B) Circos representation of the average transposon

insertions of genes within the GAC gene cluster. All genes within this cluster, except for

gacI, *gacJ*, and *gacH*, were previously identified as essential [40]. For *gacI* and *gacH*, the

fold change shown is significant (BH corrected $p < 0.05$), whereas for *gacJ* the fold change

is not significant (BH corrected $p = 0.16$).

Fig 3. Mutation of *dltA* and *lytR* renders GAS more susceptible to hGIIA, whereas

mutation of *gacI* increases hGIIA resistance in multiple GAS serotypes. Deletion of (A)

dltA or (B) *lytR* increases GAS susceptibility to hGIIA-mediated killing in a concentration-

dependent manner. Deletion of *gacI* renders GAS more resistant to hGIIA-mediated killing

as shown for (C) 5448 and (D) other tested GAS serotypes. Data represent mean +/- SD of

three independent experiments. *, $p > 0.05$; **, $p > 0.01$; ***, $p > 0.001$.

Fig 4. Lack of the GAC GlcNAc side chain delays hGIIA cell wall translocation. (A)

Detection of hGIIA binding to GAS 5448 by flow cytometry. (B) Deletion of *gacI* affects

surface charge of GAS as determined in cationic cytochrome c binding assay. (C)

Localization of hGIIA (H48Q) on GAS 5448 was analyzed by fluorescence microscopy (+)

and quantified based on analysis of 10 fields including 307 stained cells from two separate

experiments. As control, H48Q hGIIA was omitted (-). hGIIA was detected with a mouse

anti-human hGIIA monoclonal antibody or an IgG1 isotype as negative control.

Representative cells are shown. (D) Effect of hGIIA on GAS membrane potential after 2 hour

incubation. Decreased PerCP signal indicates a disrupted membrane potential. (E) SYTOX green uptake over time by GAS strains or (F) GAS protoplasts after incubation with recombinant hGIIA. Data represent mean +/- SD of three independent experiments. ns = not significant, *, $p > 0.05$; **, $p > 0.01$; ***, $p > 0.001$.

Fig 5. GBS lacking the GBC are resistant to hGIIA. (A) HGIIA kills GBS strain NEM316 WT but not Δ*gbcO* in a concentration-dependent manner and phenotype is restored in complemented strain ΔgbcO/pTCV. The killing is represented as the percentage of CFU surviving after hGIIA exposure compared to the inoculum. GBS killing is prevented when (B) exposed to catalytically inactive hGIIA H48Q and (C) by the hGIIA-specific inhibitor LY311727. (D) Treatment of NEM316 WT with the *gbcO*-type inhibitor tunicamycin reproduces the Δ*gbcO* phenotype with regard to hGIIA-mediated killing GBS more resistant to hGIIA-mediated killing. (E) Visualization of bacteria-bound hGIIA H48Q to GBS NEM316 by fluorescence microscopy (+). As control, H48Q hGIIA mutant protein was omitted (-). hGIIA was detected with a mouse anti-human hGIIA monoclonal antibody. An irrelevant IgG1 isotype antibody served as negative control. The cell wall was labeled with Ester-350, and newly formed septa were visualized with fluorescently labeled vancomycin (Vanc-FL), which stains sites of peptidoglycan insertion. Shown are representative cells. Quantification of hGIIA binding to polar or non-polar regions of GBS are based on analysis of 12 fields including 578 stained cells from two separate experiments. For all other panels, data represent mean +/- SD of three independent experiments.

Fig S1. Endogenous hGIIA in normal human serum does not affect GAS growth. Addition of hGIIA inhibitor LY311727 does not influence GAS growth in human serum. Data represent mean +/- SD of three independent experiments, ns = not significant.

Fig S2. Biased transposon insertions in M5005_Spy_1390. Unusual high number of transposon insertions at one location in the gene M5005_Spy_1390.

Fig S3. Tn-seq results. Circos respresentation of the Tn-seq data. Each bar in the inner two circles, where blue is control and red the hGIIA treated, represent the average RKPM value of a gene. The following to circles represent the BH corrected p-value and the fold change in log of the hGIIA treated samples vs control samples. Red bars indicate a fold a respective fold decrease and green bars a respective fold increase of transposon insertions. The gene highlighted in red is M5005_Spy_1390, which showed significant fold change due to unusual high transposon insertions at one specific point in the gene.

Fig S4. Three additional time points for DiOC$_2$(3) measurements. Membrane potential after (A) 5 minutes, (B) 30 minutes and (C) 60 minutes of hGIIA stress. Data represent mean +/- SD of three independent experiments. *, $p > 0.05$; **, $p > 0.01$; ***, $p > 0.001$.

Fig S5. Kinetics of the SYTOX influx in intact GAS strains. SYTOX influx measured over 120 minutes when GAS strains are incubated with (A) 5, (B) 0.5, (C) 0.1, (D) 0.01, (E) 0.001, and (F) 0 µg/ml hGIIA. Data represent mean +/- SD of three independent experiments.

Fig S6. Kinetics of the SYTOX influx in protoplast GAS strains. SYTOX influx measured over 120 minutes when protoplast GAS strains are incubated with (A) 5, (B) 0.5, (C) 0.1, (D) 0.01, (E) 0.001, and (F) 0 µg/ml hGIIA. Data represents mean +/- SD of three independent experiments.

Acknowledgements

The authors would like to thank Dr. Michel-Yves Mistou (INRA, Jouy-en-Josas, France) for providing his *gbcO* mutant and complemented GBS strains.

Funding statement

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. N.M.v.S and V.P.v.H were supported by a VIDI grant (91713303) from the Dutch Scientific Organization (NWO).

Figure 1

A

B

Figure 2

A

B

Figure 3

Figure 4

Figure 5