Network hubs in root-associated fungal metacommunities

Hirokazu Toju[1,2*], Akifumi S. Tanabe[3], Hirotoshi Sato[4]

*Correspondence: toju.hirokazu.4c@kyoto-u.ac.jp

[1]Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2113, Japan

[2]Precursory Research for Embryonic Science and Technology (PRESTO), Japan Science and Technology Agency, Kawaguchi, Saitama 332-0012, Japan

[3]Faculty of Science and Technology, Ryukoku University, 1-5 Yokotani, Seta Oe-cho, Otsu, Shiga 520-2194, Japan

[4]Graduate School of Human and Environmental Studies, Kyoto University, Sakyo, Kyoto 606-8501, Japan

This article includes 5 Figures, 1 Table, and 9 additional files.

Abstract

Background: Although a number of recent studies have uncovered remarkable diversity of microbes associated with plants, understanding and managing dynamics of plant microbiomes remain major scientific challenges. In this respect, network analytical methods have provided a basis for exploring "hub" microbial species, which potentially organize community-scale processes of plant–microbe interactions.

Methods: By compiling Illumina sequencing data of root-associated fungi in eight forest ecosystems across the Japanese Archipelago, we explored hubs within "metacommunity-scale" networks of plant–fungus associations. In total, the metadata included 8,080 fungal operational taxonomic units (OTUs) detected from 227 local populations of 150 plant species/taxa.

Results: Few fungal OTUs were common across all the eight forests. However, in each metacommunity-scale network representing northern four localities or southern four localities, diverse mycorrhizal, endophytic, and pathogenic fungi were classified as "metacommunity hubs", which were detected from diverse host plant taxa throughout a climatic region. Specifically, *Mortierella* (Mortierellales), *Cladophialophora* (Chaetothyriales), *Ilyonectria* (Hypocreales), *Pezicula* (Helotiales), and *Cadophora* (incertae sedis) had broad geographic and host ranges across the northern (cool-temperate) region, while *Saitozyma*/*Cryptococcus* (Tremellales/Trichosporonales) and *Mortierella* as well as some arbuscular mycorrhizal fungi were placed at the central positions of the metacommunity-scale network representing warm-temperate and subtropical forests in southern Japan.

Conclusions: The network theoretical framework presented in this study will help us explore prospective fungi and bacteria, which have high potentials for agricultural application to diverse plant species within each climatic region. As some of those fungal taxa with broad geographic and host ranges have been known to increase the growth and pathogen resistance of host plants, further studies elucidating their functional roles are awaited.

Keywords: agriculture; biodiversity; ecosystem restoration; host specificity or preference; latitudinal gradients; metacommunities; microbial inoculation; network hubs; plant–fungus

47   interactions; mycorrhizal and endophytic symbiosis.

Background

Below-ground fungi in the endosphere and rhizosphere are key drivers of terrestrial

ecosystem processes [1-4]. Mycorrhizal fungi, for example, are important partners of most

land plant species, enhancing nutritional conditions and pathogen resistance of host plants

[5-7]. In reward for the essential physiological services, they receive ca. 20% of net

photosynthetic products from plants [8, 9]. Recent studies have also indicated that diverse

taxonomic groups of endophytic fungi (e.g., endophytic fungi in the ascomycete orders

Helotiales and Chaetothyriales) commonly interact with plant roots, providing soil

nitrogen/phosphorous to their hosts [10-14], converting organic nitrogen into inorganic forms

in the rhizosphere [15], and increasing plants' resistance to environmental stresses [16-18].

Because of their fundamental roles, below-ground fungi have been considered as prospective

sources of ecosystem-level functioning in forest management, agriculture, and ecosystem

restoration [17-20]. However, due to the exceptional diversity of below-ground fungi [21-23]

and the extraordinary complexity of below-ground plant–fungus interactions [24-26], we are

still at an early stage of managing and manipulating plant-associated microbiomes [27-29].

In disentangling complex webs of below-ground plant–fungus associations, network

analyses, which have been originally applied to human relations and the World-Wide Web

[30, 31], provide crucial insights. By using network analytical tools, we can infer how plant

species in a forest, grassland, or farmland are associated with diverse taxonomic and

functional groups of fungi [24, 32-34]. Such information of network structure (topology) can

be used to identify "hub" species, which are placed at the center of a network depicting

multispecies host–symbiont associations [35] (cf. [34, 36, 37]). Those hubs with broad

host/symbiont ranges are expected to play key roles by mediating otherwise discrete

ecological processes within a community [19, 24]. For example, although arbuscular

mycorrhizal and ectomycorrhizal symbioses have been considered to involve distinct sets of

plant and fungal lineages [38] (but see [39, 40]), hub endophytic fungi with broad host ranges

may mediate indirect interactions between arbuscular mycorrhizal and ectomycorrhizal plant

species through below-ground mycelial connections. As information of plant-associated

fungal communities is now easily available with high-throughput DNA sequencing

technologies [1, 21, 22], finding hub microbial species out of hundreds or thousands of

species within a network has become an important basis for understanding and predicting

ecosystem-scale phenomena.

Nonetheless, given that fungi can disperse long distances with spores, conidia, and

propagules [41-44], information of local-scale networks alone does not provide thorough

insights into below-ground plant–fungus interactions in the wild. In other words, no forests,

grasslands, and farmlands are free from perturbations caused by fungi immigrating from other

localities [45-49]. Therefore, to consider how local ecosystem processes are interlinked by

dispersal of fungi, we need to take into account "metacommunity-scale" networks of plant–

fungus associations [35]. Within a dataset of multiple local communities (e.g., [25]), fungal

species that occur in multiple localities may interlink local networks of plant–fungus

associations. Among them, some species that not only have broad geographic ranges but also

are associated with diverse host plant species would be placed at the core positions of a

metacommunity-scale network [35]. Such "metacommunity hub" fungi would be major

drivers of the synchronization and restructuring of local ecosystem processes (*sensu* [50]),

and hence their functional roles need to be investigated with priority [35]. Moreover, in the

screening of mycorrhizal and endophytic fungi that can be used in agriculture and ecosystem

restoration programs [17, 20, 51], analytical pipelines for identifying metacommunity hubs

will help us explore species that are potentially applied (inoculated) to diverse plant species

over broad geographic ranges of farmlands, forests, or grasslands. Nonetheless, despite the

potential importance of metacommunity hubs in both basic and applied microbiology, few

studies have examined metacommunity-level networks of plant–symbiont associations.

By compiling Illumina sequencing datasets of root-associated fungi [52], we herein

inferred a metacommunity-level network of below-ground plant–fungus associations and

thereby explored metacommunity hubs. Our metadata consisted of plant–fungus association

data in eight forest localities across the entire range of the Japanese Archipelago, including

150 plant species/taxa and 8,080 fungal operational taxonomic units (OTUs) in temperate and

subtropical regions. Based on the information of local- and metacommunity-level networks,

each of the fungal OTUs was evaluated in light of its topological positions. We then

examined whether fungal OTUs placed at the core of local-level plant–fungus networks could

play key topological roles within the metacommunity-level network. Overall, this study

uncover how diverse taxonomic groups of mycorrhizal and endophytic fungi can form

metacommunity-scale networks of below-ground plant–fungus associations, providing a basis

for analyzing complex spatial processes of species-rich host–microbe systems.

Methods

Terminology

While a single type of plant–fungus interactions is targeted in each of most mycological

studies (e.g., arbuscular mycorrhizal symbiosis or ectomycorrhizal symbiosis), we herein

analyze the metadata including multiple categories of below-ground plant–fungus

associations [52]. Because arbuscular mycorrhizal, ectomycorrhizal, and endophytic fungi, for

example, vary in their microscopic structure within plant tissue [38], it is impossible to

develop a general criterion of mutualistic/antagonistic interactions for all those fungal

functional groups. Therefore, we used the phrase "associations" instead of "interactions"

throughout the manuscript when we discuss patterns detected based on the Illumina

sequencing metadata of root-associated fungi. Consequently, our results represented not only

mutualistic or antagonistic interactions but also neutral or commensalistic interactions [24, 53,

54]. Our aim in this study is to gain an overview of the metacommunity-scale plant–fungus

associations, while the nature of respective plant–fungus associations should be evaluated in

future inoculation experiments.

Data

We compiled the Illumina (MiSeq) sequencing data collected in a previous study [52], in

which community-scale statistical properties of below-ground plant–fungus associations were

compared among eight forest localities (four cool-temperate, one warm-temperate, and three

subtropical forests) across the entire range of the Japanese Archipelago (45.042–24.407 ºN;

Fig. 1) (DDBJ Sequence Read Archives accession: DRA006339). In each forest, 2-cm

segment of terminal roots were sampled from 3-cm below the soil surface at 1-m horizontal

intervals [52]. Those root samples were collected irrespective of their morphology and

mycorrhizal type: hence, the samples as a whole represented below-ground relative

abundance of plant species in each forest community. Based on the sequences of the genes

encoding the large subunit of ribulose-1,5-bisphosphate carboxylase (*rbcL*) and the internal

transcribed spacer 1 (ITS1) of the ribosomal RNA region, host plant species were identified,

although there were plant root samples that could not be identified to species with the *rbcL*

and ITS1 regions [52].

The Illumina sequencing reads of the fungal ITS1 region were processed as detailed in

the data-source study [52]. The primers used were designed to target not only Ascomycota

and Basidiomycota but also diverse non-Dikarya (e.g., Glomeromycota) taxa [55]. In most

studies analyzing community structure of Ascomycota and Basidiomycota fungi, OTUs of the

ITS region are defined with a cut-off sequence similarity of 97% [22, 56, 57] (see also [58]).

Meanwhile, Glomeromycota fungi generally have much higher intraspecific ITS-sequence

variation than other taxonomic groups of fungi [59]. Consequently, we used 97% and 94%

cut-off sequence similarities for defining non-Glomeromycota and Glomeromycota fungal

OTUs, respectively [52]. The OTUs were then subjected to reference database search with the

query-centric auto-*k*-nearest-neighbor algorithm [60, 61] and subsequent taxonomic

assignment with the lowest common ancestor algorithm [62]. Based on the inferred taxonomy,

the functional group of each fungal OTU was inferred using the program FUNGuild 1.0 [63].

After a series of bioinformatics and rarefaction procedures, 1,000 fungal ITS reads were

obtained from each of the 240 samples collected in each forest locality (i.e., 1,000 reads × 240

samples × 8 sites). A sample (row) × fungal OTU (column) data matrix, in which a cell entry

depicted the number of sequencing reads of an OTU in a sample, was obtained for each local

forest ("sample-level" matrix) (Additional file 1: Data S1). Each local sample-level matrix

was then converted into a "species-level" matrix, in which a cell entry represented the number

of root samples from which associations of a plant species/taxa (row) and a fungal OTU

(columns) was observed: 17–55 plant species/taxa and 1,149–1,797 fungal OTUs were

detected from the local species-level matrices (Additional file 2: Data S2). In total, the

matrices included 150 plant species/taxa and 8,080 fungal OTUs (Additional file 3: Data S3).

Local networks

Among the eight forest localities, variation in the order-level taxonomic compositions were

examined with the permutational analysis of variance (PERMANOVA; [64]) and the

permutational analysis for the multivariate homogeneity of dispersions (PERMDISP; [65])

with the "adonis" and "betadisper" functions of the vegan 2.4-3 package [66] of R 3.4.1 [67],

respectively. The β-diversity values used in the PERMANOVA and PERMDISP analyses

were calculated with the "Bray-Curtis" metric based on the sample-level matrices (Additional

file 1: Data S1). Note that the "Raup-Crick" β-diversity metric [68], which controls

α-diversity in community data but requires computationally intensive randomization, was not

applicable to our large metadata. Geographic variation in the compositions of fungal

functional groups was also evaluated by PERMANOVA and PERMDISP analyses.

For each of the eight local forests, the network structure of below-ground plant–fungus

associations was visualized based on the species-level matrix (Additional file 2: Data S2)

using the program GePhi 0.9.1 [69] with the "ForceAtlas2" layout algorithm [70]. Within the

networks, the order-level taxonomy of fungal OTUs was highlighted.

To evaluate host ranges of each fungal OTU in each local forest, we first calculated the *d'*

metric of interaction specificity [71]. However, estimates of the *d'* metric varied considerably

among fungal OTUs observed from small numbers of root samples (Additional file 4; Figure

S1) presumably due to overestimation or underestimation of host preferences for those rare

OTUs. Therefore, we scored each fungal OTU based on their topological positions within

each local network by calculating network centrality indices (degree, closeness, betweenness,

and eigenvector centralities metrics of network centrality; [31]). Among the centrality metrics,

betweenness centrality, which measures the extent to which a given nodes (species) is located

within the shortest paths connecting pairs of other nodes in a network [72], is often used to

explore organisms with broad host or partner ranges [35]. Thus, in each local network, fungal

OTUs were ranked based on their betweenness centrality scores (local betweenness).

Metacommunity-scale network

By compiling the species-level matrices of the eight local forests, the topology of the

metacommunity-scale network of plant–fungus associations was inferred. In general, species

interaction (association) networks of local communities can be interconnected by species that

appear in two or more local networks, thereby merged into a metacommunity-scale network

[35]. In our data across the eight local forests, 2,109 OTUs out of the 8,080 fungal OTUs

appeared in two or more localities. Therefore, we could infer the topology of a

metacommunity-scale network, in which the eight local networks were combined by the

2,109 fungal OTUs. In the metacommunity-scale network, plant species/taxa observed in

different localities were treated as different network nodes because our purpose in this study

was to explore fungi that potentially play key roles in synchronizing local ecosystem

processes [35]. In total, 227 plant nodes representing local populations of 150 plant

species/taxa were included in the metacommunity-scale network.

We then screened for fungal OTUs with broad geographic and host ranges based on the

betweenness centrality scores of respective fungal OTUs within the metacommunity network

(metacommunity betweenness, B_{meta}). In general, species with highest metacommunity

betweenness scores not only occur in local communities over broad biotic/abiotic

environmental conditions but also are associated with broad ranges of host/partner species

[35]. Possible relationship between local- and metacommunity-scale topological roles was

then examined by plotting local and metacommunity betweenness scores (B_{local} and B_{meta}) of

each fungal OTUs on a two-dimensional surface. To make the betweenness scores vary from

0 to 1, betweenness centrality of a fungal OTU i was standardized in each of the local- and

metacommunity-scale networks as follows:

$$B'_{\mathrm{local},i} = \frac{B_{\mathrm{local},i} - \min(B_{\mathrm{local}})}{\max(B_{\mathrm{local}}) - \min(B_{\mathrm{local}})} \quad \text{and} \quad B'_{\mathrm{meta},i} = \frac{B_{\mathrm{meta},i} - \min(B_{\mathrm{meta}})}{\max(B_{\mathrm{meta}}) - \min(B_{\mathrm{meta}})},$$

where $B_{\text{local},i}$ and $B_{\text{meta},i}$ were raw estimates of local- and metacommunity-scale betweenness of a fungal OTU i, and min() and max() indicated minimum and maximum values, respectively. For local betweenness of each OTU, a mean value across local networks was subsequently calculated ($\overline{B}'_{\text{local},i}$): the local communities from which a target OTU was absent was omitted in the calculation of mean local betweenness. On the two-dimensional surface, the OTUs were then classified into four categories: metacommunity hubs having high betweenness in both local- and metacommunity-scale networks ($\overline{B}'_{\text{local},i} \geq 0.5$; $B'_{\text{meta},i} \geq 0.5$), metacommunity connectors that had broad geographic ranges but displayed low local betweenness ($\overline{B}'_{\text{local},i} < 0.5$; $B'_{\text{meta},i} \geq 0.5$), local hubs that had high betweenness in local networks but not in the metacommunity-scale network ($\overline{B}'_{\text{local},i} \geq 0.5$; $B'_{\text{meta},i} < 0.5$), and peripherals with low betweenness at both local and metacommunity levels ($\overline{B}'_{\text{local},i} < 0.5$; $B'_{\text{meta},i} < 0.5$) [35]. Approximately, 1–2% of fungal OTUs show betweenness scores higher than 0.5 in each local or metacommunity network, while the threshold value can be changed depending on the purpose of each study [35].

In addition to metacommunity hubs within the metacommunity-scale network representing all the eight localities, those within the metacommunity-scale network representing northern (sites 1–4) or southern (sites 5–8) four localities were also explored. This additional analysis allowed us to screen for fungal OTUs that potentially adapted to broad ranges of biotic and abiotic environments within northern (cool-temperate) or southern (warm-temperate or subtropical) part of Japan.

Results

Local networks

Among the eight forest localities, order-level taxonomic compositions of fungi varied significantly (PERMANOVA; $F_{\text{model}} = 35.7$, $P < 0.001$), while the differentiation of community structure was attributed at least partly to geographic variation in among-sample dispersion (PERMDISP; $F = 13.2$, $P < 0.001$) (Fig. 2a). Compositions of fungal functional groups were also differentiated among the eight localities (PERMANOVA; $F_{\text{model}} = 34.9$, $P <$

0.001), while within-site dispersion was significantly varied geographically (PERMDISP; $F =$

9.2, $P < 0.001$) (Fig. 2b). The proportion of ectomycorrhizal fungal orders, such as Russulales,

Thelephorales, and Sebacinales, was higher in temperate forests than in subtropical forests,

while that of arbuscular mycorrhizal fungi increased in subtropical localities (Fig. 2). The

proportion of the ascomycete order Helotiales, which has been known to include not only

ectomycorrhizal but also endophytic, saprotrophic, and ericoid mycorrhizal fungi [73], was

higher in northern localities. In contrast, Diaporthales, which has been considered as

predominantly plant pathogenic taxon [74] (but see [75]), was common in subtropical forests

but not in others.

In each of the eight local networks depicting plant–fungus associations, some fungal

OTUs were located at the central positions of the network, while others are distributed at

peripheral positions (Additional file 5; Figure S2). Specifically, fungal OTUs belonging to the

ascomycete orders Chaetothyriales (e.g., *Cladophialophora* and *Exophiala*) and Helotiales

(e.g., *Rhizodermea*, *Pezicula*, *Rhizoscyphus*, and *Leptodontidium*) as well as some *Mortierella*

OTUs had high betweenness centrality scores in each of the cool-temperate forests (Fig. 3a-b).

In contrast, arbuscular mycorrhizal fungi (Glomeromycota) were common among OTUs with

highest betweenness scores in subtropical forests (Fig. 3a-c). Some fungi in the ascomycete

order Hypocreales (e.g., *Trichoderma*, *Ilyonectria*, *Simplicillium*, and *Calonectria*) also had

high betweenness scores in some temperate and subtropical forests (Fig. 3b).

Metacommunity-scale network

In the metacommunity-scale network representing the connections among the eight local

networks, not only arbuscular mycorrhizal but also saprotrophic/endophytic fungi were placed

at the central topological positions (Fig. 4; Additional file 6; Figure S3). Among

non-Glomeromycota OTUs, *Mortierella* (Mortierellales), *Cryptococcus* (Trichosporonales;

the Blast top-hit fungus in the NCBI database was recently moved to *Saitozyma*

(Tremellales); [76]), *Malassezia* (Malasseziales), *Oidiodendron* (incertae sedis), *Trichoderma*

(Hypocreales), and a fungus distantly allied to *Melanconiella* (Diaporthales) displayed highest

metacommunity betweenness (Table 1). Among the OTUs with high metacommunity

betweenness, only a *Mortierella* OTU was designated as a metacommunity hub (i.e., $\overline{B}'_{\text{local},i}$

≥ 0.5; $B'_{\text{meta},i} \geq 0.5$) and others had low betweenness scores at the local community level

($\overline{B}'_{\text{local},i} < 0.5$; Fig. 5a).

In the metacommunity-scale network representing the four cool-temperate forests (sites

1–4), many saprotrophic/endophytic fungal OTUs were associated with diverse plant

species/taxa, located at the central topological positions within the network topology

(Additional file 7; Figure S4; Fig. 5b). The list of these fungi with high metacommunity

betweenness involved OTUs in the genera *Mortierella*, *Cladophialophora* (Chaetothyriales),

Pezicula (Helotiales), and *Oidiodendron* as well as OTUs allied to *Ilyonectria protearum*

(Nectriales) and *Cadophora orchidicola* (Helotiales) (Table 1). Most of those fungal OTUs

also had high metacommunity betweenness, designated as metacommunity hubs (Fig. 5b).

In the metacommunity-scale network consisting of the warm-temperate and subtropical

forests (sites 5–8), arbuscular mycorrhizal and saprotrophic/endophytic fungi were placed at

the hub positions (Additional file 8; Figure S5; Fig. 5c). The list of non-Glomeromycota

OTUs with highest metacommunity betweenness included *Saitozyma* (*Cryptococcus*),

Mortierella, *Trichoderma*, and *Tomentella* as well as OTUs allied to *Cladophialophora*,

Scleropezicula (Helotiales), *Melanconiella* (Diaporthales), and *Rhexodenticula* (incertae

sedis) (Table 1). Among the taxa, *Saitozyma* and *Mortierella* included OTUs classified as

metacommunity hubs (Fig. 5c; Table 1). In an additional analysis of a metacommunity-scale

network including only the three subtropical forests (sites 6-8), similar sets of fungal taxa

were highlighted (Additional file 9; Table S1). The detailed information of the network index

scores examined in this study is provided in Data S3 (Additional file 3: Data S3).

Discussion

Based on the metadata of root-associated fungi across the Japanese Archipelago, we herein

inferred the structure of a network representing metacommunity-scale associations of 150

plant species/taxa and 8,080 fungal OTUs. Our analysis targeted diverse functional groups of

fungi such as arbuscular mycorrhizal, ectomycorrhizal, ericoid-mycorrhizal,

saprotrophic/endophytic, and pathogenic fungi, which have been analyzed separately in most

previous studies on plant–fungus networks. The comprehensive analysis of below-ground

plant–fungus associations allowed us to explore metacommunity hub fungi, which not only

occurred over broad geographic ranges but also had broad host ranges in respective local

communities. Consequently, this study highlights several taxonomic groups of fungi

potentially playing key roles in synchronizing metacommunity-scale processes of temperate

and/or subtropical forests.

In the metacommunity-scale network representing all the eight local forests (Fig. 4),

fungi in several saprotrophic or endophytic taxa showed higher betweenness centrality scores

than other fungi (Table 1). *Mortierella* is generally considered as a saprotrophic lineage [77]

but it also includes fungi contributing to the growth and pathogen resistance of plants [78-80].

A phosphate solubilizing strain of *Mortierella*, for example, increases shoot and root growth

of host plants under salt stress, especially when co-inoculated with an arbuscular mycorrhizal

fungus [78]. In addition, polyunsaturated fatty acids produced by some *Mortierella* species

are known to increase resistance of plants against phytopathogens [79, 80]. Fungi in the genus

Trichoderma are commonly detected and isolated from the rhizosphere [77, 81]. Many of

them inhibit the growth of other fungi, often used in the biological control of phytopathogens

[82-84]. Some of them are also reported to suppress root-knot nematodes [85] or to promote

root growth [86]. The analysis also highlighted basidiomycete yeasts in the genus *Saitozyma*

or *Cryptococcus* (teleomorph = *Filobasidiella*), which are often isolated from soil [22, 87] as

well as both above-ground and below-ground parts of plants [88-91].

Along with those possibly saprotrophic or endophytic taxa, ericoid mycorrhizal and

phytopathogenic taxa of fungi displayed relatively high betweenness scores within the

metacommunity-scale network representing all the eight local forests (Table 1). Specifically,

Oidiodendron (teleomorph = *Myxotrichum*) is a taxon represented by possibly ericoid

mycorrhizal species (*O. maius* and *O. griseum*) [92, 93], although fungi in the genus are found

also from roots of non-ericaceous plants and soil [94]. On the other hand, fungi in the family

Nectriaceae are known to cause black foot disease [95], often having serious damage on

economically important woody plants [96, 97]. Although we collected seemingly benign roots

in the study forests, some samples may be damaged by those pathogens. Alternatively, some

lineages of Nectriaceae fungi may be associated with plant hosts non-symptomatically,

having adverse effects context-dependently.

Although these fungi were candidates of metacommunity hubs, which are characterized

by broad geographic ranges and host plant ranges, none except but a *Mortierella* OTU had

high betweenness scores at both local and metacommunity levels (Fig. 5a). This result

suggests that even if some fungi have broad geographic ranges across the Japanese

Archipelago, few played important topological roles in each of the local networks

representing plant–fungus associations. In other words, fungi that can adapt to biotic and

abiotic environments in forest ecosystems throughout cool-temperate, warm-temperate, and

subtropical regions are rare.

Therefore, we also explored fungi with broad geographic and host ranges within the

metacommunities representing northern (cool-temperate) and southern (warm-temperate and

subtropical) regions of Japan. In the metacommunity consisting of the four cool-temperate

forests (Additional file 7; Figure S4), fungal OTUs in the genera *Mortierella*,

Cladophialophora, and *Pezicula* as well as those allied to *Ilyonectria* and *Cadophora* had

highest betweenness at both local and metacommunity levels, classified as metacommunity

hubs (Fig. 5b; Table 1). Among them, *Cladophialophora* is of particular interest because it

has been known as a lineage of "dark septate endophytes" [98-100] (*sensu* [14, 15, 101]). A

species within the genus, *C. chaetospira* (= *Heteroconium chaetospira*), to which

high-betweenness OTUs in our data were closely allied, has been known not only to provide

nitrogen to host plants but also to suppress pathogens [12, 16, 102]. Likewise, the Helotiales

genus *Pezicula* (anamorph = *Cryptosporiopsis*) includes endophytic fungi [103-105], some of

which produce secondary metabolites suppressing other microbes in the rhizosphere [106,

107]. Our finding that some of *Cladophialophora* and *Pezicula* fungi could be associated with

various taxonomic groups of plants over broad geographic ranges highlights potentially

important physiological and ecological roles of those endophytes at the community and

metacommunity levels.

In the southern metacommunity networks consisting of warm-temperate and subtropical

forests (Additional file 8; Figure S5), some arbuscular mycorrhizal OTUs and *Saitozyma*

(*Cryptococcus*) and *Mortierella* OTUs had high betweenness scores at both local and

metacommunity levels, designated as metacommunity hubs (Fig. 5c; Table 1). Given the

above-mentioned prevalence of fungal OTUs allied to *Cladophialophora chaetospira* in the

cool-temperate metacommunity, the contrasting list of metacommunity hubs in the southern

(warm-temperate–subtropical) metacommunity implies that different taxonomic and

functional groups of fungi play major metacommunity-scale roles in different climatic regions.

This working hypothesis is partially supported by previous studies indicating endemism and

vicariance in the biogeography of fungi and bacteria [108, 109], promoting conceptual

advances beyond the classic belief that every microbe is everywhere but the environment

selects microbes colonizing respective local communities [110].

The roles of those metacommunity hubs detected in this study are of particular interest

from the aspect of theoretical ecology. Hub species connected to many other species in an

ecosystem often integrate "energy channels" [111] within species interaction networks,

having great impacts on biodiversity and productivity of the ecosystems [35]. The concept of

"keystone" or "foundation" species [112, 113] can be extended to the metacommunity level,

thereby promoting studies exploring species that restructure and synchronize ecological (and

evolutionary) dynamics over broad geographic ranges [35]. Given that below-ground plant–

fungus symbioses are key components of the terrestrial biosphere [1, 2], identifying fungal

species that potentially have great impacts on the metacommunity-scale processes of such

below-ground interactions will provide crucial insights into the conservation and restoration

of forests and grasslands. We here showed that the list of metacommunity hubs could involve

various lineages of endophytic fungi, whose ecosystem-scale functions have been

underappreciated compared to those of mycorrhizal fungi. As those endophytic fungi are

potentially used as inoculants when we reintroduce plant seedlings in ecosystem restoration

programs [20, 51], exploring fungi with highest potentials in each climatic/biogeographic

region will be a promising direction of research in conservation biology.

The finding that compositions of metacommunity hubs could vary depending on climatic

regions also gives key implications for the application of endophytes in agriculture. Although

a number of studies have tried to use endophytic fungi and/or bacteria as microbial inoculants

in agriculture [17, 18, 114], such microbes introduced to agroecosystems are often

outcompeted and replaced by indigenous (resident) microbes [115, 116]. Moreover, even if an

endophytic species or strain increases plant growth in pot experiments under controlled

environmental conditions, its effects in the field often vary considerably depending on biotic

and abiotic contexts of local agroecosystems [17] (see also [117]). Therefore, in the screening

of endophytes that can be used in broad ranges of biotic and abiotic environmental conditions,

the metacommunity-scale network analysis outlined in this study will help us find promising

candidates out of thousands or tens of thousands microbial species in the wild. Consequently,

to find promising microbes whose inocula can persist in agroecosystems for long time periods,

exploration of metacommunity hubs needs to be performed in respective climatic or

biogeographic regions.

For more advanced applications in conservation biology and agriculture, continual

improvements of methods for analyzing metacommunity-scale networks are necessary. First,

while the fungal OTUs in our network analysis was defined based on the cut-off sequence

similarities used in other studies targeting "species-level" diversity of fungi [57, 59],

physiological functions can vary greatly within fungal species or species groups [14, 118].

Given that bioinformatic tools that potentially help us detect single-nucleotide-level variation

are becoming available [119], the resolution of network analyses may be greatly improved in

the near future. Second, although some computer programs allow us to infer functions of

respective microbial OTUs within network data [63, 120], the database information of

microbial functions remains scarce. To increase the coverage and accuracy of automatic

annotations of microbial functions, studies describing the physiology, ecology, and genomes

of microbes should be accelerated. With improved reference databases, more insights into the

metacommunity-scale organization of plant–fungus associations will be obtained by

reanalyzing the network data by compiling enhanced information of fungal functional groups.

Third, as the diversity and compositions of plant–fungus associations included in a network

can depend on how we process raw samples, special care is required in the selection of

methods for washing and preparing root (or soil) samples. By sterilizing root samples with

NaClO [121], for example, we may be able to exclude fungi or bacteria that are merely

adhering to root surfaces. Meanwhile, some of those fungi and bacteria on root surfaces may

play pivotal physiological roles in the growth and survival of plants [122]. Accordingly, it

would be productive to compare network topologies of plant–microbe associations among

different source materials by partitioning endosphere, rhizoplane, and rhizosphere microbial

samples with a series of sample cleaning processes using ultrasonic devices [123]. Fourth,

although this study targeted fungi associated with roots, our methods can be easily extended

to network analyses involving other groups of microbes. By simultaneously analyzing the

prokaryote 16S rRNA region [123-125] with the fungal ITS region, we can examine how

bacteria, archaea, and fungi are involved in below-ground webs of symbioses. Fifth, not only

plant–microbe associations but also microbe–microbe interactions can be estimated with

network analytical frameworks. Various statistical pipelines have been proposed to infer how

microbes interact with each other in facilitative or competitive ways within host

macroorganisms [37, 126, 127]. Overall, those directions of analytical extensions will

enhance our understanding of plant microbiome dynamics in nature.

Conclusions

By compiling datasets of below-ground plant–fungus associations in temperate and

subtropical forest ecosystems, we explored metacommunity-hub fungi, which were

characterized by broad geographic and host ranges. Such metacommunity-scale analyses are

expected to provide bird's-eye views of complex plant–microbe associations, highlighting

plant-growth-promoting microbes that can be applied to diverse plant taxa in various

environments. Given that endophytic fungi promoting the growth and pathogen resistance of

host plants can be isolated from forest soil (e.g., *Cladophialophora chaetospira* [99]), the list

of metacommunity-hub endophytic fungi featured in this study itself may include prospective

species to be used in agriculture. By extending the targets of such network analyses to diverse

types of plant-associated microbes (e.g., phyllosphere fungi and bacteria [75, 124, 128]) in

various climatic/biogeographic regions, a solid basis for managing plant microbiomes will be

developed.

Abbreviations

DDBJ: DNA Data Bank of Japan; ITS: internal transcribed spacer; OTU: Operational

taxonomic unit; PERMANOVA: permutational analysis of variance; PERMDISP:

permutational analysis for the multivariate homogeneity of dispersions; rRNA: ribosomal

ribonucleic acid.

Acknowledgements

We thank Teshio Experimental Forest (Hokkaido University), Tomakomai Experimental

Forest (Hokkaido University), Sugadaira Research Station (Tsukuba University), Yona Field

(Ryukyu University), Tropical Biosphere Research Center (Ryukyu University), and Forestry

Agency of Japan for the permission of fieldwork.

Funding

This work was financially supported by JSPS KAKENHI Grant (26711026), JST PRESTO

(JPMJPR16Q6), and the Funding Program for Next Generation World-Leading Researchers

of Cabinet Office, the Government of Japan (GS014) to HT.

Availability of data and materials

The Illumina sequencing data were deposited to DNA Data Bank of Japan (DDBJ Sequence

Read Archive: DRA006339). The raw data of fungal community structure and the fungal

community matrices analyzed are available with the source study [52] and Additional files

1-3, respectively.

Authors' contributions

HT designed the work. HT, AST, and HS conducted fieldwork. HT performed the molecular

experiments. HT wrote the manuscript with AST and HS.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable

Ethics approval and consent to participate

Not applicable

Additional files

Additional file 1: Data S1. Sample-level matrices of the eight forests examined.

Additional file 2: Data S2. Species-level matrices of plant–fungus associations.

Additional file 3: Data S3. Information of 8080 fungal OTUs analyzed.

Additional file 4: Figure S1. Number of sequencing reads, interaction specificity, and local

betweenness.

Additional file 5: Figure S2. Structure of plant–fungus networks in each local forest.

Additional file 6: Figure S3. Locality information within the full metacommunity-scale

network.

Additional file 7: Figure S4. Metacommunity-scale network of cool-temperate forests.

Additional file 8: Figure S5. Metacommunity-scale network of warm-temperate and

subtropical forests.

Additional file 9: Table S1. Top-10 list of non-Glomeromycota OTUs with highest

betweenness within the subtropical metacommunity network.

References

1.   Peay KG, Kennedy PG, Talbot JM: Dimensions of biodiversity in the Earth
501   mycobiome. Nat. Rev. Microbiol. 2016;14:434-447.

502   2.   van der Heijden MG, Bardgett RD, van Straalen NM: The unseen majority: soil
503   microbes as drivers of plant diversity and productivity in terrestrial ecosystems. Ecol.
504   Lett. 2008;11:296-310.

505   3.   van der Heijden MG, Klironomos JN, Ursic M, Moutoglis P, Streitwolf-Engel R,
506   Boller T et al: Mycorrhizal fungal diversity determines plant biodiversity, ecosystem
507   variability and productivity. Nature 1998;396:69-72.

508   4.   Wardle DA, Bardgett RD, Klironomos JN, Setälä H, van der Putten WH, Wall DH:
509   Ecological linkages between aboveground and belowground biota. Science
510   2004;304:1629-1633.

511   5.   Nara K: Ectomycorrhizal networks and seedling establishment during early primary
512   succession. New Phytol. 2006;169:169-178.

513   6.   Finlay R, Read D: The structure and function of the vegetative mycelium of
514   ectomycorrhizal plants. II. The uptake and distribution of phosphorus by mycelial
515   strands interconnecting host plants. New Phytol. 1986;103:157-165.

516   7.   Klironomos JN, McCune J, Hart M, Neville J: The influence of arbuscular
517   mycorrhizae on the relationship between plant diversity and productivity. Ecol. Lett.
518   2000;3:137-141.

519   8.   Högberg MN, Högberg P: Extramatrical ectomycorrhizal mycelium contributes
520   one-third of microbial biomass and produces, together with associated roots, half the

dissolved organic carbon in a forest soil. New Phytol. 2002;154:791-795.

9.   Högberg P, Nordgren A, Buchmann N, Taylor AF, Ekblad A, Högberg MN et al:
Large-scale forest girdling shows that current photosynthesis drives soil respiration.
Nature 2001;411:789-792.

10.   Almario J, Jeena G, Wunder J, Langen G, Zuccaro A, Coupland G et al:
Root-associated fungal microbiota of nonmycorrhizal *Arabis alpina* and its
contribution to plant phosphorus nutrition. Proc. Natl. Acad. Sci. USA.
2017;114:E9403–E9412.

11.   Hiruma K, Gerlach N, Sacristán S, Nakano RT, Hacquard S, Kracher B et al: Root
endophyte *Colletotrichum tofieldiae* confers plant fitness benefits that are phosphate
status dependent. Cell 2016;165:464-474.

12.   Hashiba T, Narisawa K: The development and endophytic nature of the fungus
Heteroconium chaetospira. FEMS Microbiol. Lett. 2005;252:191-196.

13.   Addy H, Piercey M, Currah R: Microfungal endophytes in roots. Can. J. Bot.
2005;83:1-13.

14.   Jumpponen A: Dark septate endophytes-are they mycorrhizal? Mycorrhiza
2001;11:207-211.

15.   Newsham KK: A meta-analysis of plant responses to dark septate root endophytes.
New Phytol. 2011;190:783-793.

16.   Narisawa K, Tokumasu S, Hashiba T: Suppression of clubroot formation in Chinese
cabbage by the root endophytic fungus, *Heteroconium chaetospira*. Plant Pathol.
1998;47:206-210.

17.   Calvo P, Nelson L, Kloepper JW: Agricultural uses of plant biostimulants. Plant Soil
2014;383:3-41.

18.   Porras-Alfaro A, Bayman P: Hidden fungi, emergent properties: endophytes and
microbiomes. Ann. Rev. Phytopathol. 2011;49:291-315.

19.   Mandyam K, Jumpponen A: Seeking the elusive function of the root-colonising dark
septate endophytic fungi. Studies Mycol. 2005;53:173-189.

20. Fuchs B, Haselwandter K: Red list plants: colonization by arbuscular mycorrhizal
fungi and dark septate endophytes. Mycorrhiza 2004;14:277-281.

21. Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R et al: Global
diversity and geography of soil fungi. Science 2014;346:1256688.

22. Buée M, Reich M, Murat C, Morin E, Nilsson R, Uroz S et al: 454 Pyrosequencing
analyses of forest soils reveal an unexpectedly high fungal diversity. New Phytol.
2009;184:449-456.

23. Öpik M, Vanatoa A, Vanatoa E, Moora M, Davison J, Kalwij J et al: The online
database MaarjAM reveals global and ecosystemic distribution patterns in arbuscular
mycorrhizal fungi (Glomeromycota). New Phytol. 2010;188:223-241.

24. Toju H, Guimarães PR, Jr, Olesen JM, Thompson JN: Assembly of complex plant–
fungus networks. Nat. Commun. 2014;5:5273.

25. Toju H, Guimarães PR, Jr, Olesen JM, Thompson JN: Below-ground plant–fungus
network topology is not congruent with above-ground plant–animal network topology.
Sci. Adv. 2015;1:e1500291.

26. Kadowaki K, Yamamoto S, Sato H, Tanabe AS, Hidaka A, Toju H: Plant−soil
feedbacks between arbuscular- and ecto-mycorrhizal communities. bioRxiv
2017:https://doi.org/10.1101/228387

27. Schlaeppi K, Bulgarelli D: The plant microbiome at work. Mol. Plant-Microbe Int.
2015;28:212-217.

28. Berendsen RL, Pieterse CM, Bakker PA: The rhizosphere microbiome and plant
health. Trends Plant Sci. 2012;17:478-486.

29. Busby PE, Soman C, Wagner MR, Friesen ML, Kremer J, Bennett A et al: Research
priorities for harnessing plant microbiomes in sustainable agriculture. PLOS Biol.
2017;15:e2001793.

30. Albert R, Jeong H, Barabási AL: Error and attack tolerance of complex networks.
Nature 2000;406:378-382.

31. Newman MEJ: Networks: an introduction. Oxford: Oxford University Press; 2010.

32.   Toju H, Tanabe A, Ishii H: Ericaceous plant–fungus network in a harsh alpine–
subalpine environment. Mol. Ecol. 2016;25:3242-3257.

33.   Bennett AE, Daniell TJ, Öpik M, Davison J, Moora M, Zobel M et al: Arbuscular
mycorrhizal fungal networks vary throughout the growing season and between
successional stages. PLoS One 2013;8:e83241.

34.   Bahram M, Harend H, Tedersoo L: Network perspectives of ectomycorrhizal
associations. Fungal Ecol. 2014;7:70-77.

35.   Toju H, Yamamichi M, Guimarães Jr. PR, Olesen J, Mougi A, Yoshida T et al:
Species-rich networks and eco-evolutionary synthesis at the metacommunity level.
Nat. Ecol. Evol. 2017;1.

36.   Agler MT, Ruhe J, Kroll S, Morhenn C, Kim S-T, Weigel D et al: Microbial hub taxa
link host and abiotic factors to plant microbiome variation. PLOS Biol.
2016;14:e1002352.

37.   Toju H, Yamamoto S, Tanabe AS, Hayakawa T, Ishii HS: Network modules and hubs
in plant-root fungal biome. J. R. Soc. Interface 2016;13:20151097.

38.   Smith SE, Read DJ: Mycorrhizal symbiosis. New York: Academic press; 2008.

39.   Dickie IA, Koide RT, Fayish AC: Vesicular-arbuscular mycorrhizal infection of
Quercus rubra seedlings. New Phytol. 2001;151:257-264.

40.   Plattner I, Hall I: Parasitism of non-host plants by the mycorrhizal fungus *Tuber*
melanosporum. Mycol. Res. 1995;99:1367-1370.

41.   Hallenberg N, Kuffer N: Long‐distance spore dispersal in wood‐inhabiting
basidiomycetes. Nordic J. Bot. 2001;21:431-436.

42.   Nagarajan S, Singh D: Long-distance dispersion of rust pathogens. Ann. Rev.
Phytopathol. 1990;28:139-153.

43.   Brown JK, Hovmøller MS: Aerial dispersal of pathogens on the global and continental
scales and its impact on plant disease. Science 2002;297:537-541.

44.   Hovmøller M, Justesen A, Brown J: Clonality and long‐distance migration of
Puccinia striiformis f. sp. tritici in north‐west Europe. Plant Pathol. 2002;51:24-32.

45.   Jousimo J, Tack AJ, Ovaskainen O, Mononen T, Susi H, Tollenaere C et al:
Ecological and evolutionary effects of fragmentation on infectious disease dynamics.
Science 2014;344:1289-1293.

46.   Parker MA: Mutualism in metapopulations of legumes and rhizobia. Am. Nat.
1999;153:S48-S60.

47.   Thrall PH, Burdon JJ: Evolution of virulence in a plant host-pathogen metapopulation.
Science 2003;299:1735-1737.

48.   Lekberg Y, Koide RT, Rohr JR, ALDRICH‑WOLFE L, Morton JB: Role of niche
restrictions and dispersal in the composition of arbuscular mycorrhizal fungal
communities. J. Ecol. 2007;95:95-105.

49.   Peay KG, Schubert MG, Nguyen NH, Bruns TD: Measuring ectomycorrhizal fungal
dispersal: macroecological patterns driven by microscopic propagules. Mol. Ecol.
2012;21:4122-4136.

50.   Gouhier TC, Guichard F, Gonzalez A: Synchrony and stability of food webs in
metacommunities. Am. Nat. 2010;175:E16-E34.

51.   Miller R, Jastrow J: The application of VA mycorrhizae to ecosystem restoration and
reclamation. New York: Springer; 1992.

52.   Toju H, Sato H, Yamamoto S, Tanabe AS: Structural diversity across arbuscular
mycorrhizal, ectomycorrhizal, and endophytic plant–fungus networks. bioRxiv
2018:https://doi.org/10.1101/269563

53.   Caruso T, Rillig MC, Garlaschelli D: On the application of network theory to
arbuscular mycorrhizal fungi–plant interactions: the importance of basic assumptions.
New Phytol. 2012;194:891-894.

54.   Thompson JN: The geographic mosaic of coevolution. Chicago: University of
Chicago Press; 2005.

55.   Toju H, Tanabe AS, Yamamoto S, Sato H: High-coverage ITS primers for the
DNA-based identification of ascomycetes and basidiomycetes in environmental
samples. PLOS ONE 2012;7:e40863.

56.   Smith DP, Peay KG: Sequence depth, not PCR replication, improves ecological

inference from next generation DNA sequencing. PLOS ONE 2014;9:e90234.

57.   O'Brien HE, Parrent JL, Jackson JA, Moncalvo J-M, Vilgalys R: Fungal community
analysis by large-scale sequencing of environmental samples. Appl. Env. Microbiol.
2005;71:5544-5550.

58.   Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M et al:
Towards a unified paradigm for sequence-based identification of fungi. Mol. Ecol.
2013;22:5271-5277.

59.   Thiéry O, Vasar M, Jairus T, Davison J, Roux C, Kivistik PA et al: Sequence variation
in nuclear ribosomal small subunit, internal transcribed spacer and large subunit
regions of *Rhizophagus irregularis* and *Gigaspora margarita* is high and isolate‐
dependent. Mol. Ecol. 2016;25:2816-2832.

60.   Tanabe AS: Claident v0.2.2016.07.05, a software distributed by author at
http://www.fifthdimension.jp/. 2016.

61.   Tanabe AS, Toju H: Two new computational methods for universal DNA barcoding: a
benchmark using barcode sequences of bacteria, archaea, animals, fungi, and land
plants. PLOS ONE 2013;8:e76910.

62.   Huson DH, Auch AF, Qi J, Schuster SC: MEGAN analysis of metagenomic data.
Genome Res. 2007;17:377-386.

63.   Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J et al: FUNGuild: an
open annotation tool for parsing fungal community datasets by ecological guild.
Fungal Ecol. 2016;20:241-248.

64.   Anderson MJ: A new method for non-parametric multivariate analysis of variance.
Austral Ecol. 2001;26:32-46.

65.   Anderson MJ: Distance-based tests for homogeneity of multivariate dispersions.
Biometrics 2006;62:245-253.

66.   Oksanen J, Blanachet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB et al: Vegan:
community ecology package. R package version 2.0-3 available at
http://CRAN.R-project.org/package=vegan. 2012;10:2008.

67.   R-Core-Team: R 3.4.1: A language and environment for statistical computing

available at http://www.R-project.org/. In. Vienna, Austri: R Foundation for
Statistical Computing; 2015.

68.   Chase JM, Kraft NJ, Smith KG, Vellend M, Inouye BD: Using null models to
666   disentangle variation in community dissimilarity from variation in α-diversity.
667   Ecosphere 2011;2:1-11.

668   69.   Bastian M, Heymann S, Jacomy M: Gephi: an open source software for exploring and
669   manipulating networks. Proceedings of the 3rd International AAAI Conference on
670   Weblogs and Social Media 2009;8:361-362.

671   70.   Jacomy M, Heymann S, Venturini T, Bastian M: ForceAtlas2, a continuous graph
672   layout algorithm for handy network visualization designed for the Gephi Software.
673   PLOS ONE 2011;96:e98679.

674   71.   Blüthgen N, Menzel F, Blüthgen N: Measuring specialization in species interaction
675   networks. BMC Ecol. 2006;6:9.

676   72.   Freeman LC: A set of measures of centrality based on betweenness. Sociometry
677   1977;40:35-41.

678   73.   Tedersoo L, Pärtel K, Jairus T, Gates G, Põldmaa K, Tamm H: Ascomycetes
679   associated with ectomycorrhizas: molecular diversity and ecology with particular
680   reference to the Helotiales. Env. Microbiol. 2009;11:3166-3178.

681   74.   Rossman AY, Farr DF, Castlebury LA: A review of the phylogeny and biology of the
682   Diaporthales. Mycoscience 2007;48:135-144.

683   75.   Sieber TN: Endophytic fungi in forest trees: are they mutualists? Fungal Biol. Rev.
684   2007;21:75-89.

685   76.   Liu X-Z, Wang Q-M, Göker M, Groenewald M, Kachalkin A, Lumbsch HT et al:
686   Towards an integrated phylogenetic classification of the Tremellomycetes. Studies
687   Mycol. 2015;81:85-147.

688   77.   Watanabe T: Pictorial atlas of soil and seed fungi: morphologies of cultured fungi and
689   key to species. Boca Raton: CRC press; 2010.

690   78.   Zhang H, Wu X, Li G, Qin P: Interactions between arbuscular mycorrhizal fungi and
691   phosphate-solubilizing fungus (Mortierella sp.) and their effects on Kostelelzkya

virginica growth and enzyme activities of rhizosphere and bulk soils at different
salinities. Biol. Fertil. Soils 2011;47:543.

79. Eroshin V, Dedyukhina E: Effect of lipids from *Mortierella hygrophila* on plant
resistance to phytopathogens. World J. Microbiol. Biotechnol. 2002;18:165-167.

80. Melo IS, Santos SN, Rosa LH, Parma MM, Silva LJ, Queiroz SC et al: Isolation and
biological activities of an endophytic *Mortierella alpina* strain from the Antarctic
moss *Schistidium antarctici*. Extremophiles 2014;18:15-23.

81. Papavizas G: *Trichoderma* and *Gliocladium*: biology, ecology, and potential for
biocontrol. Ann. Rev. Phytopathol. 1985;23:23-54.

82. Datnoff L, Nemec S, Pernezny K: Biological control of *Fusarium* crown and root rot
of tomato in Florida using *Trichoderma harzianum* and *Glomus intraradices*. Biologi.
Control 1995;5:427-431.

83. Bailey B, Bae H, Strem M, Crozier J, Thomas S, Samuels G et al: Antibiosis,
mycoparasitism, and colonization success for endophytic *Trichoderma* isolates with
biological control potential in Theobroma cacao. Biologi. Control 2008;46:24-35.

84. Bae H, Roberts DP, Lim H-S, Strem MD, Park S-C, Ryu C-M et al: Endophytic
Trichoderma isolates from tropical environments delay disease onset and induce
resistance against Phytophthora capsici in hot pepper using multiple mechanisms. Mol.
Plant-Microbe Int. 2011;24:336-351.

85. AL-Shammari TA, Bahkali AH, Elgorban AM, El-Kahky MT, Al-Sum BA: The use
of Trichoderma longibrachiatum and *Mortierella alpina* against root-knot nematode,
Meloidogyne javanica on tomato. J. Pure Appl. Microbiol. 2013;7:199-207.

86. Ming Q, Su C, Zheng C, Jia M, Zhang Q, Zhang H et al: Elicitors from the endophytic
fungus *Trichoderma atroviride* promote *Salvia miltiorrhiza* hairy root growth and
tanshinone biosynthesis. J. Exp. Bot. 2013;64:5687-5694.

87. Mašínová T, Bahnmann BD, Větrovský T, Tomšovský M, Merunková K, Baldrian P:
Drivers of yeast community composition in the litter and soil of a temperate forest.
FEMS Microbiol. Ecol. 2016;93:fiw223.

88. Nutaratat P, Srisuk N, Arunrattiyakorn P, Limtong S: Plant growth-promoting traits of

epiphytic and endophytic yeasts isolated from rice and sugar cane leaves in Thailand.

Fungal Biol. 2014;118:683-694.

89.   Renker C, Blanke V, Börstler B, Heinrichs J, Buscot F: Diversity of *Cryptococcus* and

Dioszegia yeasts (Basidiomycota) inhabiting arbuscular mycorrhizal roots or spores.

FEMS Yeast Res. 2004;4:597-603.

90.   Gai CS, Lacava PT, Maccheroni W, Glienke C, Araújo WL, Miller TA et al: Diversity

of endophytic yeasts from sweet orange and their localization by scanning electron

microscopy. J. Basic Microbiol. 2009;49:441-451.

91.   Cloete KJ, Valentine AJ, Stander MA, Blomerus LM, Botha A: Evidence of symbiosis

between the soil yeast *Cryptococcus laurentii* and a sclerophyllous medicinal shrub,

Agathosma betulina (Berg.) Pillans. Microb. Ecol. 2009;57:624-632.

92.   Couture M, Fortin J, Dalpe Y: Oidiodendron griseum Robak: An endophyte of ericoid

mycorrhiza in *Vaccinium* spp. New Phytol. 1983;95:375-380.

93.   Vohník M, Albrechtová J, Vosátka M: The inoculation with *Oidiodendron maius* and

Phialocephala fortinii alters phosphorus and nitrogen uptake, foliar C: N ratio and root

biomass distribution in Rhododendron cv. Azurro. Symbiosis 2005;40:87-96.

94.   Rice AV, Currah RS: *Oidiodendron*: A survey of the named species and related

anamorphs of *Myxotrichum*. Studies Mycol. 2005;53:83-120.

95.   Lombard L, Van Der Merwe NA, Groenewald JZ, Crous PW: Lineages in

Nectriaceae: re-evaluating the generic status of *Ilyonectria* and allied genera.

Phytopathol. Mediterr. 2014;53:515-532.

96.   Whitelaw‑Weckert M, Rahman L, Appleby L, Hall A, Clark A, Waite H et al: Co‑

infection by Botryosphaeriaceae and Ilyonectria spp. fungi during propagation causes

decline of young grafted grapevines. Plant Pathol. 2013;62:1226-1237.

97.   Vitale A, Aiello D, Guarnaccia V, Perrone G, Stea G, Polizzi G: First report of root rot

caused by *Ilyonectria* (= *Neonectria*) macrodidyma on avocado (*Persea americana*) in

Italy. J. Phytopathol. 2012;160:156-159.

98.   Usuki F, Narisawa K: Formation of structures resembling ericoid mycorrhizas by the

root endophytic fungus *Heteroconium chaetospira* within roots of *Rhododendron*

obtusum var. *kaempferi*. Mycorrhiza 2005;15:61-64.

99.   Narisawa K, Hambleton S, Currah RS: *Heteroconium chaetospira*, a dark septate root
endophyte allied to the Herpotrichiellaceae (Chaetothyriales) obtained from some
forest soil samples in Canada using bait plants. Mycoscience 2007;48:274-281.

100.   Vohník M, Albrechtová J: The co-occurrence and morphological continuum between
ericoid mycorrhiza and dark septate endophytes in roots of six European
Rhododendron species. Folia Geobotanica 2011;46:373-386.

101.   Jumpponen A, Trappe JM: Dark septate endophytes: a review of facultative biotrophic
root-colonizing fungi. New Phytol. 1998;140:295-310.

102.   Usuki F, Narisawa K: A mutualistic symbiosis between a dark septate endophytic
fungus, *Heteroconium chaetospira*, and a nonmycorrhizal plant, Chinese cabbage.
Mycologia 2007;99:175-184.

103.   Wang W, Tsuneda A, Gibas CF, Currah RS: *Cryptosporiopsis* species isolated from
the roots of aspen in central Alberta: identification, morphology, and interactions with
the host, in vitro Botany 2007;85:1214-1226.

104.   Kowalski T, Bartnik C: *Cryptosporiopsis radicicola* sp. nov. from roots of *Quercus*
robur. Mycol. Res. 1995;99:663-666.

105.   Sigler L, Allan T, Lim SR, Berch S, Berbee M: Two new *Cryptosporiopsis* species
from roots of ericaceous hosts in western North America. Studies Mycol.
2005;53:53-62.

106.   Schulz B, Sucker J, Aust H, Krohn K, Ludewig K, Jones P et al: Biologically active
secondary metabolites of endophytic *Pezicula* species. Mycol. Res.
1995;99:1007-1015.

107.   Strobel GA, Miller RV, Martinez-Miller C, Condron MM, Teplow DB, Hess W:
Cryptocandin, a potent antimycotic from the endophytic fungus *Cryptosporiopsis* cf.
quercina. Microbiology 1999;145:1919-1926.

108.   Talbot JM, Bruns TD, Taylor JW, Smith DP, Branco S, Glassman SI et al: Endemism
and functional convergence across the North American soil mycobiome. Proc. Natl.
Acad. Sci. USA. 2014;111:6341-6346.

109.   Martiny JBH, Bohannan BJ, Brown JH, Colwell RK, Fuhrman JA, Green JL et al:
Microbial biogeography: putting microorganisms on the map. Nat. Rev. Microbiol.
2006;4:102-112.

110.   De Wit R, Bouvier T: 'Everything is everywhere, but, the environment selects'; what
did Baas Becking and Beijerinck really say? Env. Microbiol. 2006;8:755-758.

111.   Rooney N, McCann K, Gellner G, Moore JC: Structural asymmetry and the stability
of diverse food webs. Nature 2006;442:265-269.

112.   Ellison AM, Bank MS, Clinton BD, Colburn EA, Elliott K, Ford CR et al: Loss of
foundation species: consequences for the structure and dynamics of forested
ecosystems. Front. Ecol. Env. 2005;3:479-486.

113.   Paine RT: Food web complexity and species diversity. Am. Nat. 1966;100:65-75.

114.   Waller F, Achatz B, Baltruschat H, Fodor J, Becker K, Fischer M et al: The
endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance,
disease resistance, and higher yield. Proc. Natl. Acad. Sci. USA.
2005;102:13386-13391.

115.   Castro-Sowinski S, Herschkovitz Y, Okon Y, Jurkevitch E: Effects of inoculation with
plant growth-promoting rhizobacteria on resident rhizosphere microorganisms. FEMS
Microbiol. Lett. 2007;276:1-11.

116.   Streeter JG: Failure of inoculant rhizobia to overcome the dominance of indigenous
strains for nodule formation. Can. J. Bot. 1994;40:513-522.

117.   Hartman K, van der Heijden MG, Roussely-Provent V, Walser J-C, Schlaeppi K:
Deciphering composition and function of the root microbiome of a legume plant.
Microbiome 2017;5:2.

118.   Jumpponen A, Trappe JM: Performance of *Pinus contorta* inoculated with two strains
of root endophytic fungus, *Phialocephala fortinii*: effects of synthesis system and
glucose concentration. Can. J. Bot. 1998;76:1205-1213.

119.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP: DADA2:
high-resolution sample inference from Illumina amplicon data. Nat. Methods
2016;13:581-583.

120.   Langille MG, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA et al:
809   Predictive functional profiling of microbial communities using 16S rRNA marker
810   gene sequences. Nat. Biotech. 2013;31:814-821.

811   121.   Vohník M, Mrnka L, Lukešová T, Bruzone MC, Kohout P, Fehrer J: The cultivable
812   endophytic community of Norway spruce ectomycorrhizas from microhabitats lacking
813   ericaceous hosts is dominated by ericoid mycorrhizal *Meliniomyces variabilis*. Fungal
814   Ecol. 2013;6:281-292.

815   122.   Richardson AE, Barea J-M, McNeill AM, Prigent-Combaret C: Acquisition of
816   phosphorus and nitrogen in the rhizosphere and plant growth promotion by
817   microorganisms. Plant Soil 2009;321:305-339.

818   123.   Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S et al:
819   Defining the core *Arabidopsis thaliana* root microbiome. Nature 2012;488:86-90.

820   124.   Bai Y, Müller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M et al: Functional
821   overlap of the *Arabidopsis* leaf and root microbiota. Nature 2015;528:364-369.

822   125.   Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S et al:
823   Structure, variation, and assembly of the root-associated microbiomes of rice. Proc.
824   Natl. Acad. Sci. USA. 2015;112:E911-E920.

825   126.   Faust K, Raes J: Microbial interactions: from networks to models. Nat. Rev. Microbiol.
826   2012;10:538-550.

827   127.   Kurtz ZD, Mueller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA: Sparse and
828   compositionally robust inference of microbial ecological networks. PLOS Comp. Biol.
829   2015;11:e1004226.

830   128.   Arnold AE, Mejía LC, Kyllo D, Rojas EI, Maynard Z, Robbins N et al: Fungal
831   endophytes limit pathogen damage in a tropical tree. Proc. Natl. Acad. Sci. USA.
832   2003;100:15649-15654.

Table 1 Top-10 list of non-Glomeromycota OTUs with highest betweenness within the metacommunity networks. In each of the three metacommunity-scale networks examined (full, cool-temperate, and warm-temperate/subtropical), fungal OTUs were ranked based on their betweenness centrality scores. As taxonomic information of Glomeromycota OTUs with high betweenness scores was redundant (e.g., *Glomus* spp. or Glomeraceae spp.), the top-10 list of non-Glomeromycota OTUs is shown. Taxonomy information of each OTU was inferred based on the query-centric auto-*k*-nearest-neighbor algorithm of reference database search [60, 61] and subsequent taxonomic assignment with the lowest common ancestor algorithm [62]. The results of the NCBI nucleotide Blast are also shown. For simplicity, the functional groups of fungi inferred with the program FUNGuild [63] were organized into several categories. See Data S3 (Additional file 3) for details of the categories and for full results including Glomeromycota and other fungal OTUs.

OTU	Phylum	Class	Order	Family	Genus	Category	NCBI Blast top hit	Accession	Cover	Identity
Full (8sites)										
F_0042*	-	-	Mortierellales	Mortierellaceae	*Mortierella*	Saprotroph/Endophyte	*Mortierella humilis*	KP714537	100%	100%
F_0381	Basidiomycota	Tremellomycetes	Trichosporonales	Trichosporonaceae	*Cryptococcus*	Others_Unknown	*Saitozyma podzolica†*	KY320605	92%	99%
F_0079	Ascomycota	Sordariomycetes	Hypocreales	Nectriaceae	-	Saprotroph/Endophyte	*Ilyonectria protearum*	NR_152890	99%	100%
F_0489	-	-	Mortierellales	Mortierellaceae	*Mortierella*	Saprotroph/Endophyte	*Mortierella* sp.	KM113754	100%	100%
F_0010	Ascomycota	Leotiomycetes	-	Myxotrichaceae	*Oidiodendron*	Ericoid_Mycorrhizal	*Oidiodendron maius*	LC206669	100%	100%
F_0368	Basidiomycota	Malasseziomycetes	Malasseziales	Malasseziaceae	*Malassezia*	Others_Unknown	*Malassezia restricta*	KT809059	100%	100%
F_0623	-	-	Mortierellales	Mortierellaceae	*Mortierella*	Saprotroph/Endophyte	*Mortierella gamsii*	KY305027	100%	100%
F_1188	Basidiomycota	Tremellomycetes	Trichosporonales	Trichosporonaceae	*Cryptococcus*	Others_Unknown	*Saitozyma podzolica†*	KY320605	92%	99%
F_0007	Ascomycota	Sordariomycetes	Diaporthales	Melanconidaceae	*Melanconiella*	Saprotroph/Endophyte	*Melanconiella elegans*	KJ173701	100%	85%
F_0485	Ascomycota	Sordariomycetes	Hypocreales	Hypocreaceae	*Trichoderma*	Saprotroph/Endophyte	*Trichoderma* sp.	HG008760	100%	100%

Northen 4 sites (cool-temperate)

ID	Phylum	Class	Order	Family	Genus	Guild	Species	Accession	%	%
F_0042*	-	-	Mortierellales	Mortierellaceae	*Mortierella*	Saprotroph/Endophyte	*Mortierella humilis*	KP714537	100%	100%
F_0034*	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	*Cladophialophora*	Saprotroph/Endophyte	*Cladophialophora chaetospira*	KF359558	100%	99%
F_0079*	Ascomycota	Sordariomycetes	Hypocreales	Nectriaceae	-	Saprotroph/Endophyte	*Ilyonectria protearum*	NR_152890	99%	100%
F_0015*	Ascomycota	-	-	-	-	Others_Unknown	*Cadophora orchidicola*	KX611558	100%	99%
F_0202*	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	*Cladophialophora*	Saprotroph/Endophyte	*Cladophialophora chaetospira*	HQ871875	100%	99%
F_0195*	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	*Cladophialophora*	Saprotroph/Endophyte	*Cladophialophora chaetospira*	EU035405	100%	100%
F_0181*	Ascomycota	Leotiomycetes	Helotiales	Dermateaceae	*Pezicula*	Endophyte	*Pezicula melanigena*	LC206665	100%	99%
F_0010	Ascomycota	Leotiomycetes	-	Myxotrichaceae	*Oidiodendron*	Ericoid_Mycorrhizal	*Oidiodendron maius*	LC206669	100%	100%
F_0103*	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	*Cladophialophora*	Saprotroph/Endophyte	*Cladophialophora chaetospira*	EU035403	100%	97%
F_0489*	-	-	Mortierellales	Mortierellaceae	*Mortierella*	Saprotroph/Endophyte	*Mortierella* sp.	KM113754	100%	100%

Southern 4 sites (warm-temperate and subtropical)

ID	Phylum	Class	Order	Family	Genus	Guild	Species	Accession	%	%
F_0381*	Basidiomycota	Tremellomycetes	Trichosporonales	Trichosporonaceae	*Cryptococcus*	Others_Unknown	*Saitozyma podzolica*†	KY320605	92%	99%
F_0042*	-	-	Mortierellales	Mortierellaceae	*Mortierella*	Saprotroph/Endophyte	*Mortierella humilis*	KP714537	100%	100%
F_0610*	Ascomycota	Sordariomycetes	Hypocreales	Hypocreaceae	*Trichoderma*	Saprotroph/Endophyte	*Trichoderma spirale*	KU948158	100%	100%
F_1188*	Basidiomycota	Tremellomycetes	Trichosporonales	Trichosporonaceae	*Cryptococcus*	Others_Unknown	*Saitozyma podzolica*†	KY320605	92%	99%
F_0029	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	-	Others_Unknown	*Cladophialophora* sp.	LC189029	100%	99%
F_0017	Ascomycota	-	-	-	-	Others_Unknown	*Scleropezicula* sp.	KT809119	100%	98%
F_0007	Ascomycota	Sordariomycetes	Diaporthales	Melanconidaceae	*Melanconiella*	Saprotroph/Endophyte	*Melanconiella elegans*	KJ173701	100%	85%
F_0485	Ascomycota	Sordariomycetes	Hypocreales	Hypocreaceae	*Trichoderma*	Saprotroph/Endophyte	*Trichoderma* sp.	HG008760	100%	100%
F_0112	Basidiomycota	Agaricomycetes	Thelephorales	Thelephoraceae	*Tomentella*	Ectomycorrhizal	*Tomentella stuposa*	KR019860	100%	98%
F_0073	Ascomycota	Sordariomycetes	-	-	-	Others_Unknown	*Rhexodenticula acaciae*	KY173442	94%	95%

*Fungal OTUs classified as metacommunity hubs (mean local betweenness > 0.5; metacommunity betweenness > 0.5)

†Synonym, *Cryptcoccus podzolica*

Fig. 1 Study sites examined in this study. Across the entire range of the Japanese Archipelago, root samples were collected in four cool-temperate forests (sites 1–4), one warm-temperate forest (site 5), and three subtropical forests (sites 6–8).

Fig. 2 Compositions of fungal taxa and functional groups in each forest. **a** Order-level taxonomic composition of fungal OTUs in each locality. The number of fungal OTUs detected is shown in a parenthesis for each forest. **b** Functional-group composition. The fungal functional groups were inferred by the program FUNGuild [63].

Fig. 3 Fungal OTUs with highest local betweenness. **a** Order-level taxonomic composition of top-20 OTUs with highest local betweenness in each forest. See Data S3 (Additional file 3) for betweenness scores of all fungal OTUs in respective local forests. **b** Genus-level taxonomic composition of top-20 OTUs with highest local betweenness. **c** Functional-group composition of top-20 OTUs with highest local betweenness.

- ● Plant
- ● Arbuscular mycorrhizal
- ● Ectomycorrhizal
- ● Ericoid mycorrhizal
- ● Saprotroph / endophyte
- ● Plant pathogen
- ● Other / Unknown

Fig. 4 Metacommunity-scale network including all the eight local forests. The size of circles roughly represents relative scores of betweenness centrality. The functional groups of fungi inferred with the program FUNGuild [63] were organized into six categories: i.e., arbuscular mycorrhizal (bue), ectomycorrhizal (red), ericoid mycorrhizal (skyblue), saprotrophic/endophytic (yellow), plant pathogenic (purple), and other/unknown fungi (grey) (Additional file 3; Data S3). For plant species/taxa (green), the geographic information of source populations is indicated in Additional file 6 (Figure S3).

Fig. 5 Relationship between local- and metacommunity-level betweenness. **a** Full meatcommunity. On the horizontal axis, the mean values of betweenness centrality scores across all the eight local forests are shown for respective fungal OTUs. On the vertical axis, the betweenness scores within the metacommunity-scale network consisting of the eight localities (Fig. 4) are shown for respective OTUs. **b** Metacommunity of cool-temperate forests. For the sub-dataset consisting of the four cool-temperate forests (Additional file 7: Figure S4), mean local betweenness and metacommunity betweenness are shown on the horizontal and vertical axes, respectively. **c** Metacommunity of warm-temperate and subtropical forests. For the sub-dataset consisting of the warm-temperate forest and the three subtropical forests (Additional file 8: Figure S5), mean local betweenness and metacommunity betweenness are shown on the horizontal and vertical axes, respectively.