Parentage assignment with genotyping-by-sequencing data

Andrew Whalen*, Gregor Gorjanc, and John M Hickey

The Roslin Institute and Royal (Dick) School of Veterinary Studies, The University of Edinburgh,

Midlothian, Scotland, UK

*Corresponding author

Email addresses:

AW: awhalen@roslin.ed.ac.uk

GG: gregor.gorjanc@roslin.ed.ac.uk

JMH: john.hickey@roslin.ed.ac.uk

Abstract

In this paper we evaluate using genotype-by-sequencing (GBS) data to perform parentage

assignment in lieu of traditional array data. The use of GBS data raises two issues: First, for low-

coverage GBS data, it may not be possible to call the genotype at many loci, a critical first step for

detecting opposing homozygous markers. Second, the amount of sequencing coverage may vary

across individuals, making it challenging to directly compare the likelihood scores between

putative parents. To address these issues we extend the probabilistic framework of Huisman (2017)

and evaluate putative parents by comparing their (potentially noisy) genotypes to a series of

proposal distributions. These distributions describe the expected genotype probabilities for the

relatives of an individual. We assign putative parents as a parent if they are classified as a parent

(as opposed to e.g., an unrelated individual), and if the assignment score passes a threshold. We

evaluated this method on simulated data and found that (1) high-coverage GBS data performs

similarly to array data and requires only a small number of markers to correctly assign parents and

(2) low-coverage GBS data (as low as 0.1x) can also be used, provided that it is obtained across a

large number of markers. When analysing the low-coverage GBS data, we also found a high

number of false positives if the true parent is not contained within the list of candidate parents, but

that this false positive rate can be greatly reduced by hand tuning the assignment threshold. We

provide this parentage assignment method as a standalone program called AlphaAssign.

Introduction

In this paper we evaluate the performance of using genotype-by-sequence (GBS) data to

perform parentage assignment in commercial plant and animal breeding settings. Having accurate

parentage information is important for many routine breeding applications, such as reducing the

cost of genotyping through pedigree-based imputation (Huang et al., 2012), reducing the bias of

genomic estimates of breeding values (Solberg et al., 2009), and combining genotyped and non-

genotyped individuals into a joint analysis (Legarra et al., 2009). When the parents of an individual

are not recorded, parentage assignment algorithms can use genetic data to reconstruct parent-child

relationships. Much of the previous work on parentage assignment has focused on the case where

the genetic data was generated from microsatellite markers or more recently from SNP arrays

(Rohrer et al., 2007; Fisher et al., 2009; Riester et al., 2009; Tokarska et al., 2009). In the case of

SNP arrays between 50 and 700 markers are required to accurately assign parents and rule out

false assignments (Rohrer et al., 2007; Strucken et al., 2016; Fisher et al., 2009; Tortereau et al.,

2017). GBS is a flexible alternative to arrays, particularly for species that may not have a well-

established reference genome, or where a suitable array has not been developed. However, the

performance of using GBS data for parentage assignment – to our knowledge – is not well

understood.

The primary challenge for using GBS data is the potentially high uncertainty in the true

genotype of an individual based on the observed genetic data. In a GBS platform, a restriction

enzyme is used to cut DNA into fragments that are then sequenced (Baird et al., 2008; Davey et

al., 2011; Elshire et al., 2011). This means that unlike arrays, which produce called genotypes,

GBS produces read counts for the reference and alternative alleles. For high-coverage GBS data

the underlying genotype can easily be called from the read counts. For low-coverage GBS data

calling genotypes is more difficult, particularly on loci which only receive a few reads.

Distinguishing between heterozygous and homozygous loci is particularly challenging. If GBS

produces two reads for the reference alleles and zero reads for the alternative allele, this could

indicate that the individual is homozygous for the reference allele, or the individual could be

heterozygous and their reference allele was sequenced twice. The difficulty in calling homozygous

loci makes parentage assignment particularly difficult because many parentage assignment

algorithms, either explicitly or implicitly, rely on finding opposing homozygous loci to filter out

putative parents. In addition, the lack of opposing homozygous loci may increase false positive

rate of parentage assignment if the true parent is not in the list of putative parents, since full sibs

or half sibs of the true parent may appear to be more related to the individual than expected by

chance (Meagher and Thompson, 1986).

Likelihood based methods (e.g., Kalinowski et al., 2007; Riester et al., 2009) are one

solution to handle genetic data with high uncertainty. In a likelihood based method, parentage

assignment is based on the likelihood of an individual's genotype conditioned on the putative

parent's genotype. If the genotypes of either the individual or the putative parent cannot be

assessed accurately, this likelihood score can be calculated by marginalizing over possible

genotypes. Likelihood methods work well in cases where all individuals have the same amount of

genetic data (e.g., same number of markers or sequencing coverage), but may break down when

individuals are genotyped at a different number of markers or at different coverage levels. An

example of this could be two putative parents with array data. Suppose the first putative parent

was genotyped at 50 markers that overlap with the child, and the second was genotyped at 1,000

markers that overlap with the child. If both parents were heterozygous at all loci and we assume

that the loci are not linked, then the likelihood value for the first parent would be $.5^{50}$ (each allele

having a 50% chance of being transmitted), whereas the likelihood value for the second parent

would be $.5^{1000}$. These likelihood values are hard to compare against each other because they are

calculated on different sets of markers. This problem can be solved by selecting a subset of markers

that are genotyped in all putative parents (which may drastically reduce the amount of information

available), or using the population allele frequency for the genotype at missing markers (which

disadvantages individuals with missing values).

A third option, that may be more appealing for GBS data, is to instead change the parentage

assignment problem into a relationship classification problem. With this framing, the goal of the

algorithm is to classify the relationship between each putative parent and the focal individual (e.g.,

parent, grandparent, sibling, child). A putative parent is then assigned as the parent, if they are

classified as a parent, pass an assignment threshold, and are the highest scoring parent out of the

list of putative parents (Huisman, 2017; Riester et al., 2009). One of the main advantages of this

approach is that the classification task (which is able to filter out most putative parents) only relies

on the genetic information available for an individual and a putative parent and does not require

direct comparison to other putative parents. This property is particularly appealing for GBS data

where the amount of information on each individual may differ depending on the genotyping

resources spent and the allele frequency of the loci with sequence reads.

In this paper we extend the parentage assignment method of Huisman (2017) to explicitly

handle GBS data. We then evaluated its performance in a simulated animal breeding population.

We found that, similar to array data, it is possible to obtain accurate parent assignment with a fairly

small number of sequence reads (e.g., 0.1x coverage), but that ruling out false positives is harder,

and that a sizeable number of false positives could occur for medium coverage (0.5-2x) GBS data

on a large number of linked markers.

Materials and Methods

Here we describe our approach for parentage assignment with GBS data. This work builds

closely on the probabilistic framework of Huisman (2017), but we present the full model for

completeness. To assign parents we first construct a series of proposal distributions for each

putative parent based on the genotypes of a focal individual and it's known relatives. These

proposal distributions describe the expected genotypes for a relative as a function of their

relationship with the focal individual (e.g., parent, full sib of the parent, unrelated). We then

classify each putative parent into one of these relationships, and if it is classified as a parent, and

the assignment score passes a threshold, we assign it as the parent. If there are multiple possible

parents, the highest scoring individual is assigned. Although this algorithm was originally designed

in the context of animals, it also works for diploid and allopolyploid plants.

To simplify the language, we assume that we are attempting to assign the father of a focal

individual. For a given focal individual i and its mother m we calculate the probability that the

putative parent f is the true father by:

$$p(h = father | g_f, g_i, g_m) = \frac{p(g_f | g_i, g_m, h = father) p(h = father)}{\sum_{h'} p(g_f | g_i, g_m, h') p(h')},$$ (1)

where g_x is the genotype of individual x, h is the relationship between the focal individual i and

the putative parent f, and the denominator is enumerated over the set of possible relationships h'.

In the case where the genotypes of the mother are unknown we assume that her genotype

probabilities are derived from Hardy-Weinberg Equilibrium.

In this paper we consider four possible relationships: that the putative parent is the true

father, a full sib of the true father, a half sib of the true father, or unrelated. The conditional

probability distributions for alternative relationships can be constructed via the generative

framework we provide below. To simplify calculations, we assume that *p(h')* is uniform over all

possible relationships. In addition, we assume all markers segregate independently allowing

$p(g_f|g_i, g_m, h)$ to be calculated as the product of the probability of the putative parent's genotype

at each marker k:

$$p(g_f|g_i, g_m, h) = \prod_k p(g_{f,k}|g_{i,k}, g_{m,k}, h).$$ (2)

131 In the case of array data, and particularly GBS data, our assessment of the true genotypes,

132 g_f, g_i, and g_m may be noisy. To account for this noise we marginalize across possible genotypes

133 based on observed genetic data $d = (d_i, d_f, d_m)$:

$$p(d_{f,k}|d_{i,k}, d_{m,k}, h) = \sum_{g_{f,k}} \sum_{g_{m,k}} \sum_{g_{i,k}} p(g_{f,k}|g_{i,k}, g_{m,k}, h)p(g_{f,k}|d_{f,k})p(g_{m,k}|d_{m,k})p(g_{i,k}|d_{i,k}).$$ (3)

This model requires the calculation of two terms: (1) the genotype probabilities conditional

on the observed data $p(g_{x,k}|d_{x,k})$ and (2) the proposal distribution for an individual's genotype

based on their relationship with the focal individual $p(g_{f,k}|g_{i,k}, g_{m,k}, h)$. We outline how to

calculate both terms below.

Evaluating genotype probabilities conditional on the observed data

In this model we assume that each marker is biallelic and has four possible phased

genotypes, aa, aA, Aa, AA. With observed array data for marker k, $d_{x,k}$, the conditional

probabilities for each genotype $g_{x,k}$ are:

$$p(g_{x,k}|d_{x,k}) = \begin{cases} 1 - \dfrac{3e}{4} & \textit{if } g_{x,k} = aa \textit{ and } d_{x,k} = 0 \\[2mm] 1 - \dfrac{3e}{4} & \textit{if } g_{x,k} = AA \textit{ and } d_{x,k} = 2 \\[2mm] .5 - \dfrac{e}{4} & \textit{if } g_{x,k} = aA \textit{ or } Aa \textit{ and } d_{x,k} = 1 \\[2mm] \dfrac{e}{4} & \textit{otherwise,} \end{cases}$$ (4)

where e is the assumed genotyping error rate. This evaluation of individual genotype probabilities

differs from Huisman (2017), where it is assumed that errors can only occur between homozygous

and heterozygous states (and not between opposing homozygote states) and distinction is not made

between two heterozygous genotypes. The genotype probabilities above correspond more closely

to those commonly used in peeling (e.g., Whalen et al., 2017) and allow inferences to be made

even when the genotyping error rate is high.

With observed GBS data for marker k, $d_{x,k}$, the conditional genotype probabilities are:

$$p(g_{x,k}|d_{x,k}) \propto \begin{cases} (1-e)^{n_{ref}} e^{n_{alt}} & if\ g_{x,k} = aa \\ \dfrac{.5^{n_{ref}+n_{alt}}}{2} & if\ g_{x,k} = aA\ or\ Aa \\ (1-e)^{n_{alt}} e^{n_{ref}} & if\ g_{x,k} = AA, \end{cases} \qquad (5)$$

where e is the sequencing error rate, n_{ref} is the number of sequence reads supporting the reference

allele and n_{alt} is the number of sequence reads supporting the alternative allele. The genotype

probabilities in Equation 5 do not sum to one, and so the probabilities need to be normalized for

each allele. Equation 4 is consistent with previous work on parentage assignment with array data

(Kalinowski et al., 2007; Huisman, 2017), while Equation 5 is consistent with previous work on

imputation with GBS-like data (Li et al., 2010; VanRaden et al., 2015; Whalen et al., 2017).

Generating proposal distributions via single locus peeling

We generate proposal distributions $p(g_{f,k}|g_{i,k}, g_{m,k}, h)$ for the genotype probabilities of

each relationship via single locus peeling (Elston and Stewart, 1971). Single locus peeling provides

a rich generative model for estimating the genotype probabilities of un-genotyped relatives based

on the genotypes of an individual and a known parent. Although our presentation differs from

Huisman (2017) it results in the same distributions. Under this framework, we calculate the

genotype probabilities for three relatives: the father, a full-sib of the father, and a half-sib of the

father. These probabilities are calculated by first estimating the genotype probabilities for the

father, peeling up to the paternal grandparents, and finally peeling down to the full sib and the half

sib of the father (Figure 1).

Given genetic data on the focal individual d_i and a mother d_m, we can construct a proposal

distribution for the father via:

$$p(g_f|d_m, d_i) \propto \sum_{g_m} \sum_{g_i} T(g_i|g_f, g_m)p(g_i|d_i)p(g_f|d_f), \qquad (6)$$

173 where $p(g_i|d_i)$ is given by Equation 4 or 5 above, and $T(g_i|g_f, g_m)$ is the probability that the

174 individual inherited genotype g_i conditional on their parents having genotypes g_f and g_m, e.g.,

175 $T(g_i = aA|g_f = aA, g_m = AA) = 0.5$ (Marshall et al., 2003).

176 Using Equation 6, we can peel up to construct a joint distribution for the genotypes of the

177 paternal grandparents (gf, gm):

$$p(g_{gf}, g_{gm}|d_i, d_f) \propto \sum_{g_f} T(g_f|g_{gf}, g_{gm})p(g_f|d_i, d_f), \qquad (7)$$

179 where $p(g_f|d_i, d_f)$ is given in Equation 6, above. We can then peel down to generate the proposal

180 distributions for a full sib and a half sib of the father. The proposal distributions differs in whether

181 the full joint distribution of both grandparents is used (full sib, fs), or if only one of the grandparents

182 is used and the other parent assumed to have genotypes based on Hardy Weinberg Equilibrium

183 (half sib, hs):

$$p(g_{fs,}|g_{gf}, g_{gm}, d_i, d_f) = \sum_{g_f} T(g_{fs}|g_{gf}, g_{gm})p(g_{gf}, g_{gm}|d_i, d_f), \qquad (8)$$

$$p(g_{hs,}|g_{gf}, g_{gm}, d_i, d_f) = \sum_{g_{null}} \sum_{g_{gd}} T(g_{hs}|g_{gf}, g_{null})p(g_{gf}, g_{gm}|d_i, d_f)p(g_{null}), \qquad (9)$$

186 where $p(g_{null})$ represents the probability of having a genotype if that genotype was drawn at

187 random from the population.

188   The proposal distribution for an unrelated individual simply assumes that their genotypes

189   are drawn at random from the population according to Hardy Weinberg Equilibrium:

190                                   $$p(g_{unrelated}) = p(g_{null}).$$ (9)

191   To assign a parent we calculated an assignment score for each putative parent:

192                          $$score = -\log{(1 - p(h = father|d_f, d_i, d_m))}.$$ (10)

The score will be close to 0 if the individual is unlikely to be the father, and tends towards positive

infinity with increasing evidence that the individual is the father. A putative parent was assigned

as the true parent if its assignment score was the highest of the putative parents considered, and

was higher than a pre-defined threshold.

Although the described process may seem computationally intensive, there are two features

which simplify calculations. First, because the proposal distributions depend only on the focal

individual and its known parent, the proposal distributions only need to be calculated once and can

be re-used for all putative parents of the focal individual. Second, peeling can be performed

efficiently as a series of tensor operations on the genotypes of focal individual and its known

parent, filtered through the inheritance matrix T, which allows us to take advantage of linear

algebra libraries.

Simulated data

The simulated data modelled a livestock population. We initially sampled a set of genomes

with 20 chromosomes using the Markovian coalescent simulator MaCS (Chen et al., 2009). For

this we assumed that each chromosome is 10^8 bp long, a per site mutation rate is 2.5×10^{-8}, a per

site recombination rate is 1.0×10^{-8}, and that effective population size changed over time. Based

on estimates for the Holstein cattle population (Villa-Angulo et al., 2009), we set the effective

population size to 100 in the final generation of the coalescent simulation and to 1,256, 4,350, and

43,500 at respectively 1000, 10,000, and 100,000 generations ago, with linear changes in between.

We then used the sampled chromosomes to initiate a population of 1,000 animals with equal sex

proportions. We bred this population for 5 generations. In each generation, we selected 10 males

and mated them at random to 100 females. Each potential focal individual therefore had 1 true

father, 4 male full sibs of the father, and 45 male half sibs of the father. All individuals were

genotyped at 50,000 markers. Subsets of these markers were used in different simulations as

described below. Array data were simulated without any errors, due to the low error rate for

modern SNP genotyping arrays (<1%; e.g., Kalinowski et al., 2007). In addition to array data, we

generated low-coverage GBS data for the last two generations of individuals. We assumed that the

GBS method targeted the same loci as the genotyping array and was performed at coverage levels

between 0.1x to 10x. For each coverage level, the number of sequence reads at a given marker was

generated via a Poisson distribution with mean equal to the coverage level. Each read randomly

sampled one of the two alleles at a marker. The read sampling process also included a small

sequencing error rate of 0.1%. We generated the simulated data using the R package AlphaSimR

(Gaynor et al.), which is available at www.alphagenes.roslin.ed.ac.uk/AlphaSimR.

Scenarios

227   We evaluated the accuracy of parent assignment for the last generation of 1,000 individuals

across 4 different scenarios. In the first scenario (a) we analysed the accuracy of performing parent

assignment when:

•   the mother was known and genotyped,

•   all of the male full- and half-sibs along with 50 other individuals (total of 100 potential

parents) were putative parents,

• and either both the parents and progeny had array data, the parents had array data and the

progeny had GBS data, or both the parents and the progeny had GBS data.

These sub-scenarios span a spectrum of possible practical settings. The sub-scenario where the

parents had array data but the progeny had GBS data may represent either the case where the

progeny are initially genotyped with a low-cost GBS platform and any selected parents are re-

genotyped with an array, or it may represent the case where pedigree information was used to

impute and accurately call parental genotypes. In the remaining scenarios we focused on the case

where both parents and progeny had GBS data and analysed (b) the impact of knowing and

genotyping the known alternative parent, (c) the impact of restricting the pool of putative parents

to either 100 unrelated individuals, 45 half sibs, or the 4 full sibs, and (d) examined how the false

positive rate changed depending on the threshold used for assignment (see below).

In each scenario we performed three evaluations. To evaluate the overall accuracy, we

assumed the true parent was included in the list of putative parents, and evaluated accuracy by the

number of times the top parent was the true parent. To evaluate the true positive rate we included

the true parent in the list of putative parents, but assigned the top scoring parent only if it passed

an assignment threshold. To evaluate the false positive rate, we excluded the true parent from the

list of putative parents and counted the number of times the top scoring parent passed the

assignment threshold. The first evaluation represented a case where we know the true parent is

included in the list of putative parents (e.g., groups of females cohabitating with multiple males or

artificial insemination using polyspermic matings). The second and third evaluations were

designed to assess performance when we are not sure whether or not the true parent is included in

the list of potential parents (e.g., natural service sires or wild populations).

Software

Parentage   assignment   was   performed   using   AlphaAssign

(http://www.alphagenes.roslin.ed.ac.uk/alphasuite-softwares/alphaassign/) which, implements the

described algorithm. AlphaAssign has three run-time parameters: (i) an assumed genotyping error

rate for array data, (ii) an assumed sequencing error rate for GBS data, and (iii) an assignment

threshold to determine the required score to assign a putative parent as a parent. Throughout this

paper we assumed a 1% genotyping error rate, a 0.1% sequencing error rate, an assignment

threshold of 10 (determined via pilot simulations) although we varied the assignment threshold in

the last set of simulations.

Results

Parent assignment with array and GBS data

First we examined the number of markers required for accurate parentage assignment when both

parents and progeny were genotyped with array data. If the true parent was included in the list of

putative parents (and an assignment threshold was used), 100 markers were required to obtain

100% parentage assignment accuracy. If the true parent was excluded from the list of putative

parents, the false positive rate was less than 0.1% if there were between 50 to 350 markers, and

there were no false positives when there were more than 500 markers.

Unlike array data where the number of markers can be more easily varied, for GBS data

the number of markers is usually determined by the choice of restriction enzymes while the amount

of coverage obtained on each individual can be varied. Because of this we focused on the required

coverage level to accurately assign parents based on a fixed number of markers. Figure 2 shows

the accuracy and false positive rates based on the amount of coverage allocated to each progeny,

stratified by the number of markers that this coverage is spread over. Because performance with

array data was nearly identical to that with 10x GBS data we did not include array data in Figure

2.

We evaluated the performance of parentage assignment when the parents were genotyped

with array data and the progeny were genotyped with GBS data. If the true parent was included in

the list of putative parents, a coverage of 0.4x was required to obtain 100% accuracy when there

were 50,000 GBS markers. The required coverage increased to 1x for 5,000 markers, and to 2x for

1,000 markers. If the true parent was excluded from the list of putative parents, we found that the

false positive rate was less than 0.2% in all cases.

The accuracy of parentage assignment decreased when both the parents and progeny had

GBS data. If the true parent was included in the list of putative parents, a coverage of 0.4x was

required to obtain 100% accuracy when there were 50,000 GBS markers. The required coverage

increased to 2x for 5,000 markers, and to 5x for 1,000 markers. If the true parent was excluded

from the list of putative parents, we found that the false positive rate was as high as a 60%. These

false positives were clustered on low to medium coverage GBS data (0.1 - 3x) with a large number

of markers (>1000).

False positive assignments by relationship

Figure 3 stratifies the false positive rate based on whether unrelated individuals, half-sibs

of the true parent, or full-sibs of the true parent were included in the list of putative parents. In line

with expectations we found a high false positive rate (as high as 60% in some conditions) when

only the full-sibs of the true parent were included as putative parents. This decreased to at most

35% when only the half-sibs of the true parent were included and to under 20% when only

unrelated individuals were included. As seen previously, most of the false positives were occurred

when there were a large number of markers and low to medium coverage GBS data.

Parent assignment when neither parent is known

Figure 4 compares the performance of parentage assignment when one of the parents is

known and genotyped compared to when neither parent is known or genotyped. We found that

having one parent known and genotyped increased the accuracy of parentage assignment and

decreased the number of false positives in all cases. The benefit was largest when both the progeny

and parents had high-coverage GBS data.

Controlling false assignments by modifying the threshold

Figure 5 shows the true positive rates and false positive rates for when sequencing

resources were spread over 50,000 markers, as a function of the threshold used to assign a putative

parent as the parent. We found that, compared to the results in Figures 2 and 3, it was possible to

substantially reduce the false positive rate by increasing the assignment threshold, but that the ideal

threshold depends on the total coverage. The relationship between the false positive and true

positive rate is given as a receiver operating characteristic in Figure 5(c).

Timing

The algorithm took 3 minutes and 54 seconds to assign parents for 1000 progeny, each

with 100 putative parents. The progeny and their parents were genotyped using GBS data across

5,000 markers. The algorithm scales linearly with the number of markers and the number of

putative parents per individual.

Discussion

In this paper we extended the parentage assignment method of Huisman (2017) to account

for low-coverage sequence data and analysed the performance of parentage assignment when

genotyping is performed via sequencing instead of the traditional genome-wide arrays. We found

that high-coverage GBS data (i.e., 10x or higher) has the same performance as array data. We also

found that low-coverage GBS data (as low as 0.1x) can be used to perform parentage assignment

as long as it is obtained on a sufficiently large number of markers, but that there may be a large

number of false assignments if the true parent is not included in the list of putative parents. The

number of false positives could be reduced by modifying the threshold used to call assignments.

In light of these results, we will discuss (1) the accuracy of parentage assignment, (2) potential

extensions to control the false positive rate, and (3) the use of peeling to construct the proposal

distributions in more detail.

Parentage assignment accuracy with GBS data

A goal of this work was to quantify the amount of GBS data required to accurately perform

parentage assignment. We found that, similar to array data, the total amount of data required is

relatively low. For example, when using high-coverage GBS data between 100 to 200 markers are

required to accurately assign parents. This is in line with previous estimates for array data (Rohrer

et al., 2007; Strucken et al., 2016; Fisher et al., 2009; Tortereau et al., 2017), where between 50-

700 markers were required. The differences in the exact number of markers required (100-200

compared to 50-700) is likely due to the structure of the underlying genetic data (i.e., number of

chromosomes, minor allele frequency of the markers), and the assumption in this study that one of

the parents was already known and genotyped.

In addition to being able to use high-coverage GBS data to perform parentage assignment,

we found that low-coverage GBS data could also be used, provided it was spread across a larger

number of markers. The increase in required number of markers is due to the lower information

content at an individual loci for low-coverage GBS data, requiring the data to be pooled across a

larger number of markers to achieve the same level of accuracy.

The results of this study suggest that GBS data – either high-coverage data on a small

number of markers, or low-coverage data on a large number of markers – is an effective alternative

to array data for performing parentage assignment. This result is particularly important given the

emerging importance of GBS as an alternative for SNP array data, both in species where SNP

arrays are available (e.g., De Donato et al., 2013; Brouard et al., 2017) and in those where SNP

arrays have not been constructed (e.g., Robledo et al., 2017; Palaiokostas et al., 2018).

Controlling the false positive rate

During our analysis of low-coverage data, we found an inflation of false positives when

both the parents and the progeny had GBS data. These false positives were likely due to the fact

that with between 1-3x coverage GBS data we were able to determine that two animals are

genetically similar, but were not able to obtain a sufficient number of loci with precisely inferred

genotyped to find opposing homozygous loci.

Consistent with previous work, we found that using a hand-tuned assignment threshold

could reduce the number of false positives (Huisman, 2017; Riester et al., 2009). An alternative

approach would be to adaptively determine the assignment threshold via introspection of the

underlying data (Grashei et al., 2018). In the majority of the simulations, a fixed threshold of 10

was used based on pilot simulations with array data. As we demonstrate in Figure 5, substantially

raising the threshold for assignment could reduce the false-positive rate even for 50,000 markers

and low-coverage sequence data, although at the cost of a decreased true-positive rate. The optimal

threshold value for assignment depends on the overall sequencing coverage, making it challenging

to use a fixed threshold in cases where individuals are sequenced at different coverages. We believe

that automating this process is an area for future research, and may depend on the exact breeding

program structure, the exact GBS system deployed (e.g., Baird et al., 2008; Davey et al., 2011;

Elshire et al., 2011), and reason that parentage information is required.

Furthermore, we believe that the issue of false parent-assignments may be less of an issue

in the context of commercial agricultural populations compared to wild populations for two

reasons. First, most of the false assignments that we observed were cases where the true parent

was not included in the pedigree and a full- or half-sib of the true parent was included and wrongly

assigned as a parent. In the context of many animal breeding programs, the routine use of pairs of

sibs as parents may not commonly arise because of explicit efforts to manage diversity and

inbreeding (e.g., Woolliams et al., 2015). Second, due to the genetic similarity between the full-

sib of the true parent and the true parent, using the full-sib of the true parent as a "proxy" parent

for the progeny may have limited impact on downstream applications such as estimation of

breeding values. Further research is required to quantify the impact of such false positives in

downstream applications.

Constructing proposal distributions via peeling

In this paper, we closely followed the approach of Huisman (2017) for performing

parentage assignment, with two differences. First, we modified the genotype probability function

to handle sequence data. Second, we recast the construction of proposal distributions for relatives

as a series of peeling operations on artificial pedigrees. We believe the later development is of

more interest. Peeling provides a rich and computationally efficient framework for estimating the

genotypes of a relative based on the genotypes of individuals in an existing pedigree. In this paper

we focused on a small number of possible relationships, but this framework can be easily extended

to consider a wider and potentially complex class of relatives (e.g., siblings of the focal individual,

cousins of the parent, or grandparents), or could be altered to assess alternative relationships (e.g.,

performing grandparent assignment instead of parentage assignment). Use of these additional

relationship classes may depend on the purpose of a particular application.

Conclusion

In conclusion, we extended the algorithm of Huisman (2017) to perform parentage

assignment with sequence data, and evaluated the performance of using low-coverage GBS data

for parentage assignment. We found that low-coverage GBS data could be used for accurate

parentage assignment, but that there may be concerns with false positives if the true parent is not

included on the list of putative parents. Such false positives might be mitigated on a case-by-case

basis by tuning the assignment criteria used. These results suggest that GBS data can be used as

an alternative to array data for parentage assignment.

Conflicts of interest

The authors declare they have no competing interests.

Acknowledgements

The authors acknowledge the financial support from the BBSRC ISPG to The Roslin

Institute BB/J004235/1, from Genus PLC and from Grant Nos. BB/M009254/1, BB/L020726/1,

BB/N004736/1, BB/N004728/1, BB/L020467/1, BB/N006178/1 and Medical Research Council

(MRC) Grant No. MR/M000370/1. This work has made use of the resources provided by the

Edinburgh Compute and Data Facility (ECDF) (http://www.ecdf.ed.ac.uk).

References

Baird, N.A., Etter, P.D., Atwood, T.S., Currey, M.C., Shiver, A.L., Lewis, Z.A., Selker,

E.U., Cresko, W.A., and Johnson, E.A. (2008). Rapid SNP discovery and genetic mapping using

sequenced RAD markers. PLoS ONE *3*, e3376.

Brouard, J.-S., Boyle, B., Ibeagha-Awemu, E.M., and Bissonnette, N. (2017). Low-depth

genotyping-by-sequencing (GBS) in a bovine population: strategies to maximize the selection of

high quality genotypes and the accuracy of imputation. BMC Genet. *18*, 32.

Chen, G.K., Marjoram, P., and Wall, J.D. (2009). Fast and flexible simulation of DNA

sequence data. Genome Res. *19*, 136–142.

Davey, J.W., Hohenlohe, P.A., Etter, P.D., Boone, J.Q., Catchen, J.M., and Blaxter, M.L.

(2011). Genome-wide genetic marker discovery and genotyping using next-generation

sequencing. Nat. Rev. Genet. *12*, 499–510.

De Donato, M., Peters, S.O., Mitchell, S.E., Hussain, T., and Imumorin, I.G. (2013).

Genotyping-by-sequencing (GBS): A novel, efficient and cost-effective genotyping method for

cattle using next-generation sequencing. PLoS One *8*, e62137.

Elshire, R.J., Glaubitz, J.C., Sun, Q., Poland, J.A., Kawamoto, K., Buckler, E.S., and

Mitchell, S.E. (2011). A robust, simple genotyping-by-sequencing (GBS) approach for high

diversity species. PLoS One *6*, e19379.

Elston, R.C., and Stewart, J. (1971). A general model for the genetic analysis of pedigree

data. Hum. Hered. *21*, 523–542.

Fisher, P.J., Malthus, B., Walker, M.C., Corbett, G., and Spelman, R.J. (2009). The

number of single nucleotide polymorphisms and on-farm data required for whole-herd parentage

testing in dairy cattle herds. J. Dairy Sci. *92*, 369–374.

Gaynor, R.C., Gorjanc, G., Wilson, D.L., Money, D., and Hickey, J.M. AlphaSimR: An

R Package for Breeding Program Simulations. Manuscr. Prep.

Grashei, K.E., Ødegård, J., and Meuwissen, T.H.E. (2018). Using genomic relationship

likelihood for parentage assignment. Genet. Sel. Evol. *50*, 26.

Huang, Y., Hickey, J.M., Cleveland, M.A., and Maltecca, C. (2012). Assessment of

alternative genotyping strategies to maximize imputation accuracy at minimal cost. Genet. Sel.

Evol. *44*, 25.

Huisman, J. (2017). Pedigree reconstruction from SNP data: parentage assignment,

sibship clustering and beyond. Mol. Ecol. Resour. *17*, 1009–1024.

Kalinowski, S.T., Taper, M.L., and Marshall, T.C. (2007). Revising how the computer

program cervus accommodates genotyping error increases success in paternity assignment. Mol.

Ecol. *16*, 1099–1106.

Legarra, A., Aguilar, I., and Misztal, I. (2009). A relationship matrix including full

pedigree and genomic information. J. Dairy Sci. *92*, 4656–4663.

Li, Y., Willer, C.J., Ding, J., Scheet, P., and Abecasis, G.R. (2010). MaCH: using

sequence and genotype data to estimate haplotypes and unobserved genotypes. Genet.

Epidemiol. *34*, 816–834.

Marshall T. C., Slate J., Kruuk L. E. B., and Pemberton J. M. (2003). Statistical

confidence for likelihood-based paternity inference in natural populations. Mol. Ecol. *7*, 639–

655.

Meagher, T.R., and Thompson, E. (1986). The relationship between single parent and

parent pair genetic likelihoods in genealogy reconstruction. Theor. Popul. Biol. *29*, 87–106.

Palaiokostas, C., Cariou, S., Bestin, A., Bruant, J.-S., Haffray, P., Morin, T., Cabon, J.,

Allal, F., Vandeputte, M., and Houston, R.D. (2018). Genome-wide association and genomic

prediction of resistance to viral nervous necrosis in European sea bass (Dicentrarchus labrax)

using RAD sequencing. Genet. Sel. Evol. *50*, 30.

Riester, M., Stadler, P.F., and Klemm, K. (2009). FRANz: reconstruction of wild multi-

generation pedigrees. Bioinformatics *25*, 2134–2139.

Robledo, D., Palaiokostas, C., Bargelloni, L., Martínez, P., and Houston, R. (2017).

Applications of genotyping by sequencing in aquaculture breeding and genetics. Rev. Aquac. *10*,

670–682.

Rohrer, G.A., Freking, B.A., and Nonneman, D. (2007). Single nucleotide

polymorphisms for pig identification and parentage exclusion. Anim. Genet. *38*, 253–258.

Solberg, T.R., Sonesson, A.K., Woolliams, J.A., Ødegard, J., and Meuwissen, T.H.

(2009). Persistence of accuracy of genome-wide breeding values over generations when

including a polygenic effect. Genet. Sel. Evol. *41*, 53.

Strucken, E.M., Lee, S.H., Lee, H.K., Song, K.D., Gibson, J.P., and Gondro, C. (2016).

How many markers are enough? Factors influencing parentage testing in different livestock

populations. J. Anim. Breed. Genet. *133*, 13–23.

Tokarska, M., Marshall, T., Kowalczyk, R., Wójcik, J.M., Pertoldi, C., Kristensen, T.N.,

Loeschcke, V., Gregersen, V.R., and Bendixen, C. (2009). Effectiveness of microsatellite and

SNP markers for parentage and identity analysis in species with low genetic diversity: the case of

European bison. Heredity *103*, 326.

Tortereau, F., Moreno, C.R., Tosser-Klopp, G., Servin, B., and Raoul, J. (2017).

Development of a SNP panel dedicated to parentage assignment in French sheep populations.

BMC Genet. *18*, 50.

VanRaden, P.M., Sun, C., and O'Connell, J.R. (2015). Fast imputation using medium or

low-coverage sequence data. BMC Genet. *16*, 82.

Villa-Angulo, R., Matukumalli, L.K., Gill, C.A., Choi, J., Tassell, C.P.V., and

Grefenstette, J.J. (2009). High-resolution haplotype block structure in the cattle genome. BMC

Genet. *10*, 19.

Whalen, A., Ros-Freixedes, R., Wilson, D.L., Gorjanc, G., and Hickey, J.M. (2017).

Hybrid peeling for fast and accurate calling, phasing, and imputation with sequence data of any

coverage in pedigrees. BioRxiv.

Woolliams, J.A., Berg, P., Dagnachew, B.S., and Meuwissen, T.H.E. (2015). Genetic

contributions and their optimization. J. Anim. Breed. Genet. *132*, 89–99.

Figures

Figure 1. A graphical representation of the peeling order for the proposal distributions. The arrows

represent the direction in which the peeling operations should be performed. Hardy-Weinberg

equilibrium is used to generate the genotype distributions for the unrelated individual, the mother

of the half sib, and if unknown, the mother's genotype. Although this graphic assumes the mother

is known and the father unknown, a symmetric picture could be constructed when the mother is

unknown and father known.

Figure 2. Parentage assignment performance when array or GBS data was available for the parents and GBS data was available for the progeny. The left panels give the number of correct assignments (for 1000 progeny) when the true parent was on the list of putative parents and no assignment threshold was used – the top scoring parent was assigned. The middle panels give the number of correct assignments when the true parent was on the list of putative parents and assignment threshold was used. The right panels give the number of incorrect assignments when the true parent was excluded from the list of putative parents.

Figure 3. Number of false positive parentage assignments (for 1000 progeny) when GBS data was available for parents and progeny, the parent was excluded from the list of putative parents, assignment threshold was used, and the list of putative parents contained either 100 unrelated individuals (left panel), 45 half sibs of the true parent (middle panel), or 4 full sibs of the true parent (right panel).

Figure 4. A comparison between the parentage assignment performance with one parent known and genotyped and no parent known at different GBS coverage levels (left and middle panes compare true positives while the right pane compares false positives).

527   Figure 5. The rate of true positives, false positives, and the relationship between then when varying

528   the total amount of coverage and the calling threshold.