Evidence for sex microchromosomes in a species with temperature-

influenced sex determination (*Crotaphytus collaris*)

Authors: Jodie M. Wiggins[1], Enrique Santoyo-Brito[1], Jon B. Scales[2], Stanley F. Fox[1]

1. Department of Integrative Biology, Oklahoma State University, OK 74078, USA

2. Department of Biology, Midwestern State University, TX 76308, USA

Keywords

Sex-determining mechanism, genotypic sex determination, temperature-dependent sex

determination, gene-dosage, sex microchromosome, reptile, Crotaphytus collaris

Abstract

The characteristics of a species' evolution can be powerfully influenced by its mode of sex

determination and, indeed, sex determination mechanisms vary widely among eukaryotes. In

non-avian reptiles, sex was long thought to be determined bimodally, either by temperature or

genetics. Here we add to the growing evidence that sex determining mechanisms in reptiles fall

along a continuum rather than existing as a mutually exclusive dichotomy. Using qPCR, we

demonstrate that the lizard *Crotaphytus collaris* possesses sex-based gene dosage consistent with

the presence of sex michrochromosomes, despite that extreme incubation temperatures can

influence hatchling sex ratio. Our results suggest a temperature override that switches genotypic

females to phenotypic males at high and low temperatures.

Introduction

Mode of sex determination has far reaching consequences affecting evolutionary processes such as speciation (Haldane 1922) adaptive capability and heterozygosity (Bull 1983; Shine et al. 2002; Burt and Trivers 2006), maternal capacity for sex ratio manipulation (Kratochvíl et al. 2008), extent of secondary sexual characteristics (Reeve and Pfennig 2003) and capacity to respond to climate change (Mitchell and Janzen 2010). Among eukaryotes, the array of sex determination mechanisms (SDMs) is diverse (Bull 1983; Charlsworth 1996; Bachtrog et al. 2014). Environmental sex determination (ESD) is characterized by a mode of sex determination entirely dependent on environmental factors such as temperature encountered during embryogenesis (Merchant-Larios and Díaz-Hernández 2013), photoperiod (Walker 2005; Guler et al. 2012), or social cues during subsequent development (Bull 1983; Janzen and Paukstis 1991; Valenzuela and Lance 2004; Bachtrog et al. 2014) . Conversely, sex in GSD species is determined at conception by genes, and is uninfluenced by environment (Bull 1983; Sarre et al. 2004).

Long held was the belief that within reptilian lineages there existed a single dichotomy of mutually exclusive SDMs (Bull 1983; Janzen and Paukstis 1991). Either a species' sex was thought to be determined by the environment (environmental sex determination; ESD) or by sex chromosomes (genotypic sex determination; GSD) (Bull 1983; Janzen and Paukstis 1991; Bachtrog et al. 2014). In fact, reptilian species do utilize both male and female heterogamety (GSD) (King 1977) and temperature-dependent sex determination (TSD; a specific type of ESD) (Ewert and Nelson 1991). However, in recent years, the line between ESD and GSD has become decidedly blurred, and an increasingly complex picture is emerging in which GSD and TSD are the ends of a continuum of SDMs in reptiles. Examples of intermediate SDMs include species in

which different populations utilize different sex determining mechanisms (Pen et al. 2010),

temperature-dependent sex reversal of a species with a ZZ/ZW GSD system in the wild (Quinn

et al. 2007; Holleley et al. 2015), and revelations of the genetic underpinnings of TSD in

alligators and turtles (Spotila et al. 1998; Smith et al. 1999; Kettlewell et al. 2000; Western and

Sinclair 2001). This shifting landscape provides an exceptional study system for better

understanding sex determination in vertebrates (Sarre et al. 2004).

The collared lizard, *Crotaphytus collaris*, is a widespread, oviparous species in which sex

chromosomes have not been identified (Gorman 1973; De Smet 1981). Yet, *C. collaris* has been

classified as a GSD species based on phylogenetic analyses (Pokorna and Kratochvíl 2009).

However, as mentioned above, even members of the same species can utilize different sex

determining mechanisms (Pen et al. 2010). Thus, classifying *C. collaris* solely based on

phylogeny may provide an incorrect or incomplete picture (Viets et al. 1994). Compellingly, in

an investigation seeking to determine if *C. collaris* utilizes TSD, an inverse TSD type II pattern

was identified in which the percentage of female offspring declined as constant incubation

temperatures or treatments approached high and low extremes (Santoyo-Brito et al. 2017). While

the authors of this study point out that their sample size was low, this indicates a

temperature influence on sex determination in *C. collaris*. However, even at low and high

treatments this study did not find ratios of either sex nearing 100%. These findings hint at a more

complex mechanism for sex determination than pure TSD or pure GSD in *C. collaris,* as

suggested in other reptile species (Quinn et al. 2007; Radder et al. 2008; Holleley et al. 2015). In

species that possess XY sex chromosomes, the heterogametic sex is expected to have half the

dosage of X-linked genes (Rovatsos et al. 2014a). Indeed, sex-specific gene dosage at two loci in

the closely related *Crotaphytus insularis* points to heterogamety (males are heterogametic) and,

thus, to the possibility of GSD in Crotaphytids (Rovatsos et al. 2014b). The apparent conflict

between the findings of the above studies compels us to determine if *C. collaris* demonstrates

gene dosage akin to that identified in *C. insularis*.

Methods

DNA isolation and PCR

Blood was collected from the toes of twenty wild caught lizards (10 male, 10 female)

upon capture at Sooner Lake Dam, Pawnee Co., Oklahoma. Blood was immediately preserved

on Whatman FTA classic cards. DNA was later extracted from the cards by excising a 3-mm

square of blood-saturated card using sterile scissors then following the GE Healthcare extraction

protocol using Chelex® 100 resin. We tested for gene dosage in the genes ATP2A2

(sarcoplasmic reticulum calcium ATPase 2), TMEM (transmembrane protein 123D), and PEBP1

(phosphatidylethanolamine binding protein 1) (Table 1). Primer sequences for the genes TMEM,

PEBP1, and ATP2A2 were obtained from Rosavatos et al (2014; Table 1). The single copy gene

EF1α was used for gene dosage normalization. PCRs were assembled in 25-μl final volumes

containing 12.5 μl 2X Bullseye EvaGreen qPCR master mix buffer (MidSci, St. Louis, MO), 25

ng genomic DNA, and 200 pMoles forward and reverse primers. The thermal profile was an

initial denaturation of 10 min at 94˚C followed by 40 cycles of 94˚C for 30 sec, 55˚C for 30 sec,

72˚ C for 45 sec. Amplification via qPCR was executed in a Stratagene MX3005 thermocycler.

Gene Dosage Calculations

Final gene dosage ratios were calculated for the two male and two female lizards whose

DNA consistently and reliably amplified across three replicates. Crossing point values were

calculated in MaxPro (Stratagene) then normalized to EF1α. Gene dosage was calculated as in

Rosavotas et al. (2014b) with: $R = [2^{Cp\ gene}/2^{Cp\ EF1\alpha}]^{-1}$ and r (relative gene dosage ratio) =

R_{male}/R_{female}. It was expected that male *C. collaris* are the heterogametic sex based on results in

the closely related *Crotaphytus insularis* (Rovatsos et al. 2014b), thus an r = 0.5 is expected for

single copy, X-linked genes while r = 1.0 is expected for autosomal genes.

Results

Results demonstrate sex-linked differences for all three analyzed genes (ATP2A2,

TMEM, and PEBP1; Fig 1). In each case, the average female r value is exactly one. This result is

consistent with females having two copies of the gene of interest and, thus, being the

homogametic sex. For each gene an average value near 0.5 was obtained in the males (ATP2A2

= 0.59, TMEM = 0.61, PEBP1 = 0.59). This result is consistent with males having a single gene

copy and being heterogametic.

Discussion

Though sex chromosomes have not been detected in Crotaphytids (Gorman 1973; De

Smet 1981) our results are consistent with heterogamety and point to the existence of GSD in *C.

collaris* as in *C. insularis* (Rovatsos et al. 2014b). While GSD is common among lizards, *C.

collaris* has been shown to experience changes in sex ratios with varying incubation

temperatures (Santoyo-Brito et al. 2017). *Crotaphytus collaris* may be another in the emerging

examples of a reptilian species with GSD that can be overridden by temperature extremes (Shine

et al. 2002; Holleley et al. 2015). We agree with Santoyo-Brito et al. (2017) that *C. collaris*

likely possess sex microchromosomes in an XX/XY pattern and that extreme incubation

temperatures alter the sex determining pathway such that XX individuals develop as phenotypic

males. We further hypothesize that gravid *C. collaris* females select nest sites such that GSD will

function without temperature interference as evidenced by field hatchling ratios near 50/50 as

expected in GSD populations (Wiggins unpublished data).

Evidence continues to emerge that some extant reptile species employ multimodal sex

determining mechanisms (Shine et al. 2002; Valenzuela et al. 2003; Sarre et al. 2004; Quinn et

al. 2007; Radder et al. 2008; Holleley et al. 2015). These species may be at a transition point

from GSD to TSD. *C. collaris* demonstrates gene dosage consistent with that of 28 species of

Pleurodont iguanian lizards (Pleurodonta) spanning 11 genera and including *A. carolinensis* and

C. insularis (Rovatsos et al. 2014a,b). The absence of genes on the Y that are present on the X,

coupled with the chromosomal looping in the *A. carolinensis* XY synaptonemal complex, points

to a degenerate Y microchromosome in this species and, subsequently, those with the same sex-

based gene dosage (Alföldi et al. 2011; Bachtrog et al. 2014; Rovatsos et al. 2014a,b; Lisachov et

al. 2017). Thus, there exists the strong possibility that *C. collaris* also possesses a degenerate Y

microchromosome. Results from Santoyo-Brito et al. (2017) show a decline in number of

females hatched at extreme high and low temperatures, pointing to a temperature override that

converts XX individuals into phenotypic males (XXm). If these XXm are viable, they may mate

and reproduce (as males). While this scenario will lead to an increase in the proportion of genetic

females, the possibility of producing offspring who possess both degenerate chromosomes (YY)

would be avoided as in *Bassiana duperreyi* (Shine et al. 2002). As a result, GSD and TSD could

conceivably co-exist in *C. collaris* without a definitive transition to one or the other. However, as

global temperatures rise, frequent shifts of XXf to XXm could eliminate the Y

microchromosome, driving this species to pure TSD and increasing its vulnerability to extinction

by fixation of homogamety and absence of XY individuals (Mitchell and Janzen 2010).

In summary, our data convincingly point to the presence of as-yet unidentified sex

microchromosomes in *C. collaris* as suggested by Santoyo-Brito et al. (2017) and add to the

growing evidence that SDMs in non-avian reptiles are not bimodal (Shine et al. 2002; Holleley et

al. 2015). Further inquiry into the effects of extreme temperature incubations on *C. collaris* sex

determination is warranted (Santoyo-Brito et al. 2017); specifically, investigating whether some

of the individuals incubated at either high or low temperature extremes are genotypically female

but phenotypically male as suggested by Santoyo-Brito et al. (2017) and as shown in free-

ranging *P. vitticeps* (Holleley et al. 2015), if such individuals are reproductively viable, and if

they exist in natural populations.

Literature Cited

Alföldi, J., F. Di Palma, M. Grabherr, C. Williams, L. Kong, E. Mauceli, P. Russell, C. B. Lowe,

R. E. Glor, J. D. Jaffe, D. A. Ray, S. Boissinot, A. M. Shedlock, C. Botka, T. A. Castoe, J.

K. Colbourne, M. K. Fujita, R. G. Moreno, B. F. ten Hallers, D. Haussler, A. Heger, D.

Heiman, D. E. Janes, J. Johnson, P. J. de Jong, M. Y. Koriabine, M. Lara, P. A. Novick, C.

L. Organ, S. E. Peach, S. Poe, D. D. Pollock, K. de Queiroz, T. Sanger, S. Searle, J. D.

Smith, Z. Smith, R. Swofford, J. Turner-Maier, J. Wade, S. Young, A. Zadissa, S. V

Edwards, T. C. Glenn, C. J. Schneider, J. B. Losos, E. S. Lander, M. Breen, C. P. Ponting,

and K. Lindblad-Toh. 2011. The genome of the green anole lizard and a comparative

analysis with birds and mammals. Nature 477:587–91.

Bachtrog, D., J. E. Mank, C. L. Peichel, M. Kirkpatrick, S. P. Otto, T.-L. Ashman, M. W. Hahn,

J. Kitano, I. Mayrose, R. Ming, N. Perrin, L. Ross, M. N. Valenzuela, and J. C. Vamosi.

2014. Sex Determination: Why So Many Ways of Doing It? PLoS Biol. 12:1–13.

Bull, J. J. 1983. Evolution of sex determining mechanisms. Benjamin/Cummings Publishing

Company, Menlo Park, California.

Burt, A., and R. Trivers. 2006. Genes in Conflict. Harvard University Press, Cambridge,

Massachusetts.

Charlsworth, B. 1996. The evolution of chromosomal sex determination and dosage

compensation. Curr. Biol. 6:149–162.

De Smet, W. 1981. Description of the orcein stained karyotypes of 27 lizard species (Lacertilia,

Reptilia) belonging to the families Iguanidae, Agamidae, Chamaeleontidae and Gekkonidae

(Ascala- bota). Acta Zool. Pathol. Antverp. 76:35–72.

Ewert, M. A., and C. E. Nelson. 1991. Sex determination in turtles: patterns and some possible

186 adaptive values. Copeia 1:50–69.

187 Gorman, G. 1973. The chromosomes of the Reptilia, a cytotaxonomic interpretation. Pp. 349–

188 424 *in* A. Chiarelle and E. Cpanna, eds. Cytotaxonomic and Vertebrate Evolution.

189 Academic Press, Inc., USA.

190 Guler, Y., S. Short, P. Kile, and A. Fort. 2012. Integrating field and laboratory evidence for

191 environmental sex determination in the amphipod, Echinogammarus marinus. Mar. Biol.

192 159:2885–2890.

193 Haldane, J. B. S. 1922. Sex ratio and unisexual sterility in hybrid animals. J. Genet. 12:101–109.

194 Holleley, C. E., D. O'Meally, S. D. Sarre, J. A. Marshall-Graves, T. Ezaz, K. Matsubara, B.

195 Azad, X. Zhang, and A. Georges. 2015. Sex reversal triggers the rapid transition from

196 genetic to temperature-dependent sex. Nature 523:79–82.

197 Janzen, F. J., and G. Paukstis. 1991. Environmental sex determination in reptiles: ecology,

198 evolution, and experimental design. Q. Rev. Biol. 66:149–179.

199 Kettlewell, J. R., C. S. Raymond, and D. Zarkower. 2000. Temperature-dependent expression of

200 turtle Dmrt1 prior to sexual differentiation. Genesis 26:174–178.

201 King, M. 1977. Evolution and Reproduction. Pp. 55–60 *in* J. Calaby and H. Tyndale-Biscoe, eds.

202 Evolution and Reproduction. Australian Academy of Science, Canberra.

203 Kratochvíl, L., L. Kubicka, and E. Landova. 2008. Does the mechanism of sex determination

204 constrain the potential for sex manipulation? A test in geckos with contrasting sex-

205 determining systems. Naturwissenschaften 95:209–215.

206 Lisachov, A. P., V. A. Trifonov, M. Giovannotti, M. A. Ferguson-Smith, and P. M. Borodin.

207 2017. Heteromorphism of "Homomorphic" Sex Chromosomes in Two Anole Species

208 (Squamata, Dactyloidae) Revealed by Synaptonemal Complex Analysis. Cytogenet.

Genome Res. 151:89–95.

Merchant-Larios, H., and V. Díaz-Hernández. 2013. Environmental sex determination

mechanisms in reptiles. Sex Dev. 7:95–103.

Mitchell, N. J., and F. J. Janzen. 2010. Temperature-Dependent Sex Determination and

Contemporary Climate Change. Sex. Dev. 4:129–140.

Pen, I., T. Uller, B. Feldmeyer, A. Harts, G. M. While, and E. Wapstra. 2010. Climate-driven

population divergence in sex-determining systems. Nature 468:436–438. Nature Research.

Pokorna, M., and L. Kratochvíl. 2009. Phylogeny of sex-determining mechanisms in squamate

reptiles: are sex chromosomes an evolutionary trap? Zool. J. Linn. Soc. 156:168–183.

Quinn, A. E., A. Georges, S. D. Sarre, F. Guarino, T. Ezaz, and J. A. M. Graves. 2007.

Temperature Sex Reversal Implies Sex Gene Dosage in a Reptile. Science. 316:411.

Radder, R. S., A. E. Quinn, A. Georges, S. D. Sarre, and R. Shine. 2008. Genetic evidence for

co-occurrence of chromosomal and thermal sex-determining systems in a lizard. Biol. Lett.

4:176–8.

Reeve, H. K., and D. W. Pfennig. 2003. Genetic biases for showy males: Are some genetic

systems especially conducive to sexual selection? PNAS 100:1089–1094.

Rovatsos, M., M. Altmanová, M. Pokorná, and L. Kratochvíl. 2014a. Conserved sex

chromosomes across adaptively radiated anolis lizards. Evolution (N. Y). 68:2079–2085.

Rovatsos, M., M. Pokorná, M. Altmanová, L. Kratochvíl, and M. Pokorna. 2014b. Cretaceous

park of sex determination : sex chromosomes are conserved across iguanas. Biol. Lett.

10:1–4.

Santoyo-Brito, E., M. Anderson, and S. F. Fox. 2017. Incubation Temperature Modifies Sex

Ratio of Hatchlings in Collared Lizards, Crotaphytus collaris. J. Herpetol. 51:197–201.

Sarre, S. D., A. Georges, and A. E. Quinn. 2004. The ends of a continuum: Genetic and

temperature-dependent sex determination in reptiles. BioEssays 26:639–645.

Shine, R., M. J. Elphick, and S. Donnellan. 2002. Co-occurrence of multiple, supposedly

incompatible modes of sex determination in a lizard population. Ecol. Lett. 5:486–489.

Smith, C., P. McClive, P. Western, K. Reed, and A. Sinclair. 1999. Evolution- Conservation of a

sex-determining gene. Nature 402:601–602.

Spotila, L. D., J. R. Spotila, and S. E. Hall. 1998. Sequence and Expression Analysisof Wt1 and

Sox9 in the Red-Eared Slider Turtle, Trachemys Scripta. J. Exp. Zool. 281:417–427.

Valenzuela, M. N., D. C. Adams, and F. J. Janzen. 2003. Pattern Does Not Equal Process:

Exactly When Is Sex Environmentally Determined? Am. Nat. 161:676–683.

Valenzuela, M. N., and V. Lance. 2004. Temperature Dependent Sex Determination in

Vertebrates. Smithsonian Books, Washington, DC.

Viets, B. E., M. A. Ewert, L. G. Talent, and C. E. Nelson. 1994. Sex-Determining Mechanisms

in Squamate Reptiles. J. Exp. Zool. 56:45–56.

Walker, G. 2005. Sex determination in the larvae of the parasitic barnacle Heterosaccus lunatus:

an experimental approach. J. Exp. Mar. Bio. Ecol. 318:31–38.

Western, P., and A. Sinclair. 2001. Sex, genes, and heat: Triggers of diversity. J. Exp. Zool.

290:624–631.

255   **Table 1. Genes and primer sequences used to determine relative gene dosage through**

256   **qPCR in *Crotaphytus collaris***

Gene short name	Gene name	Forward Primer	Reverse Primer	Amplicon size
EF1a	Elongation factor 1	CCTTATTGTTGCT GCTGGTGTT	GTGCTAACTTCTT TGACGATTTCC	189
TME	Transmembrane protein 132D	TATCCGAGCAGA CCCAAAGTCC	AAGGAGACCCAA CTCAGCCAC	183
ATP	Sarcoplasmic/ endoplasmic reticulum calcium ATPase 2	CAAAGCAGCGG GCATTTAGG	ATCACTGGGGAC AACAAGGG	160
PEPB	Phosphatidylethoan olamine-binding protein 1	GACAGGGCTCCA TCGCTAC	CATAGTCATCCCA CTCCGCC	188

Figure 1. Relative gene dosage at genes PEBP, ATP, and TME for the two males and two females whose DNA amplified consistently and reliably across three replicates. For each gene, the female gene dosage is exactly 1 and the male gene dosage is near 0.5. Each of these genes maps to the *Anolis carolinensis* X microchromosome, thus, this pattern is consistent with male heterogamety.