A Genetic Program Mediates Cold-warming Response and Promotes Stress-induced Phenoptosis in *C. elegans*

Wei Jiang[1, 2]†, Yuehua Wei[1]†, Yong Long[1, 3]†, Arthur Owen[4], Bingying Wang[1], Xuebing Wu[5], Shuo Luo[5], Yongjun Dang[2], Dengke K. Ma[1]*

[1]Cardiovascular Research Institute and Department of Physiology, University of California San Francisco, San Francisco, CA 94158

[2]Key Laboratory of Molecular Medicine, Ministry of Education and Department of Biochemistry and Molecular Biology, School of Basic Medical Sciences, Shanghai Medical College, Fudan University, Shanghai, 200032, China

[3]State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430072, China

[4]Department of Molecular Cell Biology, University of California Berkeley, Berkeley, CA 94720

[5]Department of Biology, Massachusetts Institute of Technology, 9 Cambridge Center, Cambridge, MA 02142

†These authors contributed equally to this work

*For Correspondence: Dengke.Ma@ucsf.edu

Abstract

How multicellular organisms respond to and are impacted by severe hypothermic

stress is largely unknown. From *C. elegans* screens for mutants abnormally

responding to cold-warming stress, we identify a molecular genetic pathway

comprising ISY-1, a conserved uncharacterized protein, and ZIP-10, a bZIP-type

transcription factor. ISY-1 binds to RNA and gatekeeps the ZIP-10 transcriptional

program by regulating the microRNA *mir-60*. Downstream of ISY-1 and *mir-60*,

***zip-10* levels rapidly and specifically increase upon cold-warming stress.**

Consequently, ZIP-10 up-regulates Cathepsin protease-encoding genes to

promote the organismic death, or phenoptosis, of *C. elegans*. *zip-10* deficiency

confers enhanced resistance to prolonged cold-warming stress, more

prominently in adults than larvae. We conclude that the ZIP-10 genetic program

mediates cold-warming response and may have evolved to promote wild-

population kin selection under resource-limiting and thermal stress conditions.

Introduction

Temperature shifts pervasively affect numerous biological processes in all

organisms. Heat shock stimuli activate expression of many heat-shock inducible genes

through the sigma-32 factor and the evolutionarily conserved transcription factor HSF

(Heat Shock Factor) in bacteria and eukaryotes, respectively (Gomez-Pastor et al.,

2017; Yura et al., 1993). Coordinated expression of heat shock-induced chaperone

proteins facilitates cellular proteostasis and adaptation to temperature upshift (Mahat et

al., 2016; Solís et al., 2016). In contrast to heat shock response, how organisms

respond to cold shock is still largely unknown (Al-Fageeh and Smales, 2006; Choi et al.,

2012; Yenari and Han, 2012; Zhu, 2016). Although extensive RNA expression profiling

studies have identified many protein-coding genes and non-coding RNAs that are

regulated by cold shock via both transcriptional and post-transcriptional mechanisms

(Al-Fageeh and Smales, 2006; Giuliodori et al., 2010; Kandror et al., 2004; Zhou et al.,

2017), master regulators of cold shock response and cold-regulated genes

(counterparts of HSF) have long been elusive and mechanisms of cold shock response

in multicellular organisms remain poorly characterized.

At the organismic level, warm-blooded mammals normally keep body

temperature at about 37°C and initiate multiple homeostatic mechanisms to maintain

body temperature upon exposure to hypothermia (Morrison, 2016; Tansey and

Johnson, 2015). In humans, therapeutic hypothermia (32-34°C) has been widely used

to treat ischemic disorders and proposed to activate multifaceted cellular programs to

protect against ischemic damages (Choi et al., 2012; Polderman, 2009; Yenari and Han,

2012). By contrast, cold-blooded animals including most invertebrates experience

varying body temperature depending on the environment, but can nonetheless elicit

stereotypic behavioral, physiological and transcriptional response to chronic

hypothermia or transient cold shock (Al-Fageeh and Smales, 2006; Garrity et al., 2010).

Like many other types of stress, prolonged severe hypothermia can lead to the death of

organisms, in most cases likely because of failure in adapting to the stress, or

alternatively through stress-induced phenoptosis, namely genetically programed

organismic death (Longo et al., 2005; Skulachev, 1999, 2002). Although phenoptosis

has been phenotypically documented in many cases, its evolutionary significance and

genetic mechanisms remain unclear and debated (Longo et al., 2005; Sapolsky, 2004).

We previously discovered a *C. elegans* genetic pathway that maintains cell

membrane fluidity by regulating lipid desaturation in response to moderate hypothermia

(10-15°C) (Fan and Evans, 2015; Ma et al., 2015). Expression of the gene *fat-7*, which

encodes a lipid desaturase, is transcriptionally induced by 10-15°C but not by more

severe hypothermia (i.e. cold shock at 0-4°C), which impairs *C. elegans* reproduction

and growth, and elicits distinct physiological and behavioral responses (Garrity et al.,

2010; Lyons et al., 1975; Ma et al., 2015; Murray et al., 2007). However, as severe

hypothermia arrests most of cell biological processes, strong transcriptional responses

to cold shock e.g. 0-4°C likely only manifest during the organismic recovery to normal

ambient temperature. We thus hypothesize that a genetic pathway differing from that

operating under moderate hypothermia exposure controls the transcriptional response

to severe hypothermia/cold shock followed by warming in *C. elegans*.

In this work, we performed transcriptome profiling to first identify genes that are

regulated by exposure to cold shock followed by recovery at normal temperature. We

then used GFP-based transcriptional reporters in large-scale forward genetic screens to

identify a genetic pathway consisting of *isy-1* and *zip-10*, the latter of which responds to

cold-warming (CW) and mediates transcriptional responses to CW. Unexpectedly, we

found *zip-10* induction promotes organismal death while deficiency of *zip-10* confers

resistance to prolonged CW stress, more prominently in adults than young larvae. We

propose that CW activates a ZIP-10 dependent genetic program favoring *C. elegans*

phenoptosis and postulate that such programmed organismic death may have evolved

to promote wild-population kin selection under thermal stress conditions.

Results

To identify new mechanisms of *C. elegans* response to severe hypothermia, we

performed RNA sequencing (RNA-seq) of wild-type *C. elegans* populations after 2-hr

exposure to 4°C cold shock followed by recovery at 20°C for 1 hr. We used such CW

conditions in an attempt to identify genes that specifically and rapidly respond to CW

rather than those that respond to general organismic deterioration after long cold

exposure. After differential expression analyses of triplicate samples, we identified 604

genes that are significantly up- or down-regulated by such CW conditions (Table 1,

Figure 1A and Figure 1-figure supplement 1A). Gene ontology analysis indicated that

the CW-regulated genes are involved in biological processes including lipid

metabolisms, autophagy, proteostasis and cell signaling (Figure 1-figure supplement

1B). We generated transgenic *C. elegans* strains in which *GFP* is driven by promoters

of the top-ranked CW-inducible genes. In this work, we focus on *asp-17* as a robust

CW-inducible reporter gene owing to its low baseline expression level and high-fold

induction by CW, features that permitted facile isolation of full-penetrance mutants after

random mutagenesis (see below) with both abnormal *asp-17p::GFP* expression and

altered organismic tolerance to prolonged cold stress.

C. elegans asp-17 encodes an aspartyl-like protease with unknown molecular

functions. Like other CW-inducible genes, *asp-17* up-regulation is more prominently

induced by severe than moderate hypothermia followed by recovery from cold shock

(Figures 1B and 1C). Among the aspartyl-like protease family members, we found that

only *asp-17* was robustly and specifically induced by CW (Figure 1D). The up-regulation

of endogenous *asp-17* by CW can be recapitulated by an integrated GFP reporter

driven by the endogenous *asp-17* promoter, indicating transcriptional regulation of *asp-*

17 by CW (Figure 2-Figure supplement 1A). We varied CW treatment conditions and

found that the induction of *asp-17* strictly required the warming phase after cold shock

(Figure 1E). However, heat shock at 32°C did not increase *asp-17* expression (Figure

1E), consistent with previous large-scale transcriptome profiling studies in *C. elegans*

(Brunquell et al., 2016). Single-molecule fluorescent in-situ hybridization (smFISH)

identified the CW-induced *asp-17* predominantly in intestinal cells (Figure 1F). Since

CW activates numerous other genes in addition to *asp-17*, we sought to use *asp-*

17p::GFP as a robust readout reporter to identify the upstream genetic pathway and

transcriptional regulators that control *asp-17* induction by CW.

We performed a forward genetic screen using EMS-induced random

mutagenesis of a parental strain carrying a genome-integrated *asp-17p::GFP* reporter

and isolated over 30 mutants with constitutive *asp-17p::GFP* expression in the absence

of CW (Figure 2-figure supplement 1B and 1C). We molecularly cloned one mutant

dma50 that exhibited fully penetrant and constitutively strong expression of *asp-*

17p::*GFP* (Figure 2A,B and Figure 2-figure supplement 1D-1F). Compared with wild

type, *dma50* strongly up-regulated *asp-17*::*GFP* in the intestine (Figure 2B). By single

nucleotide polymorphism-based linkage analysis of the intestinal *asp-17p*::GFP

phenotype, we mapped *dma50* to a genetic interval on Chromosome V and used whole-

genome sequencing to identify candidate causal gene mutations (Figure 2-figure

supplement 1D,E). Based on phenocopying by feeding RNAi and transformation rescue

of the *asp-17p*::GFP phenotype, *dma50* defines a previously uncharacterized gene *isy-1*

(Figures 2A-2F and Figure 2-figure supplement 1F). *isy-1* (*I*nteractor of <u>SY</u>F1 in yeast)

encodes a protein with strong sequence similarity to an evolutionarily highly conserved

family of RNA-binding proteins in eukaryotes (Figure 2A and Figure 2-figure supplement

2A-2C) (Dix et al., 1999; Du et al., 2015).

dma50 caused substitution of a negatively charged glutamate, which is

completely conserved in the ISY protein family, to a positively charged lysine in the

predicted coiled-coil region of *C. elegans* ISY-1 (Figure 2A). An *isy-1p::isy-1::GFP*

translational reporter indicated a rather ubiquitous distribution of ISY-1::GFP in many

tissues including intestinal nuclei (Figure 2C). The strong intestinal *asp-17p*::GFP

expression caused by *dma50* was fully rescued by transgenic expression of wild-type

isy-1(+), single-copy integration of a *mCherry*-tagged *isy-1(+)* allele, or *isy-1*(+)

expression driven by the intestine-specific *ges-1* promoter (Figures 2E and 2F). In

addition, the *ges-1*-driven transgenic expression of sense plus antisense *isy-1* RNAi

fully recapitulated the *dma50* phenotype (Figure 2D). Endogenous expression of *asp-17*

was also drastically up-regulated in *isy-1* mutants (Figure 2-Figure supplement 2D).

Thus, these results identify *isy-1* as a causal cell-autonomous regulator of *asp-17*.

Human ISY1 is critical for certain microRNA processing while yeast ISY1 is a

likely component of the spliceosome (Dix et al., 1999; Du et al., 2015; Galej et al.,

2016). We found that CW-induced *asp-17* up-regulation was further enhanced in *isy-1*

mutants compared with wild type (Figure 3A), suggesting that ISY-1 normally restricts

transcriptional activity of *asp-17*. To determine the mechanism by which ISY-1 regulates

transcription of *asp-17*, we sought to identify transcription factors (TF) that meet two

criteria: a), its mRNA or protein products are altered in *isy-1* mutants, and b), it is

genetically epistatic to *isy-1*, i.e. its loss-of-function (LOF) can suppress *isy-1* LOF (thus

also likely required for *asp-17* induction by CW). We performed RNA-seq from triplicate

samples of wild-type and *isy-1* mutants, from which we analyzed differentially expressed

TF genes in *isy-1* mutants and found that a bZIP-type transcription factor-encoding

gene *zip-10* met both criteria (Figure 3B, Table 2). *zip-10* mRNA was drastically up-

regulated in *isy-1* mutants, whereas levels of closely related bZIP-family genes, such as

zip-11, were unaffected (Figure 3C). Importantly, genetic deletion of *zip-10* completely

abrogated the ability of *isy-1* RNAi to activate *asp-17p::GFP* (Figure 3D). These results

indicate that ISY-1 regulates *asp-17* by controlling the level of *zip-10* mRNAs.

Next, we examined how the ISY-1/ZIP-10/ASP-17 pathway is regulated by CW.

CW did not apparently alter levels of endogenous *isy-1* mRNAs or mCherry-tagged ISY-

1 proteins under the endogenous *isy-1* promoter (Figure 2-Figure supplement 2D and

2E). By contrast, we found that CW induced drastic up-regulation of ZIP-10 proteins

from a tagged *zip-10p::zip-10::EGFP::FLAG* allele in an integrated transgenic strain

(Figure 3E). Although EGFP fluorescence was invisible in animals carrying such

transgenes (likely because it is sandwiched by *zip-10* and FLAG), the striking induction

of ZIP-10::EGFP::FLAG was completely blocked by RNAi against *zip-10* or *GFP*,

confirming the transgene specificity (Figure 3F). The baseline level of ZIP-

10::EGFP::FLAG was close to the detection limit of Western blot under normal

conditions, but nonetheless is strongly up-regulated upon RNAi against *isy-1* (Figure

3F). Similar to that of *asp-17*, the induction of *zip-10p::zip-10::EGFP::FLAG* strictly

required the warming phase of CW and occurred rapidly but transiently after warming

during CW (Figure 3G). CW strongly up-regulated *asp-17* expression in both wild type

and *isy-1* mutants, which exhibited abnormally high *zip-10* mRNA levels (Figure 3H).

Furthermore, *zip-10* deletion completely abrogated the up-regulation of *asp-17* levels by

CW (Figure 3I). We also examined the ZIP-10 dependency of other CW-inducible genes

identified by RNA-seq and found that at least *cpr-3* also required ZIP-10, but other CW-

inducible genes including *srr-6* and *F53A9.1*, did not (Figure 3I). These results

demonstrate that ISY-1 suppresses *asp-17* by decreasing *zip-10* levels whereas CW

up-regulates ZIP-10 protein abundance to promote *asp-17* expression.

How is *zip-10* regulated by ISY-1 and CW? Loss of ISY-1 function affected

neither general intron splicing based on an intronic GFP reporter assay, nor specific

splicing of *zip-10*, although both CW and *isy-1* mutations strongly up-regulated *zip-10*

mRNA levels (Figure 3-Figure supplement 1A-1E). We constructed a GFP

transcriptional reporter driven by the endogenous *zip-10* promoter and found it was

markedly up-regulated by *isy-1* RNAi (Figure 3-Figure supplement 2A). While non-

thermal stresses such as hypoxia and starvation did not increase ZIP-10 levels, CW

drastically increased abundance of ZIP-10 in both cytosol and nucleus without affecting

abundance of other house-keeping proteins, including HSP90, tubulin and histone H3

(Figure 3-Figure supplement 2B and 2C). CW up-regulation of ZIP-10 required warming

and was enhanced by more prolonged cold shock (Figure 3-Figure supplement 2D).

Since CW can markedly increase *zip-10* mRNA levels but to a lesser extent than the

isy-1 mutation (Figure 3-Figure supplement 1D and 1E), we tested whether ZIP-10

proteins might be regulated by CW through translational control and mRNA stability.

RNAi against genes encoding eIF5 and a component of the Ccr4-Not complex did not

apparently alter ZIP-10 levels (Figure 3-Figure supplement 2E). Together, these results

indicate that CW and ISY-1 regulate *zip-10* primarily at the transcriptional level.

Human ISY1 facilitates the processing of primary transcripts encoding certain

families of microRNAs (Du et al., 2015). Both *zip-10* and *asp-17* are up-regulated in a

mutant *C. elegans* strain deficient in the microRNA *mir-60* (Kato et al., 2016). We thus

tested whether *mir-60* mediates the regulation of *zip-10* by ISY-1. Immunoprecipitation

of mCherry-tagged ISY-1 followed by QPCR revealed specific binding of primary

transcripts encoding *mir-60* as well as a protein-coding gene *cebp-1* (Figure 4A-4C).

Although neither *isy-1* nor *mir-60* levels were affected by CW, we found CW slightly but

significantly increased *mir-60* binding to ISY-1, perhaps as a feedback mechanism to

limit over-activation of *zip-10*-dependent genes after CW treatment (Figure 4B).

Importantly, mature *mir-60* levels were drastically decreased in *isy-1* mutants while loss

of *mir-60* led to up-regulation of *zip-10* and *zip-10*-dependent subset of CW-inducible

genes, including *asp-17* and *cpr-3,* but not many other CW-inducible genes (Figure 4D,

4E and Figure 4-figure supplement 1). The 3' untranslated region (Utr) of *zip-10*

appeared not to be regulated by CW or *isy-1* RNAi (Figure 4F). However, *isy-1* RNAi

caused an abnormally high baseline level of ZIP-10 in the absence of CW and enabled

further heightened ZIP-10 up-regulation in response to CW, followed by its down-

regulation over an extended period of warming (Figure 4G). These results indicate that

CW regulates transcription of *zip-10* (and thereby that of *asp-17*), while ISY-1 controls

expression of *zip-10* via *mir-60*, likely through microRNA processing and regulation of

additional upstream transcriptional *zip-10* regulators that respond to CW.

We compared the genes differentially regulated by CW and those by *isy-*

1(dma50) mutants and found 246 genes, including the two ZIP-10 dependent targets

asp-17 and *cpr-3*, that are commonly regulated by both conditions (Figure 5A, Table 3).

Global transcriptome changes between these two conditions are also significantly

correlated (Figure 5B) (correlation coefficient *R* as 0.54, significance *P* value as 0). We

used the bioinformatics tool MEME (Bailey et al., 2009) to identify motifs present in the

promoters (~600 bp upstream of transcription start sites) of the commonly regulated

gene subset and identified a single enriched motif characterized by AT-rich sequences

(Figure 5C). The gene most enriched with this motif is *asp-17*, the promoter of which

contains 16 such motifs (Figure 5C). ZIP-10 is a bZIP-type transcription factor predicted

to contain N-terminal low sequence-complexity domains and a C-terminal DNA-binding

and glutamine-rich transactivation domain (Figure 3-Figure supplement 2F-2H). To test

whether the *asp-17* promoter with the identified AT-rich motifs can be bound directly by

ZIP-10, we performed chromatin immunoprecipitation (ChIP) experiments and detected

asp-17 promoter sequences in the FLAG-tagged ZIP-10 chromatin complex only under

CW conditions (Figure 5D). These results indicate that ZIP-10 directly binds to and

activates the *asp-17* promoter in the genetic program regulated by ISY-1 and CW.

The striking regulation of *zip-10* and *asp-17* by CW and ISY-1 prompted us to

examine the organismic phenotype of various mutants upon prolonged CW stress. A

majority of wild-type *C. elegans* adults died upon prolonged CW stress (e.g. 2-4^{o}C for

over 24 hrs) (Ohta et al., 2014). We found that *asp-17* or *zip-10* loss-of-function mutants

exhibited markedly higher survival rates than wild type under the same prolonged CW

stress condition (Figure 5E). Consistent with a role of wild-type *zip-10* in promoting

organismic death, inducible *zip-10* over-expression by mild transient heat shock,

mediated by the *hsp-16* promoter, promoted animal death even in the absence of CW

(Figure 5F). By contrast, other ectopically induced *zip* genes including *zip-11* and *zip-2*

did not affect animal death, while a mutation specifically disrupting the glutamine-rich

transactivation domain of ZIP-10 abolished the death-promoting effect (Figures 5F and

Figure 3-Figure supplement 2H). Although *zip-10* is genetically epistatic to *isy-1* in the

regulation of *asp-17*, we found that *isy-1* mutants are also markedly resistant to

prolonged cold stress. This paradox was resolved after we observed that many

downstream target genes of the stress-coping transcription factors HIF-1, HSF-1 and

DAF-16 are up-regulated in *isy-1* mutants, and loss-of-function of at least *daf-16* could

partly suppress cold tolerance by *isy-1* RNAi (Figures 5G and 5H). Since ISY-1

regulates *zip-10* via *mir-60* (Figure 4) supporting a role of ISY-1 in specific microRNA

processing (Du et al., 2015), we performed small RNA library sequencing of wild-type

and *isy-1* mutants and identified specific members of microRNAs that were differentially

regulated, including *mir-60* and additional microRNAs predicted to target stress-coping

TFs (Figure 5-Figure supplement 1A-1F). Thus, *isy-1* mutants likely exhibit pleiotropic

phenotypes caused by abnormal activation of multiple TFs in addition to ZIP-10. In

contrast to ZIP-10 dependent genes (*asp-17* and *cpr-3*), the HIF-1/HSF-1/DAF-16 target

genes were not apparently induced by CW (Table 1). Furthermore, unlike HSF-1 or

DAF-16 induction by other types of stress stimuli, ZIP-10 induction by CW appears

present specifically in adults but not in larvae (Figure 5-figure supplement 2C).

Consistently, the phenotypic difference in cold tolerance between wild type and *zip-10*

mutants manifested more prominently in developmentally more mature-stage and older

animals (Figure 5-figure supplement 2D). These results indicate that CW specifically

activates a ZIP-10-driven and developmental stage-modulated transcriptional genetic

program to promote the organismic death, or phenoptosis, of *C. elegans* (Figure 5I).

Discussion

From a genetic screen for *C. elegans* mutants with altered transcriptional

response to CW, we identified *isy-1* and subsequently discovered the CW and ISY-1-

regulated transcription factor ZIP-10 as a key mediator of the transcriptional response to

CW. A thermal stress-responding TF might be expected to promote adaptation of

animals towards the stressor, causing its LOF mutants to be sensitive to the stress.

Unexpectedly, we found *zip-10* mutants are markedly resistant to prolonged cold stress.

However, unlike other stress-responding TFs that activate genes largely beneficial for

physiological homeostasis and thus animal health under stress conditions (Baird et al.,

2014; Dempersmier et al., 2015; Hwang and Lee, 2011; Kandror et al., 2004; Kumsta et

al., 2017; Landis and Murphy, 2010), identified transcriptional targets of ZIP-10 include

at least two Cathepsin-type proteases, CPR-3 and ASP-17 (Figure 4E). In contrast to

aspartyl-type proteases which are largely unknown in cellular functions, caspase-type

proteases are well-known apoptotic cell death executioners while CPR-4, a Cathepsin

CPR-3 paralogue, has been shown to inhibit cell deaths in *C. elegans* (Metzstein et al.,

1998; Peng et al., 2017; Peter, 2011). Ectopic expression of *zip-10* and its targets

promotes organismic deaths, in contrast to the effect of *zip-10* or *asp-17* deficiency on

cold tolerance (Figure 5E and 5F). As duration of cold shock affects levels of ZIP-10

and transient CW does not trigger phenoptosis, the pro-death role of the *zip-10* genetic

program likely depends on multiple factors, including the duration and severity of cold

exposure. Notably, apoptotic cell death-promoting effects have also been described for

specific members of mammalian bZIP TFs (Chüeh et al., 2017; Hartman et al., 2004;

Ritchie et al., 2009). The specific and robust induction of ZIP-10 by CW, the opposing

cold-tolerance phenotypes caused by *zip-10* loss-of-function and gain-of-function

genetic manipulations, as well as the pro-death roles of ZIP-10 targets support the

notion that the *zip-10* pathway is activated by CW to promote phenoptosis.

How do ISY-1 and CW regulate the *zip-10* pathway? We found that the *zip-10*

promoter activity responds to the loss of ISY-1, which normally maintains *mir-60* levels

and thereby regulates *zip-10* transcription likely through the processing of small RNAs.

Severe cold stress also leads to accumulation of another class of small RNA risiRNA,

which is important for maintaining rRNA homeostasis (Zhou et al., 2017). Whether ISY-

1 might also affect risiRNA processing remains to be characterized. Constitutive up-

regulation of ZIP-10 targets in *isy-1* mutants and the lack of evidence for regulation of

ISY-1 by CW supports ISY-1 as a gate-keeper for the ZIP-10-driven transcriptional

response to CW (Figure 5I). Regulation of *zip-10* is primarily transcriptional based on

evidence we present in this study; further studies are required to discern to what extent

mir-60 might directly act at the *zip-10* locus or more indirectly impact the transcription of

zip-10, e.g. by post-transcriptionally inhibiting translation of a transcriptional activator.

Up-regulation of the activity of the *zip-10* promoter by CW indicates that additional cold-

responding sensors and effectors upstream of ZIP-10 remain to be identified, by

signaling mechanisms perhaps similar to the well-characterized cold-responding

pathways found in other organisms (Dempersmier et al., 2015; Kandror et al., 2004;

Zhu, 2016). Precisely how *zip-10* is regulated by CW in coordination with ISY-1 to

promote *C. elegans* death under prolonged CW stress awaits further investigation.

The roles of ZIP-10 and a dedicated genetic program in promoting organismic

death are surprising but would make sense in light of the evolutionary kin selection

theory. Kin selection refers to the evolutionary process in favor of the reproductive

success of an organism's kin despite a cost to the organism's own reproduction

(Hamilton, 1963; Smith, 1964). Dedicated genetic programs may have evolved to

promote kin selection at the population level. Although the concept and potential

mechanisms of programed organismic death, or phenoptosis, are debated, examples of

kin selection and stress-induced organismic deterioration have been widely documented

in many organisms (Longo et al., 2005; Sapolsky, 2004; Skulachev, 1999, 2002).

Laboratory conditions for hermaphroditic *C. elegans* clearly no longer exert

selection pressure for genetic programs underlying phenoptosis or kin selection.

However, our mathematic modeling of an exemplar situation of population growth for

wild-type and *zip-10* deficient animals under food-limiting and CW stress conditions

supports the phenoptosis or kin selection hypothesis for the *zip-10* pathway (Figure 5-

Figure supplement 2A and 2B). Experimentally, we found that both the CW-induced *zip-*

10 expression and the death-promoting effect of ZIP-10 occurred more prominently in

older adults than in larvae (Figure 5-Figure supplement 2C and 2D). Extending from the

kin selection theory, we postulate that the evolutionary advantage of programmed

organismic death might manifest in the wild, where resources for growth and

reproduction are limited and environments can change drastically. As such, the

selective death of adult animals would benefit young and reproductively more privileged

populations to facilitate the spreading of genes by young populations under resource-

limiting and high-stress conditions. Our work provides an unprecedented example of

stress-induced phenoptosis in *C. elegans* and identify a specific transcription factor in a

genetic program that likely evolved to promote kin selection during animal evolution.

These findings therefore bear broad implications for understanding thermal stress

response, programmed organismic death (phenoptosis) and evolutionary biology.

Experimental Procedure

<u>*C. elegans* strains and genetic manipulations</u>

C. elegans strains were maintained with standard procedures unless otherwise

specified. The N2 Bristol strain was used as the reference wild type, and the

polymorphic Hawaiian strain CB4856 was used for genetic linkage mapping and SNP

analysis (Brenner, 1974; Davis et al., 2005). Forward genetic screen for constitutive

asp-17p::GFP reporter-activating mutants after ethyl methanesulfonate (EMS)-induced

random mutagenesis was performed as described previously (Ma et al., 2012, 2015).

Single-copy integration of *isy-1p::isy-1::mCherry* transgene was generated using the

MosSCI method (Frøkjaer-Jensen et al., 2008). To generate *asp-17* null alleles in *C.*

elegans, we used CRISPR-Cas9 to induce double stranded breaks and subsequent

non-homologous end joining caused a deletion of *asp-17*. Feeding RNAi was performed

as previously described (Kamath and Ahringer, 2003). Transgenic strains were

generated by germline transformation as described (Mello et al., 1991). Transgenic

constructs were co-injected (at 10 - 50 ng/µl) with dominant *unc-54p*::mCherry or *rol-6*

markers, and stable extrachromosomal lines of mCherry+ or roller animals were

established. Genotypes of strains used are as follows: *daf-16(mu86) I, mir-60(n4947) II*;

isy-1(dma50) V, zip-10(ok3462) V, asp-17(dma99) V, dmaIs10[asp-17p::GFP; unc-

54p::mCherry] X, dmaIs21[zip-10p::GFP; unc-54p::mCherry]; wgIs634[zip-10p::zip-

10::EGFP::FLAG + unc-119(+)], oxTi302 [eft-3p::mCherry::tbb-2 3'UTR + Cbr-unc-

119(+)], dmaSi1[isy-1p::isy-1::mCherry, unc-119(+)], dmaEx95[ges-1p::isy-1(+); rol-

6(+)], dmaEx99[isy-1 genomic DNA (2ng/ul); rol-6(+)], nEx102[ges-1p::isy-1(+); rol-

6(+)], nEx103[ges-1p::isy-1(+); rol-6(+)], dmaEx104[ges-1p::mCherry::3utr(zip-10), rol-

6(+)], dmaEx123[hsp-16p::zip-10; rol-6(+)], dmaEx124[hsp-16p::zip-10; rol-6(+)],

dmaEx131[zip-10p::GFP; unc-54p::mCherry].

<u>Sample and library preparation for RNA sequencing</u>

Control N2 animals and the *isy-1* mutants were maintained at 20°C. For cold stress, N2

animals were exposed to 4°C for 2 hrs followed by 1 hr recovery at 20°C. Upon sample

collection, the animals were washed down from NGM plates using M9 solution and

subjected to RNA extraction using the RNeasy Mini Kit from Qiagen. 1 µg total RNA from

each sample was used for sequencing library construction. Each treatment included 3

biological replicates. The NEBNext® rRNA Depletion Kit was used for rRNA depletion.

After rRNA depletion, the Agencourt RNAClean XP Beads from Beckman Coulter were

used for RNA purification. Then, the NEBNext® Ultra™ Directional RNA Library Prep Kit

for Illumina® was used for RNA fragmentation, first strand and second strand cDNA

synthesis and double-stranded cDNA end repair. Double strand cDNAs were purified

using the Agencourt AMPure XP from Beckman Coulter and ligated to adaptors of the

NEBNext Multiplex Oligos for Illumina. Finally, the Q5 Hot Start HiFi PCR Master Mix was

used for PCR enrichment of the adaptor-ligated DNA. The concentration and quality of

the constructed sequencing libraries were measured by using the Agilent High Sensitivity

DNA Kit and a Bioanalyzer 2100 from Agilent Technologies. The libraries were submitted

to 100 bp paired-end high throughput sequencing using Hiseq-3000 by the Center for

Advanced Technology (CAT) of the University of California, San Francisco.

RNA-seq data analysis was performed using a super computer system equipped

with multiple processors. The raw reads were trimmed and filtered by the prinseq-lite

software (0.20.4) (Schmieder and Edwards, 2011). Reads longer than 30 bp and with a

minimum quality score higher than 15 were kept and used for subsequent analyses. The

filtered left and right read sets were compared by the Pairfq script to separate paired and

single reads. The clean reads were mapped to the *C. elegans* genome sequence using

Hisat2 (2.0.5)(Kim et al., 2015) with default parameters. The number of mapped reads

were counted by featureCounts from the Subread package (1.5.0) (Liao et al., 2014).

Differential gene expression analysis was performed using the DESeq2 package (Love

et al., 2014). Adjusted P-value ≤ 0.05 was used as the threshold to identify the

differentially expressed genes. Gene ontology and KEGG pathway enrichment analyses

for the differentially expressed genes were conducted using the Cytoscape plugins

BiNGO (Maere et al., 2005) and ClueGO (Bindea et al., 2009), respectively. Plots for the

mapped reads were generated by IGVtools (Thorvaldsdóttir et al., 2013).

<u>Quantitative RT-PCR</u>

50 µl pellet animals were resuspended in 250 µl lysis buffer of Quick-RNA

MiniPrep kit (Zymo Research, R1055) then lysed by TissueRuptor (Motor unit "8" for 1

min). Total RNA was extracted following the instruction (Zymo Research, R1055). 2 µg

RNA/sample was reverse transcribed into cDNA (BioTools, B24408). Real-time PCR

was performed by using Roche LightCycler®96 (Roche, 05815916001) system and

SYBR Green (Thermo Fisher Scientific, FERK1081) as a dsDNA-specific binding dye.

qRT-PCR condition was set to 95°C for denaturation, followed by 45 cycles of 10s at

95°C, 10s at 60°C, and 20s at 72°C. Melting curve analysis was performed after the

final cycle to examine the specificity of primers in each reaction. Relative mRNA was

calculated by ΔΔCT method and normalized to actin. Primers for qRT-PCR: *asp-17*

(Forward, ATGTTCCGCTGACTGCGAAG; Reverse, TTTCATTCATTTCATCCCAC),

F53A9.1, Forward, ACTACGGAAACGGAGGATAC; Reverse,

TGGCCGTGATGATGATGATG), *srr-6*   (Forward, CTCCAAGTCCTGAAGTCGTG;

Reverse, GTAGGGATGGATTGAACTCG), *isy-1* (Forward,

AGATGCTGAGCGATTCAGAC; Reverse, CTTTCGATAGTCCGTACCAC), *zip-10*

(Forward, TCGAGATGCTCTTCAACTG; Reverse, CTAACTGCTTGCCGGAG), *cpr-3*

(Forward, GTAGTGGAGCAGTAACAGGTG; Reverse,

CAGTTTGAATTTCGGTGACGG), *act-3* (Forward, TCCATCATGAAGTGCGACAT;

Reverse, TAGATCCTCCGATCCAGACG).

<u>Sample preparation and Western blot of proteins</u>

Transgenic (*isy-1p::isy-1::mCherry* and *zip-10p::zip-10::EGFP::FLAG*) animals

were cold shocked (4°C) for 0, 1, 2 or 4 hrs, followed by recovery at 25°C for 1 h. Animals

were harvested and washed 3 times with M9 and 20 µl pellet animals were lysed directly

in Laemmli Sample Buffer and used for Western blot analysis. Proteins were resolved by

15% SDS-PAGE (Bio-Rad, 4561084) and transferred to a nitrocellulose membrane (Bio-

Rad, 1620167). Proteins were detected using antibodies against Flag (Sigma, F3165),

mCherry (M11217, Life Technologies), Tubulin (Sigma, T5168), H3 (Abcam, ab1791) or

HSP90 (Proteintech, 13171-1-AP).

For subcellular fractionation, 50 µl pellet animals were resuspended in 150 µl 1 X

cell lysis buffer (Cell Signaling Technology, 9803S) with protease inhibitor cocktail

(BioTools, B14002) and 10 µM PMSF, and incubated for 10 min on ice. Animals were

lysed by TissueRuptor (Qiagen, 9001271) with Motor unit "6" for 30 sec on ice. After

incubation on ice for 5 minutes and centrifugation at 5,000 rpm at 4°C for 2 min, the

supernatant was collected as the cytoplasmic part. The nuclear pellet was washed three

times with lysis buffer and resuspended in 150 µl RIPA buffer (Thermo Fisher Scientific,

P89900) for 30 min on ice, spun at 12,000 rpm for 15 min, and the supernatant was

collected as nuclear extract. Tubulin and H3 were separately used as cytoplasm and

nuclear loading control. For RNAi experiments, *zip-10p::zip-10::EGFP::FLAG* animals

were bleached and the eggs were laid onto RNAi plates. Animals were harvested as

L4/young adults and subject to Western blot analysis as described above.

<u>Chromatin and RNA immunoprecipitation (ChIP-qPCR and RIP-qPCR)</u>

ChIP-qPCR assay was carried out as before with modifications. Briefly, CW-

treated animals (4°C for 4 hrs, recovered at 25°C for 1 hr) and control (25°C) animals

were harvested and washed by 1 X PBS. The pellet animals were resuspended in cross-

linking buffer (1% formaldehyde in 1 X PBS) followed by homogenization using

TissueRuptor with Motor unit "4" for 1 min at room temperature. The process was then

stopped by addition of glycine (125 mM final concentration). After washing and discarding

the supernatant, the pellet was resuspended in lysis buffer and lysed by TissueRuptor

with Motor unit "6" for 1 min on ice, with lysate kept on ice for additional 3 min, and then

repeated 3 times. The lysate was centrifuged to collect the supernatant and one percent

of the aliquot was used as "Input". Lysate was precleared by adding salmon sperm

DNA/protein-A agarose beads (Bioworld, 20182011-1), rotating at 4°C for 1 hour. After

centrifugation, supernatant was divided equally and added with 50 µg Flag antibody

(Sigma, F3165) and mouse IgG (Santa Cruz Biotechnology, sc-2025) respectively. The

samples were incubated and rotated overnight at 4°C. Next, salmon sperm DNA/protein-

A agarose beads were added again for 2 hrs at 4°C. The beads-antibody-TF-DNA

complex was washed extensively and the complex and input were diluted with proteinase

K buffer. The samples were then incubated at 55°C for 4 hrs and then at 65°C overnight

to reverse crosslink. DNA was extracted by phenol-chloroform-isoamylalcohol (Sigma-

Aldrich, 77617). *asp-17* promoter was measured by qRT-PCR and calculated by percent

input method. Primers for ChIP-qPCR: *asp-17* promoter (Forward,

TTCGCTGCACCTATATGTTG; Reverse, CCGCTAATACCCTTATCAC).

RNA immunoprecipitation (RIP)-qPCR assay was carried out as before with

modifications to accommodate our reagents (Kershner and Kimble, 2010). Briefly,

synchronous day-1 *isy-1p::isy-1::mcherry* animals were divided into two groups. One

group is control (25°C) and the other is cold-warming (4°C for 4 hours, recovered at 25°C

for 1 hour). Animals were harvested and washed by M9 buffer until the supernatant was

clear, and then washed once in buffer A and twice in lysis buffer. About 250 µl worm

pellets were frozen in liquid nitrogen twice and homogenized using TissueRuptor with

Motor unit "4" for 1 min on ice. The lysate was kept on ice for 15 min and centrifuged to

collect the supernatant and 1% of the aliquot was kept as "Input". Equal amount of

supernatant was added with RFP-Trap®_MA (Chromotek) and rotated for 4 hrs at 4°C.

IP magnetic agarose beads were washed and 10% of IP beads were boiled for 6 min in

2X Laemmli Sample Buffer. RNA was eluted from remaining beads using 200 µl lysis

buffer of Quick-RNA MiniPrepkit (Zymo Research, R1055) and extracted following the

instruction. RNA was quantified with a Nanodrop device. 500 ng RNA was reverse

transcribed into cDNA and qPCRwas performed as before. Relative mRNA was

calculated by percent input method. Primers for RIP-qPCR: Primary *mir-60* Forward

TCGAAAACCGCTTGTTCTTG, Reverse CGATTTCTCAAGTCTTGAACTAG; *cebp-1*

Forward GATCCTTCGCAAGACAAGAC, Reverse CACATTGTCGGTAGGAACGTC.

Cold tolerance assay

Animals were cultured under non-starved conditions for at least 4 generations at

25°C before cold tolerance assay. For cold tolerance assay of L1-stage animals, bleach-

synchronized populations were kept at 4°C for 96 hrs and then recovered for 4 hrs at

25°C. For cold tolerance assay of adults, animals were raised at 25°C from hatching with

excessive bacteria food on agar plates. Well-fed L4 stage animals were transferred to

new plates and kept at 25°C overnight to reach day-1 adulthood. To cold shock the

animals, agar plates were spread with equal distance on a thin plastic board and

transferred to a constant 4°C-cold room for the indicated duration. After cold shock,

animals were then moved to 25°C for recovery for 4 hrs before counting the survival rate.

Animals were scored as dead if they showed no movement upon light touch.

Imaging and fluorescence quantification

smFISH of *C. elegans* and imaging were performed as previously described (Ji

and Oudenaarden, 2005). For Nomarski and fluorescence imaging, spinning-disc

confocal and digital automated epifluorescence microscopes (EVOS, Life Technologies)

were used to capture images of animals after RNAi or CW treatments. Synchronous

population of worms were randomly picked and treated with 1 mM levamisole water

solution to paralyze the animals. The animals were mounted on an agar pad on a slide

and aligned for imaging. Identical conditions and settings were used for both control and

test groups. For quantification of fluorescence images, the animals in the images were

outlined and signals were quantified by ImageJ software. The intensity of an individual

animal was obtained by dividing the total signal by the area of that animal. The average

intensity of the control group was set to be 1.0, to which all other intensities were

normalized. Graphpad Prism software was used to plot the data.

<u>Small RNA-seq and bioinformatics</u>

For small RNA sequencing, total RNA was isolated by the Quick-RNA MiniPrep kit

(Zymo Research, R1055) that yields total RNA including small RNAs ranging 17-200 nt.

RNA samples extracted from triplicate N2 animals and *isy-1* mutants were submitted to

Beijing Genomics Institute for small RNA library construction and sequencing. The low-

quality reads were filtered and clean reads were mapped to the *C. elegans* genome using

Bowtie2 program (Langmead and Salzberg, 2012). MiRDeep2 (Friedländer et al., 2012)

was used to characterize known and predict novel miRNAs. The small RNA expression

level was calculated as TPM (transcript per million). Differentially expressed small RNAs

were detected by DESeq2 (Love et al., 2014). The threshold for differentially expressed

sRNAs was adjusted P-value ≤ 0.05 and the absolute value of Log2ratio ≥ 1. Targets of

miRNAs were predicted by TargetScan (Jan et al., 2011), RNAhybrid (Krüger and

Rehmsmeier, 2006) and miRanda (John et al., 2004) using default parameters.

<u>Statistical analysis</u>

Data were analyzed using GraphPad Prism Software (Graphpad, San Diego, CA) and

presented as means ± S.D. unless otherwise specified with p values calculated by

unpaired Student's t-tests, one-way or two-way ANOVA (comparisons across more than

two groups) and adjusted with Bonferroni's corrections.

517 **Acknowledgements**

518 We thank the *Caenorhabditis* Genetics Center, National BioResource Project in Japan

519 and the Million Mutation Project for *C. elegans* strains. The work was supported by NIH

520 grants R01GM117461, R00HL116654, ADA grant 1-16-IBS-197, Pew Scholar Award,

521 Alfred P. Sloan Foundation Fellowship, and Packard Fellowship in Science and

522 Engineering (D.K.M), R01AG032435 (Y.W.) and a CPSF postdoctoral fellowship (W.J.).

524 **References**

Al-Fageeh, M.B., and Smales, C.M. (2006). Control and regulation of the cellular responses to
cold shock: the responses in yeast and mammalian systems. Biochem. J. *397*, 247–259.

Bailey, T.L., Boden, M., Buske, F.A., Frith, M., Grant, C.E., Clementi, L., Ren, J., Li, W.W., and
Noble, W.S. (2009). MEME SUITE: tools for motif discovery and searching. Nucleic Acids Res.
37, W202-208.

Baird, N.A., Douglas, P.M., Simic, M.S., Grant, A.R., Moresco, J.J., Wolff, S.C., Yates, J.R.,
Manning, G., and Dillin, A. (2014). HSF-1-mediated cytoskeletal integrity determines
thermotolerance and life span. Science *346*, 360–363.

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W.-H.,
Pagès, F., Trajanoski, Z., and Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher
functionally grouped gene ontology and pathway annotation networks. Bioinforma. Oxf. Engl.
25, 1091–1093.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71–94.

Brunquell, J., Morris, S., Lu, Y., Cheng, F., and Westerheide, S.D. (2016). The genome-wide
role of HSF-1 in the regulation of gene expression in Caenorhabditis elegans. BMC Genomics
17.

Choi, H.A., Badjatia, N., and Mayer, S.A. (2012). Hypothermia for acute brain injury--
mechanisms and practical aspects. Nat. Rev. Neurol. *8*, 214–222.

Chüeh, A.C., Tse, J.W., Dickinson, M., Ioannidis, P., Jenkins, L., Tögel, L., Tan, B., Luk, I.,
Dávalos-Salas, M., Nightingale, R., et al. (2017). ATF3 repression of BCL-XL determines
apoptotic sensitivity to HDAC inhibitors across tumour types. Clin. Cancer Res.
clincanres.0466.2017.

Davis, M.W., Hammarlund, M., Harrach, T., Hullett, P., Olsen, S., and Jorgensen, E.M. (2005).
Rapid single nucleotide polymorphism mapping in C. elegans. BMC Genomics *6*, 118.

Dempersmier, J., Sambeat, A., Gulyaeva, O., Paul, S.M., Hudak, C.S.S., Raposo, H.F., Kwan,
H.-Y., Kang, C., Wong, R.H.F., and Sul, H.S. (2015). Cold-inducible Zfp516 activates UCP1
transcription to promote browning of white fat and development of brown fat. Mol. Cell *57*, 235–
246.

Dix, I., Russell, C., Yehuda, S.B., Kupiec, M., and Beggs, J.D. (1999). The identification and
characterization of a novel splicing protein, Isy1p, of Saccharomyces cerevisiae. RNA N. Y. N *5*,
360–368.

Du, P., Wang, L., Sliz, P., and Gregory, R.I. (2015). A Biogenesis Step Upstream of
Microprocessor Controls miR-17~92 Expression. Cell *162*, 885–899.

Fan, W., and Evans, R.M. (2015). Turning up the heat on membrane fluidity. Cell *161*, 962–963.

Friedländer, M.R., Mackowiak, S.D., Li, N., Chen, W., and Rajewsky, N. (2012). miRDeep2
accurately identifies known and hundreds of novel microRNA genes in seven animal clades.
Nucleic Acids Res. *40*, 37–52.

Frøkjaer-Jensen, C., Davis, M.W., Hopkins, C.E., Newman, B.J., Thummel, J.M., Olesen, S.-P.,
Grunnet, M., and Jorgensen, E.M. (2008). Single-copy insertion of transgenes in Caenorhabditis
elegans. Nat. Genet. *40*, 1375–1383.

Galej, W.P., Wilkinson, M.E., Fica, S.M., Oubridge, C., Newman, A.J., and Nagai, K. (2016).
Cryo-EM structure of the spliceosome immediately after branching. Nature *537*, 197–201.

Garrity, P.A., Goodman, M.B., Samuel, A.D., and Sengupta, P. (2010). Running hot and cold:
behavioral strategies, neural circuits, and the molecular machinery for thermotaxis in C. elegans
and Drosophila. Genes Dev. *24*, 2365–2382.

Giuliodori, A.M., Di Pietro, F., Marzi, S., Masquida, B., Wagner, R., Romby, P., Gualerzi, C.O.,
and Pon, C.L. (2010). The cspA mRNA is a thermosensor that modulates translation of the cold-
shock protein CspA. Mol. Cell *37*, 21–33.

Gomez-Pastor, R., Burchfiel, E.T., and Thiele, D.J. (2017). Regulation of heat shock
transcription factors and their roles in physiology and disease. Nat. Rev. Mol. Cell Biol.

Hamilton, W.D. (1963). The Evolution of Altruistic Behavior. Am. Nat. *97*, 354–356.

Hartman, M.G., Lu, D., Kim, M.-L., Kociba, G.J., Shukri, T., Buteau, J., Wang, X., Frankel, W.L.,
Guttridge, D., Prentki, M., et al. (2004). Role for Activating Transcription Factor 3 in Stress-
Induced β-Cell Apoptosis. Mol. Cell. Biol. *24*, 5721–5732.

Hwang, A.B., and Lee, S.-J. (2011). Regulation of life span by mitochondrial respiration: the
HIF-1 and ROS connection. Aging *3*, 304–310.

Jan, C.H., Friedman, R.C., Ruby, J.G., and Bartel, D.P. (2011). Formation, regulation and
evolution of Caenorhabditis elegans 3'UTRs. Nature *469*, 97–101.

Ji, N., and Oudenaarden, A. van (2005). Single molecule fluorescent in situ hybridization
(smFISH) of C. elegans worms and embryos (WormBook).

John, B., Enright, A.J., Aravin, A., Tuschl, T., Sander, C., and Marks, D.S. (2004). Human
MicroRNA targets. PLoS Biol. *2*, e363.

Kamath, R.S., and Ahringer, J. (2003). Genome-wide RNAi screening in Caenorhabditis
elegans. Methods San Diego Calif *30*, 313–321.

Kandror, O., Bretschneider, N., Kreydin, E., Cavalieri, D., and Goldberg, A.L. (2004). Yeast
adapt to near-freezing temperatures by STRE/Msn2,4-dependent induction of trehalose
synthesis and certain molecular chaperones. Mol. Cell *13*, 771–781.

Kato, M., Kashem, M.A., and Cheng, C. (2016). An intestinal microRNA modulates the
homeostatic adaptation to chronic oxidative stress in C. elegans. Aging *8*, 1979–2005.

Kershner, A.M., and Kimble, J. (2010). Genome-wide analysis of mRNA targets for
Caenorhabditis elegans FBF, a conserved stem cell regulator. Proc. Natl. Acad. Sci. U. S. A.
107, 3936–3941.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low
memory requirements. Nat. Methods *12*, 357–360.

Krüger, J., and Rehmsmeier, M. (2006). RNAhybrid: microRNA target prediction easy, fast and
flexible. Nucleic Acids Res. *34*, W451-454.

Kumsta, C., Chang, J.T., Schmalz, J., and Hansen, M. (2017). Hormetic heat stress and HSF-1
induce autophagy to improve survival and proteostasis in C. elegans. Nat. Commun. *8*, 14337.

Landis, J.N., and Murphy, C.T. (2010). Integration of diverse inputs in the regulation of
Caenorhabditis elegans DAF-16/FOXO. Dev. Dyn. Off. Publ. Am. Assoc. Anat. *239*, 1405–1412.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat.
Methods *9*, 357–359.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. Bioinforma. Oxf. Engl. *30*, 923–930.

Longo, V.D., Mitteldorf, J., and Skulachev, V.P. (2005). Programmed and altruistic ageing. Nat.
Rev. Genet. *6*, 866–872.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

Lyons, J.M., Keith, A.D., and Thomason, I.J. (1975). Temperature-induced phase transitions in
nematode lipids and their influence on respiration. J. Nematol. *7*, 98–104.

Ma, D.K., Vozdek, R., Bhatla, N., and Horvitz, H.R. (2012). CYSL-1 interacts with the O2-
sensing hydroxylase EGL-9 to promote H2S-modulated hypoxia-induced behavioral plasticity in
C. elegans. Neuron *73*, 925–940.

Ma, D.K., Li, Z., Lu, A.Y., Sun, F., Chen, S., Rothe, M., Menzel, R., Sun, F., and Horvitz, H.R.
(2015). Acyl-CoA Dehydrogenase Drives Heat Adaptation by Sequestering Fatty Acids. Cell
161, 1152–1163.

Maere, S., Heymans, K., and Kuiper, M. (2005). BiNGO: a Cytoscape plugin to assess
overrepresentation of gene ontology categories in biological networks. Bioinforma. Oxf. Engl.
21, 3448–3449.

Mahat, D.B., Salamanca, H.H., Duarte, F.M., Danko, C.G., and Lis, J.T. (2016). Mammalian
Heat Shock Response and Mechanisms Underlying Its Genome-wide Transcriptional
Regulation. Mol. Cell *62*, 63–78.

Mello, C.C., Kramer, J.M., Stinchcomb, D., and Ambros, V. (1991). Efficient gene transfer in
C.elegans: extrachromosomal maintenance and integration of transforming sequences. EMBO
J. *10*, 3959–3970.

Metzstein, M.M., Stanfield, G.M., and Horvitz, H.R. (1998). Genetics of programmed cell death
in C. elegans: past, present and future. Trends Genet. TIG *14*, 410–416.

Morrison, S.F. (2016). Central control of body temperature. F1000Research *5*.

Murray, P., Hayward, S.A.L., Govan, G.G., Gracey, A.Y., and Cossins, A.R. (2007). An explicit
test of the phospholipid saturation hypothesis of acquired cold tolerance in Caenorhabditis
elegans. Proc. Natl. Acad. Sci. U. S. A. *104*, 5489–5494.

Ohta, A., Ujisawa, T., Sonoda, S., and Kuhara, A. (2014). Light and pheromone-sensing
neurons regulates cold habituation through insulin signalling in *Caenorhabditis elegans*. Nat.
Commun. *5*, ncomms5412.

Peng, Y., Zhang, M., Zheng, L., Liang, Q., Li, H., Chen, J.-T., Guo, H., Yoshina, S., Chen, Y.-Z.,
Zhao, X., et al. (2017). Cysteine protease cathepsin B mediates radiation-induced bystander
effects. Nature *547*, 458–462.

Peter, M.E. (2011). Programmed cell death: Apoptosis meets necrosis. Nature *471*, 310–312.

Polderman, K.H. (2009). Mechanisms of action, physiological effects, and complications of
hypothermia. Crit. Care Med. *37*, S186-202.

Ritchie, A., Gutierrez, O., and Fernandez-Luna, J.L. (2009). PAR bZIP-bik is a novel
transcriptional pathway that mediates oxidative stress-induced apoptosis in fibroblasts. Cell
Death Differ. *16*, 838–846.

Sapolsky, R.M. (2004). Why Zebras Don't Get Ulcers, Third Edition (New York: Holt
Paperbacks).

Schmieder, R., and Edwards, R. (2011). Quality control and preprocessing of metagenomic
datasets. Bioinforma. Oxf. Engl. *27*, 863–864.

Skulachev, V.P. (1999). Phenoptosis: programmed death of an organism. Biochem. Biokhimiia
64, 1418–1426.

Skulachev, V.P. (2002). Programmed death phenomena: from organelle to organism. Ann. N. Y.
Acad. Sci. *959*, 214–237.

Smith, J.M. (1964). Group Selection and Kin Selection. Nature *201*, 1145–1147.

Solís, E.J., Pandey, J.P., Zheng, X., Jin, D.X., Gupta, P.B., Airoldi, E.M., Pincus, D., and Denic,
V. (2016). Defining the Essential Function of Yeast Hsf1 Reveals a Compact Transcriptional
Program for Maintaining Eukaryotic Proteostasis. Mol. Cell *63*, 60–71.

Tansey, E.A., and Johnson, C.D. (2015). Recent advances in thermoregulation. Adv. Physiol.
Educ. *39*, 139–148.

Thorvaldsdóttir, H., Robinson, J.T., and Mesirov, J.P. (2013). Integrative Genomics Viewer
(IGV): high-performance genomics data visualization and exploration. Brief. Bioinform. *14*, 178–
192.

Yenari, M.A., and Han, H.S. (2012). Neuroprotective mechanisms of hypothermia in brain
ischaemia. Nat. Rev. Neurosci. *13*, 267–278.

Yura, T., Nagai, H., and Mori, H. (1993). Regulation of the heat-shock response in bacteria.
Annu. Rev. Microbiol. *47*, 321–350.

Zhou, X., Feng, X., Mao, H., Li, M., Xu, F., Hu, K., and Guang, S. (2017). RdRP-synthesized
antisense ribosomal siRNAs silence pre-rRNA via the nuclear RNAi pathway. Nat. Struct. Mol.
Biol. *24*, 258–269.

Zhu, J.-K. (2016). Abiotic Stress Signaling and Responses in Plants. Cell *167*, 313–324.

Figure legends

Figure 1: *asp-17* is robustly and specifically induced by cold-warming. A, Volcano

plot of RNA-seq showing differentially regulated genes (up-regulated genes in red;

down-regulated genes in green; *asp-17* was indicated by purple dot) by cold-warming in

wild type *C. elegans*. **B**, RNA-seq reads at the *asp-17* locus showing consistent up-

regulation of *asp-17* transcript levels in triplicate samples after cold-warming. **C**, QPCR

measurements of gene expression levels showing up-regulation of three representative

CW inducible genes, including *asp-17*, after shifting from 25°C to different degrees of

hypothermia (4, 10, 15 and 22°C) lasting 2 hrs followed by recovery at 25°C for 0.5 hr.

D, Quantification of RNA-seq reads indicating specific up-regulation of *asp-17* but not

other members of the *asp* family genes (only expressed *asp* genes are shown). **E**,

QPCR measurements of *asp-17* levels under conditions of indicated durations of cold

and warming (with or without 25°C recovery for "yes/no"). **F**, smFISH images showing

asp-17 up-regulation (signals indicated by yellow) by CW predominantly in the intestine

of wild type but not *asp-17* KO animals. n ≥ 20 total animals for each group with N ≥ 3

independent biological replicates; *** indicates P < 0.001. Scale bar: 50 μm.

Figure 2: A forward genetic screen identifies *C. elegans isy-1* as a causal

regulator of *asp-17*::GFP expression. A, Schematic of the *C. elegans* ISY-1 protein

showing the domain structure predicted by SMART (top) and a multiple sequence

alignment of ISY-1 homologues from major metazoans showing the conservation of the

glutamate residue substituted to lysine by the *dma50* mutation isolated from EMS

screens. **B**, Nomarski and fluorescence images showing the phenotype of intestinal

asp-17p::GFP in wild type and *dma50* mutants. **C**, Fluorescence images showing the

distribution of ISY-1::GFP driven by the endogenous *isy-1* promoter. Arrows indicate

neuronal, hypodermal and intestinal nuclei. **D**, Nomarski and fluorescence images

showing intestinal *asp-17p*::GFP in a transgenic strain expressing RNAi against *isy-1*

specifically in intestine. **E**, Schematic of the *C. elegans isy-1* gene structure with

mCherry tagged at the C-terminus (top); Nomarski and fluorescence images showing

rescue of *dma50* by various transgenes (below). **F,** Graph quantification of fluorescence

intensities showing rescue of *dma50* in *asp-17p*::GFP activation. N ≥ 3 independent

biological replicates; *** indicates P < 0.001. Scale bar: 20 μm.

Figure 3: ZIP-10 acts downstream of ISY-1 and mediates transcriptional response

to CW. A, QPCR measurements of *asp-17* levels induced by CW in wild type and *isy-*

1(dma50) mutants. **B**, Volcano plot of RNA-seq showing differentially regulated genes

(up-regulated genes in red; down-regulated genes in green) in *isy-1* mutants compared

with wild type. **C**, RNA-seq measurements of expression levels for indicated genes in

wild type and *isy-1(dma50)* mutants. **D**, Nomarski and fluorescence images showing

asp-17p::*GFP* induction by *isy-1* RNAi was blocked in *zip-10* mutants. **E**, Western blots

of the integrated *zip-10p::zip-10::EGFP::FLAG* strain showing time-dependent protein

induction by CW. **F**, Western blots of the integrated *zip-10p::zip-10::EGFP::FLAG* strain

showing its up-regulation by *isy-1* RNAi and down-regulation by *GFP* or *zip-10* RNAi.

Both short- and long-exposure blots are shown. **G**, Western blots of the integrated *zip-*

10p::zip-10::EGFP::FLAG strain showing its up-regulation strictly required warming after

cold shock. **H**, Western blots of the integrated *zip-10*p::*zip-10::EGFP::FLAG* strain

showing its up-regulation by CW was further enhanced by *isy-1* RNAi. **I**, QPCR

measurements of gene expression levels showing ZIP-10 dependent up-regulation of

asp-17 and *cpr-3* but not *srr-6* or *F53B9.1* after cold for indicated durations and 1-hr

warming. n ≥ 20 total animals for each group with N ≥ 3 independent biological

replicates; *** indicates P < 0.001. Scale bar: 20 μm.

Figure 4: ISY-1 regulates *zip-10* via *mir-60*. A, Western blot of mCherry/RFP-trapped

RNA immunoprecipitates in animals treated with or without cold-warming. **B,** QPCR

measurements of the percent input for primary *mir-60* transcripts from mCherry/RFP-

trapped RNA immunoprecipitates in animals treated with or without cold-warming. **C,**

QPCR measurements of the percent input for *cbp-1* transcripts from mCherry/RFP-

trapped RNA immunoprecipitates in animals treated with or without cold-warming. **D,**

QPCR measurements of the mature *mir-60* transcript levels from wild type, *isy-

1(dma50)* and *mir-60(n4947)* deletion mutants. **E,** QPCR measurements of the levels of

CW-inducible gene transcripts in animals with indicated genotypes and conditions. **F,**

Western blot of lysates from animals carrying the array *ges-1p::mCherry::3'utr(zip-10)*

reporters with CW or *isy-1* RNAi. No change of reporter activity was observed. **G,**

Western blot of lysates from animals carrying *zip-10p::zip-10::EGFP::FLAG* reporters

with various indicated CW and RNAi conditions. n ≥ 20 total animals for each group with

N ≥ 3 independent biological replicates; *** indicates P < 0.001; ** indicates P < 0.01.

Figure 5: ISY-1 and ZIP-10 regulate a genetic program to promote organismic

death. A, Venn diagram indicating numbers of genes commonly regulated by CW and

isy-1(dma50) mutants. **B**, Scatter plot depicting the correlation between the

transcriptome response to cold (y-axis) and the transcriptome response to *isy-1*

mutation (x-axis). Shown are the log2 fold changes compared with corresponding

controls. Pearson correlation coefficient and the associated P-value were calculated

using R functions. **C**, AT-rich motif identified by MEME enriched among the CW and *isy-*

1(dma50) regulated genes, with a table listing numbers of the motif present in top-

ranked 4 genes. **D**, ChIP-QPCR measurements of ZIP-10::FLAG binding to the *asp-17*

promoter. **E**, Survival rates of indicated genotypes after prolonged CW (4°C for 48 hrs

followed by 4 hrs of warming). **F**, Organismic death rates of indicated genotypes after

heat shock (32°C) induction of *zip-10* wild type, mutant with defective transactivation C-

terminus, *zip-2* and *zip-11* (left), without CW. Nomarski image (right) indicate

morphologies of normal and dead animals with induction of *zip-10*. **G**, RNA-seq

measurements of gene targets of indicated TFs. **H**, Table showing the *asp-17p::GFP*

and cold tolerance phenotypes of animals with indicated genotypes. **I**, model for the role

and regulation of the ZIP-10 pathway. n ≥ 20 total animals for each group with N ≥ 3

independent biological replicates; *** indicates P < 0.001.

Figure 1-figure supplement 1: RNA-seq identified genes up-regulated by cold-

warming. A, Hierarchical clustering analysis of genes up- and down-regulated genes by

CW. **B**, Gene ontology analysis of genes up- and down-regulated genes by CW.

Figure 2-figure supplement 1: EMS screens identified *isy-1* as a regulator of *asp-*

***17*. A**, Fluorescence images showing up-regulation of *asp-17p::GFP* but not the co-

injection marker *unc-54p::mCherry* by CW. **B**, Schematic illustrating the workflow of

EMS mutagenesis to isolate mutants with abnormal *asp-17p::GFP* expression. **C**, Table

showing the mutants isolated from the screen, including *dma50* that defines the gene

F53B7.3, named as *isy-1*. **D**, Chromosome genetic maps showing DraI SNP sites and

the mapped interval for *dma50*. **E**, Table listing candidate genes with mutations

identified by WGS, with two genes highlighted in yellow within the mapped interval. **F**,

fluorescence images showing that RNAi of *isy-1* phenocopies *dma50*. N ≥ 3

independent biological replicates; *** indicates P < 0.001. Scale bar: 20 μm.

Figure 2-figure supplement 2: ISY-1 is a *C. elegans* member of evolutionarily

conserved protein family and its level is not regulated by cold-warming. A,

Multiple sequence alignment (Cluster Omega, visualized by Jalview) of ISY-1 family

members in major metazoan species. **B**, Cladogram of ISY family proteins. **C**, Sanger

sequencing confirmation of *isy-1(dma50)*. **D,** RNA-seq measurements of *isy-1* and *asp-*

17 levels under CW conditions and in *isy-1(dma50)* mutants. **E**, Western blot of *isy-*

*1*p::ISY-1::mCherry showing no regulation by CW. N ≥ 3 independent biological

replicates; *** indicates P < 0.001.

Figure 3-figure supplement 1: ISY-1 does not affect general splicing of a GFP

reporter nor specific splicing of *zip-10*. A, Nomarski and fluorescence images

showing that animals with *isy-1* RNAi are not defective in general intron splicing of an

eft-3p::GFP (intron-containing) reporter. **B**, Read plot from RNA-seq of wild type at the

zip-10 locus. **C**, Read plot from RNA-seq of *isy-1(dma50)* mutants at the *zip-10* locus

indicating the intact splicing of the *zip-10* intron despite increased mRNA levels. **D,**

Normalized QPCR measurements of mRNA expression levels of *isy-1* and *zip-10* from

wild type and *isy-1(dma50)* mutants. **E,** Normalized QPCR measurements of mRNA

expression levels of *isy-1* and *zip-10* from wild type and CW-treated animals.

Figure 3-figure supplement 2: Mechanisms of regulation and function of *zip-10*. A,

Nomarski and fluorescence images showing the *zip-10p::GFP* reporter activated by *isy-*

1 RNAi. **B**, Western blot of lysates from *zip-10p::zip-10::EGFP::FLAG* transgenic

animals showing its increased abundance after CW but not by hypoxia (0.5% O_2) or

starvation for 24 hrs. **C**, Western blot of fractionated lysates from *zip-10p::zip-*

10::EGFP::FLAG transgenic animals after CW showing its increased abundance in

cytosol and nucleus. **D**, Western blot of lysates from *zip-10p::zip-10::EGFP::FLAG*

transgenic animals showing its increased abundance after CW when warming was

permitted and cold exposure was prolonged. **E**, Western blot of lysates from *zip-*

10p::zip-10::FLAG transgenic and various RNAi treated animals showing its increased

abundance after CW in *isy-1* RNAi treated animals but was unaffected by RNAi against

C37C3.2, encoding the *C. elegans* orthologue of translation initiation factor 5 (eIF5), or

T26A8.4, encoding the *C. elegans* orthologue of *Saccharomyces cerevisiae* Caf120, a

component of the Ccr4-Not deadenylase RNA-degrading complex. **F**, ZIP-10 domain

organization based on analysis by SMART (smart.embl-heidelberg.de) and modeled

structure of ZIP-10's bZIP domain (swissmodel.expasy.org). **G**, Sequence alignment of

ZIP-10's bZIP domain and the Maf transcription factor as a template of the modelled

ZIP-10 structure. **H**, Schematic of the *zip-10* locus showing the wild type, EGFP::FLAG

tagged allele, and the wild-type and C-terminal transactivation mutant ZIP-10 protein

sequences. The mutated residues (from Q and N to V and A, respectively) at the ZIP-10

C-terminus are completely conserved in nematodes.

Figure 4-figure supplement 1: A specific subset of CW-inducible genes is

dependent on ZIP-10. Shown is a table of fold induction for gene expression levels

determined by QPCR measurements of randomly selected CW-inducible genes in wild

type and *zip-10* mutant animals.

Figure 5-figure supplement 1: Identification of differentially regulated microRNAs

in wild type and *isy-1(dma50)* mutants. A, Summary table showing the read statistics

for the small RNA-seq experiments using three biologically triplicate samples in wild

type and *isy-1(dma50)* mutants**. B**, Small RNA length distribution plots for wild type and

isy-1(dma50) mutants. **C**, Distribution graphs for various types of small RNAs identified

by small RNA sequencing for wild type and *isy-1(dma50)* mutants. **D**, Volcano plot

showing the differentially regulated microRNAs, including *mir-60*, in the *isy-1(dma50)*

mutants (three replicates each). **E**, Schematic of two examples illustrating the TF-

encoding genes, *skn-1* and *daf-16* at 3' UTRs, targeted by two microRNAs, *mir-51* and

mir-359, respectively, which are down-regulated in *isy-1(dma50)* mutants. **F**,

Hierarchical clustering analysis of microRNAs up- or down-regulated in *isy-1(dma50)*

mutants. *mir-51* and *mir-359* down-regulation were indicated by arrows.

Figure 5-figure supplement 2: ZIP-10 constitutes a genetic program promoting kin selection of *C. elegans* under resource-limiting and stress conditions. A, Mathematical model illustrating the advantage of wild-type (red line) versus *zip-10* KO animals (blue line) at the population level under growth resource-liming and thermal stress conditions. Individual fitness distribution (x-axis) after cold-warming thermal stress is set to be exponential (thin red line) and linear (thin blue line) to fit the phenotypic differences in death rates of wild type and *zip-10* KO animals and to simplify comparison and calculation. Given the same period of time and the same amount of growth resources (integral differences between thin and thick lines), wild-type population yields more thermal stress-adapted reproducing animals than *zip-10* deficient animals. The parameters for integrals were adjusted so that the "growth resources" are the same despite that the population growth modes are different for wild type and *zip-10* KO. **B**, Schematic illustrating an exemplar situation of the mathematical model in which 5 post-reproduction adults and 5 larvae are living in a food-limiting condition. After cold-warming thermal stress, wild-type adults die out, leaving food for young larvae to grow into reproductive adults whereas *zip-10* KO post-reproduction adults do not die, consuming the limited food and competing out the young larvae so that fewer reproductively active *zip-10* KO animals emerge from such conditions. **C**, Experimental evidence indicating the specific induction of *zip-10p::zip-10::EGFP::FLAG* by CW as measured by Western blot in adults but not in larvae. **D**, Survival rates of wild type and *zip-10* KO (*ok3462*) animals at different stages after prolonged CW thermal stresses (96-hr cold shock for L1 while 48-hr for others) showing that ZIP-10's pro-death

effects are more prominent in old animals. n ≥ 20 total animals for each group with N ≥

3 independent biological replicates; *** indicates P < 0.001, ** indicates P < 0.01.

Table 1: Lists of genes up- and down-regulated by CW with adjusted P < 0.05 and

log2FoldChange from biological triplicate samples of wild-type *C. elegans*.

Table 2: Lists of genes up- and down-regulated by the *isy-1(dma50)* mutation with

adjusted P < 0.05 and log2FoldChange from biological triplicate samples of wild-type

and *isy-1(dma50)* mutant *C. elegans*.

Table 3: Lists of genes commonly regulated by CW and the *isy-1(dma50)* mutation with

adjusted P < 0.05 and log2FoldChange from biological triplicate samples of wild-type

and *isy-1(dma50)* mutant *C. elegans*.

A

Coiled coil   Coiled coil   Low complexity

Human
Mouse
Zebrafish
Drosophilia
C. elegans

isy-1(dma50)
GAA -> AAA [Glu -> Lys]

B

asp-17p::GFP   merge

dmaIs10[asp-17p::GFP]

asp-17p::GFP   merge

isy-1(dma50); dmaIs10[asp-17p::GFP]

C   *Ex[isy-1p::isy-1::GFP]*

neuronal
nuclei

intestinal
nuclei

hypodermal
nuclei

D

normaski

asp-17p::GFP

control RNAi   *ges-1p::isy-1* RNAi

E Rescue by single-copy MosSCI insertion:

100 bp

isy-1 promoter   *isy-1* coding exon   mCherry::tbb-2 3'utr

Chr2: *ttTi5605* locus

normaski

asp-17p::GFP

no   Mos   *isy-1p::*   *ges-1p::*
rescue   SCI   *isy-1*   *isy-1*

dmaIs10   *isy-1; dmaIs10*

F

GFP intensity (A.U.)

*** **

dmaIs10   no rescue MosSCI *isy-1::isy-1 ges-1::isy-1*

dmaIs10   *isy-1; dmaIs10*

A CW-induced genes / *isy-1(dma50)*-induced genes: 182, 246, 632

B R = 0.53; Log2 (CW vs control) vs Log2 (*isy-1* vs control)

C
	Numbers of motifs
asp-17p	16
F08G2.5p	5
clec-66p	2
M02H5.8p	1

D *asp-17* promoter ChIP; Percent input (%); IgG, Flag (M2); Control, Cold-warming; ***

E Survival rate after CW (%); Wild type, *asp-17* KO, *zip-10* KO; ***

F Organismic death rate (%); HS only; *hsp-16*p driven; ***
1, *zip-10* (+)
2, *zip-10* transactivation mutant
3, *zip-2* (+)
4, *zip-11* (+)
hsp-16p::zip-10; Dead, Alive, Dead

G target genes controlled by; Normalized levels by RNAseq; Control, *isy-1(dma50)*; HIF-1: *cysl-2*, *nhr-57*; HSF-1: *hsp-16.2*, *hsp-16.41*; DAF-16: *dod-19*, *dod-22*; **

H
Genotype	*asp-17p*::GFP	Cold tolerance (n > 100)
Wild type	+	32%
isy-1(dma50) or isy-1 RNAi	+++++	57%
zip-10(ok3462); isy-1 RNAi	+	82%
zip-10(ok3462)	+	78%
mir-60(n4947)	++++	29%
mir-60(n4947); zip-10(ok3462)	+	51%
asp-17(dma99)	+	68%
asp-17(dma99); isy-1 RNAi	+++++	72%
daf-16(mu86)	+	10%
daf-16(mu86); isy-1 RNAi	+++++	38%

I Mild other stress; ISY-1 and co-factors ⊣ HIF-1 DAF-16 etc. TFs → Organismic Health; *mir-60*; Cold shock - warming → ZIP-10 → *asp-17 cpr-3* etc. → Organismic Death (adults)

GOID	GOTerm	Term P-Value
KEGG:00071	Fatty acid degradation	0.068035515
KEGG:00330	Arginine and proline metabolism	0.012703481
KEGG:00564	Glycerophospholipid metabolism	0.027845609
KEGG:02010	ABC transporters	6.87E-07
KEGG:04068	FoxO signaling pathway	0.099454727
KEGG:04136	Autophagy	0.010025137
KEGG:04137	Mitophagy	0.002142392
KEGG:04140	Autophagy	0.007908727
KEGG:04141	Protein processing in endoplasmic reticulum	0.129746247
KEGG:04144	Endocytosis	0.005199927
KEGG:04310	Wnt signaling pathway	0.050156699
KEGG:04350	TGF-beta signaling pathway	0.034403422

Figure 4-Figure Supplement 1: Relative mRNA levels of CW-inducible genes in wild-type and *zip-10* mutants.

Gene		1	2	3	4	*P* value
asp-17	WT	1.20±0.22	9.20±0.07	20.38±12.77	45.45±12.47	
	zip-10 m	0.37±0.04	0.46±0.12	0.71±0.29	0.43±0.10	0.025
cpr-3	WT	1.01±0.18	2.80±0.16	3.67±0.10	9.75±1.43	
	zip-10 m	1.24±0.05	1.31±0.48	1.99±0.40	1.30±0.52	0.005
ZK896.4	WT	1.01±0.20	3.56±1.27	6.31±1.13	12.22±2.00	
	zip-10 m	1.19±0.07	0.76±0.07	1.14±0.07	1.03±0.21	0.009
F53A9.1	WT	0.80±0.29	1.27±0.48	2.78±1.07	5.51±2.29	
	zip-10 m	0.69±0.19	1.50±0.42	2.60±0.62	4.95±1.56	0.748
srr-6	WT	1.09±0.29	4.68±2.84	18.31±15.68	55.37±33.32	
	zip-10 m	1.74±0.53	37.31±22.32	136.49±46.46	188.14±64.10	0.050
ceh-37	WT	1.04±0.35	1.75±0.15	2.36±0.56	5.52±1.12	
	zip-10 m	0.76±0.40	1.76±0.65	2.58±0.57	3.53±0.59	0. 072
fis-2	WT	1.09±0.57	0.94±0.36	2.92±0.09	8.57±2.12	
	zip-10 m	2.19±0.47	3.55±0.44	4.57±0.24	11.93±1.98	0. 115
CYP14A5	WT	0.98±0.14	2.53±0.46	7.54±2.90	26.04±5.55	
	zip-10 m	3.99±1.29	6.77±0.77	10.12±0.78	28.57±5.66	0. 610
sqst-1	WT	0.98±0.29	2.59±1.15	4.95±1.08	13.59±4.28	
	zip-10 m	2.68±0.54	5.81±0.25	8.13±0.59	12.85±2.95	0. 820
best-5	WT	0.97±0.31	1.79±0.40	5.34±3.24	18.21±6.83	
	zip-10 m	1.37±0.97	2.94±0.26	6.06±2.23	13.35±4.41	0. 368
tsp-1	WT	1.05±0.41	6.78±2.86	18.03±6.33	72.47±13.72	
	zip-10 m	3.04±2.34	12.53±4.52	23.45±2.73	50.02±9.17	0. 108
dod-3	WT	1.08±0.54	9.42±4.50	18.04±2.08	33.69±6.53	
	zip-10 m	2.64±1.17	8.11±2.47	13.19±3.75	28.77±6.74	0. 415
oac-20	WT	1.00±0.09	4.19±0.56	7.92±0.49	19.54±3.53	
	zip-10 m	1.38±0.08	2.90±1.17	5.05±1.01	16.20±4.87	0. 396
F37C4.5	WT	1.01±0.17	2.10±0.38	2.79±1.84	8.94±2.34	
	zip-10 m	1.43±0.18	1.49±0.27	2.19±0.33	5.28±1.69	0. 100
dod-19	WT	1.08±0.45	0.95±0.01	0.67±0.02	1.41±0.37	
	zip-10 m	1.33±0.07	0.85±0.09	0.97±0.18	1.75±0.23	0. 262
dod-22	WT	1.15±0.74	1.52±0.1.	1.27±0.22	2.76±0.50	
	zip-10 m	1.41±0.25	1.29±0.34	2.55±0.24	3.41±0.59	0. 226

oac-6	WT	1.01±0.12	3.07±0.77	5.36±1.28	10.94±3.72	
	zip-10 m	1.79±0.25	2.34±1.30	4.80±0.98	8.22±1.61	0. 336
K09D9.1	WT	1.05±0.38	1.05±0.58	4.01±2.82	24.56±3.64	
	zip-10 m	3.31±1.14	7.38±1.61	11.55±3.07	27.66±8.46	0. 605
clec-41	WT	1.08±0.49	0.83±0.38	1.37±0.17	2.31±0.54	
	zip-10 m	1.63±0.27	0.35±0.21	0.68±0.26	1.26±0.19	0. 065
F48E3.8	WT	1.17±0.72	0.87±0.42	2.07±0.97	5.30±1.14	
	zip-10 m	0.99±0.27	0.60±0.49	1.22±0.08	5.66±0.66	0. 666
cebp-1	WT	1.04±0.34	1.44±0.22	2.84±0.34	6.80±3.05	
	zip-10 m	2.21±0.69	2.07±0.37	3.01±0.14	4.58±0.90	0. 335

WT: wild type;
zip-10 m: *zip-10* mutant (*ok3462*);
1. 25^0C;
2. 4^0C-1h, 25^0C-1h;
3. 4^0C-2h, 25^0C-1h;
4. 4^0C-4h, 25^0C-1h.
n=3 biological replicates.

$$Growth\ Resource = \int_0^{100} (\beta_{wt} - 1) * \left(\frac{x}{100}\right)^{10} * 100\ dx = \int_0^{100} (\gamma_{ko} - 1) * 0.2 * x dx$$