A bHLH-PAS protein regulates light-dependent diurnal rhythmic processes in the marine diatom *Phaeodactylum tricornutum*

Rossella Annunziata[a,b,1], Andrés Ritter[a,1], Antonio Emidio Fortunato[a,c,1], Soizic Cheminant-Navarro[a], Nicolas Agier[a], Marie J. J. Huysman[a,d,e*], Per Winge[f], Atle Bones[f], François-Yves Bouget[h], Marco Cosentino Lagomarsino[a], Jean Pierre Bouly [a], and Angela Falciatore[a,2].

[a]Sorbonne Université, CNRS, Institut de Biologie Paris-Seine, Laboratory of Computational and Quantitative Biology, F-75005, Paris, France.

[b]Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli, Italy.

[c]ImmunRise Technologies, 75006 Paris, France.

[d]Ghent University, Department of Plant Biotechnology and Bioinformatics, Technologiepark 927, B-9052 Ghent, Belgium.

[e]VIB Center for Plant Systems Biology, Technologiepark 927, B-9052 Ghent, Belgium.

[f]Department of Biology, NTNU Norwegian University of Science and Technology, 7491 Trondheim, Norway.

[h]Sorbonne Université, UMR 7621, Laboratoire d'Océanographie Microbienne, Observatoire Océanologique, Banyuls-sur-Mer, France.

[1]These authors contributed equally to this work.

*Disclaimer: The author (MJJH) is currently an employee of the European Research Council Executive Agency. The views expressed are purely those of the writer and may not in any circumstances be regarded as stating an official position of the European Commission.

[2]Corresponding Author:

Angela Falciatore

Diatom Functional Genomics Team

Laboratory of Computational and Quantitative Biology (LCQB)

Institut de Biologie Paris-Seine (IBPS)

Jussieu Campus – Sorbonne Université

4, Place Jussieu, 75005 Paris, France

Tel: +33 0144278142

Email: angela.falciatore@upmc.fr

Keywords: Diatoms, diurnal rhythms, transcription factors, cell division, gene expression.

ABSTRACT

Periodic light-dark cycles govern the timing of basic biological processes in organisms inhabiting land as well as the sea, where life evolved. Although prominent marine phytoplanktonic organisms such as diatoms show robust diurnal rhythms in growth, cell cycle and gene expression, the molecular foundations controlling these processes are still obscure. By exploring the regulatory landscape of diatom diurnal rhythms, we unveil the function of a *Phaeodactylum tricornutum* bHLH-PAS protein, *Pt*bHLH1a, in the regulation of light-dependent diurnal rhythms. Peak expression of *PtbHLH1a* mRNA occurs toward the end of the light period and it adjusts to photoperiod changes. Ectopic over-expression of *Pt*bHLH1a results in lines showing a phase shift in diurnal cell fluorescence, compared to the wild-type cells, and with altered cell cycle progression and gene expression. Reduced oscillations in gene expression are also observed in overexpression lines compared to wild-type in continuous darkness, showing that the regulation of rhythmicity by *Pt*bHLH1a is not directly dependent on light inputs and cell division. *Pt*bHLH1a homologs are widespread in diatom genomes which may indicate a common function in many species. This study adds new elements to understand diatom biology and ecology and offers new perspectives to elucidate timekeeping mechanisms in marine organisms belonging to a major, but underinvestigated branch of the tree of life.

SIGNIFICANCE STATEMENT

Most organisms experience diurnal light-dark changes and show rhythms of basic biological processes such that they occur at optimal times of the day. The ocean harbours a huge diversity of organisms showing light-dependent rhythms, but their molecular foundations are still largely unknown. In this study, we discover a novel protein, *Pt*bHLH1a that regulates cell division, gene expression and the diurnal timing of these events in the marine diatom *Phaedoactylum tricornutum*. The identification of *PtbHLH1a*-like genes in many diatom species suggests a conserved function in diurnal rhythm regulation in the most species-rich group of algae in the ocean. This study unveils critical features of diatom biology and advances the field of marine rhythms and their environmental regulation.

INTRODUCTION

The Earth's rotation means that life evolved under a 24h diurnal cycle of alternate light and dark periods. Most living organisms have developed daily rhythms of many fundamental biological processes, ranging from physiology to behaviour, such that they occur at optimal times of the day (1) which can enhance fitness (2, 3). These rhythms are the product of the coordinated action of signals from endogenous timekeepers, together with environmental and metabolic inputs (4, 5). Robust diel rhythms in growth, cell cycle, gene expression, pigment synthesis, phototactic movements and bioluminescence have been also observed in a variety of phytoplanktonic organisms, including diatoms (6-13). Diatoms represent the most species-rich group of algae in the ocean and populate a wide range of aquatic environments (14, 15). These algae of the Stramenopile lineage show peculiar genomic, metabolic and cellular features, and are evolutionarily distant from the most studied model organisms in the field of biological rhythms (16-20). Diatoms have an impressive capacity to deal with environmental changes thanks to sophisticated acclimation mechanisms (21-25). Recent genome-wide analyses also showed that 25% of the diurnal transcriptome is influenced by light-dark cycles in the centric diatom *Thalasiossira pseudonana* (26). Moreover, detailed diurnal studies in the pennate diatom *Phaeodactylum tricornutum* highlighted a strict temporal separation of transcriptional gene networks (27, 28), as previously observed in other algae (29). Tight diurnal control of the *P. tricornutum* cell cycle has also been observed (27, 30, 31), with light onset triggering cell cycle progression through the transcriptional activation of the diatom-specific cyclin dsCYC2 by the blue light sensor Aureochrome-1a (30). Furthermore, the transcription factor bZIP14 has recently been identified as a diurnal activator of the tricarboxylic acid (TCA) cycle, a process restricted to the late afternoon in diatoms (32). Together, these studies illustrate complex regulation of diurnal cellular activities in *P. tricornutum*. However, the molecular mechanisms orchestrating diurnal processes are still unknown in diatoms and many other phytoplanktonic organisms. Notably, no orthologs of the circadian clock components discovered in bacteria, fungi, animals or plants have been found in the diatom genomes except for cryptochromes (33). Nonetheless, a number of proteins containing bHLH-PAS domains, which feature in genes involved in the regulation of rhythmic processes in animals (34), have been identified in diatom genomes (35).

In this work, we integrate transcriptomic, physiological and functional analyses to explore the regulatory landscape of *P. tricornutum* diurnal rhythms. We uncover the function of a bHLH-PAS protein (*Pt*bHLH1a) in the regulation of critical light-dependent rhythmic processes, such as cell cycle and diurnal transcription. Phylogenetic analyses reveal that

99 bHLH1a homologs are widely distributed in diatoms, thus we speculate a common function in
100 many diatom species. These results open the way to new exploration of diatom genomes in
101 search of their elusive molecular timekeepers.
102
RESULTS
Transcriptome profiling identifies potential regulators of diurnal rhythms in *P.*
tricornutum
To identify potential regulators of cellular rhythmicity in *P. tricornutum*, a publicly available
diurnal transcriptomic dataset (27) was analyzed. One hundred and four genes with robust diel
oscillating expression were selected, of which eight were photoreceptors (30, 33, 36, 37) and
66 were transcription factors (TFs) (35), which might be involved in the generation of light-
dependent rhythms. The remaining 30 genes selected were potential output genes implicated
in diel rhythmic processes (pigment synthesis, cell cycle regulation and photosynthesis)
(Table S1). The transcriptional dynamics of the selected genes were examined in a 16:8-h
light:dark (L:D) photocycle for 32h. Hierarchical clustering (HCL) analysis of the resulting
expression profiles revealed 4 clusters of co-expressed genes, termed A-D (Fig. 1A), with
peak expression at different times between dawn and dusk (Fig. 1B). Cluster A phased at
dawn, suggesting a transcriptional anticipation of the light onset (Fig. 1A-B). This cluster
comprised 18 genes including 14 TFs, mostly belonging to the Heat Shock Transcription
Factor family (HSF), two DNA repair enzymes CPD photolyases (*CPD2* and *CPD3*) and one
carotenoid synthesis enzyme (*PDS1*). Cluster B phased around 7h Zeitgeber Time (hours after
illumination, ZT) and encompassed 36 genes (Fig. 1A-B), including the ds*CYC2* gene
controlling the onset of cell division (30). Cluster B also contained 18 TFs, of which eight
were sigma factors putatively involved in the regulation of chloroplast transcription, three
genes implicated in photoprotection (*LHCX1*, *Vdl2* and *Zep1*) and the chlorophyll synthesis
POR1 gene. Such active transcription of genes involved in chloroplast activity during the light
period has been shown previously (21, 38). The blue light sensors *Aurochrome1b* and the
cryptochromes *CPF1* and *CryP-like* also belonged to cluster B and show a strong expression
following light onset, in accordance with previous observations (39-41). Cluster C phased
around ZT9 (Fig. 1A-B) and comprises 9 TFs and 10 metabolism-related genes, including
genes encoding photosynthetic apparatus. Finally, cluster D phased before dusk and included
23 TF genes including the TCA cycle regulator *bZIP14* (27, 32), likely contributing to
preparing cells for light to dark transition (Fig. 1A-B). Cluster D also contained the *CPF4* and

the Far-Red light sensing phytochrome (*DPH1*) whose peak expression at the end of the light

period has been observed previously (36, 39).

Together these results underline the existence of tight transcriptional programs phasing at

discrete moments of the day which potentially control the timing of cellular activities along

the diurnal cycle.

PtbHLH1a expression is adjusted in a photoperiod-dependent manner

Our analysis identified two TFs, *Pt*bHLH1a (Phatr3_J44962) belonging to cluster D and

*Pt*bHLH1b (Phatr3_J44963) belonging to cluster C, which each have a Per-ARNT-Sim (PAS)

domain in conjunction with a bHLH DNA-binding domain. Because bHLH-PAS proteins

have been shown to be involved in the regulation of rhythmic processes in animals (4, 34, 42),

the expression profiles of *PtbHLH1a* and *PtbHLH1b* were examined in *P. tricornutum* cells

growing under different photoperiods. *PtbHLH1a* expression peaked at ZT8 in the 12L:12D

photoperiod and at ZT12 in the 16L:8D photoperiod, 4 hours before the end of the light period

in both cases, then gradually decreased to below detection limits at ZT0 (Fig. 1C).

Transcription of *PtbHLH1b* appeared to start earlier than that of *PtbHLH1a*. In cells entrained

in 12L:12D cycles, *PtbHLH1b* expression peaked at ZT8, whereas it peaked between

ZT8/ZT12 in 16L:8D photoperiods (Fig. 1C). Thus, *PtbHLH1b* expression onset almost

coincided in the two photoperiods although transcription dramatically dropped after ZT8 in

12L:12D, while remained at maximum levels up to ZT12 in long days (Fig. 1C).

The robustness of *PtbHLH1a* and *PtbHLH1b* diurnal expression profiles was further

examined under stress conditions using recent transcriptome datasets from *P. tricornutum*

cells grown in 12L:12D cycles in iron replete and deplete conditions (28). Iron homeostasis is

diurnally regulated in phytoplankton (43) and it affects rhythmic processes such as cell cycle

progression and diurnal gene expression in *P. tricornutum* (28). Interestingly, *PtbHLH1a* and

PtbHLH1b expression profiles showed similar patterns in both control and iron depleted

conditions, with peaks of expression before dusk at ZT9 (Fig. S1), similar to our observations

(Fig. 1C).

Altogether these results demonstrate robust control of *PtbHLH1a* and *PtbHLH1b* diurnal

expression timing, which is adjusted in a photoperiod-dependent manner and unaffected by

iron depletion. The involvement of *Pt*bHLH1a in the regulation of diurnal light-dependent

rhythmic processes was hypothesized considering a possible role in dusk anticipation.

PtbHLH1a ectopic expression determines phase shifts in cellular rhythmicity

To determine *Pt*bHLH1a's function, cell lines were generated expressing HA-tagged bHLH1a under the regulation of the *Light harvesting complex protein family F2* promoter (*Lhcf2p*) (Fig. 2A), which activates transcription 3h after light onset (44). Gene expression analysis allowed the selection of three independent lines, hereafter named OE-1, OE-2 and OE-3, showing over-expression of the *PtbHLH1a* gene (Fig. 2B) and earlier expression timing compared to the wild type (Wt) strain (Fig. 4A and S2). Next, daily cellular rhythmicity was analyzed using the flow cytometer channel FL3 (excitation 488nm, emission 655-730 nm) that estimates chlorophyll a cellular content (8, 38), over the course of three days (Fig 2C). Cellular fluorescence displayed highly oscillating rhythms in 16L:8D grown cultures with a periodicity of approximately 24h (Fig. 2C, Table S2). Cell fluorescence in Wt cultures increased during daytime to peak around ZT13 (Fig. 2D and Table S2), and then started to decrease before night onset. This fall in fluorescence was concomitant with an increase in cell concentration ((38) and Fig. S4), likely reflecting chloroplast partitioning to daughter cells during cell division (7). Fluorescence progressively declined during the night period, reaching a trough in the early morning (Fig. 2C). Despite maintaining rhythmicity in the cellular fluorescence dynamics, OE-1 (Fig 2C), as well as OE-2 and OE-3 lines (Fig. 2D), displayed a remarkable phase shift of around 1-2 h in the maximum fluorescence timing compared to the Wt (Table S2). Cellular fluorescence phase responses were further investigated in resetting experiments. Wt and OE-1 cultures, that show the strongest phase shift phenotype, were grown in 16L:8D photoperiods, then transferred to 8L:16D and monitored for another 6 days. After the transfer to 8L:16D photocycles, the timing of maximum cell fluorescence in Wt cells was maintained for two days and then re-synchronized to the newly imposed photoperiod, peaking at ZT7.99±1.88 starting from the third day (Fig. 2E-F). In contrast, after 3 days of re-entrainment in 8L:16D, the OE-1 line showed a 3-hour phase delay (ZT 11.49±2.87) compared to Wt (Fig. 2E-F). Together, these results indicate that diatom cellular rhythmicity is entrained in a photoperiod-dependent manner and that *Pt*bHLH1a deregulation alters the capacity of cells to set diurnal phase pattern.

PtbHLH1a regulates diurnal cell cycle progression

The altered rhythm of fluorescence upon *PtbHLH1a* over-expression described above may reflect delayed or asynchronous cell division dynamics. To get further insights into the effect of *PtbHLH1a* overexpression on rhythmic processes, cell cycle dynamics in the WT and OE-1 lines were thoroughly analysed. Cell cultures were synchronized by 40h of dark treatment and

harvested on an hourly basis for 12h after re-illumination. At T0, total DNA content measurements showed comparable proportions of cells in G1 phase in all samples indicating effective synchronization of cell cultures (Fig. 3A). Starting after 3h of illumination, a progressive reduction of G1 cell number was observed in the Wt, with a minimum number reached after 10h of illumination. After 10h, the percentage of Wt cells in G1 increased with the emergence of daughter cells (Fig. 3A). Interestingly, compared to Wt, OE-1 cells showed a slower exit from the G1 phase, with the percentage G1 cells continuing to decrease over the entire 12h of illumination studied (Fig. 3A).

To further characterize the cell cycle deregulation caused by *PtbHLH1a* over-expression, the expression profiles of specific cell cycle phase marker genes (31) were analyzed in dark-synchronized Wt and OE-1 lines illuminated for 12h. The G1 phase gene markers *CDKA1* and *CDKD1* showed similar expression profiles in both lines until 4h from the onset of illumination (Fig. 3B). Starting from this time point, transcript levels of both genes were consistently higher in the OE-1 line compared to Wt except for at the end of the time course when they converged. This illustrates that the G1 phase duration of the two cell lines is different. Conversely, the G2/M marker *CYCB1* showed lower expression in OE-1 compared to the Wt between 4 and 8h after the onset of illumination (Fig. 3B). The expression of another G2/M phase marker, *CYCA/B1*, also resulted deregulated in OE-1, presenting reduced amplitude and peaking 2h later compared to Wt. Together, these results suggest that the deregulation of *PtbHLH1a* affects cell cycle, possibly by altering transition from G1 to S or G2/M phases.

PtbHLH1a regulates pace of diel gene expression

The effect of *Pt*bHLH1a de-regulation on gene expression was investigated since the expression of many *P. tricornutum* genes phase diurnally (27). To this end, Wt and OE-1 lines were grown in 16L:8D photocycles and sampled every 3 hours over 25 hours. For this analysis, genes with strong diurnal transcription oscillation were selected, including TFs (*bHLH1a*, *bHLH1b* and *bHLH3*) and rhythmic genes putatively involved in chlorophyll and carotenoid synthesis (*NADPH:protochlorophyllide oxidoreductase 2*, *Por2*, and *Violaxanthin de-epoxidase-related*, *Vdr*) (Fig. 1A, (27, 38)). Total *PtbHLH1a* transcript levels, including endogenous and transgenic mRNAs, were shown to be higher in OE-1 cells compared to the Wt, and the expression peaking at ZT7 in the OE-1 line and ZT10 in the Wt (Fig. 4A). A decrease of endogenous *PtbHLH1a* transcripts was observed in the OE-1 line compared to the Wt, possibly reflecting negative feed-back mechanism of *Pt*bHLH1a regulating its own

transcription (Fig. 4A). A similar pattern was also observed for the *PtbHLH1b* gene, suggesting that *PtbHLH1a* and *PtbHLH1b* transcription is controlled by the same regulatory circuit. In addition, the *bHLH3* gene showed earlier phases of expression in OE-1 compared to the Wt (Fig. 4A). Similar deregulations of *PtbHLH1a, PtbHLH1b PtbHLH3* were also observed in the OE-2 and OE-3 lines at ZT10 (Fig. S3). Besides TFs, the Chlorophyll biosynthesis gene *Por2* was also anticipated and the *Vdr* gene presented increased amplitude of expression in the OE-1 line compared to the Wt (Fig. 4A).

Altered gene expression observed in *Pt*bHLH1a overexpression cell lines could be the consequence of the deregulation of cell cycle progression (Fig. 3). To test this hypothesis, gene expression was analyzed in dark conditions, when the cell cycle is arrested ((31) and (Fig. S5)). Because information about transcription dynamics in this condition was limited, an initial survey of expression of the previously selected 104 *P. tricornutum* diurnal rhythmic genes (Fig. 1A) was performed in cells exposed to continuous dark for 30 hours. Comparative analysis of transcript profiles revealed that around 20% of the genes show persistent oscillation of expression in D:D, although in some cases they displayed reduced amplitudes and/or shifted phases of expression compared to the 16L:8D condition. In particular, 19 genes were identified which showed the highest amplitude of expression in both L:D and D:D (for details see Materials and Methods), consisting of 16 putative TFs and 4 pigment-related enzymes (Fig. S6). Among the analyzed transcripts, genes that were severely affected by the absence of light were also found, being strongly down-regulated or over-expressed when compared to the L:D condition (Fig. S7). The expression of some of these genes was further analyzed in constant darkness in Wt and *Pt*bHLH1a OE-1 cells for a period of 24h. In the Wt, the analyzed genes showed comparable transcript profiles in D:D and 16L:8D conditions (Fig. 4B, Fig. S8). Conversely, 10 out of 13 tested genes displayed reduced amplitudes and shifts in the phase of expression in OE-1 compared to Wt in D:D (Fig. 4B, Fig. S8). It is worth mentioning that two of the analysed genes, *HSF1d* and *bZIP5*, showed almost overlapping profiles in OE-1 and Wt lines (Fig. S8), excluding global deregulation of transcription by modulation of bHLH1a expression. Taken together, these results suggest that *Pt*bHLH1a contributes to define timing of diurnal gene expression and that its activity is independent of direct light inputs and cell division.

PtbHLH1a-like proteins are widely represented in the genome of marine algae

bHLH-PAS proteins were thought to be restricted to the animal (Opistokonta) lineage (45) until genome and transcriptome sequencing projects revealed bHLH-PAS family members in

diatoms (35) and other microalgae (46). Interestingly, when compared to animal bHLH-PAS, diatom proteins show peculiar features including a single predicted PAS domain (Fig. 5A), whereas animal bHLH-PAS proteins have two, and a N-ter extension that is absent in the animal counterparts. Available transcriptomic and genomic databases of marine algae and animals were searched for bHLH-PAS proteins and ≈90 novel bHLH-PAS proteins were discovered from Rhodophyta, Cryptophyta, Stramenopila, Alveolata and basal Opistokonta organisms (Table S3). With one exception, all the newly identified proteins showed a single predicted PAS domain, short C-ter extensions and N-ter regions of variable length, similar to the predicted structure of diatom bHLH-PAS proteins (Fig. 5A). Notably, the only bHLH-PAS possessing two PAS domains like the animal proteins was identified in *Galdieria sulphuraria* (Rhodophyta) and represents the first TF of this family identified in Archaeplastida. All the identified sequences, including selected bHLH-PAS from Opistokonta lineages, were used to perform a detailed phylogenetic analysis of the protein family using the bHLH and PAS domains. This analysis revealed at least three clades of algal bHLH-PAS proteins clearly separated from their Opistokonta counterparts (Fig. 5B). Interestingly, domain organization and branching positions of proteins from basal Opistokonta (*Monosiga brevicollis*) and microalgae (*Guillardia theta* (Cryptophyta) and *Nannochloropsis* (Stramenopila)) (Fig. 4B) support a possible common origin for this TF family, from an ancestor featuring single bHLH and PAS domains. However, the possible contribution of horizontal gene transfer and convergent evolution to the proliferation and diversification of this family cannot be excluded, and may be supported by the features of the *G. sulphuraria* bHLH-PAS protein, likely independently acquired by this alga. Based on our analysis, the majority of microalgal bHLH-PAS proteins fall into three separate clades: the first containing 9 TFs from diatoms and *Ectocarpus siliculosus*, the second comprising *Pt*bHLH1a together with 35 proteins from diatoms and Alveolata (Dinoflagellata), and the third comprising 41 proteins from Alveolata (Ciliophora and Dinoflagellata) and diatoms, including *Pt*bHLH1b (Fig. 5B).

Our results highlight diversification and widespread distribution of bHLH-PAS family members in different groups of algae. Moreover, the presence of bHLH1a-like genes in the genome of dinoflagellate and diatom species suggests that these proteins may share similar functions in microalgae. The similarities in diel transcript regulation and timing of expression between *Pt*bHLH1a and the centric diatom *Thalassiosira pseudonana* ortholog, *TpHLH1* (26), further reinforce this hypothesis.

DISCUSSION

The diurnal cycle is characterised by profound periodic light and temperature changes which have shaped the evolution of most ecosystems on Earth. In most organisms, biological rhythms are controlled by interconnected transcriptional-translational feed-back loops involving TFs and integrating signals from the environment (5). Although this regulatory framework is conserved among eukaryotes, the regulators responsible for the timing of events within biological rhythms seem to have emerged several times through evolution (47). Therefore, our current understanding of diurnal and circadian regulation, largely based on the study of terrestrial model organisms, is not always appropriate or relevant for evolutionarily distant marine organisms. In this study, we have shed light on the unknown regulators of diurnal patterns in diatoms, one of the most prominent phytoplanktonic groups in the Ocean. In agreement with previous studies (27, 28), we showed the existence of organized transcriptional programs defining cellular activities along the daily cycle in *P. tricornutum* and identified a number of TFs phasing at different times during the 24h cycle, as novel candidates for diatom diurnal regulation. By monitoring diurnal variations in chlorophyll fluorescence, robust regulation of diatom physiological rhythms that can be re-entrained to changing photoperiods was also unveiled. The ability to adjust the phase according to the photoperiod length constitutes one defining criterion of circadian clock-regulated mechanisms (48). Likewise, the strongly oscillating diel expression pattern of a *P. tricornutum* bHLH-PAS gene, *PtbHLH1a*, responded to photoperiod length with peak expression 4 hours before night onset in both 12h and 16h day photoperiods. The timing of *PtbHLH1a* expression is also preserved in cells under iron deficiency, in contrast to the expression of many other *P. tricornutum* genes observed previously, and despite the growth rate reductions caused by nutriment depletion (28). Functional characterization of *Pt*bHLH1a established its involvement in the regulation of *P. tricornutum* diurnal rhythms. Transgenic lines over-expressing *PtbHLH1a* using a promotor that is activated earlier in the light period than the endogenous gene maintained cellular rhythms of ∼24 h but show phase-alterations that are even more accentuated in re-entrainment experiments. This phenotype may reflect a reduced capacity of cells to synchronize to environmental light-dark cycles and adjust the phase to the new photoperiod. The participation of bHLH1a in the regulation of *P. tricornutum* cell cycle progression, reported in this study, could also explain the altered cellular fluorescence rhythmicity observed in the mutants. Altered cell division timing could derive from a delayed exit from the G1 phase in transgenic lines compared to Wt, as also supported by the altered

expression of the mitotic cyclins CYCB1 and CYCA/B1 in these lines (31). The *Pt*bHLH1a
gene could participate in gating cell divisions at night time, therefore maximizing the
energetic budget, as observed in several unicellular algae (7, 8, 30, 49). Interestingly, a similar
regulation of the cell cycle occurs in mammalian cells, where the circadian clock controls the
expression of G2 cycle-related genes to gate cell division at specific times of the day (50).

Besides cell cycle, *Pt*bHLH1a deregulation also affected diurnal rhythmicity of several
gene transcripts. This phenotype was uncoupled from cell cycle deregulation as it was
observed also in the absence of cell division, during darkness. Interestingly, the deregulation
of gene transcription was much more pronounced when analyzed in D:D compared to L:D
conditions. These results suggest on one hand that multiple regulatory inputs participate in the
regulation of diurnal rhythmic gene transcription (48), partially masking *Pt*bHLH1a
contribution to this process in cyclic environments, and, on the other hand, support *Pt*bHLH1a
involvement in the maintenance of rhythms in the absence of light inputs.

The evidence provided in this work support the hypothesis that *Pt*bHLH1a is one
component of an uncharacterized endogenous circadian clock in diatoms, either as part of a
central oscillator or as a mediator of clock inputs or outputs. With the exception of CRY (33,
39), orthologs of plant and animal circadian clock genes are absent in diatom genomes.
However, *Pt*bHLH1a contains bHLH and PAS protein domains that are also present in the
CLOCK and BMAL proteins, components of the mammalian central circadian oscillator (51,
52). Interestingly, previous studies showed that the *P. tricornutum* animal-like blue light
sensor Cpf1 can repress the transcriptional activity of these proteins in a heterologous
mammalian cell system (33), suggesting at least partial conservation in the regulatory
program generating rhythmicity in animals and diatoms. The downregulation of endogenous
PtbHLH1a and *PtbHLH1b* transcripts in the OE lines analysed in this work likely reflects a
negative feedback loop in the regulation of these genes, which is typical of circadian genetic
oscillators, and further supports a possible role for *PtbHLH1a* in diatom rhythm regulation.
Moreover, as observed for the clock components, we show that *Pt*bHLH1a contributes to set
the phase of output processes in cycling environments and to the maintenance of gene
expression rhythmicity in constant darkness. Some TFs characterized in this study and
showing altered expression patterns in *Pt*bHLH1a transgenic cells (*i.e., bHLH1b*, *bHLH3*,
bZIP5, *bZIP7*, *HSF1d*, *HSF1g*, *HSF3.3a*, *HSF4.7b*) represent direct or indirect targets of
*Pt*bHLH1a activity and possible additional components of the network participating in diel
rhythm regulation. *Pt*bHLH1a might act downstream of signal transduction cascades activated
by the diatom photoreceptors analysed in this study and elsewhere. The presence of a PAS

domain in *Pt*bHLH1a also suggests that this protein might have its own light-sensing ability
(53).

Although further analyses under prolonged free running conditions (for example
continuous darkness and continuous illumination) will be necessary to conclusively assess the
involvement of *Pt*bHLH1a in circadian regulation, this protein constitutes a promising entry
point for the characterization of diatom molecular timekeepers. Finally, the discovery of the
wide distribution of bHLH-PAS domain-containing proteins in diatoms, as well as in other
algae, has the potential to shed new light on the evolution of biological rhythms. bHLH-PAS
proteins might have independently acquired their function in rhythm regulation by convergent
evolution. However, the existence of this function in an ancient heterotrophic marine ancestor
that subsequently acquired plastids via endosymbiosis events (54) and prior to colonization of
land cannot be excluded. Regulators of cellular rhythmicity such as *Pt*bHLH1a may have
played a critical role for diatom prominence in marine ecosystems, by synchronizing cellular
activities in optimal temporal programs and maximizing diatoms' ability to anticipate and
adapt to cyclic environmental variations.

Biological rhythms are still poorly understood at molecular and mechanistic levels in
marine algae, despite their fundamental significance to these organisms' biology and ecology.
Further characterization of *Pt*bHLH1a homologs in diatoms and other algae is expected to
provide new insights into biological rhythms in marine organisms.

METHODS

Culture conditions

Wild-type *P. tricornutum* (*Pt*1 8.6; CCMP2561) cells and transgenic lines were grown at
18°C with shaking at 100 rpm in F/2 medium (55) without silica and illuminated at 40 µmol
photons $m^{-2} s^{-1}$ of white light (Philips TL-D De Luxe Pro 950). Detailed information is in SI.

Cell cycle analysis

Cells were synchronized in the G1 phase by 40h of darkness. After re-illumination, samples
were collected every hour for 12h. Details are in SI.

RNA extraction and gene expression analyses

Total RNA was extracted and qRT-PCR performed as described in (36). Codeset information and raw nCounter data are available from the GEO database (Series GSE112268). Detailed information is in SI.

Selection of rhythmic transcripts and clustering analysis

To select genes with rhythmic expression in the light-dark cycle, we used microarray data from (27). To select rhythmic transcripts in D:D, standard deviation from the average expression were calculated and used as selective criteria. Detailed information is in SI.

Generation of the *PtbHLH1a* overexpressing lines

Diatom transgenic lines were obtained by co-transformation of the pDEST-C-HA-*PtbHLH1a* plasmid together with Nourseothricin resistance plasmid. Details are described in SI.

Data mining, protein sequence and phylogenetic analysis

Detailed information about data mining, protein sequence and phylogenetic analysis is provided in SI.

ACKNOWLEDGEMENTS

We thank M. Jaubert and L. De Veylder for critical suggestions, A. Manzotti for help with characterization of OE lines, P. Oliveri for support with nCounter technology, D. Petroutsos and G. Finazzi for support in monitoring cell physiology. This work was funded by the HFSP research grant (#RGY0082/2010), EMBRIC, EMBRC-FR and a grant from the Gordon and Betty Moore Foundation (GBMF 4966) to A.F.

REFERENCES

1. Pittendrigh CS (1993) Temporal Organization: Reflections of a Darwinian Clock-Watcher. *Annual Review of Physiology* 55(1):17-54.
2. Dodd AN, et al. (2005) Plant circadian clocks increase photosynthesis, growth, survival, and competitive advantage. *Science* 309(5734):630-633.
3. Woelfle MA, Ouyang Y, Phanvijhitsiri K, & Johnson CH (2004) The adaptive value of circadian clocks: an experimental assessment in cyanobacteria. *Curr Biol* 14(16):1481-1486.
4. Dunlap JC (1999) Molecular bases for circadian clocks. *Cell* 96(2):271-290.
5. Bell-Pedersen D, et al. (2005) Circadian rhythms from multiple oscillators: lessons from diverse organisms. *Nat Rev Genet* 6(7):544-556.
6. Chisholm SW (1981) Temporal Patterns of Cell-Division in Unicellular Algae. Can B Fish Aquat Sci (210):150-181.
7. Moulager M, et al. (2007) Light-dependent regulation of cell division in Ostreococcus: evidence for a major transcriptional input. *Plant physiology* 144(3):1360-1369.
8. Ragni M & D'Alcala MR (2007) Circadian variability in the photobiology of Phaeodactylum tricornutum: pigment content. *J Plankton Res* 29(2):141-156.
9. Naylor E (2010) Chronobiology of marine organisms (Cambridge University Press, Cambridge; New York) pp x, 242 p.
10. Tessmar-Raible K, Raible F, & Arboleda E (2011) Another place, another timer: Marine species and the rhythms of life. *Bioessays* 33(3):165-172.
11. Ottesen EA, et al. (2013) Pattern and synchrony of gene expression among sympatric marine microbial populations. *P Natl Acad Sci USA* 110(6):E488-E497.
12. Brierley AS (2014) Diel vertical migration. *Current Biology* 24(22):R1074-R1076.
13. Hafker NS, et al. (2017) Circadian Clock Involvement in Zooplankton Diel Vertical Migration. *Current Biology* 27(14):2194-2201.
14. Field CB, Behrenfeld MJ, Randerson JT, & Falkowski P (1998) Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* 281(5374):237-240.
15. Malviya S, et al. (2016) Insights into global diatom distribution and diversity in the world's ocean. *Proc Natl Acad Sci U S A* 113(11):E1516-1525.
16. Baldauf SL (2003) The Deep Roots of Eukaryotes. *Science* 300(5626):1703-1706.
17. Bowler C, et al. (2008) The Phaeodactylum genome reveals the evolutionary history of diatom genomes. *Nature* 456(7219):239-244.
18. Moustafa A, et al. (2009) Genomic footprints of a cryptic plastid endosymbiosis in diatoms. *Science* 324(5935):1724-1726.
19. Flori S, et al. (2017) Plastid thylakoid architecture optimizes photosynthesis in diatoms. *Nat Commun* 8:15885.
20. Raible F & Falciatore A (2014) It's about time: Rhythms as a new dimension of molecular marine research. *Mar Genomics* 14:1-2.
21. Bailleul B, et al. (2010) An atypical member of the light-harvesting complex stress-related protein family modulates diatom responses to light. *P Natl Acad Sci USA* 107(42):18214-18219.
22. Allen AE, et al. (2011) Evolution and metabolic significance of the urea cycle in photosynthetic diatoms. *Nature* 473:203.
23. Goss R & Lepetit B (2015) Biodiversity of NPQ. *Journal of Plant Physiology* 172:13-32.
24. Morrissey J, et al. (2015) A novel protein, ubiquitous in marine phytoplankton, concentrates iron at the cell surface and facilitates uptake. *Curr Biol* 25(3):364-371.
25. McQuaid JB, et al. (2018) Carbonate-sensitive phytotransferrin controls high-affinity iron uptake in diatoms. *Nature* 555(7697):534-537.
26. Ashworth J, et al. (2013) Genome-wide diel growth state transitions in the diatom Thalassiosira pseudonana. *P Natl Acad Sci USA* 110(18):7518-7523.
27. Chauton MS, Winge P, Brembu T, Vadstein O, & Bones AM (2013) Gene regulation of carbon fixation, storage, and utilization in the diatom Phaeodactylum tricornutum acclimated to light/dark cycles. *Plant physiology* 161(2):1034-1048.

28. Smith SR, et al. (2016) Transcriptional Orchestration of the Global Cellular Response of a
Model Pennate Diatom to Diel Light Cycling under Iron Limitation. *PLoS Genet*
12(12):e1006490.
29. Zones JM, Blaby IK, Merchant SS, & Umen JG (2015) High-Resolution Profiling of a
Synchronized Diurnal Transcriptome from Chlamydomonas reinhardtii Reveals Continuous
Cell and Metabolic Differentiation. *Plant Cell* 27(10):2743-2769.
30. Huysman MJ, et al. (2013) AUREOCHROME1a-mediated induction of the diatom-specific
cyclin dsCYC2 controls the onset of cell division in diatoms (Phaeodactylum tricornutum).
Plant Cell 25(1):215-228.
31. Huysman MJ, et al. (2010) Genome-wide analysis of the diatom cell cycle unveils a novel
type of cyclins involved in environmental signaling. *Genome Biol* 11(2):R17.
32. Matthijs M, et al. (2017) The transcription factor bZIP14 regulates the TCA cycle in the
diatom Phaeodactylum tricornutum. *The EMBO journal* 36(11):1559-1576.
33. Coesel S, et al. (2009) Diatom PtCPF1 is a new cryptochrome/photolyase family member with
DNA repair and transcription regulation activity. *EMBO Rep* 10(6):655-661.
34. Kewley RJ, Whitelaw ML, & Chapman-Smith A (2004) The mammalian basic helix-loop-
helix/PAS family of transcriptional regulators. *Int J Biochem Cell Biol* 36(2):189-204.
35. Rayko E, Maumus F, Maheswari U, Jabbari K, & Bowler C (2010) Transcription factor
families inferred from genome sequences of photosynthetic stramenopiles. *New Phytol*
188(1):52-66.
36. Fortunato AE, et al. (2016) Diatom Phytochromes Reveal the Existence of Far-Red-Light-
Based Sensing in the Ocean. *Plant Cell* 28(3):616-628.
37. Jaubert M, Bouly JP, Ribera d'Alcala M, & Falciatore A (2017) Light sensing and responses in
marine microalgae. *Curr Opin Plant Biol* 37:70-77.
38. Hunsperger HM, Ford CJ, Miller JS, & Cattolico RA (2016) Differential Regulation of
Duplicate Light-Dependent Protochlorophyllide Oxidoreductases in the Diatom
Phaeodactylum tricornutum. *PLoS One* 11(7):e0158614.
39. Fortunato AE, Annunziata R, Jaubert M, Bouly JP, & Falciatore A (2015) Dealing with light:
the widespread and multitasking cryptochrome/photolyase family in photosynthetic
organisms. *J Plant Physiol* 172:42-54.
40. Oliveri P, et al. (2014) The Cryptochrome/Photolyase Family in aquatic organisms. *Mar
Genomics* 14:23-37.
41. Banerjee A, et al. (2016) Allosteric communication between DNA-binding and light-
responsive domains of diatom class I aureochromes. *Nucleic Acids Res* 44(12):5957-5970.
42. Young MW & Kay SA (2001) Time zones: a comparative genetics of circadian clocks. *Nat
Rev Genet* 2(9):702-715.
43. Botebol H, et al. (2015) Central role for ferritin in the day/night regulation of iron homeostasis
in marine phytoplankton. *Proc Natl Acad Sci U S A* 112(47):14652-14657.
44. Russo MT, Annunziata R, Sanges R, Ferrante MI, & Falciatore A (2015) The upstream
regulatory sequence of the light harvesting complex Lhcf2 gene of the marine diatom
Phaeodactylum tricornutum enhances transcription in an orientation- and distance-independent
fashion. *Mar Genomics* 24 Pt 1:69-79.
45. Yan J, Ma Z, Xu X, & Guo AY (2014) Evolution, functional divergence and conserved exon-
intron structure of bHLH/PAS gene family. *Mol Genet Genomics* 289(1):25-36.
46. Thiriet-Rupert S, et al. (2016) Transcription factors in microalgae: genome-wide prediction
and comparative analysis. *BMC Genomics* 17:282.
47. Bhadra U, Thakkar N, Das P, & Bhadra MP (2017) Evolution of circadian rhythms: from
bacteria to human. *Sleep Med* 35:49-61.
48. Merrow M, et al. (2006) Entrainment of the Neurospora circadian clock. *Chronobiol Int* 23(1-
2):71-80.
49. Noordally ZB & Millar AJ (2015) Clocks in algae. *Biochemistry* 54(2):171-183.
50. Matsuo T, et al. (2003) Control mechanism of the circadian clock for timing of cell division in
vivo. *Science* 302(5643):255-259.
51. Bunger MK, et al. (2000) Mop3 is an essential component of the master circadian pacemaker
in mammals. *Cell* 103(7):1009-1017.

52. Gekakis N, et al. (1998) Role of the CLOCK protein in the mammalian circadian mechanism.
Science 280(5369):1564-1569.
53. Taylor BL & Zhulin IB (1999) PAS domains: internal sensors of oxygen, redox potential, and
light. Microbiol Mol Biol Rev 63(2):479-506.
54. Archibald JM (2015) Endosymbiosis and Eukaryotic Cell Evolution. *Current biology*
25(19):R911-921.
55. Guillard RRL (1975) Culture of Phytoplankton for Feeding Marine Invertebrates. In Culture
of Marine Invertebrate Animals, D.R. Smith and M.H. Chanley, eds. Springer US.
542
543
544
545

     FIGURE LEGENDS

Fig. 1. Diurnal expression analysis of selected rhythmic *P. tricornutum* genes. A)

nCounter expression analysis of 104 selected genes in cells grown under a 16L:8D

photoperiod and sampled every 4 hours for 32 hours. Four major groups of co-regulated genes

(A-D) are shown based on hierarchical clustering. Expression values were normalized using

PtRPS, *PtTBP* and *PtACTIN12* as reference genes and represent the average of three

biological replicates. The *PtbHLH1a* and *PtbHLH1b* gene expression profiles are indicated

with arrowheads. B) Polar plot and table showing the average phases of expression (intended

as expression peaks during the analyzed period) of the four gene clusters, calculated using the

MFourfit averaged method. Plot petal length is proportional to the standard deviation of

phases over biological replicates. In the table, N: number of genes within each cluster; Av.

Phase: average periods and phases of expression in hours. C) Diurnal expression profiles of

PtbHLH1a and *PtbHLH1b* in Wt cells grown under 12L:12D (blue line) and 16L:8D (red

line) photocycles analysed by qRT-PCR. Expression values were normalized using *PtRPS* and

PtTBP as reference genes and represent the average of three biological replicates ± s.e.m

(standard error of the mean; black bars).

For each gene, the expression value is relative to its maximum expression (maximum

expression=1). Light and dark periods are represented by white and grey regions respectively.

Fig. 2. *PtbHLH1a* over-expression determines phase shifts in cellular rhythmicity.

A) Schematic representation of the *Lhcf2p:bHLH1a-3xHA:Lhcf1t* construct used to generate

*Pt*bHLH1a over-expressing lines. *Lhcf2p*: Light Harvesting Complex F2 promoter; 3HA:

triple hemagglutinin tag; *Lhcf1t*: *Lhcf1* terminator. B) Quantification of total *PtbHLH1a*

transcripts in OE-lines and Wt grown in 16L:8D photocycles and sampled at the ZT10. qRT-

PCR expression values were normalized using *PtRPS* and *PtTBP* as reference genes and

represent the average of 3 biological replicates (n=3) ± s.e.m (black bars). C) Diurnal

oscillation of chlorophyll fluorescence (FL-3 parameter) in Wt and OE lines entrained under

16L:8D over three days. D) Diurnal phase time calculation of the FL-3 value in Wt and OE-1,

OE-2 and OE-3 lines. Values represent the average of at least 4 biological replicates (n ≤ 4)

from at least 2 independent experiments ± s.e.m. E) Phase re-entrainment analysis of

fluorescence rhythms after photoperiod change from 16L:8D to 8L:16D in Wt (yellow) and

OE-1 (red). The FL-3 parameter was monitored in cultures grown in 16L:8D photocycles and

then transferred to 8L:16D for 6 days. Values represent the average of three biological

replicates ± s.e.m. F) Bar plot representation of the re-entrained FL-3 phases in Wt and OE-1

cultures after three 8L:16D photocycles. Values represent the average of 3 biological

replicates ± s.e.m. (black bars).

*P<0.05, **P<0.01, ***P<0.001, t-test.

Fig. 3. *PtbHLH1a* over-expression affects cell cycle progression. A) Cell cycle progression

dynamics of Wt (yellow) and OE-1 (red) lines shown as the proportion of cells in the G1

phase measured by flow cytometry each hour over 12h of illumination following dark

synchronization. Results are representative of three biological replicates ± s.e.m (black bars);

t-test significance is indicated by *: P<0.05, **: P<0.01, t-test. B) qRT-PCR expression

profiles of G1 (*PtCDKA1*, *PtCDKD1*) and G2/M phase marker genes (*PtCYCB1*,

PtCYCA/B1) in synchronized Wt and OE-1 cell lines over 12h of illumination. Expression

values were normalized using *PtRPS* and *PtTBP* as reference genes and represent the average

of two independent biological replicates. For each gene, the expression value is relative to its

maximum expression (maximum expression=1).

Fig. 4 *PtbHLH1a* over-expression alters rhythmic diel gene expression. A) qRT-PCR

diurnal expression analysis of *PtbHLH1a, PtbHLH1b*, *PtbHLH3*, *PtPor2* and *PtVdr*

transcripts in 16L:8D entrained Wt (yellow) and OE-1 (red) cultures. *bHLH1a-total*

represents the sum expression of the *bHLH1a* endogenous and the transgene transcripts;

PtbHLH1a-endogenous refers to the endogenous gene only. Expression values represent the

average of 2 biological replicates ± s.e.m. (black bars). B) nCounter expression analysis of

PtbHLH1a, PtbHLH1b, *PtbHLH3*, *PtPor2* and *PtVdr* transcripts during 24 hours of dark free-

running period in the Wt (yellow) and OE-1 (red) lines. Cells were previously entrained in

16L:8D cycles. Data represent the average of 3 biological replicates ± s.e.m (black bars).

Grey rectangles represent dark periods. Expression values were normalized using *PtRPS* and

PtTBP as reference genes. For each gene, the expression value is relative to its maximum

expression (maximum expression=1).

Fig. 5. bHLH-PAS protein family structure and phylogeny. A) Schematic representation

of bHLH-PAS protein domain architecture across Eukaryotes. Segmented line indicates

possible absence of the second PAS domain in some Opistokonta species. The grey patterns

before the bHLH domain and after the PAS domain represent the variations in N-ter and C-ter

length in different organisms. B) Maximum Likelihood (ML) phylogenetic tree of the bHLH-

PAS family. The Opistokonta clade is used as the outgroup. Numbers refer to bootstrap values

of the basal nodes using ML (RAxML, 1000 bootstraps) and Bayesian Inference (MrBayes,

2.5M generations, 25% burn-in). The asterisk, the arrows and the square indicate the position

of *Monosiga brevicollis* bHLH-PAS, *P. tricornutum* HLH1a and bHLH1b, and *Thalassiosira*

pseudonana bHLH1, respectively. The colour code indicates the lineage corresponding to

each bHLH-PAS protein, shown in Fig 5A.

Figure 1 - Annunziata et al.

623 **Figure 2 - Annunziata et al.**

626 **Figure 3 - Annunziata et al.**

629   **Figure 4 - Annunziata et al.**

631
632

Figure 5 - Annunziata et al.

SUPPORTING INFORMATION

Culture conditions

For experiments in different photoperiods, cultures were pre-adapted to the different L:D cycles for 2 weeks before starting the experiment. For experiments in continuous darkness cells were pre-adapted in 16L:8D photocycles for 2 weeks, then transferred to D:D at the start of the experiment. Growth measurements were performed using a MACSQuant Analyser flow cytometer (Miltenyi Biotec, Germany) by counting the cells based on the R1-A (630nm excitation, 670-700nm emission) versus the R1-H parameters. Phase rhythmicity assays were carried out by measuring the Chlorophyll fluorescence using the flow cytometer FL-3 parameter (488 nm excitation, 670-700 nm emission). For the re-entrainment experiment, cultures were initially entrained under 16L:8D photocycles at 40 µmol $m^{-2} s^{-1}$ of white light, then transferred to 8L:16D photocycles at 80 µmol $m^{-2} s^{-1}$ of white light for 6 days. All phase time and period calculations were performed using the MFourfit curve-fitting method using the Biodare2 tool (biodare2.ed.ac.uk, (1)).

Cell cycle analysis

For cell cycle analysis cells were pelleted by centrifugation (4000 rpm, 15 minutes, 4°C), fixed in 70% EtOH and stored in the dark at 4°C until processing. Fixed cells were then washed three times with 1×PBS, stained with 4',6-diamidino-2-phenylindole (at a final concentration of 1 ng/ml) on ice for 30', then washed and resuspended in 1xPBS. After staining, samples were immediately analyzed with a MACsQuant Analyser flow cytometer (Miltenyi Biotec, Germany). For each sample, 30,000 cells were analysed and G1 and G2 proportions were inferred by calculating the 2c and 4c peak areas at 450 nm (V1-A channel) using the R software. A peak calling method was applied to the resulting histogram, based on a 1[st] derivative approach (2). The locations of G1 and G2 peaks were first determined using G1 and G2 reference samples and then used to identify G1 and G2 cells in the experimental samples. The area under each peak was used as a proxy for the proportion of cells in each population.

RNA extraction and gene expression analyses

For qRT-PCR analysis *PtRPS* and *PtTBP* were used as reference genes. Each independent replica of the qRT-PCR data was normalized against the maximum expression value of each gene (*i.e.,* gene expression range lies between 0 and 1 across the time series). Average expression and standard error was then calculated and plotted. The full list of oligonucleotides used in this work can be found in Table S4. For the nCounter analysis, gene specific probes (Table S1) were designed and screened against the *P. tricornutum* annotated transcript database (JGI, genome version 2, Phatr2) for potential cross-hybridization. Total RNA extracts (100 ng) from three biological replicates were used for hybridization. Transcript levels were measured using the nCounter analysis system (Nanostring Technologies) at the UCL Nanostring Facility (London, UK) and at the Institut Curie technical platform (Paris, France) as previously described (3). Expression values were first normalized against the internal spike-in controls, then against the geometric mean of the 3 reference genes *PtRPS*, *PtTBP* and *PtACTIN12*.

Selection of rhythmic transcripts and clustering analysis

For the selection of genes with rhythmic expression in the light-dark cycle, we used microarray data from (4). First, we identified all the genes belonging to the TFs, photoreceptors, cell cycle and metabolism-related categories (pigment synthesis and photosynthesis). Then, transcripts were ranked based on a defined t-value for each time point (mean gene expression of the replica/(1+s.d.)) and those showing t-value >+0.7 or <-0.7 across the time series, were retained. The nCounter expression data was normalized against the maximum expression value of each gene, in a similar way to qRT-PCR expression data. This normalization was applied to the 3 replicas independently and for each condition (L:D and D:D) with the average expression value used for the clustering analysis. Hierarchical clustering analysis was performed with MeV 4.9 (5) using Pearson correlation. Peak analysis was performed using the MFourfit curve-fitting method defining average expression phases for each cluster (biodare2.ed.ac.uk, (1)).

For the selection of rhythmic transcripts in D:D, expression values were normalized using as reference genes *PtRPS, PtTBP* and *PtACTIN12* and the genes with the highest values of standard deviation from the average expression (M value) over the two time courses (16L:8D and D:D) were selected. A threshold equal to 1 was set using the published *P. tricornutum* diurnal microarray dataset (4) as background. Gene expression profiles were further empirically examined and false positives eliminated.

Generation of the *PtbHLH1a* overexpressing lines

701
Transformed cells were tested for the presence of the transgene by PCR and qRT-PCR
analysis (see Table S4 for oligonucleotide sequences). The full length *PtbHLH1a* coding
sequence was obtained by PCR amplification with the specific oligonucleotides *Pt*bHLH1a-
DraI-Fw and *Pt*bHLH1a-XhoI-Rv on cDNA template using the Phusion high fidelity DNA
polymerase (Thermo Fisher, USA). The PCR fragment was inserted into the pENTR1A
vector (Invitrogen, USA) using the DraI/XhoI restriction sites, and recombined with the
pDEST-C-HA vector (6).

Data mining, protein sequence and phylogenetic analysis

The *P. tricornutum* bHLH1a (Phatr3_J44962) protein sequence was used as the query for
blastP analyses on the JGI, NCBI and MMETSP public database (7). Searches of the pfam
database for proteins possessing both the HLH and PAS domains were also performed. The
retrieved sequences were analyzed using the batch search tool on the CDD (Conserved
Domain Database) NCBI server to retrieve proteins presenting at least one HLH and one PAS
domain only. We identified 100 HLH-PAS proteins from 71 marine algal species which were
aligned using MAFFT (8), along with 22 HLH-PAS proteins from relevant metazoan (*Homo
sapiens*, *Mus musculus* and *Drosophila melanogaster*) and unicellular Opistokonta (*Monosiga
brevicollis* and *Capsaspora owczarzaki*). Preliminary phylogenies were produced with MEGA
7 (9) to eliminate ambiguously aligned sequences, refining the alignment to 107 sequences
and a final length of 198 aa (<5% gap per position). The best aminoacidic model to fit the
data was estimated with ProtTest 3.4.2 (10). Phylogenetic analyses were performed with
RAxML (1000 bootstraps) and MrBayes 3.2.6 (2.5 million generations, 2 runs, 25% burn-in)
on the CIPRESS gateway (11). The final tree was edited in FigureTree 1.4
(http://tree.bio.ed.ac.uk/software/Figure tree/). GenBank accession codes of the genes utilized
in the bHLH-PAS phylogenetic analysis are reported in Table S3.

Figure S1 – Annunziata et al.

Fig. S1. Diel expression patterns of *PtbHLH-PAS* genes under Fe-depletion conditions in

12L:12D photoperiods. Diel expression patterns of *PtbHLH1a* and *bHLH1b* in normal (400

pM Fe') and iron depletion conditions (40 and 20 pM Fe') were obtained using transcriptome

data extracted from (36). Light and dark periods are represented by white and black

rectangles. Expression values are given relative to the maximum expression for each gene,

where '1' represents the highest expression value of the time series.

Figure S2 – Annunziata et al.
742
743

744
745

Fig. S2. The *Lhcf2p:bHLH1a-3xHA:Lhcf1t* construct drives over-expression and
anticipation of *PtbHLH1a*. Quantification of total *PtbHLH1a* transcripts in two independent
*Pt*bHLH1a over-expressing lines (OE-1, OE-2) compared to the wild type (Wt) strain. Cells
were grown in 16L:8D photocycles and sampled at the ZT7 and ZT12 time points. qRT-PCR
expression values were normalized using *PtRPS* and *PtTBP* as reference genes and represent
the average of two biological replicates (n=2) ± s.e.m (black bars).
752

Figure S3 – Annunziata et al.

Fig. S3. *PtbHLH1a* over-expression alters rhythmic diel gene expression in three independent lines. Quantitative gene expression analysis by qRT-PCR of *PtbHLH1a-endogenous, PtbHLH1b* and *PtbHLH3*, transcripts in 16L:8D entrained Wt, OE-1, OE-2 and OE-3 cultures. Samples were harvested at ZT10. *PtbHLH1a-endo* refers to the transcript levels of the endogenous gene only. Expression values represent the average of 3 biological replicates ± s.e.m. (black bars) and were normalized using *PtRPS* and *PtTBP* as reference genes. For each gene, expression values are represented as relative to its maximum expression corresponding in the graph to '1'. *P<0.05, **P<0.01, ***P<0.001, t-test.

Figure S4 – Annunziata et al.

Fig. S4. Diurnal cell growth dynamics in *P. tricornutum*. Diel cell number measurements in

wild type cultures grown under 16L:8D photoperiods. Values represent the mean counts of

three independent biological replicates ± s.e.m. (black bars).

Figure S5 – Annunziata et al.

Fig. S5. Cell division is arrested in continuous dark conditions in *P. tricornutum*. Diurnal
cell growth measurements of the wild type cultures entrained in 16L:8D and then transferred
to continuous darkness condition (D:D). Values represent the mean counts of three
independent experiments ± s.e.m. (black bars).

Figure S6 – Annunziata et al.

Fig. S6. nCounter expression analysis of genes maintaining rhythmic expression in D:D conditions and 16L:8D condition in Wt cells. Data represent the average expression of biological triplicates ±SD and are normalized using the *PtRPS*, *PtTBP* and *PtACTIN12* reference genes. For each gene, the expression value is relative to its maximum expression corresponding in the graph to '1'. Results for cells grown in 16L:8D cycle are shown in orange (L:D); Results for cells in constant darkness (following 16L:8D adaptation) are shown in grey (D:D).

 Figure S7 – Annunziata et al.

Fig. S7. nCounter expression analysis of selected genes with altered rhythmic expression
in Wt cells in D:D conditions compared to 16L:8D condition. Expression values represent
the average of three biological triplicates ±SD and are normalized using the *PtRPS*, *PtTBP*
and *PtACTIN12* reference genes. For each gene, the expression value is relative to its
maximum expression corresponding in the graph to '1'. Results for cells grown in 16L:8D
cycle are shown in orange (L:D); Results for cells in constant darkness (following 16L:8D
adaptation) are shown in grey (D:D).

Figure S8 – Annunziata et al.

Fig. S8. nCounter analysis of selected rhythmic gene expression profiles in continuous darkness in Wt and *Pt*bHLH1a OE-1 cells. Cells were entrained in 16L:8D cycles, then transferred to D:D and collected every 3 hours for 24h. Expression values represent the average of three biological replicates (n=3) ± s.e.m (black bars) and have been normalized using *PtRPS* and *PtTBP*.

 Table S1: Accession codes and sequence probes used in the nCounter analysis.

Category	Gene Name	NCBI Accession	Phatr3 Accession	nCounter Probe Target Sequence
Reference Gene	Actin12	XM_002182185.1	Phatr3_J29136	ATGTAAGCCTGCAGCCACTGAGGACTTGTGCTCGTAACCCTGATTTCGATATTCCAAAATGCGGCTCACTAAAAAGACATCGTAGTCCAGTGCCAGTCCC
Photoreceptor	Aureo1b	#N/A	Phatr3_J49458	TTATTGCCGAGTGCTGTACCTCGAGTCCGTTGTTTGAAGAAATGGACGGCATAGATCAAACGAAAGGGGCCAATCTGGAACGTGCCGATTTTTCATTGAT
Photoreceptor	Aureo2	XM_002183279.1	Phatr3_J15468	AGACGTTACTGTCACTCTCATGAACTACACTGCCGATGGTACACCATTTTGGAACAAGCTCTTTATTGCCGCATTGCGTGACGCGCAAAATAACATTGTC
TF	bHLH1a_PAS	XM_002179032.1	Phatr3_J44962	TACTTTGGTCAATATCAAAGTCAGTTTGGTACGAACAGCTCAGCATAGCCCTCGGTTTTTCAATGTGGCGTTGGTGCCATCAGACGATGCAGCGAAGCTG
TF	bHLH1b_PAS	XM_002178812.1	Phatr3_J44963	AGGAATTGACTACCGTGCTGTTTTCAATCATTGTCCATATGCCATGGGGGTGGCTTCATTGGACGGAAGAATACTGGTCTGCAACAGCTCTTTCGAATCT
TF	bHLH3	XM_002176560.1	Phatr3_J42586	ATCACTAGCGACACATTCAGATTCATCTGCGGGAGAAGGACGGTCTATTCCGTCAGTAGTTGGTACAAGTGTTAGCGGTTCAAGTCGCTCTGGGAACTGC
TF	bHLH5	XM_002177858.1	Phatr3_J43365	ACCCTCTTGCAGCGCAATCATTCGCTCCCGCATACGACTATGGGTCGATTTCTCAATGAACGCCATTCCTTTTCCTCAGACCTTTCGCTCTCGCTGGCGT
Cell cycle	BUB1/MAD3	XM_002178639.1	Phatr3_J10954	CGTGTGCTGCATGTATGCGGACAAAACCGATCGTCCACTGGAAGTTTTCCAGCATCTACATCAGCAAAGGATCGGGAGCGATATTGCCGTGTTCTGGATG
TF	bZIP10	XM_002177776.1	Phatr3_J43744	AAGCTGCTCGGGAATCGCGTCGGCGGAAGAAGGTTATGATTGAAGAACTTCAACGCAGCGTGATTTTCTTCTCGCGTGCCAACGGAACCCTCAAACAACA
TF	bZIP13	XM_002181635.1	Phatr3_J47278	CGCGGCGGCTTCGCGTGCTAAGATTCGATGCCGCATCACCGAACTTGAAACCGAGGTTAGTGGTTGGAAGGATAAATACACACAAGCAATGGAACGCTTG
TF	bZIP14	XM_002179477.1	Phatr3_EG02108	ATGCAGAGCTTGGAGGGTCGAGCTACGGATCTCAAGCATGAGCAAATTCGTTTAAAACAGATCATTAACGAAAAGAACACGGCGAACATTCTGGTGGGGC
TF	bZIP16	XM_002181462.1	Phatr3_J47279	CTTTTGGAAGCAGTCACAAATCCGCAAACATGAACAGTTCCAACGCCCAGGAAGTCGCTCGTCAACTGAACGAAGATCCTGCGCTCGCTCGGAAGCGTCG
TF	bZIP18	XM_002176556.1	Phatr3_J42577	ACTCTTCTATAGCGGAATCCGAGGCCGATTTCAAAACTATAGCTCAGGCTGCTGTTTCGAATCTGATCATGTCGGCTGGAACTACCAAGGTTGAATCTGG
TF	bZIP19	XM_002183101.1	Phatr3_J48701	CCTGCACCTTCTCAAATCAGCTGTGCAACGAGTGCAACAAGAAGAAATATTGGCTGCCGAGGCATTACCTACACCTTTTATAGGCGTTTTGCGATTTGGC
TF	bZIP24	XM_002184633.1	Phatr3_J49887	TTGTCACGGGGCCAACAGACTATCTGGAAAGATCACTTCTTGCCCAAAGAGAAGCGGAAAGAACTCGTTCGGCTTTGTCAAGCATCGCGCTTCGGGGGAA
TF	bZIP25	XM_002185584.1	Phatr3_J46647	TCGTCGTTCACGATGGCATCTGTACCAACTCACCTAGTAGGAAATACTGTCATCACTGATACCAGTTATCAGCCAGTTTCGTCGCTATTACCTGCAGCGG
TF	bZIP26	XM_002182641.1	Phatr3_EG02494	GATTCTTTGACGGACTCAAGGAGCGACAACTTGTCATCGACATCTCAGGAGCTTTCGGTGCGACAGGAAAAGGTCGAAGCCGCTTTGAAATCGAAACCTC
TF	bZIP5_PAS	XM_002179145.1	Phatr3_J45142	CCCTTGTTTCCGAACAGCGCATTCGATCCGCAAAACAACGCTAGCAACAACGCACAATCACAGTCGCAAATGATGATGCAGAACGTGGCTATTATGAGTG
TF	bZIP6_PAS	XM_002184884.1	Phatr3_J50039	CGCGCAATCTAATACACAATATCAGCTAGACAGCCCACAACATCCCGAAAATACGATGTCGCAGTGTAGACCGCAGCAGGGTTCAAATAATTCGGCGAAC
TF	bZIP7_PAS	XM_002183297.1	Phatr3_J48800	CGTGAGAGACCTTGCCATGCGTAATCAAGGATTAGAGGCATTAGCGGCGATTGCGACAGCAAGCTCGTCTTCAGGGCACACGTTTAGTACCAACACGAAT
Metabolism	CaThioredoxin	XM_002184396.1	Phatr3_J49634	CTGGAGAGCCCTACCGTGCTAGTGTCCGTGTATTGGAAAAATGTCGTTGTTTTGCGTTCAGCGTAGAAGACATTCCTGACAGCTCCATTCTGAGTCTGAA
TF	CCCH4	XM_002181104.1	Phatr3_J13664	CATGTGCCGCCTTGGAGCGTCGCACATTAACATGTCTACTGGCGAAAACATTCGAAAGGACATTGATGGTCCGAAGCCCAAGCCAATCTTAAATACTCTA
TF	CCCH8	XM_002186091.1	Phatr3_J44042	TCCGAAAAATATGTGCGAAGAGTCCCCACTTGTCGCGTATGTTCTTGGCTCATACGAGTATTGCGCCGACGGAAGTGGAAGAGATTGAGTTTGCGGGTGAA
TF	CCHH1_CCCH20	XM_002179821.1	Phatr3_EG02317	TAACTACAACAACGTTCCTCAGCTAAACGGTTTGAACCATCTGCCTTCTAAGAACCTATCCGACTTACATTCCCACTATCAAGATCATAGGAGTCCAAAC
TF	CCHH14	#N/A	Phatr3_J34600	GCCCAGTTTCTGCGGAGGCATCCCGTTCAATCCGCCGATCTGAGTTACGAAACGCACTACATGGCGTACCAAAAAGTTTACACTGATGCCCGAGTTGTTC
TF	CCT3	XM_002185983.1	Phatr3_J43850	AAACGCTGCCGCCGCGTTTGGAATAAGAAGATTCGCTACGGTTGCCGTAAAAATTTAGCAGACCGCCGGTTGCGCGTGAAGGGGCGATTCGTGAAACGTT
TF	CCT4	XM_002178133.1	Phatr3_J44285	TATGAATGCAGTCGACCGGGAATTGAGTTTCCATCAAACAACAGAGCTAAGCTGGTCGAGCACCAGCACGGCCTTTCGCCCATTCCTTTTCACGATAGTG
Cell cycle	cdc20	XM_002180546.1	Phatr3_J12783	TTACGACGGAATTGCTTTGAAGAAAATCAGAACACTTCATGGGCACACAGGCCGAATATCGTCGTTGGGATGGAACCAACACTGGTTGAGTTCAGGCGCA
Photoreceptor	Cpd2	XM_002179343.1	Phatr3_J54342	TCTGGTGTGCGGCAATACGGCGGACAGTGTTTCGGGAGCTCTGTAAAATTGCTTCTGAGATTGGCGCGAGTGCAGTCTATTGGAATCGCGAGATGACACCT
Photoreceptor	Cpd3	XM_002180035.1	Phatr3_J51952	CTCTGATTTTTCTCCGATACGCGAATACCGTCAATGGATGGAACTCCAGGCCGTACCTATCTTGGAGGAAGCGAAGGTCCCGTTTTATCAGGTTGATGCC
Photoreceptor	Cpf1	XM_002180059.1	Phatr3_J27429	GGATGCGTGCCAAAAATCCGAATCACTCCTTCCGATTTACGTAGTCGACCCTGAATTTCCCTTCGCGCAAACTGCTGGGTGCCGCGCCGGTACAATTCGT
Photoreceptor	Cpf4	XM_002184521.1	Phatr3_J55091	CTCGATCCCTTGTTGCGGCGCAAACGAGACTGCATTTCCGTAATGGGTCTACCTAATGATTTGTAGACTCCATTGTCGAGGCAGCGTTTGAAGCAGCTG
Photoreceptor	CryP-like	XM_002178853.1	Phatr3_J34592	CACTTTATGGTGGCGAAACAGCCGGACTCGCACGATTGTTTGAAGGATTACTTCGAAACGCGCAACGGGATGCTGGGGCCGAACTATTCGACCAAATTCA
TF	CSF2	XM_002185522.1	Phatr3_J41601	GAACAGAACGAAAAGATTCGGGACGCTTATGCATTCCAACAGACGAAAATCGACGAGGCTCATCAACTTATTGTCAATTTGGGCATGGACGTTGCAGACT
Cell cycle	CYCB1	XM_002180361.1	Phatr3_J46095	TTAGAGAAAAGTCAATCGCAAATCCTGAGCTCCGAGCGGTGAATAAAAAGTACAGCGGTCATCGATACGGCGGAGTTGCTTCGACCGTTCTGGTATTTGA
Cell cycle	CYCH1	XM_002185709.1	Phatr3_J36892	TTTGCGATGGCTCGCGTTTGACGTTTCGGTTTCCCATCCGCACCGAGGAGTTGCCGCAATCGTCAACCATCTCGTGGATTGCTCACCTCAATGGTTTGGC
Photoreceptor	Dph	XM_002179026.1	Phatr3_J54330	CAACAATTCCATTACAACCAAAGAGCTGACGGAATGTGATCGTGAGCCTGTGCACTTGATCGCAAACGTACAAGGGGGTACCGGCCATTTGTTGTTCATT
Cell cycle	dsCYC2	XM_002179247.1	Phatr3_J34956	GATATTGCCGTTACCGAACGCGAAATAATGACTGCTCTTTCCTGGCACTTGCATCCTCCAACCGCTATTGGCTTTTCACGAATGTACTGGTCTCTGCTGG
TF	E2F_Dp2b	XM_002181452.1	Phatr3_EG02016	TGATTTGAATAGAGCTGTGCAGGAAATGAGGGTACAAAAGCGCCGGATTTATGACATTACCAACGTTTTGGAAGGTATTGGGCTGATTACCAAGGATAGC
Cell cycle	E2F1	XM_002176842.1	Phatr3_J43065	GTGTGGGTCTCATCGAAAAACGATCCAAAAACACAGTCGCTTGGAAAGGGAGTGAGCTTCTTCTTGGCTCGTCCTTTTCAAGTGCCGCCAAGCAGAGAAT
Cell cycle	FtsZ	XM_002185088.1	Phatr3_J42361	GAAAAAGGAGAATGAAGCAAGCGATTGCCGCGATTGAGAGGTTGCGAGAGAATGTGGATACGGTCATCGTAGTGTCCAACGACAGACTGCTCGAGATTAT
Metabolism	GapC1	XM_002182255.1	Phatr3_J22122	TGAAGGGATTCCTCGGATACTCCGACGAACCGTTGGTCTCCACCGACTTTGAAGGTGACTTGCGCTCCTCCATCTTTGATGCCGATGCCGGTATCATGCT

Metabolism	Gsat	XM_002180931.1	Phatr3_J36347	ATTCGTTTGGTCCGCGCACACACGGGTCGGGAAAAGGTCATCAAGTTCGAAGGATGCTACCACGGACACGCCGATTCATTTTTGGTCCAAGCCGGATCCG
TF	Hox1	XM_002184935.1	Phatr3_EG02213	AACAAACATCGGTTCACATACTGCGTCATACCGGACAAGTCCCGACAGTCGAAGATGTTACGATCCTTACCAATGTTCCGGCGGAACGCATTCTCCGAAC
TF	HSF1.3b	XM_002179894.1	Phatr3_J35419	CTCCAGGGAACAACTCGATACCAGAGTTCCTCTATCAGCTAACAAAGATGCTGACGGATAACAACCGAGATATTATTGAATGGACAAATGGCAAGATTGA
TF	HSF1a	XM_002177362.1	Phatr3_J43051	CCACCAACGTCGGATAGCTCCGACTGGGGTTCTCCAATCCAATCATCCAAGCCAGCTGTTGATCTCTTAACGAGGAGTATGCCGCAATGGACAAATATAC
TF	HSF1b	XM_002181395.1	Phatr3_J47181	TTGTAGTGGACACTTCGATGCTCGAAGAGGTCATCAACGAAACGCAGCCACTCGTTGGAGACAAAGGTAGCTTCTCCCTCTTGACAAATAGAGCACCAGT
TF	HSF1d	XM_002178673.1	Phatr3_J44750	GTCACCATTCCAGCATCGGATGCGGCTCCTTACGCCATTTCTCCCGAAACCTCACCGAGAGACTTGACTCTGAGACACAACGAAATTCCCAACCTTGATG
TF	HSF1g	XM_002177016.1	Phatr3_J42514	ATAGAGAATCTGCAAGCACTAGGGACCATGACGACTACTTCATCGAAGCGTAGCATTACAGAGAGCGAAGATAGTTCCCTGTCAGACTCAGCATCAGGAG
TF	HSF2.1a	XM_002181681.1	Phatr3_J47360	AGCCCTTGCCGCTGGAATTTTGGGCTCGTTGTCGCCTCAAGTCAACCAACCACAGCCGTGGATGATCCAACACGCCCTGTCAATCGCTCAAGCGTTAGTT
TF	HSF2.2c	XM_002185405.1	Phatr3_J50481	GGACTGGAAACTGTAAGCGCGACATATTTCCGCAGCGCCTTCTAGCGATTTTGAGCGAATCATCCCTATCTGACATTATCACTTGGCTTCCACATGGACG
TF	HSF3.2b	XM_002186124.1	Phatr3_J44099	TTGACTCCTCGCGACCAGCAAGCACTTAGCGCGTTTCAACAGTCCCTCGGTGCATCCGAGAGTCAGTTCAAGTCTATGAGCTTTTCCACCACAACTCCTC
TF	HSF3.2e	XM_002179404.1	Phatr3_J45206	TTGGAACGGGAACGCTTTCCCTCAGATTAGCAATCCGCAACCTTCGCCTACTCCTCCAAGTTCGACGATGGTAAATACTTCGGACTTAGCAAGACTACTT
TF	HSF3.3a	XM_002179738.1	Phatr3_J45393	TCTGGGCAAATCTTTCGTATGGAAGGTTTGGCGACGCCGAGTAGTCAGGGAGTTGGCGTAACCACAGGCAGTTGCCGCTTCGAGACCAAGACGCTGTTA
TF	HSF3.3f	XM_002179735.1	Phatr3_J45389	CATGCCAATTGAACACCCAATTCAGCCATACATGGAAGAGCTTAAGAGGGTCCATCTTGCAGCGGCGGCGGGAAGTCTCGAAAATGCGCTTCGAATTTTG
TF	HSF4.2c	XM_002182762.1	Phatr3_J48361	CGCACGTATATCACGATTTCGCACAAGTGCGGCCCGACGCGGATCCCATGATGATCGTCCGGAAAAAGACTGGAGGAGTCACTCAGCCATTTCCCGAAAA
TF	HSF4.2j	XM_002177548.1	Phatr3_J43363	GTCGGCGGACGGGCCTTTATCATTCACAAGCCTGACAAGTTCTTTAAGGATATAGTGCCGCTATATTTTCGCCAGTCGCGGCTGAGTTCATTCAAGCGGC
TF	HSF4.3a	XM_002184368.1	Phatr3_J49594	CGCGAACACTCGCGTTCCAATGTCGCGGAAGAACCTTCCAAGCGAACCTTTGTTCAGCATATTTACCATGACCATATGAACGACCCTGAGCAGGTGGATG
TF	HSF4.3b	XM_002183012.1	Phatr3_J48558	CTGTCTTGAGCCTATCGCAGCAGTGGCGCCGAATTTCTACGCTATGCCAACTACTGAAGGTGGAGACAAAGACCAATTATTCTCGATCCAAAATACGTGG
TF	HSF4.3c	XM_002183081.1	Phatr3_J48667	AAACTGTCAGTCATGCTTGACCATGTAAAAGCCGCAGGATTGGACGACGTTATTTCTTGGGCCAGTCATGGCCGCTGTTTTAGCATTCACAACCCAGACC
TF	HSF4.4b	XM_002184371.1	Phatr3_J55070	CCCATTCCGGTAAACCCGGTTTCCAACTTGGGCGTATCCAGAGCGCCTAGTTTTGGTAAGCTCGAACCAGGCCAATTCGCTGCTTATGGCGGTTGTATGA
TF	HSF4.6a	XM_002184347.1	Phatr3_J49557	CTCAAACACTGGCGATTGCCGTATCGCCTCGTTCCGCCATCCAGTGCACGCCAACATCGTGAATCCTTCGCACTTTCGCTCGTCTCCAATAACGGGCAC
TF	HSF4.7a	XM_002185271.1	Phatr3_EG00092	CGGACATCCCGGGGCGGATTTGAGATCGACGCGATAGTGTCATTCTTGCGTTCCAAGAGTATTTCGCAACTCTACATTATCGGTGGTGACGGTACTCACC
TF	HSF4.7b	XM_002184277.1	Phatr3_J49596	CTCGAGGTGGTGTGGGAACTCCTTTTCCACTCAAGCTGCACGAAATGCTTCAAAATGTCGTACAAGACGGCTACGCTCACATTGTATCGTGGCAGCCTCA
Metabolism	Lhcx1	XM_002179724.1	Phatr3_J27278	AATCCTTGAGAATCTTCAGGGTTAAAGAGTGCATCCATCCAAAACTGGAGGTATGGCTGGTCACATGCACAGAGTTAACAACCAATCCCTTTTCAAGCGA
TF	Myb1R_SHAQKYF4	XM_002178425.1	Phatr3_J44331	CGTATCTGCCGAATCGGTCTCTGGTGCAAATCAAGTCCCACGCCCAAAAAGTGCTCAAGCGCATCGATCAAGGCGAACACGTCTTCCGACGTATCGAGGA
TF	Myb1R_SHAQKYF5	XM_002181623.1	Phatr3_J47256	ACGCTTCACCTGGCCCGAAGACTTACACCGGGATTTTGTATCGGCAATTTTTGATGTCGGCTTGAAACAGTCGTCACCGTCCACCATTTTGGAAACCATG
TF	Myb1R_SHAQKYF6	XM_002180835.1	Phatr3_J46535	TAACGACGAAGCGCAAATCCTTACCGACAAACTTGCCGTCCCGCAAAAAGGGTAAGACACGAAAGTCTGCCGCCTTGGTACAGCGTAATACTTCAGTTTC
TF	Myb1R10	XM_002181361.1	Phatr3_J37257	CATATCCCTCTTCGGGGCCCTCACTTCCCTGTACAAGTGAATCTTTGGGGGTGAAAGCGAATCGTAGAAAGAAACGTGCTAAGGACGAGACCTCGGTTCT
TF	Myb1R4	XM_002181410.1	Phatr3_J47205	AAAGTTCGTTGCAGGTTTAGACGGAACGTCTACCGCATTGTCAATGGGAACAATCCGGCGAGTAGGATTCACCCATATGCGATTGAACTTCTGGAAGCCG
TF	Myb2R3a	XM_002181911.1	Phatr3_J47726	TCCATCTTTCCTTTGAGCGACTGTGACGACCTGTGGAGGGAGAGTGATTTTTGTCACTGTGAAGTTGTCGGTACTCCGGCACTCGGCAGGTAAAATACTC
TF	Myb2R3b	XM_002182076.1	Phatr3_EG02275	GCGCACAACGGTTTAAAGTGATTCGGGATTCATTCACAACAGTCAATGGCTAATTGGCCTCCATCGTTCACTGGTTGACATTTGCGATCGGCCCAACACT
TF	Myb3R1	XM_002182738.1	Phatr3_J15016	GTATGATCCTGGAGTGTCATTTGACGTTGGGAAACCGCTGGGCCGAAATTGCCAAGCGTCTACCGGGAAGAACTGATAACGCTATTAAAAATCACTGGAA
TF	Myb3R2	XM_002180449.1	Phatr3_J6839	CGAATTCTCATAGGAGCACAGGACACAATGGGAAATCGATGGGCTGAGATTGCCAAGCGGCTGCCAGGACGTACGGACAATGCGATAAAGAATCGCTGGA
TF	Myb3R4	XM_002180741.1	Phatr3_J36337	GACAGATGCCGAAGACGCAATTGTAATGGATGCGGTCAATTCCAGCTCGGAGCAACCATTTACTCGTTGGTCAGACTTGGCGCAACGTCTACCCGGACGT
TF	Myb3R5	XM_002184658.1	Phatr3_J7959	GACGTCGTTATCTGAATTGCTTGGACCCAACAATCCGAAAAGACGAATGGACGGAGCAAGAGAAAGAAGCTTTGTTTGATGCACAGCGGCGCATGGGAAA
TF	Myb5R	XM_002185245.1	Phatr3_J50365	TCAACACCACATAAACGCTCGTTTCAGGCTTTGGATGTGTTAGTGAACGCGGTTGGTTTTCATCGAGAACCAAATGACCAAATGAAGCAGACGAAGATGT
Cell cycle	PCNA	XM_002182225.1	Phatr3_J29196	AATGCAGGAACCGGTCGAACTCACCTTTGCCCTGCGCTACCTCAACTTTTTTACCAAGGCGACTCCGTTGAGTGGACACGTCATTATCTCCATGGCACCG
Metabolism	Pds1	XM_002180135.1	Phatr3_J35509	AGCATGGCCAAGGCACTCGATTTCATTGATCCCGACAAATTGAGTATGACGGTCGTTTTGACCGCCATGAATCGCTTTTTGAACGAAGACAACGGCCTCC
Metabolism	petA	GI:118411009	#N/A	TCTCTGGATCAGGAGCCATAACAGGGAAAATTAACTCTTGATGAGTCTTTCCTGCAATTGGTCCAACTACTAAGATATTATCAAATTCACTACTATACGG
Metabolism	Pgr5	XM_002178672.1	Phatr3_J44748	GTAACGTTCGCGCTTCTGATCGCTACCGCTGCCGCGTTTCGCGTGGCGGGCGGAGCCACCGGGGGTTCTTCCTTTGTGCTTTCCCGACGAACCAGCATCG
Metabolism	PgrL	XM_002177043.1	Phatr3_J42543	CCACAACAAGTAGCGTTTCCAATTTAGACAAGACTACGAACAAACGATGCGAGTCAACTCTGTGGTATTTGGCCTTTTGGTGTTCTCGGCCGTCAGCGAT
Metabolism	Por1	XM_002179653.1	Phatr3_J12155	TCGGGTGGTGAGATTTTTGAGAATCAGCAGTCCGACGCAGTACGGGATCTTCCTACCGCCAAGAAAATGTGGAAACTGAGTAGGGAAGCAGTTGGTCTTT
Metabolism	Por2	XM_002180956.1	Phatr3_J13001	AATCACCACTATTCCGTGAAAAGCGCCCGTGGTTCCGTAAATACTTTCCCATCTTTATGAAGTTTATTACGGGAGGATACGTTGGAGAGCACGAAGCCGG
Metabolism	Por3	XM_002177432.1	Phatr3_J43164	CGCGTTCTCAACACGAGGAAAAGCGGTGAGACTCATGATTCGTGATCCGACCAGTAGTCGTATCGAGGCACACCGTAGCGTGGGAGACGTAGACGTTCCC
Metabolism	psaA	GI:118410964	#N/A	GCTAAGAAGAACGCCCAAGTAGTTCCAATACCACCCAATAAATAATGAGCTAAACCAACTGCACGACCCTGAGTAATGCTTAATGCACGAGGTTGGATTG
Metabolism	psaD	GI:118410999	#N/A	TTAAATGTACGTAATTGGGTCCCAATGCTAAACATTGCTCTTTGCGCGCTAAGTATAATAAATTTTCTCCACTACGCATGATAGCTGCTCCACCAATTG

Metabolism	psbC	HQ912250.1	#N/A	GGCCCAACAGGTCCTGAGGCTTCACAAGCACAAGCATTCACGTTTTTAGTTCGAGATCAACGTTTAGGTGCTAATGTTTCATCAGCACAGGGTCCAACTG
Metabolism	Psy1	XM_002178740.1	Phatr3_EG02349	ACCTGTCGTCGTGGGAATTGCGATTGGAGCGTTTGTGGCAATACGGACAAGTACAGGATGTCTTTGACCTGTGCTTGCTAGATCTTCGAATCCAGTATCC
Reference Gene	Rps	XM_002178225.1	Phatr3_J10847	CTCCGCGCATTTTTGCCCGGTTCCCATTTGACCGGACAGTTGCCCGACGAAGATCTCATTGGAAACAACTTGCAGCTTAAATTCCTCGAAGTCAACCAGG
TF	Sigma70.1a	XM_002182291.1	Phatr3_J14599	GACAATTAGGGGTATCTCGGGATCGGATTCGCCTTGTCGAAACCCGAGCACTGAACAAATTGCGACACCCGCAACAAAATTACAAGCTGCAGTCTTACGT
TF	Sigma70.1b	XM_002182124.1	Phatr3_J3388	TTCTTGGTCATTGGAAGAAACGGCCAAACAACTAGGGATGTCTCGGGATCGGATTCGCCTTGTCGAAGCCCGAGCACTGAATAAATTGCGACGTCCACAA
TF	Sigma70.2	XM_002178682.1	Phatr3_J5537	ACGTGGATTGCGATTCGCCACCTACGCCACTTACTGGGTCACCAATTCCGTACGCTTCTGCTTCCAAACCGCATCCACCGGATGTTTGCGGGTACCGGTA
TF	Sigma70.3	XM_002185461.1	Phatr3_J17029	TATGATGGTTCCTGATCTAAAGAAGCAAATTCGTCGCTCGCTTCGCGCCAAGGCTGCCCTGACGGAATCAAATATTCGACTCGTGGTATCCATTGCGAAA
TF	Sigma70.4	XM_002177232.1	Phatr3_J9312	AACCCGAAAGATCTCGGTCACGAAAGTCACGATTTGCTAGGGGACACAATCTCCGCCAGTAGTGCTCTTTTTGACGAAAGCACACCCGAAAAACGAGTCG
TF	Sigma70.5	XM_002182854.1	Phatr3_J14908	AAGAACTCATACAGGAAGGCAGTTTGGGATTGTTGCGCGCGGCCGAATTGTTTGATCCCGAGCGAGGACTGCGATTCAGCACTTACGCGGTAGTGTGGAT
TF	Sigma70.6	XM_002178042.1	Phatr3_J9855	CTGGAAAAATCAATATGGAAGCTATTCGGCAGACCATTGAAGAAGGACTGGAAGCCAAAAATCAGCTAGTCACTTCGAATCTGCGGATGGTACAGAGTGT
TF	Sigma70.7	XM_002185047.1	Phatr3_J50183	AATGATGACGCTCCTCTCGCATACGGTGGAACTCCGTCGTATCAAGCAAATCGAAGCTGATCTTGCTTTGACGTATCCATCTTCCCGGAAACCCACTACT
TF	Taz2	XM_002178935.1	Phatr3_J44807	CCTAATGTAGAGACTGTAGTGGTAAAAGCGAAGCGCGCAGCCTCGACTATTTGGACTTTACTTCATGCACAGAATTGCGTTTTGGGGGACAGATGCCCAC
Reference Gene	Tbp	XM_002186285.1	Phatr3_J10199	CTGATTTTTTCCAGCGGTAAAATGTGCGTGACCGGAGTCAAGAGCACACACAACGCCAATCTGGCGGCGAAAAAGTTTGCCTACATTGTGGAGCGCGTCG
Metabolism	Vde	XM_002178607.1	Phatr3_J51703	GGCGATGACGGATCATATCCCGTTCCCGCACCCGAAATCGTCGTACCCAAGTTTGACACCAAGTTCTTCGATGGACGACTCTACATCTCAGCCGGACAAA
Metabolism	Vdl1	XM_002180599.1	Phatr3_J36048	AAGGCTCTCTTTAGCAATCCGCGTGGAATCAAAGGTGTCTCCTGTTTGGGACGCTGCAAGGGCGAGCAATCGTGTGCGACGCGGTGTTTTGCCGAATTCG
Metabolism	Vdl2	XM_002180015.1	Phatr3_J45846	ACACACCGCCTTCGGCAATGAATATCCAATTTCCGGAGCCAGACGAATCACTCCATGTCTGGGACCACGTACGAAACATCGGTAAAGTGTGTACCGGGTT
Metabolism	Vdr	XM_002177477.1	Phatr3_J43240	GTCGCTTGCGGTGCCAACGAAGCCTATGATAAGTTTCCTTCGCAGAACCAACTCTTCTATCCCGCTGCTCGCGGTCGAGATTTGTGGTACGATCCCGTTT
Metabolism	Zds	XM_002184417.1	Phatr3_J30514	CGGCGTCAACGAATCCATGCGGGACACCGCCAACGTACTCGCCCGTTTTAATTCACTCATTCTCGCCCGAGTCTTTGGACACGCCGACGTCACGGAACTC
Metabolism	Zep1	XM_002179586.1	Phatr3_J45485	TTCCCAATGAAGTTTTCTACCACGGTGTCATCGGCACTGTTCCTGATCGCGTCGGTATCGACCACGACTTCCTTCACACCCGTCCAGTCCTTTGGCGTGC
Metabolism	Zep3	XM_002178331.1	Phatr3_J10970	CCAAGGTTTTGACCAAGATGCCCACCATGGATGTTACCGTACTGGAACAAACGTCCGAATTCAAACGGTTCGGTGGACCCATCCAGCTCGCTAGTAACGC

825 **Table S2:** Calculated phases, amplitudes and periods of the Wt and *Pt*bHlH1a OE lines growing
826 in 16L:8D. (P., Period; Ph., phase; Amp., amplitude; Std, standard deviation; t-test, t-test P-
827 value between Wt and the indicated line).

Data	N	Period	P.Std	P. t-test	Phase	Ph.Std	Ph. t-test	Amp.	Amp.Std	Amp. t-test
Wt	8	23.7	0.32		13.02	0.54		8.30E-01	1.20E-01	
OE-1	5	23.37	0.51	0.168894675	15.93	1.83	0.001213017	5.60E-01	2.60E-01	0.02447126
OE-2	6	23.38	0.48	0.159824505	15.19	1.52	0.002642208	6.10E-01	7.00E-02	0.001606661
OE-3	4	23.89	0.22	0.323356278	13.94	0.51	0.017556508	8.40E-01	1.20E-01	0.841165049

830 **Table S3:** Accession numbers of the proteins utilized in the bHLH-PAS phylogenetic analysis.

SPECIES	ABBREVIATION	ACCESSION	ACCESSION(Alt)
Amphiprora sp., Strain CCMP467	Amph	CAMPEP_0168730192	
Asterionellopsis glacialis, Strain CCMP134	Agla2	CAMPEP_0195290872	
Asterionellopsis glacialis, Strain CCMP1581	Agla3	CAMPEP_0197142484	
Astrosyne radiata, Strain 13vi08-1A	Arad1	CAMPEP_0116831938	

Astrosyne radiata, Strain 13vi08-1A	Arad2	CAMPEP_0116837632	
Attheya septentrionalis, Strain CCMP2084	Asep	CAMPEP_0198303966	
Capsaspora owczarzaki ATCC 30864	Cowc1	XP_004345696	
Capsaspora owczarzaki ATCC 30864	Cowc2	XP_004343694	
Chaetoceros debilis, Strain MM31A-1	Cdeb	CAMPEP_0194099558	
Chaetoceros dichaeta, Strain CCMP1751	Cdic	CAMPEP_0198277646	
Chaetoceros neogracile, Strain CCMP1317	Cneo2	CAMPEP_0195415676	
Chaetoceros neogracile, Strain CCMP1317	Cneo3	CAMPEP_0195453836	
Chaetoceros sp., Strain GSL56	Chae	CAMPEP_0176495412	
Corethron hystrix, Strain 308	Chys	CAMPEP_0113330172	
Corethron pennatum, Strain L29A3	Cpen	CAMPEP_0194280324	
Cyclophora tenuis, Strain ECT3854	Cten1	CAMPEP_0116540656	
Cyclophora tenuis, Strain ECT3854	Cten2	CAMPEP_0116579710	
Cyclotella meneghiniana, Strain CCMP 338	Cmen	CAMPEP_0172279412	
Cylindrotheca closterium	Cclo	CAMPEP_0113624952	
Dactyliosolen fragilissimus	Dfra	CAMPEP_0184871870	
Detonula confervacea, Strain CCMP 353	Dcon	CAMPEP_0172306268	
Ditylum brightwellii, Strain Pop2 (SS10)	Dbri	CAMPEP_0181012646	
Drosophila melanogaster	dmClock	AAC62234	FBpp0099478
Drosophila melanogaster	dmCycle	NP_524168	FBpp0074693
Drosophila melanogaster	dmSim	AAC64519	FBpp0082178
Drosophila melanogaster	dmTango	NP_731308	FBpp0081483
Drosophila melanogaster	dmTrh	AAA96754	FBpp0293065
Durinskia baltica, Strain CSIRO CS-38	Dbal	CAMPEP_0170381008	
Ectocarpus siliculosus	Esil1	Ec-02_004780 PAS domain (681) ;mRNA; f:5082087-5086119	
Eucampia antarctica, Strain CCMP1452	Eant	CAMPEP_0197830682	
Extubocellulus spinifer, Strain CCMP396	Espi1	CAMPEP_0178587872	
Extubocellulus spinifer, Strain CCMP396	Espi2	CAMPEP_0178723458	
Fragilariopsis kerguelensis, Strain L26-C5	Fkue1	CAMPEP_0170867208	
Fragilariopsis kerguelensis, Strain L26-C5	Fkue2	CAMPEP_0170907188	
Fragylariopsis cylindrus	FcbHLH1a	jgi|Fracy1|204663	
Fragylariopsis cylindrus	FcbHLH4	jgi|Fracy1|241747	
Fragylariopsis cylindrus	FcbHLH7	jgi|Fracy1|172104	
Galdieria sulphuraria	Gsul	XP_005709404	
Glenodinium foliaceum, Strain CCAP 1116/3	Gfol1	CAMPEP_0167841300	
Grammatophora oceanica, Strain CCMP 410	Goce	CAMPEP_0194036250	
Guillardia theta CCMP2712	Gthe1	XP_005835918	
Homo sapiens	hsArnt	AAA51777	
Homo sapiens	hsbMal1	NP_1284651	
Homo sapiens	hsClock	AAB83969	
Homo sapiens	hsNpas1	AAH39016	
Homo sapiens	hsSim2	NP_5060	
Kryptoperidinium foliaceum, Strain CCMP 1326	Kfol1	CAMPEP_0176099000	

Leptocylindrus danicus var. danicus, Strain B650	Ldan1	CAMPEP_0116039394	
Leptocylindrus danicus var. danicus, Strain B651	Ldan2	CAMPEP_0116053596	
Leptocylindrus danicus, Strain CCMP1856	Ldan3	CAMPEP_0196809694	
Licmophora paradoxa, Strain CCMP2313	Lpar	CAMPEP_0202478474	
Minutocellus polymorphus, Strain RCC2270	Mpol	CAMPEP_0185804732	
Monosiga brevicollis	Mbre2	jgi|Monbr1|26507|	
Mus musculus	mmArnt	AAA56717	
Mus musculus	mmbMal1b	BAD26600	
Mus musculus	mmbMal2	ABC50103	
Mus musculus	mmClock	AAC53200	
Mus musculus	mmNpas1	NP_32744	
Mus musculus	mmSim2	AAA91202	
Nannochloropsis gaditana CCMP526	Ngad_2	Naga_100016g14	
Nannochloropsis gaditana CCMP526	Ngad_3	Naga_100165g4	
Nannochloropsis oceanica CCMP1779	Noce	NannoCCMP1779_9983-mRNA-1	
Nannochloropsis oceanica CCMP1779	Noce_2	NannoCCMP1779_1600-mRNA-1	
Nitzschia punctata, Strain CCMP561	Npun	CAMPEP_0178827522	
Nitzschia sp.	Nitz	CAMPEP_0113465330	
Phaeodactylum tricornutum	PtbHLH1a	jgi|Phatr2|44962	Phatr3_J44962
Phaeodactylum tricornutum	PtbHLH1b	jgi|Phatr2|44963	Phatr3_J44963
Phaeodactylum tricornutum	PtbHLH2	jgi|Phatr2|54435	Phatr3_J54435
Proboscia alata, Strain PI-D3	Pala	CAMPEP_0194393728	
Pseudo-nitzschia arenysensis, Strain B593	Pare	CAMPEP_0116139930	
Pseudo-nitzschia australis, Strain 10249 10 AB	Paus1	CAMPEP_0168282354	
Pseudo-nitzschia australis, Strain 10249 10 AB	Paus2	CAMPEP_0168310394	
Pseudo-nitzschia delicatissima, Strain B596	Pdel1	CAMPEP_0116091770	
Pseudo-nitzschia delicatissima, Strain B596	Pdel2	CAMPEP_0116094064	
Pseudo-nitzschia fraudulenta, Strain WWA7	Pfra1	CAMPEP_0201229664	
Pseudo-nitzschia fraudulenta, Strain WWA7	Pfra2	CAMPEP_0201232896	
Pseudo-nitzschia heimii, Strain UNC1101	Phei1	CAMPEP_0197189528	
Pseudo-nitzschia heimii, Strain UNC1101	Phei2	CAMPEP_0197199168	
Pseudo-nitzschia multiseries	PmbHLH1a	jgi|Psemu1|26622	
Pseudo-nitzschia multiseries	PmbHLH7	jgi|Psemu1|228145	
Pseudo-nitzschia pungens, Strain cf. pungens	Ppun	CAMPEP_0172413298	
Rhizosolenia setigera, Strain CCMP 1694	Rset	CAMPEP_0178942232	
Skeletonema costatum, Strain 1716	Scos1	CAMPEP_0113408658	
Skeletonema dohrnii, Strain SkelB	Sdoh	CAMPEP_0195221452	
Skeletonema japonicum, Strain CCMP2506	Sjap	CAMPEP_0201725346	
Skeletonema marinoi, Strain FE7	Smar1	CAMPEP_0180846198	
Skeletonema marinoi, Strain UNC1201	Smar2	CAMPEP_0197231700	
Skeletonema menzelii, Strain CCMP793	Smen	CAMPEP_0183679140	
Stauroneis constricta, Strain CCMP1120	Scon1	CAMPEP_0119548954	
Stauroneis constricta, Strain CCMP1120	Scon2	CAMPEP_0119572288	

Staurosira complex sp., Strain CCMP2646	Stau	CAMPEP_0202487246	
Synedropsis recta cf, Strain CCMP1620	Srec1	CAMPEP_0119009808	
Synedropsis recta cf, Strain CCMP1620	Srec2	CAMPEP_0119029558	
Thalassionema frauenfeldii, Strain CCMP 1798	Tfra1	CAMPEP_0178899126	
Thalassionema frauenfeldii, Strain CCMP 1798	Tfra2	CAMPEP_0178923098	
Thalassionema nitzschioides, Strain L26-B	Tnit1	CAMPEP_0194218124	
Thalassionema nitzschioides, Strain L26-B	Tnit2	CAMPEP_0194240022	
Thalassiosira antarctica, Strain CCMP982	Tant	CAMPEP_0202006238	
Thalassiosira gravida, Strain GMp14c1	Tgra	CAMPEP_0201684800	
Thalassiosira miniscula, Strain CCMP1093	Tmin	CAMPEP_0183747574	
Thalassiosira oceanica	Toce2	EJK65393	
Thalassiosira pseudonana	TpbHLH1	jgi|Thaps3|24007	
Thalassiosira pseudonana	TpbHLH2	jgi|Thaps3|20899	
Thalassiosira pseudonana	TpbHLH7	jgi|Thaps3|23208	
Thalassiosira sp., Strain FW	Tfw	CAMPEP_0172354094	
Thalassiosira weissflogii, Strain CCMP1010	Twei	CAMPEP_0203515806	
Thalassiothrix antarctica, Strain L6-D1	Txnt	CAMPEP_0194143228	
Tiarina fusus, Strain LIS	Tfus1	CAMPEP_0117003500	
Tiarina fusus, Strain LIS	Tfus3	CAMPEP_0117046192	

833   **Table S4:** List of the oligonucleotides used in this work.

Gene	Phatr3 Accession	Oligo name	Sequence	Type
bHLH1a	Phatr3_J44962	bHLH1a-endogenous-QFw	TTATGTCTTTCGGCGACGGG	QPCR
		bHLH1a-endogenous-QRv	AGCAACGAATGCATGCAAGG	
		bHLH1a-total-QFw	ATTCTTGGTCCCACCCGGTA	QPCR
		bHLH1a-total-QRv	ACGCCACATTGAAAAACCGAG	
		Pt-bHLH1a-DraI-Fw	GATTTTAAAATGAATAAGCCAGGACAGCG	Full lenght cloning
		Pt-bHLH1a-XhoI-Rv	TTGCTCGAGCACAGCTTCGCTGCATCGTC	
bHLH1b	Phatr3_J44963	bHLH1b-QFw	TGCGATCTCAACGGCTAATA	QPCR
		bHLH1b-QRv	CGCAAAACGAGGCTAATTTC	
bHLH3	Phatr3_J42586	bHLH3-QFw	CACTCTCATCATGCGGGAAT	QPCR
		bHLH3-QRv	GCGCGTTGTCTTCCTCTATC	
bZIP7	Phatr3_J48800	bZIP7-QFw	CCTTATTGATATTCAAGATTCCAAGG	QPCR
		bZIP7-QRv	GTTTCGGAACCTGCATAGGA	
CDKA1	Phatr3_J20262	CDKA1-QFw	AGCGGTATCAAAGGATGGAAAAG	QPCR
		CDKA1-QRv	CTTCATCTTCGGCTTCAAGGC	
CDKD	Phatr3_J10160	CDKD-QFw	ATTACTTCTGCGGAGACCATTC	QPCR
		CDKD-QRv	GCGGTAAAGATTTCGTCAAAGG	
CYCA/B;1	Phatr3_J17135	CYCA/B1-QFw	TACACCGCCACTCCAAGAC	QPCR
		CYCA/B1-QRv	TCGGAGGACGGGATGGG	
CYCB1	Phatr3_J46095	CYCB1-QFw	TCCTGGTCCGCTACTTGAAAG	QPCR
		CYCB1-QRv	GCTGGCTGGGAAGATAACGC	
Por2	Phatr3_J13001	Por2-QFw	CCTGGTTGCATTGCCGAATC	QPCR
		Por2-QRv	TCTCCAACGTATCCTCCCGT	
Rps	Phatr3_J10847	Rps-QFw	CGAAGTCAACCAGGAAACCAA	QPCR
		Rps-QRv	GTGCAAGAGACCGGACATACC	
Tbp	Phatr3_J10199	Tbp-QFw	ACCGGAGTCAAGAGCACACAC	QPCR
		Tbp-QRv	CGGAATGCGCGTATACCAGT	
VDR	Phatr3_J43240	VDR-QFw	TTTCCTTCGCAGAACCAACT	QPCR
		VDR-QRv	TTGTCGAGCACTGAAAATCG	

References

1. Zielinski T, Moore AM, Troup E, Halliday KJ, & Millar AJ (2014) Strengths and limitations of period estimation methods for circadian data. *PLoS One* 9(5):e96462.
2. Agier N & Fischer G (2016) A Versatile Procedure to Generate Genome-Wide Spatiotemporal Program of Replication in Yeast Species. *Methods Mol Biol* 1361:247-264 .
3. Geiss GK, et al. (2008) Direct multiplexed measurement of gene expression with color-coded probe pairs. *Nature biotechnology* 26(3):317-325.
4. Chauton MS, Winge P, Brembu T, Vadstein O, & Bones AM (2013) Gene regulation of carbon fixation, storage, and utilization in the diatom Phaeodactylum tricornutum acclimated to light/dark cycles. *Plant physiology* 161(2):1034-1048.
5. Saeed AI, et al. (2003) TM4: a free, open-source system for microarray data management and analysis. *Biotechniques* 34(2):374-378.
6. Siaut M, et al. (2007) Molecular toolbox for studying diatom biology in Phaeodactylum tricornutum. *Gene* 406(1-2):23-35.
7. Keeling PJ, et al. (2014) The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): Illuminating the Functional Diversity of Eukaryotic Life in the Oceans through Transcriptome Sequencing. *Plos Biol* 12(6).
8. Katoh K & Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30(4):772-780.
9. Kumar S, Stecher G, & Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33(7):1870-1874.
10. Darriba D, Taboada GL, Doallo R, & Posada D (2011) ProtTest 3: fast selection of best-fit models of protein evolution. *Bioinformatics* 27(8):1164-1165.
11. Miller MA, Pfeiffer, W., and Schwartz, T. (2010) Creating the CIPRES Science Gateway for Inference of Large PhylogeneticTrees. In Proceedings of the Gateway Computing Environments Workshop (GCE) (New Orleans, LA):1-8.