1   Early Holocene Morphological Variation in Hunter-Gatherer Hands and Feet
2

3   Kara C. Hoover[1][*]
4   J. Colette Berbesque [2]
5
6   [1]Department of Anthropology, University of Alaska Fairbanks, Fairbanks Alaska USA,
7   kchoover@alaska.edu
8   [2]Centre for Research in Evolutionary, Social and Inter-Disciplinary Anthropology,
9   University of Roehampton, London UK Colette.Berbesque@roehampton.ac.uk
10
11 *Corresponding author

13 Running title: Hunter-Gatherer Hands and Feet

18 Abstract

The Windover mortuary pond dates to the Early Archaic period (6,800-5,200 years ago)
and constitutes one of the earliest archaeological sites with intact and well-preserved
human remains in North America. Unlike many prehistoric egalitarian hunter-gatherers,
the Windover people did not practice a sex-based division of labor, rather they shared the
load. We explore how mobility and subsistence, as reconstructed from archaeological
data, influenced the morphology of hand and foot bones of this population. Using length-
width ratios to account for trait size and taking the natural log of the ratio to account for
sexual dimorphism, we analyzed 10 variables (4 carpal and 6 tarsal. We identified
differences in the rear-and midfoot regions that suggest the Windover occupation was
engaging in heavy impact mobility. In particular, females exhibit greater variation and
higher mean averages than males for size of shock-absorbing bones. We identified sex-
based differences in carpal bones, with females exhibiting greater variation and higher
mean ratios in most variables compared to males. Despite widespread behavioural
evidence on contemporary populations for human lateralization for hands and feet, we
found no evidence of either handedness or footedness in Windover. The lack evidence for
footedness was expected due its minimal impact on walking gait but the lack of evidence
for handedness is surprising given that modern ethnographic studies have shown strong
handedness in hunter-gatherers during tool and goods manufacture. If the archaeological
evidence for a shared division of labor is true, we have potentially uncovered evidence
for a heavier activity load in females.

1.INTRODUCTION

A sex-based division of labor is seen across most human societies for the majority of evolutionary history and contributes to size-based morphological variation between the sexes [1] [2]. The wealth of ethnographic data on extant hunter-gatherers provides insights into activities ascribed to each of the sexes. Applying what we know about extant hunter-gatherer behavior to the bioarchaeological record, we can link muscular-skeletal markers to habitual activities. Physical markers of activity include geometric variation in bone structure and function due to biomechanical loading [3], diaphyseal structure [4-8], degenerative joint disease (DJD), osteoarthritis, muscular-skeletal markers (MSM) [9], and dental wear patterns relative to tool manufacture [10, 11].

1.1 Mobility Activity

Hunter-gatherer mobility can be described as residential (moving camp to a new location as in seasonal occupation of resource rich areas) and logistical (individuals and/or smaller groups temporarily split from the main group for shorter foraging trips or longer hunting trips) [12, 13]—this is particularly true for mobile hunter-gatherers that specialize in terrestrial resources [14-16]. There is some evidence for sex-based variation in hunter-gatherer mobility; modern Hadza hunter-gatherer males engage in greater daily walking distances at faster speeds than females [17, 18].

Distal limbs have been implicated to a greater degree than upper limbs in reflecting habitual activity due to the biomechanical forces arising from locomotive substrate (i.e., terrains on which activities are conducted), distances travelled in a day, and relative speed of locomotion [7, 9, 17-30]. Of interest to this study is how biomechanical stress from mobility and footedness might affect the tarsals. The ground reaction force generated by the bare (or minimally shod) foot contacting the ground is transmitted through the subtalar skeleton, with peak forces at heel-strike through the calcaneus and at heel-off through the metatarsophalangeal articulations [31]. Thus, during normal locomotion, the typical bipedal heel strike transmits body mass from the tibia to the rear foot (talus and calcaneus) to the ground [32] while the shock of impact is absorbed by the midfoot (navicular, cuboid, and cuneiform bones) [32]. The calcaneus is most affected by the rear heel strike and calcaneal tuber length (as a proxy for Achilles tendon moment arm length) is correlated with running economy (long calcaneal tuber = greater energy cost) [33] Most data from extent hunter-gatherers [30, 34] and barefoot populations suggest a rear heel strike [35] is preferred among experienced runners. Dorsal spurs on the calcaneus are linked to increased activity while plantar spurs are linked to standing, inactivity, and excess weight [27]. While footedness in humans develops in late childhood (11-12 years old) with a right skew [36, 37], its influence on walking gait is not significant and is unlikely to affect the muscular-skeletal system in the absence of other evidence of lateralization [38]. Thus, regions of the foot may be differential shaped by daily logistical mobility that emphasizes either slow walking and stationary weight-bearing activity (such as might occur when foraging) or rapid locomotion such as brisk walking or running that requires greater shock absorption. And, if subsistence based activities are assigned via sex [2, 6], there may be sex-differences in how these regions of the feet vary.

85
1.2 Domestic economies

Domestic economies include subsistence and tool manufacture, which supports subsistence activities. Subsistence covaries with biological factors (e.g., habitat, reproduction, health) and cultural factors (e.g., social organization, sedentism, mobility). A comprehensive analysis of 229 hunter-gatherer diets, eco-environmental data, and plant-to-animal dietary ratios found that most populations consume similar amounts of carbohydrates (30-35% of the diet) except in more extreme environments (i.e., increases in desert and tropical grasslands and decreases in higher latitudes) [39]. Indeed, there is a strong clinal pattern of variation in male and female caloric contributions to diet. Subsistence contributions by sex are inversely correlated with effective temperature, a combined measure of the intensity and annual distribution of solar radiation [12, 40]. Higher latitudes and colder climates rely more on male caloric contributions from big game hunting while temperate and tropical regions rely more heavily on female caloric contributions across the spectrum (e.g., small game, fishing, and plants) [39]. In general, males tend to increase foraging activities in more stable productive habitats [41] and females tend to decrease labor in subsistence activities when males are hunting big game (focusing instead on activities like weaving and cordage) [42].

Ethnographic data indicate that domestic economies such as tool making, child care and carrying, butchering/food preparation, textiles, and carrying firewood and water create physical strain [14, 43-45]. Increased reliance on tools is linked to evolutionarily more gracile bodies [46] but tool manufacturing and use are detectable on the body through increased upper limb robusticity [25] and lateralization, or side preference [47]. Unimanual activities (e.g., spear use) leave a distinct mark of directional asymmetrical in the upper limbs compared to bimanual activities such as grinding or rowing [48]. Indeed, extant hunter-gatherers exhibit strong hand preference specifically when making and using tools [49, 50] which suggests mosaic progression to extreme lateralization we see in modern populations [47]. And, there is evidence in the archaeological record of sidedness varying between the sexes [22, 51]. While domestic economies vary across groups, they tend to be sex-based and more frequently involve lateralized repetitive stress compared to subsistence and mobility [48]. Thus, bioarchaeological evidence of subsistence activities and tool manufacture may found in repetitive stress to the musculo-skeletal system and result in lateralized MSM, DJD, and osteoarthritis of the limbs involved. Carrying loads may place additional weight-bearing biomechanical stress on the foot and the domestic economies may serve to differentiate the wrists.

1.3 Aims

The focus of this paper is to explore whether or not logistical mobility and domestic economies (subsistence and tool manufacture) are archaeologically visible in the feet (tarsals) and hands (carpals) and, if so, whether or not they reflect a sex-based division of labor. Much attention has been paid to sexual dimorphism in carpal and tarsal elements in forensic contexts and with applications to sexing skeletons in bioarchaeology; these studies have had varied success with most pointing to the calcaneus and sometimes talus as the most diagnostic elements [52-64]—for a review see [65]. Carpals and tarsals were chosen for this study for two reasons. First, they are heavily implicated in the daily

activities of hunter-gatherers (e.g., mobility, use of weapons, tool-making, domestic
economies). Second, they are less likely to be influenced by the noise created from
conflicting signals of genetics and lifestyle that obfuscates differentiation of ultimate and
proximate causes of variation that plagues the long bones [21]. Third, dense and small
bones, such as carpals and tarsals, are among the better-preserved elements in the human
body (Henderson 1987; Mann 1981; Waldron 1987) and if daily activities are visible in
these elements, we are potentially able to capture data that might otherwise be lost in less
well-preserved skeletons.
139
We examined carpal and tarsal data from the Florida Early Archaic Windover Site which
contains the remains of a population of hunter-gatherers experiencing a major climate
change in North America and, along with it, a change in domestic economies. Warmer
climate was driving big game north and broad-spectrum foraging was emerging as the
primary subsistence economy, de-emphasizing the dietary contribution of males and
increasing the contributions of females. We generated research questions based on
general cross-cultural patterns identified in hunter-gatherers (Section 1) and specific
patterns identified at the Windover Site (Section 2). As such, research questions are
placed at the end of Section 2.
149
150 **2. MATERIALS (BIOARCHAEOLOGICAL CONTEXT)**
151
The Windover mortuary pond archaeological site (8BR46) is a National Historic
Landmark dating to the Early Archaic Period (7500-5000 BCE) with calibrated radio-
carbon dates from 9000 to 7929 BP. In general, the North American Archaic Period
(8000 to 1000 BC) is characterized by hunting-gathering subsistence economies with
dietary staples including nuts, seeds, and shellfish [66]. The Florida Archaic Period
follows the same pattern (e.g., broad spectrum hunting, fishing, and plant gathering and
use of freshwater resources) with increased exploitation of coastal shellfish and marine
resources. The comparatively wetter climate [67] created an abundance of resources and
and subsistence strategies were no longer dominated by big game. The broad spectrum
foraging strategy that emerged is reflected in more complex tool kits [66, 68].
162
Windover was used as a mortuary pond with 5-6 short periods of activity, peaking at
7450 BP [68]. Burials furthest from the pond edge at time of excavation dated to the
earliest period of mortuary pond use and those closest, more recent. Roughly 100 burials
were undisturbed with fully articulated bones; ages ranged from infancy to over sixty-
five, with 52% classified as subadults [69]. Most individuals were buried within 24 to 48
hours after death (Doran 1992; Dickel 1988, Hauswirth 1991) in a flexed position, on the
left side with heads oriented to the west, and pinned by sharpened stakes approximately
1m below the surface of the peat [70]. The nearly neutral pH of the pond (6.1-6.9) created
ideal conditions for preservation of both skeletal and soft tissues; allowing researchers to
sequence DNA from preserved brain matter [70], reconstruct diet from preserved
stomach contents [71], and study textile industries [72]. The population exhibited
predominantly good health and included individuals of extremely advanced age (50+) for
hunter-gatherer groups which reflects local resource abundance [73] and medical
practices [71, 74-77]. Common to hunter-gatherer populations, adults of both sexes

exhibited a high incidence of osteoarthritis [78], frequent enamel defects [79, 80], and
skeletal trauma [76]. Overall, female health was worse than males [77, 81]
179
2.1 Mobility
The bog was strategically located between the Indian River coastal lagoon system and the
St. John's River, an area rich in marine, fresh water, and terrestrial resources, which
indicates that the population did not have to travel long distances between seasons [74]
and did not fission into smaller groups between visits to the pond [77]. Seasonal mobility
is indicated by analysis of preserved stomach contents which were from plants and fruits
maturing during July and October. In addition, growth ring data recovered from mortuary
stakes indicated the wood was harvested in late summer/early fall [82]. Ultimately,
however, the residential mobility of the Windover population was limited to a
constrained geographic area around the bog with most evidence pointing to emergence of
sedentism [77].
191
2.2 Subsistence economies
Paleodietary analysis from carbon and nitrogen bone-collagen values and
archaeobotanical information suggest exploitation of riparian resources rather than the
more common Florida Archaic use of marine mammals or terrestrial fauna such as deer
or rabbit [71]. Males and females did not have significantly different isotope values for
major dietary components [77, 81]. Based on ethnographic data, the resource rich
environment fostered by a milder wetter climate suggests greater reliance on female
caloric contributions to diet [39]and this is supported by grave goods--both males and
females were found with materials for hunting small mammals, reptiles, and fish [75].
There is evidence of some specialization because only male graves atlatl components,
spears, lithic projectiles, and hollow point awls (for making fishing nets) and only
females and subadult graves contained direct evidence for food processing (e.g.,
butchered bone, mortar and pestle, containers) [72, 75].
205
2.3 Tool economies
Analysis of tools found amongst the grave goods suggests tool material choice (not type)
was sex-based; females preferred organic materials (e.g., shell, carapace) while males
preferred antler bone [74, 75]. Few tool types were exclusive to one sex which suggests
few activities were specific to one sex [75]. Females also tended to have more decorative
items [75] and were found exclusively with materials for plant-based medicine [71, 74,
75]. Interestingly, stone tools only play a cameo in the story of tools at Windover [74]
while female goods (textiles, baskets, containers, medicines) have a starring role across
the history of the pond. The absence of internment with stone tools suggests a cultural
emphasis on the labor of women and products from both men and women in the domestic
economy rather than an emphasis on male big-game hunting [74]. This may be an
outcome of climate change [66, 68] rapidly altering domestic roles from the Paleoindian
to the Early Archaic periods.
219
2.4 Activity Reconstruction
DJD at Windover has been analyzed in two separate studies [78, 83] using different
methods but both finding high rates of DJD consistent with prehistoric hunter-gatherer

lifestyles [78, 83]. DJD frequency in the cervical spine is particularly high in females and
may be explained by food or palm leaf (for textile fibers) processing activities [83] or
carrying heavy loads [78] both of which are supported by grave good evidence [72, 74,
75]. Males exhibit greater DJD in the lumbar region which they were carrying heavy
loads (perhaps game or goods during seasonal camp relocation) or stressed from
repetitive motions related to hide processing [83]. Elbows were commonly affected
which might be interpreted as male atlatl throwing but females exhibited more DJD in
wrist elbows and shoulders than males [83] which suggests a shared activity [78]. Wrists
exhibited little evidence of DJD (3/97 left and right wrists affected) but females had more
hand trauma [83] and males had a higher frequency of severe DJD in both hands (18% of
the sample) [78]. Significant bilateral degeneration of talar-calcaneal articular facets was
noted in 37% of individuals [78], which might reflect high mobility [27], possibly
running [33, 35, 84, 85]. There were some overall sex-based patterns in DJD with males
exhibiting more knee and hip damage on the left and females exhibiting more severe
change on the right [78], which might suggest footedness and increased mobility in males
with more shock to the feet.
239
An examination of muscle insertion sites [86] found low levels of habitual stress
(indicative of stressful repetitive activity) but muscle insertion sites were fairly robust. As
with the DJD results, there is much overlap between the sexes in scores further
supporting the notion that most activities were shared. The lack of asymmetry in MSM,
particularly in males due to the use of the atlatl, suggests a lack of repetition in this
activity or other activities which exert symmetrical force on the upper limbs and obscure
the lateralization of spear-throwing—possibly kayaking [86]. Analysis of fractures
suggests interpersonal violence (affecting male crania and, less frequently, the post
cranial skeleton) but the majority of trauma came from accidents with females slightly
more affected than males. Ribs (often on the right side) were the most fractured in both
sexes with ulnar fractures in second place. The vertebra exhibit evidence of compression
fractures (more frequent in females and equal to ulna fractures in incidence) consistent
with falls when landing in an upright position or carrying heavy loads. Overall, fracture
patterns suggest accidents related to logistical mobility in the uneven intercoastal terrain
and heavy underbrush [76].
255
2.5 Research Questions
1. Mobility
a) Are there differences between weight-bearing and shock absorbing tarsals as
suggested by previous research indicating heavy load carrying and high
mobility or running?
b) Is there asymmetry in tarsals as suggested by asymmetrical DJD in lower
limbs and fracture patterns?
2. Subsistence
a) Is there a between-sex difference in carpal bones based on previous research
showing greater hand trauma and domestic economy production in females?
b) Is there asymmetry in carpal bones reflecting handedness in complex tasks for
making tools and textiles?
268

269 **3. METHODS**
270
Carpal and tarsal elements with standard anatomical reference points intact were included
from adults with well-defined features used in sex assessment (given the focus on sex-
based morphological variation) resulting in a sample of 44 (27 males, 17 females), but
sample size varies by measurement. All measurements were taken on right and left sides,
when available. Length and width were measured on four carpal bones, four tarsal bones,
and the weight-bearing areas of the talus (trochlea) and calcaneus (load arm). Raw data
were explored for significant outliers using Grubb's statistic.
278
279 **Table 1: Measurements**

Element	Measurement	Orientation	Description
Scaphoid	Length	proximal	scaphoid tubercle-lateral most point
	Width	palmar medial	bisection of scaphoid ridge
Capitate	Length	palmer	proximal-distal end
	Width	lateral	thinnest point
Lunate	Length	proximal	proximal-distal end
	Width	proximal	medial-lateral sides
Hamate	Length	lateral	proximal end-distal ridge (between metacarpal facets)
	Width	lateral	most medial to most lateral side of the facet
Calcaneus	Length	lateral	(Steele and Bramblett 1988)
	Width	superior	(Steele and Bramblett 1988)
	Load arm length	superior	(Steele and Bramblett 1988)
	Load arm width	superior	(Steele and Bramblett 1988)
Talus	Length	superior	(Steele and Bramblett 1988)
	Width	superior	(Steele and Bramblett 1988)
	Trochlea tibia length	superior	(Steele and Bramblett 1988)
	Trochlea tibia width	superior	(Steele and Bramblett 1988)
Navicular	Length	distal	medial tuberosity to lateral cuneiform facet
	Width	inferior	between intermediate and medial cuneiform facets
Int. Cuneiform	Length	superior	proximal and distal midpoint
	Width	superior	thickest middle portion

280
There are two major methodological concerns for this size-based morphological study:
sexual dimorphism [87-89] and trait size variation [90, 91]. The inherent difference
between two variables may be confounded by relative difference in size—comparing the
intermediate cuneiform to the talus or comparing male to female. A common resolution
to sexual dimorphism in size-based studies is to take a ratio of two variables, such as
length and width [87-93]. The issue of trait size variation is one common to asymmetry
studies [94-96] and often resolved by scaling data to the natural log [92, 95]. The natural
log of ratio dataset is symmetric, homoscedastic, and retains the original linear scale of
standard deviation to the mean (the spread or variation of the data) [97]. Thus, we have

taken the natural log, $\log_n$, of the absolute value of length-to-width ratio, *L:W*, per bony
element, *x*:
$\log_n|x(L:W)|$
This resulted in a total of four carpal variables (capitate, hamate, lunate, scaphoid) and
six tarsal variables (calcaneus, calcaneus load arm, intermediate cuneiform, navicular,
talus, talus, trochlea tibia). These data are included as Supplementary Tables 1 and 2.
296
Log transformed ratio data were tested for the normal distribution. A two-way (sex by
class) univariate General Liner Model (GLM) assessed differences between tarsal classes
(weight bearing, shock absorbing). A multivariate GLM assessed differences in carpal
bones. A two-way (sides by individual) univariate GLM was used to assess the presence
of directional asymmetry in both tarsal and carpal bones [94-96]. All results were
evaluated relative to confidence intervals, power, and estimated effect size.
303
304 **4. RESULTS**
305
306 Outliers identified by Grubb's statistic were inspected and removed if determined to be
307 data entry error. Extreme values not identified as outliers were retained. Data were
308 transformed to mitigate the confounding effect of trait size and sexual dimorphism as
309 previously described. All variables were normally distributed (Supplementary Table 3).
310 Descriptive data for all variables by sex are found in Supplementary Table 4.
311
Are there differences between weight-bearing and shock absorbing tarsals as suggested
by previous research indicating heavy load carrying and high mobility or running?
Exploratory analysis suggested that females have larger shock-absorbing bones than
males and shock-absorbing bones were larger than weight-bearing bones (Figure 1, left).
The two-way univariate GLM (Levene=001, p=1.00) indicated no significant between-
sex difference or interaction between sex and bone class but the statistical significance
for differences between shock-absorbing and weight-bearing tarsals is high (Table 2,
Figure 1, right).
320

321
322 **Figure 1: Confidence intervals for weight-bearing and shock absorbing mean pooled**
323 **sex (left) and by sex (right)**
324
325 **Table 2: Univariate GLM Analysis of Tarsal Scaled Ratio Data**

							Mean	SE	n	95% CI Lower	Upper
Test	**F**	**df**	**Sig**	**ηp**	**Power**		**Mean**	**SE**	**n**	**Lower**	**Upper**
Sex(S)	1.83	1	0.41	0.65	0.09	M	0.33	0.03	237	0.27	0.39
						F	0.36	0.04	142	0.28	0.44
Class(C)	108.06	1	0.06	0.99	0.59	S	0.45	0.04	129	0.37	0.53
						W	0.25	0.03	250	0.19	0.31
S*C	0.14	1	0.71	0.00	0.07	S-M	0.42	0.05	82	0.33	0.52
						W-M	0.24	0.04	155	0.17	0.52
						S-F	0.47	0.07	47	0.34	0.60
						W-F	0.25	0.05	95	0.16	0.34

327 Is there asymmetry in tarsals as suggested by asymmetrical DJD in lower limbs and
328 fracture patterns? The two-way univariate GLM did not indicate any lateralization of the
329 foot bones, or 'footedness' and no further testing was indicated (Supplementary Table 5).
330

Is there a between-sex difference in carpal bones based on previous research showing
greater hand trauma and domestic economy production in females? Exploratory plots
(Figure 2) show that females have larger carpal ratios than males except for the hamate.
Results of a multivariate GLM suggest potentially significant differences between the
sexes (Pillai's Trace F=2.811, p=0.059, ηp=0.0.398). The univariate individual tests
generated by the multivariate GLM (Table 3) suggest the lunate and scaphoid contribute
most to the potentially significant differences. The visualization of 95% confidence
intervals (Figure 3) indicates that females have greater ranges that encompass most male
values.
340

341
342 **Figure 2: Means plots for carpals**

344 **Table 3: Multivariate GLM Analysis of Carpal Ratio Scaled Data**

								Mean	SE	n	95% CI Lower	Upper
Levene	**Sig**		**F**	**df**	**Sig**	**ηp**	**Power**	**Mean**	**SE**	**n**	**Lower**	**Upper**

								²					
A	3.22	0.09	2.85	1	0.11	0.13	0.36	M	0.60	0.02	28	0.56	0.63
								F	0.65	0.02	21	0.60	0.70
B	2.15	0.16	1.11	1	0.31	0.05	0.17	M	0.28	0.02	21	0.24	0.32
								F	0.25	0.02	14	0.20	0.30
C	4.32	0.05	2.71	1	0.12	0.12	0.35	M	-0.08	0.03	27	-0.15	-0.01
								F	0.01	0.04	20	-0.08	0.10
D	0.22	0.64	1.10	1	0.31	0.05	0.17	M	0.85	0.03	22	0.79	0.92
								F	0.91	0.04	19	0.82	1.00

345 [1]A-Capitate; B-Hamate; C-Lunate; D-Scaphoid
346 [2]M, male; F, female
347

348
349 **Figure 3: Carpal ratio scaled data 95% confidence intervals by sex**

350
Is there asymmetry in carpal bones reflecting handedness in complex tasks for making
tools and textiles? The two-way univariate GLM did not indicate any lateralization of the
hand bones, or 'handedness' and no further testing was indicated (Supplementary Table
5).

5. DISCUSSION

Archaeological data reconstructing the environment surrounding the mortuary pond
suggest that the larger region of the Windover population was rich with riparian and
terrestrial resources due to climate warming at the start of the Early Archaic. As
megafauna distributions increased in latitude, Windover subsistence strategies
increasingly relied less on big-game hunting more on broad spectrum foraging [67, 82].
Grave good analysis and activity reconstructions suggest heavy overlap between the
sexes in domestic economies rather than a strict sex-based division of labor [75-78, 86].

5.1 Tarsal Variation
Based on previous analysis, we speculated whether load carrying might influence
differences between weight-bearing and shock absorbing tarsals. A univariate two ways
(sex by class) GLM indicated no significant between-sex differences or significant
interaction between sex and class (shock-absorbing, weight-bearing). The power for both

these tests is extremely low and confidence interval data suggests there may be a
statistically undetected significant between-sex difference. This potential difference is not
consistent with sexual dimorphism because shock-absorbing bone summary data suggest
higher female mean ratios and a greater range than males while the weight-bearing bone
summary data suggest no differences (same mean, females contained within the male
interval). This female specific pattern is reflected in the results of the pooled sex test for
difference between weight-bearing and shock absorbing bones. Reaching the threshold
for statistical significance was likely prevented by the observed power (59%) which
likely was not high enough to capture the difference. Thus we feel confident in
interpreting these results as biological significant. Females have larger mean ratios for
shock-absorbing bones than males. We expected the reverse in both sexes—greater stress
on weight-bearing tarsals (talus and trochlea, calcaneus and load arm) due to heavy load-
carrying, as evidenced by the increased bilateral degeneration of the talar-calcaneal
articular surfaces [78], cervical DJD in females [78], and presence of lumbar DJD in
males [78]. The rate of fractures attributed to accidents suggested high mobility in a
rough terrain (which certainly describes intercoastal regions) and our results may reflect
this as well.
385
Clinical evidence of the effect on the foot due to barefoot walking and running during
growth and development suggests it lengthens the foot at the expense of width and
flattens the arches [35, 84, 85]—this is true even when controlling for potentially
confounding effects of demographic variation [84]. The navicular is a key stone bone in
the arches of the foot that is impinged during foot strike by the talus and other
cuneiforms. Structurally linking the rear- and mid-foot, it bears the transmission force of
weight during the push-off phase of locomotion and experiences highly localized stress in
middle one-third of the bone, which makes it prone to fracture in highly athletic
individuals [98-101]. Anatomically, the intermediate cuneiform articulates with the
navicular proximally and second metatarsal distally. The second metatarsal-intermediate
cuneiform joint is a highly stable key stone joint with limited mobility. Injury to the joint
occurs via direct force from load applied to the base of the foot or indirect forces from a
longitudinal force applied to a plantarflexed foot [102, 103]:1826, which is particularly
pronounced in barefoot populations [30, 35, 84, 85, 104]. Bone alterations resulting from
these biomechanical stresses might be the source of the higher length to width ratios in
shock-absorbing tarsals observed in this study. We posit that females may have navigated
a wider range of terrains than males (given their preference for shell tools), might have
foraged at greater distances for raw materials (for tools, plant-based medicines, or
textiles), or engaged in faster walking and running speeds to cover greater distances.
Future avenues of research might be to examine the tubers and spurs on the calcaneus to
identify possible differences in locomotion versus weight-bearing activities [27, 33].
407
5.2 Foot Lateralization
Despite evidence suggesting footedness would not be visible [38, there was
bioarchaeological evidence suggesting lateralization in the lower limbs that may have
affected the tarsals. In particular, males had greater DJD on the left with knees and hips
implicated while females had greater DJD on the right {Smith, 2008 #5573}. These
lateral patterns did not translate to differences in the tarsals bones but possible other areas

to consider in future studies would be the hallux due to its involvement in the extreme
plantarflexion associated with barefoot locomotion [85].
416
5.3 Carpal Variation
We postulated there might be between-sex differences in carpal bones due to increased
hand trauma in females [83] and cervical vertebra DJD possibly linked to food and textile
processing activities [72, 74, 83] compared to increased DJD in males [78]. We found
evidence to support activity-based variation between the sexes—females had higher mean
ratios and greater variation than males. These differences cannot be explained by biology
and sexual dimorphism because an analysis of modern humans suggests, that despite
anatomical differences in wrist function and rotation, between-sex variation in carpal
bones is attributable entirely to relative size which can be accounted for through scaling
procedures [105], as we have done. The larger size in females is further evidence that this
defies the pattern expected due to male-female size differences. In all but the capitate, the
range of female values contains all male values and female ratio means exceed males in
all but the hamate.
430
Does the extreme variation in female carpal bone sizes may reflect inter-individual
differences in activity intensity or workloads compared to males? This interpretation is
challenging to other studies that have argued for a shared activity load. As reviewed
previously, cultures in which male contributions to diet dominate tend to emphasize
female labor in domains other than diet (e.g., weaving, cordage) [42]. Perhaps females
were previously engaged in many non-subsistence domestic tasks during the Paleoindian
period, which was characterized by big game hunting [66]. Then, the climate change in
the Early Archaic undermined male contributions to diet as megafauna migrated north
[67, 82] and female activities (including a return to intense foraging) dominated.
440
The absence of stone tools as grave goods is highly uncharacteristic in societies that value
male hunting contributions and the emphasis at Windover on artifacts typically associated
with female labor (e.g., medicine preparation, containers for carrying goods, decorative
items, textiles) lend support to this [74]. Males were associated with tools that suggest
their involvement in this domestic labor and activity reconstructions suggest a heavy
overlap between the sexes in labor [75-78, 86] but there is also evidence for sex-based
division of labor (males and big game hunting, females and plant-based medicine).
Indeed, the Windover population is characterized by obvious skeletal markers for health
care (e.g., post-fracture bone alignment, surviving childhood stressors) [77, 81, 83] and
this care was likely provided by females. Further, the sex-based preference for tool
materials suggests a gendered ideology surrounding tasks [75]. While we cannot infer too
deeply from the site because it is a mortuary pond and only reflects internment ritual, we
can argue that our findings muddy the water a little in terms of the shared load model and
point to some increased burden of activity on females, some specialization within females
to specific tasks, or some female specific activities.
456
5.4 Hand Lateralization
Our final area of inquiry was lateralization in the carpal bones due to previous research
suggesting that hunter-gatherers exhibited strong handedness when engaged in complex

tasks [47, 49, 50]. The complex tool kit at Windover would have provided an avenue for
handedness to be archaeological visible but, neither male nor female carpal bones
exhibited significant directional asymmetry. We might have expected some lateralization
in males due to spear-throwing but the DJD and MSM patterns were shared between the
sexes which suggests that males may not have been regularly engaged in spear-throwing,
or perhaps the markers of spear-throwing were offset by changes in other aspects of wrist
anatomy [106]:238. A previous study on a population with clear handedness identified
only two bones exhibiting directional asymmetry, the lunate and trapezium [107] which
suggests handedness is not likely to be archaeologically visible.
469
470 **6. CONCLUSION**
471
Windover females exhibit greater variation (Supplementary Table 4) in seven out of 10
ratio variables and larger means in eight out of ten variables. Clearly, this is not expected
were sexual dimorphism were to have influenced the results despite our data
transformations to mitigate this effect. Thus, we argue that the observed patterns (in both
descriptive data and statistical tests) indicate marking of the female body by activity more
so than marking of the male body. A previous study found that Windover females
experienced worse health than males [77, 81] and this might be related to their heavier
burden of labor. Even if grave goods suggests a shared load, a few sex-based patterns
exist with males still associated with big game hunting and females associated with
medicine. Further, the raw material for tools is partitioned by sex and would differentiate
mobility patterns associated with acquiring tool materials. In addition, females are found
exclusively with containers and may have dominated container production (including
acquisition of raw materials) or used containers to transport resources to the habitation
site—yet to be identified in the archaeological record.
486
While we might have expected some hand [49, 50] and possibly some foot lateralization
[36, 37], we did not find it even if DJD and MSM patterns suggesting there might be
some. The lack evidence for footedness was expected due its minimal impact on walking
gait [38]. The lack of evidence for handedness is unexpected since modern ethnographic
studies on hunter-gatherers have shown strong handedness during tool and goods
manufacture [49, 50]. But, handedness in these populations overall have been found to be
much less pronounced when compared to modern industrial societies [49]. Although
some archaeologists are concerned by the 'tyranny of ethnography' (Wobst 1978), we
believe predictions about behavior are an essential part of investigating archaeological
populations. Certainly, as demonstrated in this paper, some aspects of behavior are more
archaeologically visible than others.
498
499 Overall, we can conclude that lateralization in the wrist and foot is not archaeologically
500 visible in this population. Further, we conclude that previous assessments of the high
501 value placed on female labor at Windover [74] is embodied and potentially reflects an
502 asymmetric workload or retention of some sex-based activities. Ultimately, the Windover
503 females tell a much more interesting bioarchaeological story than the males.

ACKNOWLEDGEMENTS
We wish to acknowledge Dr. Frank L. Williams (Georgia State University) for
suggesting the frame of the study.

FUNDING STATEMENT
This research did not receive any specific grant from funding agencies in the public,
commercial, or not-for-profit sectors.

<div align="center">

7. REFERENCES

</div>

1.   Frayer DW, Wolpoff MH. Sexual Dimorphism. Annual Review of Anthropology.
1985;14(1):429-73. doi: 10.1146/annurev.an.14.100185.002241.

2.   Frayer DW. Sexual dimorphism and cultural evolution in the Late Pleistocene and
Holocene of Europe. J Hum Evol. 1980;9(5):399-415. doi: https://doi.org/10.1016/0047-
2484(80)90050-0.

3.   Ruff C, Holt B, Trinkaus E. Who's afraid of the big bad Wolff?: "Wolff's law"
and bone functional adaptation. Am J Phys Anthropol. 2006;129(4):484-98. doi:
10.1002/ajpa.20371.

4.   Ruff CB. Structural allometry of the femur and tibia in Hominoidea and Macaca.
Folia Primatol. 1987;48:9–49.

5.   Schaffler MB, Burr DB, Jungers WL, Ruff CB. Structural and mechanical
indicators of limb specialization in primates. Folia Primatologic. 1985;45:61–75.

6.   Ruff CB. Sexual dimorphism in human lower limb bone structure: relationship to
subsistence strategy and sexual division of labor. J Hum Evol. 1987;16:391–416.

7.   Bridges PS. Skeletal biology and behavior in ancient humans. Evolutionary
Anthropology: Issues, News, and Reviews. 1995;4(4):112-20. doi:
10.1002/evan.1360040403.

8.   Bridges PS. Changes in activities with the shift to agriculture in the Southeastern
United States. Current Anthropology. 1989;30:385-94.

9.   Eshed V, Gopher A, Galili E, Hershkovitz I. Musculoskeletal stress markers in
Natufian hunter-gatherers and Neolithic farmers in the Levant: The upper limb. Am J
Phys Anthropol. 2004;123(4):303-15. doi: 10.1002/ajpa.10312.

10.   Berbesque JC, Marlowe FW, Pawn I, Thompson P, Johnson G, Mabulla A. Sex
differences in Hadza hunter-gatherer dental wear: A preliminary report. . Human Nature.
2012; 23:270-82. doi: 10.1007/s12110-012-9145-9.

11.   Estalrrich A, Rosas A. Division of labor by sex and age in Neandertals: an
approach through the study of activity-related dental wear. J Hum Evol. 2015;80:51-63.

<div align="center">

</div>

12.   Binford LR. Willow Smoke and Dogs' Tails: Hunter-Gatherer Settlement
Systems and Archaeological Site Formation. American Antiquity. 1980;45(1):4-20. doi:
10.2307/279653.

13.   Kelly RL. Hunter-gatherer mobility strategies. Journal of Anthropological
Research. 1983;39(3):277-306. doi: 10.1086/jar.39.3.3629672.

14.   Sahlins MD. Notes on the original affluent society. In: Lee RB, DeVore I, editors.
Man the Hunter. Chicago: Aldine; 1968. p. 85-9.

15.   Marlowe FW. Hunter-gatherers and human evolution. Evol Anthr. 2005;14:54-67.

16.   Panter-Brick C. Sexual division of labor: energetic and evolutionary scenarios.
Am J Hum Biol. 2002;14(5):627-40.

17.   Raichlen DA, Pontzer H, Harris JA, Mabulla AZP, Marlowe FW, Josh Snodgrass
J, et al. Physical activity patterns and biomarkers of cardiovascular disease risk in hunter-
gatherers. Am J Hum Biol. 2017;29(2):e22919-n/a. doi: 10.1002/ajhb.22919.

18.   Berbesque JC, Wood BM, Crittenden AN, Mabulla A, Marlowe FW. Eat first,
share later: Hadza hunter–gatherer men consume more while foraging than in central
places. Evolution and Human Behavior. 2016;37(4):281-6. doi:
https://doi.org/10.1016/j.evolhumbehav.2016.01.003.

19.   Shaw CN, Stock JT. Intensity, repetitiveness, and directionality of habitual
adolescent mobility patterns influence the tibial diaphysis morphology of athletes. Am J
Phys Anthropol. 2009;140(1):149-59. doi: 10.1002/ajpa.21064.

20.   Ruff CB. Body size, body shape, and long bone strength in modern humans. J
Hum Evol. 2000;38(2):269-90. doi: https://doi.org/10.1006/jhev.1999.0322.

21.   Pearson Osbjorn M. Activity, Climate, and Postcranial Robusticity: Implications
for Modern Human Origins and Scenarios of Adaptive Change. Current Anthropology.
2000;41(4):569-607. doi: 10.1086/317382.

22.   Bridges PS. Degenerative joint disease in hunter–gatherers and agriculturalists
from the Southeastern United States. Am J Phys Anthropol. 1991;85(4):379-91. doi:
10.1002/ajpa.1330850403.

23.   Malina RM, Little BB. Physical activity: The present in the context of the past.
Am J Hum Biol. 2008;20(4):373-91. doi: 10.1002/ajhb.20772.

24.   Stock JT. Hunter-gatherer postcranial robusticity relative to patterns of mobility,
climatic adaptation, and selection for tissue economy. Am J Phys Anthropol.
2006;131(2):194-204. doi: 10.1002/ajpa.20398.

25. Carlson KJ, Grine FE, Pearson OM. Robusticity and sexual dimorphism in the
postcranium of modern hunter-gatherers from Australia. Am J Phys Anthropol.
2007;134(1):9-23. doi: 10.1002/ajpa.20617.

26. Venkataraman VV, Kraft TS, Desilva JM, Dominy NJ. Phenotypic plasticity of
climbing-related traits in the ankle joint of great apes and rainforest hunter-gatherers.
Hum Biol. 2013;85(1-3):309-28. doi: 10.3378/027.085.0315.

27. Weiss E. Calcaneal spurs: Examining etiology using prehistoric skeletal remains
to understand present day heel pain. The Foot. 2012;22(3):125-9. doi:
https://doi.org/10.1016/j.foot.2012.04.003.

28. Lieverse AR, Weber AW, Bazaliiskiy VI, Goriunova OI, Savel'ev NA.
Osteoarthritis in Siberia's Cis-Baikal: Skeletal indicators of hunter-gatherer adaptation
and cultural change. Am J Phys Anthropol. 2007;132(1):1-16. doi: 10.1002/ajpa.20479.

29. Ruff C. Sexual dimorphism in human lower limb bone structure: relationship to
subsistence strategy and sexual division of labor. J Hum Evol. 1987;16(5):391-416. doi:
https://doi.org/10.1016/0047-2484(87)90069-8.

30. Pontzer H, Suchman K, Raichlen DA, Wood BM, Mabulla AZP, Marlowe FW.
Foot strike patterns and hind limb joint angles during running in Hadza hunter-gatherers.
Journal of Sport and Health Science. 2014;3(2):95-101. doi:
https://doi.org/10.1016/j.jshs.2014.03.010.

31. Trinkaus E, Shang H. Anatomical evidence for the antiquity of human footwear.
Journal of Archaeologial Science. 2008;35:1928–33.

32. Nordin M, Frankel VH. Basic Biomechanics of the Musculoskeletal System. 4th
ed. Baltimore, MD: Lippincott Williams & Wilkins; 2012.

33. Raichlen DA, Armstrong H, Lieberman DE. Calcaneus length determines running
economy: Implications for endurance running performance in modern humans and
Neandertals. J Hum Evol. 2011;60(3):299-308. doi:
https://doi.org/10.1016/j.jhevol.2010.11.002.

34. Hatala KG, Dingwall HL, Wunderlich RE, Richmond BG. Variation in Foot
Strike Patterns during Running among Habitually Barefoot Populations. PLoS One.
2013;8(1):e52548. doi: 10.1371/journal.pone.0052548.

35. Fredericks W, Swank S, Teisberg M, Hampton B, Ridpath L, Hanna JB. Lower
Extremity Biomechanical Relationships with Different Speeds in Traditional, Minimalist,
and Barefoot Footwear. J Sports Sci Med. 2015;14(2):276-83.

36. Gabbard C. Foot Laterality in Children, Adolescents, and Adults. Laterality:
Asymmetries of Body, Brain and Cognition. 1996;1(3):199-206. doi:
10.1080/713754236.

37.   Gentry V, Gabbard C. Foot-Preference Behavior: A Developmental Perspective.
The Journal of General Psychology. 1995;122(1):37-45. doi:
10.1080/00221309.1995.9921220.

38.   Zverev YP. Spatial parameters of walking gait and footedness. Ann Hum Biol.
2006;33(2):161-76. doi: 10.1080/03014460500500222.

39.   Hiatt B. Woman the gatherer. In: Gale F, editor. Woman's Role in an Aboriginal
Society. Canberra: Australian Institute of Aboriginal Studies; 1978. p. 4-15.

40.   Bailey H. A method of determining warmth and temperateness of climate.
Geografiska Annaler. 1960;43:1-16.

41.   Marlowe FW. Hunting and gathering: The human sexual division of foraging
labor. Cross-Cultural Research. 2007;41(2):170–95.

42.   Waguespack N. The organization of male and female labor in foraging societies:
Implications for early Paleoindian archaeology. American Anthropologist. 2005;107:666-
76.

43.   Bentley GR. Hunter-Gatherer Energetics and Fertility: A Reassessment of the
!kung San. Hum Ecol. 1985;13(1):79-109.

44.   Cowlishaw G. The determinants of fertility among Australian Aborigines.
Mankind. 1981;13(1):37-55.

45.   Hurtado A, Hawkes K, Hill K, Kaplan H. Female subsistence strategies among
Ache hunter-gatherers of Eastern Paraguay. Hum Ecol. 1985;13(1):1-28.

46.   Trinkaus E. Neandertal postcrania and the adaptive shift to modern humans. In:
Trinkaus E, editor. The Mousterian legacy: Human biocultural changes in the Upper
Pleistocene. 164. Oxford: British Archaeological Reports International Series; 1983. p.
165–200.

47.   Stock JT, Shirley MK, Sarringhaus LA, Davies TG, Shaw CN. Skeletal evidence
for variable patterns of handedness in chimpanzees, human hunter-gatherers, and recent
British populations. Ann N Y Acad Sci. 2013;1288:86-99. doi: 10.1111/nyas.12067.

48.   Weiss E. Sex differences in humeral bilateral asymmetry in two hunter-gatherer
populations: California Amerinds and British Columbian Amerinds. Am J Phys
Anthropol. 2009;140(1):19-24. doi: 10.1002/ajpa.21025.

49.   Cavanagh T, Berbesque JC, Wood B, Marlowe F. Hadza handedness: Lateralized
behaviors in a contemporary hunter-gatherer population. Evolution and Human Behavior.
2016;37:202-9. doi: http://dx.doi.org/10.1016/j.evolhumbehav.2015.11.002.

50.   Robira B, Pouydebat E, San-Galli A, Meulman EJM, Aubaile F, Breuer T, et al.
Handedness in gestural and manipulative actions in male hunter-gatherer Aka pygmies

from Central African Republic. Am J Phys Anthropol. 2018;10.1002/ajpa.23435:n/a-n/a.
doi: 10.1002/ajpa.23435.

51.   Bridges PS. Vertebral arthritis and physical activities in the prehistoric
southeastern United States. Am J Phys Anthropol. 1994;93:83-93.

52.   Bostanci E. A Biometrical and Morphological Study of the Astragalus and
Calcaneus of the Roman People of Gordium in Anatolia: An Introduction to the
Evolution of the Human Foot. Ankara: Turk Tarih Kurumu Basimevi; 1962.

53.   Bunning P. Some observations on the West African calcaneus and the associated
talo-calceaneal interosseous ligamentous apparatus. Am J Phys Anthropol. 1964;37:373-
88.

54.   Steele DG, McKern TW. A method for assessment of maximum long bone length
and living stature from fragmentary long bones. Am J Phys Anthropol. 1969;31:215-28.

55.   Steele D. The estimation of sex on the basis of the talus and calcaneous. Am J
Phys Anthropol. 1976;45:581-8.

56.   Steele D. The talus and calcaneous: discriminant functions for estimation of sex
among American Whites and Negroes. Lawrence: University of Kansas; 1970.

57.   Harris SM, Case DT. Sexual Dimorphism in the Tarsal Bones: Implications for
Sex Determination. J Forensic Sci. 2012;57(2):295-305. doi: 10.1111/j.1556-
4029.2011.02004.x.

58.   Riepert T, Drechsler T, Schild H, Nafe B, Mattern R. Estimation of sex on the
basis of radiographs of the calcaneus. Forensic Sci Int. 1996;77(3):133-40.

59.   Mastrangelo P, De Luca S, Sanchez-Mejorada G. Sex assessment from carpals
bones: Discriminant function analysis in a contemporary Mexican sample. Forensic Sci
Int. 2011;209(1-3):196.e1-.e15.

60.   Mastrangelo P, Luca S, Alemán I, Botella M. Sex assessment from the carpals
bones: Discriminant function analysis in a 20th century Spanish sample. Forensic Sci Int.
2011;206(1-3):216.e1-.e10.

61.   Kidd RS, Oxnard CE. Patterns of morphological discrimination in selected human
tarsal elements. Am J Phys Anthropol. 2002;117(2):169-81. doi: 10.1002/ajpa.20017.

62.   Gualdi-Russo E. Sex determination from the talus and calcaneus measurements.
Forensic Sci Int. 2007;171(2):151-6. doi: https://doi.org/10.1016/j.forsciint.2006.10.014.

63.   Introna F, Jr., Di Vella G, Campobasso CP, Dragone M. Sex determination by
discriminant analysis of calcanei measurements. J Forensic Sci. 1997;42(4):725-8.

64.   Hoover K. Carpals and Tarsals: Discriminant Functions for the Estimation of Sex.
Tallahassee: The Florida State University; 1997.

65.   Davies CM, Hackman L, Black SM. The foot in forensic human identification – A
review. The Foot. 2014;24(1):31-6. doi: https://doi.org/10.1016/j.foot.2013.12.001.

66.   Milanich JT. Archaeology of Precolumbian Florida. Gainesville, Florida: The
University Press of Florida; 1994.

67.   Halligan JJ, Waters MR, Perrotti A, Owens IJ, Feinberg JM, Bourne MD, et al.
Pre-Clovis occupation 14,550 years ago at the Page-Ladson site, Florida, and the
peopling of the Americas. Science Advances. 2016;2(5). doi: 10.1126/sciadv.1600375.

68.   Doran GH, Dickel DN. Radiometric chronology at the Archaic Windover Site.
Florida Anthropologist. 1988;41:365-80.

69.   Purdy B. The Art and Archaeology of Florida's Wetlands. Boca Raton: CRC
Press; 1991.

70.   Hauswirth WW, Dickel CD, Lawlor DA. DNA analysis of the Windover
population. In: Herrmann B, Hummel S, editors. Ancient DNA: Recovery and Analysis
of Genetic Material from Paleontological, Archaeological, Museum, Medical, and
Forensic Specimens: Springer; 1994. p. 104-21.

71.   Tuross N, Fogel ML, Newsom L, Doran GH. Subsistence in the Florida Archaic:
The Stable-Isotope and Archaeobotanical Evidence from the Windover Site. American
Antiquity. 1994;59(2):288-303. doi: 10.2307/281933.

72.   Adovasio JM, Andrews RL, Hyland DC, Illingworth JS. Perishable industries
from the Windover Bog: An unexpected window into the Florida Archaic. North
American Archaeologist. 2001;22(1):1-90. doi: https://doi.org/10.2190/BX11-VDQY-
LG5N-P94F.

73.   Klingle D. Burial in Florida: Culture, Ritual, Health, and Status: The Archaic to
Seminole Periods. Tallahassee, Florida: The Florida State University; 2006.

74.   Adovasio JM, Soffer O, Page J. The Invisible Sex: Uncovering the True Roles of
Women in Prehistory. New York City: Harper-Collins e-books; 2009.

75.   Hamlin C. Sharing the load: Gender and task division at the Windover Site. In:
Arnold B, Wicker NL, editors. Gender and the Archaeology of Death. AltaMira Press:
Lanham, Maryland; 2001.

76.   Smith RK. The Analysis of Skeletal Fractures From Windover (8Br246) and
Their Inference Regarding Lifestyle. Tallahassee, Florida: The Florida State University;
2003.

77.   Wentz RK. A Bioarchaeological Assessment of Health from Florida's Archaic:
Application of the Western Hemisphere Health Index to the Remains from Windover
(8Br246). Tallahasee, FL: The Florida State University; 2006.

78.   Smith HM. Degenerative Join Disease in the Windover Population. Tallahassee,
Florida: The Florida State University; 2008.

79.   Berbesque JC, Doran GH. Brief communication: physiological stress in the
Florida Archaic-enamel hypoplasia and patterns of developmental insult in early North
American hunter-gatherers. Am J Phys Anthropol. 2008;136(3):351-6. doi:
10.1002/ajpa.20816.

80.   Berbesque JC, Hoover KC. Frequency and developmental timing of linear enamel
hypoplasia defects in Early Archaic Texan hunter-gatherers. PeerJ. 2018;6:e4367. doi:
10.7717/peerj.4367.

81.   Wentz RK, Tucker B, Krigbaum J, Doran GH. Gauging differential health among
the sexes at Windover (8BR246) using the Western Hemisphere Health Index. Mem Inst
Oswaldo Cruz, Rio de Janeiro. 2006;101(II):77-83.

82.   Doran G, Dickel D. Multidisciplinary investigations at the Windover Site. In:
Purdy B, editor. Wet Site Archaeology. Caldwell: Telford Press; 1988.

83.   Wentz RK. Patterns of degenerative joint disease among males and females at
Windover (8BR246) tha their relationship to grave goods. The Florida Anthropologist.
2010;32(1):6-10.

84.   Hollander K, de Villiers JE, Sehner S, Wegscheider K, Braumann K-M, Venter R,
et al. Growing-up (habitually) barefoot influences the development of foot and arch
morphology in children and adolescents. Sci Rep. 2017;7:8079. doi: 10.1038/s41598-
017-07868-4.

85.   Franklin S, Grey MJ, Heneghan N, Bowen L, Li F-X. Barefoot vs common
footwear: A systematic review of the kinematic, kinetic and muscle activity differences
during walking. Gait Posture. 2015;42(3):230-9. doi: 10.1016/j.gaitpost.2015.05.019.

86.   Hagaman K. Activity-Induced Musculoskeletal Stress Marker Analysis of the
Windover Population. Tallahassee, FL: The Florida State University; 2009.

87.   Fairbairn DJ. Allometry for Sexual Size Dimorphism: Pattern and Process in the
Coevolution of Body Size in Males and Females. Annu Rev Ecol Syst. 1997;28(1):659-
87. doi: 10.1146/annurev.ecolsys.28.1.659.

88.   Smith RJ, Cheverud JM. Scaling of sexual dimorphism in body mass: A
phylogenetic analysis of Rensch's Rule in primates. Int J Primatol. 2002;23(5):1095-135.

89.   Jungers WL. Aspects of size and scaling in primate biology with special reference
to the locomotor skeleton. Am J Phys Anthropol. 1984;27(S5):73-97. doi:
10.1002/ajpa.1330270505.

90.   Lewontin RC. On the measurement of relative variability. Syst Zool.
1966;15(2):141-2.

91.   Huxley JS, Tessier G. Terminology of relative growth. Nat Biotechnol.
1936;137:780-1.

92.   Mobb GE, Wood BA. Allometry and sexual dimorphism in the primate
innominate bone. Am J Anat. 1977;150:530-8.

93.   Jungers WL, Falsetti AB, Wall CE. Shape, relative size, and size-adjustments in
morphometrics. Yearbook of Physical Anthropology. 1995;38:137-61.

94.   Palmer AR. Fluctuating Asymmetry Analyses: A Primer. In: Markow TA, editor.
Developmental Instability: Its Origins and Evolutionary Implications. Dordrecht,
Netherlands: Kluwer; 1994. p. 335-64.

95.   Palmer AR, Strobeck C. Fluctuating asymmetry analysis revisited. In: Polak M,
editor. Developmental Instability (DI): Causes and Consequences. Oxford: Oxford
University Press; 2003. p. 279-319.

96.   Palmer AR, Strobeck C. Fluctuating asymmetry: measurement, analysis, patterns.
Annu Rev Ecol Syst. 1986;17:391-421.

97.   Sokal RR, Rohlf FJ. Biometry: The Principles and Practice of Statistics in
Biological Research. 3rd ed. New York: W.H. Freeman and Company; 1995.

98.   Coris EE, Lombardo JA. Tarsal navicular stress fractures. Am Fam Physician.
2003;67(1):85-90.

99.   Khan KM, Brukner PD, Kearney C, Fuller PJ, Bradshaw CJ, Kiss ZS. Tarsal
navicular stress fracture in athletes. Sports Med. 1994;17(1):65-76.

100.   de Clercq PF, Bevernage BD, Leemrijse T. Stress fracture of the navicular bone.
Acta Orthop Belg. 2008;74(6):725-34.

101.   Shakked RJ, Walters EE, O'Malley MJ. Tarsal navicular stress fractures. Curr Rev
Musculoskelet Med. 2017;10:122-30.

102.   Liu GT, Lavery LA, Schenck RC, Lanctot DR, Zhu CF, Athanasiou KA. Human
articular cartilage biomechanics of the second metatarsal intermediate cuneiform joint.
The Journal of Foot and Ankle Surgery. 1997;36(5):367-74. doi:
https://doi.org/10.1016/S1067-2516(97)80039-7.

103.   Rodgers MM. Dynamic biomechanics of the normal foot and ankle during
walking and running. Phys Ther. 1988;68(12):1822-30.

104.   Hollander K, van der Zwaard BC, de Villiers JE, Braumann K-M, Venter R, Zech
A. The effects of being habitually barefoot on foot mechanics and motor performance in
children and adolescents aged 6–18 years: study protocol for a multicenter cross-sectional
study (Barefoot LIFE project). Journal of Foot and Ankle Research. 2016;9(1):36. doi:
10.1186/s13047-016-0166-1.

105.   Crisco JJ, Coburn JC, Moore DC, Upal MA. Carpal bone size and scaling in men
versus in women. The Journal of Hand Surgery. 2005;30(1):35-42. doi:
https://doi.org/10.1016/j.jhsa.2004.08.012.

106.   Maki JM. The Biomechanics of Spear Throwing: An Analysis of the Effects of
Anatomical Variation on Throwing Performance, with Implications for the Fossil Record.
St. Louis: Washington University in St. Louis; 2013.

107.   Reina N, Cavaignac E, Trousdale WH, Laffosse JM, Braga J. Laterality and grip
strength influence hand bone micro-architecture in modern humans, an HRpQCT study. J
Anat. 2017;230(6):796-804.
795