A Signature of Passivity? An Explorative Study of the N3 Event-Related Potential

Component in Passive Oddball Tasks

Boris Kotchoubey [1], Yuri G. Pavlov [1, 2]

[1]Institute of Medical Psychology and Behavioral Neurobiology, University of Tübingen, Germany

[2]Department of Psychology, Ural Federal University, Ekaterinburg, Russian Federation,

pavlovug@gmail.com

Corresponding author:

Boris Kotchoubey

Email: boris.kotchoubey@uni-tuebingen.de

Address: Institute of Medical Psychology and Behavioral Neurobiology, Silcherstr. 5, Tübingen, 72076,

Germany

Phone: +49 7071 29 74221

Short title: N3

Abstract

Background

Many passive oddball experiments show a sharp negative deflection N3 after P3b, peaking between 400

and 500 ms, but this wave has never been analyzed properly. We conducted five passive oddball

experiments, in which the number of deviants (i.e., one or two), their alleged meaning, and their

distinctiveness varied.

Results

Mastoid- or common-referenced waveforms showed a fronto-central N3 in all experiments. The data

were CSD (Current Source Density) transformed and underwent a Principal Component Analysis (PCA).

The PCA revealed N3 containing two subcomponents with very stable peak latencies of about 415 and

455 ms, respectively. Both topography of the subcomponents and their variation with experimental

conditions were very similar, indicating a midfrontal sink and a posterior temporal source. An analysis of

P3a and P3b components replicated previously known effects.

Conclusions

We discuss the similarities and differences between the passive N3 and other components including the

MMN, N1, late positive Slow Wave, and reorienting negativity. We also make general hypotheses about

a possible functional meaning of N3; on this basis, specific hypotheses are formulated and further

experiments are suggested to test these hypotheses.

Key words: Cortical Preparation, Current Source Density, Oddball, Principal Component Analysis (PCA),

Uncertainty

Introduction

More than half a century has passed since we first learned that infrequent stimuli, presented among

highly frequent ones, elicit in the EEG a slow late positive wave with a typical peak latency slightly longer

than 300 ms (Sutton et al., 1965). Already the first systematic review on this effect is almost four

decades old (Donchin, 1981). Later on, this late positivity of the event-related potential (ERP), called

P300 or P3, was found to entail two distinct components, P3a and P3b (e.g., Courchesne et al., 1975).

Moreover, even P3b was shown to be not a unitary wave but consists of further subcomponents (e.g.,

Falkenstein et al., 1994).

The eliciting condition, referred to as the "oddball task" (Duncan-Johnson & Donchin, 1977) standardly

involved an active task to respond to the rare stimuli or to both rare and frequent stimuli. If subjects

have to respond to rare stimuli, the factors of rarity (or probability) and task-relevance are confounded.

This fact underlay the idea to obtain the same effect without any active task requirement (Polich, 1987).

It was shown that such a passive oddball paradigm also yields the P3 ERP component to the rare

stimulus, although its amplitude is considerably smaller than in the active oddball paradigm (Lang et al.,

1997; Polich, 1989). Although the exact component structure of this passive P3 remains unclear, the

data indicate that it is not just a frontal P3a but contains at least a portion of the typical parieto-central

P3b (e.g., Polich and McIsaak, 1994; Bennington and Pollich, 1999; Lang et al., 1997).

The interest to the passive P3 paradigm was particularly stimulated by clinical consideration, specifically

by the idea to use cognitive ERP components for the analysis of cortical functions in behaviorally

unresponsive patients, e.g., in vegetative state, minimally conscious state, in the late stages of

neurodegenerative diseases, or in patients with Guillain-Barré syndrome. As these patients do not have

any behavior, their cognitive abilities can only be examined by means of direct neurophysiological

techniques, and the oddball ERP paradigm appears to be a convenient one. However, for the same

reason of behavioral unresponsiveness the patients cannot be given an overt task – and if a task is

covert (e.g., to count a rare stimulus), we cannot know whether they really perform it.

Several control studies aimed at the development of the passive oddball paradigm for severely brain

damaged patients confirmed the presence of P3b, but the waveforms presented in these studies also

indicated a later negative deflection with a peak latency between 400 and 500 ms. Bostanov &

Kotchoubey (2006) used a continuous wavelet transform and clearly showed this wave in both time and

time-frequency analyzed responses. Erlbeck et al. (2014) and Morlet et al. (2017) directly compared

ERPs in the oddball paradigm (a) with an active instruction to respond to stimuli, (b) with active

distraction of attention away from stimuli, and (c) with mind wandering (Morlet et al., 2017) or without

any task (Erlbeck et al., 2014). Both studies showed a clear negativity following P3 to deviant stimuli in

the last condition, but not in the other two. Unfortunately, the hypotheses these authors had developed

for their oddball experiments concerned the Mismatch Negativity (MMN) and P3 components, thus the

later components were not analyzed. A large, rather short-lasting, fronto-central negative wave after

P3b can be seen clearly in the results obtained in passive oddball conditions with broadly various kinds

of stimulation (e.g., Oades et al., 1995a, 1995b; Potts, 2004; Justen and Herbert, 2016). However, most

of these studies were concentrated on P300 and/or MMN and did not analyze or even notice the post-

P3 negativity. For example, O'Donnell et al. (1992) carefully described their ERPs as consisting of two

negative-positive complexes N1-P2 and N2-P3 but ignored a clear negativity that followed P3; this was

one of the first studies were such negativity can be seen in Figures.

Barry et al. (2006, 2009) were probably the first who explicitly mentioned this ERP component and

termed it N3. However, the principal interest of this group was the relationship between the phase-

locked ERPs and the underlying EEG activity. Therefore, even they did not perform a systematic analysis

of N3 as compared with other ERP components even though they noticed some regularities of this

component. Mueller et al. (2008) carefully investigated the life-span development of oddball ERP

components (including N3) from early childhood to late adulthood. However, they did not clearly

distinguish this rather sharp negative wave with a peak latency between 400-500 ms from the yet later,

very slow component (peaking after 600 ms) that was also negative at frontal leads but inverted its

polarity at parietal leads.

We found N3 accidently in Experiment I described below. This experiment, as well as some other studies

showing a clear N3 (e.g., Justin & Herbert, 2016) contained an oddball related to learning and employed

potentially significant stimuli. Therefore, we tested (i) the learning hypothesis in Experiment II (in which

no learning took place altogether) and (ii) the significance hypothesis in Experiment IV (in which

potentially significant stimuli were presented with increased salience) and in Experiment III (in which

stimuli lost their significance). Also, because all these and many other similar experiments in the

literature were three-stimulus oddballs, in Experiment V we asked (iii) whether a two-stimulus oddball

would also yield N3.

The main aim of this explorative study was to show that N3 is not an occasional observation but a real

and consistent ERP component, whose timing and topography are different from other known

negativities recorded in oddball paradigms. Finally, we hypothesized that N3, being a late wave

following the P3 complex, would demonstrate two important features common with this complex:

firstly, N3 was expected to be larger to deviants than to standards; secondly, it was expected to

decrease with repeated stimulation like P3 does (e.g., Polich, 1987; Polich and McIsaak, 1994).

To our best knowledge, the present study is the first ERP report specifically devoted to N3. For this

reason, the study is largely exploratory. Of course, this is a serious limitation, which restraints the

opportunities to design experiments on the basis of exact hypotheses. The reason is, however, that facts

must be accumulated before precise hypotheses can be formulated.

Methods: General

Participants

Participants of all experiments were healthy individuals between 19 and 35 years, have had no present

or prior diseases of the nervous system or hearing disorders. All of them reported being well aroused

and in a good mood at the beginning of the experiment. None reported the use of any medicaments

during the last weeks before the experiments. They were seated in comfortable chair and asked to close

their eyes during EEG recording and to listen attentively to the stimuli presented through earphones. No

other instruction was given.

The experiments were approved by the Ethical Commission of the University of Tübingen Medical

Faculty. All participants gave their informed consent.

Stimulation

Stimuli were chords each consisting of five harmonic frequencies and lasting for 200 ms. The intensity of

all auditory stimuli was kept about 70 dB SPL. They were presented binaurally by means of pneumatic

earphones (3M E-A-RTONE). All experiments contained a phase of passive oddball, in which one stimulus

was presented frequently (Standard), and another one or two stimuli were rare deviants (Figure 1). Only

the results of this phase (common to all experiments) will be reported. Stimuli were presented with SOA

(onset-to-onset) varying between 950 and 1050 ms. The order of presentation was random except that

one and the same deviant could not be delivered more than twice in a row.

/Figure 1 about here/

Recording

The EEG was recorded using 64 active ActiCHamp electrodes (Easycap GmbH, Herrsching, Germany)

located according to the extended 10-20 system. In addition to the EEG, the vertical and horizontal

electrooculograms were recorded. The impedance was below 25 kOhm. During recording, the EEG was

referenced to Cz, the digitalization rate was 1000 Hz, and the low pass filter was 500 Hz.

Preprocessing

The off-line inspection of the recordings revealed in some traces poor data quality in one or two of the

64 channels. After re-referencing to common average, these channels were replaced with interpolation

of the adjacent electrodes. After this, the data were filtered within a band from 0.1 to 30 Hz. An

Independent Component Analysis (ICA) was employed to separate and remove activity related to ocular

artifacts according the AMICA algorithm (Palmer et al., 2012). The EEG was then broken into segments

starting from 200 ms prior to tone onset and ending 800 ms after tone onset. The segments that still

contained artifacts notwithstanding the preceding ICA correction were dismissed. The ERPs were

averaged in relation to a baseline from -200 ms to 0 ms.

Current Source Density (CSD)

We computed reference-free CSD estimates ($\mu V/cm2$ units; 10-cm head radius) by mean of the spherical

spline surface Laplacian (Perrin et al., 1989) with the following parameters: 50 iterations; m = 4;

smoothing constant $\lambda = 10-5$ (for detailed description of the procedure see Kayser and Tenke, 2006).

CSD transformation helps to avoid problems associated with the choice of a recording reference. The

CSD pattern of sources and sinks creates a clear image of the current generators of ERP components.

This is especially important for auditory ERP when the use of linked-mastoids reference can lead to

cancelling out potentials generated by sources in the temporal (auditory) cortex. Figure 2 shows a

comparison between CSD and two other kinds of reference in Experiment II.

/Figure 2 about here/

Principal Component Analysis

After preprocessing, data were imported into the ERP PCA Toolkit (Dien, 2010). A temporal PCA with a

Promax rotation was conducted to the combined dataset of CSD-ERPs averaged across trials where time

intervals, conditions, electrodes sites and participants were considered as observations and time points

as variables. In order to determine the number of factors to retain, the parallel test (Horn, 1965) was

applied. The covariance matrix and Kaiser normalization were used for the PCA. The same procedure

was applied, first, to a combined dataset of Experiment I to IV and then for Experiment V data

separately. As a result of the parallel test, 55 temporal factors were extracted for rotation for

Experiments I to IV and 44 for Experiment V. The waveforms for each factor were then reconstructed by

multiplying factor scores by their corresponding loadings and standard deviations. CSD of reconstructed

waveforms representing principal components of interest were averaged in 50 ms windows around peak

values (±25 ms) in ROIs, time intervals and conditions. These data were used for statistical calculations.

The first eight principal components (PCs) with eigenvalues > 1 revealed close similarity with visually

detected peaks (see Figure 3 for scalp distribution of the PCs). Particularly, PC1 with the peak latency of

93 ms was interpreted as N1; PC2 (412 ms), as an earlier peak of N3; PC3 (294 ms), as N2b; PC4 (456

ms), as a second subcomponent of N3; PC5 (215 ms), as P3a; PC6 (138 ms), as P2; and PC7 (334 ms), as

P3b. Note that our N2b was recorded between P3a and P3b, while several studies reported N2b

preceding P3a (e.g., Barry and De Blasio, 2015; Getzmann et al., 2015). Therefore, the reader may doubt

whether it is the same component. We share this doubt but retain the name "N2b" because, first, no

clear negative peak between P2 and P3b was manifested in our data; second, we were not interested in

the true nature of N2b and analyzed this negativity only to compare it with N3.

Because we had no prediction about the dynamics of P2, and to avoid unnecessary increase of the

number of analyses, we did not analyze P2. Other components will be reported below with different

number of details: (A) The N3 component will be reported in all details because, firstly, it is the main

object of the study and, secondly, it has not been extensively studied in the literature, see above. (B)

Other negative components N1 and N2b were analyzed not for their own sake but to compare them

with N3. Finally, (C), the positive components P3a and P3b are, in contrast to N3, very well investigated

in various kinds of oddball paradigm. Their analysis in the current study aimed solely at replication of

already known effects.

/Figure 3 about here/

Statistical Analysis

As ERP components in oddball experiments are most typically investigated at the midline, our first

analysis included only midline locations: frontal (mean of AFz, Fz and FCz), central (Cz), centroparietal

(CPz) and parietal (Pz) leads. To explore the hypothesized changes with time, the responses were

averaged separately for the first, second, and third thirds of the whole sequence of 400 stimuli. Each

average included at least 18 (usually 20) deviants of each kind. The three levels of the factor Time will be

referred to as T1 (stimuli 1 – 133), T2 (stimuli 134 – 266), and T3 (stimuli 267 – 400). The repeated-

measures ANOVA included the factors Stimulus (3 levels), Site (4 levels), and Time (3 levels).

To analyze the activity over the hemispheres, five regions of interests (ROIs) were selected for each

hemisphere: frontal (Fro), frontocentral (FC), central (Ce), posterior (Pos), and temporoparietal (TP), as

shown in Figure 4. The corresponding repeated-measures ANOVA included the factors Stimulus (3

levels), Area (i.e., ROI, 5 levels), Hemisphere (2 levels), and Time (3 levels).

/Figure 4 about here/

The large total number of analyses might result in false positive effects (e.g., Luck & Gaspelin, 2017), and

to date no exact method of correction exists. As a first approximation, we regarded the obtained

Principal Components (PCs) as independent repetitions. Because in each experiment we analyzed six

PCs, we shall generally report effects as "statistically significant" when $p < 0.05/6 = 0.0083$. However,

the effects with $.05 > p > .0083$ will also be reported if they are repeatedly presented in similar

conditions.

Experiment I

Methods

Twenty-three subjects (14 females) aged 19 to 33 (mean 25.2) participated in Experiment I. The

experiment entailed two phases: Acquisition and Test. Chord 1, consisting of the frequencies 200, 400,

800, 1600, and 3200 Hz, was paired 21 times in the Acquisition phase in subjects with odd numbers with

the own name of the corresponding participant (SON). Chord 2 consisted of 130, 260, 520, 1040, and

2080 Hz and was randomly paired 21 times with three other familiar names (OFN). Chord 3, consisting of

330, 660, 1320, 2640, and 5280 Hz, was presented 21 times without any relation to other stimuli. In

subjects with even numbers, Chord 1 was paired with OFN, and Chord 2 with SON. The order of

presentation was randomized but neither Chord 1 nor Chord 2 was allowed to appear more than 3 times

in a row. The SOA within a pair chord-name was 300 ms. The interval after a pair was 1700-1800 ms, and

the SOA after Chord 3 was 1150-1250 ms. Preliminary testing with eight healthy individuals showed that

none of them had any difficulty to distinguish between the tones.

The average duration of the own name and the other names was 669 ms (SD = 9 ms) and 676 ms (SD =

12 ms) respectively (t = 0.78, p = .44). Other names originated from the same pool of the most frequent

German names used for each subject's own name, and always contained the same number of syllables

as the own name.

In the Test phase (passive oddball) Chord 1 was presented 280 times, and Chord 2 and Chord 3, 60 times

each. Name stimuli were not presented any longer. In other words, Test phase was a typical three-

stimulus oddball paradigm with Chord 3 as a standard and Chords 1 and 2 as deviants.

Results

Figure 5 presents the overview of the obtained ERP waveforms. Table 1 shows the main statistical

results across components and experiments for midline recording sites. The ANOVA results for lateral

electrode sites are shown in Table 2, and the statistics obtained in each ROI are summarized in Table 3.

In the following text only sporadic statistical results are explained that are not resumed in the Tables.

/Figure 5 about here/

N3, the main object of the present study, revealed in the PCA as entailing two principal components,

PC2 and **PC4**, with peak latencies of 412 and 456 ms, respectively. Both PCs were negative at frontal

sites and positive at Pz, resulting in a main effect of Site. A similar anterior-posterior negativity-positivity

slope was also significant in lateral sites. In addition, strong Stimulus x Area interactions indicated that

this slope was much stronger for deviants than standards, thus demonstrating an oddball effect (see

Table 3).

The component **N1 (PC1)** had a frontal maximum at the midline (main effect of Site, see Table 1). Lateral

amplitudes were characterized by strong regional differences with positive values over TP areas and

negative values in all other areas (main effect of Area) and more negative amplitudes in response to

deviants than standards (main effect of Stimulus). This latter general effect was, however, inverted over

TP areas where the amplitudes were more positive to deviants than standards, resulting in significant

Stimulus x Area interactions.

Over posterior areas, where no stimulus effect was found, the amplitudes were more negative on the

left than right side (t(22) = 3.56, p < .001, η^2 = .47). The opposite holds true for TP areas (t(22) = 2.91, p

= .008, η^2 = .40), yielding a significant Area x Hemisphere interaction.

N2b (PC3) was characterized by positive values in response to deviants but negative to standards,

resulting in a main effect of Stimulus at midline. This effect was modified by Time, as Chord 2 elicited

positive PC3/N2b values at T3 like the standard. Also at the lateral sites the same effects of Stimulus

(i.e., positivity to deviants, negativity to standards) and topographic differences (i.e., significantly

positive amplitudes in the Pos and TP areas, and slightly negative amplitudes in the Fro and FC areas)

were found.

P3a (**PC5)** was larger to deviants than standards (main Stimulus effect). This difference had its maximum

at Cz and minimum at Pz (Stimulus x Site interaction). Moreover, the amplitude was generally largest at

Cz (main effect of Site). Further, the amplitude of P3a habituated from T1 to T2 and T3 (i.e., main effect

of Time).

Over the lateral areas P3a to standards increased in the posterior direction and reached its maximum in

TP areas. The amplitude to deviants was, in contrast, maximal in FC areas and minimal in TP areas,

resulting in a strong Stimulus x Area interaction (see Table 2). In general, the amplitude was more

positive to deviants than standards (main effect of Stimulus). In TP areas the amplitude (particularly in

response to standards) was more positive on the right than left side, which manifested itself in the

interaction between Area and Hemisphere.

The amplitude of **P3b (PC7)** was larger to deviants than standards at Pz (Stimulus x Site interaction in the

midline ANOVA) and at the lateral sites (main effect of Stimulus in the lateral ANOVA). The maximal

positive deviant-standard difference was found in Pos and TP areas, whereas in Fro and FC areas the

amplitudes were even more negative to deviants than standards. Anterior negativities were larger on

the right than left side (Area x Hemisphere interaction: $F(4,88) = 3.15$, $p = .036$, $\eta^2 = .13$).

Table 1. Results of ANOVA Site x Stimulus x Time in midline electrodes

	Experiment I	Experiment II	Experiment III	Experiment IV	Experiment V
N3 early					
Site	$F(3,66) = 5.77$, p = .015, $\eta^2 = .21$	$F(3,57) = 17.73$, $p < .001$, $\eta^2 = .48$	$F(3,60) = 21.87$, $p < .001$, $\eta^2 = .52$	$F(3,63) = 20.45$, $p < .001$, $\eta^2 = .49$	$F(3,63) = 42.94$, $p < .001$, $\eta^2 = .67$
Stimulus		$F(2,38) = 6.53$, $p = .004$, $\eta^2 = .26$			
Stimulus:Site		$F(6,114) = 2.85$, $p = .039$, $\eta^2 = .13$	$F(6,120) = 6.55$, $p = .001$, $\eta^2 = .25$		$F(3,63) = 15.39$, $p < .001$, $\eta^2 = .42$
Stimulus:Time				$F(6,126) = 4.38$, $p = .003$, $\eta^2 = .17$	
N3 late					
Site	$F(2,44) = 14.53$, $p < .001$, $\eta^2 = .42$	$F(3,57) = 14.82$, $p < .001$, $\eta^2 = .44$	$F(3,60) = 17.76$, $p < .001$, $\eta^2 = .47$	$F(3,63) = 36.25$, $p < .001$, $\eta^2 = .63$	$F(3,63) = 44.29$, $p < .001$, $\eta^2 = .68$
Stimulus:Site		$F(6,114) = 2.94$, $p = .025$, $\eta^2 = .12$	$F(6,120) = 3.20$, $p = .026$, $\eta^2 = .15$		$F(3,63) = 5.98$, $p = .004$, $\eta^2 = .22$
N1					
Site	$F(3,66) = 6.70$, $p = .006$, $\eta^2 = .23$	$F(3,57) = 5.99$, $p = .006$, $\eta^2 = .24$	$F(3,60) = 8.54$, $p = .004$, $\eta^2 = .30$	$F(3,63) = 34.17$, $p < .001$, $\eta^2 = .62$	$F(3,63) = 18.50$, $p < .001$, $\eta^2 = .47$
Stimulus			$F(2,40) = 5.87$, $p = .006$, $\eta^2 = .23$		
Stimulus:Site					$F(3,63) = 7.04$, $p = .003$, $\eta^2 = .25$
N2b					
Site	$F(3,66) = 3.89$, $p = .028$, $\eta^2 = .15$		$F(3,60) = 4.98$, $p = .010$, $\eta^2 = .20$	$F(3,63) = 19.86$, $p < .001$, $\eta^2 = .46$	$F(3,63) = 9.73$, $p = .001$, $\eta^2 = .32$
Stimulus		$F(2,38) = 21.63$, $p < .001$, $\eta^2 = .53$	$F(2,40) = 5.79$, $p = .009$, $\eta^2 = .23$	$F(2,42) = 8.98$, $p = .001$, $\eta^2 = .30$	$F(1,21) = 19.21$, $p < .001$, $\eta^2 = .48$
Stimulus:Site				$F(6,126) = 6.03$, $p < .001$, $\eta^2 = .22$	
Stimulus:Time	$F(4,88) = 4.27$, $p = .008$, $\eta^2 = .16$				

P3a					
Site	$F_{(2,44)} = 9.57$, $p < .001$, $\eta^2 = .30$	$F_{(3,57)} = 4.53$, $p = .025$, $\eta^2 = .19$			
Stimulus	$F_{(2,44)} = 15.83$, $p < .001$, $\eta^2 = .42$	$F_{(2,38)} = 8.86$, $p = .002$, $\eta^2 = .32$	$F_{(2,40)} = 20.22$, $p < .001$, $\eta^2 = .50$	$F_{(2,42)} = 37.80$, $p < .001$, $\eta^2 = .60$	$F_{(1,21)} = 23.00$, $p < .001$, $\eta^2 = .52$
Time	$F_{(2,44)} = 3.88$, $p = .049$, $\eta^2 = .15$		$F_{(2,40)} = 5.52$, $p = .009$, $\eta^2 = .22$	$F_{(2,42)} = 10.55$, $p < .001$, $\eta^2 = .33$	$F_{(2,42)} = 3.55$, $p = .039$, $\eta^2 = .15$
Stimulus:Site	$F_{(6,132)} = 3.77$, $p = .020$, $\eta^2 = .15$	$F_{(6,114)} = 6.06$, $p = .004$, $\eta^2 = .24$	$F_{(6,120)} = 5.17$, $p = .001$, $\eta^2 = .21$	$F_{(6,126)} = 9.39$, $p < .001$, $\eta^2 = .31$	$F_{(3,63)} = 13.01$, $p < .001$, $\eta^2 = .39$
P3b					
Site			$F_{(3,60)} = 9.44$, $p < .001$, $\eta^2 = .33$	$F_{(3,63)} = 9.81$, $p = .001$, $\eta^2 = .32$	
Stimulus					$F_{(1,21)} = 9.12$, $p = .007$, $\eta^2 = .30$
Time					$F_{(2,42)} = 4.23$, $p = .022$, $\eta^2 = .17$
Stimulus:Time				$F_{(4,84)} = 3.16$, $p = .033$, $\eta^2 = .13$	
Stimulus:Site	$F_{(6,132)} = 3.36$, $p = .020$, $\eta^2 = .13$		$F_{(6,120)} = 2.92$, $p = .037$, $\eta^2 = .13$		$F_{(3,63)} = 3.34$, $p = .045$, $\eta^2 = .14$

270   Table 2 – Results of ANOVA Area x Hemisphere x Stimulus x Time in lateral electrodes

	Experiment I	Experiment II	Experiment III	Experiment IV	Experiment V
N3 early					
Area	$F(4,88) = 36.26$, $p < .001$, $\eta^2 = .62$	$F(4,76) = 48.48$, $p < .001$, $\eta^2 = .72$	$F(4,80) = 45.48$, $p < .001$, $\eta^2 = .70$	$F(4,84) = 39.54$, $p < .001$, $\eta^2 = .65$	$F(4,84) = 66.25$, $p < .001$, $\eta^2 = .76$
Stimulus		$F(2,38) = 4.56$, $p = .018$, $\eta^2 = .19$		$F(2,42) = 4.41$, $p = .018$, $\eta^2 = .17$	
Stimulus:Area	$F(8,176) = 5.34$, $p = .004$, $\eta^2 = .20$	$F(8,152) = 5.90$ $p < .001$, $\eta^2 = .24$	$F(8,160) = 6.55$, $p = .001$, $\eta^2 = .25$	$F(8,168) = 5.09$, $p = .001$, $\eta^2 = .19$	$F(4,84) = 21.88$, $p < .001$, $\eta^2 = .51$
N3 late					
Area	$F(4,88) = 48.25$, $p < .001$, $\eta^2 = .69$	$F(4,76) = 27.34$, $p < .001$, $\eta^2 = .59$	$F(4,80) = 28.32$, $p < .001$ $\eta^2 = .59$	$F(4,84) = 45.84$, $p < .001$, $\eta^2 = .69$	$F(4,84) = 59.92$, $p < .001$, $\eta^2 = .74$
Stimulus:Area	$F(8,176) = 7.08$, $p < .001$, $\eta^2 = .24$	$F(8,152) = 3.54$, $p = .005$, $\eta^2 = .16$	$F(8,160) = 4.12$, $p = .003$, $\eta^2 = .17$	$F(8,168) = 5.81$, $p = .001$, $\eta^2 = .22$	$F(4,84) = 7.24$, $p < .001$, $\eta^2 = .26$
N1					
Area	$F(4,88) = 38.78$, $p < .001$, $\eta^2 = .64$	$F(4,76) = 28.85$, $p < .001$, $\eta^2 = .60$	$F(4,80) = 24.65$, $p < .001$, $\eta^2 = .55$	$F(4,84) = 59.46$, $p < .001$, $\eta^2 = .74$	$F(4,84) = 50.00$, $p < .001$, $\eta^2 = .70$
Stimulus	$F(2,44) = 33.72$, $p < .001$, $\eta^2 = .61$	$F(2,38) = 69.35$, $p < .001$, $\eta^2 = .79$	$F(2,40) = 31.20$, $p < .001$, $\eta^2 = .61$	$F(2,42) = 28.25$, $p < .001$, $\eta^2 = .64$	$F(1,21) = 73.22$, $p < .001$, $\eta^2 = .78$
Stimulus: Area	$F(8,176) = 11.48$, $p < .001$, $\eta^2 = .34$	$F(8,152) = 9.78$, $p < .001$, $\eta^2 = .34$	$F(8,160) = 10.07$, $p < .001$, $\eta^2 = .34$	$F(8,168) = 6.48$, $p < .001$, $\eta^2 = .24$	$F(4,84) = 34.15$, $p < .001$, $\eta^2 = .62$
Area:Hemisphere	$F(4,88) = 4.53$, $p = .007$, $\eta^2 = .17$				
N2b					
Area	$F(4,88) = 27.79$, $p < .001$, $\eta^2 = .56$	$F(4,76) = 10.12$, $p = .001$, $\eta^2 = .35$	$F(4,80) = 17.38$, $p < .001$, $\eta^2 = .46$	$F(4,84) = 37.84$, $p < .001$, $\eta^2 = .64$	$F(4,84) = 50.31$, $p < .001$, $\eta^2 = .71$
Stimulus	$F(2,44) = 20.90$, $p < .001$, $\eta^2 = .49$	$F(2,38) = 24.50$, $p < .001$, $\eta^2 = .56$	$F(2,40) = 27.21$, $p < .001$, $\eta^2 = .58$	$F(2,42) = 6.18$, $p = .005$, $\eta^2 = .22$	$F(1,21) = 45.47$, $p < .001$, $\eta^2 = .68$
P3a					
Area					$F(4,84) = 6.02$, $p =. 005$, $\eta^2 = .22$
Stimulus	$F(2,44) = 6.43$, $p = .008$, $\eta^2 = .23$				

Area:Hemisphere	$F(4,76) = 3.33$, $p = .029$, $\eta^2 = .15$		$F(4,80) = 4.68$, $p = .008$, $\eta^2 = .19$	$F(4,84) = 3.31$, $p = .026$, $\eta^2 = .14$	
Stimulus: Area	$F(8,146) = 14.13$, $p < .001$, $\eta^2 = .39$	$F(8,152) = 6.28$, $p < .001$, $\eta^2 = .25$	$F(8,160) = 6.70$, $p < .001$, $\eta^2 = .25$	$F(8,168) = 14.06$, $p < .001$, $\eta^2 = .40$	$F(4,84) = 9.13$, $p <. 001$, $\eta^2 = .30$
P3b					
Area	$F(4,88) = 17.77$, $p < .001$, $\eta^2 = .45$	$F(4,76) = 6.82$, $p = .001$, $\eta^2 = .26$	$F(4,80) = 16.58$, $p < .001$, $\eta^2 = .45$	$F(4,84) = 19.55$, $p < .001$, $\eta^2 = .48$	$F(4,84) = 13.97$, $p <. 001$, $\eta^2 = .38$
Stimulus	*$F(2,44) = 12.04$, $p < .001$, $\eta^2 = .35$*	$F(2,38) = 5.99$, $p = .011$, $\eta^2 = .24$	$F(2,40) = 10.18$, $p = .001$, $\eta^2 = .34$	$F(2,42) = 3.47$, $p = .048$, $\eta^2 = .14$	$F(1,21) = 35.33$, $p <. 001$, $\eta^2 = .63$
Time	$F(2,44) = 9.56$, $p < .001$, $\eta^2 = .30$		$F(2,40) = 3.97$, $p = .042$, $\eta^2 = .17$	$F(2,42) = 4.58$, $p = .021$, $\eta^2 = .18$	
Time:Area	$F(8,176) = 3.12$, $p = .013$, $\eta^2 = .12$	$F(8,152) = 3.55$, $p = .023$, $\eta^2 = .16$			
Stimulus: Area	$F(8,176) = 5.76$, $p = .001$, $\eta^2 = .21$	$F(8,152) = 3.75$, $p = .009$, $\eta^2 = .16$	$F(8,160) = 3.11$, $p = .018$, $\eta^2 = .13$	$F(8,168) = 6.88$, $p < .001$, $\eta^2 = .25$	$F(4,84) = 6.57$, $p =. 003$, $\eta^2 = .24$

Discussion

In addition to the expected oddball effects on P3a and P3b, the experiment showed a robust N3 wave

following P3b. Also this wave was characterized by a strong oddball effect, and its spatial distribution

was different from other negativities (N1, N2b).

The experiment employed a passive oddball paradigm with potentially significant (previously associated

with SON and OFN) stimuli. An obvious hypothesis (which could also be supported by similar data from

the literature, see Introduction), was that N3 is related to the significance of deviants or to the learning

process preceding oddball. On this basis the following three experiments were designed. Specifically,

Experiment III was designed to reduce the significance, and Experiment IV was, in contrast, expected to

enhance the salience of the stimuli (thus enhancing the ERP effects). Experiment II was expected to

eliminate preceding learning.

284 Table 3 – Analysis of Area x Stimulus interactions for negative ERP components at lateral sites: ANOVA
285 results

Area	Experiment I	Experiment II	Experiment III	Experiment IV	Experiment V
N3 early					
Fro D—		$F(2,38) = 4.34$, p = .021, η^2 = .22;	$F(2,40) = 3.99$, p = .036, η^2 = .14	$F(2,42) = 9.87$, p < .001, η^2 = .32	$F(1,21) = 25.47$, p < .001, η^2 = .52
FC D—	$F(2,44) = 10.49$, p < .001, η^2 = .32	$F(2,38) = 4.47$, p = .030, η^2 = .23	$F(2,40) = 4.51$, p = .018, η^2 = .18	$F(2,42) = 11.59$, p < .001, η^2 = .36	$F(1,21) = 28.63$, p < .001, η^2 = .58
Ce D—				.	
TP D+		(F(2,38) = 4.31, p = .020, η^2 = .18	$F(2,40) = 3.74$, p = .045, η^2 = .16		$F(1,21) = 11.71$, p = .003, η^2 = .36
N3 late					
Fro D—	$F(2,44) = 11.23$, p < .001, η^2 = .34	$F(2,38) = 3.40$, p = .011, η^2 = .18	$F(2,40) = 8.24$, p = .003, η^2 = .48	$F(2,42) = 12.40$, p < .001, η^2 = .37	$F(1,21) = 10.09$, p = .005, η^2 = .32
FC D—	$F(2,44) = 8.11$, p = .001, η^2 = .27	$F(2,38) = 13.32$, p < .001, η^2 = .41		$F(2,42) = 5.90$, p = .006, η^2 = .22	$F(1,21) = 5.27$, p = .032, η^2 = .20
Ce D—	$F(2,44) = 6.51$, p = .004, η^2 = .23 (§)	$F(2,38) = 6.70$, p = .003, η^2 = .26			
TP D+	$F(2,44) = 9.43$, p = .001, η^2 = .30	$F(2,38) = 10.87$, p < .001, η^2 = .26		.	
N1					
Fro D—	$F(2,44) = 16.53$, p < .001, η^2 = .43	$F(2,38) = 16.45$, p < .001, η^2 = .46	$F(2,40) = 7.44$, p = .013, η^2 = .27	$F(2,42) = 19.21$, p < .001, η^2 = .48	
FC D—	$F(2,44) = 20.95$, p < .001, η^2 = .49	$F(2,38) = 37.82$, p < .001, η^2 = .67		$F(2,42) = = 13.42$, p < .001, η^2 = .39	$F(1,21) = 11.73$, p = .003, η^2 = .36
Ce D—	$F(2,44) = 31.19$, p < .001, η^2 = .59	$F(2,38) = 20.00$, p < .001, η^2 = .51	$F(2,40) = 8.38$, p = .001, η^2 = .30 (*)	$F(2,42) = 11.03$, p < .001, η^2 = .34	$F(1,21) = 52.45$, p < .001, η^2 = .69
TP D+	$F(2,44) = 11.27$, p < .001, η^2 = .35	$F(2,38) = 9.63$, p < .001, η^2 = .34		$F(2,42) = 4.80$, p = .021, η^2 = .19	$F(1,21) = 13.42$, p = .001, η^2 = .39

The table presents main effects of Stimulus at single areas when the Stimulus x Area interaction was
significant. Fro, frontal areas; FC, fronto-central areas; Ce, central areas; TP, lateral temporoparietal
areas. Posterior areas are omitted because no effect attained p ≤ .05 there. D— means that the
amplitude was more negative to deviants than standards. D+ means that the amplitude was more
positive to deviants than to standards. Empty cells mean that, although Stimulus x Area interactions
were significant, the effect of Stimulus in a given area was not.
(§) Chord 2 more negative than Chord 1: t(22) = 2.90, p = .008, η^2 = .28
(*) This was a Stimulus x Time interaction with deviants being significantly more negative than
standards at time intervals T2 and T3

Experiment II

Methods

Experiments II, III, and IV were carried out with the same participants, always in the same order. The

resulting samples were not identical, however. Specifically, the instruction to close eyes resulted in the

almost complete absence of ocular activity. On the other hand, two participants exhibited signs of sleep

stages N1 or even N2 (AASM sleep criteria) in their EEG, and their data had to be dismissed. The data of

seven males and 13 females, aged 22 to 29 (mean 25.6) were analyzed. None of the participants of

Experiments II to IV participated in Experiments I or V.

The design was fully identical to the Test phase of Experiment I. However, no Acquisition phase

preceded it. The only instruction was to keep the eyes closed and to listen to stimuli.

Results

In addition to the Site effect like in Experiment I, the amplitude of both **N3** subcomponents was more

negative to deviants than standards at frontal sites (F(2,40) = 6.46, p = .004, η^2 = .24, 3.95, p = .032, η^2 =

.17 for PC2 and PC4, respectively), and more positive to deviants than standards at Pz (F(2,40) = 4.51, p =

.023, η^2 = .18, and 3.33, p = .051, η^2 = .15 for PC2 and PC4, respectively). At the lateral areas, strong

effects of Area and Stimulus x Area interactions indicated the same differential effects of standards and

deviants as in Experiment I above. The exact statistical results are shown in Tables 1 – 3.

The effects on **N1** were almost identical to those in Experiment I. Regarding **N2b**, additionally to the

effects like those in Experiment I, lateral frontal and fronto-central negativities significantly decreased

with Time (main effect of Time: F(4,76) = 10.12, p = .001, η^2 = .35).

Also the effects on the positive components were similar to Experiment I with the following differences:

the Area x Hemisphere interaction for P3a was not significant; the oddball effect at the midline for P3b

(i.e., the Stimulus x Site interaction) was not significant. The Area x Hemisphere interaction for P3b was,

again, significant: $F(4,76) = 5.00$, $p = .004$, $\eta^2 = .21$.

Experiment III

Methods

After exclusion of one data set for technical reasons, data of twenty-one subjects (12 females, aged 22 –

29, mean 25.7) were analyzed.

The design was identical to Experiment I including both Acquisition and Test phases. However, the

words presented during Acquisition were made completely unrecognizable at the same time preserving

all auditory features that SON and OFN had in Experiment I. Note that a simple reversion of the audio

track would not be enough for this sake because correctly pronounced and inverted names have quite

different dependence of intensity on time. Therefore, a more complex masking procedure was

employed. First, the first 25% of time points of an original name were multiplied by a linearly spaced

vector of coefficients from 1.5 to 0, and the remaining 75% points were set to 0. Second, the first 25% of

time points of the same name played backwards were multiplied by a linearly spaced vector of

coefficients from 0 to 1.5, and the last 75% time points remained unchanged. Third, the two files were

added. The processing was done in Matlab. A pilot experiment with 40 medical students (not

participating in the EEG experiments) revealed than none of them was able to recognize the presented

words, including each participant' masked own name.

Results

The effects on **N3** at midline were similar to Experiment I. PC2 was again more negative to deviants than

standards at frontal sites ($F(2,40) = 6.46$, $p = .004$, $\eta^2 = .24$), and this effect was inverted at Pz ($F(2,40) =$

4.51, $p = .023$, $\eta^2 = .18$). PC4 was more negative to deviants than standards at Cz ($F(2,40) = 4.11$, $p =$

.024, $\eta^2 = .17$) with a similar non-significant frontal tendency ($F(2,40) = 3.12$, $p = .057$, $\eta^2 = .14$) and the

lack of Stimulus effect at CPz and Pz. Lateral effects were identical for PC2 and PC4, i.e., the amplitudes

were significantly negative (and more negative to deviants than standards) over the Fro and FC areas,

and significantly positive (and more positive to deviants than standards) over the TP areas (see Tables 2

and 3).

The amplitude of **N1** was significantly larger to Chord 1 (previously linked with masked SON) than Chord

2 (previously linked with masked OFN), while the standard stimulus elicited an intermediate amplitude

between the two deviants. At the lateral sites, deviants elicited larger negativities in frontal areas

($F(1,20 = 7.44$, $p = .013$, $\eta^2 = .27$) and in central areas at T2 and T3 (Stimulus x Time interaction with

$F(2,40) = 8.38$, $p = .001$, $\eta^2 = .30$). The standard-deviant differences in other areas were not significant.

The effects on **N2b** were quite like Experiment II, including the decrease of lateral negativities with Time

($F(2,40) = 4.48$, $p = .020$, $\eta^2 = .18$) that was lacking in Experiment I. The effects on **P3a** and **P3b** were

similar to Experiment I including the larger anterior negativities on the right than left side (Area x

Hemisphere interaction: $F(4,80) = 3.70$, $p = .016$, $\eta^2 = .16$).

Discussion

Contrary to our hypothesis, changes in N3 in Experiments II and III were almost the same as in

Experiment I although the significance of the eliciting stimuli substantially decreased. In Experiment III

the tones had previously been associated not with participants' own names and other familiar names (as

in Experiment I) but with unrecognizable acoustical sequences. In Experiment II the tones were

completely neutral. Their significance was so low that two of the 22 participants felt asleep, and the

others complained of boredom. In accordance with these observations, participants' reports after

Experiments II and III were not exact, as many of them reported having heard "two or three" different

sounds. Even though the same effects in Experiment II were, generally, weaker than in Experiment I, this

fact can most plausibly be attributed to the lower alertness but not to some specific effects of learning.

The idea of Experiment IV was, in contrast to Experiments II and III, to increase the (possibly insufficient)

salience of auditory stimuli. This was done by means of (i) monaural presentation, and (ii), partial

reinforcement. In addition, we wanted to rule out the supposition that the effects might be related with

the particular set of stimuli used in Experiments I to III.

Experiment IV

Methods

Participants were 13 females and 9 males, aged 22 – 29, mean 25.7. The design was similar to

Experiment I with three differences. First, different tones were used. Chord 1 (associated with SON)

contained the harmonic components of 300, 600, 120, 2400, and 4800 Hz; chord 2 (associated with

OFN) contained the harmonics of 195, 390, 780, 1560, and 3120 Hz; and the standard contained the

components of 495, 990, 1980, 3960, and 7920 Hz. Second, Chords 1 and 2 were presented monaurally,

one to the left ear and the other one to the right ear, with the side of presentation being

counterbalanced among participants. Third, the Test phase included partial reinforcement. The tone

related to SON in the Acquisition phase was also followed by SON on randomly selected nine (of the 60)

trials in the Test phase; on the remaining 51 trials the tone was presented alone like in Experiment I,

Likewise, the tone related to OFN in the Acquisition phase was also followed by OFN on 9 trials in the

Test phase. The partial reinforcement aimed at the refreshment of the association between tones and

names.

 Results

Only the main Site effect on **N3** at midline replicated the corresponding effect in the previous

experiments, but Stimulus x Site interactions did not attain the corrected significance level of .0083. PC2

revealed, in addition, a significant Site x Time interaction, indicating the increasing positivity at Pz at T3

as compared with T1 and T2 (main Time effect at Pz: $F(2,42) = 3.46$, $p = .038$, $\eta^2 = .15$). Interestingly, the

Stimulus x Site interaction for PC2 was significant when only two deviant stimuli were compared ($F(3,63)$

$= 6.09$, $p = .006$, $\eta^2 = .26$) indicating a significantly more negative frontal amplitude to the OFN-

associated tone than to the SON-associated tone ($t(21) = 2.57$, $p = .018$, $\eta^2 = .24$).

In the lateral areas the effects of Area and Stimulus x Area interactions were similar to the other

experiments, as can be seen in Tables 2 and 3. In addition to the data of the Tables, the OFN-associated

tone elicited a larger FC negativity (main effect of Stimulus for PC2: $F(1,21) = 7.22$, $p = .014$, $\eta^2 = .26$)

and a larger TP positivity at T1 (Stimulus x Time interaction: $F(2,42) = 3.62$, $p = .036$, $\eta^2 = .15$) than the

SON-associated tone. Note, however, that these effects were "significant" only with the *uncorrected*

significance level.

The effects on **N1** were the same as in Experiment I. In addition to already depicted effects on **N2b**, a

significant Stimulus X Site interaction ($F(6,126) = 6.03$, $p < .001$, $\eta^2 = .22$) indicated that the anterior-

posterior inversion from negative to positive polarity took place for deviants at more anterior sites than

for standards.

The effects on **P3a** were identical to those in Experiments I and III, including the habituation of the

component (i.e., main effect of Time). There was no oddball effect on **P3b** at Pz (like Experiment II, but

differing from Experiments I and III). In contrast, the significant Stimulus x Time interaction was due to

the fact that at T2 and T3 the deviant previously linked with OFN elicited larger responses than the

previously linked with SON (main effect of Stimulus at T3: $F(2,42) = 5.74$, $p = .007$, $\eta^2 = .22$, with the

mean amplitudes of 1.47, 2.97, and 0.26 $\mu V/cm^2$ to standard, the former deviant, and the latter deviant,

respectively).

Discussion

At a behavioral level, the manipulation yielded the expected improvement of participants' reports as

compared with the preceding experiments. All participants correctly reported having heard three

different tones, two of which were associated with their own names or with other familiar names. ERP

results were, nevertheless, quite similar to the results of all previous experiments. The following two

analyses were intended to check this similarity.

Three experiments

Methods

Nineteen participants (11 females, aged 22-29, mean 25.5$\pm$0.48) yielded analyzable data in all three

Experiments (II, III and IV). Therefore, we performed additional ANOVAs with the results of these

subjects for the three experiments together. The analyses included the same repeated measures factors

plus the additional factor Experiment (3 levels).

Results

The analysis of **N3** revealed only one interaction that approached (but not reached) the corrected

significance level of .0083. The interaction resulted from topographic differences for PC4 being larger in

Experiment IV than in Experiments I and III (Experiment x Area interaction for lateral electrodes:

$F(8,144) = 3.43$, $p = .009$, $\eta^2 = .16$). Figure 6 illustrates the comparative dynamics of different ERP

components at the sites of their maximal expression.

/Figure 6 about here/

As regards **P3a**, both the mean component amplitude and its topographical differences were larger in

Experiment IV than in the two other experiments (main effect of Experiment: $F_{(2,36)} = 21.24$, $p < .001$,

$\eta^2 = .54$; Experiment x Site interaction: $F_{(6,108)} = 3.55$, $p = .022$, $\eta^2 = .17$). As regards **P3b**, the

topographical differences between the lateral areas (i.e., Fro and FC negativities, Pos and TP positivities)

were significantly smaller in Experiment II than in Experiments III and IV (Experiment x Area: $F_{(8,144)} =$

4.33, $p = .003$, $\eta^2 = .19$).

Comparison between N1 and N3

Methods

Given the apparent similarities in the distribution of N1 and N3, we conducted additional ANOVAs to

find differences between these negative components. These ANOVAs were similar to those reported

above but included an additional factor Principal Component (PC) having three levels (PC1, PC2, and

PC4). Since absolute differences in magnitude between the components are not very informative, only

interactions of PC with other factors will be reported.

Results

The results were almost identical in all experiments. Highly significant interactions in lateral areas were

obtained for the factors PC and Stimulus (Experiment I: $F_{(4,88)} = 13.57$, $\eta^2 = .38$; Experiment II: $F_{(4,76)} =$

8.09, $\eta^2 = .30$; Experiment III: $F_{(4,80)} = 12.81$, $\eta^2 = .39$; Experiment IV: $F_{(4,84)} = 8.69$, $\eta^2 = .29$; all $p <$

$.001$), PC and Area (Experiment I: $F_{(8,176)} = 4.12$, $p = .022$, $\eta^2 = .16$; Experiment II: $F_{(8,152)} = 5.39$, $p =$

$.002$, $\eta^2 = .22$; Experiment III: $F_{(8,160)} = 3.71$, $p = .035$, $\eta^2 = .16$; Experiment IV: $F_{(8,168)} = 22.64$, $p <$

$.001$, $\eta^2 = .52$), as well as between PC, Stimulus, and Area (Experiment I: $F_{(16,352)} = 2.45$, $p = .039$, $\eta^2 =$

$.10$; Experiment II: $F_{(16,304)} = 2.85$, $p = .002$, $\eta^2 = .13$; Experiment III: $F_{(16,320)} = 3.13$, $p = .007$, $\eta^2 =$

.14; Experiment IV: $F(16,336) = 2.36$, $p = .010$, $\eta^2 = .10$). An inspection of these interactions showed that

the difference between standards and deviants was significantly larger for N1 than for both N3

subcomponents. Particularly, significant PC x Stimulus interactions (all $p \leq .002$) were obtained in FC and

Ce areas, where both the absolute negative values and the differences between standards and deviants

were substantially larger for N1 than for N3. When PC1/N1 was excluded from the analysis (that is, the

factor PC was taken with two levels), no effect of PC whatsoever approached significance.

Significant PC x Time interactions were obtained in Experiments II ($F(4,76) = 5.39$, $p = .002$, $\eta^2 = .22$) and

IV ($F(4,84) = 4.31$, $p = .005$, $\eta^2 = .17$). The N1 amplitude significantly decreased with time (linear trend:

$F(1,21) = 8.25$, $p = .009$, $\eta^2 = .28$; and $F(1,21) = 15.87$, $p = .001$, for Experiments II and IV, respectively),

but the amplitudes of N3 did not. Finally, Experiment IV showed a significant interaction between PC

and Site *at midline* ($F(6,126) = 5.40$, $p = .001$, $\eta^2 = .20$). While all negativities were characterized by the

common trend to smaller negative (or larger positive) values in the posterior direction, the exact

distribution was different: the negative-to-positive inversion took place already at Cz for N3, but only at

Pz for N1.

Experiment V

Methods

Participants were 13 females and nine males, aged 19-33 (mean 25.0). Experiment V was, in fact,

designed for a different purpose (investigation of affective conditioning), but included into the present

report for a simple reason. Experiments I to IV all employed three stimuli and the similarity of their

results might be attributed to this fact. In Experiment V, however, ERPs were recorded to two stimuli,

and we wondered whether the same N3 effects would be obtained in this condition too.

This experiment also entailed an Acquisition phase and a Test phase. Five negative and 5 positive

emotional exclamations were used. The stimuli and their effect on EEG and magnetoencephalographic

(MEG) responses were reported in Kotchoubey et al. (2003), Bostanov and Kotchoubey (2004), and

Kotchoubey et al. (2009). Emotionally positive and negative exclamations did not differ from each other

in terms of the average duration and fundamental frequency (F0), and their average loudness was

equalized as well. The Acquisition phase consisted of 50 presentations of Chord 1 paired with five

positive exclamations (10 times each exclamation) and 50 presentations of Chord 2 paired with five

negative exclamations (10 times each). The two chords were 440 + 880 + 1760 + 3520 + 7040 Hz and 150

+ 300 + 600 + 1200 + 2400 Hz, and their linkage to positive and negative emotional exclamations was

counterbalanced among participants. Tone duration was 200 ms, SOA between the tone and the

exclamation 300 ms, and SOA between pairs randomly varied from 2000 to 2280 ms. The following Test

phase was an oddball paradigm: one of the chords presented during Acquisition was randomly selected

as the standard (presented 300 times), and the other one, as the deviant (presented 60 times).

Emotional exclamations were not presented in the Test phase anymore. SOA in the Test phase randomly

varied between 950 and 1050 ms.

Table 4. Principal components selected in the experiments with three stimuli (Experiment I to IV) and in
Experiment V with two stimuli. Note very close peak latencies of the PCs obtained with different
stimulation

Peak name		N1	P2	P3a	N2b	P3b	N3early	N3late
Exp I-IV	Principal component	PC1	PC6	PC5	PC3	PC7	PC2	PC4
	Latency, ms	96	136	215	294	334	412	456
Exp V	Principal component	PC4	PC3	PC6	PC2	PC5	PC1	PC8
	Latency, ms	90	129	207	294	336	416	459

The same PCA procedure was applied to Experiment V data separately from the other four experiments.

44 temporal factors were extracted for rotation. Despite numerous methodological differences between

the experiments, the PCA resulted in surprisingly similar distribution of the main components (see

Figure 7). The only major difference was the presence of PC7 (peak latency 376 ms) that could not be

identified as a peak in the ERP waveform. As shown in Table 4, other PCs were almost identical to those

in Experiments I – IV, only their order was changed.

/Figure 7 about here/

Results

The main object of the present study, **N3**, was again represented as two subcomponents, PC1 and PC8.

At midline, both were negative at frontal and central sites but positive at Pz (main Site effect). Both

fronto-central negativities and parietal positivities were about twice larger to deviants than to standards

(Stimulus x Site interactions). The analysis over lateral sites revealed negative amplitudes in Fro and FC

areas, and positive amplitudes in Pos and TP areas (main effect of Area). The topographical differences

were larger for deviants than standards (Table 5), resulting in highly significant Stimulus x Area

interactions. The amplitude of PC1 (but not PC8) appeared to decrease with time, particularly at frontal

sites on midline and lateral Fro and FC areas, but neither the main effect of Time, nor the Time x Site

and Time x Area interaction attained the corrected significance level.

N1 (**PC4)** had its midline maximum at frontal and fronto-central sites with a decreasing amplitude

toward Pz. The amplitude to deviants was larger than to standards at all sites except Pz, resulting in a

significant interaction between Stimulus and Site. At lateral sites, the amplitude was negative (and more

negative to deviants than standards) in the Fro and FC areas, and positive (and more positive to deviants

than standards) in the TP areas, leading to strong main effects of Area, Stimulus and their interaction.

N2b (PC2) was positive related to the baseline at Cz, CPz, and Pz and close to zero frontally. At lateral

sites its amplitude became more positive in the posterior direction. Also, the amplitude was generally

positive to deviants and negative to standards.

Table 5. Anterior-posterior differences in the amplitudes (in $\mu V/cm^2$) of two Principal Components (PCs)
reflecting N3. Fro – frontal, TP – temporoparietal. "Lateral" means averaged values for the left and right
hemispheres, because the factor Hemisphere was not significant.

		Fro midline	Pz	Fro lateral	TP lateral
PC1	standard	-2.9	-9.8	-4.2	-11.1
	deviant	1	4.8	4.7	4.4
PC8	standard	-2.7	-5.3	-4.8	-6
	deviant	1.9	3.7	2.9	5.1

Both positive components, **P3a (PC6)** and **P3b (PC5)**, decreased at midline with Time and were more

positive to deviants than standards. The stimulus effect on P3a was best expressed at Cz, and the

stimulus effect on P3b, at Pz (but Stimulus x Site interaction reached significance only for P3a).

The lateral distribution of the positive components strongly differed between standards and deviants.

Although both stimuli elicited negative **P3a** amplitudes in Fro areas, the maximum positivity to standard

was observed in TP areas, but the maximum positivity to deviant was much more anterior, in FC areas.

P3b amplitudes to standards were slightly negative in almost all areas (Fro, FC, Ce, Pos), but positive in

TP areas, whereas P3b amplitudes to deviants were negative in FC areas, zero in Fro areas, and positive

in Ce, Pos and TP areas. Thus both components displayed the main Area effect and the Stimulus x Area

interaction. Averaged across all areas, mean amplitudes were positive to deviants and slightly negative

to standards.

Like in the previous experiments, we directly compared N1 with N3 in Experiment V because

superficially they appear to display similar dynamics. PC1 was taken as the best representative of N3,

and PC4, as the representative of N1. Thus the factor PC had 2 levels.

In the lateral ANOVA the factor PC strongly interacted with Stimulus ($F_{(1,21)}$ = 30.13, p <. 001, η^2 = .59)

and Area ($F_{(4,84)}$ = 9.09, p <. 001, η^2 = .30), also yielding a three way PC x Stimulus x Area interaction

($F_{(4,84)}$ = 4.71, p =. 006, η^2 = .18). Generally, the difference between standards and deviants was

significantly larger for PC4/N1 than for PC1/N3. The topographic distributions were different: N1 was

negative in the Fro, FC, and Ce areas, zero in the Pos areas, and positive only in the TP areas, and the

mean difference between the most negative Fro areas and the most positive TP areas was 8.8 $\mu V/cm^2$.

N3 was negative in the Fro and FC areas, zero in the Ce areas, and positive in the Pos and TP areas, and

the difference between the most negative and most positive areas was 12.3 $\mu V/cm^2$. The maximal effect

of stimulus (> 4 $\mu V/cm^2$ difference between deviant and standard) was obtained in the Ce areas for N1

but in the FC and TP areas for N3.

The analysis at midline replicated the interactions with topography (PC x Site: $F_{(3,63)}$ = 8.19, p =. 001, η^2

= .28; PC x Stimulus x Site: $F_{(3,63)}$ = 4.84, p =. 016, η^2 = .19). Like at the lateral sites described above, the

transition from negative to positive values happened more posteriorly for PC1/N3 than for PC4/N1, and

the difference between maximal negative and maximal positive values was larger for the former than

the latter (5.1 and 8.1 $\mu V/cm^2$ for PC4/N1 and PC1/N3, respectively). The largest effect of stimulus was

at Cz for PC4/N1 but at frontal leads for PC1/N3. Figure 8 presents the comparison between N3 and N1

in three experiments.

/Figure 8 about here/

For control, this analysis was repeated using PC8 instead of PC2. According to our inspection, PC8

reflected a later subcomponent of the same component N3. In fact the analysis of PC1 versus PC8 did

not reveal any significant interactions of the factor PC.

Discussion

The analyses of Experiments II, III, and IV together, as well as the results of Experiment V, confirm the

stability of the N3 component and its relative independence of the design of oddball experiments. The

effects of the methodical differences between the experiments were very weak if any. The data further

dissolve the view of the apparent similarity between N3 and N1: when the two components are included

in an ANOVA as a factor, substantial differences between them are revealed. In contrast, when two N3

subcomponents are included in an ANOVA as a factor, this factor does not interact with any other

indicating that the two subcomponents reflect basically the same thing.

Mastoid references

Because many studies standardly report ERP data using average mastoid reference, we additionally

carried out the same PCA for mastoid-referenced data to enhance comparability with the literature. The

results were very similar for all experiments. Therefore, here we report the results of an analysis

including the data of all five experiments together. In the experiments with two deviant stimuli,

responses to all deviants were averaged together to obtain one standard and one deviant waveform.

The PCA yielded a total of 41 Factors. The following analysis was performed like for CSD data.

As can be seen in Figure 9, principal component peaks had latencies very similar to CSD-extracted

components: 96 ms for N1 (PC1), 215 ms for P3a (PC5), 334 ms for P3b (PC7), 412 and 456 ms for two

subcomponents of N3 (PC6 and PC9, respectively). The main difference between CSD and mastoid

referenced N3 is that in the latter case, N3 decreased in the posterior direction but did not reverse its

polarity; that is, it remains negative at all sites.

/Figure 9 about here/

We omit the results of the statistical analysis of mastoid-referenced data (which was performed in

exactly the same manner as for CSD data) because this would considerably increase the volume of the

article without any added information value.

General Discussion

Reliability

To our best knowledge, this is the first systematic investigation of the N3 wave in the oddball paradigm.

Being the first, it is largely exploratory in character. We conducted a large number of analyses, some of

which with up to five factors. Such analyses can yield false positive results. We are aware both of this

problem and of the fact that it still does not have a satisfactory solution. Thus we admit that some

effects described in the Results section may later turn out to be spurious. Nevertheless, we have several

reasons to be convinced that at least the most important findings are reliable. First, many of them

attained very high significance level with $p \leq .001$, which would survive any reasonable correction. But

the second reason for our optimism is much more important than the formal level of significance: the

major effects are consistently *replicated* from experiment to experiment notwithstanding considerable

differences between experimental conditions. This pleasantly surprising replicability already starts with

the results of PCA. The latencies of principal components found in different experiments differed, on

average, in 4.8 ms; the latencies of the components reflecting N3 (the target component of the present

study) differed in 2.7 ms only. In other words, the structure of ERP was virtually identical in all

experiments.

N3 is obviously not an artifact. In our experiments all kinds of artifacts were carefully controlled.

Participants listened to stimuli with eyes closed, which strongly diminished ocular effects. Bad channels

were replaced by interpolation, non-EEG components were removed by means of the ICA, and finally, a

few remaining artifact trials were excluded from averaging. Also in the literature we could find no

indication of the artificial nature of N3.

The PCA showed that N3 consistently entailed two subcomponents only weakly correlating with each

other, with peak latencies of 412-416 and 456-459 ms, respectively. All experimental effects, however,

were virtually the same on the two subcomponents and revealed only quantitative differences (mostly,

the earlier subcomponent exhibited stronger effects than the later one). Both subcomponents also had

the same topography.

Differentiation from other phenomena

The strong similarity between the earlier and the later N3 subcomponents can be compared with the

significant differences between them and the other ERP negativities. Although a superficial glance might

see a similar distribution of N3 and N1 (negativities in frontal areas, positivities in temporo-parietal

areas), their direct comparative analysis demonstrated substantial differences. The point of the

negative-to-positive inversion was much more anterior for N3 than for N1. The maximum of frontal

negativity was more lateralized for N1 than N3. The distribution of N1 CSD is compatible with a fronto-

temporal sink, whereas the distribution of N3 CSD is rather in line with a midfrontal sink. Further, N1

demonstrated a tendency to habituation, but N3 did not.

As all stimuli in our experiments had a duration of 200 ms, one might even ask whether N3 was an N1 of

the *offset* response. But this hypothesis is, of course, wrong just because other studies (including our

own: Kotchoubey and Pavlov, 2017) used stimuli of broadly varying duration, yet the peak latency of N3,

according to the corresponding figures, was always between 400 and 450 ms from stimulus *onset*.

Likewise, N3 is obviously not a mismatch negativity. CSD data of the MMN demonstrate a completely

different topographical distribution (see, for example, Deouell et al., 1998, Fig. 3b; Shalgi and Deouell,

2007, Fig. 5; Kayser and Tenke, 2015, Fig. 12). Furthermore, the number of stimuli in the present

experiments was rather small. N3 could be seen even after as few deviants as 20-50. Although a few

studies showed the MMN after 50-70 deviants, most typical MMN experiments use hundreds of

deviants to demonstrate consistent effects.

The decrease of the N3 amplitude in the posterior direction might remind on the Positive Slow Wave

(PSW, e.g., Squires et al., 1975; Sutton and Ruchkin, 1984; Dien, 1998; Spencer et al., 1999; Dien et al.,

2004) that can also have negative amplitudes in frontal leads. However, the analysis of the data with

average mastoid reference (Figure 8) shows that, contrary to the PSW, N3 does not change its polarity in

the sagittal dimension, i.e., it remains negative also in the parietal areas. This is in line with CSD data

indicating that the electrical source of the N3 dipole is not parietal but posterior temporal, in the vicinity

of the mastoid electrodes. Furthermore, Kayser et al. (1998) identified N3 and the PSW as two different

principal components.

Other two late components that can be mentioned *post hoc* are the O-wave (Näätänen et al., 1982) and

the reorienting negativity (RON: Schröger & Wolff, 1999; Berti, 2008). We did not think on these

components in advance because the conditions of our experiments have nothing in common with those

in which O-wave and the RON are ordinarily obtained. The former occurs in attended conditions, usually

when a motor response is required to the next expected stimulus. It is long lasting and can (like the PSW

above) be inverted at parietal sites, while N3 is recorded in the passive condition, relatively short-lasting

and always negative except over posterior temporal regions.

The RON, as one of its discoverers emphasized, "was observed after distraction *only in situations when a*

reorientation on the task at hand was required"; therefore, "the RON was interpreted as the correlate of

a switch of attention onto the task-relevant information" (Berti, 2008; p.609; italics added). Most

usually, a RON follows P3a without a P3b, contrary to N3 that follows P3b. Sometimes, however, the

positivity prior to a RON contains two components called "early P3a" and "late P3a" (e.g., Correa-Jaraba

et al., 2016, 2018), and there is no strong argument as of why the latter cannot, in fact, be a P3b. In

some exceptional cases one can see a clear P3b before the RON (e.g., Scheer et al., 2016).

In respect of time course and morphology, the RON is much more similar to the passive N3 than any

other late components such as O-wave or PSW. Moreover, a few studies using CSD analysis also indicate

a surprising similarity in the topography (e.g., Rämä et al., 2018). To sum up, there is an astonishing

contradiction between the obvious similarities of N3 and the RON, on the one hand, and the drastic

differences in the experimental conditions in which these two waves are elicited. The contradiction can,

from our point of view, be solved in one of two ways. Either a more comprehensive analysis will detect

substantial differences between the RON and N3 as two independent components, or, if they are

identical, the present-day interpretation of RON (including its processing specificity) will be seriously

challenged, because the typical conditions of N3 do not involve any reorientation process.

Furthermore, N3 does not appear to depend on the alleged significance of stimuli, at least within the

range tested in this study. Tones, which elicited N3, had been previously related to participants' own

names in Experiments I and IV, to emotional stimuli in Experiment V, to unconceivable sounds in

Experiment III, and were completely neutral in Experiment II. No significant differences between the

results of these experiments were observed. Likewise, N3 did not substantially depend on the number of

stimuli in the oddball experiment, as it was equally recorded in two- and three-stimuli paradigms.

Finally, N3 demonstrated an oddball effect, being larger to deviants than standards. In contrast to N1

and P3, this effect was rather manifested as a Stimulus x Topography interaction than as a main effect of

Stimulus. This means that N1 was consistently more negative, and P3a and P3b were more positive to

deviants than standards, although these amplitude increases also had their local maxima for each

component. N3, in contrast, was typically more negative to deviants than standards over anterior

regions but less negative to deviants than standards over posterior regions.

Passive N3 and its hypothetical functional meaning.

Table 5 shows an overview of the presence of N3 in the previously published passive oddball reports.

The search was performed on PUBMED using the terms "passive oddball" or "passive P3". Because we

assume that N3 is a sharp negativity following P3b, we included only studies whose results indicate a

P3b (and not only the complex MMN-P3a). Another inclusion criterion was that ERP waveforms at least

at the three midline sites (Fz, Cz, Pz) were clearly presented.

Table 6. Experiments designated as "passive oddballs", in which figures do or do not show an N3
following P3b.

	Study	Stimulus modality	Instruction	N3 yes/no	comment
1	Polich 1987	auditory	To daydream	No	
2	Polich 1989 Exp1	auditory	To ignore	Yes	
3	Polich 1989 Exp2	auditory	To ignore	No	Stimulus sequences were
4	Polich 1989 Exp2	auditory	To solve a puzzle	No	not random, i.e., deviants might be predicted
5	O'Donnell 1992	visual	To relax and ignore	Yes	N3 in participants from 20 to 31 years old, no N3 in

					older participants
6	Polich 1994	auditory	Just to listen	Yes	
7	Oades 1995a	auditory	Unclear	Yes	
8	Oades 1995b	auditory	Unclear	Yes	
9	Lang 1997	visual	to pay attention w/out a task	No	In the corresponding active experiments participants had to pay attention *and count* rare stimuli
10	Lang 1997	auditory	to pay attention w/out a task	No	
11	Bennington 1999	visual	Just to observe	No	Deviants might be predicted like in Polich 1989 above (No. 3, 4)
12	Bennington 1999	auditory	Just to listen	Yes	
13	Zenker 1999	auditory	Just to listen	Yes	Participants were children from 6 to 14, N3 observed only in the oldest group
14	Potts 2004	visual	Just to observe	Yes	
15	Holeckova 2006	auditory	Watching TV	Yes	Stimuli were own name and other names
16	Barry 2006	auditory	Just to listen	Yes	
17	Bostanov 2006	auditory	Just to listen	Yes	Confirmed by a time-frequency analysis
18	Brinkman 2008	auditory	Viewing a	Yes	N3 larger in children than

			cartoon		adults
19	Wronka 2008	auditory	Visual task	No	Three-stimuli paradigm
20	Mueller 2008	auditory	Just to listen	Yes	In older children and young adults
21	Barry 2009	auditory	Just to listen	Yes	
22	Eichenlaub 2012	auditory	Viewing a video	No	Stimuli were own name and other names
23	Fisher 2014	auditory	Viewing a video	Yes	Three-stimuli paradigm, N3 in both healthy subjects and (delayed) in schizophrenic patients
24	Erlbeck 2014	auditory	Just to listen	Yes	
25	Erlbeck 2014	auditory	Visual task	No	
26	Kayser 2014	auditory	No particular task	Yes	
27	Kühnis 2014	auditory	Visual task	No	Vowels presented to musicians and lay persons
28	Marhöfer 2014	visual	Visual distraction	No	Four-stimuli oddball (3 deviants)
29	Brown 2015	auditory	Solving a puzzle	No	
30	Brown 2015	auditory	Reading a book	No	
31	Rogenmoser 2015	auditory	Visual task	No	Auditory stimuli difficult

					to distinguish
32	Justen 2016	auditory	No particular task	Yes	Simple tones
33	Justen 2016	auditory	No particular task	No	Individual finger snapping sounds
34	Scheer 2016	auditory	Steering task	Yes	Environmental sounds
35	Morlet 2017	auditory	Mind wandering	Yes	
36	Morlet 2017	auditory	Active distraction	No	
37	Justen 2018	auditory	No particular task	Yes	Simple tones

References to the table: (Barry et al., 2006; Barry et al., 2009; Bennington and Pollich, 1999; Bostanov

and Kotchoubey, 2006; Brinkman and Stauder, 2008; Brown et al., 2015; Eichenlaub et al., 2012; Erlbeck

et al., 2014; Fisher et al., 2014; Holeckova et al., 2006; Justen and Herbert, 2016; Justen and Herbert,

2018; Kayser et al., 1997; Kühnis et al., 2014; Lang et al., 1997; Marhöfer et al., 2014; Morlet et al., 2017;

O'Donnell et al., 1997; Oades et al., 1995a; Oades et al., 1995b; Polich, 1987; Polich, 1989; Polich and

McIsaak, 1994; Potts, 2004; Rogenmoser et al., 2015; Scheer et al., 2016; Wronka et al., 2008; Zenker

and Barajas, 1999).

The table lists a total of thirty-seven experimental conditions, N3 being observed in twenty-one of them.

Of these 21, the condition described as "passive" included an additional distraction task in four

experiments: viewing a cartoon in Brinkman and Stauder (2008), viewing a video in Holeckova et al.

(2006) and Fisher et al. (2014), and a demanding steering task in Scheer et al. (2016). Inspecting these

four experiments, one can see that Holeckova et al. (2006) presented participants' own names and

similar names as the to-be-ignored auditory stimuli, and one may suppose that the distraction task did

not completely distract participants from the processing of the highly significant words. In Scheer et al.

(2016) a strong N3 after P3b was elicited by highly attractive environmental stimuli such as laughing, but

a rather small N3 (without a preceding P3b) was also observed in response to tones. Participants of

Brinkman & Stauder (2008) were adolescents, and those of Fisher et al. (2014), schizophrenic patients.

However, in the latter study N3 was also observed in the control group. In the other 17 experiments

showing N3, the authors implicitly (Oades et al., 1995a, b) or explicitly (the rest of studies) indicated the

lack of any task: participants should simply view visual stimuli or listen to auditory ones.

Of the 16 data sets that do *not* show an N3, active distraction was used in nine experiments. Moreover,

the participants in both visual and auditory experiments of Lang et al. (1997) were instructed to actively

follow stimuli, albeit without a motor task. Polich (1987) instructed his subjects to daydream, which may

or may not be regarded as distraction. In his later works (Polich, 1989) a passive condition without any

distraction was presented twice, and N3 was recorded in Experiment 1 but not in Experiment 2. The

sequence of auditory stimuli in this study and in Bennington and Polich (1999) was not random or

pseudorandom, but consisted of ten-stimuli trains, and deviants could only appear in the second halves

of the trains. This means that subjects could guess deviants more or less successfully. In an extreme

case, after a series of 9 standards, the deviant could even be predicted with certainty. Finally, Justen and

Herbert (2016) showed an extremely large N3 in a completely passive condition with tonal stimuli, but

very small (if any) in the same oddball with sounds of snapping fingers. Again, such sounds (also

including snapping of a participant's own fingers) are highly idiosyncratic.

Of course, passive conditions present a principal psychological problem, because the less exact is

participants' instruction, the less do we really know what they were doing during stimulation. This

general difficulty is also valid for the experiments summarized in Table 5. However, the above analysis

results in a rough differentiation between the two kinds of conditions that can be designated as

"passive-attentive", in which participants' attention was directed either to a different class of stimuli or

to the presented stimuli (but without any response requirement), and "passive-inattentive", in which

participants' attention was allowed to freely move around. If this classification is correct, N3 was shown

in 17 "passive-inattentive" conditions and 4 "passive-attentive" conditions. No N3 was shown in 5

"passive-inattentive" conditions and 11 "passive-attentive" conditions. The resulting difference is highly

significant (Fisher Exact Test: p = .006).

This leads to a hypothesis that N3 may be a result of an intermediate level of stimulus processing that

differs from both very superficial and very deep processing. Generally, slow cortical negativities reflect

preparatory activation of apical dendrites, and thus anticipatory "warming up" large populations of

cortical neurons (Rockstroh et al., 1989; Birbaumer et al., 1990; Mitzdorf, 1991). Because such

preparation to a further activity is usually followed by the phase of the realization of the prepared

potential, the vast majority of cortical ERPs can be presented as a sequence of negative-positive

complexes (i.e., cycles of preparation and realization: Kotchoubey, 2006). In some cases, however, a

cortical response ends with a negative rather than positive phase, such as the occipital P1-N1 complex of

visual ERPs or the N400 to semantically unrelated word pairs (e.g., *table-mouse*) presented on the

background of related word pairs (e.g., *cat-mouse*). Such an ending negativity is usually a sign of

preparation that did not find its realization (Kotchoubey, 2006).

According to this hypothesis, N3 should be elicited neither during a goal-directed activity, when the

eliciting stimulus is regarded as task relevant, and the corresponding task is performed, nor in a situation

of attentional distraction when the stimulus is regarded as fully irrelevant and processing resources are

directed to a completely different task. In the former case, there are two options. With simple task-

relevant stimuli, such as in most oddball experiments, the processing is closed with a P3b whose

meaning has been a subject of long discussions (e.g., Johnson, 1986; Donchin and Coles, 1988; Verleger,

1988; Verleger et al., 1994; Kotchoubey et al., 1997). Very complex or highly emotional stimuli elicit,

beyond this, late slow wave activity that reflects sustaining processing. In the latter case again two

options appear: if an irrelevant stimulus matches the current expectancies, its processing ends with the

P2 component some 180-250 ms after stimulus onset; if it mismatches the expectancies, the complex of

the MMN and P3a follows. None of these four cases requires an additional negative deflection N3 after

P3b.

The situation is different when the stimulus is regarded by the brain as potentially relevant, yet its

supposed relevance cannot be linked to any adaptive activity. It appears to mean something, but no

behavioral consequence can be inferred from this meaning. The result is processing that commences but

does not come to a closure.

If this hypothesis is true, further experiments can be designed to vary the precision of the meaning of

presented stimuli. A starting paradigm may be the design of Erlbeck et al. (2014) or Morlet et al. (2017)

who employed an active condition (attention to the stimuli), a distraction condition (attention away

from the stimuli), and a passive condition (free attention). However, future experiments on this basis

should employ more gradual conditions, such as attentional tasks with high and low motivation

(assuming that motivation mobilizes attentional resources), instructions to pay attention to the stimuli

without any overt or covert response, or just presentation of stimuli without any instruction. Also,

though the results of the above studies *appear* to be in line with the hypothesis (i.e., a manifested N3 is

obtained in the passive condition only), nothing can be said with confidence because the authors were

primarily interested in earlier components such as the MMN and P3.

Moreover, if this hypothesis is true, N3 might also be found in some *active* oddball experiments, e.g.,

when the experimental instruction was (intentionally or not) formulated with less precision. For

example, N3 in active oddball was recorded in Polich (1989) and Kayser et al. (1998), and in Müller et al.

(2008) it was even larger in the active than passive paradigm, which is in strong contrast with the data of

Erlbeck et al. (2014), Brown et al. (2015) and Morlet et al. (2017). However, the hypothesis about a link

between the active N3 and the precision of the experimental instruction can only be tested in a

systematic review of the voluminous literature about ERPs in active oddball, which should be a subject

of a separate study.

Funding

The study was supported by the German Research Society (Deutsche Forschungsgemeinschaft), Grant

KO-1753/13-1.

Acknowledgement

The authors thank Janna Holst and Ines Armbruster for their help in carrying out the experiments.

REFERENCES

Barry RJ, Rushby JA, Smith JL, Clarke AR, Croft RJ. (2006): Dynamics of narrow-band EEG

phase effects in the passive auditory oddball task. Europ J Neurosci 24:291-304.

Barry RJ, Rushby JA, Smith JL, Clarke AR, Croft RJ. (2009): Brain dynamics in the auditory

oddball task as a function of stimulus intensity and task requirements. Internal J

Psychophysiol 73:313-325. doi: 10.1016/j.ijpsycho.2009.05.003.

Barry RJ, De Blasio FM. (2015). Performance and ERP components in the equiprobable go/no-

go task: Inhibition in children. Psychophysiol 52(9):1228-1237. doi: 10.1111/psyp.12447

Bennington JY, Polich J. (1999): Comparison of P300 from passive and active tasks for auditory

and visual stimuli. Internal J Psychophysiol 34:171-177.

Berti S. (2008): Cognitive control after distraction: Event-related brain potentials (ERPs)

dissociate between different processes of attentional allocation. Psychophysiology 45,

608-620. doi: 10.1111/j.1469-8986.2008.00660.x

Birbaumer N, Elbert T, Canavan AGM, Rockstroh B. (1990): Slow potentials of the cerebral

cortex and behavior. Physiol Rev 70:1-41.

Bostanov V, Kotchoubey B. (2004): Recognition of affective prosody: Continuous wavelet

measures of event-related brain potentials to emotional exclamations. Psychophysiol

41:259-268.

Bostanov V, Kotchoubey B. (2006): The t-CWT: A new ERP detection and quantification

method based on the continuous wavelet transform and Student's t-statistics. Clin

Neurophysiol 117:2627-2644.

Brinkman MJ, Stauder JE. (2008). The development of passive auditory novelty processing.

Internat J Psychophysiol 70:33-39. doi: 10.1016/j.ijpsycho.2008.04.005.

Brown SBRE, Van der Wee NJA, Van Noorden MS, Giltay EJ, Neuwenhuis S. (2015):

Noradrenergic and cholinergic modulation of late ERP responses to deviant stimuli.

Psychophysiol 52:1620–1631.

Correa-Jaraba KS, Cid-Fernández S, Lindín M, Díaz F. (2016). Involuntary capture and

voluntary reorienting of attention decline in middle-aged and old participants. Front Hum

Neurosci 10:129.

Correa-Jaraba KS, Lindín M, Díaz F. (2018). Increased amplitude of the P3a ERP component as

a neurocognitive marker for differentiating amnestic subtypes of mild cognitive

impairment. Front Aging Neurosci 10:19.

Courchesne E, Hillyard SA, Courchesne R. (1975): Stimulus novelty, task relevance and the

visual evoked potential in man. EEG Clin Neurophysiol 39:131-143.

Deouell LY, Bentin S, Giard MH. (1998): Mismatch negativity in dichotic listening: Evidence

for interhemispheric differences and multiple generators. Psychophysiol 35:355-365.

Dien J. (2010): The ERP PCA Toolkit: An open source program for advanced statistical analysis

of event-related potential data. J Neurosci Meth 187:138–145.

https://doi.org/10.1016/j.jneumeth.2009.12.009.

Dien J, Spencer KM, Donchin E. (2004). Parsing the "Late Positive Complex": Mental

chronometry and the ERP components that inhabit the neighborhood of the P300.

Psychophysiol 41(5):665-678.

Donchin E. (1981): Surprise! … Surprise? Psychophysiol 18:493-513.

Donchin E, Coles MGH. (1988): Is the P300 component a manifestation of context updating?

Behav Brain Sci 11:357-374.

Duncan-Johnson CC, Donchin E. (1977): On quantifying surprise: The variation of event-related

potentials with subjective probability. Psychophysiol 14:456-467.

Eichenlaub JB, Ruby P, Morlet D. (2012): What is the specificity of the response to the own

first-name when presented as a novel in a passive oddball paradigm? An ERP study.

Brain Res 1447:65-78. doi: 10.1016/j.brainres.2012.01.072.

Erlbeck H, Kübler A, Kotchoubey B, Veser S. (2014): Task instructions modulate the attentional

mode affecting the auditory MMN and the semantic N400. Front Hum Neurosci 8:654.

Falkenstein M, Hohnsbein J, Hoormann J. (1994): Effects of choice complexity on different

subcomponents of the late positive complex of the event-related potential. EEG Clin

Neurophysiol 92:148-160.

Fisher DJ, Smith DM, Labelle A, Knott VJ. (2014): Attenuation of mismatch negativity (MMN)

and novelty P300 in schizophrenia patients with auditory hallucinations experiencing

acute exacerbation of illness. Biol Psychol 100:43-49.

Getzmann S, Falkenstein M, Wascher E. (2015). ERP correlates of auditory goal-directed

behavior of younger and older adults in a dynamic speech perception task. Behav Brain

Res 278:435-445. doi: 10.1016/j.bbr.2014.10.026

Holeckova I, Fischer C, Giard MH, Delpuech C, Morlet D. (2006): Brain responses to a subject's

own name uttered by a familiar voice. Brain Res 1082:142-152.

Horn JL. (1965): A rationale and test for the number of factors in factor analysis. Psychometrika

30:179–185. https://doi.org/10.1007/BF02289447.

Johnson R. (1986): A triarchic model of P300 amplitude. Psychophysiol 23:367-384.

Justen C, Herbert C. (2016): Snap your fingers! An ERP/sLORETA study investigating implicit

processing of self- vs. other-related movement sounds using the passive oddball

paradigm. Front Hum Neurosci 10:465.

Justen C, Herbert C. (2018): The spatio-temporal dynamics of deviance and target detection in

the passive and active auditory oddball paradigm: A sLORETA study. BMC Neurosci 19,

25. doi: 10.1186/s12868-018-0422-3

Kayser J, Tenke CE. (2006): Principal components analysis of Laplacian waveforms as a generic

method for identifying ERP generator patterns: I. Evaluation with auditory oddball tasks.

Clin Neurophysiol 117:348–368. https://doi.org/10.1016/j.clinph.2005.08.034.

Kayser J, Tenke CE. (2015): Issues and considerations for using the scalp surface Laplacian in

EEG/ERP research: A tutorial review. 97:189-209.

Kayser J, Tenke CE, Croppmann CJ, Alschuler DM, Fekri S, Ben-David S, Corcoran CM,

Bruder GE. (2014): Auditory event-related potentials and alpha oscillations in the

psychosis prodrome: Neuronal generator patterns during a novelty oddball task. Internat J

Psychophysiol 91:104-120.

Kayser J, Tenke CE, Nordby H, Hammerborg D, Hugdahl K, Erdmann G. (1997): Event-related

potential (ERP) asymmetries to emotional stimuli in a visual half-field paradigm.

Psychophysiol 34:414-426.

Kayser J, Tenke CE, Bruder GE. (1998). Dissociation of brain ERP topographies for tonal and

phonetic oddball tasks. Psychophysiol 35:576-590.

Kotchoubey B. (2006): Event-related potentials, cognition, and behavior: A biological approach.

Neurosci Biobehav Rev 30:42-65.

Kotchoubey B, Grözinger B, Kornhuber AW, Kornhuber HH. (1997): Electrophysiological

analysis of expectancy: P3 in informed guessing. Internat J Neurosci 91:105-122.

Kotchoubey B, Kaiser J, Bostanov V, Lutzenberger W, Birbaumer N. (2009): Recognition of

affective prosody in brain-damaged patients and healthy controls: A neurophysiological

study using EEG and whole-head MEG. CABN 9:153-167.

Kotchoubey B, Lang S, Bostanov V, Birbaumer N. (2003): Cortical processing in Guillain-Barre

syndrome after years of total immobility. J Neurol 250:1121-1123.

Kotchoubey B, Pavlov YG. (2017): Name conditioning in event-related brain potentials.

Neurobiol Learn Mem, doi: https://doi.org/10.1016/j.nlm.2017.09.009

Kühnis J, Elmer S, Jäncke L. (2014): Auditory evoked responses in musicians during passive

vowel listening are modulated by functional connectivity between bilateral auditory-

related brain regions. J Cogn Neurosci 26:2750-2761. doi: 10.1162/jocn_a_00674.

Lang S, Kotchoubey B, Lutz A, Birbaumer N. (1997): Was tut man, wenn man nichts tut?

Kognitive EKP-Komponenten ohne kognitive Aufgabe. Z Exper Psychol XLIV:138-162.

Luck SJ, Gaspelin N. (2017): How to get statistically significant effects in any ERP experiment

(and why you shouldn't). Psychophysiol 54:146–157.

Marhöfer DJ, Bach M, Heinrich SP. (2014): Faces are more attractive than motion: evidence

from two simultaneous oddball paradigms. Doc Ophthalm 128:201-209. doi:

10.1007/s10633-014-9434-1.

Mitzdorf U. (1991): Physiological sources of evoked potentials. EEG Clin Neurophysiol Suppl.

42:47-57.

Morlet D, Ruby P, André-Obadia N, Fischer C. (2017). The auditory oddball paradigm revised to

improve bedside detection of consciousness in behaviorally unresponsive patients.

Psychophysiology 54(11), 1644–1662. Doi: 10.1111/psyp.12954

Mueller V, Brehmer Y, von Oertzen T, Li SC, Lindenberger U. (2008): Electrophysiological

correlates of selective attention: A lifespan comparison. BMC Neurosci 9, 18. doi:

10.1186/1471-2202-9-18

Näätänen R, Simpson M, Loveless NE. (1982): Stimulus deviance and evoked potentials. Biol

Psychol 14, 53-98.

Oades RD, Dittmann-Balcar A, Zerbin D. (1995a): The topography of 4 subtraction ERP-

waveforms derived from a 3-tone auditory oddball task in healthy young adults. Internat J

Neurosci 81:265-281.

Oades RD, Zerbin D, Dittmann-Balcar A. (1995b): The topography of event-related potentials in

passive and active conditions of a 3-tone auditory oddball test. Internat J Neurosci

81:249-264.

O'Donnell BF, Friedman S, Swearer JM, Drachman DA. (1992): Active and passive P3 latency

and psychometric performance: Influence of age and individual differences. Internat J

Psychophysiol 12:187-195.

Palmer JA, Kreutz-Delgado K, Makeig S. (2012): AMICA: An adaptive mixture of independent

component analyzers with shared components. Swartz Center for Computational

Neuroscience, University of California San Diego, Tech. Rep.

https://sccn.ucsd.edu/~jason/amica_a.pdf.

Polich J. (1987): Comparison of P300 from a passive tone sequence paradigm and an active

discrimination task. Psychophysiol 24:41-46.

Polich J. (1989): P300 from a passive auditory paradigm. EEG Clin Neurophysiol 74:312-320.

Polich J, McIsaak HK. (1994): Comparison of auditory P300 habituation from active and passive

conditions. Internat J Psychophysiol 17:25-34.

Potts GF. (2004): An ERP index of task relevance evaluation of visual stimuli. Brain Res 56:5-

13.

Rämä P, Leminen A, Koskenoja-Vainikka S, Leminen M, Alho K, Kujala T. (2018). Effect of

language experience on selective auditory attention: An event related potential study.

Internat J Psychophysiol 127:38-45.

Rockstroh B, Elbert T, Canavan A, Lutzenberger W, Birbaumer N. (1989): Slow Cortical

Potentials and Behavior. Baltimore: Urban & Schwarzenberg.

Rogenmoser L, Elmer S, Jäncke L. (2015): Absolute pitch: evidence for early cognitive

facilitation during passive listening as revealed by reduced P3a amplitudes. J Cogn

Neurosci 27:623-637.

Scheer M, Bülthoff HH, Chuang LL. (2016): Steering demands diminish the early-P3, late-P3

and RON components of the event-related potential of task-irrelevant environmental

sounds. Front Hum Neuroscie 10, 73. doi: 10.3389/fnhum.2016.00073

Schröger E, Wolff,C. (1998).: Attentional orienting and reorienting is indicated by human event-

related brain potentials: NeuroReport, 9(15), 3355–3358.

https://doi.org/10.1097/00001756-199810260-00003

Shalgi S, Deouell LY. (2007): Direct evidence for differential roles of temporal and frontal

components of auditory change detection. Neuropsychologia 45:1878–1888.

Spencer KM, Dien J, Donchin E. (1999). A componential analysis of the ERP elicited by novel

events using a dense electrode array. Psychophysiol 36:409-414.

Squires NK, Squires KC, Hillyard SA. (1975). Two varieties of long-latency positive waves

evoked by unpredictable auditory stimuli in man. EEG Clin Neurophysiol 38: 387-401.

Sutton S, Braren M, Zubin J, John ER. (1965): Evoked-potential correlates of stimulus

uncertainty. Science 150:1187-1188.

Sutton S, Ruchkin DS. (1984). The late positive complex: advances and new problems. Ann NY

Acad Sci 425:1-23.

Verleger R. (1988): Event-related potentials and cognition: A critique of the context updating

hypothesis and an alternative interpretation of P3. Behav Brain Sci 11:343-356.

Verleger R, Jaskowsky P, Wauschkuhn B. (1994): Suspense and surprise: On the relationship

between expectancies and P3. Psychophysiol 31:359-369.

Wronka E, Kaiser J, Coenen AM. (2008): The auditory P3 from passive and active three-

stimulus oddball paradigm. Acta Neurobiol Exper 68:362-372.

Zenker F, Barajas JJ. (1999): Auditory P300 development from an active, passive and single-

tone paradigms. Internat J Psychophysiol 33:99-111.

Figure legends

Figure 1. Experimental design.

Figure 2 An illustration of the effect of electrical reference, for the results of Experiment II at three

midline electrodes Fz, Cz, and Pz. Black dotted line, standards. Light red line, Deviant 1. Dark red line,

Deviant 2. Positivity in this and the following figures is plotted upwards.

Figure 3. Topographic representation of each principal component and its peak latency, calculated for

Current Source Density data. Because the results were nearly identical in Experiments I - IV, they are

presented here averaged across the four three-stimuli experiments.

Figure 4. Regions of interest (ROIs) selected for statistical analysis.

Figure 5. Average CSD ERP waveforms in each of the five experiments, at selected locations.

Figure 6. Mean amplitude of each ERP component in the region of its maximal expression (N1,

frontocentral right; P3a, Cz; N2b, temporoparietal left; P3b, temporoparietal left; N3a (early) and N3b

(late), frontocentral right). White bars, standard; light blue bars, Deviant 1 (or the only deviant in

Experiment V); dark blue, Deviant 2. Bars show standard errors of means.

Figure 7. Topographic representation of each principal component at their peak latencies for Experiment

V.

Figure 8. Comparison between the distributions of N1 and N3 over lateral areas. FC, frontocentral areas;

TP, temporoparietal areas. To save space only the data of three experiments are presented, but the

distributions were not qualitatively different also in the other two. Because of only minimal hemispheric

differences, left and right areas are averaged, as well as Deviant1 and Deviant 2 in Experiments with two

deviants. Amplitudes are in $\mu V/cm^2$. Bars show standard errors of means.

Figure 9. Results of mastoid-referenced data. Top: ERP waveforms averaged for all five experiments,

referred to average mastoids, for standards (black) and deviants (red). In Experiments I – IV two deviants

were taken together. Most prominent peaks are indicated by arrows (P2, which did not differ between

standards and deviants, is not shown). Bottom: topographical distributions for PCA data.

CSD   Common average   Mastoids

······ **Standards** —— **Deviants1** —— **Deviants2**

	St	Dev1	Dev2
N1 (PC1, 96 ms)			
P3a (PC5, 215 ms)			
N2b (PC3, 294 ms)			
P3b (PC7, 334 ms)			
N3a (PC2, 412 ms)			
N3b (PC4, 456 ms)			

µV/cm²

-20   0   20