1 Bacterial cyclopropane fatty acid synthase mRNA is targeted by activating and
2 repressing small RNAs
3

5 Colleen M. Bianco[1], Kathrin S. Fröhlich[2], and Carin K. Vanderpool[1*]

6 [1]Department of Microbiology, University of Illinois, Urbana, IL 61801, USA

7 [2]Microbiology, Ludwig-Maximilians-University, 82152, Munich, Germany
8

9
*Corresponding author
Carin K. Vanderpool, Ph.D.
Department of Microbiology
University of Illinois at Urbana-Champaign
B213 CLSL, MC-110
601 S. Goodwin Ave.
Urbana, IL 61801
(t) 217-333-7033
(f) 217-244-6697
cvanderp@illinois.edu
20
21
22

23 Multiple sRNAs regulate CFA Synthase
24

Abstract

Altering membrane protein and lipid composition is an important strategy for maintaining membrane integrity during environmental stress. Many bacterial small RNAs (sRNAs) control membrane protein production, but sRNA-mediated regulation of membrane fatty acid composition is less understood. The sRNA RydC was previously shown to stabilize *cfa* (cyclopropane fatty acid synthase) mRNA, resulting in higher levels of cyclopropane fatty acids in the cell membrane. Here, we report that three additional sRNAs, ArrS, CpxQ, and GadF also regulate *cfa* post-transcriptionally. RydC, ArrS, and GadF act through masking an RNase E cleavage site in the *cfa* mRNA 5' untranslated region (UTR), and all three sRNAs post-transcriptionally activate *cfa*. In contrast, CpxQ binds to a different site in the *cfa* mRNA 5' UTR and represses *cfa*. Alteration of membrane lipid composition is a key mechanism for bacteria to survive low pH environments, and we show that *cfa* translation increases in an sRNA-dependent manner when cells are subjected to mild acid stress. This work suggests an important role for sRNAs in the acid stress response through regulation of *cfa* mRNA.

Keywords:

Cyclopropane fatty acids synthase/ Fatty acid modification/ Hfq/ post-transcriptional gene regulation/ RNase E

Introduction

Bacteria modify the biophysical properties of their membranes to adapt to changing environmental conditions such as pH, temperature, and pressure fluctuations (Zhang & Rock, 2008). Membrane properties can be altered by changing the type or abundance of proteins embedded in the membrane or by modifying the relative proportions of different types of phospholipids. Membrane fluidity is a crucial biophysical property as it affects membrane-associated functions such as permeability to solutes, solute transport, and protein-protein interactions. The length and saturation of the acyl chains in the phospholipids determines the fluidity of the membrane, and bacteria adjust the ratio of saturated to unsaturated fatty acids (UFAs) to adapt to their environment. For example, bacteria can increase resistance to toxic compounds, such as antimicrobial peptides, by increasing the production of saturated fatty acids (Zhang & Rock, 2008), which will form a tightly packed and less fluid membrane compared to a membrane rich in unsaturated fatty acids.

While *de novo* production of fatty acids is an important adaptation mechanism, bacteria may encounter abrupt environmental changes that require rapid modification of the fatty acids already incorporated in the membrane. One post-synthetic modification is the conversion of a pre-existing UFA to a cyclopropane fatty acid (CFA) by the enzyme cyclopropane fatty acid synthase (encoded by *cfa*). CFAs are formed by the addition of a methylene group, derived from S-adenosyl methionine, across the double bond of an UFA incorporated in a phospholipid (Grogan & Cronan, 1997, Law, 1971). CFAs occur in the phospholipids of many species of bacteria, but although widely studied, their physiological function remains unclear. One hypothesis is that the formation of CFAs may reduce membrane fluidity and permeability (Grogan & Cronan, 1997), including the permeability to protons (Shabala & Ross, 2008). Indeed, *Escherichia coli cfa* mutants are unable to survive acid shock, a rapid shift from pH 7 to pH 3 (Brown, Ross et al.,

1997, Chang & Cronan, 1999). In addition, pathogenic *E. coli* strains contain more CFAs

and are more resistant to acid than nonpathogenic *E. coli* strains (Brown et al., 1997).

The formation of CFAs has been further linked to bacterial virulence as inactivation of a

cfa homologue that introduces a cyclopropane ring in the major cell envelope component

(α-mycolates) of *Mycobacterium tuberculosis* leaves the bacteria unable to establish a

persistent infection (Glickman, Cox et al., 2000).

Production of CFAs is regulated at multiple levels. In *E. coli* and *Salmonella*, *cfa*

transcription is driven by two promoters (Kim, Kim et al., 2005, Wang & Cronan, 1994).

The distal promoter is σ^{70}-dependent and yields a long *cfa* transcript with a 212-nt 5′-

untranslated region (UTR). The proximal promoter is controlled by the general stress

response σ factor (σ^{s}) encoded by *rpoS* and produces a short *cfa* transcript with a 34-nt

5′ UTR (Fig. 1A). The σ^{70}-dependent promoter is functional throughout growth, whereas

transcription from the σ^{s} promoter occurs only during stationary phase.

Post-transcriptional regulation by small non-coding RNAs (sRNAs) constitutes a

major layer of gene expression control in bacteria. In many cases, a ubiquitous RNA

chaperone, Hfq, is required for sRNA stability and to mediate base-pairing interactions

between an sRNA and its cognate target mRNAs. A recent study applied a methodology

based on ligation of Hfq-associated RNAs and sequencing of chimeric fragments to

globally map RNA interactions *in vivo* (Melamed, Peer et al., 2016). This approach

(termed RIL-seq) uncovered *cfa* mRNA interactions with multiple sRNAs in *E. coli*: ArrS,

CpxQ, GadF, and RydC. RydC was previously reported to stabilize *cfa* mRNA, which in

turn increases levels of cyclopropane fatty acid (CFA) synthase and results in

proportionally more CFAs in membrane lipids (Fröhlich, Papenfort et al., 2013). RydC, a

64 nucleotides (nt) long sRNA, folds as a characteristic pseudoknot structure which

exposes a stretch of highly conserved nucleotides at the very 5' end of the RNA. Binding

of RydC within the long *cfa* mRNA masks a recognition site for the major endonuclease

RNase E and inhibits transcript degradation. In addition, RydC has been reported to

regulate *yejABEF* (Antal, Bordeau et al., 2005), *csgD* (Bordeau & Felden, 2014)*, pheA*

and *trpE* (King, Vanderpool et al., 2019) mRNAs, though the conditions and signals

stimulating RydC production and the physiological role of RydC in bacterial physiology

remain unknown. In contrast, ArrS, CpxQ and GadF sRNAs were previously linked to

cell envelope or acid stress responses. Production of the CpxQ sRNA depends on the

Cpx two-component system that responds to cell envelope stress (Chao & Vogel, 2016).

CpxQ is processed by RNase E from the 3' UTR of *cpxP* mRNA, which encodes a

chaperone of the Cpx system (Chao & Vogel, 2016). Together with Hfq, CpxQ represses

translation of multiple targets many of which encode inner membrane proteins (Chao &

Vogel, 2016, Grabowicz, Koren et al., 2016). ArrS is an antisense sRNA that is encoded

upstream of *gadE* (encoding the major acid resistance transcription factor) and is

complementary to the 5' UTR of the longest of three *gadE* transcripts (Aiso, Murata et

al., 2011). ArrS expression is induced by low pH through σ^s and GadE, and

overexpression of ArrS increases survival of cells exposed to acidic pH (Aiso, Kamiya et

al., 2014). A second sRNA associated with the *gadE* locus is GadF, which is liberated

from the 3' UTR of *gadE* mRNA by endonucleolytic cleavage (Melamed et al., 2016). To

date, GadF has not been extensively studied, however the enrichment of acid response-

associated genes in its RIL-seq profile suggests a role in this pathway.

In this study, we sought to elucidate the mechanisms of regulation of *cfa* mRNA

by the four sRNAs, RydC, ArrS, GadF, and CpxQ, and to gain insight into the

physiological importance of sRNA-mediated regulation of membrane CFA composition in

the model organism *E. coli.* Our results indicate that RydC, ArrS and GadF activate *cfa*

to varying degrees, and all three require overlapping binding sites in the long 5' UTR of

cfa mRNA. We found that RydC is responsible for increasing *cfa* expression at acidic

pH, and that cells exhibit an acid stress resistance phenotype that is dependent on

sRNA-mediated post-transcriptional activation of *cfa.* Unlike RydC, ArrS and GadF, the

sRNA CpxQ represses *cfa* post-transcriptionally. CpxQ binds at a distinct site on *cfa*

mRNA, upstream from the site bound by activating sRNAs. Our results suggest that

CpxQ binding at this site modulates the stability of *cfa* mRNA by promoting its

degradation by RNase E. Altogether, our work suggests that several distinct sRNAs,

presumably responding to different environmental conditions, can modulate membrane

fatty acid composition via activation or repression of *cfa* mRNA.

Results

Multiple sRNAs regulate CFA synthase mRNA in an isoform-specific manner

At the transcriptional level, expression of *cfa* is controlled by the activity of two

promoters that yield mRNAs with a 212-nt UTR (distal σ^{70}-dependent promoter) or a 34-

nt UTR (σ^{S}-dependent promoter), respectively (Wang & Cronan, 1994) (Fig. 1A). Only

the long mRNA isoform is subject to post-transcriptional regulation by the sRNA RydC,

which stabilizes the transcript by base-pairing at an RNase E recognition site within the

5' UTR (Fröhlich et al., 2013). An experimental study based on the ligation of Hfq-

associated RNAs (Melamed et al., 2016) confirmed the previously observed interaction

between *cfa* mRNA and RydC sRNA, and in addition revealed the potential interactions

of the *cfa* transcript to ArrS, CpxQ, and GadF sRNAs. To verify that ArrS, CpxQ, and

GadF sRNAs alter *cfa* expression post-transcriptionally *in vivo*, we constructed two

translational *cfa'-'lacZ* fusions under the control of the arabinose-inducible P_{BAD} promoter

(King et al., 2019). One fusion contains the 212-nt 5' UTR and the first 36 codons of *cfa*

(*cfa'-'lacZ*-Long, Fig. 1B), while the second has only the short 34-nt 5' UTR (*cfa'-'lacZ*-

Short, Fig. 1C). We ectopically induced each sRNA from a plasmid (P_{Lac} promoter), and

determined the effect on the two reporter fusions. RydC and ArrS increased *cfa'-'lacZ*–

Long activity 18- and 10- fold, respectively, while CpxQ repressed the long fusion ~3-fold

(Fig. 1B). Expression of GadF resulted in a very minor increase (1.3-fold) of *cfa'-'lacZ*–

Long activity (Fig. 1B). No sRNA had an effect on *cfa'-'lacZ*–Short activity (Fig. 1C).

These results indicate that the long isoform of *cfa* mRNA can be post-transcriptionally

activated or repressed by multiple sRNAs, while the short isoform is not affected by

these regulatory molecules.

Furthermore, RIL- seq also revealed the potential interactions of the *cfa* transcript

to OxyS and GcvB sRNAs. We tested *cfa'-'lacZ*–Long and *cfa'-'lacZ*–Short for regulation

by OxyS and GcvB (Fig. EV1). Expression of OxyS increased *cfa'-'lacZ*–Long activity

2.1-fold while expression of GcvB decreased *cfa'-'lacZ*–Long activity 1.5-fold (Fig.

EV1A). Neither sRNA significantly affected *cfa'-'lacZ*–Short activity (Fig. EV1B). OxyS is

expressed under oxidative stress and represses *rpoS* expression by an unknown

mechanism (Altuvia, WeinsteinFischer et al., 1997, Zhang, Altuvia et al., 1998). GcvB

mutants are acid sensitive and this sensitivity may be due to reduced RpoS levels,

however how GcvB activates RpoS expression is not known (Jin, Watt et al., 2009). Due

to the potential regulation of RpoS by OxyS and GcvB (which could indirectly regulate

cfa'-'lacZ–Long by modulating expression from the σ^S-dependent promoter), and the

observed minor effect of these sRNAs on regulation of *cfa*, we did not further

characterize regulation by OxyS and GcvB.

Activating and repressing sRNAs bind distinct sites on *cfa* mRNA

RydC, ArrS, GadF, and CpxQ all regulated the *cfa* translational fusion containing the *cfa*

212-nt 5'-UTR but not the *cfa* translational fusion containing the short 34-nt 5'-UTR.

Given that sRNAs typically regulate their targets by engaging in Hfq-dependent

formation of base-pairing interactions, we assumed that each sRNA recognizes a site

within the upstream region of *cfa* mRNA. RydC has previously been shown to base-pair

at an RNase E recognition site located within this region (Fröhlich et al., 2013). We used

IntaRNA (Mann, Wright et al., 2017) to predict interactions between ArrS, GadF, and

CpxQ with *cfa* mRNA, and determined that all three sRNAs potentially bound the target

in proximity to the previously determined RydC site (Figs. 2A, EV2A). To genetically test

these base pairing interactions we first constructed a *cfa'-'lacZ* fusion with a 133-nt

deletion from the beginning of the 212-5′ UTR, which contains predicted binding sites of

all of the sRNAs (*cfa'-'lacZ*-ΔsRNABS, Fig. EV2B), including the RNase E recognition

site that RydC protects. This fusion only contains the proximal σ^S-dependent promoter

and 79-nt upstream of this promoter. Of note, *cfa'-'lacZ*-ΔsRNABS had higher basal

activity compared to *cfa'-'lacZ*-Long (Figs. 1B, EV2C), presumably because *cfa'-'lacZ*-

ΔsRNABS fusion does not contain a known RNase E recognition site, inherently

stabilizing the transcript. When each individual sRNA was ectopically expressed from a

plasmid, none of the sRNAs affected *cfa'-'lacZ*-ΔsRNABS activity (Fig. EV2C), indicating

each sRNA base pairs upstream of the -79 nt position. To specifically analyze where

each sRNA binds *cfa* mRNA, we completed structure probing experiments (Fig. 2). We

digested a ~140-nt fragment of *cfa* mRNA (-212 to -72 relative to the start codon) in the

presence of Hfq and each of the four individual candidate sRNAs with ribonuclease T1

or lead (II) acetate (Fig. 2B). We found that RydC (lanes 5, 10), and ArrS (lanes 6, 11)

bind sites that overlap, whereas CpxQ (lanes 8,13) binds upstream at a distinct site.

Protection by GadF was weak (lane 7, T1 digest) or not observed (lane 12, lead (II)

acetate cleavage), consistent with *in vivo* observations that GadF is a very weak

regulator of *cfa* mRNA. We did not characterize GadF-*cfa* mRNA interactions further.

The results of the *in vitro* structure probing experiment were in line with our

computational predictions of base-pairing sites, suggesting that the two activating

sRNAs, ArrS and RydC, shared a common binding site whereas CpxQ employed a

distinct site to repress *cfa* mRNA.

To test base-pairing predictions with each sRNA directly, we did a mutational

analysis. We first introduced a point mutation in the two activating sRNAs, ArrS and

RydC, that were predicted to disrupt base pairing to *cfa'-'lacZ*-Long (C7G for *arrS* M1),

and C5G for *rydC* M1 (Fig. 3A). These mutated sRNA variants, ArrS M1 and RydC M1,

could no longer activate *cfa'-'lacZ*-Long (Fig. 3B). We made a compensatory mutation

(G101C) to create *cfa'-'lacZ*-Long M1, to restore putative base pairing interactions with

the mutant sRNAs (Fig. 3A, B). Wild-type ArrS and RydC could not activate the mutant

cfa'-'lacZ-Long M1fusion (Fig. 3B), consistent with the idea that this residue in the *cfa* 5'-

UTR is critical for interactions with both sRNAs. In contrast, the mutants RydC M1 and

ArrS M1 activated the *cfa'-'lacZ*-Long M1 fusion (Fig. 3B), with fold-activation restored to

levels similar to the wild-type mRNA-sRNA pairs (Fig. 3B).

To characterize CpxQ-mediated regulation of *cfa* mRNA, we mutated the

predicted CpxQ binding site (Fig. 2A, 3C). We introduced three point mutations in CpxQ

that should disrupt base pairing to *cfa'-'lacZ*-Long (C5G, C6G, U7A, called CpxQ M2,

Fig. 3C). This CpxQ mutant lost most of the repression activity for the *cfa'-'lacZ*-Long

(Fig. 3D). Mutations in the *cfa* fusion (U103A, C102G, G101C, called *cfa'-'lacZ*-Long M2)

that disrupt base pairing interactions with wild-type CpxQ, likewise eliminated the

repression by wild-type CpxQ (Fig. 3D). We noted that the point mutations in the *cfa*

fusion that disrupt the interaction with CpxQ strongly reduce the basal levels of reporter

activity (~5-fold). This suggests that these nucleotides are important for stability of *cfa*

mRNA. Wild-type levels of repression were not fully restored in the compensatory

mutant pair – CpxQ M2 and *cfa'-'lacZ*-Long M2 (Fig. 3D), suggesting that the mutation in

the fusion already serves to make *cfa* mRNA more sensitive to degradation such that

CpxQ binding has no further impact. Nevertheless, the loss of regulation caused by

individual mutations (Fig. 3D) and the clear CpxQ footprint on *cfa* mRNA (Fig. 2B)

strongly support that CpxQ binds at a site on *cfa* mRNA substantially upstream of the

binding site of the activating sRNAs.

Activating and repressing sRNAs modulate RNase E-dependent degradation of *cfa*

mRNA

Previous studies determined that the RydC-*cfa* mRNA base pairing prevents

RNase E-mediated decay and stabilizes the mRNA to allow increased translation

(Fröhlich et al., 2013). We hypothesize, because ArrS pairs with the same region of *cfa*

mRNA as RydC, that ArrS regulates *cfa* mRNA stability by the same mechanism.

However, the repressive effect of CpxQ must be based on a different mechanism. CpxQ

has been shown to repress other mRNA targets using one of two conserved seed

regions (Chao & Vogel, 2016). One of these seed regions matches the region of CpxQ

that base pairs with *cfa* mRNA (Fig. 3C). CpxQ represses its other targets by base

pairing near the Shine-Dalgarno sequence to prevent ribosome binding and directly

inhibit translation initiation, or by base pairing within the coding region and stimulating

mRNA decay by RNase E (Chao & Vogel, 2016). Since the CpxQ-*cfa* interaction site

within the *cfa* 5'-UTR is far upstream of the Shine-Dalgarno sequence CpxQ must not

repress translation initiation directly. Instead, we wanted to determine if CpxQ caused

cfa mRNA decay by stimulating RNase E activity. When CpxQ is produced from the

native chromosomal locus, it carries a 5' monophosphate (5'P) end because it is a

processed product of the *cpxP* mRNA and not a primary transcript, which would carry a

5' triphosphate (5'PPP). It has been shown that the 5'P end of an sRNA can stimulate

RNase E activity and lead to rapid degradation of the paired mRNA (Bandyra, Said et

al., 2012). It was demonstrated *in vitro* that RNase E-dependent degradation of another

CpxQ target, *nhaB* mRNA, was faster in the presence of 5'P CpxQ compared to 5'PPP

CpxQ (Chao & Vogel, 2016). To determine whether the phosphorylation status of the

CpxQ 5' end impacts its ability to regulate *cfa* mRNA, we designed CpxQ plasmids that

would allow for the production of processed 5'P CpxQ. Three plasmids were used (Fig.

EV3A). All three plasmids contained DNA regions encompassing mature CpxQ and its ρ-

independent terminator. The transcription start site for each plasmid varied (Fig. EV3A).

The CpxQ plasmid (the same as used in experiments described above) placed the +1 of

CpxQ at the native processing site. The CpxQ1 plasmid contained a larger region,

beginning in the *cpxP* coding region (just after the *cpxP* start codon). The CpxQ2

plasmid contained a region starting 30-nt upstream of the processing site in the 3' UTR

of *cpxP*. We expected the CpxQ plasmid would produce CpxQ containing a 5'PPP while

the CpxQ1 and CpxQ2 plasmids would produce a *cpxQ* transcript that would be

processed by RNase E and yield CpxQ with a 5'P. We tested how each plasmid

regulated *cfa'-'lacZ*-Long (Fig. EV3B) and found that both CpxQ1 and CpxQ2 repressed

cfa'-'lacZ-Long activity to the same extent as CpxQ. A northern blot probing for CpxQ

confirmed that CpxQ1 and CpxQ2 were processed to produce the 58nt CpxQ and that

all three plasmids produced the same amount of the 58nt CpxQ sRNA (Fig. EV3C). In

addition, deletion of *rppH* (encoding the 5' pyrophosphatase which converts 5' PPP-

CpxQ to 5' P-CpxQ) in the *cfa'-'lacZ*-Long fusion background did not affect regulation by

CpxQ (Fig. EV3D). These results suggest that the status of the CpxQ 5' end is not

critical for its regulatory activity on *cfa* mRNA.

We anticipated that CpxQ modulates the stability of *cfa* mRNA by making it more

susceptible to cleavage by RNase E. The RNase E cleavage site protected by RydC is

far downstream of the CpxQ binding site. Given the AU-rich nature of the *cfa* 5' UTR, we

hypothesized that there might be additional RNase E cleavage sites that are normally

hidden in secondary structure but are made available to be recognized by RNase E

upon CpxQ binding. We completed a primer extension experiment to profile the mRNA

cleavage products of the *cfa* 5' UTR in response to RydC and CpxQ (Fig. 4A). The two

cfa transcription start sites are the most dominant signals. RydC strongly activates the

longer *cfa*1 mRNA but has no effect on the shorter *cfa*2 transcript (lanes 7-9).

Furthermore, the mRNA cleavage product that corresponds to the RydC-*cfa* interaction

site disappears when RydC is expressed (compare lanes 7 and 9), agreeing with

previous work that this area contains an RNase E cleavage site that is masked upon

formation of the RydC-*cfa* RNA duplex (Fröhlich et al., 2013). CpxQ repressed the

longer *cfa*1 mRNA and had no effect on the shorter *cfa*2 transcript (lanes 10-12). No

other differences in band signals were detected, indicating that we were unable to detect

new CpxQ-dependent mRNA cleavage products.

Another possibility is that CpxQ modulates the stability of *cfa* mRNA by making it

more susceptible to cleavage at the known RNase E recognition site that is protected by

RydC. To test whether RNase E-dependent cleavage is involved in CpxQ-mediated

repression of *cfa*, we made mutations in putative RNase E sites in the *E. coli cfa* 5' UTR

(Fig. 4B). A global study of RNase E cleavage sites predicted a 5-nt minimal RNase E

consensus sequence of "RN↓WUU" (with R as G/A, W as A/U, N as any nucleotide and

↓ as the cleavage site) (Chao, Li et al., 2017). Moreover, in *Salmonella*, mutation of a U

residue 2-nt downstream of the mapped RNase E cleavage site (Fig. EV4) significantly

reduced the cleavage of *cfa* by RNase E (Chao et al., 2017). This uridine is conserved in

E. coli cfa mRNA, so we mutated this U to a G (a non-preferred nucleotide for RNase E

recognition) in the *cfa'-'lacZ*-Long ("G1" mutant, Fig. 4B, C,) and tested for regulation by

RydC and CpxQ. Mutation of the U alone in the G1 mutant did not reduce the basal level

of the fusion compared to WT, suggesting that in *E. coli,* the mutation did not

substantially alter RNase E cleavage of *cfa* mRNA (Fig. 4C). CpxQ and RydC were able

to regulate the G1 mutant similar to the wild-type fusion (Fig. 4C).

In some cases, uridines at the second and third position downstream of an

RNase E cleavage site are highly conserved and mutation of both U residues is required

to fully inhibit cleavage by RNase E (Chao et al., 2017). The sequence surrounding the

RNase E cleavage site in the 5' UTR of *cfa* differs between *Salmonella* and *E. coli* (Fig.

EV4). In *Salmonella*, the sequence surrounding the RNase E cleavage site is GG↓AUA

while in *E. coli* it is GG↓AUU. Changing both U residues to Gs (the G2 mutant, Fig. 4B,

C) increased the basal activity of *cfa'-'lacZ*-Long (~2-fold compared to WT) suggesting

that these mutations inhibit cleavage by RNase E (Fig. 4C). The G2 mutation did not

impair activation by RydC, but substantially impaired CpxQ-mediated repression (Fig.

4C). Mutation of the next two residues downstream of the RNase E cleavage site yielded

"G3" and "G4" mutant fusions (Fig. 4B). The basal levels of activity were further

increased in these fusions, and both activation by RydC and repression by CpxQ were

abrogated (Fig. 4C). These results strongly suggest that the regulation of *cfa* mRNA by

both activating and repressing sRNAs involves modulation of accessibility of a single

RNase E cleavage site that is adjacent to the site for base pairing by activating sRNAs,

but far from the binding site of the repressing sRNA.

Role of Hfq in sRNA-dependent regulation of *cfa* mRNA

In the previous study of RydC-mediated regulation of *cfa* mRNA, it was shown

that swapping the RydC binding site on *cfa* mRNA for the binding site of another Hfq-

dependent sRNA (e.g., RybB or RyhB) reprogrammed the *cfa* mRNA to be activated by

the corresponding sRNA (Fröhlich et al., 2013). However, swapping the RydC binding

site with an Hfq-independent sRNA binding site (i.e., IsrA) only had a small activating

effect (~two-fold vs. ten-fold by RydC) (Fröhlich et al., 2013). This observation suggested

that Hfq plays a key role in sRNA-dependent regulation of *cfa* mRNA stability. An Hfq

binding site at the very 5' end of the UTR of *Salmonella cfa* mRNA (Fig. 5A) was

identified by CLIP-seq (Holmqvist, Wright et al., 2016). To further characterize the

effects of this Hfq binding site on sRNA-dependent regulation of *cfa* mRNA, we deleted

the first 50-nt of the 5'-UTR (-212 to -163, Fig. 5A, B, called *cfa'-'lacZ*-ΔHfqBS) which

includes the Hfq binding site. This fusion had lower basal activity compared to *cfa'-'lacZ*-

Long (~2-fold decrease; Fig. 5B), suggesting that the Hfq binding site located in the -212

to -163 region of the *cfa* mRNA plays a role in *cfa* mRNA structure or stability.

Regulation of *cfa'-'lacZ*-ΔHfqBS by both activating and repressing sRNAs was strongly

impaired even though the fusion retains both sRNA binding sites (Fig. 5B).

To better understand the importance of the -212 to -163 region and Hfq binding

site for sRNA-mediated regulation of *cfa* mRNA, we conducted mutational analysis.

CLIP-seq captured a U to C crosslinking-induced mutation for *Salmonella* Hfq-*cfa*

mRNA interaction as indicated in the "H1" mutant in Fig. 5A (Holmqvist et al., 2016). The

E. coli fusion with the single U to C mutation, *cfa'-'lacZ*-Long H1, had a similar level of

basal activity to wild-type, and was regulated normally by CpxQ and RydC (Fig. 5C). The

sequences of *Salmonella* and *E. coli cfa* mRNAs are not identical in this region (Fig.

EV4). Notably, there is a run of U residues in this region in *E. coli* but not in *Salmonella*.

To test whether the additional U residues in *E. coli cfa* mRNA contribute to regulation by

sRNAs, we also made the "H4" mutant with a total of four U to C substitutions (Fig. 5A).

Basal activity of the *cfa'-'lacZ*-Long H4 fusion was reduced compared to the wild-type

and H1 fusions (Fig. 5C). Moreover, CpxQ no longer regulated the *cfa'-'lacZ*-Long H4

fusion, but regulation by RydC remained at near wild-type levels (Fig. 5C). This result

suggests that Hfq binding at this upstream site may be particularly important for CpxQ-

mediated repression and less important for RydC-mediated activation of *cfa*.

Post-transcriptional regulation of *cfa* mRNA by sRNAs changes membrane lipid

composition

We next wanted to assess the consequences of multiple sRNAs regulating cfa

expression with regard to the physiology of the bacterium. CFA synthase forms CFAs by

transferring a methylene group from S-adenosyl methionine to the double bond of an

UFA of a mature phospholipid that is already incorporated in the membrane. Specifically,

palmitoleic acid (C16:1) is converted into methylene-hexadecanoic acid (C17CFA) and

vaccenic acid (C18:1) is converted into methylene-octadecanoic acid (C19CFA). To test

how post-transcriptional regulation of *cfa* by sRNAs alters the proportion of CFAs in

membrane fatty acids, we conducted gas chromatography on membrane lipids isolated

from strains where each sRNA was ectopically expressed (Fig. 6A, Table S1). The FA

composition of all strains was similar except for CFA and UFA content (Fig. 6A, Table

S1). In strains carrying P$_{lac}$-*cfa,* we detected reduced levels of 16:1 and 18:1 UFAs and

~6-fold increase in levels of C17CFA, compared to the strain carrying the vector control.

Similarly, ArrS-producing cells had ~5-fold higher levels of C17CFA and reduced 16:1

and 18:1 UFAs compared to vector control. RydC-producing cells likewise had reduced

levels of 16:1 and 18:1 UFAs and ~8-fold increased levels of C17CFA compared to the

control strain. CpxQ-producing strains had modestly reduced C17CFA (~2-fold reduced

compared to vector control). These data show that regulation of *cfa* mRNA translation by

each sRNA is correlated with changes in membrane CFA content, implying that the

regulation by sRNAs could contribute to meaningful changes in cell membrane structure

and function.

Role of sRNAs in surviving acid shock

Because CFAs have been implicated in resistance to acidic pH (Brown et al.,

1997, Chang & Cronan, 1999), we next investigated whether sRNAs promote survival

after acid shock. Survival of wild-type, Δ*cfa,* Δ*rydC,* Δ*arrS,* Δ*gadF* and Δ*cpxQ* strains

and complemented mutants was measured after rapid shift of cultures from pH 7 to pH

3. Survival was determined by the ratio of CFUs recovered after acid shock compared to

CFUs before acid shock (Fig. 6B). Compared to the wild-type strain, the Δ*rydC,* and Δ*cfa*

strains had lower survival rates after the acid shock. The Δ*cfa* mutant had the lowest

survival rate of all the strains tested (Δ*cfa* 1.0 ± 0.3 % survival compared to WT 4.3 ± 0.5

%). The Δ*rydC* strain had a survival rate intermediate between that of Δ*cfa* and wild-type

(Δ*rydC,* 2.5 ± 0.4 %) suggesting that RydC might be important for activating *cfa* or other

targets to promote resistance to acid shock. The Δ*arrS,* and Δ*cpxQ* strains showed the

same survival rates as wild-type, implying that these two sRNAs do not play an

important role in regulating *cfa* or other targets to promote acid shock resistance under

these conditions.

Complementation of the *cfa, rydC* and *arrS* mutations by expression of the

corresponding gene from a plasmid resulted in enhanced survival of acid shock

compared to the wild-type strain (Fig. 6B, "striped" bars). Enhanced acid resistance for

ArrS-producing strains was observed previously and was attributed to ArrS positively

regulating *gadE* expression (Aiso et al., 2014). Our data suggest that ArrS-dependent

activation of *cfa* might also contribute to enhanced acid resistance of ArrS-producing

cells*.* The complemented *cpxQ* strain showed survival rates comparable to the wild-type

strain (Fig. 6B). We have shown that ArrS and RydC both activate *cfa* post-

transcriptionally and increase the CFA content in cell membranes but only deletion of

rydC renders otherwise wild-type cells more susceptible to acid shock. These results

suggest that at least RydC-mediated activation of *cfa* translation promotes cell survival

during an acid shock.

Small RNA-dependent regulation of *cfa* at acidic pH

To further investigate sRNA-dependent regulation during acid stress, we

examined sRNA-dependent regulation of *cfa* translational fusions at neutral and acidic

pH. Cells carrying *cfa'-'lacZ*-Long (Fig. 7), *cfa'-'lacZ*-Short or *cfa'-'lacZ*-ΔsRNABS (Fig.

EV5) were grown at either pH 7 or pH 5 and assayed for β-galactosidase activity. For

cfa'-'lacZ-Long, activity was higher at pH 5 compared to pH 7 (Fig. 7). Activity of both

cfa'-'lacZ-Short and *cfa'-'lacZ*-ΔsRNABS was similar at both pH 5 and pH 7 (Fig. EV5).

These observations indicate that activation of *cfa* mRNA in response to acidic pH occurs

post-transcriptionally, and the loss of regulation of a fusion lacking the sRNA binding

sites suggested that one or more sRNAs could be responsible for regulation under these

conditions.

To further probe the effects of sRNA regulation of *cfa* mRNA at acidic pH, the

experiment was performed in wild-type and sRNA mutant backgrounds (Fig. 7). Activity

of *cfa'-'lacZ*-Short was not affected by changes in pH or by deletion of any of the sRNAs

(Fig. EV5). For *cfa'-'lacZ*-Long, wild-type and Δ*arrS* strains all had similar activities at pH

7 and pH 5, with activity at pH 5 higher than at pH 7 (Fig. 7). The Δ*cpxQ* mutant

compared to wild-type had higher *cfa'-'lacZ* activity at pH 7 (Fig. 7). Since ectopic

expression of *cpxQ* repressed *cfa* translation (Fig. 1B), the increased levels of *cfa'-'lacZ*

activity in the Δ*cpxQ* mutant suggests that at neutral pH, CpxQ is produced at sufficient

levels to repress *cfa*. In contrast, at pH 5, the Δ*cpxQ* mutant and wild-type had similar

levels of *cfa'-'lacZ*-Long activity (Fig. 7). The Δ*rydC* mutant had lower *cfa'-'lacZ* activity at

both pH 7 and 5 compared to wild-type, suggesting that RydC may be produced at

sufficient levels at both neutral and acidic pH to have an activating effect on *cfa* under

both conditions. We have shown that ectopic production of ArrS or RydC activates *cfa*

and increases the CFA content in cell membranes, but only deletion of *rydC* affects *cfa'-*

'lacZ-Long activity at pH 5. This observation fits with our previous result that deletion of

rydC renders otherwise wild-type (*cfa*⁺) cells more susceptible to acid shock (Fig. 6B).

To further investigate the interplay of the sRNAs on *cfa* mRNA translation at

different pH values, we deleted *rydC* in combination with each of the other sRNAs in the

cfa'-'lacZ-Long strain (Fig. 7) and measured β-galactosidase activity of the reporter

fusions at pH 5 and 7. Deletion of *arrS* in the Δ*rydC* background had no effect on *cfa'-*

'lacZ-Long activity compared to the Δ*rydC* parent (Fig. 7), suggesting that ArrS does not

play a role in regulation of *cfa* translation under these conditions. Compared with the

Δ*rydC* parent, the Δ*cpxQ* Δ*rydC* strain had slightly higher *cfa'-'lacZ*-Long activity at pH 7

(Fig. 7). In contrast, at pH 5, the Δ*rydC* parent and Δ*cpxQ* Δ*rydC* strain had similar levels

of *cfa'-'lacZ*-Long activity (Fig. 7). Altogether, the data are consistent with the hypothesis

that RydC exerts a positive effect on *cfa* mRNA at both neutral and acidic pH under our

growth conditions. In contrast, CpxQ has a mild repressive effect on *cfa* mRNA only at

neutral pH.

RpoS is not responsible for observed differences in *cfa* mRNA regulation

453   A previous study suggested that transcription from the σ^s-dependent *cfa*

promoter was increased when cells were grown at pH 5 compared to pH 7 (Chang &

Cronan, 1999). Our *cfa'-'lacZ*-Long reporter fusion contains the region encompassing the

σ^s-dependent promoter. To determine if the increased *cfa'-'lacZ*-Long activity we

observed during mild acid stress was dependent on RpoS, we measured *cfa'-'lacZ*–Long

activity at pH 5 and 7 in a Δ*rpoS* background (Fig. 8A). Deletion of *rpoS* had no effect on

cfa'-'lacZ-Long activity at pH 5 or 7 and we observed the same increase in activity in

response to pH 5 in both wild-type and Δ*rpoS* backgrounds (Fig. 8A), indicating that

RpoS does not impact the observed activity of the *cfa'-'lacZ*-Long fusion.

462   We showed that ectopic production of each sRNA, RydC, ArrS, and CpxQ can

regulate *cfa'-'lacZ*-Long (Fig. 1B). While the experiment above suggests that pH-

dependent regulatory effects are not mediated by RpoS, we wanted to further test

whether sRNAs could be acting indirectly on *cfa* via post-transcriptional regulation of

rpoS. We ectopically expressed RydC, ArrS, or CpxQ in wild-type and Δ*rpoS cfa'-'lacZ*-

Long strains and did not observe any differences in sRNA-dependent regulation

between wild-type and Δ*rpoS* (Fig. 8B), indicating that the sRNA-dependent regulation is

not mediated indirectly through RpoS. These data further support the idea that the

regulation of *cfa* by these sRNAs is direct.

Discussion

Microbial membranes are the first line of defense against environmental stress as

well as the site of many metabolic processes. As such, maintenance of membrane

integrity and homeostasis is key to cell survival. Small RNAs (sRNAs) are important

post-transcriptional regulators that play key roles in the response to environmental

stress. We are interested in how sRNAs can contribute to membrane modification in

response to stressors, particularly how sRNAs can lead to altered phospholipid

composition. In the present work, we investigate the roles of multiple sRNAs in post-

transcriptional regulation of *cfa* mRNA, encoding a key enzyme used to alter lipid

composition of bacterial membranes. We show that *cfa* can be post-transcriptionally

regulated by four sRNAs, all of which act on the same long isoform of *cfa* mRNA that

results from transcription from the constitutive σ^{70}-dependent promoter (Fig. 9A). RydC

and ArrS post-transcriptionally activate *cfa*, while the sRNA CpxQ post-transcriptionally

represses *cfa* (Fig.1B). RydC and ArrS bind at a site that overlaps an RNase E cleavage

site, which may prevent RNase E cleavage, increase *cfa* mRNA stability and allow for

increased *cfa* translation. How CpxQ represses *cfa* translation is still unknown. Based on

the location of the CpxQ-*cfa* mRNA base pairing interaction, it is unlikely that the

repression mechanism is similar to those that have already been characterized for

CpxQ. We hypothesized that CpxQ could be working through Hfq or RNase E (Fig. 9A).

Our mutational analyses indicate that both the RNase E recognition site that is adjacent

to the site for base pairing by activating sRNAs (Fig. 4) and an Hfq binding site (Fig. 5)

affect *cfa* mRNA stability and are necessary for CpxQ repression. We propose that

CpxQ base pairing with *cfa* mRNA makes the RNase E cleavage site more susceptible

to degradation. One possibility is that CpxQ:*cfa* base pairing induces a structural change

that makes this RNase E site more susceptible to cleavage. CpxQ may also increase

degradation at this RNase E site by directing Hfq to one binding site at the expense of

another (Fig 9A). RydC requires Hfq to stabilize *cfa* mRNA (Fröhlich et al., 2013) but

does not require the identified Hfq binding site, indicating that there may be another Hfq

binding site in the 5′ UTR of *cfa*. The sRNAs may determine the location of Hfq binding

which in turn may modulate the accessibility of the RNase E cleavage site. Further

studies on RNase E cleavage of *cfa* mRNA in the presence of CpxQ or Hfq CLIP-seq in

E. coli, will give more insight into how Hfq and RNase E affect CpxQ repression of *cfa*

mRNA.

Regulation of *cfa* mRNA by multiple sRNAs suggests that under some

condition(s) these sRNAs influence membrane lipid composition and membrane integrity

to promote fitness or stress resistance. Towards elucidating the physiological roles of

RydC, ArrS, and CpxQ in the regulation of membrane composition, we investigated the

role of each sRNA in acid stress, a condition where CFA synthase is known to play a

role (Brown et al., 1997, Chang & Cronan, 1999). We determined that *cfa* translation

was higher when cells were exposed to pH 5 compared to pH 7 (Fig. 7). When *rydC* was

deleted, *cfa* translation could not reach maximum wild-type levels at pH 5 and Δ*rydC*

mutants were more susceptible to acid shock compared to wild-type, giving us the first

phenotype associated with RydC (Figs. 6B, 7). However, it is not currently known if

activation of *cfa* at acidic pH is the major physiological role of RydC, as the conditions

and signals stimulating RydC production are not yet known.

While ectopic expression of *arrS* increased *cfa* translation and promoted higher

levels of acid resistance (Figs. 1B, 6B), deletion of *arrS* did not affect *cfa* translation or

acid shock survival (Figs. 6B, 7), indicating that ArrS may not play a role in acid stress

under the conditions tested. ArrS has been characterized as an activator of the

transcriptional activator GadE and ectopic expression of ArrS increases acid resistance

though its activation of *gadE* but a deletion phenotype has not been reported (Aiso et al.,

2014, Aiso et al., 2011). ArrS may activate *cfa* under another condition where CFAs in

the membrane contribute to stress resistance (Fig. 9B).

Expression of GadF only slightly regulated *cfa* reporter fusions (Fig. 1B). GadF is

relatively uncharacterized but was shown to activate *acrB* 12-fold compared to the

reported 2-fold activation of *cfa* (Melamed et al., 2016). (Note, in our hands, the

regulation of *cfa* by GadF is even more modest, at ~1.3-fold activation (Fig. 1B).) AcrB is

an efflux pump that uses proton motive force to pump compounds, such as antibiotics,

organic solvents, and detergents, out of the cell (Anes, McCusker et al., 2015). GadF

regulation of *cfa* mRNA may be relevant under these conditions, as CFA membrane

content has been shown to promote resistance to organic solvents (Kanno, Katayama et

al., 2013). Understanding the regulation of GadF itself would provide insight into the

importance of its regulation of *cfa* mRNA, however, it is not currently known how or in

response to what stimuli *gadE* mRNA (from which GadF is derived) is processed (Fig.

9B).

CpxQ is the only sRNA of the four we have characterized that represses *cfa* (Fig.

1B). Deletion of *cpxQ* resulted in increased *cfa* expression at pH 7 compared to wild-

type but did not affect *cfa* at pH 5 (Fig. 7). The physiological consequences of CpxQ-

mediated repression of *cfa* remain unclear. Responding to acid stress is likely not a

relevant physiological role for CpxQ since deletion of *cpxQ* did not affect acid shock

survival (Fig. 6B). However, our observation that CpxQ represses *cfa* translation at

neutral pH combined with what is known about CpxQ may provide some insight. CpxQ is

part of the Cpx two-component system that responds to cell envelope stress (Chao &

Vogel, 2016, Grabowicz et al., 2016). Both the Cpx two-component system and the

processing of CpxQ are induced when cell membrane potential is dissipated (Chao &

Vogel, 2016). The translation of the two targets of CpxQ, an inner membrane sodium-

proton antiporter NhaB and the periplasmic chaperone Skp, is repressed, possibly to

minimize membrane stress and maintain the proton motive force (PMF) across the inner

membrane (Chao & Vogel, 2016, Grabowicz et al., 2016, Grabowicz & Silhavy, 2017).

CpxQ regulation of *cfa* might also help to maintain inner membrane PMF, but the effect
of CFAs on maintenance of membrane potential is currently unknown. A previous study
showed that a lack of CFAs in cellular membranes increases H+ permeability but
decreases H+ extrusion (Shabala & Ross, 2008). Furthermore, the Cpx response is
widely thought to be important for the regulation of energy production and substrate
transport across the inner membrane (Raivio, 2014) and fatty acid composition is known
to affect inner membrane transporter activity and protein function (Lee, 2004, Tveld,
Driessen et al., 1993) but how CFAs affect these functions has not been studied.
However, there is a linear correlation between enzyme activity and membrane fluidity
(Tveld et al., 1993) and, *in vitro*, CFAs are known to increase membrane fluidity by
disrupting lipid packing (Poger & Mark, 2015).

Another possibility that has not been explored is the effect of the Cfa protein and
mechanism of catalysis on inner membrane integrity. During catalysis, one subunit of the
Cfa dimer binds to the membrane while the other subunit extracts and modifies
unsaturated lipids from the membrane (Hari, Grant et al., 2018). Given that the Cpx
response regulates events at the inner membrane and predominantly inhibits the
production of inner membrane localized protein complexes, Cfa's tight association and
interaction inside the inner membrane may trigger the Cpx response. Lastly, PMF is the
cellular energy source and the conversion of UFAs to CFAs by Cfa is an energetically
expensive reaction (Wang & Cronan, 1994) so CpxQ may repress *cfa* translation when
PMF is dissipated simply to reduce cellular ATP consumption. Clearly defining the
impact of Cfa on PMF and determining under what conditions CpxQ differentially
regulates its regulon would elucidate the physiological function of CpxQ regulation of *cfa*
translation.

The UFA to CFA conversion by CFA synthase is energetically expensive, so it is
perhaps not surprising that CFA synthase production is tightly controlled at the

transcriptional level (Kim et al., 2005, Wang & Cronan, 1994) and at the post-

transcriptional level by multiple sRNAs. Regulation of *cfa* mRNA by multiple sRNAs may

have many advantages. There are a few examples of multiple sRNAs acting on the

same target. Post-transcriptional control of the important transcription factor RpoS (σ^s)

by multiple sRNAs to mediate responses to diverse stress conditions is well documented

(Battesti, Majdalani et al., 2011). The *rpoS* gene is post-transcriptionally regulated by at

least four sRNAs in response to different conditions. sRNA regulation of *rpoS* highlights

the potential importance of regulation of the same mRNA target by multiple sRNAs:

synthesis of each sRNA is induced by a different stress which allows RpoS in turn to

properly respond to these various stress signals. We propose that expression of each

sRNA is induced by a different environmental condition that either requires more Cfa, in

the case of RydC, ArrS, and GadF, or less Cfa in the case of CpxQ (Fig. 9B). It is known

that transcription of *arrS* is induced by acid stress and that CpxQ is produced in

response to cell membrane potential dissipation, but how that relates to their regulation

of *cfa* is not understood. The regulation of *rydC* itself has not been elucidated, and it is

not currently known how *gadE* is processed or in response to what stimuli. Uncovering

the stimuli that promote synthesis of ArrS, CpxQ, GadF, and RydC would provide more

insight into how and why *cfa* mRNA is regulated by multiple sRNAs.

Extensive previous work has shown that sRNAs play an integral role in changing

membrane protein composition in response to changing environments (Grabowicz &

Silhavy, 2017, Guillier & Gottesman, 2006, Holmqvist & Wagner, 2017, Vogel &

Papenfort, 2006) but how sRNAs influence membrane fatty acid composition is not well

studied. Here, we build on the first example of an sRNA altering fatty acid composition

(Fröhlich et al., 2013) by showing regulation of *cfa* by four sRNAs. Currently, the sRNAs

studied here are the only examples of sRNAs altering the fatty acid composition of

membranes, which can consequently alter the stability and permeability of the

membrane.

Materials and Methods

Strain and plasmid construction

Strains and plasmids used in this study are listed in Table EV2. All strains used

in this study are derivatives of *E. coli* K12 DJ624 (D. Jin, National Cancer Institute).

Oligonucleotide primers used in this study are listed in Table EV3. Integrated DNA

Technologies synthesized all primers. Δ*rpoS* and Δ*cfa* mutations were made via P1 *vir*

transduction from the Keio collection (Baba, Ara et al., 2006). All other chromosomal

mutations were made using λ Red recombination (Datsenko & Wanner, 2000, Yu, Ellis

et al., 2000) and marked alleles were moved between strains by P1 *vir* transduction

(Miller & H, 1972). Kanamycin markers were removed using pCP20 (Datsenko &

Wanner, 2000).

The *lacZ* translational reporter fusions were constructed using λ Red homologous

recombination into strain PM1205 and counter-selection against *sacB,* as previously

described (Mandin & Gottesman, 2009). Transcription of the fusion is under control of

the P$_{BAD}$ promoter. The fusion contains the *cfa* 5′ UTR fragment of interest and the first

36 codons of the *cfa* open reading frame. *cfa'-'lacZ*-Long, *cfa'-'lacZ*-Short, *cfa'-'lacZ*-

ΔsRNABS, and *cfa'-'lacZ*-ΔHfqBS were constructed by PCR amplifying the *cfa* 5′ UTR

fragment of interest from DJ624 using a fusion- specific forward primer containing

flanking homology to P$_{BAD}$ and the reverse primer O-AK24R, which contains flanking

homology to *lacZ* and anneals in the *cfa* coding region (Table EV3). *cfa'-'lacZ*-Long and

cfa'-'lacZ-Short were constructed by A.M. King (King et al., 2019). *cfa'-'lacZ*-Long H1 and

H4 were constructed by using a forward primer containing the desired point mutations

and flanking homology to P$_{BAD}$ and reverse primer O-AK24R to PCR amplify the *cfa* 5′

UTR fragment of interest from DJ624 (Table EV3). *cfa'-'lacZ*-LongM1 and *cfa'-'lacZ*-

LongM2 were made using Gibson Assembly (Gibson, Young et al., 2009). pBR322 was

PCR amplified using primers with 5' homologies to the *cfa* 5' UTR containing the desired

mutations. Gibson reaction was performed using NEBuilder HiFi DNA Assembly Master

Mix (New England Biolabs) according to the manufacture's protocol. The Gibson product

was transformed into XL10 competent cells and the plasmid was purified. The mutated

cfa 5' UTR was PCR amplified from this plasmid using the same primers used to create

cfa'-'lacZ-Long. The plasmid used to make *cfa'-'lacZ*-LongM1, which contains the G101C

point mutation, was saved as pCB1 for further strain construction. The RNase E

recognition site mutant fusions *cfa'-'lacZ*-LongG1 and G2 were made by mutating pCB1

using QuikChange mutagenesis (Agilent Technologies) and oligos that restore the

G101C mutation to the WT and add the additional desired mutations. The plasmid used

to make *cfa'-'lacZ*-LongG2 was saved as pCB2. *cfa'-'lacZ*-LongG3 and G4 were made

by mutating pCB2 using QuikChange mutagenesis (Agilent Technologies) and oligos

that add the desired mutations. The mutated *cfa* 5' UTR was PCR amplified from each

plasmid using the same primers used to create *cfa'-'lacZ*-Long. All PCR products were

recombined into PM1205 using λ Red homologous recombination and counter-selection

against *sacB* as previously described (Mandin & Gottesman, 2009). All fusions were

verified using DNA sequencing.

Plasmids containing WT sRNAs, mutant sRNAs, and Cfa under the control of the

P_{LlacO} promoter were constructed by PCR amplifying each sRNA from *E. coli* DJ624

chromosomal DNA using oligonucleotides containing HindIII and BamHI restriction sites

(Table EV3). The primers used to create P_{LlacO}-*arrS M1* and P_{LlacO}-*rydC M1* contained

the desired point mutations. PCR products and vector pBRCS12 (Wadler & Vanderpool,

2009) were digested with HindIII and BamHI (New England Biolabs) restriction

endonucleases. Digestion products were ligated using DNA Ligase (New England

Biolabs) and the plasmids were transformed into XL10 competent cells and the plasmid

was purified. P$_{LlacO}$-*cpxQ M2* was created using QuikChange mutagenesis (Agilent

Technologies) of P$_{LlacO}$-*cpxQ* using primers in Table EV3. All plasmids were confirmed

by DNA sequencing.

657
Media and growth conditions:

Bacteria were cultured in LB broth medium or on LB agar plates at 37°C, unless stated

otherwise. When necessary, media were supplemented with antibiotics at following

concentrations: 100 µg ml^{-1} ampicillin (Amp) or 25 µg ml^{-1} kanamycin (Kan). Isopropyl β-

D-1-thiogalactopyranoside (IPTG) was used at 0.1 mM (final concentration) for induction

of expression from P$_{LlacO-1}$ promoter.

β-Galactosidase assays:

Ectopic expression of sRNAs:

Bacterial strains harboring translational *lacZ* reporter fusions carrying the sRNA plasmid

were cultured overnight in Terrific Broth (TB) with Amp and 0.002% L-arabinose then

subcultured 1:100 to fresh TB medium containing Amp and 0.002% L-arabinose. Strains

were grown at 37°C with shaking to early exponential phase then 0.1 mM IPTG was

added to induce sRNA expression. Samples were harvested 60 minutes later and β-

galactosidase assays were then performed as previously described (Miller & H, 1972).

Acid:

Bacterial strains harboring translational *lacZ* reporter fusions were cultured overnight in

TB medium pH 7.0 with 0.002% L-arabinose then subcultured 1:100 to fresh TB medium

pH 7.0 containing 0.002% L-arabinose. Strains were grown at 37°C with shaking for 2

hours then strains were subcultured 1:100 again into TB medium with 0.002% L-

arabinose at pH 7.0 or pH 5.0. Samples were harvested 120 minutes later and β-

galactosidase assays were then performed as previously described (Miller & H, 1972).

Northern Blot

Bacterial strains were cultured in LB medium to OD_{600}~0.4 then plasmids were induced with 0.1 mM IPTG. After 60 minutes, total RNA was extracted by the hot phenol method as described previously (Aiba, Adhya et al., 1981). RNA concentrations were measured spectrophotometrically.15 µg of RNA were denatured for 5 min at 95°C in loading buffer (containing 95% formamide), separated on 8% polyacrylamide urea gel at 100 V for 1 h using 1× Tris-acetate-EDTA (TAE), then transferred to BrightStar™-Plus Positively Charged Nylon Membrane in 0.5× TAE buffer by electroblotting at 50 V for 1 hour at 4°C. RNA was crosslinked to the membrane then membrane was prehybridized for 45 min in ULTRAhyb (Ambion) solution at 42°C. Blots were hybridized overnight with P^{32}-end-labeled primer "CpxQNB" then membranes were washed at 42°C twice with 2× SSC/0.1% SDS for 8 minutes then twice with 0.1× SSC/0.1% SDS for 15 minutes. Signal was detected using film. Membranes were stripped in boiling 0.1% SDS for 10 minutes then re-probed overnight P^{32}-end-labeled primer "5S." Ladder was prepared using the Decade Markers System from ThermoFisher, according to manufacture's instructions.

Analysis of fatty acids

Cells were cultured overnight in LB containing Amp and subcultured 1:100 to fresh LB containing Amp. Cells were grown to an OD_{600} of 0.1 then plasmids were induced with 0.1M IPTG for one hour. Cells were then spun down and the cell pellet was washed twice with MilliQ water. Fatty acids were extracted according to a previously described protocol, method 3.1 (Dionisi, Golay et al., 1999). Pentadecanoic acid methyl ester was used as an internal standard. The bacterial sample was spiked with 500 µg of the internal standard, pentadecanoic acid, and then the sample was transesterified with 2 mL of 0.5M sodium methoxide for 1 min at room temperature. Fatty acid methyl esters

705 (FAMEs) were then extracted using 2 mL of hexane. The solution was centrifugation at

706 2000 rpm for 5 min and the organic upper phase was removed and dried under nitrogen.

707 FAMEs were treated with trimethylsilyldiazomethane in methanol to ensure complete

708 methylation. FAMEs were identified on an Agilent 6890N GC with a 5973 MS and a

709 Zebron WAX column (30m x 0.25mm x 0.25µm).

710
Acid Shock Assay

Strains were grown overnight in LB with Amp then subcultured 1:100 into LB with Amp at

pH 7.0 and grown at 37°C with shaking until OD_{600}= 0.1. Plasmids were induced with

0.1mM IPTG. When strains reached an OD_{600} of 0.2, cultures were diluted 10x into LB

pH 3.0 and incubated at 37°C with no shaking for 60 minutes. Survival was determined

by the ratio of CFUs on the LB plates after acid shock to CFUs on the LB plate before

acid shock.

In vitro **RNA synthesis and structure probing**

DNA templates carrying a T7 promoter sequence for *in vitro* RNA synthesis were

generated by PCR from *E. coli* gDNA (*cfa* mRNA: KFO-0702/KFO-0703; RydC KFO-

0704/KFO-0705; ArrS KFO-0708/KFO-0709; GadF KFO-0710/KFO-0711; CpxQ KFO-

07406/KFO-0707). 200 ng of template DNA were transcribed using the AmpliScirbe T7

Flash Transcription Kit (Epicentre) following the manufacturer's recommendations; size

and integrity of the transcripts were verified on a denaturing 6% PAA gel.

RNA structure probing was carried out as described previously (Fröhlich et al.,

2013) with few modifications. In brief, 0.4 pmol 5' end-labelled *cfa* mRNA were mixed

with 0.4 pmol *E. coli* Hfq protein (provided by K Bandyra and BF Luisi, University of

Cambridge) or Hfq dilution buffer (1X structure buffer, 1% (v/v) glycerol, 0.1% (v/v) Triton

X-100) in the presence of 1X structure buffer (0.01 M Tris pH 7, 0.1 M KCl, 0.01 M

MgCl$_2$) and 1 µg yeast RNA, and samples were incubated at 37°C for 10 min.

Subsequently, unlabelled sRNA (4 pmol) or water was added, and reactions were kept at

37°C for an additional 10 min. Samples were treated with RNase T1 (0.1 U; Ambion,

#AM2283) for 2 min or with lead(II) acetate (final concentration: 5 mM; Sigma #316512)

for 1.5 min, respectively.

Primer extension

Primer extension experiments were performed as previously described (Fröhlich et al.,

2013). A 5' end-labelled primer (KFO-0920) specific to *gfp* was used for reverse

transcription of the samples, as well as for the preparation of a template specific ladder

(prepared using the USB Cycle Sequencing Kit; affymetrix USB #78500; template

amplification pZE-Cat/KFO-0920 on pKF206).

Acknowledgements

We would like to express our gratitude to Dr. Sandy Pernitzsch from Scigraphix for

creating figures. We greatly appreciate discussions and experimental advice from Dr.

Joel Belasco and Dr. Jamie Richards. Thank you to Dr. Alexander Ulanov from the Roy

J. Carver Biotechnology Center for analyzing the fatty acids. We are grateful to

Vanderpool lab members, both past and present, for strains, plasmids, discussions, and

advice. We appreciated Dr. James Slauch and the members of his lab for valuable

discussions. This work was supported by the National Institutes of Health (R01

GM092830) to C.K.V. and supported C.M.B. K.S.F. acknowledges funding by the LMU

Mentoring program of the LMU Faculty of Biology.

Author contributions

C.M.B., K.S.F., and C.K.V. designed research; C.M.B. and K.S.F. performed research;

C.M.B., K.S.F., and C.K.V. analyzed data; and C.M.B., K.S.F., and C.K.V. wrote the

paper.

Conflict of interest

The authors declare that they have no conflict of interest.

762
763
764
765   References:
766

Aiba H, Adhya S, Decrombrugghe B (1981) Evidence For 2 Functional Gal Promoters In
Intact *Escherichia coli* cells. *Journal of Biological Chemistry* 256: 1905-1910

Aiso T, Kamiya S, Yonezawa H, Gamou S (2014) Overexpression of an antisense RNA,
ArrS, increases the acid resistance of Escherichia coli. *Microbiology-Sgm* 160: 954-961

Aiso T, Murata M, Gamou S (2011) Transcription of an antisense RNA of a gadE mRNA
is regulated by GadE, the central activator of the acid resistance system in Escherichia
coli. *Genes to Cells* 16: 670-680

Altuvia S, WeinsteinFischer D, Zhang AX, Postow L, Storz G (1997) A small, stable RNA
induced by oxidative stress: Role as a pleiotropic regulator and antimutator. *Cell* 90: 43-
53

Anes J, McCusker MP, Fanning S, Martins M (2015) The ins and outs of RND efflux
pumps in Escherichia coli. *Frontiers in Microbiology* 6: 14

Antal M, Bordeau V, Douchin V, Felden B (2005) A small bacterial RNA regulates a
putative ABC transporter. *Journal of Biological Chemistry* 280: 7901-7908

Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M,
Wanner BL, Mori H (2006) Construction of Escherichia coli K-12 in-frame, single-gene
knockout mutants: the Keio collection. *Molecular Systems Biology* 2: 11

Bandyra KJ, Said N, Pfeiffer V, Gorna MW, Vogel J, Luisi BF (2012) The Seed Region of
a Small RNA Drives the Controlled Destruction of the Target mRNA by the
Endoribonuclease RNase E. *Molecular Cell* 47: 943-953

Battesti A, Majdalani N, Gottesman S (2011) The RpoS-Mediated General Stress
Response in Escherichia coli. In *Annual Review of Microbiology, Vol 65*, Gottesman S,
Harwood CS (eds) pp 189-213. Palo Alto: Annual Reviews

Bordeau V, Felden B (2014) Curli synthesis and biofilm formation in enteric bacteria are
controlled by a dynamic small RNA module made up of a pseudoknot assisted by an
RNA chaperone. *Nucleic Acids Research* 42: 4682-4696

Brown JL, Ross T, McMeekin TA, Nichols PD (1997) Acid habituation of *Escherichia coli*
and the potential role of cyclopropane fatty acids in low pH tolerance. *International*
Journal of Food Microbiology 37: 163-173

Chang YY, Cronan JE (1999) Membrane cyclopropane fatty acid content is a major
factor in acid resistance of *Escherichia coli*. *Molecular Microbiology* 33: 249-259

Chao YJ, Li L, Girodat D, Forstner KU, Said N, Corcoran C, Smiga M, Papenfort K,
Reinhardt R, Wieden HJ, Luisi BF, Vogel J (2017) In Vivo Cleavage Map Illuminates the
Central Role of RNase E in Coding and Non-coding RNA Pathways. *Molecular Cell* 65:
39-51

Chao YJ, Vogel J (2016) A 3 ' UTR-Derived Small RNA Provides the Regulatory
Noncoding Arm of the Inner Membrane Stress Response. *Molecular Cell* 61: 352-363

Datsenko KA, Wanner BL (2000) One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proceedings of the National Academy of*
Sciences of the United States of America 97: 6640-6645

Dionisi F, Golay PA, Elli M, Fay LB (1999) Stability of cyclopropane and conjugated
linoleic acids during fatty acid quantification in lactic acid bacteria. *Lipids* 34: 1107-1115

Fröhlich KS, Papenfort K, Fekete A, Vogel J (2013) A small RNA activates CFA
synthase by isoform-specific mRNA stabilization. *Embo Journal* 32: 2963-2979

Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO (2009)
Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nature*
Methods 6: 343-U41

Glickman MS, Cox JS, Jacobs WR (2000) A novel mycolic acid cyclopropane synthetase
is required for cording, persistence, and virulence of *Mycobacterium tuberculosis*.
Molecular Cell 5: 717-727

Grabowicz M, Koren D, Silhavy TJ (2016) The CpxQ sRNA Negatively Regulates Skp
To Prevent Mistargeting of beta-Barrel Outer Membrane Proteins into the Cytoplasmic
Membrane. *Mbio* 7: 8

Grabowicz M, Silhavy TJ (2017) Envelope Stress Responses; An Interconnected Safety
Net. *Trends in Biochemical Sciences* 42: 232-242

Grogan DW, Cronan JE (1997) Cyclopropane ring formation in membrane lipids of
bacteria. *Microbiology and Molecular Biology Reviews* 61: 429-&

Guillier M, Gottesman S (2006) Remodelling of the Escherichia coli outer membrane by
two small regulatory RNAs. *Molecular Microbiology* 59: 231-247

Hari SB, Grant RA, Sauer RT (2018) Structural and Functional Analysis of E-coli
Cyclopropane Fatty Acid Synthase. *Structure* 26: 1251-+

Holmqvist E, Wagner EGH (2017) Impact of bacterial sRNAs in stress responses.
Biochemical Society Transactions 45: 1203-1212

Holmqvist E, Wright PR, Li L, Bischler T, Barquist L, Reinhardt R, Backofen R, Vogel J
(2016) Global RNA recognition patterns of post-transcriptional regulators Hfq and CsrA
revealed by UV crosslinking in vivo. *Embo Journal* 35: 991-1011

Jin Y, Watt RM, Danchin A, Huang JD (2009) Small noncoding RNA GcvB is a novel
regulator of acid resistance in Escherichia coli. *Bmc Genomics* 10: 7

Kanno M, Katayama T, Tamaki H, Mitani Y, Meng XY, Hori T, Narihiro T, Morita N,
Hoshino T, Yumoto I, Kimura N, Hanada S, Kamagata Y (2013) Isolation of Butanol- and
Isobutanol-Tolerant Bacteria and Physiological Characterization of Their Butanol
Tolerance. *Applied and Environmental Microbiology* 79: 6998-7005

Kim BH, Kim S, Kim HG, Lee J, Lee IS, Park YK (2005) The formation of cyclopropane
fatty acids in Salmonella enterica serovar Typhimurium. *Microbiology-Sgm* 151: 209-218

King AM, Vanderpool CK, Degnan PH (2019) sRNA Target Prediction Organizing Tool
(SPOT) Integrates Computational and Experimental Data To Facilitate Functional
Characterization of Bacterial Small RNAs. *mSphere* 4

Law JH (1971) Biosynthesis of Cyclopropane Rings. *Accounts of Chemical Research* 4:
199-&

Lee AG (2004) How lipids affect the activities of integral membrane proteins. *Biochimica*
Et Biophysica Acta-Biomembranes 1666: 62-87

Mandin P, Gottesman S (2009) A genetic approach for finding small RNAs regulators of
genes of interest identifies RybC as regulating the DpiA/DpiB two-component system.
Molecular Microbiology 72: 551-565

Mann M, Wright PR, Backofen R (2017) IntaRNA 2.0: enhanced and customizable
prediction of RNA-RNA interactions. *Nucleic Acids Research* 45: W435-W439

Melamed S, Peer A, Faigenbaum-Romm R, Gatt YE, Reiss N, Bar A, Altuvia Y,
Argaman L, Margalit H (2016) Global Mapping of Small RNA-Target Interactions in
Bacteria. *Molecular Cell* 63: 884-897

Miller, H J (1972) *Experiments in Molecular Genetics*. Cold Spring Harbor Laboratory
Press, Cold Spring Harbor Laboratory, New York

Poger D, Mark AE (2015) A Ring to Rule Them All: The Effect of Cyclopropane Fatty
Acids on the Fluidity of Lipid Bilayers. *Journal of Physical Chemistry B* 119: 5487-5495

Raivio TL (2014) Everything old is new again: An update on current research on the Cpx
envelope stress response. *Biochimica et biophysica acta* 1844: 1529-41

Shabala L, Ross T (2008) Cyclopropane fatty acids improve *Escherichia coli* survival in
acidified minimal media by reducing membrane permeability to H+ and enhanced ability
to extrude H+. *Research in Microbiology* 159: 458-461

Tveld GI, Driessen AJM, Konings WN (1993) Bacterial solute transport proteins in their
lipid environment. *Fems Microbiology Reviews* 12: 293-314

Vogel J, Papenfort K (2006) Small non-coding RNAs and the bacterial outer membrane.
Current Opinion in Microbiology 9: 605-611

Wadler CS, Vanderpool CK (2009) Characterization of homologs of the small RNA SgrS
reveals diversity in function. *Nucleic Acids Research* 37: 5477-5485

Wang AY, Cronan JE (1994) The Growth Phase-dependent Synthesis of Cyclopropane
Fatty Acids in Escherichia Coli Is the Result of an Rpos(Katf)-dependent Promoter Plus
Enzyme Instability. *Molecular Microbiology* 11: 1009-1017

Yu DG, Ellis HM, Lee EC, Jenkins NA, Copeland NG, Court DL (2000) An efficient
recombination system for chromosome engineering in Escherichia coli. *Proceedings of
the National Academy of Sciences of the United States of America* 97: 5978-5983

Zhang AX, Altuvia S, Tiwari A, Argaman L, Hengge-Aronis R, Storz G (1998) The OxyS
regulatory RNA represses rpoS translation and binds the Hfq (HF-I) protein. *Embo
Journal* 17: 6061-6068

Zhang YM, Rock CO (2008) Membrane lipid homeostasis in bacteria. *Nature Reviews
Microbiology* 6: 222-233

Figure Legends:

Figure 1: *cfa* expression is controlled by multiple sRNAs.
(A) *cfa* has two promoters. Transcription from the distal promoter is σ^{70}-dependent and yields a longer *cfa* transcript with a 212-nt 5′ UTR. Transcription from the proximal promoter is controlled by σ^s and produces a shorter *cfa* transcript with a 34-nt 5′ UTR.

(B) A *cfa* translational fusion to *lacZ* (controlled by P_{BAD} promoter) was constructed. *cfa'-'lacZ*-Long fusion is from the distal σ^{70}-dependent promoter which contains a 212-nt 5′ UTR that includes the RydC binding site (green square labeled "RydC") and the sites predicted for the ArrS, CpxQ, and GadF. *cfa'-'lacZ*-Long carrying an empty or a P_{lac}-sRNA plasmid were grown in TB medium with 0.002% L-arabinose to early exponential phase then sRNA expression was induced with 0.1 mM IPTG. Samples were harvested 60 minutes later and assayed for β-galactosidase activity of the reporter fusion. The error bars are standard deviations, and the statistical significance was determined using the Student's *t*-test. *$P < 0.05$, **$P < 0.005$, ***$P < 0.0005$, ****$P < 0.0001$, ns is for not significant. The Student's *t*-test was performed comparing fusion activity in response to expression of each sRNA to fusion activity in vector control.

(C) A *cfa'-'lacZ*-Short fusion was constructed, which contains only proximal the σ^s-dependent promoter and consequently not the predicted sRNA binding sites. Regulation of *cfa'-'lacZ*-Short by each sRNA was determined as described in 1B.

Figure 2: Activating and repressing sRNAs bind different sites on *cfa* mRNA.
(A) Predicted base pairing sites of sRNAs on *cfa* mRNA.

(B) *In vitro* structure probing using 5′ end-labeled *cfa* mRNA with RNase T1 (lanes 4–8) and lead(II) acetate (lanes 9–13) in the presence Hfq (20 nM) and each sRNA (200 nM). RNase T1 and alkaline ladders of cfa mRNA were used to map cleaved fragments. Positions of G-residues are indicated relative to the translational start site. Each sRNA binding site is marked with a black line right of the lane.

Figure 3: Activating and repressing sRNAs bind different sites on *cfa* mRNA
(A) Predicted base pairing between *cfa* mRNA and ArrS or RydC. The red nucleotides were mutated to test base pairing.

(B) Left: Mutant sRNAs (ArrS M1, RydC M1) and WT sRNAs were tested for activation of *cfa'-'lacZ*-Long as described in Fig. 1B.

(B) Right: One point mutation (G101C) were made in *cfa'-'lacZ*-Long (called *cfa'-'lacZ*-Long M1). Mutant sRNAs (ArrS M1, RydC M1) and WT sRNAs were tested for activation of *cfa'-'lacZ*-Long M1 as described in Fig. 1B.

(C) Base pairing between *cfa* mRNA and CpxQ. The red nucleotides were mutated to test base pairing.

(D) Left: Mutant CpxQ M2 and WT CpxQ were tested for activation of *cfa'-'lacZ*-Long as described in Fig. 1B.

934 (D) Right: Three point mutations (A131T, G130C, G129C) were made in *cfa'-'lacZ*-Long
935 (called *cfa'-'lacZ*-Long M2). Mutant CpxQ M2 and WT CpxQ were tested for activation of
936 *cfa'-'lacZ*-Long M2 as described in Fig. 1B.
937
938
Figure 4: **Activating and repressing sRNAs modulate RNase E-dependent**
degradation of *cfa* mRNA
(A) Primer extension of 5′ UTR of *cfa* mRNA reveals no CpxQ- dependent mRNA
cleavage products. RNA samples were withdrawn prior to and 5 or 15 minutes after
arabinose addition to induce vector control (lanes 5-6), RydC (lanes 7-9), or CpxQ (lanes
10-12). RNA samples were used as templates for primer extension using a 5′ end-
labeled *cfa* primer. Transcripts were identified using *cfa*-specific sequencing ladders
(lanes 1-4). Transcription start site 1 (TSS1) and 2 (TSS2) and the known RNase E
cleavage site are marked with solid arrows.
948
949 B) Multiple mutations of the RNase E recognition site on *cfa* mRNA were made in the
950 *cfa'-'lacZ*-Long fusion background (called G1-G4). Black arrow indicates RNase E
951 cleavage site. CpxQ and RydC base pairing locations are indicated.
952
953 (C) Regulation of *cfa'-'lacZ*-Long (WT) or *cfa'-'lacZ*-Long G1-G4 by CpxQ and RydC was
954 tested as described in Fig. 1B.
955
956
957 **Figure 5: Mutating a putative Hfq binding site alters *cfa* mRNA regulation by CpxQ**
958 **and RydC**
959 (A) 5′ UTR of *cfa* gene with a putative Hfq binding site and sRNA base pairing sites
960 indicated.
961
962 (B) The first 50-nt of the 212-nt- 5′ UTR (-212 to -163) which includes the putative Hfq
963 binding site, was deleted in *cfa'-'lacZ*-Long fusion (called *cfa'-'lacZ*-ΔHfqBS). Regulation
964 of *cfa'-'lacZ*-Long or *cfa'-'lacZ*-ΔHfqBS by CpxQ and RydC was tested as described in
965 Fig. 1B.
966
967 (C) Mutations of the Hfq binding site on *cfa* mRNA (shown in A) were made in the *cfa'-*
968 *'lacZ*-Long fusion background (called *cfa'-'lacZ*-Long H1 and *cfa'-'lacZ*-Long H4).
969 Regulation of WT and mutant fusions by RydC and CpxQ was tested as described in
970 Figure 1B.
971
972
Figure 6: Expression of sRNAs can alter fatty acid composition and survival to
acid shock
(A) Relative qualification of fatty acids in *E. coli* in response to ectopic expression of
vector control, *cfa*, *rydC*, *arrS*, or *cpxQ*. Fatty acids are presented as a percent of total
identified fatty acids. Error represents average ± standard deviation, n=3 or *n=2. ND:
not detected. The error bars are standard deviations, and the statistical significance was
determined using the Student's *t*-test. *$P < 0.05$, **$P < 0.005$, ***$P < 0.0005$,
****$P < 0.0001$, ns is for not significant. The Student's *t*-test was performed comparing
the amount of each fatty acid in response to expression of the sRNA to fatty acid amount
in vector control.
983

(B) Survival during an acid challenge. When strains reached an OD_{600} of 0.2, cultures
were diluted into LB pH 3 and incubated for 60 minutes. Survival was determined by the
ratio of CFUs on the LB plates after acid shock to CFUs on the LB plate before acid
shock. Error bars represent average ± standard deviation of 3 technical replicates. The
error bars are standard deviations, and the statistical significance was determined using
the Student's t-test. *$P < 0.05$, **$P < 0.005$, ***$P < 0.0005$, ****$P < 0.0001$, ns is for not
significant. The Student's t-test was performed comparing acid survival of each mutant
to the survival of vector control.
992
993
Figure 7: *cfa* translation is induced at acidic pH.
Cells carrying *cfa'-'lacZ*-Long in either WT background, a background where one sRNA
is deleted, or a background where *rydC* and one other sRNA are deleted were grown in
medium at pH 7 then subcultured into medium either pH 7 (pH 7.0) or pH 5 (pH 5.0).
Samples were harvested 120 minutes later and assayed for β-galactosidase activity of
the reporter fusion. Error bars represent standard deviation for three biological
replicates. A one-way ANOVA with post hoc Tukey's test was performed; the same letter
indicates no significant differences between fusion activities ($P \leq 0.05$).
1002
1003
1004 **Figure 8: Deletion of *rpoS* does not affect *cfa*'-'*lacZ*-Long activity**
1005 (A) Cells carrying *cfa'-'lacZ*-Long in a WT or Δ*rpoS* background were grown and *cfa*
1006 translation in response to acid was assayed as described in Figure 7.
1007
1008 (B) *cfa'-'lacZ*-Long in a WT or Δ*rpoS* background carrying an empty, P_{lac}-*arrS,* P_{lac}-*cpxQ,*
1009 P_{lac}-*gadF,* or P_{lac}-*rydC* plasmid were grown as described in Figure 1B.
1010
1011
1012
Figure 9: Models of sRNAs regulation of *cfa* mRNA
(A) Working model for the role of Hfq, RNAse E and sRNAs in regulation of *cfa* mRNA.
There are multiple RNase E cleavage sites that have been mapped in *Salmonella*.
Activating sRNAs, especially RydC, require Hfq for efficient pairing and inhibition of
RNAse E cleavage at one site that may be the primary cleavage site, and the pairing
stabilizes *cfa* mRNA. CpxQ may have the opposite effect – somehow pairing of CpxQ
(facilitated by Hfq), stimulates RNAse E-dependent cleavage especially at the site RydC
is known to protect.
1021
(B) Regulation of *cfa* mRNA by these sRNAs provides a sigmaS-independent
mechanism for control of cfa, and thus a way to modulate lipid composition of the
bacterial cell envelope in response to a variety of different stresses. CpxQ is part of the
Cpx regulon, which responds to signals like disrupted IM potential or IM stress. The
conditions that stimulate RydC production are unknown. ArrS and GadF are linked to
acid stress, and CFAs have previously been implicated in acid resistance. Modulating
the composition of cyclopropanated FA in response to these stresses could alter
membrane properties like fluidity and permeability and allow the cell to adapt to the
stress.
1031
1032
1033

1034
1035   **Expanded View Figures-**
1036
1037   **Figure EV1: Testing of additional sRNAs for regulation of *cfa* expression.**
1038   (A) The *cfa'-'lacZ*-Long fusion contains the entire 212-nt 5′ UTR, including the RydC
1039   binding site (indicated by green square labeled "RydC"). Regulation of *cfa'-'lacZ*-Long by
1040   RydC, OxyS, GcvB, and CpxQ was determined as described in Fig. 1B.
1041
1042   (B) The *cfa'-'lacZ*-Short fusion contains only proximal the σ^s-dependent promoter.
1043   Regulation of this fusion by RydC, OxyS, GcvB, and CpxQ was determined as described
1044   in Fig. 1B.
1045
1046
1047   **Figure EV2: Deletion of the putative sRNA binding site removes regulation of *cfa***
1048   **translation by ArrS, CpxQ, GadF, and RydC.**
1049   (A) 5′ UTR of *cfa* gene. Arrows mark transcriptional start sites and sRNA binding sites
1050   are indicated with labeled boxes.
1051
1052   (B) A *cfa* translational fusion to *lacZ* (*cfa'-'lacZ*-ΔsRNABS) that begins immediately
1053   downstream of the predicted sRNA binding site was constructed. This fusion contains
1054   the proximal σ^s-dependent promoter and 79-nt upstream of this promoter.
1055
1056   (C) Regulation of *cfa'-'lacZ*-ΔsRNABS by ArrS, CpxQ, GadF, or RydC was tested as
1057   described in Figure 1B.
1058
1059
1060   **Figure EV3: CpxQ does not require a 5' monophosphate to repress *cfa*.**
1061   (A) Three CpxQ plasmids were constructed: CpxQ encodes CpxQ directly from the
1062   CpxQ start site, CpxQ1 starts right after the "AUG" of CpxP and CpxQ2 starts 30-nt
1063   upstream of the RNase E cleavage site in the 3' UTR of CpxP. Plasmids were designed
1064   so that CpxQ1 and CpxQ2 would be cleaved by RNase E to yield 5′P CpxQ.
1065
1066   (B) Regulation of *cfa'-'lacZ*-Long was tested for each sRNA plasmid as described in Fig.
1067   1B.
1068
1069   (C) Expression levels of CpxQ from each plasmid as described in (A) were determined
1070   by northern blot analysis of total RNA samples. 5S served as loading control.
1071
1072   (D) Regulation by CpxQ was tested for *cfa'-'lacZ*-Long in a WT or Δ*rppH* background as
1073   described in Fig. 1B.
1074
1075
1076   **Figure EV4: The *cfa* 5′ UTR differs between *E. coli* and *Salmonella***
1077   Sequence alignment of the promoter and 5′ UTR of *cfa* in *E. coli* and *Salmonella*. Gray
1078   box: Hfq binding site; Orange box: CpxQ binding site. Green box: Activating sRNA
1079   binding site. Black arrow: RNase E cleavage site.
1080
1081
1082
1083

Figure EV5: Deletion of sRNAs do not affect *cfa'-'lacZ*-Short activity.

(A) Cells carrying *cfa'-'lacZ*-Short were grown and *cfa* translation in response to acid was assayed as described in Figure 7.

(B) Cells carrying *cfa'-'lacZ*-ΔsRNABS were grown and *cfa* translation in response to acid was assayed as described in Figure 7.

(C) Cells carrying *cfa'-'lacZ*-Short in either WT background or a background where one sRNA is deleted were grown and *cfa* translation in response to acid was assayed as described in Figure 7.

(D) Cells carrying *cfa'-'lacZ*-Short (in either WT background or a background where *rydC* and one other sRNA are deleted) were grown as described in Figure 7. *ΔrydC* single mutant is included for reference. Error bars represent standard deviation for three biological replicates. A one-way ANOVA with post hoc Tukey's test was performed; the same letter indicates no significant differences between fusion activities ($P \leq 0.05$).

Supplemental Tables/ Excel files

Table S1: Expression of sRNAs can alter fatty acid composition
Relative qualification of fatty acids in *E. coli* in response to ectopic expression of *arrS*, *cfa*, *cpxQ*, *rydC* or vector control. Fatty acids are presented as a percent of total identified fatty acids. Error represents average ± standard deviation, n=3 or *n=2. ND: not detected

Table S2: Plasmids and Strains used in this study

Table S3: Oligos used in this study

A

B

cfa'-'lacZ-Long

C

cfa'-'lacZ-Short

Figure 1

Figure 2

A

```
      -105            -95
       |               |
       C
5' CGUCGGAUUAU 3'   cfa'-'lacZ-Long M1
   :|||||||||:       cfa'-'lacZ-Long
3' UUAGCCUAAUG 5'   ArrS
       G             ArrS M1
   |               |
   10              1
```

```
      -109      -99
       |         |
            C
5' UUCUACGUCGGAU 3'   cfa'-'lacZ-Long M1
   |||||:|||||         cfa'-'lacZ-Long
3' CAGAUGUAGCCUU 5'   RydC
            G         RydC M1
   |               |
   12              2
```

C

```
      -135    -128
       |       |
          UCC
5' UCGGCAAGGGU 3'   cfa'-'lacZ-Long M2
   ||||||||:         cfa'-'lacZ-Long
3' UACCGUUCCUU 5'   CpxQ
          AGG         CpxQ M2
   |           |
   11          4
```

B

cfa'-'lacZ-Long

cfa'-'lacZ-Long M1

D

cfa'-'lacZ-Long

cfa'-'lacZ-Long M2

Figure 3

Figure 4

A

B

cfa'-'lacZ-Long

C

cfa'-'lacZ-Long

Figure 5

A

B

Figure 6

Figure 7

cfa'-'lacZ-Long

B

cfa'-'lacZ-Long

Figure 8

Figure 9

A

B

Expanded View Figure 1

A

B

cfa'-'lacZ-ΔsRNABS

Expanded View Figure 2

Expanded View Figure 3

Expanded View Figure 4

A *cfa'-'lacZ*-Short

B *cfa'-'lacZ*-ΔsRNABS

C *cfa'-'lacZ*-Short

D *cfa'-'lacZ*-Short

Expanded View Figure 5

Supplemental Table 1

Fatty Acid	p_{lac}				
	arrS	*cfa**	*cpxQ*	*rydC**	*vector*
C12:0	2.5 ± 1.1	1.8 ± 0.7	1.0 ± 0.1	5.2 ± 0.5	1.29 ± 0.08
C14:0	7.4 ± 1.5	6.0 ± 1.0	3.1 ± 0.1	12.6 ± 3.8	3.6 ± 0.4
C14:1	ND	ND	0.18 ± 0.07	ND	ND
C16:0	43.9 ± 1.1	43.3 ± 4.0	30.7 ± 1.6	42.8 ± 0.3	33.6 ± 1.7
C16:0 methyl	0.8 ± 0.6	0.48 ± 0.08	0.8 ± 0.2	ND	0.9 ± 0.2
C16:1	13.5 ± 1.5	10.9 ± 0.7	19.9 ± 1.1	6.2 ± 0.4	18.9 ± 0.7
C17:0 CFA	11.8 ± 1.0	13.7 ± 1.2	1.2 ± 0.3	17.3 ± 3.8	2.2 ± 0.4
C18:0	12.2 ± 1.6	14.4 ± 2.1	25.4 ± 3.6	9.8 ± 2.0	21.7 ±0.6
C18:1	7.9 ± 0.7	9.1 ± 2.0	13.8 ± 1.7	6.0 ±0.7	14.9 ± 0.9
C18:2	ND	ND	ND	ND	ND
C19:0 CFA	ND	0.9 ± 0.3	ND	ND	ND
C20:0	ND	ND	1.1 ± 0.3	ND	0.9 ± 0.1
C22:0	ND	ND	1.1 ± 0.3	ND	0.7 ± 0.2
C24:0	ND	ND	1.1 ± 0.3	ND	0.8 ±0.2
C26:0	ND	ND	0.6 ± 0.3	ND	0.3 ± 0.2