Delegating Sex: differential gene expression in stolonizing syllids uncover the hormonal control of reproduction in Annelida

Patricia Álvarez-Campos[1, 2, 3,] *, Nathan J. Kenny[4], Aida Verdes[1, 5, 6], Rosa Fernández[7], Marta Novo[8], Gonzalo Giribet[2, †], and Ana Riesgo[4,] *, †

[1]*Departamento de Biología (Zoología), Universidad Autónoma de Madrid, Facultad de Ciencias, Cantoblanco 28049, Madrid, Spain*

[2]*Museum of Comparative Zoology, Department of Organismic and Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, MA 02138, USA.*

[3]*Present address: Stem Cells, Development and Evolution, Institut Jacques Monod, CNRS UMR 7592, Université Paris-Diderot;15 rue Hélène Brion 75205 Paris cedex 13, France*

[4]*Department of Life Sciences, The Natural History Museum of London, Cromwell Road, London SW7 5BD, United Kingdom*

[5]*Department of Biology, The Graduate Center, City University of New York, 365 5th Ave, New York, NY, 10010, USA*

[6]*Sackler Institute for Comparative Genomics, American Museum of Natural History, Central Park West at 79th Street, New York, NY 10024-5192, USA*

[7] *Bioinformatics & Genomics Unit, Center for Genomic Regulation, Carrer del Dr. Aiguader 88, 08004 Barcelona, Spain*

[8]*Environmental Toxicology and Biology Group. Departamento de Física Matemática y de Fluídos. Facultad de Ciencias, UNED. Senda del Rey 9, 28040 Madrid, Spain.*

* Corresponding author: patricia.alvarez@uam.es

[†] Senior authors

Abstract

Stolonization in syllid annelids is a unique mode of reproduction among animals. During the breeding season, a structure resembling the adult but containing only gametes, called stolon, is formed at the posterior end of the animal. When the stolons mature, they detach from the adult and the gametes are released into the water column. The process is synchronized within each species, and it has been reported to be under environmental and endogenous control, probably via endocrine regulation. To further understand the reproduction in syllids and to elucidate the molecular toolkit underlying stolonization, we generated Illumina RNA-seq data from different tissues of reproductive and non-reproductive individuals of *Syllis magdalena*, and characterized gene expression during the stolonization process. Several genes involved in gametogenesis (*ovochymase*, *vitellogenin*, *testis-specific serine/threonine-kinase*), immune response (*complement receptor 2*), neuronal development (*tyrosine-protein kinase Src42A*), cell proliferation (*alpha-1D adrenergic receptor*), and steroid metabolism (*hydroxysteroid dehydrogenase* 2) were found differentially expressed in the different tissues and conditions analyzed. In addition, our findings suggest that several neurohormones, such as methyl farnesoate, dopamine and serotonin, might trigger the stolon formation, the correct maturation of gametes and the detachment of stolons when gametogenesis is complete. The process seems to be under circadian control, as indicated by the expression patterns of *r-opsins*. Overall, our results shed light into the genes that orchestrate the onset of gamete formation, and improve our understanding of how some hormones, previously reported to be involved in reproduction and metamorphosis processes in other invertebrates, seem to also regulate reproduction via stolonization.

Introduction

Annelids in the family Syllidae have a remarkable reproductive strategy, which has attracted the attention of many biologists (e.g. Nygren 1999 and references herein). Syllids exhibit epitoky, which largely implies morphological changes associated with reproduction (Malaquin 1893), and can be further divided into a variety of reproductive modes. In all epitokous modes, there are two states: the sexually immature worm, called an "atoke", and the sexually mature worm, or "epitoke". Among the epitokous types of reproduction, one of the most common is epigamy, which is not exclusive to syllids, where the entire atoke transforms into the epitoke, developing swimming chaetae, enlarging its eyes and undergoing changes in musculature (Wissocq 1970; Daly 1975; Garwood 1991). One of the most extreme types of epitokous reproduction is squizogamy or stolonization, where only a part of the individual transforms into an epitokal sexual stage, either by generating new segments or by differentiating pre-existing ones (Franke 1999). When the breeding season approaches, the syllid atoke (or stock) starts to develop a peculiar structure at the end of its body, that resembles the adult and is known as the stolon (Agassiz 1863) (Fig. 1). The stolons possess several features similar to the stock, such eyes and antennae, but are filled with gametes (Figs. 1 and 2A–E), as their brief existence is exclusively devoted to mating, followed by death (Franke 1999). The stock produces and transfers the gametes to the stolon, which is released from the stock when mature (with developed eyes and antennae) (Figs. 1 and 2E), and swims to the surface to spawn (Potts 1911; Mesnil and Caullery 1919). The pelagic stolon releases gametes into the water column, via the nephridiopores in the case of sperm, and through rupture of the body wall for the eggs (Franke 1980; Wissocq 1966, 1970; Schroeder and Hermans 1975; Okada 1937; Durchon 1951, 1952, 1959). Finally, before or after stolon detachment (depending on the species), the stock regenerates the lost final segments (e.g., Marion and Bobretsky 1875; Michel 1898; Okada 1929) (Fig. 1 and 2F).

The molecular toolkit involved in annelid reproduction is still far from being understood, although studies in several annelid species have shed some light into the matter. For instance, genes involved in pheromone production that are essential for mate recognition and spawning, such as *Temptin* and *Attractin*, and those involved in gametogenesis or fertilization, such as *Fertilin* or *Acrosin*, have been identified in *Spirobranchus* (*Pomatoceros*) *lamarckii, Hormogaster samnitica* and *H. elisae* (e.g., Kang et al. 2002; Rivera et al. 2005; Takahashi et al. 2009; Novo et al. 2013). It is also well-known that the germline specification in the marine annelids *Alitta virens, Platynereis dumerilii, Capitella teleta* or *Hermodice carunculata* involves the expression of several genes including *vasa, nanos*, and *piwi* during embryogenesis, and that *vitellogenin* (*Vtg*) is required for yolk formation in the oocyte (Hafer et al.1992; Dill and Seaver 2008; Rebscher et al. 2007; Thamm and Seaver 2008; Giani et al. 2011; Mehr et al. 2015; Schenk et al. 2016). Interestingly, a recent study has reported the potential involvement of the

sesquiterpenoid methyl farnesoate (MF), the brain neurohormone that directly regulates *Vtg* in yolk production of *P. dumerilii* females, therefore influencing the correct development of oocytes (Schenk et al. 2016). Particularly, a decrease in MF levels in the brain of *P. dumerilii* during reproduction allowed oocyte maturation but suppressed normal somatic functions and caudal regenerative capacities (Schenk et al. 2016). In crustaceans, MF has been showed to play essential roles in development and reproduction (Xie et al. 2016), similar to the role of Juvenile Hormone (JH) in insects (Riddiford 1994; Wyatt and Davey 1996). Other hormones have also been proposed to play essential roles in annelids reproduction, such as the prostomium (i.e. first pre-oral segment of the animal) hormone 8,11,14-eicosatrienoic acid, which seems to be responsible for sperm maturation and spawning in *Arenicola marina* males (Bentley 1985; Bentley et al. 1990; Pace and Bentley 1992).

Similarly, it has been proposed that the stolonization process in syllids is under hormonal control, following endogenous circadian and circalunar rhythms influenced by exogenous factors, including annual photoperiod, temperature or moon cycles (Franke 1986a; 1999). It has been hypothesized that during the summer time, with long days and high temperatures, a stolonization-promoting hormone produced in the prostomium is secreted to control a second stolonization-suppressing hormone produced in the proventricle (i.e. specialized structure of the digestive tract), allowing the initiation of stolonization (Franke 1999). In contrast, during winter, when days are short and temperatures low at high latitudes, the proventricle is not controlled by the prostomium, and the proventricular stolonization-suppressing hormone then inhibits stolonization (e.g., Abeloos 1950; Durchon 1952, 1959; Durchon and Wissocq 1964; Franke 1980, 1981, 1983a, b, 1985, 1999; Heacox 1980; Heacox and Schroeder 1982; Franke and Pfannenstiel 1984; Verger-Bocquet 1984). Hormonal factors have also been suggested to drive the sexual differentiation of the stolon (Franke 1980; Heacox and Schroeder 1982), in particular the female stolon, given that it seems that male stolon differentiation occurs autonomously, whereas female stolon differentiation may depend on hormone release by male stolons (Franke 1999). However, no candidate hormone has been proposed to control reproduction and regeneration processes in syllids, although it seems clear that there might be several involved, not only in the brain, but also in the proventricle (e.g., Schroeder and Hermans 1975; Franke 1999; Weidhase et al. 2016)

In summary, although molecular mechanisms underlying reproduction are relatively well studied in a few annelids (e.g., Kang et al. 2002; Thamm and Seaver 2008; Giani et al. 2011; Novo et al. 2013; Schenk et al. 2016), the molecular toolkit involved in the stolonization process of syllids has not been examined yet. Thus, our aim in the present study is to provide a first glimpse into the gene expression patterns occurring during the stolonization process in the syllid species *Syllis magdalena*. To achieve this goal, we have pursued four main objectives: 1) to characterize molecularly and morphologically the stolonization process in the target species; 2) to provide a detailed description of the genes potentially involved in the triggering of stolonization

and the formation/releasing of stolons and gametes, through differential gene expression analyses of reproductive and non-reproductive individuals in different tissues; 3) to understand the evolution of selected candidate genes with major roles in the reproductive processes of the phylum Annelida; and 4) to investigate if the molecular signal that determines when to divert resources from somatic functions to reproduction is the same across annelids (i.e., synthesis of methyl farnesoate).

Results and Discussion

General morphology and ultrastructure of the stolons in Syllis magdalena

The stolons of *S. magdalena* were dicerous, with two pairs of red eyes and one pair of antennae formed at the beginning of the stolonization process (Fig. 2A–E, 3A–B), similar to the process observed in *Syllis amica* (see Wissoq 1970) but different to the late formation of head structures in *Syllis gracilis* (see Pettibone 1963) or *Syllis hyalina* (see Malaquin 1893). Natatorial capillary chaetae were not developed during the stages in which the stolon was attached to the stock. Before stolon detachment, the stock completely regenerates the final part of the body that was transformed during the stolon formation (Fig. 2F). Female stolons were purple, completely full of oocytes arranged around the through-gut (Fig. 2A, C, E, 3A–B). Male stolons were white, completely full of spermatogonia, and also arranged around the gut (Fig. 2B, D).

The epithelium of the female and male stolons was columnar, comprised by large epithelial cells (>10 μm in maximum length) with basal non-nucleolated nuclei, and large globular glandular cells with electrondense material (Fig. 3C). In both stolons, below the epithelia, there was a thick layer of muscle fibers, then the germinative epithelium, and finally the digestive epithelium (Fig. 3C–F). The muscle fibers of both female and male stolons presented the regular morphology of muscle fibers of the adults, with a double striation and 25–35 myofilaments and clusters of mitochondria near the tips (Fig. 3C, E). We did not observe the "stolonal" muscle fibers described in *Syllis amica* with the mitochondria towards the middle of the fiber (Wissoq 1967) while attached to the stock. It is possible that the reorganization of the muscle fibers takes place later in the stolonization process, but it is improbable, given that it occurs during head formation in the stolon of *S. amica* (see Wissoq 1967), a process that we observed in *S. magdalena*.

In the female germinative epithelium, large yolky oocytes (50 μm approximately) were surrounded by non-nucleolated nurse cells (Fig. 3D). Oocytes were connected by microvillar processes (Fig. 3D). The male germinative epithelium only contained two large sacs of spermatogonia in the specimens collected (Fig. 3E–F). Spermatogonia (ca. 1 μm in diameter) were densely packed and possess a non-nucleolated nucleus with chromatin condensation processes (Fig. 3E–F). The digestive epithelium was comprised of large (>10 μm in maximum length) convoluted multiflagellated cells (Fig. 3F). We did not observe digestive material in the

lumen of the stolon gut (Fig. 3F). There were no differences in the developmental stage of gametes between the anterior and posterior parts of stolons (see also differential expression results).

General characterization of the de novo transcriptomes

A total 30 libraries derived from somatic and reproductive tissues of non-reproductive (NON-REPRO) and reproductive (REPRO) specimens, were used to assemble de *novo* two transcriptomes, REFSOM and REFTOTREPRO (further details in Material and Methods section). Three somatic parts were chosen for RNA extraction were chosen from all specimens: anterior part (prostomium + first 2 segments), proventricle (all segments containing the proventricle) and final part (pygidium + 2 final segments). Two additional fragments from specimens engaged in stolonization (REPRO) were sequenced: the anterior and posterior half parts of the stolons. Assembly statistics of the reference transcriptomes are summarized in Supplementary File S1, alongside read mapping results for each tissue and specimen. The coverage of our assemblies is similar or slightly higher than those in other studies on marine invertebrates (e.g., Meyer et al. 2009; Riesgo et al. 2012; Pérez-Portela et al. 2016).

An overview of the assigned GO terms for each transcriptome (including 3 different levels: cellular component, CC, biological process, BP, and molecular function, MF) and GO enrichment analyses using Fisher's tests are shown in Supplementary File S2A. The GO enrichment results for the comparisons of both transcriptomes showed 36 GO terms overrepresented in REFSOM related to cellular organization and regulation, metabolism and binding, among others (Supplementary Figure S2B). In contrast, only 8 categories appeared enriched in REFTOTREPRO, mainly related to signaling activity (Supplementary Figure S2C). Interestingly, one of these enriched categories is the activity of G-protein coupled receptors, which bind light-sensitive compounds, pheromones, hormones, neurotransmitters and other ligands involved in secretory processes or cell development, among other functions (e.g., Li et al. 1999; Iversen et al. 2002; Hauser et al. 2006; Asahara et al. 2013). The results of several of these G-protein coupled receptor expression levels on the different tissues and conditions analyzed are discussed below.

Differential gene expression analyses

Pairwise comparisons of somatic tissues (anterior part, proventricle, final segments) between REPRO and NON-REPRO

We detected 792 differentially expressed genes in the comparison between REPRO and NON-REPRO somatic tissues, 494 of them being upregulated in REPRO (178 in females and 316 in males) and 298 in NON-REPRO (Fig. 3 and Supplementary Files S3 and S4). Of these 792 genes, only 292 (~37%) had a BLAST hit and, therefore only the putative annotations for those genes (Supplementary File S4) are discussed below. Among the pairwise comparisons of REPRO

and NON-REPRO tissues, the final segments tissue are the ones that showed more differentially expressed genes, was the final segments (Fig. 4C), with 223 differentially expressed in the comparison of female final segments and NON-REPRO final segments (152 upregulated in female) and 460 differentially expressed genes in the comparison of male final segments and NON-REPRO final segments (304 of those upregulated in male). The pairwise comparisons of anterior part and proventricle between reproductive and non-reproductive individuals showed low numbers of differentially expressed genes (Fig. 4A–B). Among those, the largest differences were in the proventricle with 7 differentially expressed genes upregulated in both females and males when compared to non-reproductive, and 20 and 36 differentially expressed genes upregulated in the proventricle of non-reproductive individuals (Fig. 4B).

In the anterior part and the proventricle of females, the genes upregulated (Supplementary File S4) were related mostly to immune processes (*complement receptor 2*) or food processing (*trefoil factor 2*, *cubilin*, *serine protease 27* and *chitinase*). Similarly, in the male anterior part and proventricle (Supplementary File S4), most genes were involved in nutrient transport (*sugar transporter STL1* and *glycogen phororylase*), as well as development of the nervous system (*tyrosine-protein kinase Src42A*).

Several genes related to gametogenesis were found differentially expressed in the final segments of female and male REPRO individuals compared to NONREPRO (Supplementary File S4), including *vitellogenin* (*Vtg*) and *ovochymase* (*OVCH*) in females, and *testis-specific serine/threonine-kinase* (*TSSK*) in males, which indicates an important role of the final segments during the gametogenesis process in both stolonizing females and males. *Vitellogenin* has been already reported to be involved in annelid gametogenesis, specifically as a yolk precursor (e.g., Hafer et al. 1992), but *OVCH*, an ovary-specific gene involved in egg development of several animals (e.g., Lindsay and Hedrick 1995; Gao and Zhang 2009; Mino and Sawada 2016), is here reported for the first time in annelids. The same occurs for *TSSK*, whose expression, confined almost exclusively to testes, has largely been studied in several mammals (Hao et al. 2004), but never in annelids. Remarkably, two hormone receptors for *relaxin* and *follistatin* were found differentially expressed in the final segments of reproductive females (Supplementary File S4). The insulin-related peptide *relaxin* is important for the growth and remodeling of reproductive tissues during mammal pregnancy (e.g., Gunnersen et al.1995; Hsu et al. 2002) and is active in the ovary and during embryogenesis of zebrafish (e.g., Donizetti et al. 2008, 2010; Wilson et al. 2009). *Relaxin* activity has also been reported in invertebrates, including in the tunicate *Ciona intestinalis* (e.g., Ivell et al. 2005; Olinski et al. 2006), and in the starfish *Asterina pectinifera* (Mita 2013; Mita et al. 2014), where it takes part in oocyte release from the ovary, but this is the first time that it is described in annelids. Likewise, *follistatin*, reported as a follicle-stimulating hormone, with several additional regulatory functions both in reproductive and non-reproductive tissues (Phillips and Kretser 1998), has been already found in the transcriptome of other annelids

such as *C. teleta* and *S. lamarckii* (Kenny et al. 2014), but without a particular association with any biological process. In our case, it seems that both *relaxin* and *follistatin* are important during oocyte development in *S. magdalena,* as they are expressed in tissues where oogenesis is taking place before oocytes are transferred into the stolon (see also results section below).

Pairwise comparisons of somatic (anterior part, proventricle, final segments) between REPRO females and males

We detected 234 genes with significant expression in the comparison between female and male somatic tissues, 85 of them being upregulated in female (0 in anterior part, 27 in proventricle, 58 in final segments) and 149 in males (only in final segments) (see details in Fig. 5A, B and Supplementary File S5). Of these 234 genes, only 84 (~35%) of transcripts were annotated (Supplementary File S5). No differential expression was found in the comparisons of the female and male anterior parts, and in the proventricle comparisons, we only found differentially expressed genes in the females (Fig. 5A and Supplementary File S5; see discussion below). Similar to the previous comparisons (see above), the somatic tissue sample that showed more differentially expressed genes was the final segments, with 149 genes upregulated in males and 58 in females (Fig. 5B and Supplementary File S5).

As in the previous comparisons (see section above), several gametogenesis-related genes, such as *vitellogenin*, *ovochymase* (*OVOCH*) in females, and *TSSK* in males, were differentially expressed in F (Fig. 5B and Supplementary File S5). In addition, we also found *NOTCH* differentially expressed in F of REPRO males (Fig. 5B and Supplementary File S5). This gene has been reported to have a role in segment formation and adult regeneration in annelids (e.g., Tham and Seaver 2008), and therefore may also be involved in segment formation of stolons and pygidium regeneration of *S. magdalena* (Fig. 2F). However, the *NOTCH* pathway has been also reported to be essential for the correct development of gametes in *D. melanogaster* and mammals (Xu et al. 1992; Hayashi et al. 2001; Murta et al. 2014), and therefore it could also be playing such role during spermatogenesis in *Syllis magdalena*.

Two different transcripts of *ovochymase* were differentially expressed in final segments (*OVOCH1*) and proventricle (*OVOCH2*) female tissues (Fig. 5A and Supplementary File S5). *Ovochymases* are involved in the oogenesis in other invertebrates, where they help avoid self-fertilization and are localized in the vitelline coat of oocytes (Mino and Sawada 2016). In the ascidian *Halocynthia roretzi*, *ovochymase* has a signal peptide, three trypsin-like serine protease domains and six CUB domains (Mino and Sawada 2016). We found 3 *ovochymases* (two DE, *OVOCH1* and *OVOCH2*, and one non-DE, *OVOCH3*) in *S. magdalena*, none of them containing a signal peptide and all containing much less trypsin-like serine protease and CUB domains (Supplementary File S6). The trypsin-like serine protease domain is not exclusive to *ovochymases*, since it also occurs in chymotrypsins (Supplementary File S6), which are digestive

enzymes. Given the digestive function of the proventricle in syllids, *OVOCH1* and *OVOCH2* may be performing different functions in *S. magdalena* F and P tissues, respectively. Our molecular phylogeny of ovochymases and chymotrypsins in animals confirmed that *OVOCH1* and *OVOCH3* are homologues of other animal *ovochymases*, whereas *OVOCH2* (the one differentially expressed in the proventricle) is, in fact, a homologue of a mollusk chymotrypsin (Supplementary File S6). *OVOCH1* in *S. magdalena* could be assisting in the maturation of the oocyte, creating an envelope that could further prevent self-fertilization during gamete release in the water column.

Pairwise comparisons of stolons between REPRO females and males.

We detected 1150 differentially expressed genes in the comparison between reproductive tissues of female and male individuals, 872 upregulated in female stolons and 278 in male stolons (Fig. 5C and Supplementary File S5). This comparison showed the largest differences, with ~75% of genes upregulated in females (872) and ~25% in males (278) (Fig. 5C and Supplementary File S5). In addition, we also compared the anterior and posterior halves of stolons, finding only 7 genes upregulated in the anterior half part (Fig. 5C and Supplementary File S5), most of them related to eye (*rhabdomeric opsin*, *retinal-binding protein*) or brain (*TRPC channel protein*) functioning.

Among the most upregulated Biological Process categories in female stolons, we found Nicotinamide metabolism (Fig. 6). Cells need to accommodate the bioenergetic demands During oogenesis, nicotinate and nicotinamide are essential for organisms as the precursors for generation of the coenzymes NAD+ and NADP+, which are fundamental in redox reactions and carry electrons from one reaction to another, being the pillars of many metabolic pathways. The gene *nicotinamide mononucleotide adenylyltransferase 1-like*, which catalyzes the formation of NAD+, was upregulated in the female stolon when compared to the male stolon (Supplementary File S5). Other metabolic pathways upregulated in the female stolons include both fructose and carbohydrate metabolism, illustrating the high energetic requirements of oogenesis (Fig. 6). In male stolons, the major upregulated process related to the high energetic demands of spermatogenesis is Purine metabolism, a pathway required for nucleotide biosynthesis (Fig. 6). Interestingly, the MAPK cascade (included in the category 'Styrene catabolism'), which is central to cell proliferation, is upregulated in female stolons (Fig. 6). Similarly, the gene *alpha-1D adrenergic receptor-like*, which also regulates cell proliferation is upregulated in female stolons.

As in the case of final segments (see section above), *Vtg* and *OVOCH* in females, and *TSSK* and *NOTCH* in males, were also differentially expressed in stolons of females and males (Fig. 5C and Supplementary File S5). These results indicate an important role of the stolons in the maturation of gametes, in contrast to what has been traditionally suggested, where the stolons are thought to be only a place to keep and later spread the gametes. However, no genes related to

gamete maturation were found to show differentially expressed in the comparison between the anterior and posterior halves of stolon, which suggest that there is no sequential anteroposterior maturation of gametes within the stolons (Fig. 5D and Supplementary File S5), in agreement with our results from the morphological and ultrastructural study.

Relaxin was also found differentially expressed in female stolons, reinforcing the hypothesis of its implication in annelid oogenesis and its potential role in the release of oocytes into the water column, as it has been suggested for *relaxin* in *A. pectinifera* (Mita et al. 2014). Other genes involved in gametogenesis of annelids (e.g., Rebscher et al. 2007; Dill and Seaver 2008; Novo et al. 2013) were also found differentially expressed in female stolons (Supplementary File S5), including the member of the DEAD-box helicase protein family, *vasa*. We found two paralogues of the gene *vasa* (the DE *vasa1* and the non-DE *vasa2*) among our transcripts, in contrast to what is found in other annelids that only present one (see Supplementary File S7). While *vasa 2* grouped with all *vasa* orthologs obtained in annelids, *vasa 1* branched out from the annelids and appeared basal to other vasa orthologs from metazoans (Supplementary File S7), being more similar to *ATP-dependent RNA helicase vasa-like* proteins in arthropods than to *vasa* proteins of annelids when blasted. These results may suggest that different paralogues may be performing different functions in *S. magdalena* (Supplementary File S7). While *vasa2* could be playing a role in the female germline determination localized in the oocytes of *S. magdalena*, *vasa1* could be participating in the maintenance of totipotency of the stem cells (Juliano and Wessel 2010), although *ATP-dependent RNA helicase vasa-like* proteins are also known to be involved in oogenesis. Interestingly, we also found the category Steroid biosynthesis upregulated in female stolons (Fig. 6). In addition, our study shows the upregulation of the gene *hydroxysteroid dehydrogenase 2 isoform X2*, that could potentially mediate steroid hormone metabolism (Seckel and Walker 2001), and suggests hormonal control over the final stages of stolonization in *S. magdalena*.

In male stolons, most of the upregulated genes were involved in the construction of the flagellar apparatus (Inaba 2011), including *dyneins*, *cilia- and the flagella-associated proteins*, *ropporin*, *radial spoke 3*, and *kinesins*). This is unsurprising, given the presence of sperm in these tissues, but is an excellent positive control.

Hormonal control of stolonization

Since methyl farnesoate (MF) was discovered to be produced by mandibular organs of numerous crustaceans, this form of the insect Juvenile Hormone (JH III), has been commonly considered as the crustacean equivalent of insect JH (Laufer and Biggers 2001; Miyakawa et al. 2014). Comparably to JH in insects, MF regulates many aspects of crustacean physiology, including reproduction (Xie et al. 2016). In this context, MF is more actively synthetized by females during vitellogenesis, and higher levels of MF are associated with large reproductive systems and

aggressive mating behavior in males of the spider crab *Libinia emarginata* (Laufer et al. 1992). In the annelid *C. teleta*, exogenous extracts of MF were found to affect larval metamorphosis and settlement (Laufer and Biggers 2001), and MF has been recently demonstrated to be directly involved in *P. dumerilii* regeneration and female sexual maturation (Schenk et al. 2016). This latter study not only showed that the decrease of MF levels in the brain induces reproduction and suppresses regenerative capacities in *P. dumerilii*, but it also reported an ortholog of the MF receptor of arthropods (*bHLH-PAS-domain-containing transcription factor methoprene-tolerant receptor*, *MTr*) in the eleocytes (coelomic cells that synthesize yolk via production of *Vtg* protein), demonstrating that this hormone is not restricted to arthropods, as it was assumed (Schenk et al. 2016). Since detection of MF is not possible in RNAseq data, in order to assess whether *S. magadalena* could use a similar molecular signal to determine when to divert resources from somatic functions to reproduction, we investigated if *S. magdalena* also possessed an ortholog of *MTr*, identified as the arthropod and lophotrochozoan sesquiterpenoid receptor (e.g., Konopova and Jindra 2007; Miyakawa et al. 2013; Jindra et al. 2015; Schenk et al. 2016). In our *de novo* transcriptomes, we identified two transcripts encoding bHLH-PAS-domain-containing transcription factor that showed strong similarity to *P. dumerilii MTr*. In fact, our molecular phylogeny of *MTr* revealed that the *S. magdalena* ortholog is closely related to *MTr* orthologs of *P. dumerilii* and *C. teleta* (Fig. 7A). In agreement with Schenk et al. (2016), our results also confirmed that annelid *MTr* is clearly an ortholog of insects and crustaceans *MTrs* (Fig. 7A). These findings allow us to suggest that MF may be one of the hormones responsible for syllid stolonization. If the MF is involved in syllid reproduction, we would expect to find differences in the levels of expression of MF receptors (*MTr*) among the stolonizing and non-stolonizing syllid samples (higher in the latter), similar to what has been reported during oocyte maturation and male reproductive behavior in crustaceans and other annelids (e.g., Laufer et al. 1992; Schenk et al. 2016). Surprisingly, higher expression levels (albeit not statistically significant) of *MTr* were found only in anterior and posterior tissues of female, therefore REPRO, individuals (Fig. 7B), but not in the NON-REPRO specimens, as it might be expected (Schenk et al. 2016). Thus, in contrast to what was found in *P.dumerilii* but similar to what has been reported for arthropods, an increase in MF (or a similar putative sesquiterpenoid) is necessary to initiate the reproductive process in stolonizing syllids (Fig. 7B) (Laufer et al. 1992; Gäde et al. 1997; Wyatt 1997; Hansen et al. 2014). The fact that the differences between conditions are not statistically significant can be explained because the NON-REPRO specimens were collected only one week before the beginning of the stolonization process, and therefore they might have already entered the initial stages of reproduction without visible morphological changes. On the other hand, as in the case of *A. marina* (e.g., Pacey and Bentley 1992), it is also possible that a non-identified hormone, sesquiterpenoid or otherwise, is orchestrating the important metamorphic changes that occur during syllid stolon development, similarly to what MF and JHs do in arthropods (e.g., Hui et al.

2010; Maruzzo et al. 2012; Wen et al. 2015). However, the presence of sesquiterpenoids is further suggested by other DE gene results, as discussed further below.

Interestingly, other neurotransmitter receptors were found to be upregulated in the posterior end of NON-REPRO specimens: *dopamine receptor* (*DAr*), belonging to the large family of G-protein coupled receptors, was downregulated in the final segments of females, and *serotonin transporter* (*SERT* or *5-HTT*), which terminates the action of serotonin, was downregulated in the final segments of males (Supplementary File S5 and Fig. 8A). Our molecular phylogeny corroborates that these proteins are orthologs of the *C. teleta DAr type 2* (*DAr2*, Fig. 8B) and *C. teleta* and *Helobdella robusta SERT* genes (Fig. 8B). Dopamine (DA) and Serotonin (SER) are biogenic amines that act as a neurotransmitters and hormones, regulating an array of important physiological functions both in vertebrates and invertebrates (e.g., Winberg et al. 1997; Neckameyer, 1998a; Gingrich et al. 2000; Wicker-Thomas and Hamann 2008; Dufour et al. 2010; Giang et al. 2011). In *D. melanogaster* DA and SER control a wide range of behavioral processes such as circadian rhythms, sleep, mating behavior, learning or aggression (e.g., Nichols 2007; Giang et al. 2011) and also stimulate fertility and female receptivity (Neckameyer 1998b; Marican et al. 2004). In *C. elegans*, male mating behavior and egg deposition are also induced by DA and SER (Sulston et al. 1975; Weinshenker et al. 1995; Dempsey et al. 2005). In addition, both hormones have been reported to be involved in larval metamorphosis in cnidarians, molluscs, and echinoderms (Couper and Leise 1996; McCauley 1997; Matsuura et al. 2009). In annelids, dopaminergic and serotonergic systems have been found in several species (Grothe et al. 1987; Dietzel and Gottmann 1988; Schlawny et al. 1991; Spörhase-Eichmann et al. 1998; Krajniak and Klohr 1999; Zaccardi et al. 2004; Lawrence and Soame 2009; Helm et al. 2014; Rimskaya-Korsakova et al. 2016; Verasztó et al. 2017; Bauknecht and Jékely 2017). However, the participation of DA and SER in annelid reproduction has only been demonstrated in a handful of studies. Although it was thought that DA played an important role in sexual differentiation in *Ophryotrocha puerilis* (Grothe et al. 1987; Pfannenstiel and Spiehl 1987; Grothe and Pfannenstiel 1986), it was later demonstrated that the catecholaminergic system of this species was involved in mechano- and/or chemoreception (Schlawny et al. 1991). In contrast, both SER and DA in nereids seem to have a positive effect on oocyte development, the first by directly inducing their maturation and the second by switching off the action of the juvenile hormone (Lawrence and Soame 2009). Similarly, in the decapod *Penaeus merguiensis* SER induces ovarian maturation through MF production (Makkapan et al. 2011). In this sense, increased levels of both hormones, as indicated by the upregulation of their receptors and/or transporters (*DAr* and *SERt*) just before the beginning of stolonization (NON-REPRO individuals), could be the stimulus required to initiate oocyte and sperm development during syllid stolonization, with a decrease in the levels afterwards during the course of gametogenesis. In addition to this suggested putative direct role in gametogenesis *per se,* DA could also be the putative hormone in the brain and/or proventricle

inducing the production of MF (or other sesquiterpenoid) to regulate stolonization in *S. magdalena*, as found for DA and the JH of nereids and decapods (Lawrence and Soame 2009; Makkapan et al. 2011). Our results thus indicate a possible role of several hormonal factors in the sexual differentiation of stolons, in agreement with previous studies (Franke 1980; Heacox and Schroeder 1982).

In addition, if DA and SER were the neurohormones regulating stolonization in syllids, our results do not support the traditional view in which male stolons differentiate autonomously and female stolons differentiate upon hormone release by the male stolon (Franke 1999). We found upregulation of the receptors of these two neurohormones in both female and male individuals at the beginning of stolonization. DA and SER have been reported to be under the influence of photoperiodic and circadian rhythms, which are essential for synchronizing several processes in animals (Andretic and Hirsh 2000; Doyle et al. 2002; Lawrence and Soame 2009). Therefore, we suggest that that both female and male stolon differentiation are triggered by environmental cues regulating the production of DA and SER. As in other annelids, the main external signals that may be controlling the synchronicity of the reproductive period in syllids are light and seawater temperature (e.g., Franke 1986b). In the Adriatic Sea, the breeding season of *Syllis prolifera* is restricted from late March to early October, when the temperature ranges from 14ºC–19ºC, and there are around 12–13h of light per day (Franke 1986b). Similar results were observed in *S. magdalena*, which seems to breed during the southern hemisphere summer (see sampling methods) with a mean seawater temperature around 15ºC and around 13h hours of light per day.

In addition to steroid hormone control, we found some differentially expressed genes in the female stolons, potentially involved in the production of pheromones (specifically the sesquiterpenoid MF; see section above): *Farnesyl pyrophosphate synthase* (*FPPS*) and several *methyl transferases* (*MTases*) (Fig. 7B, C and Supplementary File S5), which could synthetize sesquiterpenoids similar to MF and JHIII in arthropods (e.g., Tobe and Bendena 1999; Hui et al. 2010). Specifically, *FPPS* is required at the beginning of the process to catalyze the reaction, generating Farnesyl Diphosphate, the raw material for sesquiterpenoid production, which is then transformed into Farnesol, then Farnesal (via the *Farnesol oxidase/dehydrogenase*, *SDR11*), later into FA (through *Farnesal dehydrogenases*, *ALDHE3*), and, in the canonical pathway, finally into MF in crustaceans (through *Farnesoic acid methyl transferase*, FAMeT), or into JH in insects (through an *epoxidase*, *FAMeT* and *Juvenile hormone acid O-methyltransferase*, *JHAMT*) (e.g., Hui et al. 2010) (Fig. 7D).

Following Schenk et al. (2016) and given our results (including those for *methoprene tolerant receptor*, above), a similar pathway seems to occur in annelids, with the synthesis of some form of sesquiterpenoid regulating reproduction, as occurs in arthropods (Xie et al. 2016). In fact, our phylogenetic results confirmed that the differentially expressed transcripts annotated

as *FPPS* and of a variety *MTases* (Figs. 7C and Supplementary File S8) are legit orthologs, and thus the beginning and end of the synthesis cascade, and the likely bottleneck, are differentially expressed. In addition, orthologs of *FPP*, *SDR11* and *ALDHE3* of spiralians were cleary found in our samples (Supplementary File S9), although these are not differentially expressed themselves. These differentially expressed *MTases* are of a variety of annotations, with some possessing homologs across the Bilateria. None possess clear homology to known arthropod *FAMeT* or *JHAMT* sequences. However, all could potentially be performing a similar role *in vivo*, and one apparent Spiralia novelty is present, which we posit as an excellent candidate for future functional investigation.

However, despite this persuasive circumstancial evidence, we still cannot confirm that the final product of this biosynthetic pathway in *S. magdalena* is MF or another sesquiterpenoid, until functional analyses are performed to test this hypothesis. Besides the putative involvement of sesquiterpenoids in the beginning of syllid stolonization, which is reinforced by the high expression of *SDR11* and *ALDH3* in somatic tissues of both male and female individuals (Fig. 7B), it seems that in our case it may also affect later stages, since *FPPS* and *MTases* are differentially expressed in female stolons (Supplementary File S5). Thus, the increase of MF levels could also be regulating the *vitellogenin* levels necessary for yolk formation, as it commonly occurs with JH in arthropods (Laufer et al. 1992; Gäde et al. 1997; Wyatt 1997; Hansen et al. 2014). In fact, the overexpression of this hormone in stolons could be the triggering signal for the stolon release from the stock. We did not find any enzyme necessary to synthetize hormones or neuropeptides differentially expressed in the male stolons, which might indicate that the synchronicity in the release of female and male stolons might be directly controlled by the female via the production of MF, as it has been also reported during spawning in *A. marina* (Hardege and Benteley 1997).

In addition, as discussed above, MF production has been shown to be influenced by external stimuli (e.g., Shin et al. 2012; Girish et al. 2015; Toyota et al. 2015), which could trigger the stolonization process simultaneously in syllid species according to the traditional hypothesis (e.g., Franke 1999). One of these external stimulus is ambient light variation, which is detected via photosensitive pigments such as opsin proteins and represents a common mechanism mediating the synchronization of gamete release or spawning in a variety of marine invertebrates (Kaniewska et al. 2015; Siebert and Juliano 2017). We have identified several *opsin* homologs in *S. magdalena*, including a *rhabdomeric opsin* previously characterized in other annelids (e.g., Arendt et al. 2004; Randel et al. 2013; Gühmann et al. 2015), that was found differentially expressed in the anterior part of stolons (Supplementary File S5), but not in the anterior part of the stock. Our molecular phylogeny including all *opsins* found in *S. magdalena* (Fig. 9A) revealed that the differentially expressed *rhabdomeric opsin* (*r-opsin 5*) and two other non-differentially expressed *opsins* (*r-opsin 3* and *4*) are homologues of the *P. dumerilii opsin* found in larval eyes

(Arendt et al. 2002). Differences on expression levels among tissues and conditions were observed in the different opsins found in our samples (Fig. 9B), which suggest several roles of opsins at different stages of syllids development, as it has been already stablished in other marine annelids (e.g. Arendt et al. 2004). Specifically, the upregulation of *r-opsin 5* in the anterior part of the stolons, where the stolon eyes are located (Fig. 2A–B, 3A–B) suggests that this *opsin* copy in particular might be responsible for detecting the light changes that would trigger MF production, and the subsequent synchronous stolon release and spawning in *S. magdalena*. A similar mechanism has been recently demonstrated in the hydrozoan jellyfish *Clytia hemisphaerica*, in which spawning is mediated by oocyte maturation-inducing neuropeptide hormones (MIHs), whose release is triggered as a response to blue-cyan light detected by a gonad photosensory *opsin* (Artigas et al 2018).

Conclusions

Using Illumina RNA-seq data, we provide the first transcriptomic characterization of the reproductive process in a species of the family Syllidae. Here, we performed a series of pairwise comparisons of the gene expression patterns in different tissues and conditions that allowed us to identify the molecular mechanisms underlying the stolonization process of *Syllis magdalena*. We found an array of differentially expressed genes involved in immune response, neuronal development, gametogenesis, cell proliferation, and steroid metabolism playing different roles in the reproductive process of *S. magdalena*. Among the most striking results of our study was the continuous gamete maturation occurring in both the final segments and the stolons and the hormonal regulation of the reproduction. Thus, following previous hypotheses proposed for other annelids, including syllids (e.g., Franke and Pfannenstiel 1984; Pacey and Bentley 1992; Franke 1999; Lawrence and Soame 2009; Schenk et al. 2016), we suggest a multihormonal model for the control of syllid stolonization, influenced by environmental signals affecting the anterior part (prostomium) and proventricle of the animal, as it was traditionally hypothesized (e.g., Franke 1999), but also influencing the posterior end of the animals (and thus, the gonads) (Fig. 10). When the breeding season approaches, both DA and SER levels increase triggered by photoperiod and circadian rhythms (Andretic and Hirsh 1999; Lawrence and Soame 2009) and they produce a direct influence on the gonads of pre-reproductive individuals (upregulation of DAr/SERt in final segments of NON-REPRO), initiating the gamete production (Fig. 10A, B). The increase of DA and SER could also positively regulate the production of the putative brain and/or proventricle hormones (such as MF or similar), as in several other invertebrates (Couper and Leise 1996; McCauley 1997; Matsuura et al. 2009) regulating the gamete production (and the consequent metamorphosis to produce stolons), as observed in crustaceans and insects (e.g., Shin et al. 2012; Girish et al. 2015; Toyota et al. 2015). At this point, a variety of other hormones and proteins,

such as *Vtg*, *OVCH*, *relaxin*, *follistatin* and *TSSK*, play their role in the correct development of gametes (Fig. 10B) until maturation is completed. During gamete and stolon maturation, high levels of MF may be required for yolk formation (upregulation in female stolon of *Vtg, FPPS and MTases*), and the presence of MF could additionally trigger stolon release from the stock as a response to external stimuli (as indicated by the upregulation of photosensitive *r-opsins*) (Fig. 10C). We also suggest that the synchronicity of the stolon and gamete release may not only be mediated by exogenous factors such as light and water temperature, but also by chemical cues provided by the female stolons, as demonstrated in other annelids (Hardege and Bentley 1997).

Overall, our results illuminate the process of stolonization in syllids, improving our understanding of how some putative hormones and gametogenesis-related genes regulate the reproduction in stolonizing syllids. However, the transcriptomic approach adopted here does not allow us to locate the specific expression of these genes, and further functional studies are needed to provide a more complete overview of the expression patterns and the proper functioning of specific pathways during reproduction in *S. magdalena*. In addition, RNAi experiments to inhibit the expression of G-protein coupled receptors and other hormones and neuropeptides would provide promising routes to understand their role during stolonization in syllids, allowing us to elucidate once and for all how these annelids delegate sex to their stolons.

Material and methods

Sample collection and preservation

Eight individuals of *S. magdalena* were collected in intertidal algal turfs of *Ulva rigida* and *Perumytilus purpuratus* beds, in Las Cruces, Central Chile (S 33º 30' 06", W 71º 37' 55") in January 2014. Four specimens were collected during full moon, two of which were developing female stolons and the other two male stolons (REPRO specimens); the other four specimens were sampled before the full moon and were not engaged in reproduction (NON-REPRO specimens). All samples were immediately fixed in RNA*later* and stored at -80 ºC until RNA extraction. Two additional male and female stolons were preserved complete in 2.5% glutaraldehyde in 0.4M PBS for electron and confocal microscopy.

Confocal and transmission electron microscopy

Whole specimens preserved in 2.5% glutaraldehyde were mounted in slides to obtain images of auto-fluorescent tissues during stolonization with a Nikon Eclipse upright with A1-Si confocal microscope at the Image Analysis Center (IAC) of the Natural History Museum of London. No stain was applied, but images were obtained in DAPI 488, 555 and 647 channels, under gentle laser excitation. For transmission electron microscopy (TEM), specimens fixed in 2.5% glutaraldehyde were later postfixed in 1% osmium tetroxide and rinsed twice in PBS before dehydration with an increasing series of acetone (from 50% to 100%). Samples were further

embedded in epoxy resin, serially sectioned with an ULTRACUT ultramicrotome at 64 nm, post-stained with uranyl acetate and lead citrate, and observed with a JEOL JEM1010 microscope at the Serveis Científico-Tècnics (SCT) at the Universitat de Barcelona and at the Servicio Interdepartamental de Investigación (SIDI) of the Universidad Autónoma de Madrid.

RNA extraction

Each tissue sample was transferred to a microcentrifuge tube containing 500 μL of TRIzol (Invitrogen), and ground with a RNase-free plastic pestle to break down the tissue, and isolate RNA and DNA. Then, another 500 μL of TRIzol and 10 μL of glycogen were added. After 10 min incubating the mixture at room temperature (RT), 100 μL of the RNA-isolating reagent bromochloropropane (BCP) was mixed in by vortexing. After 10 min incubation at RT, samples were centrifuged at 16,000 relative centrifugal force (rcf) units for 15 min at 4 °C to separate the solution into 3 layers. The upper aqueous layer, which contained total RNA, was recovered and mixed with 500 mL of isopropanol, and incubated at –20 ºC overnight. Afterwards, the sample was centrifuged at 16,000 rcf for 15 min at 4 ºC, and the supernatant was removed. Total RNA precipitation was performed by washing the remaining pellet twice by adding 1 mL of 75% ethanol and centrifuging it at 16,000 rcf at 4 °C for 5 min. The dried pellet was eluted in 100 μL of RNA Storage solution (Invitrogen). mRNA purification was performed with a Dynabeads mRNA Purification Kit (Invitrogen), following manufacturer's instructions. After incubation of total RNA at 65 ºC for 5 min, the samples were incubated for 30 min with 200 mL of magnetic beads in a rocker and washed twice with washing buffer.

Thirteen μL of 10 mM Tris-HCl were added to the eluate, and the mixture was incubated at 80 °C for 2 min. The supernatant was immediately transferred to a 0.5 mL microcentrifuge tube and stored at -80 °C. Quality of mRNA was measured with a pico RNA assay in the Agilent 2100 BioAnalyzer (Agilent Technologies). Quantity was measured with an RNA assay in a Qubit fluorometer (Life Technologies). Further details about RNA prep protocols can be found in Fernández et al. (2014).

cDNA Library Construction and Next-Generation Sequencing

cDNA libraries were constructed from extracted mRNA in the Apollo 324 automated system using the PrepX mRNA 8 Protocol Kit (IntegenX) set to 200 base pairs (bp) and stranded mRNA, under the Library Prep Illumina setting. A Polymerase Chain Reaction (PCR) was run to amplify cDNA libraries, using the KAPA Library Amplification Kit. PCR was run as follows: denaturation (45 sec at 98 ºC), cycling (15 sec at 98 ºC, 30 sec 60 ºC, and 15 sec at 72 ºC, for 16 cycles), and final extension (1 min at 72 ºC). During the PCR process, the samples were marked with a different index to allow pooling for sequencing. cDNA library quality and size were measured through a dsDNA High Sensitivity (HS) assay in an Agilent 2100 BioAnalyzer (Agilent

Technologies). A quantitative real-time PCR (qPCR) was run to measure cDNA library concentration using the KAPA Library Quantification Kit. qPCR settings were as follows: Initial denaturation (5 min at 95 °C for 1 cycle), then denaturation (30 sec at 95 °C) and annealing/extension/data acquisition (45 sec at 60 °C) combined for 35 cycles. The libraries were then run on the Illumina HiSeq 2500 sequencing platform, with output of paired-end reads of 150 bp by the FAS Center for Systems Biology at Harvard University.

Sequence processing and de novo assembly

Demultiplexed Illumina HiSeq 2500 sequencing datasets of the 30 tissue samples, in FASTQ format, were retrieved; the quality of the raw reads was assessed and visualized using FASTQC v. 0.11.5 (www.bioinformatics.babraham.ac.uk). Adapter sequences and bases with low-quality phred scores (<30) were trimmed off, and a length filter was applied retaining sequences of >25 bases using TRIMGALORE v. 0.4.2 (www.bioinformatics.babraham.ac.uk).

Two *de novo* transcriptome assemblies for *Syllis magdalena* were constructed with the software Trinity to streamline further differential gene expression analyses (Grabherr et al. 2011; Haas et al. 2013): a reference transcriptome (REFSOM assembly) containing reads from only the somatic parts (anterior part, proventricle, final segments) of each individual of both REPRO and NON-REPRO specimens (23 libraries), and a reference transcriptome including the 5 different parts (anterior part, proventricle, final segments, anterior half part of stolon and posterior half of stolon) of each individual (13 libraries) for only the reproductive specimens (REFTOTREPRO assembly). We did not obtain enough RNA from two of the female tissue samples, proventricle of specimen 0 and anterior part of stolon of specimen 1, to build a library, and therefore conditions 'proventricle' and 'anterior half of stolon' were represented by a single library in females. Given the large number of raw reads obtained in our study (> 500 million reads), we assembled two different reference transcriptomes, since assembling a single reference transcriptome with the available computational resources would have proved computationally impossible. Raw reads have been deposited in the Sequence Read Archive (BioProject ID PRJNA434571; SRA SUB3709568).

Transcriptome characterization: Blast and Annotation

Annotation of transcriptome contigs or transcripts (containing all isoforms) for both *de novo* assemblies were done separately using BLASTX against a selection of non-redundant (nr) database from NCBI containing only proteins from Metazoa, with an expected value (E-value) cutoff of $1e^{-5}$ (Altschul et al.1997). Blast results of the two *de novo* assemblies were used to retrieve Gene Ontology (GO) terms with BLAST2GO 4.0.2 (Conesa *et al.* 2005) under the three different categories: cellular component (CC), biological processes (BP) and molecular function (MF). In addition, GO enrichment analyses using Fisher's test were done in BLAST2GO, to

assess which GO terms were significantly overrepresented in pairwise comparisons between both REFSOM and REFTOTREPRO transcriptomes. The *p*-value for the reciprocal comparisons was adjusted to a 0.05 false discovery rate (FDR) (Benjamini and Hochberg 1995). The Galaxy web-based platform (http://usegalaxy.org) was used to align the RSEM results of each sample with BLASTX results for the *de novo* assemblies for display.

Estimation of expression levels

In order to obtain expression levels, as read counts, of genes (with all isoforms collapsed) for each tissue type of *S. magdalena* specimens in both reproductive and non-reproductive conditions, trimmed paired reads after trimming were mapped against the reference transcriptome, using BOWTIE2 v. 2.2.1 (Langmead and Salzberg 2012), as implemented in Trinity (Grabherr et al. 2011). The software RSEM v. 1.2.11 (Li and Dewey 2011) was used to generate a table containing read counts.

Differential gene expression analyses

Differential gene expression analyses were computed in pairwise comparisons of different tissues and conditions using the R package DESeq2, which allows analyses to be performed with low numbers of replicates (Ander and Huber 2010). Before analyzing differential gene expression, read counts were normalized by estimating a scaling factor for each transcript in DESeq2 (Dillies et al. 2013). The significance value for multiple comparisons was FDR adjusted to 0.01 (Benjamini and Hochberg 1995). Visualization of the significant outcomes of genes differentially expressed (upregulated and downregulated) between the tissues and conditions was obtained with a heatmap performed with the 'GPLOTS' package of R (http://www.r-project.org/). Using the GO annotation results for the 'reference' transcriptome, we obtained the GO terms associated with the differentially expressed isoforms in both pairwise comparisons, which were then implemented together with their *p*-value (adjusted) associated in REVIGO web server (Supek et al. 2011), and graphically represented with the 'TREEMAP' function in R. Size of the rectangles was adjusted to reflect the *p*-value using the abs_log_pvalue option in REVIGO.

Phylogenetic analyses

The evolutionary history of specific genes that could potentially be involved in the stolonization process was also assessed through phylogenetic inference. The translated amino acid sequences of these genes were aligned with orthologues of the same genes in other metazoans obtained from GenBank using MUSCLE ver. 3.6 (Edgar 2004). The G-protein coupled receptors *DAr2* and *SERT* were analyzed together. Both *vasa* and *PL10* are *DEAD-box helicases* and were analyzed together. Other genes were examined in their individual gene families. We selected the

best-fit model of amino acid substitution (LG + Γ + G, WAG, as indicated in Figure legends) with ProtTest ver. 2.4 (Abascal et al. 2005) under the Akaike Information Criterion (Posada and Buckley 2004) and later fed into the software for phylogenetic reconstruction. Maximum likelihood analyses of all the genes were conducted in RAxML ver. 7.2.7 (Stamatakis 2006) with 500 independent searches, and 1000 bootstrap replicates (Stamatakis et al. 2008).

Acknowledgments

The authors are indebted to many members of the Giribet Lab at Harvard University for their help during sample processing, specially to Dr. Sarah Lemer and Dr. David Combosch (currently at University of Guam). Special thanks go to Dr. Greg Rouse (Scripps Institution of Oceanography, UCSan Diego) and Dr. Carlos Sentís (UAM) who provided comments and advice at the beginning of the research. The first author is also very grateful to Dr. Raquel Pérez-Palacios, Dr. Jorge Barbazán (Institut Curie, Paris) and members of Dr. Michel Vervoort lab (Institut Jacques Monod) for their support and useful comments to improve the last version of the manuscript. We also thank also to Milagros Guerra (CBM, CSIC) for his help with TEM observations at UAM. This research received funding from the European Union's (European Atomic Energy Community's) Seventh Framework Program (FP7/ 2007–2013; FP7/2007–2011) under grant agreement 227799 to P.A.-C. Sequencing and analyses were conducted with internal MCZ funds to G.G. and with the support of the Center for Systems Biology and the Research Computing group, both from the Faculty of Arts and Sciences, Harvard University.

Figure legends

Figure 1. Syllinae schizogamous reproductive cycle (stolonization) using light microscope pictures of *Syllis magdalena*.

Figure 2. Ligth microscopy pictures of *Syllis magdalena* stolonizing female (A) and male (B). Confocal micrographs of *S. magdalena* stolonizing female (C), male (D), female stolon (E) and male stolons (F). Arrows in A–D pointing to the eyes of stolons (e). Arrows in E pointing to antennae (a). Arrow in F pointing to the regeneration of the final segments in the stock (rfs).

Figure 3. Ligth and electron microscopy pictures of the anterior part of the female and male stolons of *Syllis magdalena*. (A-B) Location of antennae (a) and the two pairs of eyes (e) in the female stolon. (C) Trasmission electron micrographs of the epithelium of the female stolon showing the muscle fibers (mf), granular cells (gc) and epithelial cells (ec). (D) Developing oocytes showing nucleolate (nu) nucleus (n), ooplasm filled with yolk platelets, and microvilli (m) contacting close oocytes. Note the muscle fibers (mf) nurse cells (nc) and the digestive epithelium (dc) surrounding the germinal epithelium. (E-F) Germinal epithelium (ge) in the male stolon. The stolonal epithelium is comprised by a layer of epithelial cells (ec) with interspersed granular cells (gc), and a layer of muscle fibers (mf); spermatogonia develop in the germinal

epithelium (gc) below. The digestive cells (dc) lay below the germinal epithelium.

Figure 4. Heatmaps of differentially expressed genes (annotated and not annotated genes) from pairwise comparisons of somatic tissues between reproductive (both female and male) and non-reproductive individuals. Anterior part tissue comparisons (A), proventricle comparisons (B) and final segments comparisons (C). Different colors indicate relative expression levels (see color key and histogram on each). Similarity in expression patterns between genes and individuals is represented by clustering. Abbreviations: anterior part (A), proventricle (P), final segments (F).

Figure 5. Heatmaps based on differentially expressed genes (annotated and not annotated genes) from pairwise comparisons of somatic tissues between females and males (A, B) and reproductive tissues (stolons) (C, D). Proventricle comparisons (A), final segments comparisons (B), female and male stolons comparisons (C) and anterior and posterior parts of stolons (female and male together) (D). Different colors indicate relative expression levels (see color key and histogram on each). Similarity in expression patterns between genes and individuals is represented by clustering. Abbreviations: anterior part (A), proventricle (P), final segments (F), anterior half of stolon (AS), posterior part of stolon (FS).

Figure 6. Gene Ontology treemaps for annotated differentially expressed genes in female stolons versus male stolons. The GO terms downregulated in female stolons, are upregulated accordingly in male stolons.

Figure 7. (A) Phylogenetic reconstruction of the protein alignment for methoprene-tolerant receptor (MTr) found in our samples. (B) Heatmap showing the relative levels of expression in the different tissues and conditions analyzed of the transcripts that may be involved in the synthesis of the neurohormone *methyl farnesoate* (MF): MTr, *Farnesol oxidase/dehydrogenase* (SDR11), *Farnesal dehydrogenases* (ALDHE3), the differentially expressed transcript *Farnesyl pyrophosphate synthase* (FPP) and putative *methyl transferase* (Mtase). (C) Phylogenetic reconstruction of the differentially expressed *MTases* in the female stolon. (D) Synthesis pathway of MF and JH in arthropods. Abbreviations: anterior part (A), proventricle (P), final segments (F), anterior half of stolon (AS), posterior part of stolon (FS).

Figure 8. Phylogenetic reconstruction (A) and heatmap of relative levels of expression in all the tissues and conditions (B) of the genes *dopamine receptor* (*DAr*) and *serotonin transporter* (*SERT*). Abbreviations: anterior part (A), proventricle (P), final segments (F), anterior half of stolon (AS), posterior part of stolon (FS).

Figure 9. Phylogenetic reconstruction of the protein alignment for the different *opsin* genes (*rhabdomeric* and *ciliary*) found in our samples (A) and levels of expression of all of them in the different tissues and conditions analized (B). *Rhabdomeric opsin 5* appeared differentially expressed in the anterior part of stolons. Abbreviations: anterior part (A), proventricle (P), final segments (F), anterior half of stolon (AS), posterior part of stolon (FS).

Figure 10. Multihormonal model proposed for the control of syllid stolonization.

Supplementary Material

Supplementary File S1. General statistics of the 'reference' de novo transcriptomes REFSOM and REFTOTOREP. Total number of trimmed reads used for the assemblies (Total Reads), number of aligned reads for each tissue (Aligned Reads), percentage of aligned reads for each tissue, number of aligned transcripts (Align. Transcr.: including genes and isoforms collapsed into genes), number of genes, percentage og genes, parameter N50, number of transcripts with blast hit against proteins of metazoans (Hit_Mtz), number of annotated transcripts from metazoans (Annot), percentage of GC (GC%), median transcript length (MTL), average transcript length (ATL) and number of assembled bases expressed as Mb.

Supplementary File S2. (A) Assigned Gene Ontology (GO) terms for each transcriptome including 3 different levels: CC, BP and MF. (B) GO-term enrichment analysis based on the Fisher's tests results of pairwise comparisons between both transcriptomes, showing the 36 processes overrepresented in REFSOM, and (C) the 8 processes overrepresented in REFTOTREPRO.

Supplementary File S3. (A) Heatmap and (B) correlation matrix of differentially expressed genes from all the tissues and conditions analyzed.

Supplementary File S4. Differentially expressed genes with blast hit, and further putative annotations (GO terms), from the pairwise comparisons of somatic tissues (anterior part, proventricle, final segments) between reproductive and non-reproductive individuals.

Supplementary File S5. Differentially expressed genes with blast hit, and putative annotations (GO terms), from the pairwise comparisons of somatic tissues (anterior part, proventricle, final segments) and reproductive tissues (stolons) between female and male individuals.

Supplementary File S6. Phylogenetic reconstruction of the protein alignment for all the paralogs of the genes *ovochymase* and *chymotripsin* found in our samples. *Ovochymase 1* was differentially expressed in the final segments of reproductive females, and *ovochymase 2* (clearly a *chymotrypsin* in this hypothesis) was differentially expressed in P of reproductive females.

Supplementary File S7. Phylogenetic reconstruction of the protein alignment for the DEAD-box helicases *vasa* and *PL10* showing the two homologs of *vasa* (*vasa 1* differentially expressed) and one for *PL10*.

Supplementary File S8. Phylogenetic reconstruction of the protein alignment for the different enzymes involved in the synthesis of methyl farnesoate. *Farnesol oxidase/dehydrogenase* (SDR11) (A), *Farnesyl pyrophosphate synthase* (FPP) (B), Farnesyl phosphatase (C), and *Farnesal dehydrogenases* (ALDHE3) (D).

References

Abascal F, Zardoya R, Posada D. 2005. ProtTest: selection of best-fit models of protein evolution. *Bioinformatics.* 21(9): 2104–2105.

Abeloos M. 1950. Régénération et stolonisation épigame chez l'Annélide *Syllis prolifera* Krohn. *CR Acad Sci.* 230: 1899–1900.

Agassiz A. 1863. On alternate generation in annelids, and the embryology of *Autolytus cornutus*. *Boston Journal of Natural History* 7: 384–409.

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman, D.J. 1997. Gapped Blast and PSIBlast: a new generation of protein database search programs. *Nucleic Acids Res.* 25: 3389–3402.

Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biol.* 11: R106.

Andretic R, Hirsh J. 2000. Circadian modulation of dopamine receptor responsiveness in *Drosophila melanogaster*. *Proc NatL Acad Sci USA*. 97(4): 1873-1878.

Arendt D, Tessmar K, de Campos-Baptista MI, Dorresteijn A, Wittbrodt J. 2002. Development of pigment-cup eyes in the polychaete Platynereis dumerilii and evolutionary conservation of larval eyes in Bilateria. *Development.* 129(5):1143-54.

Arendt D, Tessmar-Raible K, Snyman H, Dorresteijn AW, Wittbrodt J. 2004. Ciliary photoreceptors with a vertebrate-type opsin in an invertebrate brain. *Science* 306(5697): 869–871.

Artigas GQ, Lapébie P, Leclère L, Takeda N, Deguchi R, Jékely G, Momose T Houliston E. 2018. A gonad-expressed opsin mediates light-induced spawning in the jellyfish *Clytia*. *eLife* 7: e29555.

Asahara S, Shibutani Y, Teruyama K, Inoue HY, Kawada Y, Etoh H, Matsuda T, Kimura-Koyanagi M, Hashimoto N, Sakahara M, Fujimoto W. 2013. *Ras-related C3 botulinum toxin substrate 1* (*RAC1*) regulates glucose-stimulated insulin secretion via modulation of F-actin. *Diabetologia* 56(5): 1088–1097.

Bauknecht P, Jékely G. 2017. Ancient coexistence of norepinephrine, tyramine, and octopamine signaling in bilaterians. *BMC Biol.* 15(1): 6.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol.* 57: 289–300.

Bentley MG. 1985. Sperm maturation response in Arenicola marina L.: An in vitro assay for sperm maturation factor and its partial purification. *International journal of invertebrate reproduction and development*. 8(3), pp.139-148.

Bentley MG, Clark S, Pacey AA. 1990. The role of arachidonic acid and eicosatrienoic acids in the activation of spermatozoa in *Arenicola marina* L.(Annelida: Polychaeta). *Biol Bull.* 178(1):1-9.

Conesa A, Götz S, García-Gómez JM, Terol J, Talon M, Robles M. 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21: 3674–3676.

Couper JM, Leise EM. 1996. Serotonin injections induce metamorphosis in larvae of the gastropod mollusc *Ilyanassa obsoleta. Biol Bull.* 191(2), 178-186.

Daly JM. 1975. Reversible epitoky in the life history of the polychaete *Odontosyllis polycera* (Schmarda, 1861). *J Mar Biol Assoc UK*. 55: 327–344.

Dempsey CM, Mackenzie SM, Gargus A, Blanco G, Sze JY. 2005. Serotonin (5HT), fluoxetine, imipramine and dopamine target distinct 5HT receptor signaling to modulate *Caenorhabditis elegans* egg-laying behavior. *Genetics* 169(3): 1425-1436.

Dietzel ID, Gottmann K. 1988 Development of dopamine-containing neurons and dopamine uptake in embryos of *Hirudo medicinalis. Dev Biol.* 128:277–283

Dill KK, Seaver EC. 2008. *Vasa* and *nanos* are coexpressed in somatic and germ line tissue from early embryonic cleavage stages through adulthood in the polychaete *Capitella* sp. I. *Dev Genes Evol.* 218: 453–463.

Dillies MA, Rau A, Aubert J, Hennequet-Antier C, Jeanmougin M, Servant N, Keime C, Marot G, Castel D, Estelle J, Guernec G, 2013. A comprehensive evaluation of normalization methods for Illumina high-throughput RNA sequencing data analysis. *Brief Bioinfot.* 14: 671–683.

Donizetti A, Grossi M, Pariante P, D'aniello E, Izzo G, Minucci S, Aniello F. 2008. Two neuron clusters in the stem of postembryonic zebrafish brain specifically express *relaxin-3* gene: First evidence of nucleus incertus in fish. *Dev Dyn.* 237(12): 3864–3869.

Donizetti A, Fiengo M, del Gaudio R, Di Giaimo R, Minucci S, Aniello F. 2010. Characterization and developmental expression pattern of the *relaxin receptor rxfp1* gene in zebrafish. *Dev Growth Differ.* 52(9): 799–806.

Doyle SE, Grace MS, McIvor WI, Menaker M. 2002. Circadian rhythms of dopamine in mouse retina: the role of melatonin. *Vis Neurosci.* 19(5):593-601.

Dufour S, Sebert ME, Weltzien FA, Rousseau K, Pasqualini C. 2010. Neuroendocrine control by dopamine of teleost reproduction. *J Fish Biol.* 76: 129–160.

Durchon M. 1951. Stolonisation et hermaphroditisme succesif chez *Syllis amica*. *Arch Zool Exp Gen.* 88: 96–100.

Durchon M. 1952. Recherches expérimentales sur deux aspects de la reproduction chez les Annélides Polychètes: L'épitoquie et la stolonisation. *Ann Sci Nat Zool Biol Anim.* 14: 117–206.

Durchon M. 1959. Contribution à l'étude de la stolonisation chez les Syllidiens (Annélides, Polychétes): I. Syllinae. *Bull Biol Fr Bel* 93: 155–219.

Durchon M, Wissocq J-C. 1964. Contribution à l'étude de la stolonisation chez les Syllidiens (Annélides Polychètes): II. Autolytinae. *Ann Sci Nat Zool Biol Anim.* 6: 159–208.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32: 1792–1797.

Fernández R, Laumer CE, Vahtera V, Libro S, Kaluziak S, Sharma PP, Pérez-Porro AR, Edgecombe GD, Giribet G. 2014. Evaluating topological conflict in centipede phylogeny using transcriptomic data sets. *Mol Biol Evol.* 31: 1500–1513.

Franke H-D. 1980. Zur Determination der zeitlichen Verteilung von Fortpflanzungsprozessen in Laborkulturen des Polychaeten *Typosyllis prolifera*. *Helgol Meeresunters* 34: 61–84.

Franke H-D. 1981. Der Einfluß von Temperatur und Tageslänge auf das endokrine System der Fortpflanzungskontrolle bei dem Polychaeten *Typosyllis prolifera*. *Verh Dtsch Zool Ges.* 1981: 190.

Franke H-D. 1983a. Endocrine mechanisms mediating light temperature effects on male reproductive activity in *Typosyllis prolifera* (Polychaeta, Syllidae). *Roux Arch Dev Biol* 192: 95–102.

Franke H-D. 1983b. Endocrine control of reproductive periodicity in male *Typosyllis prolifera* (Polychaeta, Syllidae). *Int J Invertebr Repr.* 6: 229–238.

Franke H-D. 1985. On a clocklike mechanism timing lunar rhythmic reproduction in *Typosyllis prolifera* (Polychaeta). *J Comp Physiol A.* 156: 553–561.

Franke H-D. 1986a. Resetting a circalunar reproduction rhythm with artificial moonlight signals: Phase-response curve and 'moon-off' effect. *J Comp Physiol A.* 159(4): 569–576.

Franke H. D. 1986b. The role of light and endogenous factors in the timing of the reproductive cycle of *Typosyllis prolifera* and some other polychaetes. *Am Zool.* 26(2): 433–445.

Franke, H. D. 1999. Reproduction of the Syllidae (Annelida: Polychaeta). *Hydrobiologia.* 402: 39–55.

Franke H-D, Pfannenstiel H-D. 1984. Some aspects of endocrine control of polychaete reproduction. In Fischer, A. and H.-D. Pfannenstiel, editors. Polychaete Reproduction. Fortschr. Zool. 29: 53–72.

Gäde G, Hoffmann KH, Spring JH. 1997. Hormonal regulation in insects: facts, gaps, and future directions. *Physiol Rev.* 77: 963–1032.

Gao K, Zhang S. 2009. Ovochymase in amphioxus *Branchiostoma belcheri* is an ovary-specific trypsin-like serine protease with an antibacterial activity. *Dev Comp Immunol. 33*(12): 1219–1228.

Garwood PR. 1991. Reproduction and the classification of the family Syllidae (Polychaeta). *Ophelia (Supplement)* 5: 81–87.

Giang T, Rauchfuss S, Ogueta M, Scholz H. 2011. The serotonin transporter expression in Drosophila melanogaster. *J Neurogenet.* 25(1–2): 17-26.

Giani VC, Yamaguchi E, Boyle MJ, Seaver EC. 2011. Somatic and germline expression of *piwi* during development and regeneration in the marine polychaete annelid *Capitella teleta*. *Evodevo* 2: 10.

Gingrich B, Liu Y, Cascio C, Wang Z, Insel TR. 2000. Dopamine D2 receptors in the nucleus accumbens are important for social attachment in female prairie voles (*Microtus ochrogaster*). *Behav Neurosci.* 114(1): 173.

Girish BP, Swetha CH, Reddy PS. 2015.Induction of ecdysteroidogenesis, methyl farnesoate synthesis and expression of ecdysteroid receptor and retinoid X receptor in the hepatopancreas and ovary of the giant mud crab, Scylla serrata by melatonin. *Gen Comparative Endocrinol.* 217:37–42.

Grabherr MG, Haas, BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol.* 29(7): 644-652.

Grothe C, Pfannenstiel HD. 1986. Cytophysiological study of neurosecretory and pheromonal influences on sexual development in *Ophryotrocha puerilis* (Polychaeta, Dorvilleidae). *Int Journal Invertebr Repr Dev*. 10(2): 227-239.

Gunnersen JM, Crawford RJ, Tregear GW. 1995. Expression of the *relaxin* gene in rat tissues. *Mol Cell Endocrinol.* 110(1): 55–64.

Gühmann M, Jia H, Randel N, Verasztó C, Bezares-Calderón LA, Michiels NK, Yokoyama S, Jékely G. 2015. Spectral tuning of phototaxis by a go-opsin in the rhabdomeric eyes of *Platynereis*. *Curr Biol.* 25(17): 2265–2271.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, MacManes MD. 2013. De novo transcript sequence reconstruction from RNA-Seq: reference generation and analysis with Trinity. *Nat Protoc.* 8(8): 1494–1512.

Hafer J, Fischer A, Ferenz HJ. 1992. Identification of the yolk receptor protein in oocytes of *Nereis virens* (Annelida, Polychaeta) and comparison with the locust vitellogenin receptor. *J Comp Physiol B*. 162(2): 148–152.

Hansen IA, Attardo GM, Rodriguez SD, Drake LL. 2014. Four-way regulation of mosquito yolk protein precursor genes by juvenile hormone-, ecdysone-, nutrient-, and insulin-like peptide signaling pathways. *Front Physiol.* 5:103.

Hao Z, Jha KN, Kim YH, Vemuganti S, Westbrook VA, Chertihin O, Markgraf K, Flickinger CJ, Coppola M, Herr JC, Visconti PE. 2004. Expression analysis of the human testis-specific serine/threonine kinase (TSSK) homologues. A TSSK member is present in the equatorial segment of human sperm. *Mol. hum reprod.* 10(6), 433-444.

Hardege JD, Bentley MG. 1997. Spawning synchrony in *Arenicola marina*: evidence for sex pheromonal control. *Proc R Soc Lond B Biol Sci*. 264(1384), 1041-1047.

Hauser F, Williamson M, Cazzamali G, Grimmelikhuijzen CJ. 2006. Identifying neuropeptide and protein hormone receptors in *Drosophila melanogaster* by exploiting genomic data. *Brief Funct Genomic Proteomic.* 4(4): 321–330.

Hayashi T, Kageyama Y, Ishizaka K, Xia G, Kihara K, Oshima H. 2001. Requirement of *Notch 1* and its ligand jagged 2 expressions for spermatogenesis in rat and human testes. *J Androl.* 22(6), 999-1011.

Heacox AE. 1980. Reproduction and development of *Typosyllis pulchra* (Berkeley and Berkeley) (Polychaeta: Syllidae). *Pac Sci.* 34: 245–59.

Heacox AE, Schroeder PC. 1982. The effects of prostomial and proventriculus removal on sex determination and gametogenesis in *Typosyllis pulchra* (Polychaeta: Syllidae). *Roux Arch Dev Biol.* 191: 84–90.

Helm C, Stevenson PA, Rouse GW, Bleidorn C. 2014. Immunohistochemical investigations of *Myzostoma cirriferum* and *Mesomyzostoma* cf. *katoi* (Myzostomida, Annelida) with implications for the evolution of the myzostomid body plan. *Zoomorphology*. 133(3): 257-271.

Hsu SY, Nakabayashi K, Nishi S, Kumagai J, Kudo M, Sherwood OD, Hsueh AJ. 2002. Activation of orphan receptors by the hormone relaxin. *Science 295*(5555): 671–674.

Hui JH, Hayward A, Bendena WG, Takahashi T, Tobe SS. 2010. Evolution and functional divergence of enzymes involved in sesquiterpenoid hormone biosynthesis in crustaceans and insects. *Peptides*, 31(3), 451-455.

Inaba K, 2011. Sperm flagella: comparative and phylogenetic perspectives of protein components. *Mol Hum Reprod*. *17*(8), pp.524-538.

Ivell R. Anand-Ivell R. 2005. Understanding *relaxin* in the female reproductive system. *Curr Med Chem Immunol Endocr Metab Agents*. *5*(5): 383–389.

Iversen A, Cazzamali G, Williamson M, Hauser F, Grimmelikhuijzen CJ. 2002. Molecular cloning and functional expression of a *Drosophila* receptor for the neuropeptides *capa-1 and-2*. *Biochem Biophys Res Commun*. 299(4): 628–633.

Jindra M, Uhlirova M, Charles JP, Smykal V, Hill RJ. 2015. Genetic evidence for function of the bHLH-PAS protein Gce/Met as a juvenile hormone receptor. *PLoS Genetics* 11: e1005394.

Juliano C, Wessel G. 2010. Versatile germline genes. Science 329(5992): 640-641.

Kang D, Pilon M, Weisblat DA. 2002. Maternal and zygotic expression of a *nanos*-class gene in the leech *Helobdella robusta*: primordial germ cells arise from segmental mesoderm. *Dev Biol*. 245: 28–41.

Kaniewska P, Alon S, Karako-Lampert S, Hoegh-Guldberg O, Levy O, Pyott S. 2015 Signaling cascades and the importance of moonlight in coral broadcast mass spawning. *eLife*. 4:e09991.

Konopova B, Jindra M. 2007. Juvenile hormone resistance gene Methoprene-tolerant controls entry into metamorphosis in the beetle *Tribolium castaneum*. *Proc Natl Acad Sci USA* 104:10488–10493.

Krajniak KG, Klohr RW. 1999. The effects of FMRFamide, serotonin, and acetylcholine on the isolated crop-gizzard of the earthworm, *Lumbricus terrestris*. *Comp Biochem Physiol A Molecul Integr Physiol*. 123(4): 409–415.

Langmead B, Salzberg S. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 9: 357–359.

Laufer H, Sagi A, Ahl J, Homola E. 1992. Methyl farnesoate appears to be a crustacean reproductive hormone. *Invertebr. Reprod. Dev*. 22: 17–19

Laufer H, Biggers WJ. 2001. Unifying concepts learned from methyl farnesoate for invertebrate reproduction and post-embryonic development. *American Zoologist* 41:442–457.

Li B, Dewey C. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 12: 323.

Li C, Kim K, Nelson LS. 1999. FMRFamide-related neuropeptide gene family in *Caenorhabditis elegans*. *Brain research* 848(1): 26–34.

Lindsay LL, Hedrick JL. 1995. Isolation and characterization of ovochymase, a chymotrypsinlike protease released during *Xenopus laevis* egg activation. *Developmental Biology* 167: 513–6.

Makkapan W, Maikaeo L, Miyazaki T, Chotigeat W. 2011. Molecular mechanism of serotonin via methyl farnesoate in ovarian development of white shrimp: *Fenneropenaeus merguiensis* de Man. *Aquaculture* 321(1): 101-107.

Malaquin A. 1893. Recherches sur les Syllidiens, morphologie, anatomie, reproduction, développement. *Mem Soc Sci Agricul Arts Lille.* 18: 1–477.

Marican C, Duportets L, Birman S, Jallon JM. 2004. Female-specific regulation of cuticular hydrocarbon biosynthesis by dopamine in *Drosophila melanogaster*. *Insect Biochem Mol Biol*. 34(8): 823-830.

Marion AF, Bobretsky NV. 1875. Etude des Annelides du golfe de Marseille. *Ann Sci Nat Zool Paleontol.* 2: 2–46.

Maruzzo D, Aldred N, Clare AS, Høeg, JT. 2012. Metamorphosis in the cirripede crustacean *Balanus amphitrite*. *PLoS One*: 7(5), e37408.

Matsuura H, Yazaki I, Okino, T, 2009. Induction of larval metamorphosis in the sea cucumber *Apostichopus japonicus* by neurotransmitters. *Fish Sci.* 75(3): 777-783.

McCauley DW. 1997. Serotonin Plays an Early Role in the Metamorphosis of the Hydrozoan *Phialidium gregarium*. *Dev Biol*. 190(2): 229-240.

Mehr S, Verdes A, DeSalle R, Sparks J, Pieribone V, Gruber DF. (2015). Transcriptome sequencing and annotation of the polychaete *Hermodice carunculata* (Annelida, Amphinomidae). *BMC Genomics*, 16(1), 445.

Mesnil F, Caullery M. 1919. Sur un processus normal de fragmentation, suivie de régénération chez une Annélide Polychète*, Syllis gracilis* Gn. *Ebenda* 169: *926*.

Meyer E, Aglyamova GV, Wang S, Buchanan-Carter J, Abrego D, Colbourne JK, Willis BL, Matz MV. 2009. Sequencing and *de novo* analysis of a coral larval transcriptome using 454 GSFlx. *BMC Genomics.* 10(1): 1.

Michel A. 1898. Recherches sur la régénération chex les Annélides. *Bull Sci Fr Bel.* 404 pp.

Mino M, Sawada H. 2016. Follicle cell trypsin-like protease Ovochymase: Its cDNA cloning, localization, and involvement in the late stage of oogenesis in the ascidian *Halocynthia roretzi*. *Mol Reprod Dev.* 83(4): 347–358.

Mita M. 2013. *Relaxin*-like gonad-stimulating substance in an echinoderm, the starfish: A novel relaxin system in reproduction of invertebrates. *Gen Comp Endocrinol.* 181: 241–245.

Mita M, Takeshige Y, Nakamura M. 2014. Effect of relaxin-like gonad-stimulating substance on gamete shedding and 1-Methyladenine production in starfish ovaries. In *Sexual Reproduction in Animals and Plants*. 115–122. Springer Japan.

Miyakawa H, Toyota K, Hirakawa I, Ogino Y, Miyagawa S, Oda S, Tatarazako N, Miura T, Colbourne JK, Iguchi T. 2013. A mutation in the receptor Methoprene-tolerant alters juvenile hormone response in insects and crustaceans. *Nat Commun.* 4:1856.

Murta D, Batista M, Trindade A, Silva E, Henrique D, Duarte A, Lopes-da-Costa L. 2014. In vivo *Notch* signaling blockade induces abnormal spermatogenesis in the mouse. *PloS one*. 9(11): e113365.

Neckameyer WS. 1998a. Dopamine and mushroom bodies in *Drosophila*: experience-dependent and-independent aspects of sexual behavior. *Learn Mem.* 5(1): 157-165.

Neckameyer WS. 1998b. Dopamine modulates female sexual receptivity in Drosophila melanogaster. *J Neurogenet* 12(2): 101-114.

Nichols CD. 2007. 5-HT2 receptors in *Drosophila* are expressed in the brain and modulate aspects of circadian behaviors. *Dev Neurobiol*, 67, 752 – 763.

Novo M, Riesgo A, Fernández-Guerra A, Giribet G. 2013. Pheromone evolution, reproductive genes, and comparative transcriptomics in Mediterranean earthworms (Annelida, Oligochaeta, Hormogastridae). *Mol Biol Evol.* 30 (7): 1614–1629.

Nygren, A. 1999. Phylogeny and reproduction in Syllidae (Polychaeta*). Zool J Linn Soc.* 126(3): 365–386.

Okada YK. 1929. Regeneration and fragmentation in the syllidean polychaetes. (Studies on the Syllidae II). *Wilhelm Roux Arch Entwickl.* 115: 542–600.

Okada YK. 1937. La stolonisation et les caractères sexuels du stolon chez les Syllidiens polychètes (É tudes sur les Syllidiens III). *Jpn J Zool.* 7: 441–490.

Olinski RP, Lundin L-G, Hallbook F. 2006. Conserved synteny between the *Ciona* genome and human paralogons identifies large duplication events in the molecular evolution of the *insulin-relaxin* gene family. *Mol Biol Evol.* 23: 10–22

Pacey AA, Bentley MG. 1992. The fatty acid 8, 11, 14-eicosatrienoic acid induces spawning in the male lugworm Arenicola marina. *J Exp Biol*. 1;173(1):165-79.

Pfannenstiel HD, Spiehl D. 1987Dopamine induces sex reversal in females of *Ophryotrocha puerilis* (Polychaeta). *Cell Differ*. 20:84.

Pérez-Portela R, Turon X, Riesgo A. 2016. Characterization of the transcriptome and gene expression of four different tissues in the ecologically relevant sea urchin *Arbacia lixula* using RNA-seq. *Mol Ecol Resour*. 16: 794–808.

Pettibone MH. 1963. Marine polychaete worms of the New England region. 1. Aphroditidae through Trochochaetidae. *Bull US Natl Mus*. 227: 1–346.

Phillips DJ, de Kretser DM. 1998. Follistatin: a multifunctional regulatory protein. *Front Neuroendocrinol*. 19(4):287-322.

Potts FA. 1911. Methods of reproduction in the syllids. *Ergebnisse und Fortschritte der Zoologie* 3(1): 1–72.

Posada, D. and Buckley, T. R. 2004. Model selection and model averaging in phylogenetics: advantages of Akaike information criterion and Bayesian approaches over likelihood ratio tests. *Syst Biol*. 53(5): 793–808.

Randel N, Bezares-Calderón LA, Gühmann M, Shahidi R, Jékely G. 2013. Expression dynamics and protein localization of rhabdomeric opsins in *Platynereis* larvae. *Integr Comp Biol*. 53(1): 7–16.

Rebscher N, Zelada-González F, Banisch TU, Raible F, Arendt D. 2007. *Vasa* unveils a common origin of germ cells and of somatic stem cells from the posterior growth zone in the polychaete *Platynereis dumerilii*. *Dev Biol*. 306: 599–611.

Riddiford LM. 1994. Cellular and molecular actions of juvenile hormone I. General considerations and premetamorphic actions. *Adv Insect Physiol*. 24: 213–274

Riesgo, A., Andrade, S.C.S., Sharma, P.P., Novo, M., Pérez-Porro, A.R., Vahtera, V., González, V.L., Kawauchi, G.Y., and Giribet, G. 2012. Comparative transcriptomics of newly sequenced invertebrates and efficiency estimation of genomic sampling in non-model taxa. *Front Zool* 9: 33.

Rimskaya-Korsakova NN, Kristof A, Malakhov VV, Wanninger A. 2016. Neural architecture of *Galathowenia oculata* Zach, 1923 (Oweniidae, Annelida). *Front Zool*. 13(1): 5.

Rivera AS, Gonsalves FC, Song MH, Norris BJ, Weisblat DA. 2005. Characterization of *Notch*-class gene expression in segmentation stem cells and segment founder cells in *Helobdella*

robusta (Lophotrochozoa; Annelida; Clitellata; Hirudinida; Glossiphoniidae). *Evol Dev.* 7(6): 588–599.

Schenk S, Krauditsch C, Frühauf P, Gerner C, Raible F. 2016. Discovery of methylfarnesoate as the annelid brain hormone reveals an ancient role of sesquiterpenoids in reproduction. *eLife*. 5: e17126.

Schlawny A, Hamann T, Müller MA, Pfannenstiel HD. 1991. The catecholaminergic system of an annelid (Ophryotrocha puerilis, Polychaeta). *Cell Tissue Res*. 265(1): 175-184.

Schroeder PC Hermans CO. 1975. *Annelida: Polychaeta*. *In*: Reproduction of marine invertebrates. A.C. Giese and J.S. Pearse, editors. Academic Press, New York. Vol. 3: 1–213

Siebert S, Juliano CE. 2017. Sex, polyps, and medusae: Determination and maintenance of sex in cnidarians. *Mol Reprod Dev.* 84:105–119.

Shin SW, Zou Z, Saha TT, Raikhel AS. 2012. bHLH-PAS heterodimer of methoprene-tolerant and Cycle mediates circadian expression of juvenile hormone-induced mosquito genes. *Proc Natl Acad Sci*. 109(41): 16576-16581.

Spörhase-Eichmann U, Winkler M, Schürmann FW. 1998. Dopaminergic sensory cells in the epidermis of the earthworm. *Naturwissenschaften*. 85(11): 547-550.

Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690.

Stamatakis A, Hoover P, Rougemont J. 2008. A rapid bootstrap algorithm for the RAxML web servers. *Syst Biol* 57: 758–771.

Sulston J, Dew M, Brenner S. 1975. Dopaminergic neurons in the nematode *Caenorhabditis elegans*. *J Comp Neurol*. 163(2): 215-226.

Takahashi T, McDougall C, Troscianko J, Chen W, Jayaraman-Nagarajan A, Shimeld SM, Ferrier DEK 2009. An EST screen from the annelid *Pomatoceros lamarckii* reveals patterns of gene loss and gain in animals. *BMC Evol Biol.* 9: 240.

Thamm K, Seaver EC. 2008. *Notch* signaling during larval and juvenile development in the polychaete annelid *Capitella* sp. I. *Dev Biol.* 320(1): 304–318.

Tobe SS, Bendena WG. 1999. The regulation of juvenile hormone production in arthropods: functional and evolutionary perspectives. *Ann NY Acad Sci*. 897(1):300-10.

Toyota K, Miyakawa H, Hiruta C, Furuta K, Ogino Y, Shinoda T, Tatarazako N, Miyagawa S, Shaw JR, Iguchi T. Methyl farnesoate synthesis is necessary for the environmental sex determination in the water flea Daphnia pulex. *J Insect Physiol*. 80:22-30.

Verasztó C, Ueda N, Bezares-Calderón LA, Panzera A, Williams EA, Shahidi R, Jékely G. 2017. Ciliomotor circuitry underlying whole-body coordination of ciliary activity in the *Platynereis* larva. *eLife*. 6.

Verger-Bocquet M. 1984. Mise en évidence de l'influence de la photopériode sur la stolonisation, par l'intermédiaire du prostomium chez *Syllis amica* Quatrefages (Annélide, Polychète). *Arch Biol*. 95: 301–306.

Weidhase M, Beckers P, Bleidorn C, Aguado MT. 2016. On the role of the proventricle region in reproduction and regeneration in *Typosyllis antoni* (Annelida: Syllidae). *BMC Evol Biol*. 16(1): 196.

Weinshenker D, Gian G, Thomas JH. 1995. Genetic and pharmacological analysis of neurotransmitters controlling egg laying in *C. elegans*. *J Neurosci*. 15(10): 6975–6985.

Wen D, Rivera-Perez C, Abdou M, Jia Q, He Q, Liu X, Zyaan O, Xu J, Bendena WG, Tobe SS, Noriega FG. 2015. Methyl farnesoate plays a dual role in regulating *Drosophila* metamorphosis. *PLoS Genet*. 11(3), e1005038.

Wicker-Thomas C, Hamann M. 2008. Interaction of dopamine, female pheromones, locomotion and sex behavior in Drosophila melanogaster. *J Insect Physiol*. 54(10): 1423-1431.

Wilson BC, Burnett D, Rappaport R, Parry LJ, Fletcher EK. 2009. *Relaxin-3* and *RXFP3* expression, and steroidogenic actions in the ovary of teleost fish. *Comp Biochem Physiol A Mol Integr Physiol*. 153(1): 69–74.

Winberg S, Nilsson A, Hylland P, Söderstöm V, Nilsson GE. 1997. Serotonin as a regulator of hypothalamic-pituitary-interrenal activity in teleost fish. *Neurosci Lett*. 230(2): 113-116.

Wissocq JC. 1966. Rôle du proventricule dans le déterminism de la stolonisation de *Syllis amica* Quatrefages (Annélide Polychète). *Comptes Rendus Acad Sci*. 262: 2605–2608.

Wissocq JC. 1970. Évolution de la musculature longitudinale dorsale et ventrale au cours de la stolonisation de *Syllis amica* Quatrefages (Annélide polychète): Muscles du ver asexué et muscles du stolon. I. *J Microsc*. 9: 355–358.

Wyatt GR. 1997. Juvenile hormone in insect reproduction - a paradox? *European J Entomol*. 94:323–333.

Wyatt GR, Davey KG. 1996. Cellular and molecular actions of juvenile hormone. II. Roles of juvenile hormone in adult insects. *Adv. Insect Physiol*. 26: 1–155.

Xie X, Tian T, Mingxin L, Yanqi Z, Zhiye L, Dongfa Z. 2016. The potential role of juvenile hormone acid methyltransferase in methyl farnesoate (MF) biosynthesis in the swimming crab *Portunus trituberculatus*. *Anim Reprod Sci*. 168: 40-49.

Xu T, Caron LA, Fehon RG, Artavanis-Tsakonas S. 1992. The involvement of the Notch locus in *Drosophila* oogenesis. *Development*. 115(4): 913-922.

Zaccardi ML, Traina G, Cataldo E, Brunelli M. 2004. Sensitization and dishabituation of swim induction in the leech Hirudo medicinalis: role of serotonin and cyclic AMP. *Behav Brain Res*. 153(2): 317-326.

Pelagic

Development

Spawning and Fertilization

Juvenile

Bottom migration

♀ Stolons ♂

SCHIZOGAMY (STOLONIZATION)

Benthic

Atokous form

♀

Stolon realease

Atokous form

Stolon development

Epitokous forms

♂

Final segment regeneration

FEMALE STOLON vs MALE STOLON

Up regulated in Female stolon

Up regulated in Male stolon

A

B

C

D

Bilateria-conserved
Mevalonate Synthesis
Pathway

↓

Farnesyl Diphosphate

↓

Farnesol

↓

Farnesal

↓

Farnesoic Acid

↓

Methyl Farnesoate

Juvenil Hormone
III Acid

Juvenile Hormones

- Farnesyl Pyrophosphate synthase (FPP)
- Farnesyl Phosphatase (FP)
- Farnesol Oxidase/Dehydrogenase
- Farnesal Dehydrogenase
- Methyl Transferases
- JHAMT
- INSECTS ONLY
- Epoxidase

A

Expression levels (counts)

82 — 35 — 0

REPRO ♀ | REPRO ♂ | NON-REPRO

| | SERt |
| | DAr |

A_0 F_0 AS_0 FS_0 | A_1 P_1 F_1 FS_1 | A_2 P_2 F_2 AS_2 FS_2 | A_3 P_3 F_3 AS_3 FS_3 | A_4 P_4 F_4 | A_5 P_5 F_5 | A_6 P_6 F_6 | A_7 P_7 F_7

replicate1 replicate2 | replicate1 replicate2 | replicate1 replicate2 replicate3 replicate4

B

DAr1 DAr2 α1-adrenergic α2-adrenergic DAr2 Serotonin

Syllis magdalena

Syllis magdalena 2

Biogenic amines receptors

Biogenic amines transporter
(Serotonin, SERT)

2.0

A

RHABDOMERIC OPSINS

- *Strongylocentrotus droebachiensis* Opsin_ABB89040
- *Syllis magdalena* Rhabdomeric opsin-1
- *Platynereis dumerilii* Rhabdomeric opsin-5_AKS48309
- *Platynereis dumerilii* Rhabdomeric opsin-4_AGL94565
- *Syllis magdalena* Rhabdomeric opsin-2
- *Nautilus pomilius* Rhabdomeric opsin_BAR90770
- *Syllis magdalena* Rhabdomeric opsin-5 DE
- *Syllis magdalena* Rhabdomeric opsin-3
- *Platynereis dumerilii* Rhabdomeric opsin_CAC86665
- *Syllis magdalena* Rhabdomeric opsin-4
- *Platynereis dumerilii* Rhabdomeric opsin-2_AKS48308
- *Syllis magdalena* Rhabdomeric opsin-6
- *Platynereis dumerilii* Rhabdomeric opsin-3_AGL94564
- *Helobdella robusta* Opsin C_AID66635
- *Helobdella robusta* Opsin A_AID66633
- *Helobdella robusta* Opsin B_AID66634

CILIARY OPSINS

- *Clytia hemisphaerica* Opsin-9_AUO16427
- *Syllis magdalena* Ciliary opsin-1
- *Danio rerio* Opsin-1, short-wave-sensitive 1_NP_571394
- *Mus musculus* Opsin-1_NP_031564
- *Danio rerio* Opsin-1, short-wave-sensitive 2_NP571267
- *Danio rerio* Opsin-1, green-sensitive_NP_571328
- *Platynereis dumerilii* Ciliary opsin_ASM47595
- *Platynereis dumerilii* Ciliary opsin_AAV63834
- *Syllis magdalena* Ciliary opsin-2
- *Platynereis dumerilii* Go coupled opsin-1_AKS48306
- *Platynereis dumerilii* Go coupled opsin-1_AKS48307
- *Clytia hemisphaerica* Opsin-7_AUO16426
- *Clytia hemisphaerica* Opsin-4_AUO16425

0.6

B

Expression levels (counts)
40,700   20,500   0

REPRO ♀   REPRO ♂   NON-REPRO

Rhabdomeric opsin 6
Rhabdomeric opsin 5*
Rhabdomeric opsin 4
Rhabdomeric opsin 3
Rhabdomeric opsin 2
Rhabdomeric opsin 1
Ciliary opsin 1
Ciliary opsin 2

A_0 F_0 AS_0 FS_0 A_1 P_1 F_1 FS_1 | A_2 P_2 F_2 AS_2 FS_2 A_3 P_3 F_3 AS_3 FS_3 | A_4 P_4 F_4 A_5 P_5 F_5 A_6 P_6 F_6 A_7 P_7 F_7

replicate1 replicate2 | replicate1 replicate2 | replicate1 replicate2 replicate3 replicate4