Specific enrichment of hyperthermophilic electroactive *Archaea* from

deep-sea hydrothermal vent on electrically conductive support.

Guillaume Pillot[1], Eléonore Frouin[1], Emilie Pasero[1], Anne Godfroy[2], Yannick Combet-Blanc[1],

Sylvain Davidson[1] & Pierre-Pol Liebgott[1*]

[1] Aix Marseille Université, IRD, Université de Toulon, CNRS, MIO UM 110, Marseille, France.

[2] Laboratoire de Microbiologie des Environnements Extrêmes - UMR6197 IFREMER, CNRS,

UBO Centre de Brest–CS10070/IUEM, Plouzané, France.

[*]Corresponding author: pierre-pol.liebgott@mio.osupytheas.fr; Mediterranean Institute of

Oceanography, Campus de Luminy, Bâtiment OCEANOMED, 13288 Marseille Cedex 09.

Abstract

While more and more investigations are done to isolate hyperthermophilic exoelectrogenic

communities from environments, none have been performed yet on deep-sea hydrothermal vent.

Samples of black smoker chimney from Rainbow site on the Atlantic mid-oceanic ridge have

been harvested for enriching exoelectrogens in microbial electrolysis cells under hyperthermophilic

(80°C) condition. Two enrichments have been performed: one from direct inoculation of crushed

chimney and the other one from inoculation of a pre-cultivation on iron (III) oxide. In both

experiments, a current production was observed from 2.4 A/m² to 5.8 A/m² with a set anode

potential of +0.05 vs SHE. Taxonomic affiliation of the exoelectrogen communities obtained

exhibited a specific enrichment of *Archaea* from *Thermococcales* and *Archeoglobales* orders on the

electrode, even when both inocula were dominated by *Bacteria*.

Keywords

Hyperthermophilic exoelectrogens, Deep-sea hydrothermal vent, Microbial electrolysis cell,

External Electron Transfer

1. Introduction

Since the discovery of the first deep-sea hydrothermal vent in 1977, many studies have expanded our understanding of extremophilic life forms in those environments. The "black smoker" deep-sea hydrothermal vents, located along the ridges of the Atlantic, Pacific and Indian oceans, are the result of volcanic activities that generate hydrothermal chimneys composed of polymetallic sulfide minerals (heterogeneous pyrite) (Dick et al. 2013). Those environments exhibit dynamic habitats that are characterized by large steep thermal and chemical gradients. These gradients provide a wide range of growth conditions for many extremophilic microorganisms growing as biofilms and being the base of these specific deep-sea ecosystems (Flores et al. 2011; Kristall et al. 2006). In these environments has been shown the presence of thermophilic microorganisms from the *Archaea* domain, mainly belonging to the orders *Thermococcales*, *Methanococcales*, and *Archaeoglobales*, whereas mesophilic and thermophilic microorganisms from the *Bacteria* domain belong to *Epsilon-proteobacteria* (Huber et al. 2010; Vetriani et al. 2014) and to the orders *Aquificales* and *Thermotogales* (Miroshnichenko and Bonch-Osmolovskaya, 2006). Considering the mineralogical composition of these hydrothermal chimneys (polymetallic massive sulfide), it appears likely that a significant proportion of its microbial populations is dependent on energetic metabolisms based on the dissimilatory reduction of insoluble "metals" or sulfurs compounds (Cao et al. 2014; Konn et al. 2015).

As the microbial cell envelope is neither physically permeable to insoluble minerals nor electrically conductive, microorganisms have evolved strategically to exchange electrons with insoluble extracellular minerals, a mechanism known as the extracellular electrons transfer (EET) (Hinks et al. 2017; Shi et al. 2016). The first described EET capable

bacteria were *Shewanella* and *Geobacter* (Shi et al. 2007). The role of membrane bound

electron transport chains in carrying out EET was well acknowledged, but the exact

mechanisms are still not completely understood (Kumar et al. 2017). Most of the

knowledge about EET in microbes is derived from studies in bioelectrochemical systems

(Allen and Bennetto, 1993; Logan et al. 2006; Schröder et al. 2015) which helped to define

a novel group of microorganisms called exoelectrogens. These exoelectrogenic

microorganisms are capable of extracellular electron transfer to a solid electrode, which are

used in Microbial Electrochemical Technologies (METs) such as Microbial Fuel Cells

(MFC) and microbial electrolysis cells (MEC) (Doyle and Marsili, 2015). In the past years,

exoelectrogenic activity has been reported in almost 100 microbial species which are

mostly affiliated with the bacterial phylum *Proteobacteria* (Koch and Harnisch, 2016). It

includes the extensively studied *Geobacter sp.* (Busalmen et al. 2008; Reguera et al. 2006)

and *Shewanella sp.* (Gorby et al. 2006; Marsili et al. 2008), as well as *Cyanobacteria*

(J. McCormick et al. 2011; Sekar et al. 2014). All these microorganisms are mesophilic

and grow optimally at moderate temperatures, ranging from 20°C to 45°C. To date, only

four thermophiles, *Thermincola ferriacetica* (Marshall and May, 2009), *Thermincola*

potens strain JR (Wrighton et al. 2011), *Calditerrivibrio nitroreducens* (Fu et al. 2013) and

Thermoanaerobacter pseudethanolicus (Lusk et al. 2015), all isolated from extreme natural

environments, were shown to generate electricity in MFCs operating at a temperature

higher than 50°C. Moreover, it is only very recently that two hyperthermophilic strains,

Pyrococcus furiosus and *Geoglobus ahangari,* have been shown to have the capacity to

produce electricity in MFC or MEC (Sekar et al. 2017; Yilmazel et al. 2018). This would

be the two first hyperthermophilic archaeon strains described having the capacity of EET.

Several studies reported the enrichment of mixed cultures of efficient thermophilic

exoelectrogens on the anode of METs (Ha et al. 2012; Jong et al. 2006; Mathis et al. 2007;

Wrighton et al. 2008). More recently, two exoelectrogenic biofilms have been enriched at

70°C from a high-temperature petroleum reservoir (Fu et al. 2015) and at 80°C from Red

Sea brine pools (Shehab et al. 2017) on the anode of MFC systems. To the best of our

knowledge, no exoelectrogenic biodiversity enrichment from deep-sea hydrothermal vents

has been achieved yet, while it seems likely that some microbial populations require this

type of energy metabolism to expand in these particular environments. Interestingly, a

MFC was installed at a hydrothermal vent field (Girguis and Holden, 2012) showing for

the first time, the *in situ* electricity generation in those extreme ecosystems. More recently

it has been demonstrated that there is a widespread and distant electron transfer through the

electrically conductive hydrothermal chimney by the internal oxidation of the

hydrothermal fluid coupled to the reduction of oxygenated seawater at the external side of

the chimney (Yamamoto et al. 2017). This electricity generation in deep-sea hydrothermal

systems must affect the surrounding biogeochemical processes and the development of

microbial communities through potential EET-capable microorganisms.

The aim of this study was to promote and identify a part of the exoelectrogenic microbial

community from a hydrothermal chimney of the Rainbow site on the Atlantic mid-oceanic

ridge. For this investigation, two experiments were carried out using a semiautomatic two-

chamber BioElectrochemical System Stirred-Reactor (BES-SR) prototype (graphical

abstract), operating at high temperatures (80°C). The first one was performed to enrich

microbes in our BES-SR system, directly from a chimney fragment. The second

enrichment in BES was carried out from an inoculum obtained by a microbial enrichment

culture in flask on Fe(III) oxide particles as electron acceptor using the same hydrothermal

sample. This was done in order to pre-cultivate potential exoelectrogenic microbes on iron

oxide and observing the impact on biodiversity obtained subsequently in BES. The

evolution of current production, of microbial diversity and biomass during the different

enrichments have been studied through electrochemical method and molecular biology

survey.

2. Material and methods

2.1. Sample collection and preparation

The inoculum used for all the experiments was collected on the Rainbow hydrothermal vent field

(36°14′N, MAR) by the Remote Operated Vehicle VICTOR 6000 during EXOMAR cruise in 2005

led by IFREMER (France) on board RV L'Atalante (Godfroy, 2005) . Sample (EXO8E1) was

collected by breaking off a piece of a high temperature active black smoker using the arm of the

submersible and bring back to the surface into a decontaminated insulated box. On board, chimney

fragments were anaerobically crushed into an anaerobic chamber (La Calhene, France) and stored

into flasks under anaerobic conditions (anoxic seawater at pH 7 with hydrogen sulfide and

N2:H2:CO2 (90:5:5) gas atmosphere) and kept at 4°C. Prior to the experiment, pieces of

hydrothermal chimney were removed from the sulfidic seawater flask, crushed in a sterile mortar

and pestle in an anaerobic chamber (Coy Laboratories Inc.) and distributed in anaerobic tubes for

further different experiments.

2.2. Enrichment on iron oxide in flask

To obtain enrichment of electro-active microbes on insoluble iron (III) oxide as electron acceptor,

2.5g of crushed hydrothermal chimney was inserted in 500 ml flasks under nitrogen gas

atmosphere filled with 250ml of mineral medium at pH 7 containing 30 g/l NaCl, 0.65 g/l KCl, 0.5

g/l NH_4Cl, 0.3 g/l KH_2PO_4, 0.3 g/l K_2HPO_4, 0.1 g/l $MgCl_2$, 0.1 g/l $CaCl_2$, 0.35 g/l Cysteine HCl,

0.2 g/l of yeast extract and 10ml/l of Balch trace mineral solution (Uzarraga et al. 2011). The media

was supplemented with 1g/l of iron (III) oxide, and 10mM of acetate or 4g/l of yeast extract as

electron donor and carbon source. The flasks were incubated at 80°C in static condition until a dark

coloration of iron oxide was observed. Two subcultures were performed in the same condition with

1% of inoculum from the previous culture.

   2.3. Semi-Automated Bioelectrochemical Systems

A prototype of semi-automated BioElectrochemical System in Stirred-Reactor (BES-SR) has been

developed to assess the enrichment of hyperthermophilic electroactive microorganisms. The system

was composed of a two-chamber jacketed glass reactor (Verre Labo Mula, France) with a 1.5L

working volume, thermostated with a Heating Circulator (Julabo SE 6, France) at 80°C ± 1 °C,

and separated by an Anion Exchange membrane (Membrane International Inc.). The working

electrode was a 20 cm² carbon cloth (PaxiTech SAS, France) with a 3M Ag/AgCl reference

electrode and the counter electrode a 20cm² carbon cloth coated with platinum (Hogarth, 1995).

The bioelectrochemical system was connected to a semi-automated platform previously described

(Boileau et al. 2016) to control the composition and rate of gas input (H_2, CO_2, O_2, N_2) with mass

flowmeters (Bronkhorst, Netherlands) and a continuous monitoring of output gas composition (H_2,

N_2, CH_4, O_2) using a micro-GC equipped with a catharometric detector (MS5A, SRA Instrument,

France). To ensure anaerobic condition, the culture medium was continually sparged with a 50

ml/min flow of N_2. The pH was maintained at 7 ± 0.1 by the addition of sodium hydroxide (NaOH

0.5 mmol/L) or hydrochloric acid (HCl 0.5 mmol/L). The stirring, driven by two axial impellers,

was set to 150 rpm. The measurements of pH and ORP (Mettler Toledo InPro 3253, Switzerland),

temperature (Prosensor pt 100, France), stirring, CO_2 with a CARBOCAP CO2 probe (Vaisala

GMT 221, Finland), H_2, N_2, CH_4, O_2, bioreactor liquid volume, NaOH and HCl consumption were

measured and managed by the BatchPro software (Decobecq Automatismes, France). SP-240

potentiostats and EC-Lab software (BioLogic, France) were used to poise the electrodes at a fixed

potential and measure current. All the measured currents are expressed in ampere per square meter

of electrode area.

2.4. Operating conditions in BES-SR

Before each experiment, the bioelectrochemical system was dismantled, washed, and sterilized by

autoclaving at 120 °C for 20 min. Then, the system was connected to the platform and 1,5L of

mineral medium was injected and supplemented with 10 mM acetate and 0.15 g/l of Yeast Extract

(YE). The liquid media was set in the operational condition for a few hours prior to perform a

cyclic voltammetry (CV, 20 mV/s). First, the system (BES1) was inoculated with 15 g of crushed

hydrothermal chimney in anaerobic condition. A chronoamperometry was carried out by the

potentiostat to poise the electrode at +0.05V vs Standard Hydrogen Electrode (SHE), and a

measurement of the current was taken every 10s. A cyclic voltammetry was also performed at the

end of the experiment. Secondly, a new system (BES2) was inoculated in the same condition with

1% of the last enrichments in flask on iron (III) oxide on yeast extract. An abiotic control and an

inoculated but non-polarized control have been performed in the same conditions to exhibit the

exoelectrogenic specificity of the biofilm on the polarized electrode.

2.5. Taxonomic and phylogenetic classification

Taxonomic affiliation was performed according to (Zhang et al. 2016). DNA were extracted from

1g of the crushed chimney and at the end of each experiment : from 1g of liquid from flask

enrichment, on the scrapings from half of the 20cm² of working electrode and from 50ml of the

liquid media of the BES and flask enrichments, which were centrifuged and suspended in 1ml of

sterile water. The DNA extraction was carried out using the MoBio PowerSoil DNA isolation kit

(Carlsbad, CA USA). The V4 region of the 16S rRNA gene was amplified using the universal

primers 515F (5′-GTG CCA GCM GCC GCG GTA A-3′) and 806R (5′-GGA CTA CNN GGG

TAT CTA AT-3′) with Taq&Load MasterMix (Promega) and PCR reactions were carried out using

C1000 Thermal Cycler (BioRad) with the following conditions: initial denaturation at 94°C for 3

min followed by 30 cycles of denaturation at 94°C for 30 s, primers annealing at 50°C for 30 s and

extension at 72°C for 90 s, followed by a final extension step of 5min at 72°C. The amplified gene

regions were sequenced on Illumina MiSeq 2500 platform (GeT-PlaGe, France) to generate paired-

end 150bp reads. The reads were merged using the FLASH software. The taxonomic affiliation was

performed with the QIIME software package v 1.9.1. Chimera were removed from the merged

sequenced using UCHIME Algorithm. Then, the filtered sequences were clustered into OTUs using

the RDP method with a minimum bootstrap confidence of 0.8 and OTUs were affiliated using the

Silva database as reference. To analyze the alpha diversity, the OTU tables were rarified to a

sampling depth of 10770 sequences per library and two metrics were calculated: the richness

component represented by the number of OTUs observed, and the Pielou's index representing the

evenness component of our biodiversity. The construction of clone library was performed by

amplifying the 16S rRNA genes by previously described PCR method with a FD1 (5'-

AGAGTTTGATCCTGGCTCAG-3') and R6 (5'-TACGGCTACCTTGTTACG-3') primer set

using Taq&Load MasterMix (Promega). The amplicons were cloned into the pGEM-T easy vector

(Promega) and transformed into *E. coli* JM109 competent cells that were grown overnight in LB

agar at 37°C. Fragment of 16S rRNA gene (~1300 bp) of the clones were sequenced by ABI3730xl

(GATC Biotech) sequencers.

2.6. Quantitative PCR of archaeal and bacterial 16S rRNA gene copies

Bacterial and archaeal quantification was carried out by qPCR with SsoAdvanced™ Sybr Green

Supermix on a CFX96 Real-Time System (C1000 Thermal Cycler, Bio-Rad Laboratories, CA,

USA) with the primers DGGE300F (5'-GCC TAC GGG AGG CAG CAG-3') and Univ516 (5'-

GTD TTA CCG CGG CKG CTG RCA-3') specific to Bacteria and Arc931F (5'-AGG AAT TGG

CGG GGG AGC A-3') and m1100R (5'-BTG GGT CTC GCT CGT TRC C-3') for Archaea. The

PCR program was composed of a 10s denaturation step at 94°C, a hybridization step of 10s at 55°C

(Bacteria) or 62°C (Archaea) and a 10s elongation step at 72°C, with melting curves performed at

the end of each reaction to ensure product specificity. A standard curve from 10^2 to 10^{10} 16S rRNA

gene copies was obtained by diluting pGEM-T plasmids harboring hyperthermophilic bacterial or

archaeal rRNA gene fragment obtained from microbial community of interest. The results were

expressed in copies number of 16s rRNA gene per gram of crushed chimney, per milliliter of liquid

media or per cm² of surface of electrode.

3. Results and discussion

3.1. Microbial diversity of hydrothermal chimney from Rainbow site of Atlantic Ocean

Prior to study specifically the electroactive community putatively present on a chimney of the

hydrothermal Rainbow site, an analysis of the total microbial diversity present in our crushed

chimney inoculum was done by using the Illumina method. The chimney biodiversity (figure 1)

was composed of 160 OTUs representing a high richness of species. As indicated by the

Equitability index (0.677), the sequences were distributed relatively equally in the different OTUs.

Furthermore, the taxonomic affiliation of the 160 OTUs showed that 66% were assigned to

Bacteria and 33% to *Archaea* domains. This was consistent with the quantitative Polymerase Chain

Reaction (qPCR) assays which showed a dominance of *Bacteria* (figure 2) compared to *Archaea*

(6.38 ± 0.05 vs 4.77 ± 0.19 log of 16S rRNA gene copies per gram of chimney, respectively). The

difference of ratio obtained can be explained by the bacterial quantification by 16S rRNA gene

qPCR, always overestimated because of the high number of gene copies per cells compared to

Archaea where the number of 16S rRNA gene copies is generally equal to 1 per cell (cf. rrnDB,

(Stoddard et al. 2015)). Within the bacteria domain, the OTUs were mainly assigned to 5 phyla:

Proteobacteria (58%), *Firmicutes* (3%), *Aquificae* (1%), *Bacteroidetes* (1%) and

Thermodesulfobacteria (1%) whereas the archaeal OTUs were only identified as *Euryarchaeota*

represented by *Thermococcales* (23%) and *Archaeoglobales* (11%). This taxonomic profile was

substantially similar to those previously reported on Rainbow chimneys (Cerqueira et al. 2017;

Flores et al. 2011), except for the proportion of methanogens and *Desulfurococcales* found

significantly lower in our samples. Despite a recognized stability in the conservation of

hyperthermophilic microorganisms at low temperature (Wirth, 2017), our samples have been

conserved more than 12 years at 4°C, this could mostly explain the decrease in Archaea diversity.

3.2. Enrichment of anode-respiring community in BES from crushed chimney

To assess the diversity of hyperthermophilic exoelectrogenic microbes from deep hydrothermal

vent on conductive electrode, a fraction of a crushed chimney from the Rainbow site was used to

inoculate the BioElectrochemical System (BES1). To mimic the hydrothermal vent conditions in

the BES, a synthetic seawater medium, containing acetate and yeast extract as carbon and energy

sources, was used. Indeed, in hydrothermal vent fluid, acetate can be chemically synthesized

through Fischer-Tropsch Type (FTT) reaction from H_2 and CO_2, both obtained during

serpentinization reaction (Schrenk et al. 2013). Acetate can also be biologically produced during

fermentative metabolisms. It should be noted that acetate is not fermentable and needs an

exogenous electron acceptor to be used as energy source for microorganisms' growth. On the other

hand, YE represent here the organic compounds produced by autotrophic and heterotrophic

community of hydrothermal chimney available for the fermentative/respiratory growth of

microorganisms.

While in the control experiment (sterile and polarized anode) no notable current (0.01 A/m², figure

3) was observed during the 10 days of experiment, a raise of current density was observed after 6

days of incubation in the BES1 inoculated with crushed chimney. The maximum current density

reached 5.9 A/m² at 8.7 days and remained stable for few hours. The 3 mM of acetate consumed

during this period as attested by HPLC measurements (data not shown) was in good agreement

with the current production assuming a maximum faradic efficiency at 95-100%, as previously

obtained in literature (Sengodon and Hays, 2012). After this period the current density decrease

progressively probably due to the exhaustion of growth factors present in the medium and

necessary to the biofilm metabolic activity. Indeed after 10 days of culture, the renewal of the

medium led again to a current increase of up to 6 A/m² at maximum current density. Similarly, the

current decreased after a few hours but remained stable in absence of medium renewal.

Cyclic voltammetries (figure 4) have been performed after inoculation and at the end of experiment

in polarized condition to observe the electrochemical profile of ElectroActive Biofilms (EAB) as

catalyst of the bioelectrochemical oxidation of acetate. The anode showed the apparition of an

oxidation peak with a midpoint potential at +0.06V (vs. SHE) at the end of the experimentation. No

oxidation peak was observed for the solution with a new electrode immersed into the medium

culture, suggesting EET mechanisms not involving mediators (data not shown). Moreover, the

capacitive current (vertical distance between oxidation and reduction waves) increasing between

beginning and end of the culture experiments reports an increase of the electrochemical double

layer and suggests the presence of a biofilm on the surface of the electrode. Thus, these results

suggest the development of EAB with direct EET using acetate as electron donor.

In contrast to the non-polarized culture conditions, microscopic observations over time have

revealed the presence of planktonic microorganisms (in the liquid medium) after 9 days of

incubation in the polarized electrode culture condition. In addition, the microbial quantification by

qPCR of the enrichment in BES1 shows 7.3 ± 0.18 and 9.23 ± 0.04 log of bacterial and archaeal

16S rRNA gene copies per milliliter of liquid media and 7.28 ± 0.53 and 9.73 ± 0.06 log per square

centimeter of electrode, respectively. On the contrary, in the non-polarized control, quantification

of 16S rRNA copies was under the 2 log of sensitivity threshold of the method, indicating that no

significant growth was observed under these conditions in liquid media or non-polarized electrode.

Thus, the polarization of the electrode (+ 0.05V vs SHE), which served as an obligatory electrons

acceptor to allow the oxidation of acetate or the weak amount of YE, seems essential also for the

planktonic microbial growth.

To identify the microbes present on the electrode (Electrode BES1) and in liquid medium (Liquid

BES1), a microbial diversity study through sequencing of the hypervariable V4 region of 16S

rRNA gene (figure 1) have been performed at the end of the enrichment. 61 OTUs were obtained

on Electrode 1 and 58 OTUs in Liquid Media 1, indicating of a loss of biodiversity from

environmental chimney (160 OTUs). The sequences obtained are distributed in the different OTUs

with however some OTUs more represented than in environmental sample (Equitability at 0.452 on

Electrode 1 and 0.331 in Liquid Media 1). This can be explained by our selective conditions of

enrichment with high temperature and substrate specificity. Based on average abundance analysis,

the microbial diversity of enrichment culture was dominated by *Euryarchaeota* phyla (> 70%),

either on electrode or in liquid medium. Interestingly, at the genus level, the dominating archaeal

OTUs on electrode were closely related to *Geoglobus* spp. (45.2 %) and *Thermococcus* spp.

(25.6%) whereas the planktonic microbial diversity in liquid medium is especially dominated by

uncultured *Thermococcus* spp. (64.1%) rather than *Geoglobus* spp. (13.3%). The bacterial diversity

on the electrode and in liquid media were mainly assigned to *Thermodesulfatator* spp. (15.6% and

18% respectively), while the remaining were distributed between members of *Bacillaceae* and

Micrococcaceae families. These specific enrichments of *Archaea* compared to *Bacteria*

demonstrate a shift in initial microbial community structure from the crushed chimney, surely

driven by the specific BES characteristics (i.e., polarized electrode, hyperthermophilic condition).

Due to the limitation for taxonomic affiliation at species level of the Illumina techniques on 300bp

16S fragments, we performed 16S rDNA clone libraries (figure 5) on ~1300bp 16S sequence. It

allowed identifying the *Thermococcus* spp. as closely related to *Thermococcus thioreducens*, *T.*

coalecens, *T. barossi*, *T. peptonophilus*, *T. celer* (98% similarities with them) and an unidentified

Thermococcus species (95% similarity with *Thermococcus thioreducens*). However, because of the

high levels of similarity between species of the genus *Thermococcus*, the taxonomic affiliation

through the coding gene for 16S rRNA is not sufficient to determine the *Thermococcus* species

present in enrichments.

However, among the *Geoglobus* spp. the taxonomic affiliation from clone libraries (figure 5) has

allowed to identify two species: *Geoglobus ahangari* (99% similarity) and a novel genus or species

of the *Archaeoglobaceae* family (MG694212 and MG694224: 95 % of similarity vs *Geoglobus*

ahangari). This novel taxon is exclusively found in the electrode area. Interestingly, this affiliation

has been possible only through near full 16S rRNA gene sequence (1300 pb) alignment. Indeed, the

hypervariable V4 region of the rRNA gene (~290pb), used to the study of microbial diversity by

Illumina-based analysis, did not allow to phylogenetically differentiate this novel taxon from

Geoglobus ahangari. Differences are remarkable at level of hypervariable V6 to V8 regions.

The discovery of *Geoglobus* species enrichment on the electrode is remarkable. The two species of

Geoglobus described so far (*G. ahangari* and *G. acetivorans*) are known to grow autotrophically

using H_2 or heterotropically using a large number of organic compounds with in both cases soluble

or insoluble iron III oxide as a final electrons acceptor (Manzella et al. 2013). More recently, it has

been shown that *Geoglobus ahangari* was exoelectrogenic by direct contact when placed in one-chamber microbial electrolysis cells at 80°C. In contrast to our results, the current produced was particularly weak probably due to its use as an axenic culture, and its lack of enzymatic component to resist to oxidative stress on electrode (Yilmazel et al. 2018). Similarly, the type species of *Geoglobus ahangari* tested in our laboratory had a slow and difficult growth on conductive support and was unable to grow after three subcultures under optimal culture conditions. Thus, observing the presence of *Geoglobus* spp. in our system suggests that the latter requires to be cultured in a consortium.

Phylogenetic analysis of 16S rRNA clone libraries has led to the identification of taxon *Thermodesulfatator atlanticus* among the *Thermodesulfatator spp.* previously identified in high throughput sequencing. This bacterium, that have been also isolated from a Rainbow site chimney, is known to be chemolithoautotrophic, sulfate-reducing obligate bacterium that uses H_2 as electron donor and CO_2 and peptide as carbon source (Alain et al. 2010). A close related species, *Thermodesulfobacterium commune* has already been identified in EAB on electrode of a MFC (Fu et al. 2015)

3.3. Enrichment of hyperthermophilic microorganisms on crystalline iron (III) oxide

In parallel of the first BES previously described, flask enrichments with insoluble metallic electron acceptor were done. Crushed chimney was inoculated in anaerobic mineral medium added with crystalline Fe (III) oxide (α-Fe_2O_3: hematite) as electron acceptor (Wahid and Kamalam, 1993), acetate (10 mM) and yeast extract (4 g/l) as energy and carbon source. After 2 days of static incubation at 80°C, the hematite is completely reduced to black compounds (magnetic particles) and the head space of the flask contained 32% of CO_2, 0.2% of H_2 and 10% of CH_4. A production of acetate was also observed in the liquid media, reaching 12 mM at the end of the experiment. The next subcultures reached the black coloration in less than 24 hours with similar concentration of each gas in the headspace of the flasks and same profile of metabolites in the liquid media. The presence of a thin biofilm could be observed on the surface of the agglomerated iron oxide. An

abiotic control with a mix of acetate, yeast extract, H_2, CH_4 and CO_2 didn't show black coloration of

iron oxide after one month of incubation at 80°C. These results suggest the development of a

community of fermentative organisms producing acetate and electroactive microorganisms able to

oxidize organic compounds brought by the YE (peptides, carbohydrates, etc.) and acetate in the

culture medium for the reduction of crystalline iron (III) oxide.

The biodiversity enriched in flask was composed of 29 OTUs, and the sequences were relatively

concentrated in dominant OTUs (equitability at 0.292). At the family level, the microbial diversity

was dominated by bacterial OTUs related to *Clostridiaceae* (73%), and *Thermodesulfobacteriaceae*

(3%). The archaeal OTUs represented only 24% of the biodiversity distributed between

Thermococcaceae (20.5 %), *Methanococcaceae* (3%) identified as *Methanotorris igneus* species

and *Archaeoglobaceae* (0.5%). The quantification showed an enrichment of 9 log of bacterial and 7

log of archaeal 16S rRNA gene copies which demonstrated a shift of enrichment toward the

preferential bacterial growth in opposition with BES enrichment. Interestingly, the taxonomic

affiliation showed that members of the *Clostridiaceae* OTUs were closely related in clone libraries

to *Caloranaerobacter ferrireducens* (97% of identity), fermentative microorganism which is

known to use crystalline iron (III) reduction as a minor pathway for electron flow while fermenting

sugars or amino acids to a mixture of volatile fatty acids (acetate, butyrate) and hydrogen (Zheng et

al. 2015). *Archaeoglobaceae* accounted for only 0.5% of the microbial communities in this

enrichment and have been closely related from clone libraries analysis to *Geoglobus ahangari*

(99% of identity). As mentioned above, *Geoglobus ahangari* is known to reduce amorphous iron

(III) oxide and these culture conditions with crystalline form do not seem to favor its growth

(Wahid & Kamalan, 1993). Thus, these results could suggest the organization of a kind of trophic

chain in our flask enrichment. The new iron reducing species close to *Caloranaerobacter*

ferrireducens would have used organic compounds from yeast extract to reduce crystalline iron

(III) oxide. The *Thermococcus* would have grown on peptides or carbohydrates from yeast extract

or partly produced by *Firmicutes,* and both would have produced H_2. The methanogens such as

Methanotorris igneus would then have used the H_2 to produce methane.

3.4. Solid electron acceptor determine the microbial ecology of electroactive enrichment

In this context, a new BioElectrochemical System (BES2) was inoculated with the flask enrichment on YE/acetate to assess the specificity of the microbial community obtained on the electrode. Experimental conditions were exactly the same as for BES1. Contrary to BES1, a lag time of 3 days (figure 3) has been observed before the increase of the current. After 4.5 days the current has reached 2.4 A/m² and stayed stable during about 1 day before decreasing rapidly. Phase contrast microscopic observations during the replacement of the culture medium in BES2 have shown the only presence of irregular coccoid cells. Thereafter, the current density reached a maximum of 3.4 A/m² at 4 days after the renewal of the culture medium. This phase of current production came along with a decrease of acetate as observed in the BES1 (data not shown). Cyclic voltammetry (figure 4) exhibit oxidation peak appearing between initial time and final time with a midpoint potential at +0.053 V (vs. SHE), slightly offset compared to the one obtained with BES1.

The biodiversity indexes between YE enrichment in flask and BES2 showed a loss of richness of species from 29 to 26 OTUs, and equitability from 0.292 to about 0.17, indicating a loss of biodiversity dominated only by a few OTUs. Furthermore, only *Archaea* were detected in BES2, when they were trivial in Flask enrichment. The dominating OTUs (figure 1) were closely related to *Thermococcaceae* uncultured species at 87.2% on electrode and 95.4% in liquid media of total OTUs. The remaining archaeal diversity was composed of *Archaeaoglobaceae* spp. (12.7% and 4.5%), with clones (figure 5a) affiliated to *Geoglobus ahangari* (99% identity) and unknown species belonging to *Archaeoglobaceae* family (95% identity with *Geoglobus ahangari*) similarly found on BES1 electrode. No *Caloranaerobacter* spp. or *Thermodesulfobacteriaceae* spp. growth was observed on the BES2 neither on electrode nor in liquid media. This is supported by the qPCR, which has shown a drastic difference of 4 log between archaeal (9 log) and bacterial (5 log) 16S rRNA copies per milliliter of liquid media and square meter of electrode.

It should be pointed out that the flask YE enrichment did not promote growth of *Archaeoglobales*

or *Archaea* in general, whereas they seemed to be re-enriched in the BES2 conditions. These

results suggest the specific enrichment on polarized electrode of *Archaea* belonging to

Thermococcales and *Archeoglobales* in our BES condition. It is noteworthy that the biodiversity

indexes showed a loss of biodiversity in BES compared to crushed chimney due to selective

condition of our experiment to enrich preferentially the electroactive microorganisms. However, in

each BES, the richness and the equitability are more important on polarized electrode than in liquid

media, suggesting a specific enrichment of more diverse species and better distribution of microbial

population on polarized support. According to the absence of growth in non-polarized electrode

and the more important diversity on electrode than in liquid, the microbial diversity found in liquid

media would arise from EAB which develop on the electrode. This is supported by the really low

H_2 production in BES (data not shown) - H2 being normally produced by *Thermococcus* sp. during

fermentative metabolisms - which could be explained by a lack of carbon and energy source in

liquid media. Thus, in absence of possible known metabolism for *Thermococcus* species in liquid,

we would suggest *Thermococcus* cells to be in quiescent conditions after it is released from BEA.

Its growth would thus have an obligatory dependence on the polarized electrode.

If the presence of *Geoglobus* species is consistent with its known capacity to transfer EET with

acetate as electron donor (Yilmazel et al. 2018), *Thermoccoccus* species abundance on polarized

electrode is more surprising. *Thermoccocales* are known to share an energetic peptidic or

carbohydrate metabolisms, associated necessarily or not with the reduction of elemental sulfur

(Bertoldo and Antranikian, 2006). In addition, it was shown that some species have the capacity to

growth by using Extracellular Polymeric Substances (EPS) of microbial origin (dextran, pullulan,

peptides, etc.) (Legin et al. 1998). So far, no respiration ability on electrode or direct EET

mechanism have been reported about this archaeal genera, even if some *Thermococcus* species

have been shown to reduce amorphous iron (III) oxide. However, the mechanism is still unclear

and could be mediated through electron transfer to humic substances and other extracellular

quinones (Lovley et al. 2000, Slobdokin et al. 2001). Furthermore, a recent study has highlighted

the possibility of hydroquinone for microbial electron transfer to electrically conductive minerals,

especially with pyrite composing hydrothermal chimney, or electrode (Taran, 2017). Nevertheless,

no current was obtained with replacement of electrode with spent media, indicating no mediated

electron transport in our condition. However, species of *Thermococcus* genus have been reported to

produce nanopods or nanotubes by budding of their cell envelope but their function have not been

fully understood (Marguet et al. 2013). As suggested by the authors, these archaeal nanopods or

nanotubes could be used to expand the metabolic sphere around cells, promote intercellular

communication or act in sulfur detoxifying mechanisms (Gorlas et al. 2015). We can therefore

suggest a role of these nanotubes in EET as previously described on *Shewanella oneidensis* with

the formation of conductive nanofilaments by extensions of the outer membrane. This bacterial

EET mechanism involves cytochrome-c on the external membrane, not present in *Thermococcus*

genome, allowing the electron transport along the filament to conductive support (Pirbadian et al.

2014). Thus, *Thermococcus* sp. could potentially be exoelectrogenic microorganisms through a still

unknown mechanism.

Remarkably, the obligatory presence of both Thermococcus and Geoglobus species found in each

BES on the polarized electrode did not seem fortuitous. Assuming that the *Thermococcus* species

found in our enrichment are heterotroph, after consumption of YE traces, their only carbon source

available would be the EPS or organic compounds produced by *Geoglobus* spp. present on

polarized electrode. Then, their fermentative metabolism would lead to the production of acetate,

H_2 and CO_2 which could then be used by *Geoglobus* species to grow using the electrode as the

ultimate electrons acceptor. However, recent data have shown the inability of *Geoglobus* to transfer

electrons to an electrode from H_2 (Yilmazel et al. 2018). Previous studies have shown that some

electroactive bacteria are also able to grow syntrophically with other microorganisms via direct

interspecies electron transfer (DIET) (Shrestha and Rotaru, 2014). As explained previously,

Geoglobus ahangari has shown a weak electron transfer capacity when grown in pure culture. The

higher current density obtained in our conditions would suggest that the *Geoglobus sp.* and

Thermoccocales enriched on our polarized electrodes live syntrophically to improve their growth,

and subsequently increased the quantity of electrons transferred to the electrode. Thus we would

suggest that there is a syntrophic mechanism, with potentially DIET, between *Thermococcus* and

Geoglobus in deep hydrothermal vents.

4. Conclusion

This study is the first to report on the enrichment of electroactive consortium in ex-situ conditions

that mimick the conductive chimney of a hydrothermal vent with polarized carbon cloth in

anaerobic artificial seawater at 80°C. Moreover, we demonstrate the specificity of enrichment of

Bacteria on Iron (III) oxide compared to the enrichment of *Archaea* (mainly *Thermococcus sp.* and

Geoglobus sp.) on BES, where the biodiversity is more conserved. These results comfort the

hypothesis of electroactivity as a well-represented metabolism in this type of environment by

confirming the presence of exoelectrogenic microorganisms capable of external electron transfer to

conductive support.

Acknowledgements

This work was financially supported by the National Interdisciplinary Program of CNRS "Initiative

structurante EC2CO" (MICROBIEN). The authors thank Erwan Roussel (LM2E, IFREMER Brest)

for helpful suggestions.

References

Alain, K., Postec, A., Grinsard, E., Lesongeur, F., Prieur, D., Godfroy, A., 2010.

Thermodesulfatator atlanticus sp. nov., a thermophilic, chemolithoautotrophic, sulfate-

reducing bacterium isolated from a Mid-Atlantic Ridge hydrothermal vent. Int. J. Syst.

Evol. Microbiol. 60, 33–38. https://doi.org/10.1099/ijs.0.009449-0

Allen, R.M., Bennetto, H.P., 1993. Microbial fuel-cells. Appl. Biochem. Biotechnol. 39–40, 27–40.

https://doi.org/10.1007/BF02918975

Boileau, C., Auria, R., Davidson, S., Casalot, L., Christen, P., Liebgott, P.-P., Combet-Blanc, Y.,

2016. Hydrogen production by the hyperthermophilic bacterium *Thermotoga maritima* part

I: effects of sulfured nutriments, with thiosulfate as model, on hydrogen production and growth. Biotechnol. Biofuels 9, 269. https://doi.org/10.1186/s13068-016-0678-8

Busalmen, J.P., Esteve-Núñez, A., Berná, A., Feliu, J.M., 2008. C-Type Cytochromes Wire Electricity-Producing Bacteria to Electrodes. Angew. Chem. 120, 4952–4955. https://doi.org/10.1002/ange.200801310

Cao, H., Wang, Y., Lee, O.O., Zeng, X., Shao, Z., Qian, P.-Y., 2014. Microbial Sulfur Cycle in Two Hydrothermal Chimneys on the Southwest Indian Ridge. mBio 5, e00980-13-e00980-13. https://doi.org/10.1128/mBio.00980-13

Cerqueira, T., Pinho, D., Froufe, H., Santos, R.S., Bettencourt, R., Egas, C., 2017. Sediment Microbial Diversity of Three Deep-Sea Hydrothermal Vents Southwest of the Azores. Microb. Ecol. 74, 332–349. https://doi.org/10.1007/s00248-017-0943-9

Dick, G.J., Anantharaman, K., Baker, B.J., Li, M., Reed, D.C., Sheik, C.S., 2013. The microbiology of deep-sea hydrothermal vent plumes: ecological and biogeographic linkages to seafloor and water column habitats. Front. Microbiol. 4. https://doi.org/10.3389/fmicb.2013.00124

Doyle, L.E., Marsili, E., 2015. Methods for enrichment of novel electrochemically-active microorganisms. Bioresour. Technol. https://doi.org/10.1016/j.biortech.2015.07.025

Flores, G.E., Campbell, J.H., Kirshtein, J.D., Meneghin, J., Podar, M., Steinberg, J.I., Seewald, J.S., Tivey, M.K., Voytek, M.A., Yang, Z.K., Reysenbach, A.-L., 2011. Microbial community structure of hydrothermal deposits from geochemically different vent fields along the Mid-Atlantic Ridge. Environ. Microbiol. 13, 2158–2171. https://doi.org/10.1111/j.1462-2920.2011.02463.x

Fu, Q., Fukushima, N., Maeda, H., Sato, K., Kobayashi, H., 2015. Bioelectrochemical analysis of a hyperthermophilic microbial fuel cell generating electricity at temperatures above 80 °C. Biosci. Biotechnol. Biochem. 0, 1–7. https://doi.org/10.1080/09168451.2015.1015952

Fu, Q., Kobayashi, H., Kawaguchi, H., Wakayama, T., Maeda, H., Sato, K., 2013. A Thermophilic Gram-Negative Nitrate-Reducing Bacterium, *Calditerrivibrio nitroreducens*, Exhibiting

Electricity Generation Capability. Environ. Sci. Technol. 47, 12583–12590. https://doi.org/10.1021/es402749f

Girguis, P.R., Holden, J.F., 2012. On the Potential for Bioenergy and Biofuels from Hydrothermal Vent Microbes. Oceanography.

Godfroy, A., 2005. EXOMAR cruise, RV L'Atalante [WWW Document]. URL http://dx.doi.org/10.17600/5010100

Gorby, Y.A., Yanina, S., McLean, J.S., Rosso, K.M., Moyles, D., Dohnalkova, A., Beveridge, T.J., Chang, I.S., Kim, B.H., Kim, K.S., Culley, D.E., Reed, S.B., Romine, M.F., Saffarini, D.A., Hill, E.A., Shi, L., Elias, D.A., Kennedy, D.W., Pinchuk, G., Watanabe, K., Ishii, S., Logan, B., Nealson, K.H., Fredrickson, J.K., 2006. Electrically conductive bacterial nanowires produced by *Shewanella oneidensis* strain MR-1 and other microorganisms. Proc. Natl. Acad. Sci. 103, 11358–11363. https://doi.org/10.1073/pnas.0604517103

Ha, P.T., Lee, T.K., Rittmann, B.E., Park, J., Chang, I.S., 2012. Treatment of Alcohol Distillery Wastewater Using a Bacteroidetes-Dominant Thermophilic Microbial Fuel Cell. Environ. Sci. Technol. 46, 3022–3030. https://doi.org/10.1021/es203861v

Hinks, J., Zhou, M., Dolfing, J., 2017. Microbial Electron Transport in the Deep Subsurface, in: Chénard, C., Lauro, F.M. (Eds.), Microbial Ecology of Extreme Environments. Springer International Publishing, Cham, pp. 81–102.

Hogarth, M.P., 1995. The development of the direct methanol fuel cell. (Ph.D.). University of Newcastle upon Tyne.

Huber, J.A., Cantin, H.V., Huse, S.M., Mark Welch, D.B., Sogin, M.L., Butterfield, D.A., 2010. Isolated communities of *Epsilonproteobacteria* in hydrothermal vent fluids of the Mariana Arc seamounts: *Epsilonproteobacteria* in vents of the Mariana Arc. FEMS Microbiol. Ecol. no-no. https://doi.org/10.1111/j.1574-6941.2010.00910.x

J. McCormick, A., Bombelli, P., M. Scott, A., J. Philips, A., G. Smith, A., C. Fisher, A., J. Howe, C., 2011. Photosynthetic biofilms in pure culture harness solar energy in a mediatorless

bio-photovoltaic cell (BPV) system. Energy Environ. Sci. 4, 4699–4709.

https://doi.org/10.1039/C1EE01965A

Jong, B.C., Kim, B.H., Chang, I.S., Liew, P.W.Y., Choo, Y.F., Kang, G.S., 2006. Enrichment,

Performance, and Microbial Diversity of a Thermophilic Mediatorless Microbial Fuel Cell.

Environ. Sci. Technol. 40, 6449–6454. https://doi.org/10.1021/es0613512

Koch, C., Harnisch, F., 2016. Is there a Specific Ecological Niche for Electroactive

Microorganisms?   ChemElectroChem   3,   1282–1295.

https://doi.org/10.1002/celc.201600079

Konn, C., Charlou, J.L., Holm, N.G., Mousis, O., 2015. The Production of Methane, Hydrogen,

and Organic Compounds in Ultramafic-Hosted Hydrothermal Vents of the Mid-Atlantic

Ridge. Astrobiology 15, 381–399. https://doi.org/10.1089/ast.2014.1198

Kristall, B., Kelley, D.S., Hannington, M.D., Delaney, J.R., 2006. Growth history of a diffusely

venting sulfide structure from the Juan de Fuca Ridge: A petrological and geochemical

study: DIFFUSELY VENTING SULFIDE STRUCTURE. Geochem. Geophys.

Geosystems 7, n/a-n/a. https://doi.org/10.1029/2005GC001166

Kumar, A., Hsu, L.H.-H., Kavanagh, P., Barrière, F., Lens, P.N.L., Lapinsonnière, L., V, J.H.L.,

Schröder, U., Jiang, X., Leech, D., 2017. The ins and outs of microorganism–electrode

electron transfer reactions. Nat. Rev. Chem. 1, 24. https://doi.org/10.1038/s41570-017-

0024

Legin, E., Copinet, A., Duchiron, F., 1998. Production of thermostable amylolytic enzymes by

Thermococcus hydrothermalis. Biotechnol. Lett. 20, 363–367.

https://doi.org/10.1023/A:1005375213196

Logan, B.E., Hamelers, B., Rozendal, R., Schröder, U., Keller, J., Freguia, S., Aelterman, P.,

Verstraete, W., Rabaey, K., 2006. Microbial fuel cells☐: Methodology and technology.

Environ. Sci. Technol. 40, 5181–5192.

Lusk, B.G., Khan, Q.F., Parameswaran, P., Hameed, A., Ali, N., Rittmann, B.E., Torres, C.I., 2015.

Characterization of Electrical Current-Generation Capabilities from Thermophilic

Bacterium *Thermoanaerobacter pseudethanolicus* Using Xylose, Glucose, Cellobiose, or Acetate with Fixed Anode Potentials. Environ. Sci. Technol. 49, 14725–14731. https://doi.org/10.1021/acs.est.5b04036

Marshall, C.W., May, H.D., 2009. Electrochemical evidence of direct electrode reduction by a thermophilic Gram-positive bacterium, *Thermincola ferriacetica*. Energy Environ. Sci. 2, 699. https://doi.org/10.1039/b823237g

Marsili, E., Baron, D.B., Shikhare, I.D., Coursolle, D., Gralnick, J.A., Bond, D.R., 2008. *Shewanella* secretes flavins that mediate extracellular electron transfer. Proc. Natl. Acad. Sci. 105, 3968–3973. https://doi.org/10.1073/pnas.0710525105

Mathis, B.J., Marshall, C.W., Milliken, C.E., Makkar, R.S., Creager, S.E., May, H.D., 2007. Electricity generation by thermophilic microorganisms from marine sediment. Appl. Microbiol. Biotechnol. 78, 147–155. https://doi.org/10.1007/s00253-007-1266-4

Miroshnichenko, M.L., Bonch-Osmolovskaya, E.A., 2006. Recent developments in the thermophilic microbiology of deep-sea hydrothermal vents. Extremophiles 10, 85–96. https://doi.org/10.1007/s00792-005-0489-5

Reguera, G., Nevin, K.P., Nicoll, J.S., Covalla, S.F., Woodard, T.L., Lovley, D.R., 2006. Biofilm and Nanowire Production Leads to Increased Current in *Geobacter sulfurreducens* Fuel Cells. Appl. Environ. Microbiol. 72, 7345–7348. https://doi.org/10.1128/AEM.01444-06

Schrenk, M.O., Brazelton, W.J., Lang, S.Q., 2013. Serpentinization, Carbon, and Deep Life. Rev. Mineral. Geochem. 75, 575–606. https://doi.org/10.2138/rmg.2013.75.18

Schröder, U., Harnisch, F., Angenent, L.T., 2015. Microbial electrochemistry and technology: terminology and classification. Energy Env. Sci 8, 513–519. https://doi.org/10.1039/C4EE03359K

Sekar, N., Umasankar, Y., P. Ramasamy, R., 2014. Photocurrent generation by immobilized cyanobacteria via direct electron transport in photo-bioelectrochemical cells. Phys. Chem. Chem. Phys. 16, 7862–7871. https://doi.org/10.1039/C4CP00494A

Sekar, N., Wu, C.-H., Adams, M.W.W., Ramasamy, R.P., 2017. Electricity generation by

Pyrococcus furiosus in microbial fuel cells operated at 90°C. Biotechnol. Bioeng. 114,

1419–1427. https://doi.org/10.1002/bit.26271

Sengodon, P., Hays, D.B., 2012. Microbial Fuel Cells. Future Fuel Technologies, National

Petroleum Council (NPC) Study

Shehab, N.A., Ortiz-Medina, J.F., Katuri, K.P., Hari, A.R., Amy, G., Logan, B.E., Saikaly, P.E.,

2017. Enrichment of extremophilic exoelectrogens in microbial electrolysis cells using Red

Sea brine pools as inocula. Bioresour. Technol. 239, 82–86.

https://doi.org/10.1016/j.biortech.2017.04.122

Shi, L., Dong, H., Reguera, G., Beyenal, H., Lu, A., Liu, J., Yu, H.-Q., Fredrickson, J.K., 2016.

Extracellular electron transfer mechanisms between microorganisms and minerals. Nat.

Rev. Microbiol. 14, 651–662. https://doi.org/10.1038/nrmicro.2016.93

Shi, L., Squier, T.C., Zachara, J.M., Fredrickson, J.K., 2007. Respiration of metal (hydr)oxides by

Shewanella and *Geobacter*: a key role for multihaem c-type cytochromes. Mol. Microbiol.

65, 12–20. https://doi.org/10.1111/j.1365-2958.2007.05783.x

Shrestha, P.M., Rotaru, A.-E., 2014. Plugging in or going wireless: strategies for interspecies

electron transfer. Front. Microbiol. 5, 237. https://doi.org/10.3389/fmicb.2014.00237

Stoddard, S.F., Smith, B.J., Hein, R., Roller, B.R.K., Schmidt, T.M., 2015. rrnDB: improved tools

for interpreting rRNA gene abundance in bacteria and archaea and a new foundation for

future development. Nucleic Acids Res. 43, D593-598.

https://doi.org/10.1093/nar/gku1201

Uzarraga, R., Auria, R., Davidson, S., Navarro, D., Combet-Blanc, Y., 2011. New Cultural

Approaches for Microaerophilic Hyperthermophiles. Curr. Microbiol. 62, 346–350.

https://doi.org/10.1007/s00284-010-9712-4

Vetriani, C., Voordeckers, J.W., Crespo-Medina, M., O'Brien, C.E., Giovannelli, D., Lutz, R.A.,

2014. Deep-sea hydrothermal vent *Epsilonproteobacteria* encode a conserved and

widespread nitrate reduction pathway (Nap). ISME J. 8, 1510–1521.

https://doi.org/10.1038/ismej.2013.246

Wirth, R., 2017. Colonization of Black Smokers by Hyperthermophilic Microorganisms. Trends

Microbiol. 25, 92–99. https://doi.org/10.1016/j.tim.2016.11.002

Wrighton, K.C., Agbo, P., Warnecke, F., Weber, K.A., Brodie, E.L., DeSantis, T.Z., Hugenholtz,

P., Andersen, G.L., Coates, J.D., 2008. A novel ecological role of the Firmicutes identified

in thermophilic microbial fuel cells. ISME J. 2, 1146–1156.

https://doi.org/10.1038/ismej.2008.48

Wrighton, K.C., Thrash, J.C., Melnyk, R.A., Bigi, J.P., Byrne-Bailey, K.G., Remis, J.P., Schichnes,

D., Auer, M., Chang, C.J., Coates, J.D., 2011. Evidence for Direct Electron Transfer by a

Gram-Positive Bacterium Isolated from a Microbial Fuel Cell. Appl. Environ. Microbiol.

77, 7633–7639. https://doi.org/10.1128/AEM.05365-11

Yamamoto, M., Nakamura, R., Kasaya, T., Kumagai, H., Suzuki, K., Takai, K., 2017. Spontaneous

and Widespread Electricity Generation in Natural Deep-Sea Hydrothermal Fields. Angew.

Chem. Int. Ed. 56, 5725–5728. https://doi.org/10.1002/anie.201701768

Yilmazel, Y.D., Zhu, X., Kim, K.-Y., Holmes, D.E., Logan, B.E., 2018. Electrical current

generation in microbial electrolysis cells by hyperthermophilic archaea *Ferroglobus*

placidus and *Geoglobus ahangari*. Bioelectrochemistry 119, 142–149.

https://doi.org/10.1016/j.bioelechem.2017.09.012

Zhang, L., Kang, M., Xu, J., Xu, J., Shuai, Y., Zhou, X., Yang, Z., Ma, K., 2016. Bacterial and

archaeal communities in the deep-sea sediments of inactive hydrothermal vents in the

Southwest India Ridge. Sci. Rep. 6. https://doi.org/10.1038/srep25982

Figure 1: Dominant taxonomic affiliation and biodiversity indices of microbial communities from Crushed Chimney, Flask enrichment on iron (III) oxide, Electrode and Liquid media from BES1 and BES2. OTUs representing lower than 0.5% of total sequences of the sample was grouped as Rares OTU. Biodiversity indexes of Richness and Equitability represent the number of observed OTU and the Pielou's J' evenness index respectively.

Figure 2: Quantification of 16S rRNA gene copies from Bacteria (orange) or Archaea (yellow) per gram of crushed chimney, per milliliter of liquid or per cm² of working electrode. The red line represents the minimum threshold of sensitivity of the qPCR method. Error bar represent the standard deviation obtained on triplicates.

Figure 3: Current monitoring (A/m²) of BES1 Enrichment inoculated with crushed hydrothermal chimney (blue line), BES2 Enrichment inoculated with flask subculture (green line) and controls (orange line). Arrows represent the renewal of the liquid media.

Figure 4: Cyclic Voltammetries between -0,6V and +0,6V at 20mV/s of Working Electrode at initial time (dot lines) and final time (full line) in BES1 (blue line) and BES2 (orange line).

A

Figure 5: Phylogenetic tree of archaeal (A) and bacterial (B) 16S rRNA clones library from Flask enrichment and Electrode (WE) or liquid media (LM) of BES1 and BES2. Percentages at nodes are bootstrap values based on 500 replications. Scale bar indicated substitutions per site.